

**Seesaw parametrization for  $n$  right-handed neutrinos**

Julian Heeck\*

*Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany*

(Received 31 July 2012; published 26 November 2012)

Introducing  $n$  right-handed neutrinos to the Standard Model yields, in general, massive active neutrinos. We give explicit parametrizations for the involved mixing and coupling matrices in terms of physical parameters for both the top-down and the bottom-up approach for arbitrary  $n$ . Bounds on the complex mixing angles in the bottom-up approach from perturbativity of the Yukawa couplings to charged lepton flavor violation are discussed. As a novel illustration of possible effects from  $n \neq 3$ , we extend the neutrino anarchy framework to arbitrary  $n$ ; we show that while the anarchic mixing angles are insensitive to the number of singlets, the observed ratios of neutrino masses prefer small  $n$  for the simplest linear measure.

DOI: [10.1103/PhysRevD.86.093023](https://doi.org/10.1103/PhysRevD.86.093023)

PACS numbers: 14.60.Pq, 14.60.St

**I. INTRODUCTION**

The (type-I) seesaw mechanism [1] is arguably the simplest and best motivated framework not only to give neutrinos mass but also to explain the smallness of said masses compared to the electroweak scale. The necessary right-handed neutrino partners are assumed to be heavy to suppress the active neutrino masses, which also allows for baryogenesis through leptogenesis [2] via the decay of the heavy states, thus explaining the baryon asymmetry of the Universe. While initially motivated in the context of grand unified theories, the seesaw mechanism has since been applied in its own right. The number of right-handed neutrinos, transforming as singlets of the Standard Model (SM) gauge group, is then no longer constrained by anomaly cancellations or the need to fill up an irreducible representation, but rather by a free parameter. At least two right-handed neutrinos are necessary to reproduce the observed neutrino mass-squared differences, as well as leptogenesis [3]. There is, however, no upper bound, and models with  $\mathcal{O}(10^2\text{--}10^3)$  singlets have been studied in the context of leptogenesis [4], lepton flavor violation (LFV) [5] and as a way to explain the large mixing angles of the active neutrinos [6]. Like for most things in life, there is also a motivation from string theory [7]. The number of singlets is formally infinite in extra-dimensional theories, as right-handed neutrinos are not restricted to the SM-brane and will thus lead to a tower of Kaluza-Klein excitations [8].

Still lacking, however, is a proper parametrization of the arising mixing and coupling matrices in terms of physical quantities, as needed, for example, for efficient parameter scans (as the number of redundant parameters explodes for large  $n$ ). The goal of this work is to provide exactly these parametrizations for arbitrary  $n$ , both for the most general case (top-down approach, to some degree discussed in Ref. [9]) and the seesaw limit (bottom-up).

The rest of this paper is organized as follows: we first fix our notation in Sec. II, discuss the top-down

parametrization in Sec. III, and the bottom-up parametrization in Sec. IV, deriving constraints on the involved parameters from perturbativity and LFV. To illustrate possible effects of  $n$  singlets, we discuss basis independence (neutrino anarchy) in this framework and present the resulting distributions for the neutrino masses in Sec. V. Appendix A gives a brief introduction to the group  $U(N)$  and various representations of its elements, as needed for our parametrizations. Appendix B collects and extends basis-independent measures for miscellaneous matrices necessary for the discussion of anarchy. Finally, Appendix C is devoted to the somewhat special case  $n = 2$ , which would interrupt the flow if included in the main text, which will only cover  $n \geq 3$ .

**II. FRAMEWORK AND NOTATION**

Introducing  $n$  SM singlet right-handed neutrinos  $N_j$  to the Standard Model modifies the Lagrangian by the following terms:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + i\bar{N}_j \not{\partial} N_j - \left( \bar{N}_j (Y_\nu)_{j\alpha} \tilde{H}^\dagger L_\alpha + \frac{1}{2} \bar{N}_j (\mathcal{M}_R)_{jk} N_k^c + \text{H.c.} \right), \quad (1)$$

where a sum over  $\alpha = e, \mu, \tau$  (sometimes 1, 2, 3 in the following) and  $j, k = 1, 2, \dots, n$ , is understood and will in the following often be denoted in a vector notation (e.g., as  $\bar{N} \mathcal{M}_R N^c$ ). Without loss of generality, we work in a basis where the charged lepton mass matrix is diagonal, which can be accomplished by unitary transformations of the lepton fields.  $\mathcal{M}_R$  can be diagonalized by an  $n \times n$  unitary matrix  $V$  like  $V^T \mathcal{M}_R V = \text{diag}$ . This merely redefines  $Y_\nu$ , so from now on we will work in a basis where  $\mathcal{M}_R$  is diagonal, with positive real entries  $M_i$ .<sup>1</sup> After the Higgs doublet  $H$  acquires its vacuum expectation value  $v$ , the

<sup>1</sup>This is not only convenient but also necessary to eliminate unphysical parameters. To this effect we note that a diagonal  $\mathcal{M}_R$  can store  $\mathcal{O}(n)$  parameters (together with  $Y_\nu$ ), while a nondiagonal  $\mathcal{M}_R$  has  $\mathcal{O}(n^2)$  independent entries.

\*julian.heeck@mpi-hd.mpg.de

Dirac mass matrix for the neutrinos  $m_D = vY_\nu$  is generated, leading to a  $(3 + n) \times (3 + n)$  Majorana mass matrix for the neutral fermions  $(\nu_L, N^c)^T$ :<sup>2</sup>

$$\mathcal{M}_{\text{full}} = \begin{pmatrix} 0 & m_D^T \\ m_D & \mathcal{M}_R \end{pmatrix}. \quad (2)$$

Assuming  $\mathcal{M}_R \gg m_D$  gives the low-energy neutrino mass matrix for the flavor neutrinos  $\nu_f \approx \nu_L - m_D^\dagger \mathcal{M}_R^{-1} N^c$  of seesaw form,

$$U_{\text{PMNS}} = P' \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{23}s_{13}c_{12}e^{i\delta} & c_{23}c_{12} - s_{23}s_{13}s_{12}e^{i\delta} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}s_{13}c_{12}e^{i\delta} & -s_{23}c_{12} - c_{23}s_{13}s_{12}e^{i\delta} & c_{23}c_{13} \end{pmatrix} P, \quad (5)$$

and the Majorana phase matrix  $P = \text{diag}(1, e^{i\alpha/2}, e^{i\beta/2})$ . Here we used the abbreviations  $s_{ij} \equiv \sin\theta_{ij}$  and  $c_{ij} \equiv \cos\theta_{ij}$  for the three mixing angles. The phase matrix  $P' = \text{diag}(e^{ia/2}, e^{ib/2}, e^{ic/2})$  can be absorbed by the lepton fields and is hence unphysical. The neutrino mass eigenstates are then  $\nu_m = U_{\text{PMNS}}^\dagger \nu_f$ .

At low energies the nine complex entries of the symmetric Majorana matrix  $\mathcal{M}_\nu$  decompose into the three eigenvalues  $m_i$  (neutrinos masses), three mixing angles  $\theta_{23}, \theta_{12}, \theta_{13}$  (atmospheric, solar, and reactor angle) and three  $CP$  violating phases  $\delta, \alpha, \beta$ , the latter two being unobservable in neutrino oscillations (but in principle testable in  $0\nu\beta\beta$  experiments [12]).

### III. TOP-DOWN PARAMETRIZATION

All the information about neutrino mixing is encoded in  $Y_\nu$ , the only nondiagonal matrix in the lepton sector. For the general complex  $n \times 3$  matrix  $Y_\nu$ , we can write down a singular value decomposition

$$\Omega_{ij}(\alpha_{ij}, \xi_{ij}) \equiv \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & \cos(\alpha_{ij}) & & \sin(\alpha_{ij})e^{i\xi_{ij}} & \\ & & -\sin(\alpha_{ij})e^{-i\xi_{ij}} & & \cos(\alpha_{ij}) & \\ & & & \ddots & & \\ & & & & \ddots & \\ & & & & & 1 \end{pmatrix}, \quad (8)$$

where  $\sin(\alpha_{ij})e^{i\xi_{ij}}$  sits at the  $i$ th row and  $j$ th column. See Appendix A for a derivation and a proof for the validity of this form of  $V_R$ . It is easy to check that  $V_R(V_L)$  has  $n^2(3^2)$  real parameters,  $n(n+1)/2$  (6) of which are phases. From the form of  $Y_\nu$  in Eq. (6), it is clear that the angles  $\phi_j$  either act on zeros in  $D_Y$  or can be absorbed by the phase matrix  $P_L^\dagger$  in  $V_L^\dagger$ . An

<sup>2</sup>Including a type-II seesaw contribution would fill the upper-left zero matrix. For explicit parametrizations of the unitary matrix that diagonalizes  $\mathcal{M}_{\text{full}}$  (for  $n = 3$ ), see e.g., Refs. [10,11].

$$\mathcal{M}_\nu \approx -m_D^T \mathcal{M}_R^{-1} m_D = -v^2 Y_\nu^T \mathcal{M}_R^{-1} Y_\nu. \quad (3)$$

Diagonalization of  $\mathcal{M}_\nu$  can be performed in the following way:

$$\begin{aligned} \mathcal{M}_\nu &= U_{\text{PMNS}}^* \text{diag}(m_1, m_2, m_3) U_{\text{PMNS}}^\dagger \\ &\equiv U_{\text{PMNS}}^* d_m U_{\text{PMNS}}^\dagger, \end{aligned} \quad (4)$$

with the unitary Pontecorvo-Maki-Nakagawa-Sakata matrix (PMNS matrix) in standard parametrization

$$Y_\nu = V_R \begin{pmatrix} \text{diag}(y_1, y_2, y_3) \\ \vec{0}_3 \\ \vdots \\ \vec{0}_3 \end{pmatrix} V_L^\dagger \equiv V_R D_Y V_L^\dagger, \quad (6)$$

with positive singular values  $y_i$ . Out of the  $6n$  real parameters in  $Y_\nu$ , three phases can be absorbed in the lepton fields, so only  $6n - 3$  are physical. We will confirm this with an explicit parametrization of the unitary matrices  $V_{R,L}$  below. For the  $3 \times 3$  matrix  $V_L$ , we take the PMNS parametrization from Eq. (5), while the  $n \times n$  matrix  $V_R$  can be written as

$$V_R = \left( \prod_{i=1}^n \prod_{j=i+1}^n \Omega_{ij}(\alpha_{ij}, \xi_{ij}) \right) \times \text{diag}(e^{i\phi_1}, \dots, e^{i\phi_n}). \quad (7)$$

Here,  $\Omega_{ij}(\alpha_{ij}, \xi_{ij})$  denotes matrices of the form

overall phase and  $P_L^\dagger$  can be absorbed by the lepton fields, so  $V_L$  contains only three angles and three phases. The ordering of the  $\Omega_{ij}$  in  $V_R$  is, of course, arbitrary, which allows us to move all the rotations that act on the zeros in  $D_Y$  to the right, so  $(n-3)(n-4)/2$  matrices of the type (8) drop out, leaving  $n(n-1) - (n-3)(n-4) = 6n - 12$  real parameters in  $V_R$ , half of which are phases. Consequently, we can restrict the product in Eq. (7) to  $\prod_{i=1}^3 \prod_{j=i+1}^n$ .<sup>3</sup> As expected,  $Y_\nu$  contains three (from  $y_i$ ) plus six ( $V_L$ ) plus  $6n - 12$  ( $V_R$ ) equals  $6n - 3$  real, physical parameters,  $3(n-1)$  of which are phases. This agrees with the parameter counting in Ref. [13].

The full mass matrix can then be written as

$$\mathcal{M}_{\text{full}} = \begin{pmatrix} 0 & \nu V_L^* D_Y^T V_R^T \\ \nu V_R D_Y V_L^\dagger & \mathcal{M}_R \end{pmatrix}, \quad (9)$$

with the explicit form for  $V_L$ :

$$V_L = \begin{pmatrix} c_{12}^L c_{13}^L & s_{12}^L c_{13}^L & s_{13}^L e^{-i\beta_1^L} \\ -c_{23}^L s_{12}^L - s_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & c_{23}^L c_{12}^L - s_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & s_{23}^L c_{13}^L \\ s_{23}^L s_{12}^L - c_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & -s_{23}^L c_{12}^L - c_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & c_{23}^L c_{13}^L \end{pmatrix} \text{diag}(1, e^{i\beta_2^L/2}, e^{i\beta_3^L/2}), \quad (10)$$

and the shorthand notation  $s_{ij}^L = \sin\theta_{ij}^L$ , etc. for the mixing angles. In the seesaw limit, we thus find

$$\mathcal{M}_\nu \simeq -\nu^2 V_L^* D_Y^T (V_R^\dagger \mathcal{M}_R V_R^*)^{-1} D_Y V_L^\dagger. \quad (11)$$

Note that the occurring mass matrix  $V_R^\dagger \mathcal{M}_R V_R^*$  is for  $n > 3$  no longer the most general complex symmetric  $n \times n$  matrix, because we already restricted  $V_R$  to the physical angles.

From Eq. (9) or Eq. (11) one can now extract the physical masses and mixing angles in terms of the parameters  $y_i$ ,  $\alpha_{ij}$ ,  $\xi_{ij}$ ,  $\theta_{ij}^L$  and  $\beta_i^L$ .

For completeness, let us count the total number of parameters in the lepton sector. We have three charged lepton masses,  $n$  heavy neutrino masses and  $6n - 3$  real parameters in  $Y_\nu$ , summing up to  $7n$  real parameters.

#### IV. BOTTOM-UP PARAMETRIZATION

Seeing as the seesaw formula (11) contains more parameters than measurable observables, it proves convenient to parametrize our ignorance of the high-energy sector of the seesaw model. In other words, there are infinitely many different  $Y_\nu$  and  $\mathcal{M}_R$  that lead to the same low-energy neutrino parameters via the seesaw formula  $\mathcal{M}_\nu \simeq -m_D^T \mathcal{M}_R^{-1} m_D$ . The parametrization of this ambiguity is the goal of the Casas-Ibarra-parametrization [14] of the Yukawa coupling,

$$Y_\nu = \frac{i}{\nu} \sqrt{\mathcal{M}_R} R \sqrt{d_m} U_{\text{PMNS}}^\dagger. \quad (12)$$

This  $Y_\nu$  solves

$$-\nu^2 Y_\nu^T \mathcal{M}_R^{-1} Y_\nu \simeq \mathcal{M}_\nu = U_{\text{PMNS}}^* d_m U_{\text{PMNS}}^\dagger, \quad (13)$$

as long as the complex  $n \times 3$  matrix  $R$  satisfies  $R^T R = \mathbb{1}_{3 \times 3}$ . Correspondingly, any values of  $\mathcal{M}_R$  and  $R$  in Eq. (12) lead to the same low-energy model, which means we can fix the PMNS mixing angles and masses to the observed ones and study  $Y_\nu$  independently in other processes.

We will now give an explicit parametrization of  $R$ , which has  $3n$  complex entries obeying 12 real constraints ( $R^T R = \mathbb{1}$  is symmetric); we, therefore, expect to find  $6(n-2)$  real physical parameters. This agrees, of course, with the parameter counting in the last section, as we also have three charged lepton masses, three light neutrino masses,  $n$  heavy neutrino masses, and six parameters in the PMNS matrix, summing up to  $7n$  input parameters. Since we only need three orthonormal complex vectors  $\in \mathbb{C}^n$  to define  $R$ , we can write

$$R = O_{n \times n} \begin{pmatrix} \mathbb{1}_{3 \times 3} \\ \vec{0}_3 \\ \vdots \\ \vec{0}_3 \end{pmatrix} \equiv O_{n \times n} \mathbb{1}_{n \times 3}, \quad (14)$$

with a complex orthogonal  $n \times n$  matrix  $O_{n \times n} \in O(n, \mathbb{C})$ . Similar to the  $U(n)$  parametrization used for  $V_R$  in the previous section, we can write  $O_{n \times n}$  as a product of the  $\Omega_{ij}(\gamma_{ij}, \xi_{ij}^L)$  matrices from Eq. (8), but with the arguments  $\xi_{ij}^L = 0$  and  $\gamma_{ij} \in \mathbb{C}$  in order for  $O_{n \times n}$  to be complex orthogonal,

$$O_{n \times n} = S \times \prod_{i=1}^n \prod_{j=i+1}^n \Omega_{ij}(\gamma_{ij}, 0). \quad (15)$$

<sup>3</sup>In other words, the  $U(n-3)$  subgroup of  $U(n)$ , acting on the lower right parts, can be modded out.

Here,  $S$  is a diagonal matrix with entries  $\pm 1$ , which decides the determinant of  $O_{n \times n}$  and, therefore, whether it describes a rotation or a reflection. We can once again mod out the  $O(n-3, \mathbb{C})$  subgroup that acts on the zeros in  $\mathbb{1}_{n \times 3}$  by reordering the rotations, so the product in Eq. (15) can be restricted to  $\prod_{i=1}^3 \prod_{j=i+1}^n$ . Generally speaking,  $R$  is an element of the quotient space  $O(n, \mathbb{C})/O(n-3, \mathbb{C})$ . For real angles  $\gamma_{ij}$ ,  $R$  is an element of the compact Stiefel manifold  $V_3(\mathbb{R}^n) \cong O(n, \mathbb{R})/O(n-3, \mathbb{R})$ , which admits a Haar measure and could thus be used for statistical considerations.

Note that the case  $n = 4$  is special in the sense that there are no superfluous rotations that need to be modded out. One can, therefore, take any complete parametrization for  $O_{4 \times 4}$  without having to worry about unphysical parameters.

For complex angles,  $O_{n \times n}$  and therefore  $R$  and  $Y_\nu$  can have arbitrarily large elements, because the entries  $\cos ix = \cosh x$  grow exponentially with  $x$ . As a result, the Yukawa couplings  $Y_\nu$  can be very large even for small  $\mathcal{M}_R$ , which is somewhat counterintuitive to the seesaw formula and relies on cancellations in the matrix product  $Y_\nu^T \mathcal{M}_R^{-1} Y_\nu$ . Even though these cancellations suggest fine-tuning of parameters, this is a viable possibility and deserves discussion, especially because a low  $\mathcal{M}_R$  with large Yukawa couplings makes an LHC discovery (direct or indirect) possible [15]. Popular models to obtain this fine-tuning in a natural way are for example inverse [16] and linear seesaw [17], usually based on the case  $n = 6$  with an imposed structure in  $\mathcal{M}_R$  and  $Y_\nu$  by some symmetry.

A useful constraint on the complex mixing angles  $\gamma_{ij}$  comes from the perturbativity of  $Y_\nu$ , on which many calculations involving  $Y_\nu$  rely. For this, we propose yet another, slightly different parametrization of  $R$ , which we obtain by noting that  $\Omega_{ij}(\gamma_{ij}, 0) = \Omega_{ij}(\text{Re}(\gamma_{ij}), 0) \times \Omega_{ij}(i \text{Im}(\gamma_{ij}), 0)$ , and that the ordering of the  $\Omega_{ij}$  is arbitrary,

$$R = S \times \left( \prod_{i=1}^3 \prod_{j=i+1}^n \Omega_{ij}(\eta_{ij}, 0) \right) \times \left( \prod_{i=1}^3 \prod_{j=i+1}^n \Omega_{ij}(i\rho_{ij}, 0) \right) \mathbb{1}_{n \times 3}. \quad (16)$$

The  $6(n-2)$  parameters that make up  $R$  are thus equally divided among the new real parameters  $\eta_{ij}$  and  $\rho_{ij}$ . The product involving  $\eta_{ij}$  is just an orthogonal matrix, so we can restrict  $\eta_{ij} \in [0, 2\pi]$  or  $[-\pi, \pi]$ .<sup>4</sup>

The parametrization (16) can be more useful than (14) and (15) because the  $\eta_{ij}$  drop out of the expression  $R^\dagger R$ ,

<sup>4</sup>The parameter space of the angles can be further reduced using discrete field redefinitions like in Refs. [10,18], which however lies outside the scope of this paper.

which arises from (the often occurring)  $Y_\nu^\dagger Y_\nu$  for degenerate  $\mathcal{M}_R$ .

### A. Perturbativity

Perturbativity of the Yukawa couplings gives constraints of the form [19]

$$\text{tr}(Y_\nu^\dagger Y_\nu) = \frac{1}{v^2} \sum_{i=1}^n \sum_{j=1}^3 |R_{ij}|^2 M_i m_j \lesssim \mathcal{O}(1). \quad (17)$$

This inequality is satisfied if

$$|R_{ij}| \lesssim \frac{v}{\sqrt{3nM_i m_j}} \simeq 10^6 \sqrt{\frac{3}{n}} \sqrt{\frac{1 \text{ TeV}}{M_i}} \sqrt{\frac{10^{-2} \text{ eV}}{m_j}}, \quad (18)$$

which constrains each entry in  $R$ . Note that the upper bound goes down for increasing  $n$ . Once the smallest masses are specified, the above equation can be used to give a total upper bound on the parameters  $\rho_{ij}$ . For example, the largest entry in  $R$  can be estimated as  $\cosh \rho_{\max}$  if just one  $\rho_{ij}$  is large and  $(\cosh \rho_{\max})^{3(n-2)}$  if all  $\rho_{ij}$  are large. This typically leads to upper bounds on  $|\rho_{ij}|$  of  $\mathcal{O}(1-10)$ .

### B. Lepton flavor violation

In the seesaw limit, the neutrino states  $\nu_L$  that couple to the charged current are approximately given by [20]  $\nu_L \simeq (1 - \frac{1}{2} m_D^\dagger \mathcal{M}_R^{-2} m_D) U_{\text{PMNS}} \nu_m + m_D^\dagger \mathcal{M}_R^{-1} N^c$ , with the mass eigenstates  $\nu_m$  and  $N^c$ . This leads to lepton flavor violating decays  $\ell \rightarrow \ell' \gamma$  [21], of which  $\mu \rightarrow e \gamma$  gives the strongest constraint,

$$\text{BR}(\mu \rightarrow e \gamma) \simeq \frac{3\alpha}{8\pi} \left| (m_D^\dagger \mathcal{M}_R^{-1} A \mathcal{M}_R^{-1} m_D)_{\mu e} - \frac{1}{2} (m_D^\dagger \mathcal{M}_R^{-2} m_D U_{\text{PMNS}} B U_{\text{PMNS}}^\dagger + \text{H.c.})_{\mu e} \right|^2, \quad (19)$$

with the diagonal matrices  $A_{ij} \equiv \delta_{ij} g(M_j^2/M_W^2)$  and  $B_{ij} \equiv \delta_{ij} g(m_j^2/M_W^2)$  and the monotonic decreasing loop function

$$g(x) = \frac{10 - 43x + 78x^2 - 49x^3 + 4x^4 + 18x^3 \log x}{6(x-1)^4}, \quad (20)$$

$$g(0) = \frac{5}{3}, \quad g(1) = \frac{17}{12}, \quad g(x \rightarrow \infty) = \frac{2}{3}.$$

Here we already omitted the contribution from the light neutrinos, as the corresponding term  $(U_{\text{PMNS}} B U_{\text{PMNS}}^\dagger)_{\mu e}$  is suppressed by the tiny mass-squared differences of the active neutrinos [21]. The remaining contributions are naively of order  $(m_D/\mathcal{M}_R)^2$  but can be enhanced via  $R$ . In the approximation  $M_j \gg M_W \gg m_k$ , we have  $A \propto \mathbb{1} \propto B$ , so Eq. (19) simplifies to

$$\begin{aligned} \text{BR}(\mu \rightarrow e\gamma) & \\ & \simeq \frac{3\alpha}{8\pi} |(m_D^\dagger \mathcal{M}_R^{-2} m_D)_{\mu e}|^2 \\ & \simeq \frac{3\alpha}{8\pi} |(U_{\text{PMNS}} \sqrt{d_m} R^\dagger \mathcal{M}_R^{-1} R \sqrt{d_m} U_{\text{PMNS}}^\dagger)_{\mu e}|^2. \end{aligned} \quad (21)$$

Since this branching ratio can easily exceed the current limit  $\text{BR}(\mu \rightarrow e\gamma) < 2.4 \times 10^{-12}$  [22]—for perturbative couplings—it gives constraints on the  $\rho_{ij}$ . Naively, the relevant product in Eq. (21) is  $|R|^2 m/M$ , so the LFV constraints have a different structure than the perturbativity bound (18). Note that this constraint is connected to the unitarity of the lepton mixing matrix [23], because the relevant matrix  $(\mathbb{1} - \frac{1}{2} m_D^\dagger \mathcal{M}_R^{-2} m_D) U_{\text{PMNS}}$  is no longer unitary at this order. Consequently, the above bound also ensures the validity of our parametrization for  $Y_\nu$  from Eq. (12), as the matrix  $U_{\text{PMNS}}$  in  $Y_\nu$  only corresponds to the lepton mixing matrix if  $(m_D^\dagger \mathcal{M}_R^{-2} m_D)_{ij} \ll 1$ .

While not particularly nice to look at analytically, the constraints given here should be useful in numerical scans involving  $Y_\nu$  for arbitrary many right-handed neutrinos, extending, for example, the analysis of Ref. [19].

## V. APPLICATION TO NEUTRINO ANARCHY

The idea that the large lepton mixing angles and small neutrino hierarchy are not due to some flavor symmetry but rather the absence of any distinction between the neutrino generations has been around for over a decade [24,25]. The basis independence of the neutrino mass and mixing matrices leads to a distribution of the mixing angles according to the Haar measure of the Lie group  $G$  that diagonalizes  $\mathcal{M}_\nu$  [so  $G = U(3)$  for complex,  $G = O(3)$  for real  $\mathcal{M}_\nu$ ]. Since these distributions prefer large mixing angles, this ansatz seems to fit well to the observed large leptonic mixing angles (see also, however, a critique of the anarchy approach in Refs. [26,27]). While the distribution of the mixing angles is uniquely given by the Haar measure, the distributions of the masses and Yukawa couplings

are not unique, and usually the simplest (linear) measure is used (listed in Appendix B).

Let us now discuss neutrino anarchy, i.e., basis independence, with  $n$  right-handed neutrinos. In the seesaw limit, we have the low-energy mass matrix for the active neutrinos from Eq. (11),

$$\mathcal{M}_\nu \simeq -v^2 V_L^* D_Y^T V_R^T \mathcal{M}_R^{-1} V_R D_Y V_L^\dagger, \quad (22)$$

where  $V_L$  and  $V_R$  are now assumed to be distributed according to the Haar measure of  $U(3)$  and  $U(n)$ , respectively, while we take the simplest linear measure for the eigenvalues of  $\mathcal{M}_R$  [25]

$$d\mathcal{M}_R \propto \prod_{i < j}^n |M_j^2 - M_i^2| \prod_{k=1}^n M_k dM_k. \quad (23)$$

We employ the invariant boundary  $\text{tr}(\mathcal{M}_R^\dagger \mathcal{M}_R) = \sum M_i^2 \leq M_0^2$  for the scanning region; see Fig. 1 for a visualization of the distribution. For the singular values of  $D_Y$ , we use the linear measure (see Appendix B for the derivation)

$$dD_Y \propto \prod_{m=1}^3 (y_m^2)^{n-3} \prod_{i < j}^3 (y_i^2 - y_j^2)^2 \prod_{k=1}^3 y_k dy_k, \quad (24)$$

with a similar boundary  $\text{tr}(Y_\nu^\dagger Y_\nu) = \text{tr}(D_Y^\dagger D_Y) = \sum_j^3 y_j^2 \leq y_0^2$ . The effect of large  $n$  will be a reduced hierarchy in the singular values  $y_i$  (see Fig. 2). Note that Eq. (24) is only valid for  $n \geq 3$ ; see Appendix C for the used measure for  $n = 2$ .

We will now briefly describe how the random matrices in this paper are generated. The random unitary matrices  $V_L$  and  $V_R$  in Eq. (22)—following the Haar distribution—can efficiently be obtained from a QR decomposition, as described in Refs. [28]. Drawing eigenvalues from Eq. (23) is a bit more complicated. The procedure usually employed in the literature is the creation of a symmetric matrix  $\mathcal{M}_R$ , where  $\text{Re}(\mathcal{M}_R)_{ij}$  and  $\text{Im}(\mathcal{M}_R)_{ij}$  are uniformly distributed in  $[-M_0, M_0]$ . To stay rotationally invariant, only matrices with  $\text{tr}(\mathcal{M}_R^\dagger \mathcal{M}_R) \leq M_0^2$  are then used and diagonalized




FIG. 1 (color online). Distribution of the  $\mathcal{M}_R$  eigenvalues  $M_k$ , according to the linear measure from Eq. (23) with the boundary constraint  $\sum_{k=1}^n M_k^2 \leq M_0^2$ .






FIG. 2 (color online). Distribution of the singular values  $y_k$  of the Yukawa matrix, according to the linear measure from Eq. (24) for  $n = 3$  (black) and  $n = 20$  (red/dashed) with the boundary constraint  $\sum_{k=1}^3 y_k^2 \leq y_0^2$ .

(roughly speaking, we change the scanning region from a hypercube to a hypersphere). This method works well for  $n \leq 3$ , but becomes virtually unusable for  $n \geq 4$ , because the volume of the hypercube increases sharply with  $n$  compared to the hypersphere, making it hard to accumulate statistics. For this reason, we draw the eigenvalues directly out of the probability distribution (23) using a multivariate Metropolis algorithm [29]. This Markov chain Monte Carlo method is sufficiently fast even for large  $n$  and conveniently works without knowing the normalization factor of the distribution. We use a multivariate Gaussian proposal distribution to generate the next steps in the chain and pick only every hundredth (tenth for  $n = 40$ ) valid point to reduce correlations in the samples.

The value  $v^2 y_0^2 / M_0$  will fix the overall light neutrino mass scale and can be used to fix one of the mass-squared differences  $\Delta m_{ij}^2$ . The PMNS mixing angles and phases do not change with  $n$  and are given by the Haar measure for  $U(3)$  [25] (ignoring unphysical phases),

$$dU_{\text{PMNS}} \propto ds_{12}^2 ds_{23}^2 dc_{13}^4 d\delta d\alpha d\beta, \quad (25)$$

so the only effect of  $n \neq 3$  is a change of the three eigenvalues of  $\mathcal{M}_\nu$ , i.e., the ratio  $R_\nu \equiv (m_2^2 - m_1^2) / (m_3^2 - m_1^2)$ , where we sorted the masses like  $m_1 \leq m_2 \leq m_3$ . In this notation,  $R_\nu < 1/2$  corresponds to normal hierarchy (NH) and  $R_\nu > 1/2$  to inverted hierarchy (IH), with best-fit values  $R_\nu^{\text{NH}} = 0.03$  and  $R_\nu^{\text{IH}} = 0.97$ , respectively, taken from a recent global fit [30].

We observe from Fig. 3 that for increasing  $n$ , the distribution for  $R_\nu$  shifts to larger values, while the width decreases. This reduction of hierarchy can be tracked back mainly to the Yukawa couplings, as the factor  $\prod_{m=1}^3 (y_m^2)^{n-3}$  in  $dD_Y$  pulls the  $y_i$  tightly together. An analysis without this factor shows that the behavior of  $R_\nu$  goes in the opposite direction, i.e., the maximum shifts toward small values for large  $n$  while being diluted. However, the distribution then quickly converges as the only  $n$ -dependent change comes from  $V_R^T \mathcal{M}_R^{-1} V_R$ , for which only the first dozen or so  $M_i$  are relevant, due to the hierarchy in  $\mathcal{M}_R$ . Correspondingly, to shift the maximum of  $R_\nu$  to its observed value, one has not only to omit the factor  $\prod_{m=1}^3 (y_m^2)^{n-3}$  in  $dD_Y$  but also to increase the hierarchy by inserting something like  $\prod_{i < j} (y_i^2 - y_j^2)^2$ , i.e., using a nonlinear measure (see Ref. [6] for other  $y_i$  distributions that accomplish this task). We conclude that the large  $n$  limit in the anarchy approach worsens the agreement with data. While the one data point nature provides can obviously not be used to find the overlying distribution, it should be fair to say that anarchy works best for small  $n$ .

Let us finally comment on the limit  $n \rightarrow \infty$ . As the Yukawa couplings  $y_i$  become quasidegenerate for very large  $n$ , the neutrino masses are simply given by the upper left  $3 \times 3$  submatrix of  $V_R^T \mathcal{M}_R^{-1} V_R$  (times a prefactor). Since there is no preferred direction in the  $V_R$  rotations, the entries  $(V_R)_{ij}$  have the same mean magnitude for a given  $n$ . The mean entries  $\langle (V_R^T \mathcal{M}_R^{-1} V_R)_{jk} \rangle$  then take the




FIG. 3 (color online).  $\log_{10}[R_\nu] \equiv \log_{10}[(m_2^2 - m_1^2) / (m_3^2 - m_1^2)]$  for various  $n$ . The purple/dashed vertical lines indicate the  $3\sigma$  range for  $R_\nu \approx 0.03$  [30] (for NH). The black vertical line divides the NH and IH solutions.







As a cross-check, one can plug the Hermitian matrix  $m_{\text{complex}}^\dagger m_{\text{complex}}$  into Eq. (B3) to rederive Eq. (B2). We can now give a measure for a nonquadratic  $n \times 3$  complex Dirac matrix  $M_{\text{SVD}} = U_R D_Y U_L^\dagger$  with  $U_R \in U(n)$ ,  $U_L \in U(3)$ . Considering  $M_{\text{SVD}}^\dagger M_{\text{SVD}}$  shows that  $dM_{\text{SVD}}$  should at least contain a factor

$$\prod_{i<j}^3 (y_i^2 - y_j^2)^2 \prod_{k=1}^3 y_k dy_k, \quad (\text{B4})$$

which however has too small of a mass dimension. The same procedure for the  $n \times n$  Hermitian  $M_{\text{SVD}} M_{\text{SVD}}^\dagger$  runs into problems, because  $n - 3$  of the eigenvalues are zero and would give a vanishing measure. Taking the product only over the nonvanishing factors, however, yields

$$dM_{\text{SVD}} \propto \prod_{m=1}^3 (y_m^2)^{n-3} \prod_{i<j}^3 (y_i^2 - y_j^2)^2 \prod_{k=1}^3 y_k dy_k dU_L dU_R, \quad (\text{B5})$$

which has the right mass dimension ( $6n$ ) and can, therefore, be viewed as the linear measure for  $M_{\text{SVD}}$ . The overlap of  $dU_L$  and  $dU_R$ —this time including angles, see Sec. III—should again be modded out in a given parametrization to avoid overcounting of physically equivalent configurations (similar to gauge-fixing, as pointed out in Ref. [27]).

The measure (B5) can, of course, be easily extended to a general measure over  $O(N) \times O(M)$  or  $U(N) \times U(M)$  with  $M \leq N$ ,

$$dM_{\text{real}} \propto \prod_{i<j}^M |y_i^2 - y_j^2| \prod_{k=1}^M y_k^{N-M} dy_k dO_L dO_R, \quad (\text{B6})$$

$$dM_{\text{complex}} \propto \prod_{i<j}^M (y_i^2 - y_j^2)^2 \prod_{k=1}^M y_k^{1+2(N-M)} dy_k dU_L dU_R,$$

with the  $N \times M$  matrices  $M_{\text{real}} = O_R D_Y O_L^T$  and  $M_{\text{complex}} = U_R D_Y U_L^\dagger$ . Unphysical rotations need to be modded out again. A simple check shows that the mass dimensions are right and  $N = M$  indeed leads back to Eq. (B2). Equation (B6) can actually be obtained from exactly these requirements, without the need to use Hermitian matrices in the derivation. For example,  $dM_{\text{complex}}$  should contain the known  $N = M$  factor  $\prod_{i<j}^M (y_i^2 - y_j^2)^2 \prod_{k=1}^M dy_k$ , which leaves some factor of mass dimension  $2M$  to the power of  $N - M$ . Since a factor  $\prod_{i<j}^M (y_i^k - y_j^k)^m$ —which would increase the hierarchy in the  $y_i$ —has mass dimension  $M(M - 1)km/2$ , it can only be used in special cases, not for general  $M$  (at least for integer  $k$  and  $m$ ). Thus, for general  $M$ , the only factor with mass dimension  $2M$  is  $\prod_i^M y_i^2$ , leading back to Eq. (B6).

### APPENDIX C: THE CASE $n = 2$

In this Appendix we collect the formulas for the case  $n = 2$  (see also Refs. [3,32]), which is a little different than the  $n \geq 3$  cases and would interrupt the flow if included in the main text. With the  $2 \times 2$  diagonal matrix  $\mathcal{M}_R$ , we find the top-down parametrization in the singular value decomposition for the Yukawa coupling,

$$Y_\nu = V_R \begin{pmatrix} y_1 & 0 & 0 \\ 0 & y_2 & 0 \end{pmatrix} V_L^\dagger. \quad (\text{C1})$$

Two phases of  $V_R$  can be absorbed by  $V_L^\dagger$ , three of  $V_L$  in the lepton fields. Since one of the diagonal phase rotations of  $V_L$  acts on the zeros, there are only two phases in  $V_L$  left, so we can write

$$V_R = \begin{pmatrix} \cos(\alpha_{12}) & \sin(\alpha_{12})e^{i\xi_{12}} \\ -\sin(\alpha_{12})e^{-i\xi_{12}} & \cos(\alpha_{12}) \end{pmatrix}, \quad (\text{C2})$$

$$V_L = \begin{pmatrix} c_{12}^L c_{13}^L & s_{12}^L c_{13}^L & s_{13}^L e^{-i\beta_1^L} \\ -c_{23}^L s_{12}^L - s_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & c_{23}^L c_{12}^L - s_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & s_{23}^L c_{13}^L \\ s_{23}^L s_{12}^L - c_{23}^L s_{13}^L c_{12}^L e^{i\beta_1^L} & -s_{23}^L c_{12}^L - c_{23}^L s_{13}^L s_{12}^L e^{i\beta_1^L} & c_{23}^L c_{13}^L \end{pmatrix} \text{diag}(1, e^{i\beta_2^L/2}, 1), \quad (\text{C3})$$

with the shorthand notation  $s_{ij}^L = \sin\theta_{ij}^L$ , etc., for the mixing angles. The linear measure for  $y_{1,2}$  for the anarchy framework can be derived along the same lines as in Appendix B, with the result

$$dD_Y \propto (y_1^2 - y_2^2)^2 y_1^3 y_2^3 dy_1 dy_2. \quad (\text{C4})$$

Finally, the bottom-up parametrization is given by Eq. (12) with two different  $R$  matrices depending on the low-energy neutrino hierarchy ( $\gamma \in \mathbb{C}$ ) [33],

$$R_{\text{NH}} = \begin{pmatrix} 0 & \cos\gamma & -\sin\gamma \\ 0 & \sin\gamma & \cos\gamma \end{pmatrix}, \quad R_{\text{IH}} = \begin{pmatrix} \cos\gamma & -\sin\gamma & 0 \\ \sin\gamma & \cos\gamma & 0 \end{pmatrix}. \quad (\text{C5})$$

- [1] P. Minkowski, *Phys. Lett.* **67B**, 421 (1977); M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p. 315; T. Yanagida, *Proceedings of the Workshop on Unified Theory and the Baryon Number of the Universe* (KEK, Tsukuba, 1979); R.N. Mohapatra and G. Senjanović, *Phys. Rev. Lett.* **44**, 912 (1980).
- [2] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174**, 45 (1986).
- [3] See, for example, S. F. King, *Nucl. Phys.* **B576**, 85 (2000); P.H. Frampton, S.L. Glashow, and T. Yanagida, *Phys. Lett. B* **548**, 119 (2002); S. Antusch, P. Di Bari, D.A. Jones, and S.F. King, *Phys. Rev. D* **86**, 023516 (2012); K. Harigaya, M. Ibe, and T. T. Yanagida, *Phys. Rev. D* **86**, 013002 (2012).
- [4] M. T. Eisele, *Phys. Rev. D* **77**, 043510 (2008).
- [5] J.R. Ellis and O. Lebedev, *Phys. Lett. B* **653**, 411 (2007).
- [6] B. Feldstein and W. Klemm, *Phys. Rev. D* **85**, 053007 (2012).
- [7] W. Buchmüller, K. Hamaguchi, O. Lebedev, S. Ramos-Sánchez and M. Ratz, *Phys. Rev. Lett.* **99**, 021601 (2007).
- [8] K.R. Dienes, E. Dudas, and T. Gherghetta, *Nucl. Phys.* **B557**, 25 (1999); G.R. Dvali and A. Y. Smirnov, *Nucl. Phys.* **B563**, 63 (1999); N. Arkani-Hamed, S. Dimopoulos, G.R. Dvali, and J. March-Russell, *Phys. Rev. D* **65**, 024032 (2001).
- [9] J. Schechter and J. W.F. Valle, *Phys. Rev. D* **22**, 2227 (1980); G.C. Branco, L. Lavoura, and M.N. Rebelo, *Phys. Lett. B* **180**, 264 (1986); Y. Liao, *Nucl. Phys.* **B749**, 153 (2006).
- [10] M. Blennow and E. Fernández-Martínez, *Phys. Lett. B* **704**, 223 (2011).
- [11] Z.Z. Xing, *Phys. Lett. B* **660**, 515 (2008); *Phys. Rev. D* **85**, 013008 (2012).
- [12] For recent reviews, see W. Rodejohann, *Int. J. Mod. Phys. E* **20**, 1833 (2011); J.J. Gómez-Cadenas, J. Martín-Albo, M. Mezzetto, F. Monrabal, and M. Sorel, *Riv. Nuovo Cimento* **35**, 29 (2012).
- [13] A. Broncano, M. B. Gavela, and E. E. Jenkins, *Phys. Lett. B* **552**, 177 (2003); **636**, 330(E) (2006).
- [14] J. A. Casas and A. Ibarra, *Nucl. Phys.* **B618**, 171 (2001); J. A. Casas, A. Ibarra, and F. Jiménez-Alburquerque, *J. High Energy Phys.* **04** (2007) 064.
- [15] A. de Gouvêa and W.C. Huang, *Phys. Rev. D* **85**, 053006 (2012); P.S.B. Dev, R. Franceschini, and R.N. Mohapatra, *Phys. Rev. D* **86**, 093010 (2012).
- [16] R.N. Mohapatra, *Phys. Rev. Lett.* **56**, 561 (1986); R.N. Mohapatra and J.W.F. Valle, *Phys. Rev. D* **34**, 1642 (1986).
- [17] E. K. Akhmedov, M. Lindner, E. Schnapka, and J.W.F. Valle, *Phys. Rev. D* **53**, 2752 (1996); S.M. Barr, *Phys. Rev. Lett.* **92**, 101601 (2004); C.H. Albright and S.M. Barr, *Phys. Rev. D* **69**, 073010 (2004).
- [18] A. de Gouvêa and J. Jenkins, *Phys. Rev. D* **78**, 053003 (2008).
- [19] J. A. Casas, J.M. Moreno, N. Rius, R. Ruiz de Austri, and B. Zaldivar, *J. High Energy Phys.* **03** (2011) 034.
- [20] W. Grimus and L. Lavoura, *J. High Energy Phys.* **11** (2000) 042.
- [21] T.P. Cheng and L.-F. Li, *Phys. Rev. Lett.* **45**, 1908 (1980).
- [22] J. Adam *et al.* (MEG Collaboration), *Phys. Rev. Lett.* **107**, 171801 (2011).
- [23] Thorough discussions can be found in S. Antusch, C. Biggio, E. Fernández-Martínez, M.B. Gavela, and J. López-Pavón, *J. High Energy Phys.* **10** (2006) 084; D.V. Forero, S. Morisi, M. Tórtola, and J.W.F. Valle, *J. High Energy Phys.* **09** (2011) 142.
- [24] L. J. Hall, H. Murayama, and N. Weiner, *Phys. Rev. Lett.* **84**, 2572 (2000); G. Altarelli, F. Feruglio, and I. Masina, *J. High Energy Phys.* **01** (2003) 035; A. de Gouvêa and H. Murayama, *Phys. Lett. B* **573**, 94 (2003); [arXiv:1204.1249](https://arxiv.org/abs/1204.1249).
- [25] N. Haba and H. Murayama, *Phys. Rev. D* **63**, 053010 (2001).
- [26] M. Hirsch, [arXiv:hep-ph/0102102](https://arxiv.org/abs/hep-ph/0102102).
- [27] J.R. Espinosa, [arXiv:hep-ph/0306019](https://arxiv.org/abs/hep-ph/0306019).
- [28] A. Edelman and N.R. Rao, *Acta Numer.* **14**, 233 (2005); F. Mezzadri, *Notices of the AMS* **54**, 592–604 (2007).
- [29] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller, *J. Chem. Phys.* **21**, 1087 (1953).
- [30] D. V. Forero, M. Tórtola, and J. W.F. Valle, *Phys. Rev. D* **86**, 073012 (2012).
- [31] M. Gronau, R. Johnson, and J. Schechter, *Phys. Rev. D* **32**, 3062 (1985).
- [32] Early references include A. Y. Smirnov, *Phys. Rev. D* **48**, 3264 (1993); A. Kleppe, in *Proceedings of the 3rd Tallinn Symposium on the Neutrino Physics, Lohusalu, Estonia, 8–11 October 1995*, edited by P. Helde, J. Loehmus, and I. Ots, Extending the Standard Model with Two Right-Handed Neutrinos (Estonian Academy of Sciences, Estonia, 1995), p. 118; E. Ma, D.P. Roy, and U. Sarkar, *Phys. Lett. B* **444**, 391 (1998).
- [33] A. Ibarra and G.G. Ross, *Phys. Lett. B* **591**, 285 (2004).