

Black hole formation from a null fluid in extended Palatini gravity

 Jesus Martinez-Asencio,¹ Gonzalo J. Olmo,^{1,*} and D. Rubiera-Garcia^{2,†}
¹*Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia—CSIC. Universidad de Valencia, Burjassot, Valencia 46100, Spain*
²*Departamento de Física, Universidad de Oviedo, Avenida Calvo Sotelo 18, Oviedo, Asturias 33007, Spain*
 (Received 15 September 2012; published 5 November 2012)

We study the formation and perturbation of black holes by null fluxes of neutral matter in a quadratic extension of general relativity formulated *à la* Palatini. Working in a spherically symmetric space-time, we obtain an exact analytical solution for the metric that extends the usual Vaidya-type solution to this type of theory. We find that the resulting space-time is formally that of a Reissner-Nordström black hole but with an effective charge term carrying the wrong sign in front of it. This effective charge is directly related to the luminosity function of the radiation stream. When the ingoing flux vanishes, the charge term disappears and the space-time relaxes to that of a Schwarzschild black hole. We provide two examples that illustrate the formation of a black hole from Minkowski space and the perturbation by a finite pulse of radiation of an existing Schwarzschild black hole.

 DOI: [10.1103/PhysRevD.86.104010](https://doi.org/10.1103/PhysRevD.86.104010)

PACS numbers: 04.40.Nr, 04.50.Kd, 04.70.Bw

I. INTRODUCTION

The properties and structure of black holes in extended theories of gravity have received increasing attention in the last years. In the context of $f(R)$ theories, in both their metric [1,2] and Palatini formulations [3] (see also Ref. [4]), important progress has been made in the understanding of their thermodynamical properties [5] on the existence of spherically symmetric [6,7] and rotating solutions [8], and also on the propagation of perturbations on such backgrounds [9].

A natural motivation to explore the properties of black holes in extensions of general relativity (GR) is to test the effects that new physics could have on these extreme scenarios. However, to date the success within this approach has been quite limited because very often only constant curvature solutions are analytically accessible and yield solutions which are essentially equivalent to the ones found in GR with a cosmological constant (see, however, Ref. [10]). To see this, consider the field equations of $f(R)$ theories in the metric formulation,

$$f_R R_{\mu\nu} - \frac{f}{2} g_{\mu\nu} - \nabla_\mu \nabla_\nu f_R + g_{\mu\nu} \square f_R = \kappa^2 T_{\mu\nu}, \quad (1)$$

where we denote $\kappa^2 \equiv 8\pi G$ and $f_R \equiv \partial f / \partial R$. From this equation, one finds that the Ricci scalar satisfies

$$3\square f_R + R f_R - 2f = \kappa^2 T. \quad (2)$$

This equation shows that in vacuum or with traceless sources such as electromagnetic fields, $R = R_0 = 2f(R_0)/f_R(R_0) = \text{constant}$ solutions are possible. Such solutions turn (1) into

$$R_{\mu\nu} = \frac{\kappa^2}{f_R(R_0)} T_{\mu\nu} + \frac{R_0}{4} g_{\mu\nu}, \quad (3)$$

where the last term plays the role of an effective cosmological constant, $4\Lambda_{\text{eff}} = R_0$, and $\kappa^2/f_R(R_0)$ simply rescales the relation between the bare and the observable Newton's constant, $G_{\text{obs}} = G/f_R(R_0)$. In the Palatini formulation, where metric and connection are treated as independent physical entities, the (metric) field equations take the form

$$\begin{aligned} R_{\mu\nu} - \frac{R}{2} g_{\mu\nu} &= \frac{\kappa^2 T_{\mu\nu}}{f_{\mathcal{R}}} - \frac{\mathcal{R} f_{\mathcal{R}} - f}{2f_{\mathcal{R}}} g_{\mu\nu} \\ &\quad - \frac{3}{2f_{\mathcal{R}}^2} \left[\partial_\mu f_{\mathcal{R}} \partial_\nu f_{\mathcal{R}} - \frac{1}{2} g_{\mu\nu} (\partial f_{\mathcal{R}})^2 \right] \\ &\quad + \frac{1}{f_{\mathcal{R}}} (\nabla_\mu \nabla_\nu f_{\mathcal{R}} - g_{\mu\nu} \square f_{\mathcal{R}}), \end{aligned} \quad (4)$$

where the Palatini curvature \mathcal{R} is a function of the trace T of the matter that satisfies the algebraic equation

$$\mathcal{R} f_{\mathcal{R}} - 2f = \kappa^2 T. \quad (5)$$

In vacuum or with traceless sources, this last equation yields a constant curvature $\mathcal{R} = R_0$, and (4) exactly boils down to (3).

This degeneracy of the two versions of $f(R)$ theories with GR plus cosmological constant precludes any attempt to extract new physics by considering $f(R)$ extensions of GR in scenarios with constant curvature regardless of the symmetries of the problem. This, in particular, implies that also in the dynamical scenario of a null fluid collapsing to form a black hole, $f(R)$ theories yield the same result as in GR, which has been verified recently in the metric approach [11].

Deviations from the predictions of GR in black hole space-times have been observed in the Palatini formulation

*gonzalo.olmo@csic.es
 †rubieradiego@gmail.com

of $f(\mathcal{R})$ theories coupled to Born-Infeld nonlinear electrodynamics [7] and also in Ricci-squared extensions of Palatini theories coupled to Maxwell's electromagnetism [12]. In the former case, it was found that for particular choices of the Lagrangian, $\tilde{f}(\mathcal{R}) = R + al_p^2 R^2$ with $l_p \equiv \sqrt{\hbar G/c^3}$ representing the Planck length, the intensity of the central singularity can be reduced as compared to GR, while in the latter nonsingular charged black hole solutions were obtained in exact analytical form. The emergence of new physics in the Palatini $f(\mathcal{R})$ case can be traced back to the fact that for a nonlinear theory of electrodynamics, the trace of the stress-energy tensor in four dimensions is not zero, which implies a nontrivial relation between \mathcal{R} and T via (5). In the extended Ricci-squared theory, the family of Lagrangians chosen of the form $f(\mathcal{R}, Q) = \tilde{f}(\mathcal{R}) + l_p^2 \mathcal{R}_{\mu\nu} \mathcal{R}^{\mu\nu}$, implies that $\mathcal{R} = \text{constant}$ for the electromagnetic field but $Q = \mathcal{R}_{\mu\nu} \mathcal{R}^{\mu\nu}$ turns out to be nonconstant. This contribution is crucial to change the inner structure of the black hole, which develops a central core whose area grows linearly with the charge in multiples of an elementary unit of order the Planck area. For a certain charge-to-mass ratio, which sets the core mass density precisely at the Planck scale, the space-time is completely regular. For any other choice of the charge-to-mass ratio, the core surface is singular (null-like singularity). In both the singular and nonsingular cases, the metric quickly tends to that predicted by GR as one moves away from the core surface. This puts forward the fact that the emergence of the core and the removal of the singularity are due to nonperturbative mechanisms. A similar qualitative behavior also appears for these models in the cosmological setting [13], where the big bang singularity is replaced by a cosmic bounce at the Planck scale.

The nonperturbative character of the dynamics of the Palatini $f(\mathcal{R}, Q)$ theories near the core makes it difficult to foresee if the nonsingular solutions found in Ref. [12] are robust under small perturbations or not. If a perturbative treatment were possible, under small external perturbations one would expect a small response of the parameters of the system, thus making the nonsingular solutions highly unstable. However, since there is evidence supporting a nonperturbative behavior, nearby configurations could dynamically tend to settle to the static regular solution by radiating away the excess of charge and mass. A proper answer to this question, therefore, can only be obtained by considering the full dynamical problem. In this paper we begin the analysis of this question and consider the simplest nontrivial dynamical scenario, namely, the formation and perturbation of spherically symmetric black holes by null fluxes of neutral matter in the $f(\mathcal{R}, Q)$ Palatini model considered in Ref. [12]. Due to the complicated new physics involved in the charged problem, in this work we focus on the case of Schwarzschild black holes and leave the analysis of the charged case for an independent publication. Here we will see that though the null fluid yields

vanishing \mathcal{R} and Q , modifications with respect to GR are manifest, even though the unperturbed Schwarzschild solution is the same as in GR and can be computed analytically. In fact, the usual Vaidya-type metric found in GR, namely, $ds^2 = -(1 - 2M(v)/r)dv^2 + 2dvdr + r^2 d\Omega^2$ [14], is now modified by a charginelike term with the wrong sign in front of it, i.e., $(1 - 2M(v)/r) \rightarrow (1 - 2M(v)/r - Q^2(v)/r^2)$, where $Q^2(v)$ is related to the *luminosity* of the radiation flux (the notation used here will be explained later). This result puts forward that in Palatini theories the stress-energy density of the matter fields, not just its integral, has a direct influence on the form of the space-time metric. In particular, this allows to see how a flux of radiation modifies the form of the space-time metric along its path and complements with an exact and non-perturbative solution the discussion presented in Ref. [15]. The existence of this charginelike (transient) contribution opens a new window to the recovery of the classical and quantum information stored in the black hole via Hawking radiation. This point will also be discussed in this work.

The content of this paper is organized as follows. In Sec. II we describe the main elements of Palatini formalism, work out the field equations for the metric and the connection, and finally express the relevant field equations using an auxiliary metric associated to the independent connection. In Sec. III we compute the relation between the auxiliary and physical metrics for a null fluid, and write explicitly the field equations for this problem. In Sec. IV we consider the problem of formation and perturbation of black holes in this framework, and illustrate it with two simple examples of luminosity functions. We conclude in Sec. V with a summary and discussion of some future perspectives.

II. PALATINI $f(\mathcal{R}, Q)$ THEORIES

We consider Palatini $f(\mathcal{R}, Q)$ theories defined as follows:

$$S[g, \Gamma, \psi_m] = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} f(\mathcal{R}, Q) + S_m[g, \psi_m], \quad (6)$$

where $\kappa^2 \equiv 8\pi G$, $S_m[g, \psi_m]$ represents the matter action, $g_{\alpha\beta}$ is the space-time metric, $R = g^{\mu\nu} R_{\mu\nu}$, $Q = g^{\mu\alpha} g^{\nu\beta} R_{\mu\nu} R_{\alpha\beta}$, $R_{\mu\nu} = R^\rho{}_{\mu\rho\nu}$, and $R^\alpha{}_{\beta\mu\nu} = \partial_\mu \Gamma^\alpha{}_{\nu\beta} - \partial_\nu \Gamma^\alpha{}_{\mu\beta} + \Gamma^\alpha{}_{\mu\lambda} \Gamma^\lambda{}_{\nu\beta} - \Gamma^\alpha{}_{\nu\lambda} \Gamma^\lambda{}_{\mu\beta}$. The connection $\Gamma^\alpha{}_{\beta\gamma}$ has no *a priori* relation with the metric (Palatini formalism) and must be determined by the theory through the corresponding field equations. A variation of (6) with respect to metric and connection leads to

$$f_{\mathcal{R}} R_{\mu\nu} - \frac{f}{2} g_{\mu\nu} + 2f_Q R_{\mu\alpha} R^\alpha{}_\nu = \kappa^2 T_{\mu\nu}, \quad (7)$$

$$\nabla_\beta [\sqrt{-g} (f_{\mathcal{R}} g^{\mu\nu} + 2f_Q R^{\mu\nu})] = 0. \quad (8)$$

Here, $T_{\mu\nu}$ is the stress-energy tensor of the matter sector S_m , obtained as $T_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\mu\nu}}$. In the above derivation

of the field equations two assumptions have been made, namely, vanishing torsion, $\Gamma_{[\beta\gamma]}^\alpha = 0$, and symmetric Ricci tensor, $R_{[\mu\nu]} = 0$ (see Ref. [16] for the resulting equations and implications when such constraints are relaxed). In order to solve for the connection, we must note that Eq. (8) depends on the metric $g_{\mu\nu}$ and on the independent connection $\Gamma_{\beta\gamma}^\alpha$, though the connection dependence can be eliminated in favor of the matter. This can be seen by defining a matrix \hat{P} whose components are $P_{\alpha}{}^\nu = R_{\alpha\beta}g^{\beta\nu}$, and rewriting (7) as

$$2f_Q\hat{P}^2 + f_R\hat{P} - \frac{f}{2}\hat{I} = \kappa^2\hat{T}, \quad (9)$$

where \hat{I} and \hat{T} are the matrix representation of the identity δ_μ^ν and the stress-energy tensor $T_\mu{}^\nu$, respectively. Equation (9) implies that $\hat{P} = \hat{P}(\hat{T})$ is a function of the matter, and therefore, $R = P_{\alpha}{}^\alpha$ and $Q = P_{\alpha}{}^\nu P_\nu{}^\alpha$ are also functions of the matter. To solve the equation for the independent connection (8), we look for another (auxiliary) metric $h_{\mu\nu}$ such that $\Gamma_{\beta\gamma}^\alpha$ becomes its Levi-Civita connection. To do so, we propose the following ansatz [12] $\sqrt{-g}(f_R g^{\mu\nu} + 2f_Q R^{\mu\nu}) = \sqrt{-h}h^{\mu\nu}$, which turns (8) into $\nabla_\beta[\sqrt{-h}h^{\mu\nu}] = 0$. Simple algebraic manipulations show that the relation between $g_{\mu\nu}$ and $h_{\mu\nu}$ is given by

$$\hat{h}^{-1} = \frac{\hat{g}^{-1}\hat{\Sigma}}{\sqrt{\det\hat{\Sigma}}}, \quad \hat{h} = \left(\sqrt{\det\hat{\Sigma}}\right)\hat{\Sigma}^{-1}\hat{g}, \quad (10)$$

which shows that the connection of $f(R, Q)$ theories with the constraints above can be explicitly solved in terms of the physical metric $g_{\mu\nu}$ and the matter sources. In (10) we have introduced the object

$$\Sigma_{\alpha}{}^\nu = (f_R \delta_{\alpha}{}^\nu + 2f_Q P_{\alpha}{}^\nu), \quad (11)$$

which contains all the information about the relative deformation between $g_{\mu\nu}$ and $h_{\mu\nu}$. In particular, in the case of $f(R)$ theories we have $\Sigma_{\alpha}{}^\nu = f_R \delta_{\alpha}{}^\nu$, and therefore, $g_{\mu\nu}$ and $h_{\mu\nu}$ become conformally related as $g_{\mu\nu} = \frac{1}{f_R} h_{\mu\nu}$. In terms of the metric $h_{\mu\nu}$, the metric field equation (7) can be written in a more compact and transparent form. To see this, note that (9) can be expressed as $\hat{P}\hat{\Sigma} = \frac{f}{2}\hat{I} + \kappa^2\hat{T}$, and taking into account that $\hat{P}\hat{\Sigma} = \sqrt{\det\hat{\Sigma}}R_{\mu\alpha}(h)h^{\alpha\nu}$ we arrive at

$$R_{\mu}{}^\nu(h) = \frac{1}{\sqrt{\det\hat{\Sigma}}}\left(\frac{f}{2}\delta_{\mu}{}^\nu + \kappa^2 T_{\mu}{}^\nu\right). \quad (12)$$

This form of the field equations describes the dynamics of Palatini $f(R, Q)$ theories in such a way that the right-hand side only depends on the matter sources, as both R and Q (and hence f) are functions of the stress-energy tensor. It should be noted that unlike in the usual metric formalism,

$h_{\mu\nu}$ satisfies second-order equations, and therefore this type of Palatini $f(R, Q)$ theory is ghost free.

III. FIELD EQUATIONS FOR A NULL FLUID

In this section we consider a specific form of matter that will allow us to explicitly obtain the matrix $\Sigma_{\alpha}{}^\nu$, a key element of the field equations of our theory. This will allow us to express the field equations in a form suitable for calculations. Then we will specify the $f(R, Q)$ model and write the field equations taking advantage of the simplifications offered by the assumed spherical symmetry.

A. Determination of $\Sigma_{\alpha}{}^\nu$

We shall consider the formation of black holes from an ingoing flux of pressureless null neutral matter whose stress-energy tensor is written as

$$T_{\mu\nu} = \rho_{\text{in}} l_\mu l_\nu, \quad (13)$$

where ρ_{in} is the energy density of the ingoing stream and l_μ represents a null radial vector, $l_\mu l^\mu = 0$. With this matter source, (9) can be rewritten as

$$2f_Q\left(\hat{P} + \frac{f_R}{4f_Q}\hat{I}\right)^2 = \left(\frac{f}{2} + \frac{f_R^2}{8f_Q}\right)\hat{I} + \kappa^2\hat{T}, \quad (14)$$

where $\hat{T} \equiv \rho_{\text{in}} l_\mu l^\nu$. Since to construct $\Sigma_{\alpha}{}^\nu$ we must have an expression for \hat{P} , we need to take the square root of the above equation. To proceed, we propose the following ansatz:

$$B_{\mu}{}^\alpha = \lambda \delta_{\mu}^\alpha + \Omega_{\text{in}} l_\mu l^\alpha, \quad (15)$$

as the square root of the left-hand side of (14), with λ and Ω_{in} two functions to be determined. We then find that

$$B_{\mu}{}^\alpha B_{\alpha}{}^\nu = \lambda^2 \delta_{\mu}^\nu + 2\lambda \Omega_{\text{in}} l_\mu l^\nu, \quad (16)$$

and comparing with the right-hand side of (14), we obtain

$$\lambda^2 = \frac{f}{2} + \frac{f_R^2}{8f_Q}, \quad \Omega_{\text{in}} = \frac{\kappa^2 \rho_{\text{in}}}{2\lambda}. \quad (17)$$

We thus find that

$$\begin{aligned} \hat{\Sigma} &= \frac{f_R}{2}\hat{I} + \sqrt{2f_Q}\hat{B} \\ &= \left[\frac{f_R}{2} + \sqrt{2f_Q}\lambda\right]\delta_{\mu}^\nu + \sqrt{\frac{f_Q}{2}}\frac{\kappa^2 \rho_{\text{in}}}{\lambda}l_\mu l^\nu. \end{aligned} \quad (18)$$

This equation implies that once a Lagrangian $f(R, Q)$ is given, the relation between $g_{\mu\nu}$ and $h_{\mu\nu}$ is immediately obtained, and consequently we will be able to write explicitly the solution in the metric $g_{\mu\nu}$ once its counterpart in the metric $h_{\mu\nu}$ is determined.

B. $f(R, Q)$ model and field equations

Therefore, to proceed further we must specify the $f(R, Q)$ model. In this work we will focus on the Lagrangian [12,13,17]

$$f(R, Q) = R + l_p^2(aR^2 + Q), \quad (19)$$

where a is some dimensionless constant and l_p is the Planck length. Lagrangians of this kind are expected to arise as effective descriptions or in the low-energy limit of quantum theories of gravity in which both metric and connection play relevant roles in the resulting effective geometry (see e.g., Refs. [18–20]). For this model, tracing in (7) with $g^{\mu\nu}$ leads to $R = -k^2T$, which is the same relation as in GR. Since for a null fluid $T = 0$, we have $R = 0$ and thus $f_R = 1$ and $f = l_p^2 Q$. With these data, Q can be calculated from the trace of (15),

$$\frac{1}{2\sqrt{2}l_p} = \sqrt{\frac{l_p^2 Q}{2} + \frac{1}{8l_p^2}}, \quad (20)$$

which is satisfied only if $Q = 0$. Using this result and the fact that $R = 0$ we can finally put $\hat{\Sigma}$ as

$$\Sigma_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} + 2l_p^2 \kappa^2 \rho_{\text{in}} l_{\mu} l^{\nu}. \quad (21)$$

With a little algebra one can verify that $\det \hat{\Sigma} = 1$. Collecting all these results we find that the field equations (12) boil down to

$$R_{\mu}{}^{\nu}(h) = \kappa^2 \rho_{\text{in}} l_{\mu} l^{\nu}, \quad (22)$$

which can also be written using the Einstein tensor of the metric $h_{\mu\nu}$ as

$$G_{\mu}{}^{\nu}(h) = \kappa^2 \rho_{\text{in}} l_{\mu} l^{\nu}. \quad (23)$$

The formal simplicity of this expression puts forward the great advantage of using the metric $h_{\mu\nu}$ instead of the physical metric $g_{\mu\nu}$ to deal with the field equations. To see this, note that $\Gamma_{\beta\gamma}^{\alpha}$ [the connection that defines $R_{\mu\nu}(h)$] can be expressed in terms of $g_{\mu\nu}$ as

$$\Gamma_{\beta\gamma}^{\alpha} = L_{\beta\gamma}^{\alpha} + A_{\beta\gamma}^{\alpha}, \quad (24)$$

where $L_{\beta\gamma}^{\alpha}$ is the Levi-Civita connection of $g_{\mu\nu}$, and $A_{\beta\gamma}^{\alpha}$ is a tensor given by

$$A_{\beta\gamma}^{\alpha} = \frac{h^{\alpha\rho}}{2} [\nabla_{\beta}^L h_{\rho\gamma} + \nabla_{\gamma}^L h_{\rho\beta} - \nabla_{\rho}^L h_{\beta\gamma}], \quad (25)$$

with $\nabla_{\beta}^L W_{\nu} = \partial_{\beta} W_{\nu} - L_{\beta\nu}^{\sigma} W_{\sigma}$. Expressing (23) in terms of $g_{\mu\nu}$ and the matter would lead to an expression of the form $G_{\mu\nu}(g) = \kappa^2 \rho_{\text{in}} l_{\mu} l_{\nu} +$ terms involving up to second-order derivatives of ρ_{in} and l_{ν} . Though that approach is certainly possible, it also appears unnecessarily lengthy and cumbersome.

C. Spherical space-time

We are considering the problem of the dynamical generation/perturbation of a black hole in a model with spherical symmetry from/with a pure radiation field. To this end we need first to specify the explicit form of the field equations for a spherically symmetric space-time. For this purpose it is convenient to use a coordinate system $x^{\alpha} \equiv (x^0, x^1, \theta, \phi)$, with $x = \{x^0, x^1\}$ the coordinates of the two-spaces with $(\theta, \phi) = \text{constant}$, and to write the line element of the physical metric $g_{\mu\nu}$ as

$$ds^2 = g_{ab}(x) dx^a dx^b + r^2(x) d\Omega^2, \quad (26)$$

and the line element of the auxiliary metric $h_{\mu\nu}$ as

$$d\tilde{s}^2 = h_{ab}(x) dx^a dx^b + \tilde{r}^2(x) d\Omega^2. \quad (27)$$

In the above expressions, $d\Omega^2$ is the line element of a two-sphere. While these line elements are formally equal, their relation is nontrivial due to the relative deformation between the metrics defined by the matrix $\hat{\Sigma}$, which in general implies that $r^2(x) \neq \tilde{r}^2(x)$. Our strategy will be to solve the field equations (23) involving the metric $h_{\mu\nu}$, and then use the transformation matrix $\hat{\Sigma}$ to obtain $g_{\mu\nu}$. In this sense, we note that using (21) in (10), the relation between the four-dimensional metrics $h_{\mu\nu}$ and $g_{\mu\nu}$ boils down to

$$\begin{aligned} g_{\mu\nu} &= h_{\mu\nu} + 2l_p^2 \kappa^2 \rho_{\text{in}} l_{\mu} l_{\nu}, \\ g^{\mu\nu} &= h^{\mu\nu} - 2l_p^2 \kappa^2 \rho_{\text{in}} l^{\mu} l^{\nu}. \end{aligned} \quad (28)$$

Since we are considering a radially infalling fluid, $l_{\mu} = (l_0, l_1, 0, 0)$, the above relations guarantee that in this problem $r^2(x) = \tilde{r}^2(x)$ and $g_{ab}(x) = h_{ab}(x) + 2l_p^2 \kappa^2 \rho_{\text{in}} l_a l_b$, where $l_a = \{l_0, l_1\}$, which simplifies the analysis. The tilde on top of the function $r(x)$ will thus be omitted from now on.

Using the line element (27), the components of the Einstein tensor can be written as [21]

$${}^4G_{ab}(h) = -\frac{1}{r^2} [2rr_{;ab} + (1 - 2r\Box r - r^a r_{;a})], \quad (29)$$

$$G_{\theta\theta}(h) = \sin^2\theta G_{\phi\phi}(h) = r\Box r - \frac{1}{2}r^2\bar{R}, \quad (30)$$

where \bar{R} and all covariant derivatives are computed using the two-dimensional metric h_{ab} (with latin indices to distinguish it from the four-dimensional metric $h_{\mu\nu}$). Note that these equations are completely general for any spherically symmetric metric, as we have not specified any coordinates associated to the hypersurfaces $(\theta, \phi) = \text{constant}$.

Combining the right-hand side of (23) with Eqs. (29) and (30), we find

$$2rr_{;ab} + (1 - 2r\Box r - r^a r_{;a})h_{ab} = -\kappa^2 r^2 T_{ab}, \quad (31)$$

$$\square r - \frac{1}{2} r \bar{R} = \kappa^2 r P, \quad (32)$$

where $T_{ab} = \rho_{\text{in}} l_a l_b$ and $P = T_{\theta\theta} = T_{\phi\phi} = 0$ for a pressureless fluid.

It is now convenient to introduce the scalar functions $A = A(x)$ and $m = m(x)$, such that $A(x) \equiv r^a r_{,a} = 1 - 2m/r$, in terms of which (31) becomes

$$r_{;ab} + \left(\frac{m}{r^2} - \square r \right) h_{ab} = -\frac{r\kappa^2}{2} T_{ab}. \quad (33)$$

Taking the trace of this equation, we obtain $\square r = \frac{2m}{r^2} + \frac{r\kappa^2}{2} T$, and (31) is written as

$$r_{;ab} - \frac{m}{r^2} h_{ab} = -\frac{r\kappa^2}{2} (T_{ab} - h_{ab} T). \quad (34)$$

Replacing now the $\square r$ term obtained above back into (32), one obtains

$$\bar{R} = \frac{4m}{r^3} + \kappa^2 (T - 2P). \quad (35)$$

Consider now the quantity $\partial_a A$, which establishes the following relation between $r_{;ab}$ and $m_{,a}$:

$$m_{,a} = \frac{m}{r} r_{,a} - h^{bc} r r_{,b} r_{,ac}. \quad (36)$$

Inserting (34) in this equation, we find

$$m_{,a} = \frac{\kappa^2 r^2}{2} r_{,b} (T_a^b - \delta_a^b T). \quad (37)$$

Equations (34), (35), and (37) constitute the system of equations defining our problem. We already noted that the null fluid satisfies $T = 0$, and imposed $P = 0$ for simplicity. The resulting field equations in such a scenario are

$$r_{;ab} - \frac{m}{r^2} h_{ab} = -4\pi r \rho_{\text{in}} l_a l_b, \quad (38)$$

$$\bar{R} = \frac{4m}{r^3}, \quad (39)$$

$$m_{,a} = 4\pi r^2 T_a^b r_{,b}. \quad (40)$$

IV. FORMATION OF BLACK HOLES

In this section we solve Eqs. (38)–(40) corresponding to an ingoing stream of particles represented by a pressureless null fluid. We will thus focus on a Vaidya-type metric [14] of the form

$$d\bar{s}^2 = -Ae^{2\psi} dv^2 + 2e^\psi dv dr + r^2 d\Omega^2, \quad (41)$$

where $A = 1 - \frac{2m(v)}{r}$ and ψ are functions of $x = \{v, r\}$, and the null vector is normalized as $l_a = -\partial_a v$. This type of metric is often used in the study of black hole formation and evaporation processes [22].

The r - v component of (38) gives $\psi_{,r} = 0$ and implies that $\psi = \psi(v)$. This means that we can define a new variable $V(v)$ such that $dV = e^{\psi(v)} dv$, and eliminate this function from the line element (41). As a result, the only nontrivial function in the problem is $m = m(v, r)$. Using the v - v equation of (38) or the v component of (40), we find

$$m_{,v} = \frac{\kappa^2 r^2}{2} \rho_{\text{in}}. \quad (42)$$

The r component of (40) gives $m_{,r} = 0$ and implies that $m = m(v)$, which can also be verified using the conservation equation of the matter. As a result, the function on the right-hand side of (42) can only depend on the v coordinate. We will thus introduce a function $L(v) = \frac{\kappa^2 r^2}{2} \rho_{\text{in}}$, called luminosity, to characterize the flux of incoming radiation so that $m_{,v} = L(v)$ and $m(v) = \int^v L(v') dv'$.

In order to write the solution in terms of the physical metric $g_{\mu\nu}$, let us define the corresponding Vaidya-type metric as

$$ds^2 = -Be^{2\Psi} dv^2 + 2e^\Psi dv dr + r^2 d\Omega^2, \quad (43)$$

where $B(v, r) = 1 - \frac{2M(v)}{r}$. Comparing with (41), we find that $\Psi = 0$ due to the r - v component of (38). From Eq. (28), it follows that

$$A = B + 2l_p^2 \kappa^2 \rho_{\text{in}}, \quad (44)$$

which implies

$$M(v) = m(v) + r l_p^2 \kappa^2 \rho_{\text{in}}. \quad (45)$$

Replacing this expression in the definition of B and using the definition of the luminosity given above, we obtain

$$B = 1 - \frac{2 \int^v L(v') dv'}{r} - \frac{4L(v)}{\rho_P r^2}, \quad (46)$$

where $\rho_P = \frac{c^2}{l_p^2 G} \sim 10^{96} \text{ kg/m}^3$ is the Planck density. This formula is the main result of this work. Let us analyze now its implications.

The space-time (46) resulting from the perturbation of a Schwarzschild black hole by a null fluid in the Palatini theory $f(R, Q) = R + l_p^2 (aR^2 + Q)$ is formally that of a (nonrotating) charged (Reissner-Nordström) black hole but with a charge term whose sign is the opposite to the usual one. This allows us to define an “effective charge” in (46) as

$$Q^2(v) = \frac{4L(v)}{\rho_P}, \quad (47)$$

and a mass term

$$\tilde{M}(v) = \int_{v_0}^v L(v') dv', \quad (48)$$

such that (46) becomes

$$B = 1 - \frac{2\tilde{M}(v)}{r} - \frac{Q^2(v)}{r^2}. \quad (49)$$

The horizons of this “opposite” sign charged metric, solutions of $r^2 - 2\tilde{M}(v)r - Q^2(v) = 0$, are found at

$$r_{\pm} = \tilde{M}(v) \pm \sqrt{\tilde{M}(v) + Q^2(v)}. \quad (50)$$

Since $r_- < 0$, it follows that there is a single horizon r_+ larger than the usual Schwarzschild radius, i.e., $r_+ > 2\tilde{M}$.

The dynamics of these “charged” black holes is strongly tied to the shape and temporal distribution of the ingoing matter stream. In this sense, it must be noted that a typical Dirac-delta profile for the luminosity function, i.e., $L(v) = m_0\delta(v - v_0)$, is not consistent because that would imply a divergent metric along the path $v = v_0$ due to the charge term (47). This property of the metric forces the consideration of smooth and finite luminosity profiles. Moreover, in order to guarantee a smooth geometry, the luminosity function should be continuous and differentiable at least up to order two. As an example, let us consider the following steplike radiation pulse (see Fig. 1):

$$L(v) = \frac{\tilde{M}_0}{4(v_1 - v_0)}(1 + \tanh(v - v_0))(1 - \tanh(v - v_1)), \quad (51)$$

with $v_0 = 1$ and $v_1 = 100$ in Fig. 1 signaling the (effective) beginning and end of the pulse, respectively. This flux of radiation has the following effect on the geometry. We begin with an empty Minkowski space at $v \rightarrow -\infty$. Near $v \approx v_0$, the amplitude of the flux grows up to its maximum and remains almost constant until $v \approx v_1$. During the time that the pulse is active, the mass (48) grows linearly (see Fig. 2), and the effective charge remains constant. Once the radiation pulse is off (beyond v_1), the charge disappears and the configuration becomes that of a Schwarzschild black hole with constant mass $\tilde{M} \approx \tilde{M}_0$.




FIG. 1 (color online). Smooth ingoing flux (51) mimicking a steplike profile to illustrate the formation of a Schwarzschild black hole out of Minkowski space.




FIG. 2 (color online). Behavior of the black hole mass function $\tilde{M}(v)$ due to the steplike pulse (51).




FIG. 3 (color online). $L(v)$ profile (52) for a compact support perturbation.




FIG. 4 (color online). Growth of the black hole mass function $\tilde{M}(v)$ due to the perturbation (52).

Consider now a Schwarzschild black hole of mass $\tilde{M} = \tilde{M}_0$ and a compact support perturbation of the form (see Fig. 3)

$$L(v) = \frac{\epsilon}{240} (|v - 3|^5 - 6|v - 2|^5 + 15|v - 1|^5 - 20|v|^5 + 15|v + 1|^5 - 6|v + 2|^5 + |v + 3|^5). \quad (52)$$

This function and its derivatives are continuous and differentiable up to third order and are exactly zero beyond $|v| \geq 3$ [unlike the pulse (51), which is defined over the whole real line]. This flux provides a local perturbation of an existing Schwarzschild black hole and complements the formation scenario of the previous example. The integral $\int_{-3}^3 L(v)$ is normalized to unity, and therefore $\tilde{M}(v) = \tilde{M}_0 + \epsilon$ for $v \geq 3$ (see Fig. 4).

V. CONCLUSIONS AND PERSPECTIVES

In this paper we have considered the formation and (exact) perturbation of spherically symmetric black holes in a Palatini $f(R, Q)$ theory by means of neutral fluxes of a pressureless null fluid. We have seen that once the flux of radiation is over, the resulting geometry is exactly the Schwarzschild solution [23] of GR. Though the asymptotic solution is exactly Schwarzschild, the intermediate perturbed states are not. As long as the ingoing flux is nonzero, the metric exhibits a chargelike term whose form is directly related to the luminosity function of the radiation stream. Though this term is Planck suppressed, its mere existence indicates that the energy density of the matter is leaving a direct imprint on the metric, putting forward that any fluctuation of the matter profile will induce an identical fluctuation on the space-time metric modulated by a factor

$1/r^2$, as given in Eq. (46). This shows, with an exact analytical solution, that in these Palatini theories the back-reaction of the matter on the geometry is completely sensitive to the most minute details of the matter profile, $Q^2(v)$, not just to its integrated form, \tilde{M} . This, in particular, motivates the consideration of the process of black hole evaporation via Hawking radiation in these scenarios to see whether the imprint that the infalling matter leaves on the geometry can be recovered at infinity through the interaction of the geometry with the outgoing Hawking quanta. This point naturally opens the door to the study of the interaction between the ingoing ρ_{in} and outgoing ρ_{out} fluxes to see whether a kind of Dray-'t Hooft-Redmount (DTR) [24] relation may be found in our model. A difficulty that may arise in our case in comparison with GR is that the peculiarities of our theory prevent the use of delta functions to describe the matter profile. In addition, the consideration of two (ingoing and outgoing) fluxes implies that the curvature scalar Q becomes a function of the product $\rho_{\text{in}}\rho_{\text{out}}$ [25]. The absence of that kind of terms on the right-hand side of Einstein's equations was crucial in the DTR analysis and also in the study of the mass inflation phenomenon [21] (see also Ref. [26]) in the charged case. These aspects will be studied in detail and reported elsewhere.

ACKNOWLEDGMENTS

The work of G. J. O. has been supported by the Spanish Grants No. FIS2008-06078-C03-02 and No. FIS2011-29813-C02-02, and the JAE-doc program of the Spanish Research Council (CSIC). D. R.-G. thanks the hospitality of the theoretical physics group at Valencia U., where part of this work was carried out.

-
- [1] S. Capozziello and M. De Laurentis, *Phys. Rep.* **509**, 167 (2011).
 - [2] A. De Felice and S. Tsujikawa, *Living Rev. Relativity* **13**, 3 (2010); T. P. Sotiriou and V. Faraoni, *Rev. Mod. Phys.* **82**, 451 (2010); S. Capozziello and M. Francaviglia, *Gen. Relativ. Gravit.* **40**, 357 (2008); S. Nojiri and S. D. Odintsov, *Int. J. Geom. Methods Mod. Phys.* **04**, 115 (2007).
 - [3] G. J. Olmo, *Int. J. Mod. Phys. D* **20**, 413 (2011).
 - [4] T. Harko, T. S. Koivisto, F. S. N. Lobo, and G. J. Olmo, *Phys. Rev. D* **85**, 084016 (2012).
 - [5] F. Briscese and E. Elizalde, *Phys. Rev. D* **77**, 044009 (2008); D. N. Vollick, *Phys. Rev. D* **76**, 124001 (2007); V. Faraoni, *Entropy* **12**, 1246 (2010); K. Bamba and C.-Q. Geng, *J. Cosmol. Astropart. Phys.* **06** (2010) 014.
 - [6] S. Capozziello, N. Frusciante, and D. Vernieri, *Gen. Relativ. Gravit.* **44**, 1881 (2012); L. Hollenstein and F. S. N. Lobo, *Phys. Rev. D* **78**, 124007 (2008).
 - [7] G. J. Olmo and D. Rubiera-Garcia, *Phys. Rev. D* **84**, 124059 (2011); *AIP Conf. Proc.* **1458**, 511 (2011).
 - [8] A. de la Cruz-Dombriz, A. Dobado, and A. L. Maroto, *Phys. Rev. D* **80**, 124011 (2009); **83**, 029903(E) (2011); J. A. R. Cembranos, A. de la Cruz-Dombriz, and P. J. Romero, *arXiv:1109.4519*; T. Moon, Y. S. Myung, and E. J. Son, *Gen. Relativ. Gravit.* **43**, 3079 (2011); S. Capozziello, M. De Laurentis, and A. Stabile, *Classical Quantum Gravity* **27**, 165008 (2010).
 - [9] A. de la Cruz-Dombriz, A. Dobado, and A. L. Maroto, *J. Phys. Conf. Ser.* **229**, 012033 (2010); Y. S. Myung, T. Moon, and E. J. Son, *Phys. Rev. D* **83**, 124009 (2011).
 - [10] T. Clifton and J. D. Barrow, *Phys. Rev. D* **72**, 103005 (2005).
 - [11] S. G. Ghosh and S. D. Maharaj, *Phys. Rev. D* **85**, 124064 (2012).
 - [12] G. J. Olmo and D. Rubiera-Garcia, *Eur. Phys. J. C* **72**, 2098 (2012); *Int. J. Mod. Phys. D* **21**, 1250067 (2012); *Phys. Rev. D* **86**, 044014 (2012).

- [13] C. Barragan and G.J. Olmo, *Phys. Rev. D* **82**, 084015 (2010); C. Barragan, G.J. Olmo, and H. Sanchis-Alepuz, *Phys. Rev. D* **80**, 024016 (2009).
- [14] W.B. Bonnor and P.C. Vaidya, *Gen. Relativ. Gravit.* **1**, 127 (1970).
- [15] G.J. Olmo, *J. Cosmol. Astropart. Phys.* **10** (2011) 018.
- [16] G.J. Olmo, *AIP Conf. Proc.* **1458**, 222 (2012).
- [17] G.J. Olmo, H. Sanchis-Alepuz, and S. Tripathi, *Phys. Rev. D* **80**, 024013 (2009).
- [18] L. Parker and D.J. Toms, *Quantum Field Theory in Curved Space-Time: Quantized Fields and Gravity* (Cambridge University Press, Cambridge, England, to be published); N. D. Birrel and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, England, 1982); A. Dobado, A. Gómez-Nicola, A. L. Maroto, and J. R. Peláez, *Effective Lagrangians for the Standard Model* (Springer, Berlin, 1997).
- [19] T. Ortin, *Gravity and Strings*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 2004); M. Green, J. Schwarz, and E. Witten, *Superstring Theory* (Cambridge University Press, Cambridge, England, 1987).
- [20] G.J. Olmo and P. Singh, *J. Cosmol. Astropart. Phys.* **01** (2009) 030.
- [21] E. Poisson and W. Israel, *Phys. Rev. D* **41**, 1796 (1990).
- [22] A. Fabbri and J. Navarro-Salas, *Modeling Black Hole Evaporation* (ICP-World Scientific, London, 2005).
- [23] From our derivations, it is straightforward to verify that the addition of a cosmological constant or the consideration of an $\tilde{f}(R)$ piece with nonquadratic corrections would lead to Schwarzschild-de Sitter solutions.
- [24] T. Dray and G. 't Hooft, *Commun. Math. Phys.* **99**, 613 (1985).
- [25] J. Martinez Asencio, G.J. Olmo, and D. Rubiera-Garcia (unpublished).
- [26] A. Ori, *Phys. Rev. Lett.* **68**, 2117 (1992); P. R. Brady and J. D. Smith, *Phys. Rev. Lett.* **75**, 1256 (1995); M. Banados, J. Silk, and S.M. West, *Phys. Rev. Lett.* **103**, 111102 (2009); P.P. Avelino, A.J.S. Hamilton, and C.A.R. Herdeiro, *Phys. Rev. D* **79**, 124045 (2009); S.E. Hong, D.-il. Hwang, E.D. Stewart, and D -han. Yeom, *Classical Quantum Gravity* **27**, 045014 (2010); E.G. Brown, R.B. Mann, and L. Modesto, *Phys. Rev. D* **84**, 104041 (2011).