

Adiabaticity and color mixing in tetraquark spectroscopy

J. Vijande*

*Departamento de Física Atómica, Molecular y Nuclear, Universidad de Valencia (UV) and IFIC (UV-CSIC), Valencia 46100, Spain*A. Valcarce[†]*Departamento de Física Fundamental, Universidad de Salamanca, 37008 Salamanca, Spain*J.-M. Richard[‡]*Institut de Physique Nucléaire de Lyon, IN2P3-CNRS-UCBL, Université de Lyon, 4 rue Enrico Fermi, 69622 Villeurbanne, France*

(Received 15 January 2013; published 27 February 2013)

We revisit the role of color mixing in the quark model calculation of tetraquark states, and compare simple pairwise potentials to more elaborate string models with three- and four-body forces. We attempt to disentangle the improved dynamics of confinement from the approximations made in the treatment of the internal color degrees of freedom.

DOI: [10.1103/PhysRevD.87.034040](https://doi.org/10.1103/PhysRevD.87.034040)

PACS numbers: 12.39.Jh, 12.40.Yx, 31.15.-p

I. INTRODUCTION

There is a persisting interest in the quark dynamics applied to multi-quark spectroscopy. The question is whether there exist compact hadron states beyond ordinary mesons (quark-antiquark) and baryons (three quarks).

In simple constituent models, several mechanisms have been proposed for binding multi-quarks: chiral dynamics for light quarks, chromomagnetism, clustering of heavy quarks in a chromoelectric potential, etc. In this article, we concentrate on this latter effect; i.e., multi-quark binding in a spin- and flavor-independent potential, and discuss the role of the internal color degrees of freedom. The validity of the existing models is beyond the scope of this paper. In particular, we shall not discuss the transition from color as a local gauge invariance to color as a global property of the wave function in constituent models. Still, even in its simplified version, color is a delicate ingredient of multi-quark dynamics.

The first studies on multi-quarks within constituent models were based on simple color-additive potentials, extrapolated from meson and baryon spectroscopy. Already for baryons, one can hardly justify the choice of a pairwise interaction, half as strong as the quark-antiquark potential, but a more elaborate modeling, based on a connected Y -shape flux tube linking the three quarks, does not change the results significantly.

We shall follow in this paper this picture of a minimal string linking the quarks. There are alternative nontrivial pictures of confinement—in particular, the ones based on diquarks, which have been extended from the baryon sector to the multi-quarks. See, for instance, Refs. [1–3].

The Y -shape potential of baryons has been extended to tetraquarks and higher multi-quark configurations [4].

There are multi- Y connected diagrams in which the string interaction links all quarks and antiquarks as a Steiner tree whose cumulated length is minimized. But the dynamics is dominated by the so-called “flip-flop” diagrams, with disconnected flux tubes for each of the quark-antiquark, three-quark or three-antiquark subclusters: the attraction comes from the minimum taken over all possible permutations of the quarks and of the antiquarks.

This flip-flop interaction contains three-body, four-body, and higher-order terms, and is thus more delicate to handle in variational calculations. Moreover, when two quarks are exchanged, the color wave function is modified. In the latest studies [5–7], this effect is not treated rigorously. Instead, a type of adiabatic approximation is used: for any set of coordinates for the quarks, the potential is taken as the minimum of all permutations of the quarks and antiquarks, irrespective of the color wave function, and this minimum, as a function of the coordinates, is interpreted as an effective potential leading to a few-body spectral problem in which color has disappeared. Interestingly, this strategy leads to stable multi-quarks for a large variety of constituent masses. This is at variance with the color-additive model, which binds only tetraquarks for large values of the quark-to-antiquark mass ratio.

The question is thus whether the multi-quark binding obtained in string models survives a nonadiabatic treatment of the internal color degrees of freedom. The problem is analyzed in the present paper by reformulating the string-based interaction as an operator in color space.

The outline is as follows: In Sec. II, we set the notation for the color components of tetraquarks. In Sec. III, we present various models of tetraquark confinement with coupled channels in color space, or an adiabatic approximation in which color disappears from the final wave equation. The results are presented in Sec. IV, and the conclusions in Sec. V.

*javier.vijande@uv.es

†valcarce@usal.es

‡j-m.richard@ipnl.in2p3.fr

II. COLOR STATES

The internal color structure of tetraquark states is described in several papers; see, e.g., Refs. [8–10]. We shall borrow the notation of Ref. [8]—in particular, the names “true baryonium” (T) and “mock baryonium” (M)—though the physics context is rather different in the present heavy-quark spectroscopy as compared to the color chemistry of the late 1970s.

The wave function for $(1, 2, 3, 4) = (qq\bar{q}\bar{q})$ is written as

$$\Psi = \psi_T|T\rangle + \psi_M|M\rangle, \quad (1)$$

where $|T\rangle$ denotes a color state with the two quarks in a color $\bar{3}$ state and the antiquarks in a color 3 state, while $|M\rangle$ corresponds to a color sextet in the quark sector and an antisextet in the antiquark one. There is also the possibility of building the global color state out of quark-antiquark clusters in either a color singlet or octet. More precisely, we introduce

$$\begin{aligned} |T\rangle &= |(12)_{\bar{3}}(34)_3\rangle, & |M\rangle &= |(12)_6(34)_{\bar{6}}\rangle, \\ |1\rangle &= |(13)_1(24)_1\rangle, & |8\rangle &= |(13)_8(24)_8\rangle, \\ |1'\rangle &= |(14)_1(23)_1\rangle, & |8'\rangle &= |(14)_8(23)_8\rangle. \end{aligned} \quad (2)$$

The relations between the different sets can be deduced from

$$\begin{aligned} |1\rangle &= \sqrt{\frac{1}{3}}|T\rangle + \sqrt{\frac{2}{3}}|M\rangle, & |8\rangle &= -\sqrt{\frac{2}{3}}|T\rangle + \sqrt{\frac{1}{3}}|M\rangle, \\ |1'\rangle &= -\sqrt{\frac{1}{3}}|T\rangle + \sqrt{\frac{2}{3}}|M\rangle, & |8'\rangle &= \sqrt{\frac{2}{3}}|T\rangle + \sqrt{\frac{1}{3}}|M\rangle. \end{aligned} \quad (3)$$

Accordingly, the matrix elements of the potential in any basis are related to the ones in another basis—for instance,

$$V_{11} = \langle 1|V|1\rangle = \frac{1}{3}V_{TT} + \frac{2}{3}V_{MM} + \frac{2\sqrt{2}}{3}V_{TM}, \quad (4)$$

and many similar relations.

As stressed by Lipkin [11], in the limit of a tetraquark with two units of heavy flavor ($QQ\bar{q}\bar{q}$), with a large quark-to antiquark mass ratio M/m , the ground state is an almost pure $|T\rangle$ state, with the two flavored quarks in an antitriplet state, as in ordinary (QQq) baryons, and the two antiquarks neutralizing that color, as in ($\bar{Q}\bar{q}\bar{q}$) antibaryons. In other words, the tetraquark state in the large M/m limit just uses well-probed color structures, such as the $3 \otimes 3 \rightarrow \bar{3}$ coupling of two quarks in baryons.

On the other hand, for smaller values of the mass ratio M/m , the simple models give at best a very shallow binding. Then the mixing of $|T\rangle$ and $|M\rangle$ is crucial for establishing the stability. See, e.g., Brink and Stancu [12].

III. MODELS OF TETRAQUARK CONFINEMENT

In the early days of multiquark calculations, the potential was assumed to be pairwise, with the color dependence associated with the exchange of a color octet, namely

$$V = -\frac{3}{16} \sum_{i<j} \tilde{\lambda}_i \tilde{\lambda}_j v(r_{ij}). \quad (5)$$

Here, $\tilde{\lambda}_i$ is the color operator for the i th quark, and is suitably modified for an antiquark belonging to the $\bar{3}$ representation of $SU(3)$. The normalization is such that $v(r_{ij})$ is the central part of the quarkonium potential.

This model, however crude, gives at least the possibility of studying the role of the internal color degrees of freedom inside a multiquark state.

The additive model being subject to heavy criticism, an alternative was sought, inspired by the strong coupling regime of QCD. It is referred to as the *flip-flop* model. For mesons, the potential is $V = \sigma r$, where r is the quark-antiquark distance and σ the string tension, which can be set to $\sigma = 1$ by rescaling, without loss of generality. For baryons, this is the Y -shape interaction $V_Y = \min_a(r_{1a} + r_{2a} + r_{3a})$. For a tetraquark, the potential is the minimum of the flip-flop term and a connected string, namely [4,5]

$$\begin{aligned} V_4 &= \min(V_{\text{ff}}, V_s), & V_{\text{ff}} &= \min(r_{13} + r_{24}, r_{14} + r_{23}), \\ V_s &= \min_{a,b}(r_{1a} + r_{2a} + r_{ab} + r_{3b} + r_{4b}), \end{aligned} \quad (6)$$

as pictured in Fig. 1.

For each set of quark coordinates r_i , the minimization in Eq. (6) implies a rotation in color space. This means that V_4 is an adiabatic approximation, which tends to overbind the system. More importantly, the minimization, at least as it was carried out in Ref. [5], does not account for any antisymmetrization. It just holds for distinguishable quarks or antiquarks.

As the model in Eq. (6) gives an interesting spectrum of stable tetraquarks [5]—and, if extended to higher configurations, a spectrum of bound pentaquarks and hexaquarks [6,7]—it is crucial to estimate the amount of overbinding due to the adiabatic approximation, and the changes occurring when a proper antisymmetrization is implemented.




FIG. 1 (color online). Top: Schematic picture of the quark-antiquark and three-quark confinement. Bottom: Three contributions to the tetraquark potential; the simple string model takes the minimum of these three contributions.

We aim at constructing an operator in color space that tends to Eq. (6) in the adiabatic limit, at least when one of the three terms is clearly the minimum. For instance, if the (1, 3) pair is clustered and lies far from the (2, 4) pair, which is also clustered, the potential is more easily described in the $\{|1\rangle, |8\rangle\}$ basis. However, working solely in this latter basis would require the choice of a value for the string tension between color-octet objects (there are studies within QCD; see, e.g., Ref. [13]) and an ansatz for the transition potential V_{18} . Instead, we shall combine the pieces of information coming from the singlet-singlet and triplet-antitriplet states, to deduce the full 2×2 matrix of the potential in the $\{|T\rangle, |M\rangle\}$ basis, in which the four-body problem will be solved.

More precisely, we will consider four different models:

- (1) Model A is the adiabatic limit given by Eq. (6), already used in Ref. [5]. However, the Steiner tree with two junctions, which plays a marginal role, is neglected. Hence, this is the pure flip-flop model.
- (2) Model B is a smooth version of the adiabatic approximation. We use $g(x) = 1/(1+x^n)$ and its complement $g'(x) = 1-g(x)$, with some large but finite exponent $n=5$ to soften the transition between different limiting regimes of the string model. In practice, we replace the above V_{TT} by

$$\hat{V}_{TT} = g_3 g'_3 V_{TT} + (1 - g_3 g'_3) \times \left[\frac{3V_{1'1'} + V_{11}}{4} g_1 + \frac{3V_{11} + V_{1'1'}}{4} g'_1 \right], \quad (7)$$

where

$$V_{11} = r_{13} + r_{24}, \quad V_{1'1'} = r_{14} + r_{23}, \quad (8)$$

and

$$g_3 = g\left(\frac{V_{33}}{V_{11}}\right), \quad g'_3 = g\left(\frac{V_{33}}{V_{1'1'}}\right), \quad (9)$$

$$g_1 = 1 - g'_1 = g\left(\frac{V_{11}}{V_{1'1'}}\right).$$

Then, the potential is known in any basis from V_{11} , $V_{1'1'}$, and \hat{V}_{TT} . For instance, in the $\{|1\rangle, |8\rangle\}$ basis, one uses V_{11} and

$$V_{18} = \frac{1}{4\sqrt{2}} \{V_{11} + 3V_{1'1'} - 4\hat{V}_{TT}\}, \quad (10)$$

$$V_{88} = \frac{1}{4} \{-V_{11} + 3V_{1'1'} + 2\hat{V}_{TT}\},$$

and it is readily checked that if $g_1 \rightarrow 1$, the potential becomes diagonal. The four-body problem is more conveniently solved in the $\{|T\rangle, |M\rangle\}$ basis, and besides \hat{V}_{TT} , the relevant matrix elements are

$$V_{TM} = \frac{3}{4\sqrt{2}} \{V_{11} - V_{1'1'}\}, \quad (11)$$

$$V_{MM} = \frac{1}{4} \{3V_{11} + 3V_{1'1'} - 2\hat{V}_{TT}\}.$$

- (3) Model C is the color-additive model of Eq. (5), normalized to a unit string tension for quarkonium.
- (4) Model D is the crude adiabatic limit of the previous one. This means that for any given set of positions, the 2×2 matrix consisting of V_{TT} , V_{TM} , and V_{MM} of model C is diagonalized, and the lowest eigenvalue is taken as the effective four-body potential, irrespective of any symmetrization or antisymmetrization. Of course, model D is more attractive than model C.
- (5) Several other variants have been envisaged but abandoned as leading to collapses or inconsistencies. For instance, it is tempting to use relations similar to Eq. (11) to express V_{TM} and V_{MM} from V_{11} , $V_{1'1'}$, and V_{TT} , as given by the string model. But then the Hamiltonian is not bounded below.

IV. RESULTS

The results are shown in Table I. The aim is not to provide a benchmark of four-body calculations. What really matters is how the ground-state energy evolves when going from one model to another. The variational estimate has been carried out using just a few Gaussians, and in the case of models B and C, imposing the proper symmetry.

For this purpose, we make use of a wave function with the relevant symmetry for each color component. As the color vector $|T\rangle$ ($|M\rangle$) is antisymmetric (symmetric) under both the exchange of the identical quarks and the identical antiquarks, it has to be combined with a radial wave function with the proper symmetries. The way of constructing these wave functions has been explicitly detailed in Ref. [10]. We will just draft here its main characteristics. The radial wave function is taken as a linear combination of generalized Gaussians depending on six variational parameters a_{ij} of the form

$$\Psi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \sum_{k=1}^4 \alpha_k \exp\left[-\sum_{i \geq j=1}^3 a_{ij} s_{ij}^k \mathbf{x}_i \cdot \mathbf{x}_j\right], \quad (12)$$

TABLE I. Results for the various models A, B, C, and D, as a function of the quark-to-antiquark mass ratio M/m .

M/m	A	B	C	D	Threshold
1	4.644	4.803	4.702	4.596	4.676
2	4.211	4.306	4.275	4.160	4.248
3	4.037	4.131	4.112	3.984	4.086
4	3.941	4.041	4.010	3.891	3.998
5	3.880	3.985	3.954	3.828	3.942
10	3.742	3.860	3.834	3.685	3.831

where s^k are six-component (ij) vectors made of arrangements of positive and negative signs. Once combined with the proper election of the signs α_k , they give rise to radial wave functions with the following symmetries in the radial space under the exchange of quarks and antiquarks: SS , SA , AS , or AA , where S stands for symmetric and A for antisymmetric.

The results from models A and D are very similar for the variational energies. This means that the extra attraction noticed in Ref. [5] is mainly due to the color mixing in the adiabatic approximation. Restoring the interaction as an operator in color space is less favorable for multi-quark binding, and it is readily seen that models B and C lead to comparable predictions. The color structure of the wave functions is also rather similar in models B and C, with mostly a singlet-singlet configuration.

However, if one looks at the details, it can be realized that the flip-flop and the additive models differ. In particular, the average QQ separation, $\langle r_{12} \rangle$, is smaller in the additive model than in the flip-flop model. If the mass ratio $M/m \rightarrow \infty$, this effect becomes more pronounced, with $\langle r_{12} \rangle \rightarrow 0$ in the former case and $\langle r_{12} \rangle$ evolving to a finite value (for m fixed) in the latter one. This influences the contribution of the W exchange to the weak decay of $(bc\bar{q}\bar{q})$, which also plays a role in the decay of (bcu) baryons [14].

This large M/m limit is rather interesting. A detailed numerical study would involve the computation of an effective QQ interaction, $V_{\text{eff}}(r_{12})$, the sum of the direct QQ interaction and the $\bar{q}\bar{q}$ binding energy around them, very similar to the Heitler-London potential of two protons in the adiabatic treatment of the hydrogen molecule. Already, for baryons with two heavy quarks, (QQq) , it has been stressed that the Born-Oppenheimer approximation works very well to reproduce the properties of the lowest states [15]. For $(QQ\bar{q}\bar{q})$, it is striking that the effective potentials $V_{\text{eff}}(r_{12})$ have qualitative differences in the additive model and in the flip-flop one. In the additive model, the minimum of $V_{\text{eff}}(r_{12})$ is reached at $r_{12} = 0$. The $\bar{q}\bar{q}$ contribution is stationary there. So, up to a constant, the potential is dominated by the direct QQ term, which is $r_{12}/2$ here. Thus, the average separation decreases as $M^{-1/3}$, according to the well-known scaling laws in a linear potential [16]. In the flip-flop model, $V_{\text{eff}}(r_{12})$ is minimum at some finite distance. This can be seen directly. This is also compulsory, if one wishes to




FIG. 2 (color online). Flip-flop potential for $r_{12} \rightarrow 0$ and cumulated potential of two mesons.




FIG. 3 (color online). Some weak decays of the (bcu) baryon and $(bc\bar{c})$ meson leading to a tetraquark in the final state.

understand why $(QQ\bar{q}\bar{q})$ is stable in the large M/m limit, as seen by the following *reductio ad absurdum*: Suppose that in the Born-Oppenheimer limit of the flip-flop model, $\langle r_{12} \rangle \rightarrow 0$. Then the flip-flop energy of $(QQ\bar{q}\bar{q})$ and the energy of its threshold, $(Q\bar{q}) + (Q\bar{q})$, which are pictured in Fig. 2 and given by

$$V_{\text{eff}}(r_{12}) = \min(r_{13} + r_{24}, r_{14} + r_{23}),$$

$$V_{\text{th}} = r_{13} + r_{24},$$
(13)

would coincide if $r_{12} = 0$, and the inequality among energies $(QQ\bar{q}\bar{q}) < (Q\bar{q}) + (Q\bar{q})$ would be impossible.

V. CONCLUSIONS

Our results illustrate the role of antisymmetrization in preventing a proliferation of multi-quarks. In particular, the main difference between the naive color-additive model and the crude flip-flop model comes from the treatment of color. It remains that in such a flavor-independent confining, the binding of $(QQ\bar{q}\bar{q})$ is obtained for heavy enough quarks.

More ambitious tools are obviously needed to handle the four-quark dynamics. In particular, in the double charm sector, the binding effect of the confining interaction is probably not sufficient. In addition, the spin part of the wave function can be made favorable: as stressed, e.g., in Ref. [17] and references therein, there is a light-light interaction in $(cc\bar{u}\bar{d})$ which is absent in the two mesons constituting the threshold.

The production of such tetraquarks can be accessible in B factories [17] and at proton-proton colliders. Note also that $(cc\bar{q}\bar{q})$ can be a decay product of hadrons containing

charm and beauty. From a (bcu) baryon, for instance, the Cabibbo-allowed $b \rightarrow c + W^- \rightarrow c + d + \bar{u}$, combined to a $d\bar{d}$ pair creation, leads to $(bcu) \rightarrow (cc\bar{u}\bar{d}) + (ddu)$. See Fig. 3. From $(b\bar{c})$, one could first envisage the Cabibbo-suppressed $b \rightarrow u + W^- \rightarrow u + \bar{c} + s$, and after the creation of a light quark-antiquark pair, this would monitor a decay $(b\bar{c}) \rightarrow (\bar{c}\bar{c}ud) + (s\bar{d})$. Of course, the CKM suppression factor is rather effective here. Perhaps more promising is the chain $b \rightarrow c + W^- \rightarrow c + \bar{c} + d$ giving altogether $(c\bar{c}u\bar{u}d\bar{c})$ after a $u\bar{u}$ pair creation. This could lead to $\bar{D}^- + X$, where \bar{D} is an anticharmed meson and X one of the new hidden charm resonances reviewed, e.g., in Refs. [18,19]. Another combination is $(\bar{c}\bar{c}ud) + (c\bar{u})$, with, however, a different topology of the quark diagram and thus different color and suppression by the

Okubo-Zweig-Iizuka rule factors, as discussed by Lipkin in a different context [20]. Anyhow, any heavy-quark factory should lead to the discovery of heavy tetraquarks with suitable triggers.

ACKNOWLEDGMENTS

We are very grateful to several colleagues for very useful discussions, in particular Makoto Oka, Paolo Gambino, and Qiang Zhao. This work has been partially funded by the Spanish Ministerio de Educación y Ciencia and EU FEDER under Contracts No. FPA2010-21750 and No. AIC-B-2011-0661, by the Spanish Consolider-Ingenio 2010 Program CPAN (CSD2007-00042), and by Generalitat Valenciana Prometeo/2009/129.

-
- [1] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, *Phys. Rev. D* **71**, 014028 (2005).
- [2] D. Ebert, R. Faustov, and V. Galkin, *Phys. Lett. B* **634**, 214 (2006).
- [3] S. Dubnicka, A. Z. Dubnickova, M. A. Ivanov, and J. G. Korner, *Phys. Rev. D* **81**, 114007 (2010).
- [4] J. Carlson and V. R. Pandharipande, *Phys. Rev. D* **43**, 1652 (1991).
- [5] J. Vijande, A. Valcarce, and J. M. Richard, *Phys. Rev. D* **76**, 114013 (2007).
- [6] J.-M. Richard, *Phys. Rev. C* **81**, 015205 (2010).
- [7] J. Vijande, A. Valcarce, and J.-M. Richard, *Phys. Rev. D* **85**, 014019 (2012).
- [8] H.-M. Chan and H. Høgaasen, *Phys. Lett.* **72B**, 121 (1977).
- [9] H.-M. Chan, M. Fukugita, T. H. Hansson, H. J. Hoffman, K. Konishi, H. Høgaasen, and S. Tsun Tsou, *Phys. Lett.* **76B**, 634 (1978).
- [10] J. Vijande and A. Valcarce, *Symmetry* **1**, 155 (2009).
- [11] H. J. Lipkin, *Phys. Lett. B* **172**, 242 (1986).
- [12] D. Brink and F. Stancu, *Phys. Rev. D* **49**, 4665 (1994).
- [13] G. S. Bali, *Phys. Rev. D* **62**, 114503 (2000).
- [14] J. Körner, M. Kramer, and D. Pirjol, *Prog. Part. Nucl. Phys.* **33**, 787 (1994).
- [15] S. Fleck and J. M. Richard, *Prog. Theor. Phys.* **82**, 760 (1989).
- [16] C. Quigg and J. L. Rosner, *Phys. Rep.* **56**, 167 (1979).
- [17] T. Hyodo, Y.-R. Liu, M. Oka, K. Sudoh, and S. Yasui, [arXiv:1209.6207](https://arxiv.org/abs/1209.6207).
- [18] M. Nielsen, F. S. Navarra, and S. H. Lee, *Phys. Rep.* **497**, 41 (2010).
- [19] A. Valcarce, T. F. Caramés, and J. Vijande, *Few-Body Syst.*, 1 (2012).
- [20] H. Lipkin, *Phys. Lett. B* **433**, 117 (1998).