

**Causality-violating Higgs singlets at the LHC**

Chiu Man Ho\* and Thomas J. Weiler†

*Department of Physics and Astronomy, Vanderbilt University, Nashville, Tennessee 37235, USA*

(Received 15 August 2011; published 5 February 2013)

We construct a simple class of compactified five-dimensional metrics which admits closed timelike curves (CTCs), and derive the resulting CTCs as analytic solutions to the geodesic equations of motion. The associated Einstein tensor satisfies all the null, weak, strong and dominant energy conditions. In particular, no negative-energy “tachyonic” matter is required. In extra-dimensional models where gauge charges are bound to our brane, it is the Kaluza-Klein modes of gauge singlets that may travel through the CTCs. From our brane point of view, many of these Kaluza-Klein modes would appear to travel backward in time. We give a simple model in which time-traveling Higgs singlets can be produced by the LHC, either from decay of the Standard Model Higgs or through mixing with the Standard Model Higgs. The signature of these time-traveling singlets is a secondary decay vertex preappearing before the primary vertex which produced them. The two vertices are correlated by momentum conservation. We demonstrate that preappearing vertices in the Higgs singlet-doublet mixing model may well be observable at the LHC.

DOI: [10.1103/PhysRevD.87.045004](https://doi.org/10.1103/PhysRevD.87.045004)

PACS numbers: 14.80.Bn, 04.50.-h, 04.20.Gz, 12.60.Fr

**I. INTRODUCTION**

Time travel has always been an ambitious dream in science fiction. However, the possibility of building a time machine could not even be formulated as science until the discovery of special and general relativity by Einstein. From the early days of general relativity onward, theoretical physicists have realized that closed timelike curves (CTCs) are allowed solutions of general relativity, and hence time travel is theoretically possible. Many proposals for CTCs in a familiar four-dimensional Universe have been discussed in the literature. In chronological order (perhaps not the best listing scheme for CTC proposals), proposals include van Stockum’s rotating cylinder [1] (extended much later by Tipler [2]); Gödel’s rotating universe [3]; Wheeler’s spacetime foam [4]; Kerr and Kerr-Newman’s black hole event horizon interior [5]; Morris, Thorne and Yurtsever’s traversable wormholes [6]; Gott’s pair of spinning cosmic strings [7]; Alcubierre’s warp drive [8]; and Ori’s vacuum torus [9]. More additions to the possibilities continue to unfold [10].

Common pathologies associated with these candidate CTCs are that the required matter distributions are often unphysical, tachyonic, unstable under the backreaction of the metric, or violate one or more of the desirable null, weak, strong and dominant energy conditions [11]. These common pathologies have led Hawking to formulate his “chronology protection conjecture” [12], which states that even for CTCs allowed by general relativity, some fundamental law of physics forbids their existence so as to maintain the chronological order of physical processes. The empirical basis for the conjecture is that so far the human species has not observed noncausal processes. The

logical basis for the conjecture is that we do not know how to make sense of a noncausal Universe.

The possibility of time travel leads to many paradoxes. The most famous paradoxes include the “grandfather” and “bootstrap” paradoxes [11]. In the grandfather paradox, one can destroy the necessary initial conditions that lead to one’s very existence; while in the bootstrap paradox, an effect can be its own cause. A further paradox is the apparent loss of unitarity, as particles may appear “now,” having disappeared at another time “then,” and vice versa. However, after almost two decades of intensive research on this subject, Hawking’s conjecture remains a hope that is not mathematically compelling. For example, it has been shown that there are points on the chronology horizon where the semiclassical Einstein field equations, on which Hawking’s conjecture is based, fail to hold [13]. This and related issues have led many physicists to believe that the validity of chronology protection will not be settled until we have a much better understanding of gravity itself, whether quantizable or emergent. In related work, some aspects of chronology protection in string theory have been studied in Refs. [14–17].

Popular for the previous decade has been the idea of Arkani-Hamed-Dimopoulos-Dvali (ADD) [18] that the weakness of gravity on our 4D brane might be explained by large extra dimensions. A lowered Planck mass is accommodated, with field strengths diluted by the extra dimensions as given by Gauss’s law. The hope is that low-scale gravity may ameliorate or explain the otherwise fine-tuned hierarchy ratio  $M_{\text{weak}}/M_{\text{Planck}}$ . In the ADD scenario, all particles with gauge charge, which include all of the Standard Model (SM) particles, are open strings with charged endpoints confined to the brane (our 4D space-time). Gauge singlets, which include the graviton, are closed strings which may freely propagate throughout the brane and bulk (the extra dimensions). After all, wherever

\*chiuman.ho@vanderbilt.edu

†tom.weiler@vanderbilt.edu

there is spacetime, whether brane or bulk, there is Einstein's gravity. Gauge singlets other than the graviton are speculative. They may include sterile neutrinos and scalar singlets. Due to mixing with gauge nonsinglet particles, e.g., active neutrinos or SM Higgs doublets, respectively, sterile neutrinos or scalar singlets will attain a nongravitational presence when they traverse the brane.

A generic feature in this ADD picture is the possibility of gauge singlets taking "shortcuts" through the extra dimensions [19–23], leading to superluminal communications from the brane's point of view. The extra dimensions could also be warped [24]. Of particular interest are the so-called asymmetrically warped spacetimes [25] in which space and time coordinates carry different warp factors. Scenarios of large extra dimensions with asymmetrically warped spacetimes are endowed with superluminal travel—a signal, say a graviton, from one point on the brane can take a "shortcut" through the bulk and later intersect the brane at a different point, with a shorter transit time than that of a photon traveling between the same two points along a brane geodesic. This suggests that regions that are traditionally "outside the horizon" could be causally related by gravitons or other gauge singlets. Exactly this mechanism has been invoked as a solution to the cosmological horizon problem without inflation [26]. Although this leads to an apparent causality violation from the brane's point of view, the full 5D theory may be completely causal. Superluminal travel through extra-dimensional shortcuts generally doesn't guarantee a CTC. To obtain a CTC, one needs the light cone in a  $t$ -versus- $r$  diagram to tip below the horizontal  $r$  axis for part of the path. Then, for this part of the path, travel along  $r$  is truly progressing along negative time. When the positive time part of the path is added, one has a CTC if the net travel time is negative.

Recently, there was an exploratory attempt to find a CTC using a spacetime with two asymmetrically warped extra dimensions [27]. In this work, it was demonstrated that paths exist which in fact are CTCs. However, these constructed paths are not solutions of geodesic equations. The construction demonstrated the existence of CTCs in principle for a class of extra-dimensional metrics, but did not present CTCs which would actually be traversed by particles. Since geodesic paths minimize the action obtained from a metric, the conjecture in Ref. [27] was that the same action that admits constructed paths with negative or zero time admits geodesic paths with even greater negative (or zero) time.

The ambitions of this article are threefold: First of all, we seek a class of CTCs embedded in a single compactified extra dimension. We require the CTCs to be geodesic paths, so that physical particles will become negative-time travelers. Secondly, we ensure that this class of CTCs is free of undesirable pathologies. Thirdly, we ask whether particles traversing these CTC geodesics may

reveal unique signatures in large detectors such as ATLAS and CMS at the LHC.

As we demonstrate in this article, we have successfully found a class of 5D metrics which generates exactly solvable geodesic equations whose solutions are in fact CTCs. We adopt an ADD framework where only gauge singlet particles (gravitons, sterile neutrinos, and Higgs singlets) may leave our 4D brane and traverse the CTC embedded in the extra dimension. In this way, the standard paradoxes (described below) are ameliorated, as no macro objects can get transported back in time. Scalar gauge singlets, e.g., Higgs singlets, mixed or unmixed with their gauge nonsinglet siblings, e.g., SM Higgs doublets, may be produced and detected at the LHC. The signature of negative-time travel is the appearance of a secondary decay or scattering vertex *earlier in time* than the occurrence of the primary vertex which produces the time-traveling particle. The two vertices are associated by overall momentum conservation.

Realizing that the grandfather, bootstrap, and unitarity paradoxes may be logically disturbing, we now discuss the paradoxes briefly. First of all, it bears repeating that in the ADD picture, it is only gauge-singlet particles that may travel CTCs. No claims of human or robot transport backwards through time are made. And while the paradoxes are unsettling, as was/is quantum mechanics, we think that it is naive to preclude the possibility of time travel on the grounds of human argument/preference. The paradoxes may be but seeming contradictions resulting from our ignorance of some fundamental laws of physics which in fact enforce consistency [28]. For instance, in Feynman's path integral language, one should sum over all possible globally defined histories. It is possible that histories leading to paradoxes may contribute little or nothing to this sum. In other words, while the grandfather paradox is dynamically allowed by Einstein's field equations, it may be kinematically forbidden due to the inaccessibility of self-contradicting histories in the path integral [29–34]. In the bootstrap paradox, the information, events, or objects in the causal loop do not seem to have an identifiable cause. The entities appear as if they were eternally existing, with the causation being pushed back to the infinite past. But the logic of the bootstrap paradox does not seem to preclude the possibility of time travel in any compelling manner.

The unitarity paradox is unsettling as it seems to suggest that the past can get particles from the future "for free." If nature respects unitarity as one of her most fundamental principles, she may have a consistent way (unknown at present) to implement it even in the face of causality violation. It is also conceivable that nature sacrifices unitarity. Precedent seems to exist in quantum mechanics: the "collapse" of a wave function, at the core of the Copenhagen interpretation of quantum mechanics, is not a unitary process, for such evolution has no inverse—one cannot un-collapse a collapsed wave function. The "many worlds" interpretation restores unitarity in a nonfalsifiable

way. Perhaps there is a similar point of view lurking behind CTCs. While it has been shown that when causality is sacrificed in interacting field theories, then one necessarily loses perturbative unitarity [31–33] (traceable to the fact that the time-ordering assignment in the Feynman propagator is ambiguous on spacetimes with CTCs). It has also been proposed that just this sacrifice of unitarity be made in a “generalized quantum mechanics” [35]. And again, in the class of CTCs we consider, paradox considerations, such as unitarity violation, apply only to the gauge-singlet sector.

Even readers who do not believe in the possibility of time travel may still find aspects of this article of intellectual interest. The process of exploring time travel may provide a glimpse of the ingredients needed to complete Einstein’s gravity. This completion may require a quantized or emergent theory of gravity, and/or higher dimensions, and/or other. Furthermore, we will propose specific experimental searches for time travel, and so stay within the realm of falsifiable physics.

## II. A CLASS OF METRICS ADMITTING CLOSED TIMELIKE CURVES

The success of the ADD model inspires us to think about the possibility of constructing viable CTCs by the aid of extra dimensions. With the criteria of simplicity in mind, we choose a time-independent metric and invoke only a single compactified spatial extra dimension. We consider the following form for the metric:

$$d\tau^2 = \eta_{ij} dx^i dx^j + dt^2 + 2g(u) dt du - h(u) du^2, \quad (2.1)$$

where  $i, j = 1, 2, 3$ ,  $\eta_{ij}$  is the spatial part of the Minkowskian metric, and  $u$  is the coordinate of a spatial extra dimension. For convenience, we set the speed of light  $c = 1$  on the brane throughout the entire article. As guided by the wisdom from previous proposals of CTCs, such as Gödel’s rotating universe [3], we have adopted a nonzero off-diagonal term  $dt du$  for a viable CTC. Another simplicity of the above 5D metric is that its 4D counterpart is completely Minkowskian. The determinant of the metric is

$$\text{Det} \equiv \text{Det}[g_{\mu\nu}] = g^2 + h. \quad (2.2)$$

A weak constraint arises from the spacelike nature of the  $u$  coordinate, which requires the signature  $\text{Det} > 0$  for the whole 5D metric. In turn this requires that  $g^2 + h > 0$  for all values of  $u$ , i.e.,  $h(u) > -g^2(u)$  at all  $u$ . We normalize the determinant by requiring the standard Minkowskian metric on the brane, i.e.,  $\text{Det}(u = 0) = g^2(0) + h(0) = +1$ .

Since we have never observed any extra dimension experimentally, we assume that it is compactified and has the topology  $S^1$  of a one-sphere (a circle). Due to this periodic boundary condition, the point  $u + L$  is identified with  $u$ , where  $L$  is the size of the extra dimension. We do

not specify the compactification scale  $L$  of the extra dimension at this point, as it is irrelevant to our construction of the CTCs. A phenomenologically interesting number is  $L \gtrsim 1/\text{TeV}$  since this opens the possibility of new effects at the LHC. We will adopt this choice in the discussion of possible phenomenology in Sec. VIII.

In the coordinates  $\{x^\mu, u\}$ , our compactified metric with an off-diagonal term  $g(u)$  is reminiscent of a cylinder rotating in  $u$  space, with axis parallel to the brane. Again, this geometry is reminiscent of Gödel’s construction or the van Stockum-Tipler construction. However, their “rotating cylinder” in the usual 4D spacetime is here replaced with an extra dimension having a compactified  $S^1$  topology. In our case as well as theirs, the metric is stationary but not static, containing a nonzero off-diagonal term involving both time and space components.

The elements of the metric tensor must reflect the symmetry of the compactified dimension, i.e., they must be periodic functions of  $u$  with period  $L$ . This in turn requires that  $g(u)$  and  $h(u)$  must have period  $L$ . Any function with period  $L$  can be expressed in terms of a Fourier series with modes  $\sin(\frac{2\pi nu}{L})$  and  $\cos(\frac{2\pi nu}{L})$ ,  $n = 0, 1, 2, \dots$ . Expanded in Fourier modes, the general metric function  $g(u)$  is

$$g(u) = g_0 + A - \sum_{n=1}^{\infty} \left\{ a_n \cos\left(\frac{2\pi nu}{L}\right) + b_n \sin\left(\frac{2\pi nu}{L}\right) \right\}, \quad (2.3)$$

where  $g(0) = g_0$  and  $A \equiv \sum_{n=1}^{\infty} a_n$  are constants. An analogous expansion can be written down for the metric function  $h(u)$ , but in what follows we will not need it.

Below we will demonstrate that the 5D metric we have constructed is sufficient to admit CTCs. It is worth mentioning that our 5D metric is easily embeddable in further extra dimensions.

## III. GEODESIC EQUATIONS AND THEIR SOLUTIONS

On the brane, the metric in Eq. (2.1) is completely Minkowskian. Accordingly, the geodesic equations of motion (EOMs) along the brane are simply a vanishing proper acceleration  $\ddot{\vec{r}} = 0$ , with dot derivative denoting differentiation with respect to the proper time  $\tau$ . Thus,

$$\dot{\vec{r}} = \dot{\vec{r}}_0, \quad \text{or} \quad \vec{r} = \vec{r}_0 \tau. \quad (3.1)$$

The geodesic equations for time and for the bulk direction are more interesting. Since the metric is time independent (“stationary”), there is a timelike Killing vector with an associated conserved quantity; the quantity is

$$\dot{t} + g(u)\dot{u} = \gamma_0 + g_0\dot{u}_0, \quad (3.2)$$

where on the right-hand side, we have written the constant in an initial-value form. The initial value of  $\dot{t}$ , on the brane, is just the boost factor  $\gamma_0$ . From this conserved quantity, we

may already deduce that time will run backwards, equivalently, that  $\dot{t} < 0$ , if  $g(u)\dot{u} > \gamma_0 + \dot{u}_0 g_0$  is allowed by the remaining geodesic equation. The remaining geodesic equation involving the bulk coordinate  $u$  is

$$2(g\ddot{t} - h\ddot{u}) - h'\dot{u}^2 = 0, \quad (3.3)$$

where the superscript ‘‘prime’’ denotes differentiation with respect to  $u$ .

Taking the dot derivative of Eq. (3.2), we may separately eliminate  $\ddot{t}$  and  $\ddot{u}$  from Eq. (3.3) to rewrite Eqs. (3.2) and (3.3) as

$$\ddot{t}(\tau) = \frac{1}{2} \frac{-2g'h + gh'}{g^2 + h} \dot{u}^2, \quad (3.4)$$

$$\ddot{u}(\tau) = -\frac{1}{2} \frac{2gg' + h'}{g^2 + h} \dot{u}^2 = -\frac{1}{2} \ln'(g^2 + h) \dot{u}^2. \quad (3.5)$$

The latter geodesic equation is readily solved with the substitution  $\xi \equiv \dot{u}$ , which implies that  $\ddot{u} = \dot{\xi} = (d\xi/du)(du/d\tau) = \xi(u)(d\xi/du)$ . Let us choose the initial conditions to be that at  $\tau = 0$ , we have  $u = 0$ . The solutions for  $\dot{u}$  and  $u$  are

$$\dot{u}(u) = \frac{\dot{u}_0}{\sqrt{g^2(u) + h(u)}}, \quad (3.6)$$

and

$$\int_0^{u(\tau)} du \sqrt{g^2 + h} = \dot{u}_0 \tau, \quad (3.7)$$

the latter being an implicit solution for  $u(\tau)$ . Having solved explicitly for  $\dot{u}(u)$  in Eq. (3.6), we may substitute it into the first of Eq. (3.4) to gain an equation for  $t(u)$ . Alternatively, we may solve the implicit equation in Eq. (3.7) for  $u(\tau)$ , and substitute it into Eq. (3.2) to get

$$t(\tau) = (\gamma_0 + g(0)\dot{u}_0)\tau - \int_0^{u(\tau)} dug(u). \quad (3.8)$$

The geodesic equations Eqs. (3.6) and (3.7) depend on  $\text{Det} = g^2 + h$  but not on  $g$  or  $h$  individually. It therefore proves to be simple and fruitful to fix the determinant to

$$\text{Det}(u) = g^2(u) + h(u) = 1, \quad \forall u. \quad (3.9)$$

We do so. With this choice, one readily obtains the EOM  $\ddot{u} = 0$ , which implies the solutions

$$\dot{u}(\tau) = \dot{u}_0, \quad (3.10)$$

and

$$u(\tau) = \dot{u}_0 \tau, \quad (\text{mod } L). \quad (3.11)$$

In analogy to the historical CTCs arising from metrics containing rotation, we will call the geodesic solutions with positive  $\dot{u}_0$  ‘‘co-rotating,’’ and solutions with negative  $\dot{u}_0$  ‘‘counter-rotating’’. So a co-rotating (counter-rotating) particle begins its trajectory with positive (negative)  $\dot{u}_0$ .

We note already at this point the possibility for periodic travel in the  $u$  direction with negative time. From Eqs. (3.2) and (3.10), we have

$$\dot{t} = \gamma_0 - (g(u) - g_0)\dot{u}_0, \quad (3.12)$$

and its value averaged over the periodic path of length  $L$

$$\bar{\dot{t}} = \frac{1}{L} \int_0^L d\tau \dot{t} = \gamma_0 - (\bar{g} - g_0)\dot{u}_0, \quad (3.13)$$

where

$$\bar{g} = \frac{1}{L} \int_0^L g(u) du = g_0 + A \quad (3.14)$$

is the average value of the metric element along the compact extra dimension. The latter equality follows immediately from Eq. (2.3).

Apparently, closing the path in negative time will depend on the relation between the mean value  $\bar{g}$  and the value of the element on the brane  $g_0$ , and on the relation between the velocities of the particle along the brane and along the bulk, characterized by  $\gamma_0$  and  $\dot{u}_0$ . In the next subsections we examine this possibility in detail.

### A. The CTC possibility

By definition, a CTC is a geodesic that returns a particle to the same space coordinates from which it left, with an arrival time before it left. The ‘‘closed’’ condition of the CTC can be satisfied easily in our metric due to the  $S^1$  topology of the extra dimension. Namely, if a particle created on the brane propagates into the extra dimension, it will necessarily come back to  $u = 0$  due to the periodic boundary condition.<sup>1</sup> The other condition for a CTC, the ‘‘timelike’’ condition, is that the time elapsed during the particle’s return path as measured by an observer sitting at the initial space coordinates is negative. To ascertain the time of travel, and its sign, we proceed to solve for  $t(u)$ . As indicated by Eq. (3.8), to do so we need to specify  $g(u)$ .<sup>2</sup> Our Fourier expansion of the general compactified metric function [Eq. (2.3)] is sufficient for this task.

Our remaining task is to determine  $t(u)$  and see if it can be negative. From Eqs. (3.8) and (3.11) we have

$$t(u) = \left(g_0 + \frac{1}{\beta_0}\right)u - \int_0^u dug(u), \quad (3.15)$$

where we find it useful to define the symbol

<sup>1</sup>The geodesic equations for travel along the brane are trivially just constancy of the three-vector part  $\vec{r}$  of the four-velocity. Added to the geodesic solution for  $u(\tau)$ , one gets a constant translation of the circle  $S^1$  along the brane, resulting in a helical motion which periodically intersects the brane (see Sec. VI).

<sup>2</sup>Once  $g(u)$  is specified,  $h(u)$  is given by  $h(u) = 1 - g^2(u)$ , due to the choice made in Eq. (3.9). In particular, the periodicity imposed in  $g(u)$  now automatically ensures that  $h(u)$  is periodic, too.



$$\beta_0 = \frac{\dot{u}_0}{\gamma_0} = \left( \frac{du}{dt} \right)_0, \quad (3.16)$$

for the initial velocity of the particle in the  $u$  direction as would be measured by a stationary observer on the brane. For the co-rotating particle,  $\beta_0 > 0$ , while for the counter-rotating particle,  $\beta_0 < 0$ . Performing the integral over Eq. (2.3) as prescribed in Eq. (3.15), we arrive at

$$t(u) = \left( \frac{1}{\beta_0} - A \right) u + \left( \frac{L}{2\pi} \right) \sum_{n=1}^{\infty} \left( \frac{1}{n} \right) \left[ a_n \sin\left( \frac{2\pi n u}{L} \right) + b_n \left[ 1 - \cos\left( \frac{2\pi n u}{L} \right) \right] \right]. \quad (3.17)$$

Due to the  $S^1$  topology of the compactified extra dimension, the particle returns to the brane at  $u = \pm NL$ , for integer  $N > 0$ . The plus (minus) sign holds for a co-rotating (counter-rotating) particle. Physically,  $N$  counts the number of times that the particle has traversed the compactified extra dimension. When the particle crosses the brane for the  $N$ th time, the time as measured by a stationary clock on the brane is

$$t_N \equiv t(u = \pm NL) = \pm \left( \frac{1}{\beta_0} - A \right) NL. \quad (3.18)$$

This crossing time depends on the Fourier modes only through  $A = \sum_{n=1} a_n$ , and is independent of the  $b_n$ . Thus, the potential for a CTC lies in the cosine modes of the metric function  $g(u)$ , and not in the sine modes.<sup>3</sup>

### B. The CTC realized: Negative time travel

A viable CTC is realized only if  $t_N < 0$ , i.e.,  $\pm N(\frac{1}{\beta_0} - A) < 0$ . For a co-rotating particle ( $\beta_0 > 0$  and positive signature), a viable CTC requires the conditions<sup>4</sup>

$$A > \frac{1}{\beta_0} > 1. \quad (3.19)$$

On the other hand, for a counter-rotating particle ( $\beta_0 < 0$  and negative signature), we require

$$A, \beta_0 < 0 \quad \text{and} \quad |A| > \left| \frac{1}{\beta_0} \right| > 1 \quad (3.20)$$

to realize the CTC. In either case, co-rotating or counter-rotating particles, the CTC conditions require that  $\text{sign}(A)$  be the same as  $\text{sign}(\beta_0)$ . Nature chooses the constant  $A$  with a definite sign, and so the CTC conditions for co-rotating and counter-rotating particles are incompatible. For definiteness in what follows, we will assume that it is the co-rotating particles which may traverse the CTC and

<sup>3</sup>This leads to a simple, necessary but not sufficient condition on the metric function for the existence of a CTC: for the  $\text{Det} = 1$  metric,  $g_0$  must differ from  $g(\frac{L}{2})$ .

<sup>4</sup>Note that we here assume that  $|\beta_0| < 1$  when the particle is created on the brane. We are allowed to make this assumption because our metric class will not need superluminal speeds to realize the CTCs.

not the counter-rotating particles, i.e., that  $A > \frac{1}{\beta_0} > 1$  holds for some  $\beta_0$ . The counter-rotating particles of course exist, but they move forward in time.

We note that the negative time of the CTC scales linearly with the number of times  $N$  that the particle traverses the compact  $u$  dimension. The temporal period of this march backwards in time is  $|\frac{1}{\beta_0} - A|L$ , with the natural time scale being  $L/c \sim 10(L/\text{mm})$  picoseconds. We next give a useful analysis of world-line slopes derived from our metric and their connection to time travel. Such an analysis can offer considerable insight into negative-time physics.

## IV. LIGHT-CONE/WORLD-LINE SLOPE ANALYSIS

A light-cone analysis of the metric, in a fashion similar to the one in Ref. [27], is illuminating. Here we will make the slight generalization to world lines of massive particles rather than light cones of massless particles.

It is required for the existence of CTCs that the world line tips so that evolution in the  $u$  direction occurs with backward evolution in time  $t$  as measured from the brane. Let  $\tau$  be the proper time of the massive particle in consideration. When written in terms of the slopes  $s \equiv dt/du$  for the world line in the  $\pm u$  directions, the line element becomes

$$\frac{d\tau^2}{du^2} = \frac{1}{\dot{u}^2(u)} = s^2 + 2sg(u) - h(u), \quad (4.1)$$

and we have neglected possible nonzero  $d\vec{r}/dt$  since it does not affect this discussion. The solutions of this quadratic equation are the two slopes for the co-rotating and counter-rotating world lines:<sup>5</sup>

$$s_{\pm}(u) = -g(u) \pm \sqrt{g^2(u) + h(u) + \frac{1}{\dot{u}^2(u)}}. \quad (4.2)$$

To ascertain the assignment of the two world lines to the co-rotating and counter-rotating particles, we note that by definition,  $s = \frac{dt}{du}$ , so  $s(u=0) = \frac{\gamma_0}{\dot{u}_0}$ . Thus,  $\text{sign}(s(0)) = \text{sign}(\dot{u}_0)$ , i.e.,  $s_+$  is the world line for the  $\dot{u} > 0$  co-rotating particle, and  $s_-$  is the world line for the  $\dot{u} < 0$  counter-rotating particle.

Equivalent to Eq. (4.2) are

$$s_-(u) + s_+(u) = -2g(u) \quad \text{and} \quad (4.3)$$

$$s_-(u)s_+(u) = -\left( h(u) + \frac{1}{\dot{u}^2(u)} \right).$$

If the  $u$  direction were not warped, we would expect the Minkowskian result  $s_- + s_+ = 0$  and  $s_-s_+ = -(1 + \dot{u}^{-2})$ . Instead, here we have  $s_-(0) + s_+(0) = -2g_0$  and  $s_-(0)s_+(0) = -h_0 + \dot{u}_0^{-2}$ , where

<sup>5</sup>One obtains the (massless particle) light-cone results by setting  $\frac{1}{\dot{u}^2}$  (proportional to  $d\tau^2$ ) to zero.




FIG. 1 (color online).  $g(u)$  (dashed) and  $h(u) = 1 - g^2(u)$  (solid) versus  $u/L$ , for parameter choices  $g_0 = -0.9$ ,  $a_1 = A = 1.8$ , and  $a_{n \neq 1} = b_n = 0$ .

$h_0 \equiv h(0)$ . So for  $g_0 \neq 0$  and/or  $h_0 \neq 1$ , we have a warped dimension beginning already at the brane slice  $u = 0$ . We maintain Minkowskian-like behavior at  $u = 0$  by choosing  $h_0$  to be non-negative.

For the world line to tip into the negative  $t$  region, its slope must pass through zero. This requires the product  $s_-(u)s_+(u)$  and hence  $h(u) + \dot{u}^{-2}$  to pass through zero. We label the value of  $u$  where this happens as  $u_*$ . Thus,  $h(u_*) = -\dot{u}_*^{-2}$ . Positivity of the metric determinant  $g^2 + h$  at all  $u$  then demands at  $u_*$ , where  $h(u_*) + \dot{u}_*^{-2}$  vanishes, that  $g^2(u_*) > \dot{u}_*^{-2} \neq 0$ .

Next we implement our simplifying assumption that  $g^2 + h = 1$  and its concomitant result  $\dot{u} = \dot{u}_0$ ,  $\forall u$  [Eq. (3.10)]. As a result, (i) the condition  $h_0 \geq 0$  in turn implies that  $|g_0| \leq 1$ ; (ii) we have  $g^2(u_*) = 1 - h(u_*) = 1 + \dot{u}_0^{-2}$ , so the condition  $g^2(u_*) \neq 0$  is automatically satisfied.

Importantly, time will turn negative if  $g(u)$  rises from its value  $|g_0| \leq 1$  on the brane to above  $\sqrt{1 + \dot{u}_0^{-2}}$ . Such behavior of  $g(u)$  is easy to accommodate with a metric function as general as Eq. (2.3). In Fig. 1 we show sample curves for  $g(u)$  and  $h(u) = 1 - g^2(u)$ . A quick inspection of the  $g(u)$  shown in this figure convinces one that even a simple metric function can accommodate time travel. In Fig. 2, we show explicitly the accumulation of negative time as the particle travels around and around the extra dimension.

One lesson learned from this slope analysis is that only co-rotating or only counter-rotating particles, but not both, may experience CTCs. This is because only *one* edge of the light cone tips below the horizontal axis into the negative-time half-plane. The development of co-rotating and counter-rotating geodesics in the previous section is consistent with this lesson. Another lesson learned is that CTCs may exist for large  $\dot{u}_0$ , but not for small  $\dot{u}_0$ ; i.e., there may exist a critical  $(\dot{u}_0)_{\min}$  such that CTCs exist for  $\dot{u}_0 > (\dot{u}_0)_{\min}$ , but not for  $\dot{u}_0 < (\dot{u}_0)_{\min}$ . Finally, we remark that the slope analysis presented here may be derived from a more general covariant analysis. The connection is shown in Appendix A.




FIG. 2.  $t(u)$  versus  $u/L$ , for the same parameter choices as in Fig. 1, and with  $\beta_0 = 2/3$ .

## V. COMPACTIFIED 5D CTCS COMPARED/ CONTRASTED WITH 4D SPINNING STRING

Before turning to phenomenological considerations of particles traversing CTCs, we wish to show that the compactified 5D metric admitting CTC geodesics is devoid of pathologies that plague similar 4D metrics. This class of 5D metrics resembles in some ways the well-studied metric which describes a spinning cosmic string.

### A. 4D spinning string(s)

The metric for the 4D spinning string is [36,37]

$$d\tau_{\text{spinning string}}^2 = (dt + 4GJd\theta)^2 - dr^2 - (1 - 4Gm)^2 r^2 d\theta^2 - dz^2, \quad (5.1)$$

where  $G$  is Newton's constant, and  $J$  and  $m$  are the angular momentum and mass per unit length of the cosmic string, respectively. In three spacetime dimensions, the Weyl tensor vanishes and any source-free region is flat. This means that in the region outside of the string, local Minkowski coordinates may be extended to cover the whole region. Specifically, by changing the coordinates in Eq. (5.1) to  $\tilde{t} = t + 4GJ\theta$  and  $\varphi = (1 - 4Gm)\theta$ , the metric appears Minkowskian, with the conformal factor being unity. As with  $\theta$ , the new angular coordinate  $\varphi$  is periodic, subject to the identification  $\varphi \sim \varphi + 2\pi - 8\pi Gm$ . There is a well-known wedge  $\Delta\varphi = 8\pi Gm$  removed from the plane, to form a cone. However, although this transformation appears to be elegant, in fact the new time  $\tilde{t}$  is a pathological linear combination of a compact variable  $\theta$  and a non-compact variable  $t$ : for fixed  $\theta$  (or  $\varphi$ ) one expects  $\tilde{t}$  to be a smooth and continuous variable, while for fixed  $t$ , one expects the identification  $\tilde{t} \sim \tilde{t} + 8\pi GJ$  in order to avoid a "jump" in the new variable. In effect, the singularity at  $g_{\theta\theta} = 0$ , i.e., at  $r = 4GJ/(1 - 4Gm)$ , is encoded in the new but pathological coordinate  $\tilde{t}$  [36].

The form of our metric in the  $(t, u)$  plane, viz.

$$d\tau^2 = dt^2 + 2g(u)dtdu - h(u)du^2 = (dt + g(u)du)^2 - (g^2(u) + h(u))du^2, \quad (5.2)$$

has similarities with the spinning string metric. Analogously, we may define new exact differentials<sup>6</sup>

$$d\bar{t} \equiv dt + g(u)du, \quad \text{and} \quad d\bar{u} \equiv \sqrt{g^2(u) + h(u)}du, \quad (5.6)$$

which puts our metric into a diagonal ‘‘Minkowski’’ form:

$$d\tau^2 = \eta_{ij}dx^i dx^j + d\bar{t}^2 - d\bar{u}^2. \quad (5.7)$$

Being locally Minkowskian everywhere, the entire 5D spacetime is therefore flat. This accords with the theorem which states that any two-dimensional (pseudo)Riemannian metric (whether or not it is in a source-free region)—here, the  $(u, t)$  submanifold—is conformal to a Minkowskian metric. In our case, the geometry is  $M^4 \times S^1$ , which is not only conformally flat, but flat period [the conformal factor  $\Omega(u, t)$  is unity]. However, the topology of our 5D space, like that of the spinning string, is nontrivial. The new variable  $\bar{t}$ , defined by  $d\bar{t} = dt + g(u)du$ , is ill defined globally, being a pathological mixture of a compact ( $u$ ) and a noncompact ( $t$ ) coordinate. Thus, the parallel between the metric for a spinning cosmic string and our metric is clear.

We remark that time as measured by an observer (or experiment) on our brane is just given by the coordinate variable  $t$ . This is seen by constructing the induced 4D metric. The constraint equation reducing the 5D metric to the induced 4D metric is simply  $u(x^\mu) = 0$ . Taking the differential yields  $du = 0$ . Inputting the latter result into the 5D metric of Eq. (2.1) [or into Eq. (5.7)] induces the standard 4D Minkowski metric.

Next we investigate whether or not our metric suffers from fundamental problems commonly found in proposed 4D metrics with CTCs.

## B. 4D spinning string pathologies

Deser *et al.* [36] showed that the metric for the spinning string admits CTCs. This metric has been criticized, by themselves and others, for the singular definition of spin that occurs as one approaches the string’s center at  $r = 0$ .

<sup>6</sup>This nontrivial transformation effectively defines a new time  $\bar{t}$  measured in the frame that co-rotates with the circle  $S^1$ . The integrated, global version of these new coordinates is

$$\bar{u} = \int_0^{\bar{u}} d\bar{u} = \int_0^u du \sqrt{\text{Det}(u)} = u, \quad \text{since } \text{Det} = 1, \quad (5.3)$$

$$\bar{t} = \int_0^{\bar{t}} d\bar{t} = \int_0^t dt + \int_0^{u(t)} g(u)du = t + \int_0^{u(t)} dug(u), \quad (5.4)$$

$$= t + (g_0 + A)u - \left(\frac{L}{2\pi}\right) \sum_{n=1}^{\infty} \left(\frac{1}{n}\right) \left[ a_n \sin\left(\frac{2\pi nu}{L}\right) + b_n \left[ 1 - \cos\left(\frac{2\pi nu}{L}\right) \right] \right]. \quad (5.5)$$

With our metric, there is no ‘‘ $r = 0$ ’’ in  $u$  space—the ‘‘center’’ of periodic  $u$  space is simply not part of spacetime.

An improved CTC was proposed by Gott, making use of a pair of cosmic strings with a relative velocity—spin angular momentum of a single string is replaced with an orbital angular momentum of the two-string system. Each of the cosmic strings is assumed to be infinitely long and hence translationally invariant along the  $z$  direction. This invariance allows one to freeze the  $z$  coordinate, thereby reducing the problem to an effective  $(2 + 1)$ -dimensional spacetime with two particles at the sites of the two string piercings. A CTC is found for a geodesic encircling the piercings and crossing between them.

The nontrivial topology associated with Gott’s spacetime leads to nonlinear energy-momentum addition rules. What has been found is that while each of the spinning cosmic strings carries an acceptable timelike energy-momentum vector, the two-string center-of-mass energy-momentum vector is spacelike or tachyonic [38,39]. Furthermore, it has been shown that in an open universe, it would take an infinite amount of energy to form Gott’s CTC [40].

Blueshifting of the string energy is another argument against the stability of Gott’s CTC [41]. Through each CTC cycle, a particle gets blueshifted [12,40]. Since the particle can traverse the CTC an infinite number of times, it can be infinitely blueshifted, while the time elapsed is negative. Total energy is conserved, and so the energy of the pair of cosmic strings is infinitely dissipated even before the particle enters the CTC for the first time. Hence, no CTC can be formed in the first place.

## C. Pathology-free compactified 5D CTCs

Our class of 5D metrics seems to be unburdened by the pathologies [38,39] described immediately above. One readily finds that all the components of the 5D Einstein tensor as determined by the metric in Eq. (2.1) are identically zero. Therefore, by the Einstein equation, the energy-momentum tensor  $T_{AB}$  also vanishes. This implies that our class of metrics with CTCs automatically satisfies all of the standard energy conditions.<sup>7</sup> Hawking has conjectured [12] that nature universally protects chronology (causality) with

<sup>7</sup>The standard energy conditions are, for any null vectors  $l^A$  and timelike vectors  $t^A$ ,

$$\text{Null energy condition (NEC): } T_{AB}l^A l^B \geq 0, \quad (5.8)$$

$$\text{Weak energy condition (WEC): } T_{AB}t^A t^B \geq 0, \quad (5.9)$$

$$\text{Strong energy condition (SEC): } T_{AB}t^A t^B \geq \frac{1}{2}T_A^A t^B t_B, \quad (5.10)$$

$$\text{Dominant energy condition (DEC): } T_{AB}t^A t^B \geq 0 \quad \text{and} \quad T_{AB}T_C^B t^A t^C \leq 0. \quad (5.11)$$

applications of physical laws. The protection hides in the details. Hawking has proved his conjecture for the case where the proposed CTC violates the weak energy condition (WEC). Such WEC violations typically involve exotic materials having negative energy density. Our CTCs do not violate the WEC, and in fact do not require matter at all. With our metric, it is the compactified extra dimension with the  $S^1$  topology rather than an exotic matter/energy distribution that enables CTCs. In the realm of energy conditions, our metric contrasts again with Gott's case of two moving strings. Gott's situation does not violate the WEC, but it can be shown that the tachyonic total energy-momentum vector leads to violation of all the other energy conditions.

Furthermore, particles traversing our CTCs are *not* blueshifted, unlike the particles traversing Gott's CTC. This can be seen as follows. One defines the contravariant momentum in the usual way, as

$$p^A \equiv m(\dot{t}, \dot{\vec{r}}, \dot{u}), \quad (5.12)$$

where  $m$  is the mass of the particle. Then the covariant five-momentum is

$$p_A = G_{AB}p^B = m(\dot{t} + g\dot{u}, -\dot{\vec{r}}, g\dot{t} - h\dot{u}). \quad (5.13)$$

According to Eq. (3.2), the quantity  $p_0 = m(\dot{t} + g\dot{u})$  is covariantly conserved along the geodesic on and off the brane. The conservation is a result of the time independence of the metric  $G_{AB}$ . Consequently, we identify the conserved quantity as the particle energy  $E$  and conclude that particles traversing the compactified 5D CTCs are not blueshifted. In a more heuristic fashion, one may say that energy conservation on the brane follows from the absence of an energy source;  $T_{AB}$  vanishes for our choice of metric class.

In addition to conservation of  $p_0$ , conservation of the particle's three-momentum along the brane follows immediately from the EOM and solution Eq. (3.1). The only component of the particle's covariant five-momentum that is not conserved along the geodesic is  $p_5 = m(g\dot{t} - h\dot{u})$ . This quantity may be written as  $(gE - m\dot{u})$ , where use has been made of the relation  $g^2 + h = 1$ . But even here, the factors  $E$  and [from Eq. (3.10)]  $\dot{u}$  are conserved quantities, and so it is just the factor  $g(u)$  that varies along the geodesic. However, the metric elements including  $g(u)$  are periodic in  $u = NL$ , and return to their brane values at each brane piercing. Thus, the particle's entire covariant five-momentum is conserved from the viewpoint of the brane. We conclude that the possible instability manifested by a particle's blueshift [12,40,41] does not occur in our class of 5D metric, nor do any other kinematic pathologies. (Conservation and nonconservation of the components of the particle's five-momentum are discussed from another point of view in Appendix III.)

In summary, we have just shown that while our class of 5D metrics bears some resemblance to the metric of the 4D

spinning string, the class is free from the  $r \rightarrow 0$  pathology of the spinning string, does not violate the standard energy conditions as does Gott's moving string pair, and does not present particle blueshifts (energy gains) as does Gott's metric.

## VI. STROBOSCOPIC WORLD LINES FOR HIGGS SINGLET

The braneworld model which we have adopted has SM gauge particles trapped on our  $(3 + 1)$ -dimensional brane, but gauge-singlet particles are free to roam the bulk as well as the brane. We are interested in possible discovery of negative time travel at the LHC, sometimes advertised as a "Higgs factory." The time-traveling Higgs singlets can be produced either from the decay of SM Higgs or through mixing with the SM Higgs. We discuss Higgs singlet production in Sec. VIII.

### A. Higgs singlet pre-/reappearances on the brane

The physical paths of the Higgs singlets are the geodesics which we calculated in previous sections. The geodesic EOMs for the four spatial components of the five-velocity are trivially  $\ddot{\vec{r}} = 0$  and  $\ddot{u} = 0$ . Thus, the projection of the particle's position onto the brane coordinates is

$$\vec{r}(\tau) = \dot{\vec{r}}_0\tau + \vec{r}_0 = \dot{\vec{r}}_0 \frac{u}{\dot{u}_0} + \vec{r}_0 = \frac{v_0}{\beta_0} u \hat{p}_0 + \vec{r}_0. \quad (6.1)$$

Here,  $\hat{p}_0 = \widehat{\left(\frac{d\vec{r}}{d\tau}\right)}_0 = \widehat{\left(\frac{d\vec{r}}{dt}\right)}_0$  is the unit direction vector of the particle's three-momentum as seen by a brane observer,  $v_0$  is the initial speed of the particle along the brane direction<sup>8</sup>

$$v_0 \equiv \left| \left( \frac{d\vec{r}}{dt} \right)_0 \right|, \quad (6.2)$$

and  $\vec{r}_0$  is the point of origin for the Higgs singlet particle, i.e., the primary vertex of the LHC collision.

Of experimental interest is the reappearance of the particle on the brane. Inserting  $u = \pm NL$  into Eq. (6.1), one finds that the particle crosses the brane stroboscopically; the trajectory lies along a straight line on the brane, but piercing the brane at regular spatial intervals given by

$$\vec{r}_N = \frac{v_0}{|\beta_0|} NL \hat{p}_0 + \vec{r}_0. \quad (6.3)$$

We note the geometric relation  $\frac{v_0}{\beta_0} = \frac{|\dot{\vec{r}}_0|}{\dot{u}_0} = \cot\theta_0$ , where  $\theta_0$  is the exit angle of the particle trajectory relative to the brane direction.

The result in Eq. (6.3) for the spatial intervals on the brane holds for both co-rotating particles and counter-rotating particles. The distance between successive brane

<sup>8</sup>Recall that  $\dot{\vec{r}} = \frac{d\vec{r}}{d\tau}$  is a constant of the motion, but  $\vec{v} = \frac{d\vec{r}}{dt}$  is not, due to the nontrivial relationship between  $\tau$  and  $t$ . This nontrivial relationship between proper and coordinate times is what enables the CTC.



crossings,  $Lv_0/\beta_0$ , is governed by the size  $L$  of the compactified dimension. These discrete spatial intervals are likely too small to be discerned. However, the Higgs singlet is only observed when it scatters or decays to produce a final state of high-momenta SM particles. We expect the decay or scattering rate to be small, so that many bulk orbits are traversed before the Higgs singlet reveals itself. We discuss many-orbit trajectories next.

### B. Many-orbit trajectories and causality violations at the LHC

The coordinate times of the reappearances of the particle on the brane are given by Eq. (3.18) as

$$t_N = \pm \left( \frac{1}{\beta_0} - A \right) NL. \quad (6.4)$$

We have shown earlier that the assumption  $A > 0$ , without loss of generality, forces co-rotating particles having  $A > \frac{1}{\beta_0} > 1$  to travel in negative time. Thus, for the particles traveling on co-rotating geodesics (with  $\beta_0 > 0$  and positive signature), their reappearances are in fact preappearances. The time intervals for preappearances are

$$t_N(\text{co-rotating}) = - \left( A - \frac{1}{\beta_0} \right) NL < 0. \quad (6.5)$$

For counter-rotating particles (with  $\beta_0 < 0$  and negative signature), the time intervals are

$$t_N(\text{counter-rotating}) = \left( A + \left| \frac{1}{\beta_0} \right| \right) NL > 0. \quad (6.6)$$

The counter-rotating particles reappear on the brane at regular time intervals, but do not preappear. We note that even for co-rotating and counter-rotating particles with the same  $|\beta_0|$ , the magnitudes of their respective time intervals are different; the co-rotating interval is necessarily shorter. The mean travel times for co-rotating and counter-rotating particles, respectively, are

$$\langle t \rangle(\text{co-rotating}) = - \left( A \langle N \rangle - \left\langle \frac{N}{\beta_0} \right\rangle \right) L < 0 \quad (6.7)$$

and

$$\langle t \rangle(\text{counter-rotating}) = \left( A \langle N \rangle + \left\langle \frac{N}{|\beta_0|} \right\rangle \right) L > 0. \quad (6.8)$$

In Sec. VIII we will show that these means are very large numbers, inversely related to the decay/interaction probability of the Higgs singlet.

### C. Higgs singlet apparent velocities along the brane

We may compute the observable velocity connecting the production site for the Higgs singlet to the stroboscopic pre- and reappearances by dividing the particle's apparent travel distance along the brane,  $\vec{r}_N(t) - \vec{r}_0$  given in Eq. (6.3), by the apparent travel time  $t_N$  given in Eq. (6.4).

These are the velocities which an observer on the brane, e.g., an LHC experimenter, would infer from measurement. For the co-rotating particles with  $\beta_0 > 0$ , we get

$$\vec{v}(\text{co-rotating}) = - \frac{v_0}{\beta_0 A - 1} \hat{p}_0. \quad (6.9)$$

The velocities of the co-rotating particles are negative for  $\beta_0 A > 1$  because the particles are traveling backwards in time. For the counter-rotating particles with  $\beta_0 < 0$ , we get

$$\vec{v}(\text{counter-rotating}) = \frac{v_0}{|\beta_0| A + 1} \hat{p}_0. \quad (6.10)$$

The velocities of the counter-rotating particles are positive, as the particles travel forward in time, and subluminal. For a given exit angle  $\theta_0 = \cos^{-1}(v_0/\beta_0)$ , the speeds of the co-rotating particles, traveling backwards in time, generally exceed the speeds of the counter-rotating particles.

We note that the apparent speeds of co-rotating particles can be superluminal in either forward time ( $\beta_0 A < 1$ ) or backwards time ( $\beta_0 A > 1$ ). We display the velocities in Fig. 3 with a plot of  $v/v_0$  versus the parameter combination  $\beta_0 A$ . The particle speed diverges at  $\beta_0 A = 1$ ; the value  $\beta_0 A = 1$  corresponds to the slope of the light cone passing through zero, an inevitability discussed in Sec. IV.




FIG. 3 (color online). Apparent brane velocity  $v$  as a fraction of initial brane velocity  $v_0$  versus  $\beta_0 A$ . The counter-rotating particle always moves subluminally forward in time, but the co-rotating particle may move superluminally in either time direction. Brane velocities are divergent at  $\beta_0 A = 1$ , which occurs as the light cone crosses the horizontal axis of the spacetime diagram. For  $\beta_0 A > 1$ , the co-rotating geodesic is a CTC. The regions delineated by (a), (b), (c), (d), (e) map onto the world lines of Fig. 4 with the same labels.




FIG. 4 (color online). Shown are stroboscopic piercings (dots) of our brane by a returning Higgs singlet. World lines delineated by (a), (b), (c), (d), (e) correspond to regions of Fig. 3 with the same labels. In (a), the counter-rotating particle travels forward in brane/coordinate time, within the forward light cone. The co-rotating particle travels outside the brane’s forward light cone. In (b), the world line is superluminal but moving forward in brane time  $t$ . In (c), the world line is horizontal; the particle “moves” instantaneously in brane time. In (d) and (e), the particle travels superluminally and subluminally, respectively, backwards in brane time (signifying a CTC).

The region  $\beta_0 A > 1$  is the CTC region, of interest for this article.

In Fig. 4 we show schematically the world lines on our brane for co-rotating particles with negative transit times, and for counter-rotating particles with positive ( $\beta_0 A < 1$ ) and negative ( $\beta_0 A > 1$ ) transit times.

At some point along the world line, during one of the brane piercings, the Higgs singlet decays or interacts to produce a secondary vertex. As we show in the next paragraph, during each brane piercing, the particle’s three-momentum is just that missing from the primary vertex, i.e., three-momentum on the brane is conserved. The arrows in Fig. 4 are meant to denote the three-momentum missing from the primary vertex at the origin, and reappearing or preappearing in a displaced secondary vertex. The piercings (dots) of the brane have a  $\Delta r \sim$  the brane width  $w$ , and a  $\Delta t \sim w/v_0$ . We have seen in Sec. VC that a three-momentum (in fact, a five-momentum) on the brane is conserved for time-traveling particles. Therefore, the particle slopes  $\Delta t/\Delta r \sim 1/v_0$  give no indication whether the particles are negative-time travelers. Only the preappearance of the secondary vertex with respect to the primary vertex reveals their acausal nature. Importantly, the secondary and primary vertices are correlated by the conservation of a particle’s three-momentum: exactly the momentum missing from the primary vertex preappears in the secondary vertex.

## VII. 5D AND EFFECTIVE 4D FIELD THEORY FOR TIME-TRAVELING HIGGS SINGLETS

In previous sections, we have constructed a class of 5D metrics which admits stable CTC solutions of the classical Einstein equations, and we have presented the solutions. Similar to the ADD scenario, we will assume that all the SM particles are confined to the brane while gauge singlets, such as Higgs singlets, gravitons and sterile neutrinos, can propagate into the bulk. In this section, we first construct the 5D Lagrangian for the coupled Higgs singlet-doublet system. Then we derive the 5D equation of motion (5D Klein-Gordon equation) in our spacetime, and solve it subject to the compactified fifth-dimension boundary condition. From this exercise, there results the interesting energy-momentum dispersion relation. Next we integrate out the fifth dimension to obtain the effective 4D theory. Finally, we incorporate electroweak (EW) symmetry breaking to obtain the effective theory relevant for EW-scale physics.

### A. 5D Lagrangian for the coupled Higgs singlet-doublet system

A simple and economical model involves the Higgs singlet  $\phi$  coupling/mixing only with the SM Higgs doublet  $H$ . We add the following Lagrangian density  $\mathcal{L}^{(5D)} = \mathcal{L}_0 + \mathcal{L}_I$  to the SM:

$$\mathcal{L}_0 = \frac{G^{AB}}{2} \partial_A \phi \partial_B \phi - \frac{m^2}{2} \phi^2, \quad (7.1)$$

$$\mathcal{L}_I = -\frac{\lambda_1}{\sqrt{L}} \phi - \sqrt{L} \lambda_3 \phi H^\dagger H \delta(u) - L \lambda_4 \phi^2 H^\dagger H \delta(u), \quad (7.2)$$

where  $A, B = \{\mu, 5\}$  and  $G^{AB}$  is the 5D inverse metric tensor, with entries

$$G^{00} = h(u); \quad G^{05} = G^{50} = g(u); \quad G^{55} = G^{ii} = -1, \quad (7.3)$$

and all remaining entries zero. From the 5D kinetic term, one sees that the mass dimension of  $\phi$  is  $3/2$ . Constant factors of  $\sqrt{L}$  have been inserted for later convenience, so that the mass dimensions of  $\lambda_1$ ,  $\lambda_3$  and  $\lambda_4$  have the usual 4D values of 3, 1, and 0, respectively. The appearance of the delta function  $\delta(u)$  in  $\mathcal{L}^{(5D)}$  restricts the interactions with SM particles to the brane ( $u = 0$ ), to which the SM particles (here, the SM Higgs doublet  $H$ ) are confined. A consequence of the restriction of SM particles to the brane via the delta function is that translation invariance in the  $u$  direction is broken. This means that neither the Kaluza-Klein (KK) number nor particle momentum in the  $u$  direction is conserved; overall momentum conservation is restored when the recoil momentum of the brane is included.

### 1. Nonrenormalizable, effective field theory

The model in Eq. (7.2) is power-counting renormalizable. However, the broken translational invariance in the extra dimension(s) leaves the model nonrenormalizable. For example, the operators  $\phi^4$  (of dimension six and so manifestly not renormalizable in 5D) and  $\phi^3$  (5D-renormalizable, but destabilizing the Hamiltonian until large  $\phi$  values allow the  $\phi^4$  operator to dominate) are each induced on the brane by a virtual loop of the  $H$  field, and they are increasingly divergent as the number of extra dimensions increases. The fact that  $\phi^4$  is a nonrenormalizable operator, yet necessarily induced by the operator  $\phi^2 H^\dagger H \delta(u)$  which is power-counting renormalizable, is an indication that  $\mathcal{L}^{(5D)}$  describes an effective theory on the brane, not a renormalizable theory.

The induced operators  $\phi^4$  and  $\phi^3$  on the brane do not affect the physics of interest in this paper, and so we do not consider these operators any further. However, there are further effects of the effective theory that cannot be ignored. For example, higher-order Higgs-pair operators  $(H^\dagger H)^N$  are induced by a virtual loop of the  $\phi$  field. The  $N = 1$  and  $N = 2$  operators may be renormalized by the SM counterterms, but higher-order operators introduce divergences for which there are no counterterms. If the model were renormalizable, these higher-order operators would be finite and calculable. Instead, they are divergent, as we briefly illustrate in Appendix II. Consequently, the model is an effective theory, valid up to an energy cutoff of characteristic scale  $\Lambda \sim 1/(\lambda_4 L)$ .

Interestingly, the complications on the brane do not pervade the bulk where the  $H$  field vanishes. Since  $H = 0$  in the bulk, there are no  $H$  loops, and so no new induced operators. In the bulk,  $\phi$  is described by free field theory.

We note that since  $\phi$  is a gauge singlet, its mass  $m$  is unrelated to spontaneous symmetry breaking and is best thought of as a free parameter. We further note that the Higgs singlet is largely unconstrained by known physics. For example, gauge singlets do not contribute to the  $\rho \equiv (M_W/M_Z \cos\theta_w)^2$  parameter.

### 2. Scales of validity

In order to construct a wave packet smaller than the size  $L$  of the extra dimension, we require  $L \gg 1/\sqrt{s_{\text{LHC}}}$ . Combined with the fact that our effective theory is valid only up to the cutoff  $\Lambda \sim 1/(\lambda_4 L)$ , which we want to lie above  $\sqrt{s_{\text{LHC}}}$ , we arrive at a small bounded value for  $\lambda_4$ :

$$\lambda_4 < \frac{1}{L\sqrt{s_{\text{LHC}}}} \sim 10^{-2} \left( \frac{10^{-18} \text{ m}}{L} \right) \left( \frac{10 \text{ TeV}}{\sqrt{s_{\text{LHC}}}} \right) \ll 1. \quad (7.4)$$

For the LHC energy scale to probe the extra dimension, we must assume that the size of the extra dimensions  $L$  is  $\geq 1/\sqrt{s_{\text{LHC}}}$ . Since the LHC is designed to probe electro-weak symmetry, one may equivalently write  $L \geq 1/\text{TeV} \geq 10^{-19} \text{ m}$  for the LHC reach. It is useful at this

point to briefly review the bounds on the size of extra dimensions. The strongest constraint on the ADD scenario comes from limits on excess cooling of a supernova due to KK graviton emission [42] (analogous to limits from cooling by axion emission). One extra dimension is ruled out. For two extra dimensions, the lower bound on the fundamental Planck scale is 10 TeV and the upper bound on the size of the extra dimensions is  $L \lesssim 10^{-7} \text{ m}$  if the two extra dimensions are of the same size, easily within the reach of  $L \gtrsim 10^{-19} \text{ m}$  at the LHC. Consistency with the solar system tests of Newtonian gravity also requires at least two extra dimensions [43]. While we have shown that a single extra dimension is sufficient to admit our class of CTCs, our construction does not disallow further extra dimensions.

### B. Klein-Gordon solution and energy-momentum dispersion relation

To develop the field theory of the Higgs singlet, we will need the energy dispersion relation for the  $\phi$  particle modes. The dispersion relation can easily be obtained from the equation of motion for the free  $\phi$  field:

$$G^{AB} \partial_A \partial_B \phi + m^2 \phi = 0. \quad (5D \text{ Klein-Gordon equation}) \quad (7.5)$$

In fact, an inspection of Eq. (5.7) [and the definition of  $\tilde{t}$  in the footnote Eq. (5.4)] suggests that the general solution to this 5D Klein-Gordon (KG) equation for the  $n$ th energy eigenfunction should take the form

$$\phi_n^{(\text{KG})} = e^{-iE_n[t + \int_0^u g(u) du]} e^{i\vec{p} \cdot \vec{x}} e^{i\xi u}, \quad (7.6)$$

where  $E_n$  is the energy of the  $n$ th mode (at fixed  $\vec{p}$ ) and  $\vec{p}$  is the standard three-momentum along the brane direction. Since the extra dimension is compactified, we require  $\phi_n(u + L) = \phi_n(u)$  which, in turn, requires that

$$\xi = \bar{g} E_n + \frac{2\pi n}{L} \quad \text{with } n = 0, \pm 1, \pm 2, \dots, \quad (7.7)$$

where the mean value  $\bar{g}$  is defined in Eq. (3.14).<sup>9</sup> Thus, the solution to the KG equation is given by

$$\phi_n^{(\text{KG})} = e^{-iE_n t} e^{i\vec{p} \cdot \vec{x}} e^{-iE_n \int_0^u (g - \bar{g}) du} e^{inu/R}, \quad (7.8)$$

where we have defined an extra-dimensional ‘‘radius’’  $R \equiv L/2\pi$  to streamline some notation.

To determine the energy dispersion relation, we simply need to plug Eq. (7.8) into the 5D KG equation above and solve for  $E_n$ . A bit of algebra yields the quadratic dispersion relation

<sup>9</sup>From Eq. (2.3) we also get the mean  $\bar{h} = 1 - [\bar{g}^2 + \frac{1}{2} \sum_n a_n^2 + \frac{1}{2} \sum_n b_n^2]$ . We will not need this relation in the present paper.

$$(1 - \bar{g}^2)E_n^2 - 2\bar{g}E_n \frac{n}{R} - \bar{p}^2 - \frac{n^2}{R^2} - m^2 = 0. \quad (7.9)$$

Solving for  $E_n$  then gives,<sup>10,11</sup>

$$E_n = \frac{\bar{g} \frac{n}{R} + \sqrt{(1 - \bar{g}^2)(\bar{p}^2 + m^2) + \frac{n^2}{R^2}}}{1 - \bar{g}^2}. \quad (7.10)$$

Equation (7.10) makes it clear that the mode energy  $E_n$  depends on  $\bar{p}$  as well as on  $n$ ; nevertheless, for brevity of notation, we will continue to use the “fixed  $\bar{p}$ ” notation for both  $E_n$  and  $\phi_n$ . Here we have discarded the negative root which is unphysical, and retained the positive root which is physical for all  $n$ , positive, zero, and negative. In the above derivation, we have assumed that  $|\bar{g}| = |g_0 + A| < 1$ . This ensures that  $E_n$  is always real. Recall that we have chosen  $h_0 \geq 0$  to maintain the Minkowskian signature of the metric in Eq. (2.1) at  $u = 0$ . This in turn implies that  $|g_0| \leq 1$ . Therefore, the ultimate conditions on the metric which guarantee CTC solutions are simple, and three in number: (i)  $|g_0 + A| < 1$ , (ii)  $|g_0| \leq 1$ , and from Eqs. (3.19) and (3.20), that (iii)  $|A| > 1$ . As a result,  $g_0$  and  $A$  necessarily have opposite signs in the regions allowing CTCs. The two regions are given by (I)  $-1 \leq g_0 < 0$  and  $1 < A < 2$ , and (II)  $0 < g_0 \leq 1$  and  $-2 < A < -1$ . In Fig. 5, we display the two regions in the  $g_0$ - $A$  plane that allow CTCs.

The dispersion relation for  $E_n$  is interesting in several respects. First of all, due to the time independence of the metric  $G_{AB}$  and the time-translational invariance of the Lagrangian  $\mathcal{L}^{(5D)}$ , the energy  $E_n$  of the particle is constant during its propagation over the extra dimensional path (the bulk) which forms the CTC. In other words, the energy is covariantly conserved. Secondly, it is only for the zero mode [with  $n = 0$  and  $\bar{g}$  effectively zero in Eq. (7.10)] that the dispersion relation is trivial. The KK modes ( $n \neq 0$ ) exhibit a contribution  $n^2/R^2$  to the effective 4D mass squared, a complicated dependence on  $\bar{g}$ , and a resultant “energy offset”  $(\frac{\bar{g}}{1-\bar{g}^2})\frac{n}{R}$  which arises from the off-diagonal, nonstatic nature of the metric.

Not surprisingly, the integer mode number  $n$  has a quantum interpretation. It is the number of full cycles of  $\Phi_n^{(KG)}$  commensurate with the circumference  $L = 2\pi R$  of the extra dimension. We see this in the following way: The half-cycles of  $\Phi_n^{(KG)}$  are separated from  $u = 0$  by the distance  $u_k$ , where  $u_k$  is the solution to

<sup>10</sup>We are grateful to A. Tolley for correcting an error in an earlier version of our KG equation, and providing the dispersion relation which solves the corrected equation.

<sup>11</sup>Note that  $\bar{g}$  but not  $\bar{h} = 1 - \bar{g}^2$  appears in the KG solution and in the dispersion relation. This is related to the fact that a coordinate change may bring the metric to Minkowski form with no vestigial mention of  $h$  but with a pathologic “time” containing a boundary condition depending on  $\bar{g}$ . See Eqs. (5.6) and (5.7).




FIG. 5. The two regions in the  $g_0$ - $A$  plane for which CTCs are possible.

$$-E_n \int_0^{u_k} du(g - \bar{g}) + \frac{2\pi n u_k}{L} = k\pi, \quad k = 1, 2, \dots \quad (7.11)$$

Notice that the lengths of these half-cycles are not uniform. However, the total number of half-cycles is obtained by setting  $(u_k)_{\max} = L$ , for which Eq. (7.11) becomes simply  $2\pi n = k\pi$ . Thus,  $k_{\max} = 2n$ , and the number of full cycles is  $k_{\max}/2$ , which is  $n$ , identical to the number of wavelengths commensurate with  $L$  in the usual flat space [ $g(u) \equiv 0$ ] case. We conclude that a nonzero  $g(u)$  alters the lengths of the cycles in the extra dimension, but does not alter their total number, which is  $n$  for the mode  $\Phi_n^{(KG)}$ .

The sum on mode number plays the same role in the extra dimension that  $\int d^3 \vec{p}$  plays in 4D. Consequently, an arbitrary field in the 5D spacetime can be expanded as a linear combination of mode fields:

$$\phi(x^\mu, u) = \sum_{n=-\infty}^{\infty} \int d^3 \vec{p} \tilde{\phi}_n(E_n, \vec{p}) \phi_n^{(KG)}, \quad (7.12)$$

where  $\tilde{\phi}_n(E_n, \vec{p})$  are the weight functions of  $n$  and  $\vec{p}$ . This completes the construction of the scalar field in 5D with a periodic boundary condition in the fifth dimension.

### C. Wave packet as a sum of many modes

The minimum quantum energy of the  $n$ th mode is associated with motion purely in the compactified dimension. Thus we take the  $\vec{p} \rightarrow 0$  limit of the dispersion relation to determine this minimum quantized energy. From Eq. (7.10), we have

$$E_n(\vec{p} = 0) = \left(\frac{n}{R}\right) \left( \frac{\bar{g} + \sqrt{1 + (1 - \bar{g}^2) \frac{m^2 R^2}{n^2}}}{1 - \bar{g}^2} \right) \xrightarrow{z \equiv m^2 R^2 / n^2 \ll 1} \left(\frac{n}{R}\right) \left( \frac{1}{1 - \bar{g}} + \frac{z}{2} \right) + \mathcal{O}(z^2). \quad (7.13)$$

So for  $m^2 \ll n^2/R^2$ , we find an energy spectrum rising (nearly) linearly in  $n$ . This means that the first  $n_{\max} \sim \sqrt{5} \sqrt{L_{\text{HC}} R} (1 - \bar{g})$  modes are excitable, in principle. In practice, decay of the SM Higgs to  $\phi\phi$ , or mixing of  $H$  with  $\phi$ ,



will excite many KK modes of  $\phi$ . We have  $n_{\max} \sim M_h R \sim 6 \times 10^{10} (R/10^{-7} \text{ m})(M_h/125 \text{ GeV})$ . Thus, we do not expect single or few mode excitations to be relevant.

If a single mode were excited, its wave function would span the entire compactified interval  $[0, L]$ , analogous to a plane wave in 4D. With a single mode, one would expect quantum mechanics rather than classical concepts to apply. However, when more modes are excited, which we expect to be the relevant case, their weighted sum may form a localized wave packet in  $[0, L]$ , in which case the deductions for a classical particle in the earlier sections should apply. We denote the relevant many-mode wave packet by  $\phi_{\bar{n}}$ , where  $\bar{n}$  is meant to be a typical or mean mode number of the packet.

However, even an initially localized wave packet will spread in time. Such packet spreading does no harm to our conclusion—that the secondary vertex of the co-rotating Higgs singlet will still precede the production vertex in time. The spreading of the wave function just increases the variance of the distribution in negative  $t$ . The classical equation of motion for the  $u$  direction continues to describe the group velocity of the centroid of the wave function as it travels in the  $u$  direction. The same happens for the tau and  $b$  fermions in Minkowski space, as they progress from their production vertices to their decay vertices.

To understand the variance of the distribution of times between primary and secondary vertices, we now quantify the wave function spreading. To be explicit, we adopt a Gaussian wave packet at  $t = 0$  with initial spatial spread  $\Delta L_0$  in the fifth dimension. The standard formula for wave packet spread in a single dimension is  $(\Delta L)^2 = (\Delta L_0)^2 + (\tau/m\Delta L_0)^2$ . Here  $L$  is the circumference of the extra dimension as usual, and  $\tau$  is the proper time of the wave function (*a priori* independent of the time of the observer). There are two characteristic times of interest to us. The first is the time at which the packet begins to noticeably spread, given by  $\tau_1 \equiv m(\Delta L_0)^2$ . The second is the time when the packet completely fills the compactified dimension, given implicitly by  $\Delta L(\tau_2) = L$ , and explicitly by  $\tau_2 = m(\Delta L_0)L = (\frac{L}{\Delta L_0})\tau_1$ . For an experimental energy such that the  $n$ th mode is excitable, we have shown below Eq. (7.11) that there are  $n$  full cycles within the extra dimension. Each mode is a distorted plane wave filling the compact dimension, with an initial width of roughly  $L$ . Adding more modes decreases the width. We approximate the initial width of the Gaussian wave packet with  $n$  modes to be roughly  $\Delta L_0 = L/\bar{n}$ , where again,  $\bar{n}$  is the mean mode number. We further approximate  $m \sim 2\pi\bar{n}/L$ , and arrive at  $\tau_1 \sim 2\pi L/\bar{n}$ . Finally, taking  $\bar{n} \ll L\sqrt{s_{\text{LHC}}}$ , say,  $\sim (L\sqrt{s_{\text{LHC}}}/10) \sim L \text{ (TeV)} \sim L/10^{-19} \text{ m}$ , we find  $\tau_1 \sim 100/\sqrt{s_{\text{LHC}}} \sim 10^{-27} \text{ s}$  and  $\tau_2 \sim \bar{n}\tau_1 \sim (L/10^{-19} \text{ m}) \times 10^{-27} \text{ s}$ . Lab frame time  $t$  is related to time in the wave function frame by  $t = \gamma_u \tau$ , with  $\gamma_u$  being the Lorentz factor for a boost in the  $u$  direction. However,  $\gamma_u$  is nowhere near large enough to compensate for the many

orders of magnitude needed to qualitatively change the results just obtained for wave packet spreading. We conclude that the times  $t_1$  and  $t_2$  which characterize the wave function spreading in the lab frame are much shorter than the  $\geq$  picosecond time associated with displaced vertices. Consequently, the wave packet effectively spreads linearly in time with coefficient  $(m\Delta L_0)^{-1} \sim 1/2\pi$ , creating a considerable variance in the times (negative for co-rotating Higgs singlets and positive for counter-rotating Higgs singlets) between primary and secondary vertices.

We make here a side remark that in addition to the minimum energy associated with motion in the  $u$  direction, the momentum in the  $u$  direction is also interesting. While not observable, it is of sufficient mathematical interest that we devote Appendix III to its description.

We have seen that the localized time-traveling particle is a sum over many modes. The Lagrangian describing its production, which we now turn to, is also a sum over modes, with each mode characterized by an energy  $E_n$  according to our dispersion relation, Eq. (7.10). The weight functions in the Lagrangian are all unity. That is to say, calculations begin in the usual fashion, as perturbations about a free field theory.

#### D. 4D effective Lagrangian density

The reduction of the 5D theory to an effective 4D Lagrangian density is accomplished by the integration

$$\mathcal{L}^{(4D)} = \int_0^L du (\mathcal{L}_0 + \mathcal{L}_I), \quad (7.14)$$

where  $\mathcal{L}_0$  and  $\mathcal{L}_I$  are the 5D free and interacting Lagrangian densities given in Eqs. (7.1) and (7.2). We are interested in showing how the SM Higgs interacts with the singlet Higgs' energy eigenstates  $\phi_n(x^\mu, u)$ . Thus, the explicit expression for the Lagrangian density of the free singlet  $\int_0^L du \mathcal{L}_0$  is irrelevant for the following discussions.

Now we turn to the interaction terms. Neglecting the tadpole term  $-\frac{\lambda_1}{\sqrt{L}}\phi$  (which can be renormalized away, if desired), we have for the 4D interaction Lagrangian density

$$\begin{aligned} \int_0^L du \mathcal{L}_I &= -\sqrt{L}\lambda_3 H^\dagger H \int_0^L du \delta(u) \sum_n \phi_n(x^\mu, u) \\ &\quad - L\lambda_4 H^\dagger H \int_0^L du \delta(u) \sum_{n_1, n_2} \phi_{n_1}(x^\mu, u) \phi_{n_2}(x^\mu, u) \\ &= -\lambda_3 H^\dagger H \sum_n \sqrt{L} \phi_n(x^\mu, 0) \\ &\quad - \lambda_4 H^\dagger H \sum_{n_1, n_2} \sqrt{L} \phi_{n_1}(x^\mu, 0) \sqrt{L} \phi_{n_2}(x^\mu, 0) \\ &= -\lambda_3 H^\dagger H \sum_n \bar{\phi}_n - \lambda_4 H^\dagger H \sum_{n_1, n_2} \bar{\phi}_{n_1} \bar{\phi}_{n_2}, \quad (7.15) \end{aligned}$$

where  $\bar{\phi}_n = \sqrt{L}\phi_n(x^\mu, 0)$  is the singlet field on the brane, normalized with  $\sqrt{L}$  to its 4D canonical dimension of one.

Note that since the energy is covariantly conserved, both  $\phi_n(x^\mu, u)$  and  $\phi_n(x^\mu, 0)$  will have the same energy  $E_n$ .

### E. Incorporating electroweak symmetry breaking

Electroweak symmetry breaking (EWSB) in  $\mathcal{L}^{(4D)}$  is effected by the replacement  $H^\dagger H \rightarrow \frac{1}{2}(h + v)^2$  in Eq. (7.15), where  $v \sim 246$  GeV is the SM Higgs vacuum expectation value (VEV). The result is

$$\begin{aligned} \mathcal{L}^{(4D)} = & \int_0^L du \mathcal{L}_0 - \frac{\lambda_3}{2} (2vh + h^2) \sum_n \bar{\phi}_n \\ & - \frac{\lambda_4}{2} (v^2 + 2vh + h^2) \sum_{n_1, n_2} \bar{\phi}_{n_1} \bar{\phi}_{n_2}. \end{aligned} \quad (7.16)$$

Omitted from Eq. (7.16) is a new tadpole term  $-\frac{1}{2} \lambda_3 v^2 \sum_n \bar{\phi}_n$  linear in  $\bar{\phi}_n$ . It is irrelevant for the purposes of this article, so we here assume for simplicity that it can be eliminated by fine-tuning the corresponding counterterms.<sup>12</sup> The off-diagonal terms in  $\lambda_4 \sum_{n_1, n_2} v^2 \bar{\phi}_{n_1} \bar{\phi}_{n_2}$  mix different  $\bar{\phi}_n$  fields, while the terms  $\sum_n \lambda_3 v h \bar{\phi}_n$  induce singlet-doublet mixing. We make the simplifying assumption that  $(\lambda_3 v, \lambda_4 v^2) \ll m^2 + \frac{n^2}{R^2}$ , so that upon diagonalization of the mass matrix, the mass squared of the  $n$ th KK mode remains close to  $M_n^2 \approx m^2 + (\frac{n}{R})^2$ . We emphasize that this assumption is made so that the calculation may proceed to a more complete proof of principle for acausal signals at the LHC. In fact, it seems likely to us that acausal signals are inherent in the present model even without this simplifying assumption, and probably in other models not yet explored.

We now turn to the details of Higgs singlet production and detection at the LHC. As encapsulated in Eq. (7.16), Higgs singlets can be produced either from decay of the SM Higgs or through mass mixing with the SM Higgs. We discuss each possibility in turn.

## VIII. PHENOMENOLOGY OF PREAPPEARING SECONDARY VERTICES

Motivated by the advent of the LHC, we will next discuss the production and detection at the LHC of Higgs singlets which traverse through the extra dimension and violate causality.

How would one know that the Higgs singlets are crossing and recrossing our brane? The secondary vertex may arise from scattering of the singlet, or from decay (if allowed by symmetry) of the singlet. These ‘‘vertices from the future’’ would appear to occur at random times, uncorrelated with the pulse times of the accelerator.<sup>13</sup> The

<sup>12</sup>A different theory emerges if the counterterm is chosen to allow a nonzero tadpole term. For example, the singlet  $\phi$  field may then acquire a VEV.

<sup>13</sup>Preappearing events might well be discarded as ‘‘noise.’’ We want to caution against this expediency.

essential correlation is via momentum. Exactly the three-momentum missing from the primary vertex is restored in the secondary vertex. Of course, the singlet particles on counter-rotating geodesics will arrive back at our brane at later times rather than earlier times. The secondary vertices of counter-rotating particles will appear later than the primary vertices which produced them, comprising a standard ‘‘displaced vertex’’ event.

The rate of, distance to, and negative time stamp for the secondary vertices will depend on three parameters. First is the production rate of the Higgs doublets, which is not addressed in this paper. Secondly is the probability for production of the Higgs singlet per production of the Higgs doublet, which we denote as  $P_P$ . Thirdly is the probability for the Higgs singlet to interact, either by scattering or by decaying, to yield an observable secondary vertex in a detector. Of course, for the Higgs singlet to scatter on or decay to SM particles (via coupling with the SM Higgs doublet), the singlet must be on the brane. We define  $P_D$  to be the probability for the Higgs singlet to create a secondary vertex per brane crossing.

Since  $\phi$  is a singlet under all SM groups, it will travel almost inertly through the LHC detectors. Each produced singlet wave packet  $\bar{\phi}_{\bar{n}}$  exits the brane and propagates into the bulk, traverses the geodesic CTCs, and returns to cross the brane at times  $t_N$  given by Eq. (6.4). Classically, translational invariance in the  $u$  direction is broken by the existence of our brane, and so  $u$ -direction momentum may appear nonconserved.<sup>14</sup> The classical picture that emerges is the restoration of  $u$ -momentum conservation when brane recoil is included.

It is worth noting that all equations from the first six sections of this paper are classical equations, and so are independent of mode number  $n$ . Thus, these equations apply to the complete wave packet  $\bar{\phi}_{\bar{n}}$  formed from superposing many individual modes.

The probability for the Higgs singlet, once produced with probability  $P_P$ , to survive  $N$  traversals of the extra dimension and ‘‘then’’ decay or scatter on the  $(N + 1)$ th traversal is

$$P(N + 1) = P_D (1 - P_D)^N \sim P_D e^{-NP_D}. \quad (8.1)$$

The latter expression, of Poisson form, pertains for  $P_D \ll 1$ , as here. It is seen that even small scattering or decay probabilities per crossing exponentiate over many, many crossings to become significant. For this Poissonian probability, we have some standard results: the probability for interaction after  $N$  traversals is flat up to the mean value  $\langle N \rangle = 1/P_D$  (very large), the rms deviation,  $\sqrt{\langle N^2 \rangle - \langle N \rangle^2}$ ,

<sup>14</sup>Momentum in fact is conserved in the following sense: From the 5D point of view, energy-momentum is conserved as the brane recoils against the emitted Higgs singlet. From the 4D point of view, energy-momentum is conserved when the dispersion relation of Eq. (7.10) is introduced into the 4D phase space, as is done in Appendix IV.

is again  $1/P_D$  (very wide, as befits a flat distribution), and the probability for the singlet to interact in fewer crossings than  $\langle N \rangle = 1/P_D$  is  $1 - e^{-1} = 63\%$ .

Thus, the typical negative time between the occurrence of the primary vertex and the preappearance of the secondary vertex should be, from Eq. (6.4), of order

$$|t_{\langle N \rangle}| = \langle N \rangle |t_1| \sim \frac{L/c}{P_D}. \quad (8.2)$$

The typical range of the secondary vertex relative to the production site, from Eq. (6.3), is

$$r_{\langle N \rangle} = \frac{\cot\theta_0 L}{P_D}. \quad (8.3)$$

(Recall that  $\theta_0$  is the exit angle of the singlet relative to the brane direction.) The probability (per unit SM Higgs production) for the Higgs singlet to be produced and also interact within a distance  $l$  of the production site is then

$$P(\text{range of secondary vertex} < l) \\ \approx P_P \int_0^{\frac{l}{\cot\theta_0 L} P_D} d(N P_D) e^{-(N P_D)} = P_P \left[ 1 - e^{-(l \tan\theta_0 / L) P_D} \right], \quad (8.4)$$

which provides the limiting value

$$P(\text{range of secondary vertex} < l) \approx P_P P_D \frac{l \tan\theta_0}{L}, \\ \text{for } \frac{l \tan\theta_0}{L} P_D \ll 1. \quad (8.5)$$

For the secondary vertex to occur within the LHC detectors, one requires  $l = l_{\text{LHC}} \sim 10$  m.

We will assume that  $\langle \tan\theta_0 \rangle$  is of order unity. Then the figure of merit that emerges for CTC detection is  $P_P P_D l_{\text{LHC}} / L$ . We have seen that the maximum allowed value of  $L$  for two extra dimensions is  $10^{-7}$  m, and that the reach of the LHC is  $\sim 1/\sqrt{s} \sim 10^{-19}$  m. Thus, we are interested in an extra-dimensional size  $L$  within the bounds  $[10^{-19}, 10^{-7}]$  m. Below we shall see that the acausal preappearance of the secondary vertex for the co-rotating singlet may be observable at the LHC.

The Higgs singlet production and interaction mechanisms depend on the symmetry of the Higgs singlet-doublet interaction terms in the Lagrangian. Therefore the production and detection probabilities  $P_P$  and  $P_D$ , respectively, do as well. We discuss them next. There are two possibilities for our 5D Lagrangian, with and without a  $Z_2$  symmetry  $\phi \leftrightarrow -\phi$ .

### A. Without the $\phi \leftrightarrow -\phi$ symmetry

In this subsection, we ignore the possible  $Z_2$  symmetry and keep the trilinear term  $\lambda_3 H^\dagger H \sum_n \bar{\phi}_n$  in the Lagrangian density. When the SM Higgs acquires its VEV  $v$ , we have the resultant singlet-doublet mass-mixing term  $\lambda_3 v h \sum_n \bar{\phi}_n$  in the 4D Lagrangian of Eq. (7.16). Note that

this singlet-doublet mixing can only occur when the singlet particle  $\bar{\phi}_n$  is traversing the brane, as the field  $H$  is confined to the brane.

In Refs. [44,45], it was shown that mixing of the Higgs field with higher-dimensional graviscalars enhances the Higgs invisible width  $\Gamma_{\text{inv}}$  while maintaining the usual Breit-Wigner form. The invisible width is extracted from the imaginary part of the Higgs self-energy graphs, which includes the mixing of the Higgs with the many modes. These calculations apply in an analogous way to the Higgs-many mode mixing in our model. Thus the many-mode wave function  $\bar{\phi}_{\bar{n}}$ , which we introduced in Sec. VII C, is the Fourier transform of an energy-space Breit-Wigner form. In practice, this means that the sum includes all modes within the Higgs invisible width, sculpted by the Breit-Wigner shape. Including modes with energy from  $M_h - \Gamma_{\text{inv}}/2$  to  $M_h + \Gamma_{\text{inv}}/2$ , we have a mean mode number  $\bar{n} \sim M_h R$ , and an effective coupling for  $h$ - $\bar{\phi}_{\bar{n}}$  mixing of  $-m_{\text{mix}}^2 \sim -\lambda_3 v R \Gamma_{\text{inv}}$ , since  $1/R$  is roughly the energy spacing between modes. In the model of Ref. [44], the branching ratio  $\text{BR}(h \rightarrow \text{invisible})$  is calculated and shown to vary from nearly one with two extra dimensions, to three orders of magnitude less with six extra dimensions. We may expect something similar here.

Diagonalization of the effective  $2 \times 2$  mass-mixing matrix leads to the mixing angle  $\theta_{h\bar{\phi}_{\bar{n}}}$  between  $h$  and  $\bar{\phi}_{\bar{n}}$ . We assume this angle to be small, an assumption equivalent to assuming  $m_{\text{mix}}^2 \ll M_h^2$ . We label the resulting mass eigenstates of this  $2 \times 2$  subspace  $h_2$  and  $h_1$  with masses  $M_2$  and  $M_1$ . The mass eigenstates are related to the unmixed singlet and doublet states  $\bar{\phi}_{\bar{n}}$  and  $h$  by

$$\begin{pmatrix} |h_2\rangle \\ |h_1\rangle \end{pmatrix} = \begin{pmatrix} \cos\theta_{h\bar{\phi}_{\bar{n}}} & \sin\theta_{h\bar{\phi}_{\bar{n}}} \\ -\sin\theta_{h\bar{\phi}_{\bar{n}}} & \cos\theta_{h\bar{\phi}_{\bar{n}}} \end{pmatrix} \begin{pmatrix} |h\rangle \\ |\bar{\phi}_{\bar{n}}\rangle \end{pmatrix}, \quad (8.6)$$

and the inverse transformation is

$$\begin{pmatrix} |h\rangle \\ |\bar{\phi}_{\bar{n}}\rangle \end{pmatrix} = \begin{pmatrix} \cos\theta_{h\bar{\phi}_{\bar{n}}} & -\sin\theta_{h\bar{\phi}_{\bar{n}}} \\ \sin\theta_{h\bar{\phi}_{\bar{n}}} & \cos\theta_{h\bar{\phi}_{\bar{n}}} \end{pmatrix} \begin{pmatrix} |h_2\rangle \\ |h_1\rangle \end{pmatrix}. \quad (8.7)$$

We will assume for definiteness that on the brane, the two states (in either basis) quickly decohere due to a significant mass splitting. This assumption is reasonable since the decoherence time is  $t_{\text{deco}} \sim \frac{2\pi\gamma_\mu}{\delta M} \sim 10^{-23}$  (GeV/ $\delta M$ ) s. (The differing mass peaks  $M_2$  and  $M_1$  may thus be distinguishable at the LHC.) So we consider only the classical probabilities  $\cos^2\theta_{h\bar{\phi}_{\bar{n}}}$  and  $\sin^2\theta_{h\bar{\phi}_{\bar{n}}}$  in the remaining calculation.

The electroweak interaction, which would otherwise produce the SM Higgs, will now produce both mass eigenstates  $h_2$  and  $h_1$ , in the ratios of  $\cos^2\theta_{h\bar{\phi}_{\bar{n}}}$  and  $\sin^2\theta_{h\bar{\phi}_{\bar{n}}}$ , times phase space factors. For purposes of illustration, we take these phase space factors to be the same for both modes. The  $\bar{\phi}_{\bar{n}}$  components of these mass eigenstates  $h_2$  and  $h_1$  are given by the probabilities  $\sin^2\theta_{h\bar{\phi}_{\bar{n}}}$  and  $\cos^2\theta_{h\bar{\phi}_{\bar{n}}}$ , respectively. Thus, per production of a Higgs

doublet, the probability that a singlet  $\bar{\phi}_{\bar{n}}$  is produced is  $P_P = 2\sin^2\theta_{h\bar{\phi}_{\bar{n}}}\cos^2\theta_{h\bar{\phi}_{\bar{n}}} = \frac{1}{2}\sin^2(2\theta_{h\bar{\phi}_{\bar{n}}}) \rightarrow \theta_{h\bar{\phi}_{\bar{n}}} \ll 1 \sim 2\theta_{h\bar{\phi}_{\bar{n}}}^2$ .

Upon returning to the brane, these pure  $\bar{\phi}_{\bar{n}}$  states mix again and hence split into  $h_2$  and  $h_1$  states, with respective probabilities  $\sin^2\theta_{h\bar{\phi}_{\bar{n}}}$  and  $\cos^2\theta_{h\bar{\phi}_{\bar{n}}}$ . The probabilities for these  $h_2$  and  $h_1$  states to decay or interact as a SM Higgs  $h$  are respectively given by  $\cos^2\theta_{h\bar{\phi}_{\bar{n}}}$  and  $\sin^2\theta_{h\bar{\phi}_{\bar{n}}}$ . Thus, the total probability per returning  $\bar{\phi}_{\bar{n}}$  particle per brane crossing to decay or interact as a SM Higgs  $h$  is again  $P_D = P_P = \frac{1}{2}\sin^2(2\theta_{h\bar{\phi}_{\bar{n}}}) \sim 2\theta_{h\bar{\phi}_{\bar{n}}}^2$ , a very small number. Therefore, per initial Higgs doublet production the probability for a singlet  $\bar{\phi}_{\bar{n}}$  component to be produced and to acausally interact on the  $N$ th brane crossing is approximately  $4\theta_{h\bar{\phi}_{\bar{n}}}^4$ , nearly independent of the number of brane crossings. These brane crossings happen again and again until the interaction ends the odyssey. From the initial production of the  $\bar{\phi}_{\bar{n}}$  component to its final interaction upon brane crossing, the time elapsed (as measured by an observer on the brane) is again given by  $t_N$  in Eq. (6.4).

Therefore, in the broken  $\phi \leftrightarrow -\phi$  symmetry model, we expect the probability that a preappearing secondary vertex will accompany each SM Higgs event to be

$$P(\text{range of secondary vertex} < l_{\text{LHC}}) \approx P_P P_D (10 \text{ m}) / L \sim 10^8 (2\theta_{h\bar{\phi}_{\bar{n}}}^2)^2 \left( \frac{10^{-7} \text{ m}}{L} \right). \quad (8.8)$$

Here, the negative time between the secondary and primary vertices would be

$$|t_{\langle N \rangle}| \approx \frac{L}{P_D} \sim 3 \times 10^{-16} \left( \frac{L / 10^{-7} \text{ m}}{2\theta_{h\bar{\phi}_{\bar{n}}}^2} \right) \text{ s}. \quad (8.9)$$

Observability of a negative-time secondary vertex requires that  $|t_{\langle N \rangle}|$  lies in the interval of roughly 1 picosecond to 30 nanoseconds, and that the probability per Higgs doublet  $P$  (range of secondary vertex  $< l_{\text{LHC}}$ ) exceeds roughly one per million. Manipulation of Eqs. (8.8) and (8.9) then reveals that the two observability requirements are met with any  $L$  down to  $10^{-14}$  m (as discussed in Section VII B, we require  $L < 10^{-7}$  m to avoid excessive supernova cooling), and

$$10^{-7} \sqrt{\frac{L}{10^{-7} \text{ m}}} \leq 2\theta_{h\bar{\phi}_{\bar{n}}}^2 \leq \begin{cases} 10^{-3.5} \left( \frac{L}{10^{-7} \text{ m}} \right), & \text{if } L > 10^{-8} \text{ m;} \\ 10^{-4} \sqrt{\frac{L}{10^{-7} \text{ m}}}, & \text{if } L < 10^{-8} \text{ m.} \end{cases} \quad (8.10)$$

For example, with the largest value of  $L$  allowed by SN cooling rates for two extra dimensions,  $10^{-7}$  m, one gets  $10^{-7} \leq 2\theta_{h\bar{\phi}_{\bar{n}}}^2 \leq 10^{-3.5}$ . With the smallest value of  $L$

allowed for observability in the LHC detectors,  $10^{-14}$  m, one gets  $10^{-10.5} \leq 2\theta_{h\bar{\phi}_{\bar{n}}}^2 \leq 10^{-7.5}$ .

Thus, we have demonstrated that for a range of choices for  $L$  and  $\theta_{h\bar{\phi}_{\bar{n}}}$ , or equivalently, for  $P_P$  and  $P_D$ , preappearing secondary vertices are observable in the LHC detectors.

## B. With the $Z_2$ symmetry $\phi \leftrightarrow -\phi$

If one imposes the discrete  $Z_2$  symmetry  $\phi \leftrightarrow -\phi$ , then the coupling constants  $\lambda_1$  and  $\lambda_3$  are zero<sup>15</sup> and the low mode Higgs singlets are stable, natural, minimal candidates for weakly interacting massive particle (WIMP) dark matter [47,48]. Constraints on this model from the CDMS II experiment [49] have been studied in Ref. [50,51]. The discrete symmetry  $\phi \leftrightarrow -\phi$  also forbids the Higgs singlet to acquire a VEV. This precludes any mixing of the Higgs singlet with the SM Higgs. With the  $Z_2$  symmetry imposed, SM Higgs decay is the sole production mechanism of the Higgs singlet. The decay vertex of the SM Higgs provides the primary vertex for the production of the Higgs singlet, and subsequent scattering of the singlet via  $t$ -channel exchange of a SM Higgs provides the secondary vertex.

In Eq. (7.16), each term of the form  $\lambda_4 v h \bar{\phi}_{n_1} \bar{\phi}_{n_2}$  provides a decay channel for the SM Higgs into a pair of Higgs singlet modes, if kinematically allowed. The general case  $h \rightarrow \bar{\phi}_{n_1} \bar{\phi}_{n_2}$  with  $n_1 \neq n_2$  is considered in Appendix IV. Here we exhibit the simplest decay channels to single mode states,  $h \rightarrow \bar{\phi}_n \bar{\phi}_{-n}$  and  $h \rightarrow \bar{\phi}_n \bar{\phi}_n$ . The width for  $h \rightarrow \bar{\phi}_n \bar{\phi}_{-n}$  is

$$\Gamma_{h \rightarrow \bar{\phi}_n \bar{\phi}_{-n}} = \frac{\lambda_4^2 v^2}{16\pi M_h} \beta_{n,-n}; \quad (8.11)$$

$$\beta_{n,-n} = \frac{1}{\sqrt{1-\bar{g}^2}} \sqrt{1 - \frac{4\bar{M}_n^2}{(1-\bar{g}^2)^2 M_h^2}},$$

while the width for  $h \rightarrow \bar{\phi}_n \bar{\phi}_n$  is

$$\Gamma_{h \rightarrow \bar{\phi}_n \bar{\phi}_n} = \frac{\lambda_4^2 v^2}{8\pi M_h} \beta_{n,n}; \quad (8.12)$$

$$\beta_{n,n} = \frac{1}{\sqrt{1-\bar{g}^2}} \sqrt{1 - \frac{4\bar{M}_n^2}{((1-\bar{g}^2)M_h - \frac{2\bar{g}n}{R})^2}},$$

where  $\bar{M}_n^2 = (1-\bar{g}^2)m^2 + \frac{n^2}{R^2}$ .

The above formulas apply to single mode final states. Reference [45] looked at Higgs decay to a pair of graviscalars. The authors found via a quite complicated calculation that the decay was suppressed compared to simpler Higgs-graviscalar mixing. However, their model concerned gravitational coupling, whereas our model has completely different couplings for mixing. Thus, the

<sup>15</sup>The four-dimensional counterpart of this simple  $Z_2$  model was first proposed in Ref. [46], where the  $\phi$  quanta are called ‘‘scalar phantoms.’’



techniques of Ref. [45] may apply, but the conclusions do not. We choose to finesse the hard calculation with an order of magnitude estimate. Each sum on modes is constrained by phase space (and not by  $\Gamma_{\text{inv}}$  as in the broken- $Z_2$  mixing case), and so includes roughly  $n_{\text{max}} \sim M_h R/2 \sim 3 \times 10^{10} (R/10^{-7} \text{ m})(M_h/125 \text{ GeV})$  states. A typical mode value will be  $\bar{n} \sim M_h R/4$ . Thus, from here forward, in Eqs. (8.11) and (8.12), we set  $n$  to  $\bar{n}$  taken as  $M_h R/4$ , and multiply the rhs by the mode-counting factor  $n_{\text{max}}$  for each of the final state singlets, yielding the rate-enhancing factor  $n_{\text{max}}^2 \sim 2.3 \times 10^{19} (L/10^{-7} \text{ m})^2 \times (M_h/125 \text{ GeV})^2$ .

It is illuminating to look at the ratio of decay widths to  $\bar{\phi}_{\bar{n}}$  pairs and to  $\tau$ -lepton pairs. For the  $\tau$ , the coupling  $g_Y$  to the SM Higgs is related to the  $\tau$  mass through EWSB:  $g_Y^2 = 2m_\tau^2/v^2$ . Neglecting terms of order  $(m_\tau/M_h)^2$ , the ratio can be approximated as

$$\frac{\Gamma_{h \rightarrow \bar{\phi}_{\bar{n}} \phi_{\pm \bar{n}}}}{\Gamma_{h \rightarrow \tau^+ \tau^-}} \sim \frac{\lambda_4^2 v^4 n_{\text{max}}^2}{M_h^2 m_\tau^2} \beta_{\bar{n}, \pm \bar{n}}. \quad (8.13)$$

This ratio can be much greater than unity, even for perturbatively small  $\lambda_4$ , and so  $P_P$  can be nearly as large as unity. It thus appears likely that  $\bar{\phi}_{\bar{n}}$  particles will be copiously produced by SM Higgs decay if kinematically allowed, that their KK modes will explore extra dimensions if the latter exist, and finally, that the KK modes will traverse the geodesic CTCs, if nature chooses an appropriately warped metric.

The exact  $Z_2$  symmetry of the model under consideration forbids decay of the lighter  $\phi$  singlets. The  $Z_2$  model does allow communication of the  $\phi$  with SM matter through  $t$ -channel exchange of a SM Higgs. The top-loop induced coupling of the SM Higgs to two gluons provides the dominant coupling of the SM Higgs  $h$  to SM matter. Despite the small couplings of  $h$  to the SM, and  $\lambda_4 v$  at the  $h \bar{\phi}_{\bar{n}_1} \phi_{\bar{n}_2}$  vertex, singlet scattering is enhanced by  $n_{\text{max}}$  in amplitude, and so  $n_{\text{max}}^2$  in rate. Moreover, the singlet will eventually scatter since it will circulate through the periodic fifth dimension again and again until its geodesic is altered by the scattering event. The scattering cross section is of order

$$\sigma_{\phi_{\bar{n}} N} \sim \left\{ \frac{(\lambda_4 v n_{\text{max}}) \lambda(h \rightarrow t\bar{t} \rightarrow gg) (\alpha_s/4\pi)}{M_h^2} \right\}^2, \quad (8.14)$$

where  $\lambda(h \rightarrow t\bar{t} \rightarrow gg)$  is the effective coupling of  $h$  to the nucleon  $N$  through a virtual top loop at the Higgs end and two gluons at the nucleon end. This coupling strength is of order  $\frac{\alpha_s}{4\pi} \sim 10^{-2}$ . Thus, we expect

$$\sigma_{\phi_{\bar{n}} N} \sim \left[ (5 \times 10^9 \lambda_4) \left( \frac{125 \text{ GeV}}{M_h} \right) \left( \frac{L}{10^{-7} \text{ m}} \right) \right]^2 \text{ fb}. \quad (8.15)$$

We get the scattering probability per brane crossing by multiplying this cross section by the physical length of the

brane crossing  $\sim w$ , by the fraction of time spent on the brane  $\Delta t/t \sim w/L$ , and by the target density  $\rho$ ; the brane width  $w$  is a free parameter, beyond our classical model, but presumably of order  $\sim L$ . We find

$$P_D = 3 \times 10^{-20} \left( \frac{\sigma_{\phi_{\bar{n}} N}}{\text{fb}} \right) \left( \frac{\rho}{5 \text{ g/cm}^3} \right) \left( \frac{w}{L} \right)^2 \left( \frac{L}{10^{-7} \text{ m}} \right). \quad (8.16)$$

As a scaling law, we have  $P_D \propto L w^2 / M_h^2$ , which grows linearly in  $L$ .

In summary, with the  $\phi \leftrightarrow -\phi$  symmetry, we expect the probability that a preappearing secondary vertex will accompany each Higgs event at the LHC to be  $P_P P_D (10 \text{ m}) / L \sim 10^{-12} \left( \frac{\sigma_{\phi_{\bar{n}} N}}{\text{fb}} \right) \left( \frac{\rho}{5 \text{ g/cm}^3} \right) \left( \frac{w}{L} \right)^2$  for  $P_P \sim 1$ . The negative time between the secondary and primary vertices would be  $\sim L / P_D \sim 10^4 \left[ \left( \frac{\sigma_{\phi_{\bar{n}} N}}{\text{fb}} \right) \left( \frac{\rho}{5 \text{ g/cm}^3} \right) \left( \frac{w}{L} \right)^2 \right]^{-1} \text{ s}$ . These numbers for the unbroken  $\phi \leftrightarrow -\phi$  model are encouraging or discouraging, depending on nature's choice for the compactification length  $L$ . The model with broken  $\phi \leftrightarrow -\phi$  symmetry is more encouraging.

### C. Correlation of preappearing secondary and postappearing primary vertices

Finally, we summarize the correlations between the primary vertex producing the negative-time traveling Higgs singlet and the secondary vertex where the Higgs singlet reveals itself. As we have seen above, the first correlation is the small but possibly measurable negative time between the primary and secondary vertices.

The second correlation relating the preappearing secondary vertex and the postappearing primary vertex is the conserved momentum. As with familiar causal pairs of vertices, the total momentum is zero only for the sum of momenta in both vertices. Momentum conservation can be used to correlate the preappearing secondary vertex with its later primary vertex, as opposed to the background of possible correlations of the secondary vertex with earlier primary vertices.

Thus, the signature for the LHC is a secondary vertex preappearing in time relative to the associated primary vertex. The two vertices are correlated by total momentum conservation. If such a signature is seen, then a very important discovery is made. If such a signature is not seen, then the model is falsified for the energy scale of the LHC.

## IX. DISCUSSIONS AND FURTHER SPECULATIONS

As we have just demonstrated with a simple model, it is possible to have a significant amount of KK Higgs singlets produced by decay of, or mixing with, SM Higgs at the LHC. If nature chooses the appropriate extra-dimensional metric, then these KK Higgs singlets can

traverse the geodesic CTCs and thereby undergo travel in negative time.<sup>16</sup>

One may wonder why such acausal particles, if they exist, have not been detected up to now. One possible answer is that these time-traveling particles may have been recorded, but either unnoticed or abandoned as experimental background. Another possible answer could be that there has not been sufficient volume or instrumentation available to the detectors before now to detect these events. It may be that for the first time our scientific community has built accelerators capable of producing time-traveling particles, and also detectors capable of sensing them.

One may also wonder whether an acausal theory could be compatible with quantum field theory (QFT). After all, in the canonical picture, QFT is built upon time-ordered products of operators, and the path integral picture is built upon a time-ordered path. What does “time-ordering” mean in an acausal theory? And might the wave packet of a particle traversing a CTC interfere with itself upon its simultaneous emission and arrival? We note that each of these two questions has been discussed before, the first one long ago in Ref. [55], and the second one more recently in Ref. [56]. We offer no new insights into these questions. Rather, we have been careful to paint a mainly classical picture in this paper. We are content for now to let experiment be the arbiter of whether acausality is realizable in nature.

Finally, we would like to conclude with some speculations. In special relativity, space and time are unified. However, it seems that there is still an implicit difference between space and time. The reason is that traveling backwards in space appears to be easy, while traveling backwards in time requires a superluminal velocity. So the question arises: why is there an apparently inexorable arrow of time in our Universe? The issue of chronology protection may somehow be related to the very concept of time. Further theoretical investigations are badly needed.

While string theory [57] and loop quantum gravity [58] are formulated very differently, there is a common vision between them. Namely, a true theory of quantum gravity should be somehow background independent. This implies that spacetime is actually a derived concept and hence emergent [59]. In particular, the AdS/CFT correspondence [60] suggests that gravity is emergent. As observed in Ref. [61], the crucial point is that diffeomorphism

<sup>16</sup>The idea of causality violation at the LHC is not new. For example, a causality-violating SM Higgs has been proposed in Ref. [52], by invoking an unconventional complex action. The possibility of wormhole production at the LHC has been discussed in Ref. [53]. The idea of testing the vertex displacements for the acausal Lee-Wick particles at the LHC has been proposed by Ref. [54]. Also, some suggestive and qualitative effects associated with time-traveling particles have been proposed in Ref. [34], but without any concrete LHC signatures.

invariance simply characterizes the redundancies in the description of the gravity theory. But spacetime coordinates are themselves part of the redundant description in general relativity. Thus, the emergence of a unique gravity requires the emergence of spacetime as well. If the true quantum theory of gravity is indeed background independent and hence spacetime is emergent, then the idea of CTCs or time travel is completely meaningless at the energy scale of quantum gravity, since there is no spacetime at all. In this case, one can loosely say that chronology is “trivially protected” in that time is simply undefined. The discussion of chronology protection and time travel then become intimately related with the dynamics of how spacetime emerged.

If it turns out that the fundamental Planck scale is around a TeV as proposed by ADD, then the LHC would be at the right energy scale to elucidate our understanding of extra dimensions. If it further turns out that nature chooses an extra-dimensional metric which admits CTCs, then discovery of acausal correlations at the LHC would offer a fantastic new insight into the nature of spacetime.

### ACKNOWLEDGMENTS

We sincerely thank Alan Guth, Djordje Minic, Sandip Pakvasa, Tom Rizzo, Arkady Vainshtein, and especially Andrew Tolley and the anonymous referee for useful comments. This work was supported by U.S. DOE Grant No. DE-FG05-85ER40226.

### APPENDIX A: COVARIANT APPROACH TO LIGHT-CONE ANALYSIS

If a global time coordinate can be defined for a metric, then the metric cannot contain CTC solutions. A time function  $t$  is a global time coordinate if its four-gradient  $\partial_\mu t$  is everywhere timelike, i.e., if  $|\partial_\mu t|^2 > 0$  everywhere. This covariant condition for the absence or presence of a CTC has been brought to our attention by A. Guth. He references a proof of this theorem in Ch. 8 of Wald’s textbook [62]. Here we wish to show that for our simple metric, this condition reduces to the light-cone condition of Eq. (4.1).

The condition for the absence of any CTC is that

$$|\partial_\mu t|^2 = \partial_\mu t \partial_\nu t g^{\mu\nu} = s_\mu s_\nu g^{\mu\nu} > 0 \text{ everywhere, with} \\ s_\mu \equiv \partial_\mu t. \tag{A1}$$

Each of the four  $s_\mu$  is the slope of the particle’s world line in the  $\mu$  direction. (Note that  $s_\mu$  is *not* a covariant four-vector.) In this paper, we have chosen the time function to be the coordinate time  $t$ . In addition, we have time-translation invariance along the brane directions, but an off-diagonal metric element  $g_{tu}$  in the time-bulk plane. This off-diagonal element mixes  $t$  and  $u$ , leading to a nontrivial world line  $t(u)$  [see Eq. (3.17)]. Thus, for our metric, Eq. (A1) becomes

$$|\partial_\mu t|^2 = g^{tt} + 2s_u g^{tu} - s_u^2 g^{uu}. \quad (\text{A2})$$

Recalling that  $g^{\mu\nu}$  is the matrix inverse of the metric  $g_{\mu\nu}$ , one readily finds for the  $2 \times 2$   $t$ - $u$  subspace that  $g^{tt} = -g_{uu} = +h(u)$ ,  $g^{uu} = -g_{tt} = -1$ , and  $g^{tu} = +g_{tu} = g(u)$ . Substitution of these elements into Eq. (A2) then gives

$$|\partial_\mu t|^2 = -s_u^2 + 2g(u)s_u + h(u). \quad (\text{A3})$$

CTCs are allowed iff  $|\partial_\mu t|^2$  passes through zero. The quadratic form in Eq. (A3) may be written in terms of its roots  $s_\pm$  as  $-(s_u - s_-)(s_u - s_+)$ . Comparing the two quadratic forms then gives

$$s_+ + s_- = 2g_{tu}(u) \quad \text{and} \quad s_+ s_- = -h(u). \quad (\text{A4})$$

Thus, we are led via the covariant pathway to the massless-particle analog of our intuitive light-cone slope condition for CTCs, given in Eq. (4.2). [The sign of  $g(u)$  is inconsequential since it can be reversed by simply redefining  $u \rightarrow -u$ .]

### APPENDIX B: DIVERGENCE-INDUCED OPERATORS AND THE EFFECTIVE CUTOFF

In  $D$  dimensions, the  $N$ -pair Higgs operator is proportional to

$$(H^\dagger H)_{\text{operator}}^N \propto \prod_{\lambda=1}^N \lambda_4 \int d^D x \delta^{(D-4)}(\vec{x}_\perp) \Delta_F(x-y), \quad (\text{B1})$$

with  $\vec{x}_\perp$  being the coordinates orthogonal to the brane. The delta function in the integrand puts the operator on the brane where the  $H$  field is nonzero. The interconnected spacetime propagators  $D_F$  are

$$\Delta_F(x-y) = \int d^D k \frac{e^{ik \cdot (x-y)}}{(k^2 - m^2)}. \quad (\text{B2})$$

Spacetime integrations lead to  $N$  four-dimensional delta functions, each enforcing four-momentum conservation at one of the  $N$  vertices. Finally, these 4D delta functions may be integrated away to leave a single 4D delta function enforcing overall momentum conservation, times  $\lambda_4^N$  times the following schematic product of integrals:

$$\int \frac{d^D k}{(k^2 - m^2)} \prod_{j=1}^{N-1} \int \frac{d^{(D-4)} \vec{k}_{j\perp}}{(k_j^2 - m^2 + \dots)}, \quad (\text{B3})$$

where  $\vec{k}_\perp$  are the  $\phi$ -field momentum components orthogonal to the brane, and the first four (“brane”) components of the  $k_j$ ’s are fixed by the delta functions. This integral product diverges as the  $[4 + N(D-6)]$ th power of the cutoff  $\Lambda$ . For example, in five dimensions, the divergence is quadratic for  $N=2$  and logarithmic for  $N=3$  (odd powers of divergence are removed by the symmetric integration that follows a Wick rotation). In general, with more Higgs pairs or with more space dimensions, the divergence is worse. Consequently, the model is an effective theory,

valid up to an energy cutoff of characteristic scale  $\Lambda \sim 1/(\lambda_4 L)$ .

### APPENDIX C: MOMENTUM IN THE BULK DIRECTION

In this appendix, we wish to discuss the occurrence of conserved energy and brane three-momentum for the particle, and the nonconserved particle momentum in the bulk direction,  $u$ . Although the bulk momentum  $p_5$  (and associated  $p^5 = G^{5\beta} p_\beta$ ) is neither conserved nor observable, it is mathematically interesting in its own right.

The geodesic equation may be written as

$$\dot{\xi}^A = \frac{1}{2} (\partial_A G_{BC}) \xi^B \xi^C, \quad \text{where } \xi^A \equiv \dot{x}^A \quad (\text{C1})$$

is the tangent vector. In this form, the geodesic equation makes it clear that for each Killing vector  $\partial_A$  (i.e.,  $\partial_A$  such that  $\partial_A G_{BC} = 0$ ,  $\forall B, C$ ), there is a conserved quantity  $\xi_A = \dot{x}_A$ . Note that the conserved quantity carries a covariant (lower) index, rather than a contravariant (upper) index. For the 4D Minkowski metric, this is a moot point since upper and lower indices are simply related by  $\pm 1$ . However, for a more general metric, this point is crucial.

Since our metric depends only on  $u$ , it admits four conserved quantities. These are  $\dot{x}_0 = G_{0A} \dot{x}^A = \dot{t} + g(u)\dot{u}$ , and  $\dot{x}_j = G_{jA} \dot{x}^A = -\dot{x}^j = -p^j/m$ , i.e., the three-momentum  $\vec{p}$  on the brane is conserved. The conserved quantity  $\dot{t} + g(u)\dot{u}$  resulting from the Killing vector  $\partial_0$  must be proportional to the eigenvalue of the generator of time translation, i.e., the energy operator. We derived the energy eigenvalue  $E_n$  in the main text, and now we equate the two. Using the initial value for conserved  $\dot{t} + g(u)\dot{u}$ , we have

$$E_n = m(\gamma_0 + g_0 \dot{u}_0). \quad (\text{C2})$$

That the mass  $m$  is the proportionality constant is readily determined by taking the 4D limit of this equation, i.e., setting  $n$  and  $\vec{g}$  to zero in  $E_n$  on the lhs, and  $\dot{u}_0$  to zero on the rhs. In this paper, we do not exploit the relation (C2).

The momentum in the bulk direction is not conserved, owing to the breaking of the translational invariance in the  $u$  direction by the brane. Nevertheless, we may use the relation between momentum operator and generator for infinitesimal space translations to define it. The momentum then satisfies the standard Dirac commutator with its conjugate variable,  $x_5$ . The momentum operator expressed in position space then becomes  $P_5 = -i\partial_5$ . Operating on the  $n$ th-mode KG plane wave then determines its eigenvalue to be

$$p_5 = -E_n(g(u) - \vec{g}) + \frac{n}{R}. \quad (\text{C3})$$

It is obvious that  $p_5$  is nonconserved, because  $g(u)$  varies with  $u$  whereas all other terms in  $p_5$  are constants or conserved quantities. The fact that  $p_5$  depends on the

global element  $\bar{g}$  is an expression of “awareness” of the periodic boundary condition in the  $u$  direction. It then follows that  $E_n$  is also aware of the boundary condition through  $\bar{g}$ , because  $E_n$  depends on  $p_5$  (as well as on  $\bar{p}$ ).

We conclude this appendix by noting that the value of  $p_5$  averaged over a cycle in  $u$  is

$$\bar{p}_5 \equiv \frac{1}{L} \int_0^L du p_5 = \frac{n}{R}, \quad (\text{C4})$$

as might have been expected by one familiar with compactified extra dimensions having a diagonal (possibly warped) metric. An implication of this result is that any observable on the brane will depend on  $p_5$  only through a  $\frac{n}{R}$  term. Examples are the energy eigenvalues  $E_n$ , which are invariants and so have the same value on or off the brane. Thus, they may depend on  $p_5$  only through  $\frac{n}{R}$ , and they do.

#### APPENDIX D: SM HIGGS DECAY: $h \rightarrow \bar{\phi}_{n_1} \bar{\phi}_{n_2}$

In this appendix, we calculate the decay width of a SM Higgs doublet  $h$  of mass  $M_h$  to a pair of singlets  $\phi_{n_1}$  and  $\phi_{n_2}$ . Here it is assumed that the states  $h$  and  $\phi_n$  are not mixed. Such is the case if a  $\phi_n \leftrightarrow -\phi_n Z_2$  symmetry is imposed.

Care is needed to correctly incorporate the unusual energy dispersion formula  $E_n$  of Eq. (7.10), and the compactified nature of the extra dimension. The Lorentz invariant integral  $\int \frac{d^3 \vec{p}}{2E} = \int d^4 p \delta(p^2 - m^2) \Theta(p^0)$ , appropriate for flat Minkowski space, must be promoted to a covariant integral. In principle,  $d^4 p$  is made covariant by multiplying it with  $\sqrt{|\text{Det}(G_{AB})|}$ . However, decay of the SM Higgs occurs only on the brane, so it is the determinant

of the induced 4D metric that enters here, and the induced 4D metric is nothing but the familiar Minkowski metric with  $|\text{Det}| = 1$ . Thus  $d^4 p$  is invariant. In addition, the quadratic form in the delta function is the eigenvalue of the scalar Klein-Gordon operator  $G^{AB} \partial_A \partial_B + m^2$ , so it too is invariant. Thus, the correct, invariant phase space integral is

$$\begin{aligned} & \sum_n \int d^4 p \delta\left((1 - \bar{g}^2)p_0^2 - 2\bar{g}p_0 \frac{n}{R} - \bar{p}^2 - \frac{n^2}{R^2} - m^2\right) \theta(p_0) \\ &= \sum_n \int \frac{d^3 \vec{p}}{2[(1 - \bar{g}^2)E_n - \frac{\bar{g}n}{R}]} \\ &= \sum_n \int \frac{d^3 \vec{p}}{2\sqrt{(1 - \bar{g}^2)(\bar{p}^2 + m^2) + (\frac{n}{R})^2}}. \end{aligned} \quad (\text{D1})$$

Here the argument of the delta function is just the quadratic form of the dispersion relation (the generalization of the 4D Minkowski space dispersion relation  $E^2 = \vec{p}^2 + m^2$ ) given in Eq. (7.9), and the latter equality follows from Eq. (7.10). The periodic boundary condition in the  $u$  direction enters the dispersion relation through the mean metric element  $\bar{g}$ .

Our calculation below for the decay width  $\Gamma_{h \rightarrow \bar{\phi}_{n_1} \bar{\phi}_{n_2}}$  follows the treatment given in Sec. 4.5 of Ref. [63]. Translational invariance in time (for any time-independent metric) guarantees energy conservation, and translational invariance in space along the three brane directions guarantees three-momentum conservation. Thus, the tree-level decay width in the center-of-momentum frame is given by

$$\begin{aligned} \Gamma_{h \rightarrow \bar{\phi}_{n_1} \bar{\phi}_{n_2}} &= \frac{1}{2E_{\text{cm}}} \int \frac{d^3 \vec{p}_1}{(2\pi)^3 2[(1 - \bar{g}^2)E_{n_1} - \frac{\bar{g}n_1}{R}]} \int \frac{d^3 \vec{p}_2}{(2\pi)^3 2[(1 - \bar{g}^2)E_{n_2} - \frac{\bar{g}n_2}{R}]} \\ &\quad \times |\mathcal{M}(h \rightarrow \bar{\phi}_{n_1} \bar{\phi}_{n_2})|^2 (2\pi)^4 \delta^{(3)}(\vec{p}_1 + \vec{p}_2) \delta(E_{\text{cm}} - E_{n_1} - E_{n_2}) \\ &= \frac{\lambda_4^2 v^2}{2E_{\text{cm}}} \int \frac{d|\vec{p}_1| |\vec{p}_1|^2 d\Omega}{(2\pi)^3 2[(1 - \bar{g}^2)E_{n_1} - \frac{\bar{g}n_1}{R}] 2[(1 - \bar{g}^2)E_{n_2} - \frac{\bar{g}n_2}{R}]} (2\pi) \delta(E_{\text{cm}} - E_{n_1} - E_{n_2}) \\ &= \frac{\lambda_4^2 v^2}{2E_{\text{cm}}} \int d\Omega \frac{|\vec{p}|^2}{16\pi^2 [(1 - \bar{g}^2)E_{n_1} - \frac{\bar{g}n_1}{R}] [(1 - \bar{g}^2)E_{n_2} - \frac{\bar{g}n_2}{R}]} \left( \frac{|\vec{p}|}{(1 - \bar{g}^2)E_{n_1} - \frac{\bar{g}n_1}{R}} + \frac{|\vec{p}|}{(1 - \bar{g}^2)E_{n_2} - \frac{\bar{g}n_2}{R}} \right)^{-1} \\ &= \frac{\lambda_4^2 v^2}{8\pi E_{\text{cm}}} \left( \frac{|\vec{p}|}{(1 - \bar{g}^2)E_{n_1} - \frac{\bar{g}n_1}{R} + (1 - \bar{g}^2)E_{n_2} - \frac{\bar{g}n_2}{R}} \right). \end{aligned} \quad (\text{D2})$$

Using the condition  $E_{\text{cm}} = M_h = E_{n_1}(|\vec{p}|) + E_{n_2}(|\vec{p}|)$  and our dispersion relation, we arrive at

$$\Gamma_{h \rightarrow \bar{\phi}_{n_1} \bar{\phi}_{n_2}} = \frac{\lambda_4^2 v^2}{16\pi M_h} \beta_{n_1, n_2}, \quad (\text{D3})$$

where

$$\beta_{n_1, n_2} = \frac{\sqrt{((1 - \bar{g}^2)M_h - \frac{\bar{g}(n_1 + n_2)}{R})^4 - 2((1 - \bar{g}^2)M_h - \frac{\bar{g}(n_1 + n_2)}{R})^2 (\bar{M}_{n_1}^2 + \bar{M}_{n_2}^2) + (\bar{M}_{n_1}^2 - \bar{M}_{n_2}^2)^2}}{\sqrt{1 - \bar{g}^2} ((1 - \bar{g}^2)M_h - \frac{\bar{g}(n_1 + n_2)}{R})^2}} \quad (\text{D4})$$



and  $\bar{M}_{n_j}^2 \equiv (1 - \bar{g}^2)m^2 + (\frac{n_j^2}{R})$ . Perhaps a more familiar form for  $\beta_{n_1, n_2}$ , obtained by rearrangement of terms,<sup>17</sup> is

$$\beta_{n_1, n_2} = \frac{\sqrt{\lambda[(1 - \bar{g}^2)M_h - \frac{\bar{g}(n_1 + n_2)}{R}]^2, \bar{M}_{n_1}^2, \bar{M}_{n_2}^2}}{\sqrt{1 - \bar{g}^2}((1 - \bar{g}^2)M_h - \frac{\bar{g}(n_1 + n_2)}{R})^2}, \quad (\text{D5})$$

where  $\lambda(s, m_a^2, m_b^2) = (s - m_a^2 - m_b^2)^2 - 4m_a^2m_b^2$  is the usual triangle function employed in flat space calculations.

Notice that when the final-state particles are identical, there are two possible contractions in the amplitude rather than one, and a reduction of the two-body phase space from a sphere to a hemisphere to avoid double counting of identical particles. The net result is an extra factor of  $2^2 \times \frac{1}{2} = 2$ .

There is a subtlety associated with apparent momentum nonconservation in the  $u$  direction. The existence of the brane at  $u = 0$  breaks translational invariance in the  $u$  direction, and so we should not expect particle momentum in the  $u$  direction to be conserved. For  $q \equiv (n_1 + n_2)$ , the particle momentum leaving the brane in the  $u$  direction is  $q/R$ ,  $q \equiv (n_1 + n_2)$ . The ‘‘missing momentum’’  $-q/R$  is absorbed by the recoil of the brane. This is analogous to the apparent lack of conservation of  $z$  momentum when a child jumps upward from a surface, either rigid like the Earth’s surface or elastic like a trampoline’s surface. Presumably, a form factor  $|F(q)|^2$  which characterizes the dynamic response of the brane is included in  $|M|^2$  above, and arrives as a factor in Eq. (D3). Only for  $q = 0$ , i.e., for  $n_2 = -n_1$ , does the brane not enter the dynamics, so  $|F(0)|^2 = 1$ . In this work, we adopt the rigid picture of the brane, in which the net momentum of the exiting Higgs singlet pair,  $q/R$ , is so small compared to brane tension that  $|F(q)|^2 \approx |F(0)|^2 = 1$  for all  $q$ .

Finally, we state the obvious, that the total width of the SM Higgs to singlet Higgs pairs is

$$\Gamma_{h \rightarrow \phi\phi} = \sum_{n_1, n_2} \Gamma_{h \rightarrow \bar{\phi}_{n_1} \bar{\phi}_{n_2}}, \quad (\text{D6})$$

where  $\sum_{n_1, n_2}$  includes all pairs of modes which are kinematically allowed, i.e., all pairs of mode numbers for which the  $\lambda$  function in Eq. (D5) is positive.

## APPENDIX E: HIGGS SINGLET-DOUBLET MIXING

From Eq. (7.16), the contribution of  $\int du \mathcal{L}_I$  to the mass-squared matrix mixing the Higgs doublet and tower of singlet states is

<sup>17</sup>The equivalence of the argument of the square root in Eq. (D4) to the triangle function is easily seen by noting that the former is of the form  $A^2 - 2A(B + C) + (B - C)^2$ , which when expanded explicitly displays the symmetric form of the triangle function,  $\lambda(A, B, C) = A^2 + B^2 + C^2 - 2AB - 2BC - 2AC$ . A further feature of the triangle function, useful for extracting  $\bar{p}^2(E)$ , is that  $\lambda(A, B + \bar{p}^2, C + \bar{p}^2) = 0$  implies that  $\bar{p}^2 = \frac{\lambda(A, B, C)}{4A}$ .

$$\mathcal{M}^2 = \begin{pmatrix} M_h^2 & \lambda_3 v & \lambda_3 v & \lambda_3 v & \cdots \\ \lambda_3 v & M_0^2 + \lambda_4 v^2 & \lambda_4 v^2 & \lambda_4 v^2 & \cdots \\ \lambda_3 v & \lambda_4 v^2 & M_1^2 + \lambda_4 v^2 & \lambda_4 v^2 & \cdots \\ \vdots & \vdots & \lambda_4 v^2 & \ddots & \lambda_4 v^2 \end{pmatrix}. \quad (\text{E1})$$

(We do not consider here the mixing contribution from  $\int du \mathcal{L}_0$ .) Subtracting  $\lambda \mathbb{1}$  from this matrix and taking the determinant then gives the secular equation for the mass-squared eigenvalues  $\lambda$ . We may use Schur’s determinant equation to simplify the calculation. For a matrix of the form

$$\mathbb{M} = \begin{pmatrix} \mathbb{A}_{p \times p} & \mathbb{B}_{p \times q} \\ \mathbb{C}_{q \times p} & \mathbb{D}_{q \times q} \end{pmatrix}, \quad (\text{E2})$$

the determinant of  $\mathbb{M}$  is given by  $\text{Det}(\mathbb{M}) = \text{Det}(\mathbb{D})\text{Det}(\mathbb{A} - \mathbb{B}\mathbb{D}^{-1}\mathbb{C})$ . We choose  $\mathbb{A}$  to be the first entry in the upper left corner, and work in zeroth order of  $\lambda_4 v^2$ . Schur’s form then implies that

$$\begin{aligned} 0 &= \text{Det}(\mathcal{M}^2 - \lambda \mathbb{1}) \\ &= \left[ \prod_{n=0}^{\infty} (M_n^2 - \lambda) \right] \left[ (M_h^2 - \lambda) - (\lambda_3 v)^2 \sum_{n=0}^{\infty} \frac{1}{(M_n^2 - \lambda)} \right]. \end{aligned} \quad (\text{E3})$$

If we are interested in the mixing of the Higgs doublet with a particular singlet mode  $\bar{\phi}_n$ , we may organize the secular equation as

$$\begin{aligned} 0 &= [(M_h^2 - \lambda)(M_n^2 - \lambda) - (\lambda_3 v)^2] \\ &\quad - (\lambda_3 v)^2 \left[ \sum_{q \neq n}^{\infty} \frac{(M_n^2 - \lambda)}{(M_q^2 - \lambda)} \right]. \end{aligned} \quad (\text{E4})$$

For small enough values of  $\lambda_3$ , one may argue that the mass-squared eigenvalue  $\lambda_n$  for the perturbed state  $\bar{\phi}_n$  remains sufficiently close to  $M_n^2$  that  $[\sum_{q \neq n}^{\infty} \frac{(M_n^2 - \lambda)}{(M_q^2 - \lambda)}]$  may be neglected. For this case, the mixing angle between states  $h$  and  $\bar{\phi}_n$  becomes  $\tan 2\theta_{h\bar{\phi}_n} \sim \frac{2\lambda_3 v}{|M_n^2 - M_h^2|}$ . For larger values of  $\lambda_3$ , or large values of  $\lambda_4$ , more care would be needed. To quantify these remarks, we first solve the piece of the secular equation in the first bracket of (E4) to get

$$\begin{aligned} \lambda_n &= \frac{1}{2} \left[ M_n^2 + M_h^2 + \sqrt{(M_n^2 - M_h^2)^2 + (2\lambda_3 v)^2} \right] \\ &\xrightarrow{\lambda_3 v \ll (M_n^2 - M_h^2)} M_n^2 + \frac{(\lambda_3 v)^2}{(M_n^2 - M_h^2)}. \end{aligned} \quad (\text{E5})$$

Then, we insert this perturbative result back into Eq. (E4) to evaluate the residual given by the second bracket. The result is

$$\text{residual} = (\lambda_3 v)^4 \sum_{q \neq n} \left[ \left( \frac{q^2}{R^2} - \frac{n^2}{R^2} \right) (M_n^2 - M_h^2) - (\lambda_3 v)^2 \right]^{-1}. \quad (\text{E6})$$

Thus, for  $\lambda_3 v \ll (M_n^2 - M_h^2)$ , the residual is a negligible order  $(\lambda_3 v)^4$  perturbation, and the results  $\lambda_n \approx M_n^2 + \frac{(\lambda_3 v)^2}{(M_n^2 - M_h^2)}$  and  $\tan 2\theta_{h\phi_n} \sim \frac{2\lambda_3 v}{M_n^2 - M_h^2}$  are robust.

- 
- [1] W. J. van Stockum, Proc. R. Soc. Edinburgh, Sect. A **57**, 135 (1937).
- [2] F. J. Tipler, Phys. Rev. D **9**, 2203 (1974).
- [3] K. Gödel, Rev. Mod. Phys. **21**, 447 (1949).
- [4] J. A. Wheeler, Phys. Rev. **97**, 511 (1955); Ann. Phys. (N.Y.) **2**, 604 (1957).
- [5] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Spacetime* (Cambridge University Press, New York, 1973).
- [6] M. S. Morris and K. S. Thorne, Am. J. Phys. **56**, 395 (1988); M. S. Morris, K. S. Thorne, and U. Yurtsever, Phys. Rev. Lett. **61**, 1446 (1988).
- [7] J. R. I. Gott, Phys. Rev. Lett. **66**, 1126 (1991).
- [8] M. Alcubierre, Classical Quantum Gravity **11**, L73 (1994); A. E. Everett, Phys. Rev. D **53**, 7365 (1996).
- [9] A. Ori, Phys. Rev. Lett. **95**, 021101 (2005).
- [10] O. Gron and S. Johannesen, Nuovo Cimento Soc. Ital. Fis. **125B**, 1215 (2011); New J. Phys. **10**, 103025 (2008).
- [11] M. Visser, *Lorentzian Wormholes: From Einstein to Hawking* (Springer, New York, 1996).
- [12] S. W. Hawking, Phys. Rev. D **46**, 603 (1992).
- [13] B. S. Kay, M. J. Radzikowski, and R. M. Wald, Commun. Math. Phys. **183**, 533 (1997).
- [14] E. K. Boyda, S. Ganguli, P. Horava, and U. Varadarajan, Phys. Rev. D **67**, 106003 (2003).
- [15] L. Dyson, J. High Energy Phys. **03** (2004) 024.
- [16] M. M. Caldarelli, D. Klemm, and P. J. Silva, Classical Quantum Gravity **22**, 3461 (2005).
- [17] C. V. Johnson and H. G. Svendsen, Phys. Rev. D **70**, 126011 (2004).
- [18] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B **429**, 263 (1998); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B **436**, 257 (1998); N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Rev. D **59**, 086004 (1999).
- [19] G. Kaelbermann, Int. J. Mod. Phys. A **15**, 3197 (2000).
- [20] H. Ishihara, Phys. Rev. Lett. **86**, 381 (2001).
- [21] R. R. Caldwell and D. Langlois, Phys. Lett. B **511**, 129 (2001).
- [22] H. Stoica, J. High Energy Phys. **07** (2002) 060.
- [23] E. Abdalla, A. G. Casali, and B. Cuadros-Melgar, Int. J. Theor. Phys. **43**, 801 (2004).
- [24] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999); **83**, 4690 (1999).
- [25] C. Csaki, J. Erlich, and C. Grojean, Nucl. Phys. B **604**, 312 (2001).
- [26] D. J. H. Chung and K. Freese, Phys. Rev. D **61**, 023511 (1999); **62**, 063513 (2000).
- [27] H. Pas, S. Pakvasa, J. Dent, and T. J. Weiler, Phys. Rev. D **80**, 044008 (2009).
- [28] I. D. Novikov, Phys. Rev. D **45**, 1989 (1992).
- [29] A. Carlini, V. P. Frolov, M. B. Mensky, I. D. Novikov, and H. H. Soleng, Int. J. Mod. Phys. D **04**, 557 (1995); **05**, 99(E) (1996).
- [30] F. Echeverria, G. Klinkhammer, and K. S. Thorne, Phys. Rev. D **44**, 1077 (1991).
- [31] J. Friedman, M. S. Morris, I. D. Novikov, F. Echeverria, G. Klinkhammer, K. S. Thorne, and U. Yurtsever, Phys. Rev. D **42**, 1915 (1990).
- [32] J. L. Friedman, N. J. Papastamatiou, and J. Z. Simon, Phys. Rev. D **46**, 4456 (1992).
- [33] D. G. Boulware, Phys. Rev. D **46**, 4421 (1992).
- [34] A. Mironov, A. Morozov, and T. N. Tomaras, Facta Univ. Ser. Phys. Chem. Technol. **4**, 381 (2006).
- [35] J. B. Hartle, Phys. Rev. D **49**, 6543 (1994).
- [36] S. Deser, R. Jackiw, and G. 't Hooft, Ann. Phys. (N.Y.) **152**, 220 (1984).
- [37] S. Deser and R. Jackiw, Comments Nucl. Part. Phys. **20**, 337 (1992).
- [38] S. Deser, R. Jackiw, and G. 't Hooft, Phys. Rev. Lett. **68**, 267 (1992).
- [39] G. M. Shore, Int. J. Mod. Phys. A **18**, 4169 (2003), and references therein.
- [40] S. M. Carroll, E. Farhi, A. H. Guth, and K. D. Olum, Phys. Rev. D **50**, 6190 (1994).
- [41] B. Shlaer and S. H. Tye, Phys. Rev. D **72**, 043532 (2005).
- [42] S. Cullen and M. Perelstein, Phys. Rev. Lett. **83**, 268 (1999); R. Franceschini, G. F. Giudice, P. P. Giardino, P. Lodone, and A. Strumia, J. High Energy Phys. **05** (2011) 092 (which provides less stringent constraints derived directly from new LHC data).
- [43] G. D. Kribs, arXiv:hep-ph/0605325.
- [44] G. F. Giudice, R. Rattazzi, and J. D. Wells, Nucl. Phys. B **595**, 250 (2001).
- [45] D. Dominici and J. F. Gunion, Phys. Rev. D **80**, 115006 (2009).
- [46] V. Silveira and A. Zee, Phys. Lett. B **161**, 136 (1985).
- [47] C. P. Burgess, M. Pospelov, and T. ter Veldhuis, Nucl. Phys. B **619**, 709 (2001).
- [48] M. C. Bento, O. Bertolami, R. Rosenfeld, and L. Teodoro, Phys. Rev. D **62**, 041302 (2000).
- [49] Z. Ahmed *et al.* (The CDMS-II Collaboration), Science **327**, 1619 (2010).
- [50] M. Asano and R. Kitano, Phys. Rev. D **81**, 054506 (2010).

- [51] M. Farina, D. Pappadopulo, and A. Strumia, *Phys. Lett. B* **688**, 329 (2010).
- [52] H. B. Nielsen and M. Ninomiya, *Int. J. Mod. Phys. A* **23**, 919 (2008).
- [53] I. Y. Aref'eva and I. V. Volovich, *Int. J. Geom. Methods Mod. Phys.* **05**, 641 (2008).
- [54] E. Alvarez, L. Da Rold, C. Schat, and A. Szykman, *J. High Energy Phys.* **10** (2009) 023; B. Grinstein, D. O'Connell, and M. B. Wise, *Phys. Rev. D* **77**, 025012 (2008).
- [55] J. A. Wheeler and R. P. Feynman, *Rev. Mod. Phys.* **17**, 157 (1945); **21**, 425 (1949).
- [56] D. M. Greenberger and K. Svozil, *Quo Vadis Quantum Mechanics?*, edited by A. Elitzur, S. Dolev and N. Kolenda (Springer Verlag, Berlin, 2005), Ch. 4.
- [57] J. Polchinski, *String Theory Vol. 1: An Introduction to the Bosonic String* (Cambridge University Press, Cambridge, England, 1998); *String Theory Vol. 2: Superstring Theory and Beyond* (Cambridge University Press, Cambridge, England, 1998).
- [58] C. Rovelli, *Quantum Gravity* (Cambridge University Press, Cambridge, England, 2004).
- [59] N. Seiberg, [arXiv:hep-th/0601234](https://arxiv.org/abs/hep-th/0601234).
- [60] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
- [61] H. Elvang and J. Polchinski, [arXiv:hep-th/0209104](https://arxiv.org/abs/hep-th/0209104).
- [62] R. M. Wald, *General Relativity* (University Press, Chicago, 1984).
- [63] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Westview Press, Boulder, Colorado, 1995).