

# Koide's $Z_3$ -symmetric parametrization, quark masses, and mixings

Piotr Żenczykowski\*

*Division of Theoretical Physics, The Henryk Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, Radzikowskiego 152, 31-342 Kraków, Poland*

(Received 27 February 2013; published 16 April 2013)

The sets of charged-lepton ( $L$ ) and quark ( $D, U$ ) masses may be parametrized in a  $Z_3$ -symmetric language appropriate for the discussion of Koide's formula. Experiment suggests that at the low-energy scale the relevant phase parameters  $\delta_f$  take on possibly exact values of  $\delta_L = 3\delta_D/2 = 3\delta_U = 2/9$ . For  $k_f$  (the other parameter relevant for the pattern of masses), a similarly simple expression ( $k_L = 1$ ) is known for charged leptons only. Using the Fritzsch-Xing decomposition of quark-mixing matrices, we show that the suggested pattern of low-energy quark masses is consistent with an earlier conjecture that  $k_{D,U} \approx 1$  in the weak basis.

DOI: [10.1103/PhysRevD.87.077302](https://doi.org/10.1103/PhysRevD.87.077302)

PACS numbers: 12.15.Ff

## I. THE DOUBLY SPECIAL KOIDE'S PARAMETRIZATION

The appearance of three generations of leptons and quarks and the related issue of their masses started to baffle us three quarters of a century ago. Over the years the problem has become further complicated by the presence of intergeneration mixing, as revealed in weak interactions. Fortunately, various approximate regularities have been found in the observed pattern of particle masses and mixings. Among the many possible parametrizations of these regularities, there might be some whose simplicity could help us in deciphering physics beyond the Standard Model.

One of the most interesting of such regularities is an empirical relation between the charged-lepton masses discovered by Koide [1] (for a brief review see Ref. [2]):

$$\frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{1 + k_L^2}{3}, \quad (1)$$

with  $k_L$  equal exactly 1. When the experimental  $e$  and  $\mu$  masses (here taken from Ref. [3]) are inserted into Eq. (1), this relation predicts the tauon mass within one standard deviation from its observed value:

$$m_\tau(k_L = 1) = 1776.9689 \text{ MeV}, \quad (2)$$

$$m_\tau(\text{exp}) = 1776.82 \pm 0.16 \text{ MeV}. \quad (3)$$

Discussions of this success of Koide's formula (1) are naturally formulated in a  $Z_3$ -symmetric framework by parametrizing the masses of any three given fermions  $f_1, f_2, f_3$  in terms of three parameters  $M_f, k_f, \delta_f$  as [4,5]

$$\sqrt{m_{f_j}} = \sqrt{M_f} \left( 1 + \sqrt{2} k_f \cos \left( \frac{2\pi j}{3} + \delta_f \right) \right), \quad (4)$$

$(j = 1, 2, 3).$

This choice of parametrization of masses is particularly suited to Koide's formula as not only  $M_f$  but also  $\delta_f$  drops out of the rhs of Eq. (1).

Since  $\delta_f$  is free we may assume  $m_1 \leq m_2 \leq m_3$  without any loss of generality. From Eq. (4) one then gets a counterpart of Eq. (1), in which it is now  $k_L$  that drops out of the formula:

$$\frac{\sqrt{3}(\sqrt{m_\mu} - \sqrt{m_e})}{2\sqrt{m_\tau} - \sqrt{m_\mu} - \sqrt{m_e}} = \tan \delta_L. \quad (5)$$

From the experimental values of  $e, \mu$ , and  $\tau$  masses one finds

$$\delta_L = 0.2222324, \quad (6)$$

which, as observed by Brannen and Rosen [6,7], is extremely close to

$$\delta_L = 2/9. \quad (7)$$

Conversely, assuming  $\delta_L = 2/9$ , Eq. (5) predicts the value of the tauon mass in terms of experimental  $e$  and  $\mu$  masses. Such a prediction is obviously completely independent from that given by the original Koide's formula [Eq. (1)]. Yet, just as in the case of Koide's formula, the obtained number is within one standard deviation from the measured  $\tau$  mass:

$$m_\tau(\delta_L = 2/9) = 1776.9664 \text{ MeV}. \quad (8)$$

Assuming that the Koide and the Brannen-Rosen observations do not reflect mere coincidences, the  $Z_3$ -symmetric parametrization (4) should be rightly called "doubly special." A peculiar feature of this parametrization is that the simple numbers of 1 and  $2/9$  work well at the *low-energy* scale and not at some high-mass scale. For example, taking the values of charged-lepton masses at the mass scale of  $M_Z$ , the extracted values of  $k_L$  and  $\delta_L$  deviate from their "perfect" values of 1 and  $2/9$  by about 0.2% and 0.5%, respectively. Apparently, an explanation of the success of predictions (2) and (8) should not be sought at the

\*piotr.zenczykowski@ifj.edu.pl

high-mass scale of some grand unified theory (see, e.g., Ref. [8]).

## II. EXTENDING THE SCHEME TO THE QUARK SECTOR

If there is some physical reason behind the appearance of simple numbers such as 1 (Koide) and  $2/9$  (Brannen-Rosen) in the charged-lepton sector, one would expect its analogs working in the quark and neutrino sectors as well. However, it is known that the original Koide formula (1) does not work when replacing the charged-lepton masses with those of neutrinos or quarks. For neutrinos one estimates directly from experiment that  $k_\nu \leq 0.81$  [9] (the mathematically allowed region being  $0 \leq k_f \leq \sqrt{2}$ ). For quarks, using their mass values appropriate at  $\mu = 2$  GeV, one obtains  $k_D \approx 1.08$  ( $k_U \approx 1.25$ ) for the down (up) quarks, respectively, [9,10]. If a higher energy scale  $\mu = M_Z$  is taken, even larger values are obtained, i.e.,  $k_D = 1.12$  and  $k_U = 1.29$ . Going from  $\mu = 2$  GeV towards the low-energy scale leads to smaller values of  $k_D$  and  $k_U$ . However, the top quark mass is so large that one certainly cannot bring  $k_U$  into the vicinity of 1.

On the other hand, it has been observed recently [11] from the quark sector analogs of Eq. (5) that for the old-fashioned effective quark masses, i.e., at the *low-energy* scale, the relevant phase parameters acquire approximate values:

$$\delta_U \approx 2/27 = \delta_L/3, \quad \delta_D \approx 4/27 = 2\delta_L/3, \quad (9)$$

which constitute a generalization of the Brannen-Rosen observation (7) to the quark sector. Due to the problem of quark confinement we obviously cannot check how precise the above equalities are. However, given the accuracy of their lepton counterpart [Eq. (8)] we may expect that they are fairly exact. Therefore, we will assume this from now on and *conjecture* that at the low-energy scale the charged-lepton and quark masses are characterized by Eqs. (7) and (9).

The problems then remain how to interpret the value of  $k_L = 1$  and how it should be generalized to the quark sector. Here we accept the suggestion of Ref. [12] that  $k_f = 1$  is a feature of the weak basis. Thus, according to Ref. [12], the masses of charged leptons are described by  $k_L = 1$  because for charged leptons the mass and the weak bases coincide. The lepton-mixing matrix (for simplicity we assume here Dirac neutrino masses), i.e.,

$$V_{\text{MNS}} = U_L^\dagger U_\nu, \quad (10)$$

is then wholly assigned to the contribution from neutrinos (for which  $k_\nu \neq 1$ ):

$$U_\nu = V_{\text{MNS}}, \quad U_L = 1. \quad (11)$$

In other words, if  $U_L$  were different from 1, one would not expect the simplicity of Koide's formula to persist.

Since the analogs of the charged-lepton equality  $k_L = 1$  do not hold in the  $U$  and  $D$  quark sectors, one expects that the quark counterpart of the decomposition (11) will be also modified. Thus, in the Cabibbo-Kobayashi-Maskawa (CKM) matrix

$$V_{\text{CKM}} = U_U^\dagger U_D, \quad (12)$$

both of the factor matrices  $U_U$  and  $U_D$  are expected to be different from 1.

For fermion type  $f$ , the general connection between the mass and the weak bases is

$$\text{diag}(m_{f_1}, m_{f_2}, m_{f_3}) = U_{f,\text{left}}^\dagger M U_{f,\text{right}}, \quad (13)$$

where  $M$  is the mass matrix in the weak basis and  $U_{f,\text{left}} \equiv U_f$ . Reference [12] chooses  $U_{f,\text{right}} = 1$ . The reason for this choice is obvious: one wants to study the effects of a change of basis in the sector of left-handed fields only. As a result, in the weak basis one deals with “pseudomasses”  $\tilde{m}_{f_j}$ , defined as [12]

$$\tilde{m}_{f_j} = \left| \sum_k U_f^{jk} m_{f_k} \right|. \quad (14)$$

The appearance of a sum over  $k$  amounts here to an assumption that those mass terms which are coefficients at the symmetric (“democratic”) combination

$$\tilde{q}_{f,\text{right}} = \frac{1}{\sqrt{3}} \left( \sum_{k=1}^3 q_{f_k,\text{right}} \right) \quad (15)$$

are taken into account in the definition of pseudomasses, but that the analogous coefficients at the remaining two combinations of  $q_{f_1,\text{right}}$ ,  $q_{f_2,\text{right}}$ , and  $q_{f_3,\text{right}}$  (orthogonal to  $\tilde{q}_{f,\text{right}}$ ) are not considered (for example, they might physically vanish). Since Koide's formula involves square roots of masses, which is a feature of unknown origin, a natural generalization of this formula to pseudomasses seems to be that these masses should be defined as moduli of  $\sum_k U_f^{jk} m_{f_k}$  as proposed in Ref. [12].<sup>1</sup>

For charged leptons ( $f = L$ ,  $U_L = 1$ ) these pseudomasses coincide with the observed masses, i.e.,  $\tilde{m}_{L_j} = |m_{L_j}|$ . On the other hand, for quarks ( $f = D, U$  with  $U_D, U_U \neq 1$ ) the pseudomasses are different from the observed mass values. It is for these pseudomasses that, according to the proposal of Ref. [12], the analogs of Koide's formula (1) are supposed to hold with  $k_D = k_U = 1$ .

The authors of Ref. [12] used quark masses at the  $Z$  mass scale and found that it is possible to get  $k_D = k_U \approx 1$  provided one takes a value of the strange quark mass  $m_s(Z)$  that is larger by a factor of 2.5 from the theoretical estimate at that scale. Since the Koide and the

<sup>1</sup>There is no freedom to redefine phases so as to obtain real positive values for  $\sum_k U_f^{jk} m_{f_k}$ . The whole available freedom in the choice of phases is used up when removing five unphysical phases from the general CKM matrix; see Sec. B in Ref. [13].

Brannen-Rosen observations work best at the *low-energy* scale (and not at the  $Z$  mass), and since at the low-energy scale the masses of quarks (and especially  $m_s$ ; see Ref. [11]) are naturally expected to be larger than at the  $Z$  mass, a question appears if it is possible to recover Koide's formula for quarks using effective quark masses corresponding to phases of formulas (9). This is the question asked here. Thus, the present paper constitutes a low-energy-scale study of the idea of Ref. [12].

### III. THE STRUCTURE OF $U_f$ IN THE QUARK SECTOR

As the assumption of Koide's formula for pseudomasses imposes constraints upon matrices  $U_D$  and  $U_U$  (and, consequently, upon  $V_{\text{CKM}}$ ), we have to discuss these matrices in some detail.

In Ref. [14] Fritzsch and Xing convincingly argued that the hierarchical structure of quark mass terms suggests certain particular parametrizations of  $U_D$  and  $U_U$  as "most physical" (i.e., that this structure selects one of the nine possible parametrizations of the  $V_{\text{CKM}}$  matrix [15] as probably the most suitable for the description of the quark-mixing phenomenon). The same parametrization was also advocated in Ref. [12] where the conjecture that Koide's formula holds in the weak basis was originally formulated. Consequently, we think that it is justified to accept the Fritzsch-Xing parametrization here. The relevant "natural" parametrizations of  $U_D$  and  $U_U$  are then

$$U_D = R_{23}(\phi_b, \theta_b)R_{12}(\theta_d), \quad U_U = R_{23}(\phi_t, \theta_t)R_{12}(\theta_u), \quad (16)$$

with ( $c_q \equiv \cos \theta_q$ ,  $s_q \equiv \sin \theta_q$ )

$$R_{12}(\theta_q) = \begin{pmatrix} c_q & -s_q & 0 \\ s_q & c_q & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (q = d, u), \quad (17)$$

$$R_{23}(\phi_q, \theta_q) = \begin{pmatrix} e^{-i\phi_q} & 0 & 0 \\ 0 & c_q & s_q \\ 0 & -s_q & c_q \end{pmatrix}, \quad (q = b, t). \quad (18)$$

Thus, the induced parametrization of the  $V_{\text{CKM}}$  matrix can be read from

$$V_{\text{CKM}} = R_{12}^\dagger(\theta_u)R_{23}^\dagger(\phi_t, \theta_t)R_{23}(\phi_b, \theta_b)R_{12}(\theta_d). \quad (19)$$

The product  $R_{23}^\dagger(\phi_t, \theta_t)R_{23}(\phi_b, \theta_b)$  may be written in the form of Eq. (18) with a single phase  $\phi = \phi_b - \phi_t$  and a single rotation angle  $\theta = \theta_b - \theta_t$ , (with  $c \equiv \cos \theta$ ,  $s \equiv \sin \theta$ )

$$R_{23}^\dagger(\phi_t, \theta_t)R_{23}(\phi_b, \theta_b) \equiv R_{23}(\phi, \theta) = \begin{pmatrix} e^{-i\phi} & 0 & 0 \\ 0 & c & s \\ 0 & -s & c \end{pmatrix}. \quad (20)$$

The CKM matrix is then parametrized as

$$V_{\text{CKM}} = \begin{pmatrix} c_u & s_u & 0 \\ -s_u & c_u & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{-i\phi} & 0 & 0 \\ 0 & c & s \\ 0 & -s & c \end{pmatrix} \begin{pmatrix} c_d & -s_d & 0 \\ s_d & c_d & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ = \begin{pmatrix} s_u s_d c + c_u c_d e^{-i\phi} & s_u c_d c - c_u s_d e^{-i\phi} & s_u s \\ c_u s_d c - s_u c_d e^{-i\phi} & c_u c_d c + s_u s_d e^{-i\phi} & c_u s \\ -s_d s & -c_d s & c \end{pmatrix}. \quad (21)$$

Since quark fields can be freely rephased, all of the three angles  $\theta_d$ ,  $\theta_u$ ,  $\theta$  can be arranged to lie in the first quadrant so that  $s_u$ ,  $s_d$ ,  $s$  and  $c_u$ ,  $c_d$ ,  $c$  are all positive (the phase  $\phi$  cannot be so restricted).

We use the following absolute values of the elements of the CKM matrix most relevant for our parametrization<sup>2</sup> [3]:

$$\begin{aligned} |V_{ub}| &= 0.00351 \pm 0.00015, \\ |V_{cb}| &= 0.0412 \pm 0.0008, \\ |V_{td}| &= 0.00867 \pm 0.00030, \\ |V_{ts}| &= 0.0404 \pm 0.0008, \\ |V_{tb}| &= 0.999146 \pm 0.000034. \end{aligned} \quad (22)$$

Inserting the above numbers, we find from (21)

$$\begin{aligned} \theta_u &= 4.87^\circ \pm 0.23^\circ, & \theta_d &= 12.11^\circ \pm 0.47^\circ, \\ \theta &= \theta_b - \theta_t = 2.37^\circ \pm 0.05^\circ. \end{aligned} \quad (23)$$

Since formula (14) involves absolute values, the actual sizes of  $\phi_b$  and  $\phi_t$  and, consequently, the experimentally imposed restriction on the  $CP$ -violating phase parameter  $\phi = \phi_b - \phi_t$  are irrelevant for our purposes. Indeed, from Eqs. (14) and (16), it follows that (1) the phase factor  $e^{-i\phi_b}$  enters into the definition of  $\tilde{m}_{D_1}$  only and (2) that it appears there as an overall factor (before the absolute value is taken). Thus the values of  $\tilde{m}_{D_k}$  are completely independent of  $\phi_b$ . Similar conclusion holds for the sector of up quarks.

<sup>2</sup>Calculations of the scale dependence of CKM matrix elements over the enormous range from  $10^2$  GeV to  $10^{15}$  GeV show that this dependence is negligible for the Cabibbo submatrix and is of the order of 15% for the remaining off-diagonal elements [16]. Consequently, one may expect that a similar dependence over the much smaller range of 1– $10^2$  GeV is negligible.

#### IV. IMPOSING KOIDE'S CONDITION ON PSEUDOMASSES

The values of  $\delta_D = 4/27$  and  $\delta_U = 2/27$ , together with the low-energy ratios  $m_s/m_d = 20.4$ ,  $m_u/m_d = 0.56$  (see, e.g., Ref. [17]), suffice to fix the pattern of *effective* (low-energy) quark masses up to two overall mass scales (in the up and down sectors). These scales are irrelevant for the discussion of Koide's formulas for pseudomasses. For illustrative purposes, however, one may set  $m_s = 160.0$  MeV and  $m_t = 172000$  MeV. This choice leads to the following representative values of effective quark masses (in MeV):

$$\begin{aligned} m_d &= 7.843, & m_s &= 160.0, & m_b &= 4209, \\ m_u &= 4.392, & m_c &= 1296, & m_t &= 172000, \end{aligned} \quad (24)$$

which sit right in the middle of the ranges given in Sec. 2.1 of Ref. [18]. Our results do not depend critically on the exact values of the effective masses used; hence, there is no need

to include uncertainties in these masses (e.g., as given in Ref. [18]).

For a given value of  $k_f$ , with  $\phi_b, \phi_t$  dropping from expression (14) for the pseudomasses, the condition

$$\frac{\sum_j \tilde{m}_{f_j}}{(\sum_j \sqrt{\tilde{m}_{f_j}})^2} = \frac{1 + k_f^2}{3} \quad (25)$$

imposes two constraints: between  $\theta_d$  and  $\theta_b$  in the down quark sector, and between  $\theta_u$  and  $\theta_t$  in the up quark sector. Thus,  $\theta_b$  becomes dependent on  $\theta_d$  (and  $\theta_t$  on  $\theta_u$ ). For  $k_D = k_U = 1$  there are two possible solutions for function  $\theta_b(\theta_d)$  and two possible solutions for function  $\theta_t(\theta_u)$ . They are shown in Fig. 1 with solid lines marked as  $\theta_{b,n}$  and  $\theta_{t,n}$  ( $n = 1, 2$ ). However, only the combination  $\theta_{\text{Koide}} = \theta_{b,1} - \theta_{t,1}$  is positive. Specifically, taking the central experimental values of  $\theta_u$  and  $\theta_d$ , one finds




FIG. 1. Correlations required by Eq. (25): (a)  $\theta_b \leftrightarrow \theta_d$  and (b)  $\theta_t \leftrightarrow \theta_u$  (all angles in degrees). Solid lines correspond to  $k_D = k_U = 1$ . Dashed lines denote solutions  $\theta_{b,1}$  and  $\theta_{t,1}$  for  $k_D = k_U = 1.015$ .




FIG. 2. Contour plots of  $\theta = \theta_{b,1}(\theta_d) - \theta_{t,1}(\theta_u)$  (all angles in degrees): (a)  $k_D = k_U = 1$  and (b)  $k_D = k_U = 1.015$ . Dashed lines are contours corresponding to  $\theta = 2.37^\circ \pm 0.10^\circ$ . Cross denotes experimental point  $(\theta_d, \theta_u) = (12.11^\circ \pm 0.47^\circ, 4.87^\circ \pm 0.23^\circ)$ .

$$\theta_{\text{Koide}} = 2.98^\circ, \quad (26)$$

which is only slightly different from the value given in Eq. (23).

A question thus emerges: how much must  $k_D, k_U$  depart from 1 to fit the current experimental value of  $\theta$ ? The dashed lines in Fig. 1 correspond to  $\theta_{b,1}$  and  $\theta_{t,1}$ , as obtained for  $k_D = k_U = 1.015$ . The predicted value of  $\theta$  is then  $\theta(k_D = k_U = 1.015) \approx 2.44^\circ$ , which is in good agreement with experiment. For completeness, we also show the contour plot of  $\theta$  as a function of  $\theta_d$  and  $\theta_u$  for  $k_D = k_U = 1$  [Fig. 2(a)] and  $k_D = k_U = 1.015$  [Fig. 2(b)]. Dashed lines correspond to contours  $2.37^\circ \pm 2\sigma$  (with  $\sigma = 0.05^\circ$ ).

Although it might seem that the above results indicate that one cannot obtain  $k_U = k_D = 1$  (as discussed earlier, playing with the phases  $\phi_b, \phi_t$  does not affect the pseudomasses), this is not the case. One has to remember that while parametrizations (16) have been suggested as the most appropriate ones [14], they may be naturally

modified. Indeed, the same  $V_{\text{CKM}}$  is obtained if one substitutes  $U_D \rightarrow U'_D = W_D U_D$  and  $U_U \rightarrow U'_U = W_U U_U$ , provided  $W_D$  and  $W_U$  denote the same arbitrary unitary matrix. If  $W_D \neq W_U$ , but both  $W_D$  and  $W_U$  are very close to 1, a minor modification of our results is expected. Such a natural modification of  $U_{D(U)} = R_{23}^{D(U)} R_{12}^{D(U)}$  is obtained if we put  $W_{D(U)} = R_{13}^{D(U)}(\theta_{13}^{D(U)})$  with very small  $\theta_{13}^D \neq \theta_{13}^U$ . Since the inclusion of two new parameters  $\theta_{13}^{D(U)}$  introduces additional freedom into the scheme, it does not make sense to study it here. However—keeping this freedom in mind—the parametrization of Eq. (16) and our numerical results may be viewed as capturing the dominant effects only.

In conclusion, the data are consistent with the statement that low-energy quark masses satisfy phase relations  $\delta_D = 2\delta_U = 4/27$ , while the expected Koide relations  $k_U = k_D = 1$  hold approximately for masses transformed to the weak basis, as suggested in Ref. [12]. These observations might be relevant for a future theory of mass.

- 
- [1] Y. Koide, *Lett. Nuovo Cimento* **34**, 201 (1982); *Phys. Lett.* **120B**, 161 (1983); *Phys. Rev. D* **28**, 252 (1983).  
 [2] A. Rivero and A. Gsponer, [arXiv:hep-ph/0505220](https://arxiv.org/abs/hep-ph/0505220).  
 [3] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012).  
 [4] Y. Koide, [arXiv:hep-ph/0005137](https://arxiv.org/abs/hep-ph/0005137); *J. Phys. G* **34**, 1653 (2007).  
 [5] C. Brannen, as cited in Ref. [2].  
 [6] C. Brannen, <http://brannenworks.com/MASSES2.pdf>, 2006.  
 [7] G. Rosen, *Mod. Phys. Lett. A* **22**, 283 (2007).  
 [8] R.N. Mohapatra and A.Y. Smirnov, *Annu. Rev. Nucl. Part. Sci.* **56**, 569 (2006).  
 [9] W. Rodejohann and H. Zhang, *Phys. Lett. B* **698**, 152 (2011).  
 [10] Z.-Z. Xing and H. Zhang, *Phys. Lett. B* **635**, 107 (2006).  
 [11] P. Żenczykowski, *Phys. Rev. D* **86**, 117303 (2012).  
 [12] J.-M. Gérard, F. Goffinet, and M. Herquet, *Phys. Lett. B* **633**, 563 (2006).  
 [13] A. Rašin, [arXiv:hep-ph/9708216](https://arxiv.org/abs/hep-ph/9708216).  
 [14] H. Fritzsch and Z.-Z. Xing, *Phys. Lett. B* **413**, 396 (1997).  
 [15] H. Fritzsch and Z.-Z. Xing, *Phys. Rev. D* **57**, 594 (1998).  
 [16] C. Balzereit, T. Hansmann, T. Mannel, and B. Plümper, *Eur. Phys. J. C* **9**, 197 (1999).  
 [17] S. Weinberg, *Trans. N.Y. Acad. Sci.* **38**, 185 (1977).  
 [18] H. Fritzsch and Z.-Z. Xing, *Prog. Part. Nucl. Phys.* **45**, 1 (2000).