

Gravitationally driven electromagnetic perturbations of neutron stars and black holesHajime Sotani,¹ Kostas D. Kokkotas,^{2,3} Pablo Laguna,⁴ and Carlos F. Sopuerta⁵¹*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*²*Theoretical Astrophysics, University of Tübingen, IAAT, Auf der Morgenstelle 10, 72076 Tübingen, Germany*³*Department of Physics, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece*⁴*Center for Relativistic Astrophysics and School of Physics, Georgia Institute of Technology, Atlanta, Georgia 30332, USA*⁵*Facultat de Ciències, Institut de Ciències de l'Espai (CSIC-IEEC),**Campus UAB, Torre C5 parells, Bellaterra, 08193 Barcelona, Spain*

(Received 15 February 2013; published 5 April 2013)

Gravitational perturbations of neutron stars and black holes are well-known sources of gravitational radiation. If the compact object is immersed in or endowed with a magnetic field, the gravitational perturbations would couple to electromagnetic perturbations and potentially trigger synergistic electromagnetic signatures. We present a detailed analytic calculation of the dynamics of coupled gravitational and electromagnetic perturbations for both neutron stars and black holes. We discuss the prospects for detecting the electromagnetic waves in these scenarios and the potential that these waves have for providing information about their source.

DOI: [10.1103/PhysRevD.87.084018](https://doi.org/10.1103/PhysRevD.87.084018)

PACS numbers: 04.30.-w, 04.40.Nr, 95.85.Sz

I. INTRODUCTION

Multimessenger astronomy has arrived. Already astroparticle observations (neutrinos and cosmic rays) are complementing traditional electromagnetic observations. The third pillar is almost ready with near future gravitational-wave observations by interferometric detectors like LIGO, Virgo, GEO600, and LCGT [1,2]. This new astronomy will enable multichannel observations of astrophysical phenomena such as γ -ray bursts, supernovae, or flaring magnetars, unveiling an unprecedented view of the nature of the source and its environment.

An important component in many astrophysical phenomena is strong magnetic fields, as demonstrated by the active role they play in the accretion processes of low-mass x-ray binaries and gamma-ray bursts [3]. The presence of strong magnetic fields opens up the possibility for interesting effects. Among them, which is the central topic of this work, is the coupling between electromagnetic and gravitational emissions that could yield synergistic multimessenger observations. In particular, it is important to assess the conditions in which electromagnetic and gravitational emissions influence each other. There are already hints for such scenario. It is believed that the flare activity of magnetars seems to be associated with starquakes [4]. These quakes are responsible not only for dramatic perturbations and rearrangements of the magnetic field, but also for the breaking of the neutron star crust and internal motions, possibly resulting in the emission of gravitational waves. Detailed studies of magnetar flare activity have revealed a number of features in the afterglow, which can be associated with the crust oscillations as well as with Alfvén waves propagating from the core towards the surface [5–17].

The link or coupling between electromagnetic radiation and gravitational waves has been investigated for some cases. One of them looked at the propagation of

gravitational waves linearly coupled to an external magnetic field [18]. It was shown that this configuration triggers magnetohydrodynamics waves in the plasma [19–23]. Furthermore, the linear nature of the coupling limits the electromagnetic waves to low frequencies, in the best case a few tenths of kHz, which will be easily absorbed by the interstellar medium or plasma. In order to produce high frequency and detectable electromagnetic waves, nonlinear couplings are needed, requiring much stronger gravitational waves. In most of these studies, the gravitational waves were assumed to propagate on a flat space-time background. This is a reasonable assumption when the interaction between the gravitational and electromagnetic waves takes place far from the source. There have been only very few attempts to treat the electromagnetic-gravity coupling in the strong field regime [24].

The aim of this work is to study the interaction of electromagnetic and gravitational waves in the vicinity of magnetized neutron stars or black holes immersed in strong magnetic fields using perturbation theory, paying particular attention to how gravitational modes drive the excitation of electromagnetic perturbations. Our work also includes estimates of the energy transferred between the gravitational and electromagnetic sectors. As expected, we find that the excited electromagnetic waves have roughly the same frequency as the driving gravitational waves, i.e., of the order of a few kHz. Electromagnetic waves at these low frequencies can be easily absorbed by the interstellar medium. As a consequence, one needs to associate them with secondary emission mechanisms (e.g., synchrotron radiation) in order to be able to trace the effects of gravitational waves on the strong magnetic fields. The later process can be studied following the mechanisms described in Refs. [19–23], and there is work in progress for the special case of strong gravitational fields.

This article is organized as follows: Sec. II gives details of the space-time background configuration. In Sec. III, we review the general form of the perturbation equations, their couplings, and the angular dependences of the various types of electromagnetic and gravitational perturbations. In Sec. IV, we reduce the equations to the particular case of dipole electromagnetic perturbations driven by the quadrupole gravitational mode for the case of a neutron star background. In Sec. V, we do the same as in Sec. IV but for the case of a black hole and consider both the case of axial and polar gravitational perturbations. In Sec. VI we show numerical results on dipole electromagnetic waves driven by quadrupole gravitational waves with axial parity for both neutron stars and black holes. Conclusions are given in Sec. VII. We adopt geometric units, $c = G = 1$, where c and G denote the speed of light and the gravitational constant, respectively, and the metric signature is $(-, +, +, +)$.

II. EQUATIONS FOR THE BACKGROUND

The background space-times we are considering (neutron stars and black holes) are governed by the Einstein-Maxwell equations, which read

$$G_{\mu\nu} = 8\pi(T_{\mu\nu} + E_{\mu\nu}), \quad (2.1)$$

$$(T^{\mu\nu} + E^{\mu\nu})_{;\nu} = 0, \quad (2.2)$$

$$F^{\mu\nu}_{;\nu} = 4\pi J^{\mu}, \quad (2.3)$$

$$F_{\mu\nu,\lambda} + F_{\lambda\mu,\nu} + F_{\nu\lambda,\mu} = 0. \quad (2.4)$$

The tensors that appear in these equations are the Einstein tensor $G_{\mu\nu}$, the Faraday antisymmetric tensor $F_{\mu\nu}$, the electromagnetic four-current J^{μ} , the energy-momentum tensor of the matter fluid $T_{\mu\nu}$, and the energy-momentum tensor of the electromagnetic field is $E_{\mu\nu}$. The energy-momentum tensors are explicitly given by

$$T_{\mu\nu} = (\rho + p)u_{\mu}u_{\nu} + pg_{\mu\nu}, \quad (2.5)$$

$$E_{\mu\nu} = \frac{1}{4\pi} \left(g^{\rho\sigma} F_{\rho\mu} F_{\sigma\nu} - \frac{1}{4} g_{\mu\nu} F_{\rho\sigma} F^{\rho\sigma} \right), \quad (2.6)$$

where ρ stands for the energy density, p for the pressure, and u_{μ} for the four-velocity of the matter fluid.

The presence of a magnetic field could in principle induce deformations to the neutron star or black hole we are considering. However, even for astrophysically strong magnetic fields, $B \sim 10^{16}G$, as in the case of magnetars, the energy of the magnetic field \mathcal{E}_B is much smaller than the gravitational energy \mathcal{E}_G , by several orders of magnitude. In fact, $\mathcal{E}_B/\mathcal{E}_G \sim 10^{-4}(B/10^{16}[G])^2$. Therefore, in setting up the background space-time metric, one can ignore the magnetic field. That is, the background metric has the form

$$ds^2 = -e^{\nu} dt^2 + e^{\lambda} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (2.7)$$

where the functions $\nu(r)$ and $\lambda(r)$ in the interior of a neutron star are determined by the well-known Tolman-Oppenheimer-Volkoff equations (see, e.g., Ref. [25]) and the matter fluid four-velocity $u^{\mu} = (e^{-\nu/2}, 0, 0, 0)$. In the exterior of a neutron star, and in the case of a black hole, they are determined by the standard Schwarzschild solution: $e^{-\lambda} = e^{\nu} = 1 - 2M/r$.

A. A dipole background magnetic field: Exterior region

Next, we compute the magnetic field for both the neutron star and the black hole. We consider first the exterior (vacuum) solution. In this case, the component of Maxwell equations given by Eq. (2.4) is automatically satisfied. The magnetic field is then obtained by solving the remaining Maxwell equations, Eq. (2.3), which in vacuum read

$$F^{\mu\nu}_{;\nu} = 0, \quad (2.8)$$

with $F_{\mu\nu} = A_{\nu,\mu} - A_{\mu,\nu}$. Since the background space-time is static, it is natural to assume that the magnetic field is also static. In addition, we require the magnetic field to be axisymmetric and poloidal,

$$B^{\mu(\text{ex})} = (0, e^{-\lambda/2} B_1^{(\text{ex})}(r) \cos\theta, e^{-\lambda/2} B_2^{(\text{ex})}(r) \sin\theta, 0), \quad (2.9)$$

which has a dependence on the polar coordinate, θ . From the relation between the magnetic field, the matter fluid velocity u^{μ} , and the field strength

$$B_{\mu} = \epsilon_{\mu\nu\alpha\beta} u^{\nu} F^{\alpha\beta} / 2 = \epsilon_{\mu\nu\alpha\beta} u^{\nu} A^{\alpha\beta}, \quad (2.10)$$

where $\epsilon_{\mu\nu\alpha\beta}$ is the complete antisymmetric tensor determined by the convention $\epsilon_{0123} = \sqrt{-g}$. It is not difficult to show that the only nonvanishing component of the vector potential A_{μ} is the ϕ component, which we will denote as $A_{\phi}^{(\text{ex})}$. Therefore, the vacuum Maxwell equation (2.8) in the Schwarzschild background becomes

$$r^2 \frac{\partial}{\partial r} \left[\left(1 - \frac{2M}{r} \right) \frac{\partial A_{\phi}^{(\text{ex})}}{\partial r} \right] + \sin\theta \frac{\partial}{\partial \theta} \left[\frac{1}{\sin\theta} \frac{\partial A_{\phi}^{(\text{ex})}}{\partial \theta} \right] = 0. \quad (2.11)$$

Expanding $A_{\phi}^{(\text{ex})}$ in vector spherical harmonics as

$$A_{\phi}^{(\text{ex})} = a_{l_M}^{(\text{ex})}(r) \sin\theta \partial_{\theta} P_{l_M}(\cos\theta), \quad (2.12)$$

we rewrite Eq. (2.11) as

$$r^2 \frac{d}{dr} \left[\left(1 - \frac{2M}{r} \right) \frac{da_{l_M}^{(\text{ex})}}{dr} \right] - l(l+1) a_{l_M}^{(\text{ex})} = 0. \quad (2.13)$$

The solution of this equation for the dipole case ($l_M = 1$) has the form [26]

$$a_1^{(\text{ex})} = -\frac{3\mu_d}{8M^3} r^2 \left[\ln\left(1 - \frac{2M}{r}\right) + \frac{2M}{r} + \frac{2M^2}{r^2} \right], \quad (2.14)$$

where μ_d is the magnetic dipole moment for an observer at infinity. With the solution of Eqs. (2.14) and (2.10), the coefficients of the magnetic field in Eq. (2.9) are given by

$$B_1^{(\text{ex})}(r) = \frac{2a_1^{(\text{ex})}}{r^2} = -\frac{3\mu_d}{4M^3} \left[\ln\left(1 - \frac{2M}{r}\right) + \frac{2M}{r} + \frac{2M^2}{r^2} \right], \quad (2.15)$$

$$B_2^{(\text{ex})}(r) = -\frac{a_{1,r}^{(\text{ex})}}{r^2} = \frac{3\mu_d}{4M^3 r} \left[\ln\left(1 - \frac{2M}{r}\right) + \frac{M}{r} + \frac{M}{r-2M} \right]. \quad (2.16)$$

Notice that in the limit $r \rightarrow \infty$,

$$B_1^{(\text{ex})}(r) \approx \frac{2\mu_d}{r^3} \quad \text{and} \quad B_2^{(\text{ex})}(r) \approx \frac{\mu_d}{r^4}. \quad (2.17)$$

B. A dipole background magnetic field: Interior region

We assume that the magnetic field inside the star is also axisymmetric and poloidal, with current $J_\mu = (0, 0, 0, J_\phi)$ [27,28]. The ideal MHD approximation is also adopted, i.e., infinite conductivity σ , which leads to $E_\mu = F_{\mu\nu}u^\nu = 0$, as follows from the relativistic Ohm's law

$$F_{\mu\nu}u^\nu = \frac{4\pi}{\sigma} (J_\mu + u_\mu J^\nu u_\nu). \quad (2.18)$$

Therefore, the vector potential A_μ is similar to that for the exterior magnetic field, i.e., $A_\mu = (0, 0, 0, A_\phi^{(\text{in})})$. The counterpart equation to Eq. (2.11) but for the interior is

$$e^{-\lambda} \frac{\partial^2 A_\phi^{(\text{in})}}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2 A_\phi^{(\text{in})}}{\partial \theta^2} + (\nu' - \lambda') \frac{e^{-\lambda}}{2} \frac{\partial A_\phi^{(\text{in})}}{\partial r} - \frac{1}{r^2} \frac{\cos \theta}{\sin \theta} \frac{\partial A_\phi^{(\text{in})}}{\partial \theta} = -4\pi J_\phi. \quad (2.19)$$

Expanding both, the vector potential $A_\phi^{(\text{in})}$ and the current J_ϕ , in vector spherical harmonics, one gets

$$A_\phi^{(\text{in})}(r, \theta) = a_{l_M}^{(\text{in})}(r) \sin \theta \partial_\theta P_{l_M}(\cos \theta), \quad (2.20)$$

$$J_\phi(r, \theta) = j_{l_M}(r) \sin \theta \partial_\theta P_{l_M}(\cos \theta), \quad (2.21)$$

which can be used to rewrite Eq. (2.19) as

$$e^{-\lambda} \frac{d^2 a_{l_M}^{(\text{in})}}{dr^2} + (\nu' - \lambda') \frac{e^{-\lambda}}{2} \frac{da_{l_M}^{(\text{in})}}{dr} - \frac{l_M(l_M + 1)}{r^2} a_{l_M}^{(\text{in})} = -4\pi j_{l_M}. \quad (2.22)$$

It is only feasible to obtain numerical solutions to Eq. (2.22), even for the dipole case ($l_M = 1$), since among

other things the coefficients are also computed numerically from the Tolman-Oppenheimer-Volkoff equations. In addition, when prescribing $j_1(r)$, it must satisfy an integrability condition (see Refs. [29,30] for details). We adopt a current with a functional form [28]:

$$j_1(r) = f_0 r^2 (\rho + p), \quad (2.23)$$

where f_0 is an arbitrary constant. In addition, we should impose the following regularity condition at center of the neutron star:

$$a_1^{(\text{in})} = \alpha_c r^2 + \mathcal{O}(r^4), \quad (2.24)$$

where α_c is also an arbitrary constant. These arbitrary constants f_0 and α_c are determined by the matching conditions at the surface of the star, namely that a_1 and $a_{1,r}$ are continuous across the stellar surface. Finally, once we have the numerical solution for $a_1(r)$, the magnetic field is obtained from

$$B^{\mu(\text{in})} = (0, e^{-\lambda/2} B_1^{(\text{in})}(r) \cos \theta, e^{-\lambda/2} B_2^{(\text{in})}(r) \sin \theta, 0) \quad (2.25)$$

with

$$B_1^{(\text{in})}(r) = \frac{2a_1^{(\text{in})}}{r^2} \quad \text{and} \quad B_2^{(\text{in})}(r) = -\frac{a_{1,r}^{(\text{in})}}{r^2}. \quad (2.26)$$

With the magnetic field determined both in the interior and exterior regions, the Faraday tensor for the background field becomes

$$F_{\mu\nu} = \epsilon_{\mu\nu\alpha\beta} B^\alpha u^\beta = r^2 \sin \theta \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & B_2 \sin \theta \\ 0 & 0 & 0 & -B_1 \cos \theta \\ 0 & -B_2 \sin \theta & B_1 \cos \theta & 0 \end{pmatrix}. \quad (2.27)$$

III. PERTURBATION EQUATIONS

We consider small perturbations of both the gravitational and electromagnetic fields, which can be described as

$$\tilde{g}_{\mu\nu} = g_{\mu\nu} + h_{\mu\nu}, \quad (3.1)$$

$$\tilde{F}_{\mu\nu} = F_{\mu\nu} + f_{\mu\nu}, \quad (3.2)$$

where $g_{\mu\nu}$ and $F_{\mu\nu}$ are the background quantities derived in the previous section. The tensors $h_{\mu\nu}$ and $f_{\mu\nu}$ denote small perturbations, i.e., $h_{\mu\nu} = \delta g_{\mu\nu}$ and $f_{\mu\nu} = \delta F_{\mu\nu}$. Linearization of the Einstein-Maxwell equations yields

$$\delta G_{\mu\nu} = 8\pi \delta(T_{\mu\nu} + E_{\mu\nu}), \quad (3.3)$$

$$\delta(T^{\mu\nu}{}_{;\nu} + E^{\mu\nu}{}_{;\nu}) = 0, \quad (3.4)$$

$$\partial_\nu [(-g)^{1/2} f^{\mu\nu}] = 4\pi\delta [(-g)^{1/2} J^\mu] - \partial_\nu [F^{\mu\nu} \delta(-g)^{1/2}], \quad (3.5)$$

$$f_{\mu\nu,\lambda} + f_{\lambda\mu,\nu} + f_{\nu\lambda,\mu} = 0. \quad (3.6)$$

From Eq. (3.5), we find that the electromagnetic perturbations are driven by the gravitational perturbations via the term containing $\delta(-g)^{1/2}$ in the right-hand side. On the other hand, for simplicity, we omit the backreaction of the electromagnetic perturbations on the gravitational perturbations, i.e., we set $\delta E_{\mu\nu} = \delta(E^{\mu\nu}{}_{;\nu}) = 0$ in Eqs. (3.3) and (3.4). This simplification is based on the assumption that the energy stored in gravitational perturbations is considerably larger than that in electromagnetic perturbations, which are typically driven by the former. On the other hand, in the giant flares of SGR 1806 – 20 and

SGR 1900 + 14 [31–33], whose peak luminosities are in the range of 10^{44} – 10^{46} ergs s^{−1}, the dramatic rearrangement of the magnetic field might lead to emission of gravitational waves. Nevertheless, recent nonlinear MHD simulations [34–38] do not support these expectations.

The first two perturbative equations, Eqs. (3.3) and (3.4), have been studied extensively in the past, in the absence of magnetic fields, both for stellar and black hole backgrounds (see, e.g., Refs. [39–44]). Thus, in this article we use the perturbation equations derived in earlier works, and we derive the analytic form of the perturbation equations for the electromagnetic field together with their coupling to the gravitational perturbations.

The metric perturbations $h_{\mu\nu}$ in the Regge-Wheeler gauge [39] can be decomposed into tensor spherical harmonics in the following way:

$$h_{\mu\nu} = \sum_{l=2}^{\infty} \sum_{m=-l}^l \begin{pmatrix} e^\nu H_{0,lm} & H_{1,lm} & -h_{0,lm} \sin^{-1}\theta \partial_\phi & h_{0,lm} \sin \theta \partial_\theta \\ * & e^\lambda H_{2,lm} & -h_{1,lm} \sin^{-1}\theta \partial_\phi & h_{1,lm} \sin \theta \partial_\theta \\ * & * & r^2 K_{lm} & 0 \\ * & * & 0 & r^2 \sin^2 \theta K_{lm} \end{pmatrix} Y_{lm}, \quad (3.7)$$

where $H_{0,lm}$, $H_{1,lm}$, $H_{2,lm}$, and K_{lm} are the functions of (t, r) describing the *polar perturbations*, while $h_{0,lm}$ and $h_{1,lm}$ describe the *axial* ones. On the other hand, the tensor harmonic expansion of the electromagnetic perturbations, $f_{\mu\nu}$, for the *magnetic multipoles* (or *axial parity*) are given by

$$f_{\mu\nu}^{(M)} = \sum_{l=2}^{\infty} \sum_{m=-l}^l \begin{pmatrix} 0 & 0 & f_{02,lm}^{(M)} \sin^{-1}\theta \partial_\phi & -f_{02,lm}^{(M)} \sin \theta \partial_\theta \\ 0 & 0 & f_{12,lm}^{(M)} \sin^{-1}\theta \partial_\phi & -f_{12,lm}^{(M)} \sin \theta \partial_\theta \\ * & * & 0 & f_{23,lm}^{(M)} \sin \theta \\ * & * & * & 0 \end{pmatrix} Y_{lm}, \quad (3.8)$$

while the expansion for the *electric multipoles* (or *polar parity*) can be written as

$$f_{\mu\nu}^{(E)} = \sum_{l=2}^{\infty} \sum_{m=-l}^l \begin{pmatrix} 0 & f_{01,lm}^{(E)} & f_{02,lm}^{(E)} \partial_\theta & f_{02,lm}^{(E)} \partial_\phi \\ * & 0 & f_{12,lm}^{(E)} \partial_\theta & f_{12,lm}^{(E)} \partial_\phi \\ * & * & 0 & 0 \\ * & * & 0 & 0 \end{pmatrix} Y_{lm}. \quad (3.9)$$

Hereafter, the quantities describing the magnetic- and electric-type electromagnetic perturbations will be denoted with the indices (M) and (E), respectively. We point out that $h_{\mu\nu}$ is a symmetric tensor, while both $f_{\mu\nu}^{(M)}$ and $f_{\mu\nu}^{(E)}$ are antisymmetric tensors, i.e., $f_{\mu\nu}^{(M)} = -f_{\nu\mu}^{(M)}$ and $f_{\mu\nu}^{(E)} = -f_{\nu\mu}^{(E)}$. From the perturbed Maxwell equation, Eq. (3.6), we can obtain the following relations connecting the above perturbative functions:

$$f_{12,lm}^{(M)} = \frac{1}{\Lambda} \frac{\partial f_{23,lm}^{(M)}}{\partial r} \quad \text{and} \quad f_{02,lm}^{(M)} = \frac{1}{\Lambda} \frac{\partial f_{23,lm}^{(M)}}{\partial t}, \quad (3.10)$$

$$f_{01,lm}^{(E)} = \frac{\partial f_{02,lm}^{(E)}}{\partial r} - \frac{\partial f_{12,lm}^{(E)}}{\partial t}, \quad (3.11)$$

where $\Lambda \equiv l(l+1)$. Notice that $f_{23,lm}^{(E)}$ and $\tilde{\Psi}$ defined as

$$\tilde{\Psi} = -\frac{r^2}{\Lambda} f_{01,lm}^{(E)} \quad (3.12)$$

are *gauge-invariant* variables (see Eq. (II-27) in Ref. [45] and Eq. (II-11) in Ref. [46]).

A. Perturbations of a dipole magnetic field: Exterior region

In the exterior vacuum region, we adopt the condition $\delta J^\mu = 0$. With this condition, the perturbed electromagnetic fields will be determined via the linearized form of the Maxwell equation, Eq. (3.5) (assuming that $J^\mu = \delta J^\mu = 0$),

$$\partial_\nu [(-g)^{1/2} f^{\mu\nu}] = -\frac{1}{2} \partial_\nu [(-g)^{1/2} F^{\mu\nu} g^{\alpha\beta} h_{\alpha\beta}], \quad (3.13)$$

together with the perturbed Maxwell equation (3.6). Equation (3.13) for $\mu = t$ and $\mu = r$ can be written down as

$$\sum_{l,m} \{A_{lm}^{(I,E)} Y_{lm} + \tilde{A}_{lm}^{(I,A)} \cos \theta Y_{lm} + B_{lm}^{(I,A)} \sin \theta \partial_\theta Y_{lm} + C_{lm}^{(I,P)} \partial_\phi Y_{lm}\} = 0 \quad (I = 0, 1), \quad (3.14)$$

where the indices ‘‘A’’ and ‘‘P’’ stand for *axial* and *polar* gravitational perturbative quantities, and obviously ‘‘I’’ stands for the t and r components of Eq. (3.13). The coefficients of Eq. (3.14) have the following expressions:

$$A_{lm}^{(0,E)} = \frac{1}{2} \left(\nu' + \lambda' - \frac{4}{r} \right) f_{01,lm}^{(E)} - f_{01,lm}^{(E)'} + \frac{\Lambda}{r^2} e^\lambda f_{02,lm}^{(E)} \quad (3.15)$$

$$\tilde{A}_{lm}^{(0,A)} = \frac{\Lambda}{r^2} e^\lambda B_1 h_{0,lm}, \quad (3.16)$$

$$B_{lm}^{(0,A)} = \left[-\frac{1}{2} \left(\nu' + \lambda' - \frac{4}{r} \right) B_2 + B_2' + \frac{1}{r^2} e^\lambda B_1 \right] h_{0,lm} + B_2 h_{0,lm}', \quad (3.17)$$

$$C_{lm}^{(0,P)} = -B_2 H_{1,lm}, \quad (3.18)$$

$$A_{lm}^{(1,E)} = r^2 \dot{f}_{01,lm}^{(E)} - \Lambda e^\nu f_{12,lm}^{(E)}, \quad (3.19)$$

$$\tilde{A}_{lm}^{(1,A)} = -\Lambda e^\nu B_1 h_{1,lm}, \quad (3.20)$$

$$B_{lm}^{(1,A)} = -e^\nu B_1 h_{1,lm} - r^2 B_2 \dot{h}_{0,lm}, \quad (3.21)$$

$$C_{lm}^{(1,P)} = r^2 e^\nu B_2 H_{0,lm}. \quad (3.22)$$

One can decompose the equations above for a specific mode with fixed harmonic numbers (l, m) , by multiplying with Y_{lm}^* and integrating over the two-sphere, i.e.,

$$A_{lm}^{(I,E)} + im C_{lm}^{(I,P)} + Q_{lm} [\tilde{A}_{l-1m}^{(I,A)} + (l-1) B_{l-1m}^{(I,A)}] + Q_{l+1m} [\tilde{A}_{l+1m}^{(I,A)} - (l+2) B_{l+1m}^{(I,A)}] = 0 \quad (I = 0, 1). \quad (3.23)$$

In a similar way, from the two remaining equations, i.e., Eq. (3.13) for $\mu = \theta$ and $\mu = \phi$, one gets the relations

$$\sum_{l,m} \{(\alpha_{lm} + \tilde{\alpha}_{lm} \cos \theta) \partial_\theta Y_{lm} - (\beta_{lm} + \tilde{\beta}_{lm} \cos \theta) \times (\partial_\phi Y_{lm} / \sin \theta) + \eta_{lm} \sin \theta Y_{lm} + \chi_{lm} \sin \theta W_{lm}\} = 0, \quad (3.24)$$

$$\sum_{l,m} \{(\beta_{lm} + \tilde{\beta}_{lm} \cos \theta) \partial_\theta Y_{lm} + (\alpha_{lm} + \tilde{\alpha}_{lm} \cos \theta) \times (\partial_\phi Y_{lm} / \sin \theta) + \zeta_{lm} \sin \theta Y_{lm} + \chi_{lm} \sin \theta X_{lm}\} = 0, \quad (3.25)$$

where

$$W_{lm} = \left(\partial_\theta^2 - \cot \theta \partial_\theta - \frac{1}{\sin^2 \theta} \partial_\phi \right) Y_{lm}, \quad (3.26)$$

$$X_{lm} = 2 \partial_\phi (\partial_\theta - \cot \theta) Y_{lm}. \quad (3.27)$$

These equations lead to an extra set of evolution equations for a specific mode (l, m) by multiplying with Y_{lm}^* and integrating over the two-sphere:

$$\Lambda \alpha_{lm} - im [\tilde{\beta}_{lm} + \zeta_{lm}] + Q_{lm} (l+1) [(l-2)(l-1) \chi_{l-1m} + (l-1) \tilde{\alpha}_{l-1m} - \eta_{l-1m}] - Q_{l+1m} l [(l+2)(l+3) \chi_{l+1m} - (l+2) \tilde{\alpha}_{l+1m} - \eta_{l+1m}] = 0, \quad (3.28)$$

$$\Lambda \beta_{lm} + im [(l-1)(l+2) \chi_{lm} + \tilde{\alpha}_{lm} + \eta_{lm}] + Q_{lm} (l+1) [(l-1) \tilde{\beta}_{l-1m} - \zeta_{l-1m}] + Q_{l+1m} l [(l+2) \tilde{\beta}_{l+1m} + \zeta_{l+1m}] = 0, \quad (3.29)$$

where the coefficients are given by

$$\alpha_{lm} = \frac{1}{2} (\lambda' - \nu') f_{12,lm}^{(E)} + e^{\lambda-\nu} \dot{f}_{02,lm}^{(E)} - f_{12,lm}^{(E)'} \quad (3.30)$$

$$\beta_{lm} = \frac{1}{2} (\nu' - \lambda') f_{12,lm}^{(M)} - e^{\lambda-\nu} \dot{f}_{02,lm}^{(M)} + f_{12,lm}^{(M)'} - \frac{1}{r^2} e^\lambda f_{23,lm}^{(M)}, \quad (3.31)$$

$$\tilde{\alpha}_{lm} = \left[\frac{1}{2} (\lambda' - \nu') B_1 - B_1' + B_2 \right] h_{1,lm} - B_1 h_{1,lm}' + e^{\lambda-\nu} B_1 \dot{h}_{0,lm}, \quad (3.32)$$

$$\tilde{\beta}_{lm} = e^\lambda B_1 K_{lm}, \quad (3.33)$$

$$\eta_{lm} = \frac{\Lambda}{2} B_2 h_{1,lm}, \quad (3.34)$$

$$\chi_{lm} = \frac{1}{2} B_2 h_{1,lm}, \quad (3.35)$$

$$\zeta_{lm} = \left[\frac{r^2}{2} (\lambda' - \nu') B_2 - 2r B_2 - r^2 B_2' \right] H_{0,lm} - r^2 B_2 H_{0,lm}' - e^\lambda B_1 K_{lm} + e^{-\nu} r^2 B_2 \dot{H}_{1,lm}. \quad (3.36)$$

B. Perturbations of a dipole magnetic field: Interior region

In the stellar interior, because we have adopted the ideal MHD approximation for which $F_{\mu\nu}u^\nu = 0$, the components of the perturbed electromagnetic field tensor are determined by using the perturbed Maxwell equation (3.6), i.e.,

$$f_{0\mu} = e^{\nu/2} F_{\mu\nu} \delta u^\nu, \quad (3.37)$$

where δu^μ is the perturbed fluid four-velocity defined as

$$\delta u^\mu = \left(\frac{1}{2} e^{-\nu/2} H_{0,lm}, R_{lm}, V_{lm} \partial_\theta \right. \\ \left. - U_{lm} \sin^{-1} \theta \partial_\phi, V_{lm} \sin^{-2} \theta \partial_\phi + U_{lm} \sin^{-1} \theta \partial_\theta \right) Y_{lm}. \quad (3.38)$$

From Eq. (3.37) one can get the following equations:

$$\sum_{l,m} \{ f_{01,lm}^{(E)} Y_{lm} - r^2 B_2 e^{\nu/2} (V_{lm} \partial_\phi Y_{lm} + U_{lm} \sin \theta \partial_\theta Y_{lm}) \} = 0, \quad (3.39)$$

$$\sum_{l,m} \{ (\mathcal{A}_{lm} + \tilde{\mathcal{A}}_{lm} \cos \theta) \partial_\theta Y_{lm} \\ - (\mathcal{B}_{lm} + \tilde{\mathcal{B}}_{lm} \cos \theta) (\partial_\phi Y_{lm} / \sin \theta) \} = 0, \quad (3.40)$$

$$\sum_{l,m} \{ (\mathcal{B}_{lm} + \tilde{\mathcal{B}}_{lm} \cos \theta) \partial_\theta Y_{lm} \\ + (\mathcal{A}_{lm} + \tilde{\mathcal{A}}_{lm} \cos \theta) (\partial_\phi Y_{lm} / \sin \theta) \\ + \tilde{\mathcal{C}}_{lm} (\sin \theta Y_{lm}) \} = 0, \quad (3.41)$$

where the coefficients \mathcal{A}_{lm} and \mathcal{B}_{lm} are functions of the perturbed electromagnetic fields, while $\tilde{\mathcal{A}}_{lm}$, $\tilde{\mathcal{B}}_{lm}$, and $\tilde{\mathcal{C}}_{lm}$ are functions of the perturbed matter fluid four-velocity. The expressions for these coefficients are

$$\mathcal{A}_{lm} = f_{02,lm}^{(E)}, \quad (3.42)$$

$$\mathcal{B}_{lm} = -f_{02,lm}^{(M)}, \quad (3.43)$$

$$\tilde{\mathcal{A}}_{lm} = r^2 B_1 e^{\nu/2} U_{lm}, \quad (3.44)$$

$$\tilde{\mathcal{B}}_{lm} = -r^2 B_1 e^{\nu/2} V_{lm}, \quad (3.45)$$

$$\tilde{\mathcal{C}}_{lm} = r^2 B_2 e^{\nu/2} R_{lm}. \quad (3.46)$$

By multiplying Eqs. (3.39), (3.40), and (3.41) with Y_{lm}^* and integrating over the two-sphere we can obtain the following system of equations that depends only on r :

$$f_{01,lm}^{(E)} - r^2 B_2 e^{\nu/2} [imV_{lm} + Q_{lm}(l-1)U_{l-1m} \\ - Q_{l+1m}(l+2)U_{l+1m}] = 0, \quad (3.47)$$

$$\Lambda \mathcal{A}_{lm} - im[\tilde{\mathcal{B}}_{lm} + \tilde{\mathcal{C}}_{lm}] + Q_{lm}(l-1)(l+1)\tilde{\mathcal{A}}_{l-1m} \\ + Q_{l+1m}l(l+2)\tilde{\mathcal{A}}_{l+1m} = 0, \quad (3.48)$$

$$\Lambda \mathcal{B}_{lm} + im\tilde{\mathcal{A}}_{lm} + Q_{lm}(l+1)[(l-1)\tilde{\mathcal{B}}_{l-1m} - \tilde{\mathcal{C}}_{l-1m}] \\ + Q_{l+1m}l[(l+2)\tilde{\mathcal{B}}_{l+1m} + \tilde{\mathcal{C}}_{l+1m}] = 0, \quad (3.49)$$

where

$$Q_{lm} \equiv \sqrt{\frac{(l-m)(l+m)}{(2l-1)(2l+1)}}. \quad (3.50)$$

Finally, we should compute Eqs. (3.23), (3.28), and (3.29), for the exterior region, and Eqs. (3.47), (3.48), and (3.49) for the interior region of the star. From this system of equations, we can see the specific couplings between the electromagnetic and gravitational perturbations. For example, an electromagnetic perturbation of specific parity with harmonic indices (l, m) depends on the gravitational perturbations of the same parity with (l, m) as well as the gravitational perturbations of the opposite parity with $(l \pm 1, m)$. In other words, for the special and simpler case of axisymmetric perturbations ($m = 0$), we arrive at the following conclusions: (1) Dipole electric (polar) electromagnetic perturbations will be driven by axial quadrupole gravitational perturbations, and (2) dipole magnetic (axial) electromagnetic perturbations will be driven by polar quadrupole and radial gravitational perturbations. These two types of couplings will be discussed in detail in the next sections.

C. Junction conditions for perturbed electromagnetic fields

In order to close the system of equations derived in the previous subsection, we should impose appropriate junction conditions on the stellar surface. Such junction conditions for the perturbed electromagnetic fields can be derived from the conditions

$$n^\mu \delta B_\mu^{(in)} = n^\mu \delta B_\mu^{(ex)}, \quad (3.51)$$

$$q_\mu{}^\nu \delta E_\nu^{(in)} = q_\mu{}^\nu \delta E_\nu^{(ex)}, \quad (3.52)$$

where n^μ is the unit outward normal vector to the stellar surface, while $q_\mu{}^\nu$ is the corresponding projection tensor associated with n^μ . These junction conditions lead to the following set of equations:

$$f_{23}^{(M)(in)} = f_{23}^{(M)(ex)}, \quad (3.53)$$

$$f_{02}^{(M)(in)} = f_{02}^{(M)(ex)} = 0, \quad (3.54)$$

$$f_{02}^{(E)(ex)} = 0. \quad (3.55)$$

IV. DIPOLE PERTURBATIONS OF A MAGNETIC FIELD ON A STELLAR BACKGROUND

In the previous section, we provided the general form of the perturbative equations. In order to focus on a simple case, we only consider axisymmetric perturbations ($m = 0$) in this section. In this way, the various couplings become less complicated. Under these conditions, we study the excitation of dipole electric perturbations driven by axial gravitational ones and dipole magnetic perturbations driven by polar gravitational ones. These perturbative modes are actually the most important ones from the energetic point of view.

A. Dipole electric perturbations driven by axial gravitational perturbations

Here, we consider only dipole “electric-type” perturbations driven by quadrupole axial gravitational perturbations. Since we neglect the backreaction of electromagnetic perturbations on the gravitational ones, the quadrupole axial gravitational perturbations of a spherically symmetric star can be described by a single wave equation [40,47], which is given by

$$\frac{\partial^2 X_{lm}}{\partial t^2} - \frac{\partial^2 X_{lm}}{\partial r_*^2} + e^\nu \left(\frac{\Lambda}{r^2} - \frac{6m}{r^3} + 4\pi(\rho - p) \right) X_{lm} = 0, \quad (4.1)$$

where

$$X_{lm} = \frac{e^{(\nu-\lambda)/2}}{r} h_{1,lm} \quad \text{and} \quad \frac{\partial}{\partial r} = e^{(\lambda-\nu)/2} \frac{\partial}{\partial r_*}. \quad (4.2)$$

Note that r_* is the tortoise coordinate defined as $r_* = r + 2M \ln(r/2M - 1)$. Since there are no fluid oscillations if the matter is assumed to be described as a perfect fluid (unless we introduce rotation), the space-time only contains pure space-time modes, i.e., the so-called w modes [42,47,48]. In this case, the axial component of the fluid perturbation U_{lm} has the form $U_{lm} = -e^{-\nu/2} h_{0,lm}/r^2$, while the component of $h_{0,lm}$ is computed from the equation

$$\frac{\partial}{\partial t} h_{0,lm} = e^{(\nu-\lambda)/2} X_{lm} + r \frac{\partial}{\partial r_*} X_{lm}, \quad (4.3)$$

which is used later to simplify the coupling terms between the two types of perturbations.

On the other hand, in the same way as in the case of electromagnetic perturbations in the exterior region, Eqs. (3.11) and (3.23) for $I = 1$, and Eq. (3.28) lead to three simple evolution equations for the three perturbation functions $f_{12,10}^{(E)}$, $f_{01,10}^{(E)}$, and $f_{02,10}^{(E)}$:

$$\frac{\partial f_{12,10}^{(E)}}{\partial t} = e^{-\nu} \frac{\partial f_{02,10}^{(E)}}{\partial r_*} - f_{01,10}^{(E)}, \quad (4.4)$$

$$\frac{\partial f_{01,10}^{(E)}}{\partial t} = \frac{2}{r^2} e^\nu f_{12,01}^{(E)} + S_{20}^{(1)}, \quad (4.5)$$

$$\frac{\partial f_{02,10}^{(E)}}{\partial t} = e^\nu \frac{\partial f_{12,10}^{(E)}}{\partial r_*} + \nu' e^{2\nu} f_{12,10}^{(E)} + S_{20}^{(2)}, \quad (4.6)$$

where $S_{20}^{(1)}$ and $S_{20}^{(2)}$ are the source terms describing the coupling of the electromagnetic perturbations with the gravitational ones, and are given by

$$S_{20}^{(1)} = 3Q_{20} \left[\left(\frac{1}{r} B_1 - e^\nu B_2 \right) X_{20} - r B_2 \frac{\partial X_{20}}{\partial r_*} \right], \quad (4.7)$$

$$S_{20}^{(2)} = \frac{3}{2} Q_{20} r e^\nu B_1' X_{20}. \quad (4.8)$$

In order to derive second-order wave-type equations for the electromagnetic perturbations, we introduce a new function $\Psi_{lm} = \Psi_{lm}(t, r)$ given by

$$\Psi_{lm} = e^\nu f_{12,lm}^{(E)}. \quad (4.9)$$

With this variable, the above evolution equations can be written as

$$\frac{\partial \Psi_{10}}{\partial t} = \frac{\partial f_{02,10}^{(E)}}{\partial r_*} - e^\nu f_{01,10}^{(E)}, \quad (4.10)$$

$$\frac{\partial f_{01,10}^{(E)}}{\partial t} = \frac{2}{r^2} \Psi_{10} + S_{20}^{(1)}, \quad (4.11)$$

$$\frac{\partial f_{02,10}^{(E)}}{\partial t} = \frac{\partial \Psi_{10}}{\partial r_*} + S_{20}^{(2)}. \quad (4.12)$$

From this system of evolution equations, one can construct a single wave-type equation for the “electric” perturbations

$$\frac{\partial^2 \Psi_{10}}{\partial t^2} - \frac{\partial^2 \Psi_{10}}{\partial r_*^2} + \frac{2}{r^2} e^\nu \Psi_{10} = S_{20}^{(E)}, \quad (4.13)$$

where the source term $S_{20}^{(E)}$ is given by

$$S_{20}^{(E)} = \frac{\partial S_{20}^{(2)}}{\partial r_*} - e^\nu S_{20}^{(1)}. \quad (4.14)$$

Without the coupling term, this wave equation outside the star is the well-known Regge-Wheeler equation for electromagnetic perturbations. It should be pointed out that Ψ is not a gauge-invariant quantity while the function $\tilde{\Psi}$ given by Eq. (3.12) is a gauge-invariant variable, where both variables Ψ and $\tilde{\Psi}$ can be related to each other via the evolution equation (4.11), i.e.,

$$\frac{\partial \tilde{\Psi}_{10}}{\partial t} = -\Psi_{10} - \frac{r^2}{2} S_{20}^{(1)}. \quad (4.15)$$

Finally, the electromagnetic perturbations in the interior region are determined from Eqs. (3.47), (3.48), and (3.11), i.e.,

$$f_{01,10}^{(E)} = B_2 S_{20}^{(3)}, \quad (4.16)$$

$$f_{02,10}^{(E)} = \frac{1}{2} B_1 S_{20}^{(3)}, \quad (4.17)$$

$$\frac{\partial f_{12,10}^{(E)}}{\partial t} = \frac{\partial f_{02,10}^{(E)}}{\partial r} - f_{01,10}^{(E)}, \quad (4.18)$$

where

$$S_{20}^{(3)} = -3Q_{20} r^2 e^{\nu/2} U_{20}. \quad (4.19)$$

B. Dipole magnetic perturbations driven by polar gravitational perturbations

As it was mentioned earlier in Sec. III for the case of axisymmetric perturbations, the ‘‘magnetic (axial)-type’’ perturbations of the electromagnetic field with harmonic index l are driven by polar gravitational perturbations with harmonic index $l \pm 1$. Here, we consider the axisymmetric perturbations ($m = 0$) for the dipole ($l = 1$) electromagnetic fields, which are driven by quadrupole ($l = 2$) gravitational perturbations.

For the description of the perturbations of the space-time and the stellar fluid, we adopt the formalism derived by Allen *et al.* in Ref. [44]. In this formalism, the perturbations are described by three coupled wave-type equations, in such a way that two equations describe the perturbations of the space-time and the other one the fluid perturbations. In addition to these three wave equations, there is also a

constraint equation. The two wave-type equations for the space-time variables are

$$\begin{aligned} & -\frac{\partial^2 S_{lm}}{\partial t^2} + \frac{\partial^2 S_{lm}}{\partial r_*^2} + \frac{2e^\nu}{r^3} [2\pi r^3 (\rho + 3p) \\ & \quad + m - (n+1)r] S_{lm} \\ & = -\frac{4e^{2\nu}}{r^5} \left[\frac{(m+4\pi p r^3)^2}{r-2m} + 4\pi \rho r^3 - 3m \right] F_{lm}, \end{aligned} \quad (4.20)$$

$$\begin{aligned} & -\frac{\partial^2 F_{lm}}{\partial t^2} + \frac{\partial^2 F_{lm}}{\partial r_*^2} + \frac{2e^\nu}{r^3} [2\pi r^3 (3\rho + p) + m - (n+1)r] F_{lm} \\ & = -2[4\pi r^2 (\rho + p) - e^{-\lambda}] S_{lm} + 8\pi (\rho + p) r e^\nu \left(1 - \frac{1}{C_s^2} \right) H_{lm}, \end{aligned} \quad (4.21)$$

where F_{lm} , S_{lm} , and H_{lm} are given by

$$F_{lm}(t, r) = r K_{lm}, \quad (4.22)$$

$$S_{lm}(t, r) = \frac{e^\nu}{r} (H_{0,lm} - K_{lm}), \quad (4.23)$$

$$H_{lm}(t, r) = \frac{\delta p_{lm}}{\rho + p}, \quad (4.24)$$

while δp_{lm} is the perturbation in the pressure, $n \equiv (l-1)(l+2)/2$, and C_s is the sound speed. On the other hand, the wave equation for the perturbed relativistic enthalpy H_{lm} describing the fluid perturbations is

$$\begin{aligned} & -\frac{1}{C_s^2} \frac{\partial^2 H_{lm}}{\partial t^2} + \frac{\partial^2 H_{lm}}{\partial r_*^2} + \frac{e^{(\nu+\lambda)/2}}{r^2} \left[(m+4\pi p r^3) \left(1 - \frac{1}{C_s^2} \right) + 2(r-2m) \right] \frac{\partial H_{lm}}{\partial r_*} \\ & + \frac{2e^\nu}{r^2} \left[2\pi r^2 (\rho + p) \left(3 + \frac{1}{C_s^2} \right) - (n+1) \right] H_{lm} = (m+4\pi p r^3) \left(1 - \frac{1}{C_s^2} \right) \frac{e^{(\lambda-\nu)/2}}{2r} \left(\frac{e^\nu}{r^2} \frac{\partial F_{lm}}{\partial r_*} - \frac{\partial S_{lm}}{\partial r_*} \right) \\ & + \left[\frac{(m+4\pi p r^3)^2}{r^2(r-2m)} \left(1 + \frac{1}{C_s^2} \right) - \frac{m+4\pi p r^3}{2r^2} \left(1 - \frac{1}{C_s^2} \right) - 4\pi r(3p + \rho) \right] S_{lm} \\ & + \frac{e^\nu}{r^2} \left[\frac{2(m+4\pi p r^3)^2}{r^2(r-2m)} \frac{1}{C_s^2} - \frac{m+4\pi p r^3}{2r^2} \left(1 - \frac{1}{C_s^2} \right) - 4\pi r(3p + \rho) \right] F_{lm}. \end{aligned} \quad (4.25)$$

This third wave equation (4.25) is valid only inside the star, while the first two are simplified considerably outside the star, which can be reduced to a single wave-type equation, i.e., the Zerilli equation (see Ref. [44] and Sec. VB). Finally, the Hamiltonian constraint

$$\begin{aligned} & \frac{\partial^2 F_{lm}}{\partial r_*^2} - \frac{e^{(\nu+\lambda)/2}}{r^2} (m+4\pi p r^3) \frac{\partial F_{lm}}{\partial r_*} + \frac{e^\nu}{r^3} [12\pi r^3 \rho - m - 2(n+1)r] F_{lm} - r e^{-(\nu+\lambda)/2} \frac{\partial S_{lm}}{\partial r_*} \\ & + \left[8\pi r^2 (\rho + p) - (n+3) + \frac{4m}{r} \right] S_{lm} + \frac{8\pi r}{C_s^2} e^\nu (\rho + p) H_{lm} = 0 \end{aligned} \quad (4.26)$$

can be used for setting up initial data and monitoring the evolution of the coupled system.

Regarding the quadrupole gravitational perturbations, the perturbation equation for the ‘‘magnetic-type’’ dipole in the exterior region is obtained from Eq. (3.29) as

$$\frac{\partial^2 \Phi_{10}}{\partial t^2} - \frac{\partial^2 \Phi_{10}}{\partial r_*^2} + \frac{2}{r^2} e^\nu \Phi_{10} = S_{20}^{(M)}, \quad (4.27)$$

where $\Phi_{lm} \equiv f_{23,lm}^{(M)}$ and

$$S_{20}^{(M)} = -Q_{20} e^\nu \left[(2B_2 + rB_2') r^2 S_{20} + \left(e^\nu B_2 + r e^\nu B_2' - \frac{2}{r} B_1 \right) F_{20} + r B_2 \frac{\partial F_{20}}{\partial r_*} \right]. \quad (4.28)$$

In order to derive the wave equation (4.27), we have used Eq. (3.10) and the (r, ϕ) component of the perturbed Einstein equations, i.e., $e^{-\nu} \dot{H}_1 - H_0' + K' - \nu' H_0 = 0$. We remark that the wave equation (4.27) without the source terms is the same as the one derived in Refs. [49,50]. In addition, the other components of the electromagnetic perturbations $f_{12,10}^{(M)}$ and $f_{02,10}^{(M)}$ can be determined with Φ_{10} via the relation (3.10).

Finally, from Eqs. (3.49) and (3.10), we can obtain the equation that determines the dipole ‘‘magnetic-type’’ perturbations for the interior region:

$$\frac{\partial \Phi_{10}}{\partial t} = Q_{20} r^2 e^{\nu/2} (B_2 R_{20} - 3B_1 V_{20}), \quad (4.29)$$

where the perturbations of the fluid velocity, R_{20} and V_{20} , in the source term are given by

$$\begin{aligned} \frac{\partial R_{20}}{\partial t} = e^{\nu/2-\lambda} \left[\left(-\frac{11p+3\rho}{2(p+\rho)} + \frac{3r\nu'}{2} \right) e^{-\nu} S_{20} \right. \\ \left. - \frac{3}{2} r e^{-\nu} S_{20}' + \frac{3p-\rho}{2r^2(p+\rho)} (F_{20} - rF_{20}') - H_{20}' \right], \end{aligned} \quad (4.30)$$

$$\frac{\partial V_{20}}{\partial t} = \frac{1}{2r^2} e^{\nu/2} \left[r e^{-\nu} S_{20} + \frac{\rho-3p}{p+\rho} \frac{F_{20}}{r} - 2H_{20} \right]. \quad (4.31)$$

V. PERTURBATIONS OF DIPOLE MAGNETIC FIELD ON A BLACK HOLE BACKGROUND

The perturbations of a dipole magnetic field on a Schwarzschild black hole background are described by the same set of perturbation equations as in the exterior region of the star except for the boundary conditions, i.e., the boundary conditions for the neutron star imposed on the stellar surface are Eqs. (3.53), (3.54), and (3.55), while for the black hole case one should impose the pure ingoing wave conditions at the event horizon. Then, even in the case of the black hole background, we observe the same coupling of the various harmonics of the electromagnetic and gravitational perturbations as for the neutron star background. That is, for the axisymmetric perturbations, the ‘‘electric’’ dipole ($l=1$) perturbations of the electromagnetic fields will be driven by axial quadrupole ($l=2$) gravitational perturbations, while the ‘‘magnetic’’ dipole ($l=1$) perturbations of the electromagnetic fields will be driven by polar quadrupole ($l=2$) gravitational ones. In

this specific case, our study is similar to the work in Ref. [24], although they use a different formalism.

A. Dipole electric perturbations driven by axial gravitational perturbations (BH)

The axial quadrupole ($l=2$) gravitational perturbations are described by the Regge-Wheeler equation

$$\frac{\partial^2 X_{lm}}{\partial t^2} - \frac{\partial^2 X_{lm}}{\partial r_*^2} + e^\nu \left(\frac{\Lambda}{r^2} - \frac{6M}{r^3} \right) X_{lm} = 0, \quad (5.1)$$

where

$$X_{lm} = \frac{e^\nu}{r} h_{1,lm}. \quad (5.2)$$

In accordance with the results of Sec. IV A, the perturbations of the electromagnetic fields will be described by a single wave equation, that is, the Regge-Wheeler equation for electromagnetic perturbations given by

$$\frac{\partial^2 \Psi_{10}}{\partial t^2} - \frac{\partial^2 \Psi_{10}}{\partial r_*^2} + \frac{2}{r^2} e^\nu \Psi_{10} = S_{20}^{(E)}, \quad (5.3)$$

where the source term becomes of the same form as in Sec. IV A: $\Psi_{lm} = e^\nu f_{12,lm}^{(E)}$.

B. Dipole magnetic perturbations driven by polar gravitational perturbations (BH)

The equation describing the ‘‘magnetic-type’’ perturbations driven by the gravitational perturbations is the same equation as the one derived for a neutron star background [see Eq. (4.27)], that is

$$\frac{\partial^2 \Phi_{10}}{\partial t^2} - \frac{\partial^2 \Phi_{10}}{\partial r_*^2} + \frac{2}{r^2} e^\nu \Phi_{10} = S_{20}^{(M)}, \quad (5.4)$$

where $\Phi_{lm} = f_{23,lm}^{(M)}$, and the source term is also of the same form as in Eq. (4.28). The perturbative equation for the space-time variables can be written in the form of the Zerilli equation

$$\frac{\partial^2 Z_{lm}}{\partial t^2} - \frac{\partial^2 Z_{lm}}{\partial r_*^2} + V_Z(r) Z_{lm} = 0, \quad (5.5)$$

$$V_Z(r) = \frac{2e^\nu [\Lambda_1^2 (\Lambda_1 + 1) r^3 + 3M \Lambda_1^2 r^2 + 9M^2 \Lambda_1 r + 9M^3]}{r^3 (r \Lambda_1 + 3M)^2}, \quad (5.6)$$

where $\Lambda_1 \equiv (l+2)(l-1)/2$. Meanwhile, in the same way as for the neutron star background, one can also adopt F_{lm} and S_{lm} as the perturbation variables for the space-time. In this case, the two wave equations simplify to become

$$\begin{aligned} \frac{\partial^2 S_{lm}}{\partial t^2} - \frac{\partial^2 S_{lm}}{\partial r_*^2} + e^\nu \left(\frac{\Lambda}{r^2} - \frac{2M}{r^3} \right) S_{lm} \\ = -\frac{4M}{r^5} e^\nu \left(3 - \frac{7M}{r} \right) F_{lm}, \end{aligned} \quad (5.7)$$

$$\frac{\partial^2 F_{lm}}{\partial t^2} - \frac{\partial^2 F_{lm}}{\partial r_*^2} + e^\nu \left(\frac{\Lambda}{r^2} - \frac{2M}{r^3} \right) F_{lm} = -2e^\nu S_{lm}, \quad (5.8)$$

which have to be supplemented with the Hamiltonian constraint equation

$$\begin{aligned} \frac{\partial^2 F_{lm}}{\partial r_*^2} - \frac{M}{r^2} \frac{\partial F_{lm}}{\partial r_*} - \frac{\Lambda}{r^2} e^\nu F_{lm} - r \frac{\partial S_{lm}}{\partial r_*} \\ - \frac{1}{2} (4e^\nu + \Lambda) S_{lm} = 0. \end{aligned} \quad (5.9)$$

Note that there are useful relations between the perturbation variables (F_{lm} , S_{lm}) and the Zerilli function (Z), i.e.,

$$F_{lm} = r \frac{dZ_{lm}}{dr_*} + \frac{\Lambda_1(\Lambda_1 + 1)r^2 + 3\Lambda_1 M r + 6M^2}{r(\Lambda_1 r + 3M)} Z_{lm}, \quad (5.10)$$

$$\begin{aligned} S_{lm} = \frac{1}{r} \frac{dF_{lm}}{dr_*} - \frac{(\Lambda_1 + 2)r - M}{r^3} F_{lm} \\ + \frac{(\Lambda_1 + 1)(\Lambda_1 r + 3M)}{r^3} Z_{lm}, \end{aligned} \quad (5.11)$$

which can be used in constructing initial data (since the Zerilli function is gauge invariant and unconstrained), or for the extraction of the Zerilli function.

VI. APPLICATIONS

As an application, we consider the case in which dipole “electric-type” perturbations are driven by axial gravitational ones and present numerical results. First, we study the coupling on a Schwarzschild black hole background and later on the background of spherical neutron stars, as discussed in Secs. VA and IVA, respectively. The more complicated cases that involve the driving of “magnetic-type” electromagnetic field perturbations driven by polar gravitational ones will be discussed elsewhere in the future.

A. Perturbations on a black hole background

In order to calculate the waveforms in the black hole background, we need to modify the background magnetic field near the event horizon. The reason for this is that the solution for a dipole magnetic field in vacuum diverges at the event horizon [see Eqs. (2.15) and (2.16)]. In fact, the isolated black hole cannot have magnetic fields due to

the no-hair theorem. But, according to the simulations of the accretion onto the black hole, the magnetic field can reach almost to the event horizon, because the accreting matter will fall into the black hole with infinite time [51,52]. Thus, we adopt a simple modification of the dipole magnetic field near the event horizon, that is, we set $B_1(r) = B_1(6M)$ and $B_2(r) = B_2(6M)$ for $r \leq 6M$, where the position at $r = 6M$ corresponds to the innermost stable circular orbit for a test particle around the Schwarzschild black hole. The magnetic dipole moment μ_d is identified with the normalized magnetic field strength B_{15} defined as $B_{15} \equiv B_p / (10^{15}[G])$, where B_p is the field strength at $r = 6M$ and $\theta = 0$.

We assume vanishing electromagnetic perturbations, i.e., $\Psi_{10} = \partial\Psi_{10}/\partial t = 0$ at the initial time slice $t = 0$, while the initial gravitational perturbations X_{20} are prescribed in terms of a Gaussian wave packet. Under these initial conditions, the electromagnetic waves will result from the coupling to the gravitational ones. In the numerical calculations, we adopt the iterated Crank-Nicholson method [53] with a grid choice of $\Delta r_* = 0.1M$ and $\Delta t = \Delta r_*/2$ (see Ref. [54] for the dependence of the choice of Δr_* and Δt on the waveforms).

The energy emitted in the form of either gravitational (E_{GW}) or electromagnetic waves (E_{EM}) is estimated by integrating the luminosity ($L_{\text{GW},l}^{(A)}$) for the axial gravitational waves and for electric-type electromagnetic waves ($L_{\text{EM},l}^{(E)}$), which are described by the following formulas [45,46]:

$$L_{\text{GW},l}^{(A)} = \frac{1}{16\pi} \frac{(l-2)!}{(l+2)!} \left| \frac{\partial X_{l0}}{\partial t} \right|^2, \quad (6.1)$$

$$L_{\text{EM},l}^{(E)} = \frac{1}{4\pi} \frac{(l+1)!}{(l-1)!} \left| \frac{\partial \Psi_{l0}}{\partial t} \right|^2. \quad (6.2)$$

In practice, we can find a relation between the energy emitted in gravitational and electromagnetic waves for a given initial space-time perturbation, which has the form

$$E_{\text{EM}} = \alpha B_{15}^2 E_{\text{GW}}, \quad (6.3)$$

where α is a “proportionality constant.”

In the simulation that we describe we set the magnetic field strength to the value $B_p = 10^{15}$ gauss. Figure 1 shows




FIG. 1 (color online). Gravitational waveform observed at $r = 2000M$. In the right panel, we also show the absolute value of X_{20} .




FIG. 2 (color online). Waveform of the driven electromagnetic waves observed at $r = 2000M$ for $B_p = 10^{15}$ gauss. In the right panel, we also show the absolute value of Ψ_{10} .




FIG. 3 (color online). FFT of the electromagnetic waves shown in Fig. 2. The two vertical lines correspond to the frequencies of quasinormal modes for $l = 1$ electromagnetic waves (dashed line) and for $l = 2$ gravitational waves (dot-dash line).

the waveform of the gravitational wave observed at $r = 2000M$, the amplitude is normalized to correspond to an emitted energy of $E_{\text{GW}} \approx 1.8 \times 10^{49} (M/50M_{\odot})$ ergs. On the other hand, the waveforms of electromagnetic waves driven by the gravitational waves are shown in Fig. 2. From this figure, we can observe somewhat complicated waveforms of electromagnetic waves due to the coupling with the gravitational waves. From the specific waveforms, one can estimate the value of the proportionality constant in the relation [Eq. (6.3)] to be $\alpha = 8.02 \times 10^{-6}$. This efficiency might not be very high, but the radiated energy of gravitational waves can reach $\sim 10^{51}$ ergs for a black hole formation due to the merger of a neutron stars binary (see, e.g., Ref. [55]). In this case the strength of the

magnetic field can be amplified by the Kelvin-Helmholtz instability to reach values of the order of 10^{15-17} gauss [56]. Although this is not an ideal situation for the black hole case we are considering in this paper, if one adopts the above efficiency for the case of a black hole formed after merger, one can expect that energies of the order of $\sim 10^{46-50}$ ergs can be emitted in the form of electromagnetic waves which can be potentially driven by the gravitational field perturbations.

Furthermore, in Fig. 3, we show the fast Fourier transform (FFT) of the electromagnetic waveforms shown in Fig. 2, where for comparison we also add the frequencies of the quasinormal modes for $l = 1$ electromagnetic waves (dashed line) and for $l = 2$ gravitational waves (dot-dash line) radiated from the Schwarzschild black hole [45]. From this figure, one can obviously see that the driven electromagnetic waves have two specific frequencies corresponding to the $l = 1$ quasinormal mode of electromagnetic waves and the $l = 2$ quasinormal mode of gravitational waves. This means that it might be possible to see the effect of gravitational waves via observation of electromagnetic waves. However, electromagnetic waves with such low frequencies could be coupled/absorbed by the interstellar medium (and/or accretion disk around the central object) during the propagation and it will be almost impossible to directly detect the driven electromagnetic waves. The only possible way to see the driven electromagnetic waves is the observation of indirect effects, such as synchrotron radiation.




FIG. 4 (color online). Waveforms of the gravitational waves (solid line) and the electromagnetic waves (dotted line) for the stellar model with $B_p = 10^{15}$ gauss, which are observed at $r = 300$ km. The left and right panels correspond to different stellar models for EOS with $\Gamma = 2$ and $K = 200 \text{ km}^2$.

TABLE I. Stellar parameters adopted in this article.

| Γ | K | ρ_c (g/cm ³) | M/M_\odot | R (km) | M/R |
|----------|-----|-------------------------------|-------------|----------|-------|
| 2 | 100 | 1.0×10^{15} | 0.802 | 10.8 | 0.109 |
| 2 | 100 | 1.5×10^{15} | 0.998 | 10.2 | 0.145 |
| 2 | 100 | 2.0×10^{15} | 1.126 | 9.67 | 0.172 |
| 2 | 100 | 3.0×10^{15} | 1.266 | 8.86 | 0.211 |
| 2 | 200 | 0.7×10^{15} | 1.365 | 14.6 | 0.138 |
| 2 | 200 | 0.9×10^{15} | 1.528 | 14.0 | 0.162 |
| 2 | 200 | 1.0×10^{15} | 1.592 | 13.7 | 0.172 |
| 2 | 200 | 1.5×10^{15} | 1.791 | 12.5 | 0.211 |
| 2 | 200 | 2.0×10^{15} | 1.876 | 11.7 | 0.237 |
| 2.25 | 600 | 1.0×10^{15} | 0.732 | 9.69 | 0.111 |
| 2.25 | 600 | 1.5×10^{15} | 1.008 | 9.44 | 0.158 |
| 2.25 | 600 | 2.0×10^{15} | 1.197 | 9.12 | 0.194 |
| 2.25 | 600 | 3.0×10^{15} | 1.404 | 8.48 | 0.245 |
| 2.25 | 600 | 4.0×10^{15} | 1.486 | 7.95 | 0.276 |

B. Perturbations on a neutron star background

In order to examine the coupling between the emitted gravitational and electromagnetic waves in a neutron star background, we adopt the same initial conditions as for the black hole case, i.e., the electromagnetic perturbations are set to zero and the initial gravitational perturbations are approximated by an ingoing Gaussian wave packet. In the numerical calculations, we adopt a grid spacing of $\Delta r = R/200$ and a time step $\Delta t/\Delta r = 0.05$, where R is the stellar radius. For the background stellar models, we adopt the polytropic equation of state (EOS) of the form $P = K\rho^\Gamma$. Then, one can get the waveforms of the reflected gravitational waves and the induced electromagnetic ones.

As an example, we show results for a stellar mode with $\Gamma = 2$ and $K = 200 \text{ km}^2$. Figure 4 shows the waveforms of the gravitational waves (solid line) and the electromagnetic waves (dotted line) observed at $r = 300 \text{ km}$, where we adopt two stellar models with different compactness M/R (see Table I for the stellar properties). Compared with the fast damping of gravitational waves, one can see the long-term oscillations in the electromagnetic waves, which can be driven not only by the quasinormal ringing of gravitational waves but also during the tail phase of the

gravitational waves. For the waveforms shown in Fig. 4, the FFT is plotted in Fig. 5, where the left and right panels correspond to the FFT of the gravitational and electromagnetic waves, respectively. From this figure, one can see the same features as in the case of a black hole. Namely, the FFT of the electromagnetic waves driven by the gravitational waves has two specific frequencies, i.e., one is the proper electromagnetic oscillation (first peak in the right panel of Fig. 5) and the other one is the oscillation corresponding to the gravitational waves (second peak in the right panel of Fig. 5). We remark that electromagnetic waves with such low frequencies could be absorbed by the interstellar medium and then, their direct detection is almost impossible. Namely, we should consider the secondary emission mechanism such as a synchrotron radiation. Maybe the plasma around the central object will be excited after receiving the energy from the electromagnetic waves driven by the gravitational waves and move along with the magnetic field lines. Anyway, such a secondary emission mechanism will be discussed somewhere. Furthermore, we find that as in the case for a black hole, the relationship between the emitted energies of gravitational and electromagnetic waves can be described by Eq. (6.3), even for neutron stars, if B_p is considered as the magnetic field strength at the stellar pole ($r = R$ and $\theta = 0$). In practice, for the specific stellar models in Fig. 4, the proportionality constant becomes $\alpha = 1.61 \times 10^{-5}$ and 4.37×10^{-6} for the particular stellar models with $M/R = 0.162$ and 0.237 , respectively.

In order to see the dependence on the stellar properties, we study a variety of stellar models with different stiffness of the EOS and with different central densities, radii, and masses, which are given in Table I. As a result, we find that the proportionality constant α can be written as a function of the stellar compactness, which is almost independent of the stellar models and the adopted EOS. In fact, in Fig. 6 we plot the values of α for various stellar models, where the circles, diamonds, and squares correspond to the results for the stellar models characterized by $(\Gamma, K) = (2, 100)$, $(2, 200)$, and $(2.25, 600)$. From this figure, one can see that the proportionality constant α depends strongly on the stellar compactness, as expected, with typical values ranging from 10^{-6} up to $\sim 10^{-4}$.




FIG. 5 (color online). FFT of the gravitational waves (left panel) and electromagnetic waves (right panel) shown in Fig. 4.




FIG. 6 (color online). The proportionality constant α as a function of the stellar compactness for various polytropic models. The circles, diamonds, and squares correspond to stellar models with $(\Gamma, K) = (2,100)$, $(2,200)$, and $(2.25,600)$.

VII. CONCLUSION

We have considered the coupling between gravitational and electromagnetic waves emitted by compact objects, i.e., black holes and neutron stars. We have derived a coupled system of equations describing the propagation of gravitational and electromagnetic waves. In our study we have investigated the driving of electromagnetic perturbations via their coupling to the gravitational ones. However, for simplicity, we have neglected the backreaction from the electromagnetic waves on the gravitational waves, because the magnetic energy of the compact objects, even for magnetars, was quite small as compared with the gravitational energy. We found that the electromagnetic waves of specific parity with harmonic indices (l, m) can be coupled to gravitational waves of the same parity and with harmonic indices (l, m) (for $m \neq 0$) and harmonic indices $(l \pm 1, m)$, for every value of m . In particular, our findings led to the result that, for the axisymmetric perturbations, i.e., $m = 0$, the dipole electric electromagnetic waves will be driven by axial quadrupole gravitational waves, while the dipole magnetic electromagnetic waves will be driven by polar gravitational waves.

As an application of our perturbative framework, we presented numerical calculations for the case in which dipole-electric electromagnetic waves were driven by the axial gravitational ones, both for the case of a black hole

and a neutron star background. We found that the emitted energy in electromagnetic waves driven by the gravitational waves was proportional to not only the emitted energy in gravitational waves but also to the square of the strength of the magnetic field of the central object. For the case of a black hole background, the ratio of the emitted energy of the electromagnetic waves to that of the gravitational waves was around $8 \times 10^{-6} (B_p/10^{15} \text{ G})^2$, where B_p was the magnetic field strength at $r = 6M$. On the other hand, in the case of a neutron star background, we found that this proportionality constant can be written as a function of the stellar compactness.

Although we have considered only the case of axial gravitational waves and the associated induced electromagnetic waves, the polar oscillations also play an important role in extracting the information about the neutron star structure since in the case of nonrotating stars, the matter oscillations are typically coupled to the polar gravitational waves. This is a direction that we are currently investigating.

ACKNOWLEDGMENTS

H. S. is grateful to Ken Ohsuga for valuable comments. This work was supported by the German Science Foundation support SFB/TR7 ‘‘Gravitational Wave Astronomy,’’ by Grants-in-Aid for Scientific Research on Innovative Areas through Grants No. 23105711, No. 24105001, and No. 24105008 provided by MEXT, by Grant-in-Aid for Young Scientists (B) through Grant No. 24740177 provided by JSPS, by the Yukawa International Program for Quark-Hadron Sciences, and by the Grant-in-Aid for the global COE program ‘‘The Next Generation of Physics, Spun from Universality and Emergence’’ from MEXT. C.F.S. acknowledges support from Contracts No. FIS2008-06078-C03-03, No. AYA-2010-15709, and No. FIS2011-30145-C03-03 of the Spanish Ministry of Science and Innovation, and Contract No. 2009-SGR-935 of AGAUR (Generalitat de Catalunya). P.L. acknowledges the support from NSF Grants No. 1205864, No. 0903973, and No. 0941417.

-
- [1] S. Márka (LIGO Scientific Collaboration and Virgo Collaboration), *Classical Quantum Gravity* **28**, 114013 (2011).
 - [2] N.L. Christensen (LIGO Scientific Collaboration and Virgo Collaboration), [arXiv:1105.5843](https://arxiv.org/abs/1105.5843).
 - [3] P. Ghosh, *Rotation and Accretion Powered Pulsars* (World Scientific, Singapore, 2007).
 - [4] A.L. Watts and T.E. Strohmayer, *Adv. Space Res.* **40**, 1446 (2007).
 - [5] K. Glampedakis, L. Samuelsson, and N. Andersson, *Mon. Not. R. Astron. Soc.* **371**, L74 (2006).
 - [6] Y. Levin, *Mon. Not. R. Astron. Soc.* **377**, 159 (2007).
 - [7] H. Sotani, K.D. Kokkotas, and N. Stergioulas, *Mon. Not. R. Astron. Soc.* **375**, 261 (2007).
 - [8] H. Sotani, K.D. Kokkotas, and N. Stergioulas, *Mon. Not. R. Astron. Soc.* **385**, L5 (2008).
 - [9] A. Colaiuda, H. Beyer, and K.D. Kokkotas, *Mon. Not. R. Astron. Soc.* **396**, 1441 (2009).

- [10] P. C erda-Dur an, N. Stergioulas, and J. A. Font, *Mon. Not. R. Astron. Soc.* **397**, 1607 (2009).
- [11] H. Sotani, *Mon. Not. R. Astron. Soc.* **417**, L70 (2011).
- [12] M. van Hoven and Y. Levin, *Mon. Not. R. Astron. Soc.* **410**, 1036 (2011).
- [13] M. Gabler, P. C erda-Dur an, J. A. Font, E. M uller, and N. Stergioulas, *Mon. Not. R. Astron. Soc.* **421**, 2054 (2012).
- [14] A. Colaiuda and K. D. Kokkotas, *Mon. Not. R. Astron. Soc.* **423**, 811 (2012).
- [15] H. Sotani, K. Nakazato, K. Iida, and K. Oyamatsu, *Phys. Rev. Lett.* **108**, 201101 (2012).
- [16] H. Sotani, K. Nakazato, K. Iida, and K. Oyamatsu, *Mon. Not. R. Astron. Soc.* **428**, L21 (2013).
- [17] M. van Hoven and Y. Levin, *Mon. Not. R. Astron. Soc.* **420**, 3035 (2012).
- [18] L. P. Grishchuk and A. G. Polnarev, *General Relativity and Gravitation*, edited by A. Held (Plenum Press, New York, 1980), Vol. 2, p. 416.
- [19] D. Papadopoulos, N. Stergioulas, L. Vlahos, and J. Kuijpers, *Astron. Astrophys.* **377**, 701 (2001).
- [20] M. Servin and G. Brodin, *Phys. Rev. D* **68**, 044017 (2003).
- [21] J. Moortgat and J. Kuijpers, *Astron. Astrophys.* **402**, 905 (2003).
- [22] J. Moortgat and J. Kuijpers, *Phys. Rev. D* **70**, 023001 (2004).
- [23] M. Forsberg, G. Brodin, M. Marklund, P. K. Shukla, and J. Moortgat, *Phys. Rev. D* **74**, 064014 (2006).
- [24] C. A. Clarkson, M. Marklund, G. Betschart, and P. K. S. Dunsby, *Astrophys. J.* **613**, 492 (2004).
- [25] B. F. Schutz, *Introduction to General Relativity* (Cambridge University Press, Cambridge, England, 1985).
- [26] I. Wasserman and S. L. Shapiro, *Astrophys. J.* **265**, 1036 (1983).
- [27] M. Bocquet, S. Bonazzola, E. Gourgoulhon, and J. Novak, *Astron. Astrophys.* **301**, 757 (1995).
- [28] K. Konno, T. Obata, and Y. Kojima, *Astron. Astrophys.* **352**, 211 (1999).
- [29] A. Colaiuda, V. Ferrari, L. Gualtieri, and J. A. Pons, *Mon. Not. R. Astron. Soc.* **385**, 2080 (2008).
- [30] S. Bonazzola, E. Gourgoulhon, M. Salgado, and J. A. Marck, *Astron. Astrophys.* **278**, 421 (1993).
- [31] C. Kouveliotou *et al.*, *Nature (London)* **393**, 235 (1998).
- [32] K. Hurley *et al.*, *Nature (London)* **397**, 41 (1999).
- [33] G. L. Israel, T. Belloni, L. Stella, Y. Rephaeli, D. E. Gruber, P. Casella, S. Dall’Osso, N. Rea, M. Persic, and R. E. Rothschild, *Astrophys. J.* **628**, L53 (2005).
- [34] P. D. Lasky, B. Zink, K. D. Kokkotas, and K. Glampedakis, *Astrophys. J.* **735**, L20 (2011).
- [35] R. Ciolfi, S. K. Lander, G. M. Manca, and L. Rezzolla, *Astrophys. J.* **736**, L6 (2011).
- [36] B. Zink, P. D. Lasky, and K. D. Kokkotas, *Phys. Rev. D* **85**, 024030 (2012).
- [37] P. Lasky, B. Zink, and K. D. Kokkotas, [arXiv:1203.3590](https://arxiv.org/abs/1203.3590).
- [38] R. Ciolfi and L. Rezzolla, *Astrophys. J.* **760**, 1 (2012).
- [39] T. Regge and J. A. Wheeler, *Phys. Rev.* **108**, 1063 (1957).
- [40] K. S. Thorne and A. Campolattaro, *Astrophys. J.* **149**, 591 (1967).
- [41] F. Zerilli, *Phys. Rev. D* **2**, 2141 (1970).
- [42] K. D. Kokkotas and B. F. Schutz, *Mon. Not. R. Astron. Soc.* **255**, 119 (1992).
- [43] Y. Kojima, *Phys. Rev. D* **46**, 4289 (1992).
- [44] G. Allen, N. Andersson, K. D. Kokkotas, and B. F. Schutz, *Phys. Rev. D* **58**, 124012 (1998).
- [45] C. M. Cunningham, R. H. Price, and V. Moncrief, *Astrophys. J.* **224**, 643 (1978).
- [46] C. M. Cunningham, R. H. Price, and V. Moncrief, *Astrophys. J.* **230**, 870 (1979).
- [47] S. Chandrasekhar and V. Ferrari, *Proc. R. Soc. A* **432**, 247 (1991).
- [48] K. D. Kokkotas, *Mon. Not. R. Astron. Soc.* **268**, 1015 (1994).
- [49] H. Sotani, S. Yoshida, and K. D. Kokkotas, *Phys. Rev. D* **75**, 084015 (2007).
- [50] H. Sotani, *Phys. Rev. D* **79**, 084037 (2009).
- [51] A. Tchekhovskoy, R. Narayan, and J. C. McKinney, *Mon. Not. R. Astron. Soc.* **418**, L79 (2011).
- [52] J. C. McKinney, A. Tchekhovskoy, and R. D. Blandford, *Mon. Not. R. Astron. Soc.* **423**, 3083 (2012).
- [53] S. A. Teukolsky, *Phys. Rev. D* **61**, 087501 (2000).
- [54] H. Sotani and M. Saijo, *Phys. Rev. D* **74**, 024001 (2006).
- [55] M. Shibata, K. Taniguchi, and K. Uryu, *Phys. Rev. D* **68**, 084020 (2003).
- [56] D. J. Price and S. Rossweg, *Science* **312**, 719 (2006).