

The a theorem for gauge-Yukawa theories beyond Banks-Zaks fixed pointOleg Antipin,^{*} Marc Gillioz,[†] Esben Mølgaard,[‡] and Francesco Sannino[§]*CP³-Origins and the Danish Institute for Advanced Study DIAS, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark*

(Received 19 March 2013; published 10 June 2013)

We investigate the a theorem for nonsupersymmetric gauge-Yukawa theories beyond the leading order in perturbation theory. The exploration is first performed in a model-independent manner and then applied to a specific relevant example. Here, a rich fixed point structure appears including the presence of a merging phenomenon between nontrivial fixed points for which the a theorem has not been tested so far.

DOI: [10.1103/PhysRevD.87.125017](https://doi.org/10.1103/PhysRevD.87.125017)

PACS numbers: 11.15.Bt, 11.10.Gh, 11.15.-q

I. INTRODUCTION

The recent work by Komargodski and Schwimmer [1,2] on the a theorem has attracted much interest. The a theorem is thought to be the generalization to four dimensions of Zamolodchikov's c theorem [3] which establishes the irreversibility of the renormalization group (RG) flow in two dimensions. Several extensions of the work of Zamolodchikov to four dimensions have been proposed. In particular, Cardy pointed to the function a given by the integral of the trace of the energy-momentum tensor over the four-sphere [4]. He then conjectured that a has properties similar to its two-dimensional cousin c . Cardy's conjecture was tested within the framework of perturbation theory in [5–7]. It was discovered that the a function is not monotonically decreasing along the RG flow, and an alternative function, denoted here by \tilde{a} , was devised to be monotonically decreasing in perturbation theory. A nice property of \tilde{a} is that it coincides with Cardy's a function at fixed points, where the β functions vanish.

Komargodski and Schwimmer suggested a nonperturbative proof of the a theorem by introducing a dilatonic field [1,2] whose scattering amplitude is related to the a function. In fact, the analyticity of the amplitude supports the monotonicity of the \tilde{a} function (not a) along the RG flow.¹ Establishing the existence of a function that is monotonic along the RG flow can lead to relevant constraints on the dynamics of a given gauge theory. In supersymmetry, the existing relation between a and the R -charge [10,11] found use in the a -maximization [12]. Using holographic methods, the a theorem could be tested and verified in the context of supersymmetry [13,14]. The generalization, using holography, to arbitrary space-time dimensions was explored in [15]. More recently, the relation between scale

and conformal invariance in four dimensions was elucidated in the context of the a theorem [16,17].

The goal of this work is to extend the perturbative analyses of the a theorem for nonsupersymmetric gauge theories with fermions and gauge singlet scalars, interacting via Yukawas, to the maximum known order in perturbation theory. This allows us to investigate the details of the a theorem, particularly for nonstandard fixed point structures.

The paper is structured as follows. In Sec. II we introduce the essential tools and discuss the generic expression of the \tilde{a} function relevant for its determination to three loops in the gauge coupling. Concrete examples are presented in Sec. III where we explicitly evaluate \tilde{a} at the Banks-Zaks (BZ) fixed point (FP) for a gauge-Yukawa theory to the leading and the next-to-leading order in perturbation theory. We then discover, for a certain non asymptotically free gauge theory, the appearance of a perturbative UV fixed point, and at the next-to-leading order also the appearance of another infrared (IR) fixed point. By changing the number of matter fields we observe the merging and disappearance of both fixed points. This rich structure of fixed points constitutes an interesting playground for elucidating the properties of the \tilde{a} function in gauge theories. We conclude in Sec. IV and provide a number of technical details in the Appendix.

II. THE a THEOREM BEYOND THE LEADING ORDER

In four dimensions and for a general quantum field theory the vacuum expectation value of the trace of the energy-momentum tensor for a locally flat metric $g_{\mu\nu}$ reads

$$\langle T_{\mu}^{\mu} \rangle = cW^2(g_{\mu\nu}) - aE_4(g_{\mu\nu}) + \dots, \quad (1)$$

where a and c are real coefficients, $E_4(g_{\mu\nu})$ the Euler density and $W(g_{\mu\nu})$ the Weyl tensor. The dots represent contributions coming from operators that can be constructed out of the fields defining the theory. Their contribution is proportional to the β functions of their couplings. The coefficient a is the one used in Cardy's conjecture, and for a free field theory it is [18]

^{*} antipin@cp3.dias.sdu.dk[†] gillioz@cp3.dias.sdu.dk[‡] molgaard@cp3.dias.sdu.dk[§] sannino@cp3.dias.sdu.dk¹The relation between the trace anomaly and the amplitude of the three point function of the energy-momentum tensor was already pointed out in [8,9].

$$a_{\text{free}} = \frac{1}{90(8\pi)^2} \left(n_s + \frac{11}{2} n_f + 62 n_v \right), \quad (2)$$

where n_s , n_f and n_v are, respectively, the number of real scalars, Weyl fermions and gauge bosons.

The change of a along the RG flow is directly related to the underlying dynamics of the theory via the β functions. This can be shown by exploiting the Abelian nature of the trace anomaly which leads to the Weyl consistency conditions in much the same manner as the well known Wess-Zumino consistency conditions [19]. As discussed in the introduction and following the work of Jack and Osborn [5,6], rather than using a one uses the function \tilde{a} related to it by

$$\tilde{a} = a + W_i \beta_i, \quad (3)$$

where W_i is a one-form which depends on the couplings of the theory. The Weyl consistency conditions² imply for \tilde{a}

$$\partial_i \tilde{a} = -\chi_{ij} \beta_j + (\partial_i W_j - \partial_j W_i) \beta_j, \quad (4)$$

where χ_{ij} can be viewed as a metric in the space of couplings. The positivity of the metric χ is established in perturbation theory, and therefore in this regime the function \tilde{a} is monotonic along the RG flow

$$\mu \frac{d\tilde{a}}{d\mu} = \beta_i \partial_i \tilde{a} = -\chi_{ij} \beta_i \beta_j \leq 0. \quad (5)$$

The irreversibility of the RG flow has been conjectured to be valid beyond perturbation theory.

In the next subsections, capitalizing on the work of Jack and Osborn [6], we will construct \tilde{a} for gauge theories featuring fermionic matter interacting with gauge singlet scalar fields via Yukawa interactions to the highest known order in perturbation theory.

A. Gauge-Yukawa theories

We consider the following Lagrangian skeleton:

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4g^2} F_{\mu\nu} F^{\mu\nu} + i \Psi_i^\dagger \sigma^\mu D_\mu \Psi_i + \frac{1}{2} \partial_\mu \phi_a \partial^\mu \phi_a \\ & - (y_{ij}^a \Psi_i^c \Psi_j \phi_a + \text{H.c.}) - \frac{1}{4!} \lambda_{abcd} \phi_a \phi_b \phi_c \phi_d, \end{aligned} \quad (6)$$

$$\chi = \begin{pmatrix} \frac{\chi_{gg}}{\alpha_g^2} (1 + A\alpha_g + B_1\alpha_g^2 + B_2\alpha_g\alpha_y) & B_0 & 0 \\ B_0 & \frac{\chi_{yy}}{\alpha_y} (1 + B_3\alpha_y + B_4\alpha_g) & 0 \\ 0 & 0 & \chi_{\lambda\lambda} \end{pmatrix}. \quad (12)$$

The coefficient χ_{gg} enters at the one-loop order, A and χ_{yy} at two loops, while $\chi_{\lambda\lambda}$ and the B_i 's appear only at three loops. Similarly, the one-form W takes the form

where we dropped the gauge indices for $F_{\mu\nu}$ and the Weyl fermions Ψ_i . The fermions transform according to a given representation R of the underlying gauge group. The real scalars ϕ are singlets with respect to the gauge group. The indices run over the number of matter fields. To exemplify our results we will consider gauge theories for which the Yukawa and quartic interactions depend each on a single parameter as follows:

$$y_{ij}^a \equiv y T_{ij}^a, \quad \lambda_{abcd} \equiv \lambda T_{abcd}, \quad (7)$$

where the T are tensors specified in a given theory. There are therefore three couplings in our setup: gauge g , Yukawa y and quartic λ . We define

$$\alpha_g = \frac{g^2}{(4\pi)^2}, \quad \alpha_y = \frac{y^2}{(4\pi)^2}, \quad \alpha_\lambda = \frac{\lambda}{(4\pi)^2}. \quad (8)$$

The generic structure of the associated β functions ($\beta_\alpha = \partial\alpha/\partial \ln \mu$) reads³

$$\beta_{\alpha_g} = -2\alpha_g^2 [b_0 + b_1\alpha_g + b_y\alpha_y + b_2\alpha_g^2 + b_3\alpha_g\alpha_y + b_4\alpha_y^2], \quad (9)$$

$$\begin{aligned} \beta_{\alpha_y} = & 2\alpha_y [c_1\alpha_y + c_2\alpha_g + c_3\alpha_g\alpha_y + c_4\alpha_g^2 + c_5\alpha_y^2 \\ & + c_6\alpha_y\alpha_\lambda + c_7\alpha_\lambda^2], \end{aligned} \quad (10)$$

$$\beta_{\alpha_\lambda} = d_1\alpha_\lambda^2 + d_2\alpha_\lambda\alpha_y + d_3\alpha_y^2. \quad (11)$$

The expansion to three loops in the gauge coupling, two loops in the Yukawa and one in the quartic coupling leads to a consistent expression for \tilde{a} to order α^3 . If the scalars were charged under the gauge group, terms proportional to $\alpha_g\alpha_\lambda$ would appear.

Having established the generic form of the β functions, we move to determining the metric χ and one-form W . They can be found by examining the relevant Feynman diagrams which enter the computation of the trace anomaly, as shown in the Appendix. We find

$$\begin{aligned} W_g = & \frac{1}{\alpha_g} (D_0 + D_1\alpha_g + C_1\alpha_g^2 + C_2\alpha_g\alpha_y), \\ W_y = & D_2 + C_3\alpha_y + C_4\alpha_g, \quad W_\lambda = D_3\alpha_\lambda. \end{aligned} \quad (13)$$

²It is worth mentioning that the Weyl consistency conditions used above assume that the trace of the energy-momentum tensor vanishes when all the β functions are zero simultaneously. Exceptions are known to exist [20] and in this case one modifies the consistency conditions [16,17] in order to build \tilde{a} .

³The factors of 2 in the definition of the gauge and Yukawa β functions follow from the definitions (8). One has for example $\beta_{\alpha_g}/\alpha_g = 2\beta_g/g$.

The general structure of χ confirms that it is sufficient for all our purposes to consider the Yukawa β function (10) to two-loop order and the quartic one (11) to one-loop only.

The leading coefficients χ_{gg} , χ_{yy} and $\chi_{\lambda\lambda}$ are [6]

$$\begin{aligned}\chi_{gg} &= \frac{d(G)}{128\pi^2}, & \chi_{yy} &= \frac{1}{128\pi^2} \left(\frac{1}{3} T_{ij}^a T_{ij}^{a*} \right), \\ \chi_{\lambda\lambda} &= \frac{1}{128\pi^2} \left(\frac{1}{72} T_{abcd} T_{abcd} \right),\end{aligned}\quad (14)$$

where we used the tensors T defined in Eq. (7) and $d(G)$ denotes the dimension of the adjoint representation G of the underlying gauge group, i.e., the number of gluons. A is given by [6]

$$A = 17C_2(G) - \frac{10}{3} N_R T(R), \quad (15)$$

where $C_2(G)$ is the quadratic Casimir of the adjoint, N_R the number of Weyl fermion in the representation R and $T(R)$ is the trace normalization satisfying $T(R)\delta_{ab} = \text{Tr}(R_a R_b)$, R_a being the generators of the fermions under the gauge group. With these coefficients, the metric χ is positive definite near the origin of the coupling constant space. It is however clear that in the absence of a theorem, the positivity of χ away from the origin is not guaranteed. The remaining coefficients of χ and W are yet to be determined but, as we shall show, they are not needed to determine \tilde{a} at the fixed points to the order investigated here.

B. Power of the consistency relations and the \tilde{a} function

The system of first order differential equations in (4) allows to derive the following conditions relating the different coefficients of the β functions as well as χ and W :

$$\begin{aligned}\chi_{gg} b_y &= -\chi_{yy} c_2, & \chi_{yy} c_6 &= \chi_{\lambda\lambda} d_3, & 4\chi_{yy} c_7 &= \chi_{\lambda\lambda} d_2, \\ 2\chi_{gg} b_4 + \chi_{yy} (B_4 c_1 + B_3 c_2 + c_3) &= 2(B_0 - C_2 + C_4) c_1, \\ \chi_{gg} (B_2 b_0 + A b_y + b_3) + 2\chi_{yy} (B_4 c_2 + c_4) &= (B_0 - C_2 + C_4) c_2 + 2(B_0 + C_2 - C_4) b_0.\end{aligned}\quad (16)$$

The equations in the first line above can be used to either test or determine some of the higher order coefficients of the β functions since we know the metric factors. The remaining equations can be used in a similar fashion. However, given that the B_i coefficients have not been explicitly computed we use the knowledge of the β functions to deduce, for example, B_3 and B_4 assuming that c_2 does not vanish.

For \tilde{a} to cubic order in the couplings, and using the consistency relations above, we have

$$\tilde{a} = a_{\text{free}} + \tilde{a}^{(1)} + \tilde{a}^{(2)} + \tilde{a}^{(3)} + \dots, \quad (17)$$

where a_{free} is the free-field theory value (2), and the one-, two- and three-loop coefficients are

$$\tilde{a}^{(1)} = -2\chi_{gg} b_0 \alpha_g, \quad (18)$$

$$\tilde{a}^{(2)} = -\chi_{gg} (b_1 + A b_0) \alpha_g^2 - 2\chi_{gg} b_y \alpha_g \alpha_y + \chi_{yy} c_1 \alpha_y^2, \quad (19)$$

$$\begin{aligned}\tilde{a}^{(3)} &= -\chi_{gg} \left[\frac{2}{3} (b_2 + A b_1) \alpha_g^3 + (b_3 + A b_y) \alpha_g^2 \alpha_y \right. \\ &\quad \left. + 2b_4 \alpha_g \alpha_y^2 + \frac{1}{3} \frac{c_1}{c_2} \left(4b_4 - \frac{c_1}{c_2} (b_3 + A b_y) \right) \alpha_y^3 \right] \\ &\quad + \chi_{yy} \left[\frac{2}{3} \left(c_5 - \frac{c_1}{c_2} c_3 + \left(\frac{c_1}{c_2} \right)^2 c_4 \right) \alpha_y^3 + c_6 \alpha_y^2 \alpha_\lambda \right. \\ &\quad \left. + 2c_7 \alpha_y \alpha_\lambda^2 \right] + \frac{1}{3} \chi_{\lambda\lambda} a_1 \alpha_\lambda^3 + \frac{\beta_{\alpha_g}}{\alpha_g^2} f(\alpha_i^3) \\ &\quad + \frac{\beta_{\alpha_y}^2}{\alpha_y} \frac{B_0 - C_2 + C_4}{4c_2},\end{aligned}\quad (20)$$

where we defined

$$\begin{aligned}f(\alpha_i^3) &= \chi_{gg} \left(\frac{B_1}{3} \alpha_g^3 + \frac{B_2}{2} \alpha_g^2 \alpha_y - \frac{B_2}{6} \left(\frac{c_1}{c_2} \right)^2 \alpha_y^3 \right) \\ &\quad + \frac{B_0 + C_2 - C_4}{3} \left(\frac{c_1}{c_2} \right)^2 \alpha_y^3.\end{aligned}\quad (21)$$

Remarkably, the unknown coefficients B_i and C_i appear only in the last two terms of $\tilde{a}^{(3)}$, where they are multiplied by β functions and hence vanish at fixed points.⁴ This was also observed to occur in supersymmetric theories [21].

It is instructive to calculate a to the second order using (3) and recalling that the leading coefficients entering the one-form W are $D_0 = \chi_{gg}$, $D_1 = \frac{1}{2} A \chi_{gg}$, and $D_2 = \frac{1}{2} \chi_{yy}$ [6]. We find the simple expression

$$a = a_{\text{free}} + \chi_{gg} (b_1 \alpha_g^2 + b_y \alpha_g \alpha_y) + \mathcal{O}(\alpha_i^3), \quad (22)$$

where, remarkably, the term linear in α_g , the one quadratic in α_y as well as the term linear in A canceled out. Because the signs of b_1 and b_y depend on the gauge theory, a is not generally a monotonically decreasing function along the perturbative RG flow.

C. \tilde{a} at fixed points

We can now move to the study of the fixed points and determine the variation of \tilde{a} between two of them. A convenient way to search for the zeros of the system of β functions (9)–(11) is to first solve analytically for $\beta_{\alpha_\lambda} = 0$ which permits to relate α_λ to α_y , then set to zero β_{α_y} further relating α_y to α_g , so that finally we can search for the zeros of the following effective β function in α_g :

⁴We have taken the liberty of adding higher order terms in order to rewrite the coefficients as β functions. These terms are irrelevant when computing \tilde{a} between perturbative fixed points.

$$\beta_{\alpha_g^{\text{eff}}}^{\text{eff}} = -2\alpha_g^2 [b_0 + b_1^{\text{eff}} \alpha_g + b_2^{\text{eff}} \alpha_g^2], \quad (23)$$

where

$$b_1^{\text{eff}} = b_1 - \frac{c_2}{c_1} b_y, \quad (24)$$

$$b_2^{\text{eff}} = b_2 - \frac{c_2}{c_1} b_3 + \left(\frac{c_1}{c_2}\right)^2 b_4 - \frac{b_y}{c_1} \left[c_4 - \frac{c_2}{c_1} c_3 + \left(\frac{c_2}{c_1}\right)^2 c_5^{\text{eff}} \right], \quad (25)$$

with

$$c_5^{\text{eff}} = c_5 - \frac{d_3}{d_1} c_7 - \frac{d_2}{2d_1} \left(c_6 - \frac{d_2}{d_1} c_7 \right) \left(1 \pm \sqrt{1 - \frac{4d_1 d_3}{d_2^2}} \right). \quad (26)$$

At this order in perturbation theory, there can be at most two perturbative fixed points for each sign in c_5^{eff} , if both b_0 and b_1^{eff} are tuned to be small. An example of this is provided in the following section. Using Eqs. (24) and (25), the difference in the function \tilde{a} —or equivalently a —between the UV and IR fixed points can then be written as

$$\begin{aligned} \Delta \tilde{a}_{\text{perturbative}} &\equiv (\tilde{a}^{\text{UV}} - \tilde{a}^{\text{IR}})_{\text{perturbative}} \\ &= -2\chi_{gg} \left[b_0 (\alpha_g^{\text{UV}} - \alpha_g^{\text{IR}}) \right. \\ &\quad + \frac{1}{2} (b_1^{\text{eff}} + Ab_0) ((\alpha_g^{\text{UV}})^2 - (\alpha_g^{\text{IR}})^2) \\ &\quad \left. + \frac{1}{3} (b_2^{\text{eff}} + Ab_1^{\text{eff}} + Bb_0) ((\alpha_g^{\text{UV}})^3 - (\alpha_g^{\text{IR}})^3) \right], \end{aligned} \quad (27)$$

where α_g^{UV} and α_g^{IR} denote the values of the gauge coupling at the UV and IR fixed point, respectively, and we defined

$$B \equiv B_1 - \frac{c_2}{c_1} \left(B_2 + \frac{B_0}{\chi_{gg}} \right). \quad (28)$$

This expression reduces to the case of a gauge theory without Yukawa interactions by replacing b_1^{eff} and b_2^{eff} with b_1 and b_2 .

Inspecting the effective β function, there is a perturbative fixed point for small b_0 , which reads

$$\alpha_g^{\text{BZ}} = -\frac{b_0}{b_1^{\text{eff}}} + \mathcal{O}(b_0^2). \quad (29)$$

For positive b_0 and negative b_1^{eff} , this is the usual Banks-Zaks IR fixed point. The situation in which the BZ fixed point is of UV nature is equally possible. This occurs by reverting the signs of both b_0 and b_1^{eff} . The trivial fixed point at the origin will be in the first case an UV fixed point and in the second an IR one. The finite change in \tilde{a} between the UV and IR fixed points can be computed either way, and one obtains

$$\Delta \tilde{a}^{\text{BZ}} = \mp \chi_{gg} \frac{b_0^2}{b_1^{\text{eff}}}. \quad (30)$$

Here the sign reflects the sign of b_0 , and $\Delta \tilde{a}$ is then positive for any physical fixed point. However, $\Delta \tilde{a}$ can formally become negative when the value of the coupling α_g at the fixed point is on the unphysical negative axis.

To three-loop order in the effective β function, one can have the two following physical zeros:

$$\begin{aligned} \alpha_g^{\text{BZ}} &= -\frac{b_1^{\text{eff}}}{2b_2^{\text{eff}}} \left(1 - \sqrt{1 - \frac{4b_0 b_2^{\text{eff}}}{(b_1^{\text{eff}})^2}} \right), \\ \alpha_g^{\text{BZ}} &= -\frac{b_1^{\text{eff}}}{2b_2^{\text{eff}}} \left(1 + \sqrt{1 - \frac{4b_0 b_2^{\text{eff}}}{(b_1^{\text{eff}})^2}} \right). \end{aligned} \quad (31)$$

For small values of b_0 , the solution with negative sign corresponds to the usual BZ fixed point, with the following corrections:

$$\alpha_g^{\text{BZ}} = -\frac{b_0}{b_1^{\text{eff}}} \left(1 + \frac{b_0 b_2^{\text{eff}}}{(b_1^{\text{eff}})^2} + \mathcal{O}(b_0^2) \right). \quad (32)$$

This expression holds provided $b_0/(b_1^{\text{eff}})^2$ is small. We shall see below that there are cases where this is not true. Using (32), we can compute the three-loop corrections to the variation of \tilde{a} ,

$$\Delta \tilde{a}^{\text{BZ}} = \mp \chi_{gg} \frac{b_0^2}{b_1^{\text{eff}}} \left(1 - (Ab_1^{\text{eff}} - 2b_2^{\text{eff}}) \frac{b_0}{3(b_1^{\text{eff}})^2} \right). \quad (33)$$

We now turn our attention to the second zero α_g^{BZ} . The first observation is that for a generic value of b_1^{eff} this fixed point occurs at a nonperturbative value of the coupling. This is what happens in general for gauge theories with fermionic matter in a given irreducible representation of the gauge group [22]. However, for gauge theories with Yukawa interactions and/or multiple matter representations, the possibility that both b_0 and b_1^{eff} are small exists. An explicit example is provided below. Furthermore, when $b_2^{\text{eff}} = (b_1^{\text{eff}})^2/(4b_0)$ the two fixed points merge. This phenomenon can happen within the range of perturbation theory. At the merger, one has $\alpha_{\text{merger}} = -2b_0/b_1^{\text{eff}}$, which, when plugged into Eq. (27), gives

$$\Delta \tilde{a}^{\text{BZ}}|_{\text{merger}} = \mp \chi_{gg} \frac{4}{3} \frac{b_0^2}{b_1^{\text{eff}}}. \quad (34)$$

The virtues of this expression will be studied in more detail elsewhere [23].

Having in our hands the explicit tools, we can explore the a theorem for gauge theories with interesting fixed point structures.

III. A CONCRETE EXAMPLE

We consider a $SU(N_c)$ gauge theory with N_f fundamental Dirac fermions $Q = (q, \tilde{q}^*)$, ℓ adjoint Weyl fermions λ , and a gauge singlet complex scalar H that transforms in the bifundamental representation of the $SU(N_f)_L \times SU(N_f)_R$ global symmetry of the theory. For the benefit of the reader, the field content and the quantum symmetries of the theory with $\ell = 1$ are summarized in Table I. The Lagrangian of the theory is

$$\mathcal{L} = \text{Tr} \left[-\frac{1}{2} F^{\mu\nu} F_{\mu\nu} + i\bar{\lambda}\not{D}\lambda + \bar{Q}i\not{D}Q + \partial_\mu H^\dagger \partial^\mu H + y_H \bar{Q}HQ \right] - u_1 (\text{Tr}[H^\dagger H])^2 - u_2 \text{Tr}(H^\dagger H)^2. \quad (35)$$

Tr is the trace over both color and flavor indices. D_μ is the usual covariant derivative. Throughout this section we will work with the rescaled couplings which enable a finite

Veneziano limit of the theory with ℓ fixed. That is, we let $N_c, N_f \rightarrow \infty$ while keeping $x \equiv N_f/N_c$ fixed. The appropriately rescaled couplings are

$$a_g = \frac{g^2 N_c}{(4\pi)^2}, \quad a_H = \frac{y_H^2 N_c}{(4\pi)^2}, \quad (36)$$

$$z_1 = \frac{u_1 N_f^2}{(4\pi)^2}, \quad z_2 = \frac{u_2 N_f}{(4\pi)^2}.$$

This model was introduced in [24] to investigate near-conformal dynamics, at the one-loop level, and its impact on the spectrum of the theory with special attention to the dilaton properties. The model was further investigated at the two-loop level in [25]. To compute \tilde{a} , following the previous section, we need to determine the three-loop contribution to the gauge β function.

Using [26–31] we find in the Veneziano limit

$$\beta_{a_g} = -\frac{2}{3} a_g^2 \left[11 - 2\ell - 2x + (34 - 16\ell - 13x)a_g + 3x^2 a_H + \frac{81x^2}{4} a_g a_H - \frac{3x^2(7+6x)}{4} a_H^2 + \frac{2857 + 112x^2 - x(1709 - 257\ell) - 1976\ell + 145\ell^2}{18} a_g^2 \right], \quad (37)$$

$$\beta_{a_H} = a_H \left[2(x+1)a_H - 6a_g + (8x+5)a_g a_H + \frac{20(x+\ell) - 203}{6} a_g^2 - 8xz_2 a_H - \frac{x(x+12)}{2} a_H^2 + 4z_2^2 \right], \quad (38)$$

$$\beta_{z_2} = 2(2z_2 a_H + 4z_2^2 - x a_H^2). \quad (39)$$

Here one can see that the double trace coupling z_1 does not participate in the running of the remaining couplings. In addition, using (14) and (15) the metric coefficients for this theory can be found:

$$\chi_{gg} = \frac{N_c^2}{2^7 \pi^2}, \quad \chi_{yy} = \frac{N_f^2}{3 \cdot 2^7 \pi^2}, \quad (40)$$

$$\chi_{\lambda\lambda} = \frac{N_f^2}{3 \cdot 2^6 \pi^2}, \quad A = 17 - \frac{10}{3}(x + \ell).$$

TABLE I. Field content of the example. The first three fields are Weyl spinors in the $(\frac{1}{2}, 0)$ representation of the Lorentz group. H is a complex scalar and G_μ is the gauge field. $U(1)_{\text{AF}}$ is the extra anomaly free symmetry arising due to the presence of λ .

Fields	$[SU(N_c)]$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_V$	$U(1)_{\text{AF}}$
λ	Adj	1	1	0	1
q	\square	$\bar{\square}$	1	$\frac{N_f - N_c}{N_c}$	$-\frac{N_c}{N_f}$
\tilde{q}	$\bar{\square}$	1	\square	$-\frac{N_f - N_c}{N_c}$	$-\frac{N_c}{N_f}$
H	1	\square	$\bar{\square}$	0	$\frac{2N_c}{N_f}$
G_μ	Adj	1	1	0	0

One can check that the expressions above satisfy the consistency relations given in the first line of (16), and therefore it constitutes an independent check of the correctness of the β functions. We now turn to the FP analysis of the model which will reveal an interesting perturbative structure.

A. Leading order analysis: Banks-Zaks fixed point

In order to see a physical BZ fixed point, the one-loop coefficient of the gauge β function has to be small and the signs of b_0 and b_1^{eff} have to be opposite. Therefore, our first task is to find a region in the parameter space of the model where the physical BZ fixed point exists. We use Eq. (24)

$$b_0 = \frac{1}{3}(11 - 2(\ell + x)),$$

$$b_1^{\text{eff}} = \frac{1}{3} \left(34 - 16\ell - 13x + \frac{9x^2}{(x+1)} \right). \quad (41)$$

From the asymptotic freedom (AF) boundary condition $b_0 = 0$ we obtain that $x = (11 - 2\ell)/2$. Substituting this value of x into b_1^{eff} , we have

$$b_{1\text{AF}}^{\text{eff}} = -\frac{25}{2} - \ell - \frac{3(11 - 2\ell)^2}{4\ell - 26}, \quad (42)$$




FIG. 1 (color online). (a) $\Delta\tilde{a}$ between the Gaussian and BZ fixed points normalized to χ_{gg} at leading order for the $\ell = 1$ case. The solid red (dashed blue) line corresponds to the model without (with) Yukawa interactions. In both cases the physical BZ FP is an IR one. (b) $\Delta\tilde{a}$ between the Gaussian and BZ fixed points normalized to χ_{gg} at leading order for the $\ell = 0$ case. In the absence (presence) of Yukawa interactions, the physical BZ fixed point is an IR (UV) one. The color code is the same as on the left panel.

where the last term comes from the Yukawa interactions. We immediately notice that for $\ell^* \approx 0.37$ the b_{1AF}^{eff} vanishes. Below, we will consider the cases $\ell = 1$ for which this coefficient is negative and $\ell = 0$ for which it is positive. In the first case we have a standard IR BZ fixed point, and in the second we obtain a new UV BZ fixed point. It is worth noticing that, in the absence of Yukawa interactions, b_{1AF}^{eff} in (42) is always negative and therefore the physical BZ FP can only be the standard IR fixed point.

1. $\ell = 1$ case

In this case there exists a perturbative IR fixed point regardless of whether we consider the presence of Yukawa interactions. We show in Fig. 1(a) the leading order result for the change in \tilde{a} between the Gaussian (trivial) FP and the BZ IR one at leading order, both in the presence (dashed blue line) and absence (solid red line) of Yukawa interactions. Both curves cross zero at $x^* = 9/2$, when asymptotic freedom is lost. For $x > 9/2$, where $b_0 < 0$, there is an unphysical BZ UV fixed point yielding a negative $\Delta\tilde{a}$. The Yukawa interactions in (42) imply $|b_1^{\text{eff}}| < |b_1|$, which leads to a larger $\Delta\tilde{a}$ in the case of the gauge theory with scalars.

2. $\ell = 0$ case

We now turn to the $\ell = 0$ case where, rather than having a BZ IR fixed point, the theory develops a UV fixed point when asymptotic freedom is lost, i.e., for $b_0 < 0$. Of course, this is possible only because of the presence of the Yukawa interactions. In Fig. 1(b), the leading order result for the change in \tilde{a} between the Gaussian and BZ fixed points without Yukawa interactions is shown in red (solid line), and the one with Yukawa interactions in blue (dashed line). Both curves cross zero for $x^* = 11/2$ when asymptotic freedom is lost.

B. Next-to-leading order analysis: Fixed point merger

At the next perturbative order we deal with the full system of Eqs. (37)–(39) and from now on, we concentrate only on the physical fixed points.

1. $\ell = 1$ case

We start again with the $\ell = 1$ theory and in Fig. 2(a) we display the FPs structure for the model with Yukawa and quartic interactions. We notice that at $x \approx 3.25$ the FP value of the gauge coupling vanishes. However, this happens in the region beyond applicability of perturbation theory since the two remaining coupling constants are large. In Fig. 2(b) we plot the change in the \tilde{a} function for the next-to-leading order BZ IR fixed point and compare it with the corresponding leading order results from Fig. 1(a). As a general feature, we notice that the next-to-leading order corrections reduce the value of $\Delta\tilde{a}$ in the perturbative regime. It is clear from the plots that for the theory with Yukawa interactions perturbation theory breaks down, when moving away from the critical value $x^* = 4.5$, earlier with respect to the theory without Yukawa interactions.

2. $\ell = 0$ case

We now turn to the $\ell = 0$ theory where, as discussed above, there is no BZ IR fixed point. However, in the asymptotically free regime there is a new physical IR fixed point emerging at the next-to-leading order. This non-BZ IR fixed point is present also when asymptotic freedom is lost, i.e., for $x > 5.5$. In this region there is also a BZ UV fixed point that was discussed in Sec. III A 2. The complete FP structure is shown in Fig. 3(a). The UV and IR fixed points merge around $x \approx 5.6$. In Fig. 3(b) we plot the change in the \tilde{a} function at next-to-leading order together with the corresponding leading order results from Fig. 1(b). We notice that $\Delta\tilde{a}$ becomes negative just before the merger which is incompatible with the a theorem. We interpret this effect as the breakdown of the perturbative expansion since the FP values of the couplings at the merger are quite large, as can be seen from Fig. 3(a).

So far, all our calculations of $\Delta\tilde{a}$ were for the flow connecting the trivial FP at the origin of the coupling constant space with the BZ one. However, it is relevant




FIG. 2 (color online). (a) The next-to-leading order physical FP structure for the $\ell = 1$ case with Yukawa and quartic interactions. (b) $\Delta\tilde{a}$ normalized to χ_{gg} for the $\ell = 1$ case. The solid red and dashed blue lines are leading order results from Fig. 1(a) while the dotted black and dot-dashed green lines are the next-to-leading order corrections.




FIG. 3 (color online). (a) The next-to-leading order physical FP structure for the $\ell = 0$ case with Yukawa and quartic interactions. The vertical dot-dashed green line represents the point where asymptotic freedom is lost. (b) $\Delta\tilde{a}$ normalized to χ_{gg} for the $\ell = 0$ case. The solid red and dashed blue lines are leading order results from Fig. 1(b) while the dotted black and dot-dashed green lines are next-to-leading order corrections.

also to determine $\Delta\tilde{a}$ for the branch connecting the two non-trivial FPs. In the theory with $\ell = 0$ and $x > 5.5$ this is the RG flow between the BZ UV fixed point and the non-BZ IR one. We display the change in the \tilde{a} function in Fig. 4(a). Of course, at the merger $\Delta\tilde{a}$ vanishes.

3. $\ell = 0.35$ case: The perturbative merger

Using both ℓ and x as continuous parameters it is formally possible to study the merging phenomenon within the perturbative regime. This happens around $\ell \approx 0.37$. Therefore we provide an example with $\ell = 0.35$. The change in the \tilde{a}




FIG. 4 (color online). (a) $\Delta\tilde{a}$ normalized to χ_{gg} for the RG flow between the BZ UV fixed point and the non-BZ IR fixed point. (b) $\Delta\tilde{a}$ normalized to χ_{gg} for the $\ell = 0.35$ case. The solid magenta curve corresponds to the flow between the BZ UV and the non-BZ IR fixed points. The dot-dashed green curve is the result between the Gaussian IR fixed point and the UV BZ one.








function for the two RG flows are shown in Fig. 4(b). Since for this value of ℓ perturbation theory holds, we observe a positive and well behaved $\Delta\tilde{a}$ all the way to the merger.

IV. CONCLUSIONS

We studied the a theorem to the maximum known order in perturbation theory for nonsupersymmetric gauge

theories with fermionic matter as well as gauge singlet scalar fields, where the latter two interact via Yukawa interactions. The computation involves three loops in the gauge β function, two loops in the Yukawa interactions and one loop in the quartic interactions. To this order we have first determined the general expression for the change in the function \tilde{a} between two fixed points, and then specialized it to the case of QCD with extra adjoint Weyl

TABLE II. One-, two- and three-loop vacuum polarization diagrams entering the computation of the metric (12) and the one-form (13).

Diagrams	Contributions to χ	Contributions to W
	$\frac{\beta_{\alpha_g}^2}{\alpha_g^2}$	$\frac{\beta_{\alpha_g}}{\alpha_g}$
	$\frac{\beta_{\alpha_g}^2}{\alpha_g}$	β_{α_g}
	$\frac{\beta_{\alpha_y}^2}{\alpha_y}$	β_{α_y}
	$\beta_{\alpha_g}^2$	$\alpha_g \beta_{\alpha_g}$
	$\beta_{\alpha_y}^2$	$\alpha_y \beta_{\alpha_y}$
	$\frac{\beta_{\alpha_g}^2}{\alpha_g} \alpha_y, \frac{\beta_{\alpha_y}^2}{\alpha_y} \alpha_g, \beta_{\alpha_g} \beta_{\alpha_y}$	$\alpha_y \beta_{\alpha_g}, \alpha_g \beta_{\alpha_y}$
	$\beta_{\alpha_\lambda}^2$	$\alpha_\lambda \beta_{\alpha_\lambda}$

fermions and elementary mesons interacting via Yukawa terms. We employed the Veneziano limit and determined the three-loop β functions of the theory. This has allowed us to test the a theorem beyond the lowest order in perturbation theory. We discovered that the model possesses an interesting structure of the fixed points, depending on the number of adjoint fermionic matter fields, among which we highlight the presence of a Banks-Zaks UV fixed point as well as the occurrence of a merging phenomenon for which the a theorem had not been tested before.


ACKNOWLEDGMENTS

The CP³-Origins centre is partially funded by the Danish National Research Foundation, Grant

No. DNRF90. We thank John Gracey for clarifying comments regarding the calculation of the three-loop gauge β function. O.A. thanks the Galileo Galilei Institute for Theoretical Physics for the hospitality and the INFN for partial support during the completion of this work.

APPENDIX: DIAGRAMMATIC FORM OF THE TRACE ANOMALY

The structure of the metric (12) and one-form (13) entering the trace anomaly can be determined by looking at vacuum polarization diagrams containing in addition to the usual vertices of the quantum field theory the following counterterms:



The terms entering the metric (12) are the ones proportional to two powers of the β functions while the ones with one power of β_i fix the one-form W (13). All the vacuum polarization diagrams up to three-loop order as well as the form of their contribution are shown in Table II. Note that more diagrams would be present if the scalar field were charged under the gauge group.

-
- [1] Z. Komargodski and A. Schwimmer, *J. High Energy Phys.* **12** (2011) 099.
 - [2] Z. Komargodski, *J. High Energy Phys.* **07** (2012) 069.
 - [3] A. B. Zamolodchikov, *JETP Lett.* **43**, 730 (1986).
 - [4] J. L. Cardy, *Phys. Lett. B* **215**, 749 (1988).
 - [5] H. Osborn, *Phys. Lett. B* **222**, 97 (1989).
 - [6] I. Jack and H. Osborn, *Nucl. Phys.* **B343**, 647 (1990).
 - [7] H. Osborn, *Nucl. Phys.* **B363**, 486 (1991).
 - [8] A. Cappelli, G. D'Appollonio, R. Guida, and N. Magnoli, *Proc. Sci.*, tnr2000 (2000) 005 [arXiv:hep-th/0009119].
 - [9] A. Cappelli, R. Guida, and N. Magnoli, *Nucl. Phys.* **B618**, 371 (2001).
 - [10] D. Anselmi, D. Freedman, M. T. Grisaru, and A. Johansen, *Nucl. Phys.* **B526**, 543 (1998).
 - [11] D. Anselmi, J. Erlich, D. Freedman, and A. Johansen, *Phys. Rev. D* **57**, 7570 (1998).
 - [12] K. A. Intriligator and B. Wecht, *Nucl. Phys.* **B667**, 183 (2003).
 - [13] D. Z. Freedman, S. S. Gubser, K. Pilch, and N. P. Warner, *Adv. Theor. Math. Phys.* **3**, 363 (1999).
 - [14] L. Girardello, M. Petrini, M. Porrati, and A. Zaffaroni, *J. High Energy Phys.* **12** (1998) 022.
 - [15] R. C. Myers and A. Sinha, *J. High Energy Phys.* **01** (2011) 125.
 - [16] J.-F. Fortin, B. Grinstein, and A. Stergiou, *J. High Energy Phys.* **01** (2013) 184.
 - [17] M. A. Luty, J. Polchinski, and R. Rattazzi, *J. High Energy Phys.* **01** (2013) 152.
 - [18] M. Duff, *Nucl. Phys.* **B125**, 334 (1977).
 - [19] J. Wess and B. Zumino, *Phys. Lett.* **37B**, 95 (1971).
 - [20] J.-F. Fortin, B. Grinstein, and A. Stergiou, *J. High Energy Phys.* **12** (2012) 112.
 - [21] D. Freedman and H. Osborn, *Phys. Lett. B* **432**, 353 (1998).
 - [22] C. Pica and F. Sannino, *Phys. Rev. D* **83**, 035013 (2011).
 - [23] O. Antipin, M. Gillioz, and F. Sannino, arXiv:1303.1547.
 - [24] O. Antipin, M. Mojaza, and F. Sannino, *Phys. Lett. B* **712**, 119 (2012).
 - [25] O. Antipin, S. Di Chiara, M. Mojaza, E. Molgaard, and F. Sannino, *Phys. Rev. D* **86**, 085009 (2012).
 - [26] M. E. Machacek and M. T. Vaughn, *Nucl. Phys.* **B222**, 83 (1983).
 - [27] M. E. Machacek and M. T. Vaughn, *Nucl. Phys.* **B236**, 221 (1984).
 - [28] M. E. Machacek and M. T. Vaughn, *Nucl. Phys.* **B249**, 70 (1985).
 - [29] A. Pickering, J. Gracey, and D. Jones, *Phys. Lett. B* **510**, 347 (2001).
 - [30] L. N. Mihaila, J. Salomon, and M. Steinhauser, *Phys. Rev. D* **86**, 096008 (2012).
 - [31] R. V. Harlander, L. Mihaila, and M. Steinhauser, *Eur. Phys. J. C* **63**, 383 (2009).