

Anti-BRST symmetry and background field methodD. Binosi^{1,*} and A. Quadri^{2,†}¹*European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT*) and Fondazione Bruno Kessler, Villa Tambosi, Strada delle Tabarelle 286, I-38123 Villazzano (TN), Italy*²*Dipartimento di Fisica, Università degli Studi di Milano, via Celoria 16, I-20133 Milano, Italy and INFN, Sezione di Milano, via Celoria 16, I-20133 Milano, Italy*

(Received 17 September 2013; published 28 October 2013)

We show that the requirement that a $SU(N)$ Yang-Mills action (gauge fixed in a linear covariant gauge) is invariant under both the Becchi-Rouet-Stora-Tyutin (BRST) symmetry as well as the corresponding anti-BRST symmetry, automatically implies that the theory is quantized in the (linear covariant) background field method (BFM) gauge. Thus, the BFM and its associated background Ward identity naturally emerge from anti-BRST invariance of the theory and need not be introduced as an *ad hoc* gauge fixing procedure. Treating ghosts and antighosts on an equal footing, as required by a BRST-anti-BRST invariant formulation of the theory, gives also rise to a local antighost equation that together with the local ghost equation completely resolves the algebraic structure of the ghost sector for any value of the gauge fixing parameter. We finally prove that the background fields are stationary points of the background effective action obtained when the quantum fields are integrated out.

DOI: [10.1103/PhysRevD.88.085036](https://doi.org/10.1103/PhysRevD.88.085036)

PACS numbers: 11.15.Tk, 12.38.Aw

I. INTRODUCTION

Quantization of gauge theories in the presence of background field configurations [1–12] is known to be a very useful tool, for it allows one to preserve gauge invariance with respect to external background sources after a gauge-fixing choice has been made for the quantum gauge modes of the theory.

This has led to a number of applications, both at the perturbative level—ranging from calculations in Yang-Mills theories [11,13] via the quantization of the standard model [14] to gravity and supergravity calculations [15]—as well as at the nonperturbative level—where the method has been instrumental in devising a gauge invariant truncation scheme for the Schwinger-Dyson equations of Yang-Mills theories [16–19].

A common algebraic framework has emerged over the years in order to tame the dependence of the vertex functional Γ on the background fields. In [20] it was first proposed to introduce a BRST partner Ω for the background field \hat{A} . The corresponding extended Slavnov-Taylor (ST) identity guarantees that the physics (described by the cohomology of the linearized ST operator) is not affected by the introduction of the background. This approach allows one to acquire an algebraic control over the renormalization of the theory under scrutiny [21,22] and to prove the so-called background equivalence theorem [23,24].

Eventually it was recognized in a series of papers [25–27] that the full dependence on the background field, fixed by the extended ST identity, is induced through a canonical transformation with respect to (w.r.t.) the

Batalin-Vilkovisky (BV) bracket of the theory. Such a canonical transformation is generated by the functional $\delta\Gamma/\delta\Omega$. Since the latter is in general background dependent, one cannot obtain the finite form of the canonical transformation by simple exponentiation; rather, one needs to resort to the Lie series of an appropriate functional differential operator. The derivation proceeds in close analogy with the case of parameter-dependent canonical transformations in classical mechanics [27].

In this paper this simple geometrical interpretation of the background field method will be pushed one step further, as a very deep connection between the BFM and the so-called anti-BRST symmetry will be unveiled.

Indeed, since the advent of BRST quantization, it has been known that a further symmetry exists, induced by an anti-BRST transformation [28–30] in which the antighost field takes the place of the ghost in the variation of the gauge and matter fields of the theory. Though this symmetry turned out to be a useful tool for constraining possible terms in the action and in simplifying relations between Green's functions, it has however been far from clear if there is any case in which it is indispensable, and thus its meaning has remained so far somewhat mysterious.

The anti-BRST symmetry generates an anti-ST identity, that can be shown to hold together with the ST identity induced by the BRST transformations. Moreover, the requirement of simultaneous BRST and anti-BRST invariance can be fulfilled, so that both identities hold true for the vertex functional Γ , provided that a suitable set of operators for the BRST-anti-BRST variation of the fields is introduced through the coupling to appropriate external classical sources. As we will see *the latter sources coincide precisely with the background fields introduced in the BFM*. Thus, for example, the BRST partner Ω of the

*binosi@ectstar.eu

†andrea.quadri@mi.infn.it

background gauge field \hat{A} is seen to be the antifield of the anti-BRST transformation $\bar{s}A$ for the gauge field A , while \hat{A} is the source coupled to the BRST-anti-BRST variation $s\bar{s}A$ of A .

Similar identifications hold true for all the other scalar and fermionic matter fields of the theory. In particular, we will show that for spontaneously broken gauge theories the procedure automatically yields the correct background 't Hooft gauge fixing.

In addition, in a BRST-anti-BRST invariant theory both a local ghost and a local antighost equation exist. While the former equation has been known for a long time, the latter has been derived up to now only in the background Landau gauge [31]. However, once the sources required to establish the ST and the anti-ST identities are introduced, it can be readily seen that there is nothing special in the Landau gauge choice and one can indeed construct a local antighost equation valid for a general R_ξ gauge. The usual background Ward identity arises then both as a consequence of the validity of the ST identity and the local antighost equation, as well as of the anti-ST identity and the local ghost equation.

As our analysis reveals that the background field method is naturally encoded in every theory which is both BRST and anti-BRST invariant, one might wonder whether the ST and the anti-ST identities impose some physical condition on the backgrounds. For that purpose, it is convenient to construct an effective action $\tilde{\Gamma}$ for the background fields, where the quantum modes have been fully integrated out, i.e., one keeps connected graphs with external backgrounds only (and therefore $\tilde{\Gamma}$ is one-particle reducible w.r.t. the quantum fields). This approach is motivated by several applications of the BFM, e.g., in the effective field theory of the color glass condensate [32–34].

As we will see, the ST and the anti-ST identities imply then that the background field configurations are a stationary point for $\tilde{\Gamma}$. This leads to some interesting combinatorial relations of a novel type between one-particle irreducible (1-PI) graphs involving external background sources, whose origin can be ultimately traced back to the canonical transformation of [27], which dictates the dependence of Γ on the backgrounds.

The paper is organized as follows. In Sec. II we discuss the conventional and the background (R_ξ) gauge fixing for a pure SU(N) Yang-Mills theory and derive the associated extended ST identity. In Sec. III the anti-BRST symmetry is introduced together with the sources required to define the composite operators of the BRST-anti-BRST algebra. The equivalence between the BRST-anti-BRST invariance and the BFM is then established. Next, Sec. IV is dedicated to the derivation of the local antighost equation in a generic R_ξ gauge. In addition, we show how the background Ward identity emerges from both the anti-ST identity combined with the local ghost equation or the ST identity combined with the local antighost equation. The local ghost and

antighost equations are then exploited to fully constrain the ghost two-point sector in any gauge. In Sec. V we finally construct the background effective action by integrating out all quantum fields, and show that the background fields are stationary points of this action. Our conclusions are presented in Sec. VII, with the following Appendix generalizing (some of) the main equations for the case in which scalars and fermion fields are present.

II. CONVENTIONAL AND BFM GAUGE FIXINGS

The action of a SU(N) Yang-Mills theory reads

$$S = S_{\text{YM}} + S_{\text{GF}} + S_{\text{FPG}}. \quad (2.1)$$

S_{YM} represents the Yang-Mills (gauge invariant) action, which is written in terms of the SU(N) field strength

$$S_{\text{YM}} = -\frac{1}{4g^2} \int d^4x F_{\mu\nu}^a F_a^{\mu\nu}; \quad (2.2)$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c.$$

S_{GF} and S_{FPG} represent respectively the (covariant) gauge-fixing functional and its associated Faddeev-Popov ghost term. The most general way of writing these terms is through the expressions

$$S_{\text{GF}} = \int d^4x \left[-\frac{\xi}{2} (b^a)^2 + b^a \mathcal{F}^a \right]; \quad (2.3)$$

$$S_{\text{FPG}} = - \int d^4x \bar{c}^a s \mathcal{F}^a.$$

In the formulas above \mathcal{F} represents the gauge-fixing function, which, for the class of R_ξ gauges considered throughout this paper, reads

$$\mathcal{F}^a = \partial^\mu A_\mu^a. \quad (2.4)$$

In addition, b are auxiliary, nondynamical fields (the so-called Nakanishi-Lautrup multipliers) that can be eliminated through their equations of motion, as a consequence of the validity of the b equation

$$\frac{\delta \Gamma}{\delta b^a} = -\xi b^a + \mathcal{F}^a, \quad (2.5)$$

for the full quantum effective action Γ . c (respectively, \bar{c}) are the ghost (respectively, antighost) fields, while, finally, s is the nilpotent BRST operator, which constitutes a symmetry of the gauge-fixed action Eq. (2.1), with the BRST transformations of the various fields given by

$$sA_\mu^a = \mathcal{D}_\mu^{ab} c^b; \quad s c^a = -\frac{1}{2} f^{abc} c^b c^c; \quad (2.6)$$

$$s\bar{c}^a = b^a; \quad s b^a = 0,$$

and the covariant derivative \mathcal{D} is defined according to

$$\mathcal{D}_\mu^{ab} = \partial_\mu \delta^{ab} + f^{acb} A_\mu^c. \quad (2.7)$$

We thus see that the sum of the gauge-fixing and Faddeev-Popov terms can be written as a total BRST variation:

$$S_{\text{GF}} + S_{\text{FPG}} = \int d^4x s \left[\bar{c}^a \mathcal{F}^a - \frac{\xi}{2} \bar{c}^a b^a \right]. \quad (2.8)$$

This is of course expected, for it is well known that the physical observables of a theory admit a mathematical characterization in terms of the local cohomology of the BRST operator [35–37], and the latter is not affected by total BRST variations.

The BRST symmetry of the Yang-Mills action can be most conveniently exposed through the so-called BV method, i.e., introducing a set of antifields Φ^* and coupling them to the BRST variation of the corresponding fields through the term [38]

$$S_{\text{BV}} = \int d^4x \sum \Phi^* s \Phi. \quad (2.9)$$

Then the (tree-level) vertex functional is given by the sum

$$\Gamma^{(0)} = S_{\text{YM}} + S_{\text{GF}} + S_{\text{FPG}} + S_{\text{BV}}, \quad (2.10)$$

and the BRST symmetry of the action is encoded by the ST identity

$$\int d^4x \left[\frac{\delta \Gamma}{\delta A_{\mu}^{*a}} \frac{\delta \Gamma}{\delta A_a^{\mu}} + \frac{\delta \Gamma}{\delta c_a^{*}} \frac{\delta \Gamma}{\delta c^a} + b^a \frac{\delta \Gamma}{\delta \bar{c}^a} \right] = 0, \quad (2.11)$$

where now Γ is the full quantum effective action.

Turning to the case of BFM types of gauges, traditionally one starts by splitting the gauge field into a background part (\hat{A}) and a quantum part (Q) according to

$$A_{\mu}^a = \hat{A}_{\mu}^a + Q_{\mu}^a. \quad (2.12)$$

Next, one retains the background gauge invariance of the gauge-fixed action by choosing a gauge-fixing function that transforms in the adjoint representation of $\text{SU}(N)$ through the general replacements

$$\partial_{\mu} \delta^{ab} \rightarrow \hat{\mathcal{D}}_{\mu}^{ab} \equiv \partial_{\mu} \delta^{ab} + f^{acb} \hat{A}_{\mu}^c; \quad A_{\mu}^a \rightarrow Q_{\mu}^a, \quad (2.13)$$

that is one has the background R_{ξ} gauge

$$\hat{\mathcal{F}}^a = \hat{\mathcal{D}}_{\mu}^{ab} Q_b^{\mu}. \quad (2.14)$$

Finally, in addition to the antifields Φ^* , the quantization of the theory in the BFM requires the introduction of an additional (vector) anticommuting source Ω , implementing the equation of motion of the background field at the quantum level, with [20]

$$s \hat{A}_{\mu}^a = \Omega_{\mu}^a; \quad s \Omega_{\mu}^a = 0. \quad (2.15)$$

The BRST transformation of the quantum field Q is given by

$$s Q_{\mu}^a = \mathcal{D}_{\mu}^{ab} c^b - \Omega_{\mu}^a. \quad (2.16)$$

Equation (2.15) ensures that \hat{A} and Ω are paired in a so-called BRST doublet [35,39] (as already happens for \bar{c} and b), thus preventing the background field from modifying the physical observables of the theory.

It then follows that the conventional ST identity Eq. (2.11) gets modified into the extended ST identity

$$\int d^4x \left[\frac{\delta \Gamma}{\delta A_{\mu}^{*a}} \frac{\delta \Gamma}{\delta Q_a^{\mu}} + \frac{\delta \Gamma}{\delta c_a^{*}} \frac{\delta \Gamma}{\delta c^a} + b^a \frac{\delta \Gamma}{\delta \bar{c}^a} + \Omega_{\mu}^a \left(\frac{\delta \Gamma}{\delta \hat{A}_{\mu}^a} - \frac{\delta \Gamma}{\delta Q_{\mu}^a} \right) \right] = 0. \quad (2.17)$$

By “undoing” the shift of the gauge field (2.12) the ST identity above may be cast in the somewhat more compact form

$$\int d^4x \left[\frac{\delta \Gamma}{\delta A_{\mu}^{*a}} \frac{\delta \Gamma}{\delta A_a^{\mu}} + \frac{\delta \Gamma}{\delta c_a^{*}} \frac{\delta \Gamma}{\delta c^a} + b^a \frac{\delta \Gamma}{\delta \bar{c}^a} + \Omega_{\mu}^a \frac{\delta \Gamma}{\delta \hat{A}_{\mu}^a} \right] = 0. \quad (2.18)$$

In particular, this formulation of the BFM in terms of the field variables A and \hat{A} turns out to be the most suitable for the ensuing analysis.

III. ANTI-BRST SYMMETRY AND THE BFM

In the BRST transformations the role of the ghost field c is very prominent, as it replaces the gauge transformation parameter of the conventional gauge transformations and its behavior can be understood in an intrinsic manner in terms of the cohomology of the Lie algebra (see, e.g., [35,40]). On the other hand, the antighost \bar{c} and its doublet partner b play the role of Lagrange multipliers introduced to enforce the gauge-fixing condition $\mathcal{F} = 0$ and its BRST transform $s\mathcal{F} = 0$. In addition, \bar{c} obeys an equation of motion which is different from that of c as the former is not the Hermitian conjugate of the latter field.

Though all these seemed to rule out the possibility that \bar{c} and c can be interchanged, a nilpotent “anti-BRST” transformation symmetry in which this is exactly what happens was introduced long ago [28–30]. Indeed, the anti-BRST transformations can be obtained from the BRST ones of Eq. (2.6) by exchanging the role of the ghost and antighost fields; that is, one has

$$\begin{aligned} \bar{s} A_{\mu}^a &= \mathcal{D}_{\mu}^{ab} \bar{c}^b; & \bar{s} \bar{c}^a &= -\frac{1}{2} f^{abc} \bar{c}^b \bar{c}^c; \\ \bar{s} c^a &= \bar{b}^a; & \bar{s} \bar{b}^a &= 0. \end{aligned} \quad (3.1)$$

In particular, the anti-BRST transformation of the gauge field is obtained from the gauge variation of A by replacing the gauge parameter by the antighost field \bar{c} .

In order to close the algebra the transformations above need to be supplemented with the additional transformations

$$\bar{s} \bar{b}^a = f^{abc} \bar{b}^b \bar{c}^c; \quad \bar{s} b^a = f^{abc} b^b \bar{c}^c. \quad (3.2)$$

On the other hand, as both s and \bar{s} are nilpotent, the additional (natural) requirement that their sum is also nilpotent (or that $\{s, \bar{s}\} = 0$) results in the constraint [28]

$$\bar{b}^a = -b^a - f^{abc} c^b \bar{c}^c, \quad (3.3)$$

which, upon use of the Jacobi identity, is readily seen to be consistent with Eq. (3.2).

Finally, the nontrivial BRST-anti-BRST transformations of the fields read

$$\begin{aligned} s\bar{s}A_\mu^a &= \mathcal{D}_\mu^{ab} b^b + f^{abc} (\mathcal{D}_\mu^{bd} c^d) \bar{c}^c; \\ s\bar{s}c^a &= s\bar{b}^a; \quad s\bar{s}\bar{c}^a = -\bar{s}b^a. \end{aligned} \quad (3.4)$$

At this point it is straightforward to realize that to render our theory (2.1) simultaneously BRST and anti-BRST invariant, requires, *before gauge fixing*, the introduction of eight sources: the usual antifields A^* and c^* , the anti-BRST sources $A^\#$, $c^\#$, $\bar{c}^\#$, and $b^\#$, and, finally, the BRST-anti-BRST sources \hat{A} and \hat{c} . Notice that we do not add any source associated to $s\bar{b}$, for, due to the constraint (3.3), the BRST transformation of this field can be completely recovered from the corresponding transformations of b , c , and \bar{c} . One has then that the BRST-anti-BRST invariant action reads

$$S_I = S_{\text{YM}} + \sum \int d^4x (\Phi^* s\Phi + \Phi^\# \bar{s}\Phi + \hat{\Phi} s\bar{s}\Phi), \quad (3.5)$$

where the sum extends over all the nonzero sources, and (with the exception of $b^\#$)

$$\begin{aligned} s\Phi^* &= \bar{s}\Phi^* = 0; & s\hat{\Phi} &= \Phi^\#; \\ s\Phi^\# &= \bar{s}\Phi^\# = 0; & \bar{s}\hat{\Phi} &= -\Phi^*. \end{aligned} \quad (3.6)$$

For the source $b^\#$ one has instead

$$sb_a^\# = \bar{c}_a^\#; \quad \bar{s}b_a^\# = 0. \quad (3.7)$$

Finally, the ghost charge assignments are

$$\begin{aligned} \text{gh}(\Phi^*) &= -\text{gh}(\Phi) - 1; \\ \text{gh}(\Phi^\#) &= -\text{gh}(\Phi) + 1; \\ \text{gh}(\hat{\Phi}) &= -\text{gh}(\Phi), \end{aligned} \quad (3.8)$$

where we have set $\text{gh}(c, \bar{c}) = (1, -1)$. Notice that the usual BV action [38] is recovered by setting the $\Phi^\#$ and $\hat{\Phi}$ sources to zero.

We are now ready to establish the central result of this paper. Consider, in fact, the BFM covariant gauge-fixing (2.14) with its associated Faddeev-Popov ghost action; a straightforward calculation yields

$$\begin{aligned} s \left[\bar{c}^a \hat{\mathcal{F}}^a - \frac{\xi}{2} \bar{c}^a b^a \right] &= s \left[\bar{c}^a \mathcal{F}^a - \frac{\xi}{2} \bar{c}^a b^a \right] \\ &\quad + \hat{A}_a^\mu (\mathcal{D}_\mu^{ab} b^b + f^{abc} \bar{c}^b \mathcal{D}_\mu^{cd} c^d) \\ &\quad + \Omega_a^\mu (\mathcal{D}_\mu^{ab} \bar{c}^b) \\ &= s \left[\bar{c}^a \mathcal{F}^a - \frac{\xi}{2} \bar{c}^a b^a \right] \\ &\quad + \hat{A}_a^\mu (s\bar{s}A_\mu^a) + \Omega_a^\mu (\bar{s}A_\mu^a), \end{aligned} \quad (3.9)$$

where \mathcal{F}^a is now the covariant gauge-fixing (2.4). As a result of the anticommutation relation $\{s, \bar{s}\} = 0$ and the identity

$$s(\bar{c}^a \partial^\mu A_\mu^a) = -\bar{s}(c^a \partial^\mu A_\mu^a), \quad (3.10)$$

we observe that also the first term on the right-hand side (r.h.s.) of Eq. (3.9) is both BRST and anti-BRST invariant (the term b^2 is obviously invariant under these transformations).

Then we see that by adding to the BRST-anti-BRST invariant action (3.5) the R_ξ gauge fixing and Faddeev-Popov term (2.8) we automatically obtain a theory formulated in the background R_ξ gauge, provided that the following identification is made:

$$\Omega_\mu^a \equiv A_\mu^{\#a}. \quad (3.11)$$

From Eq. (3.9) one also sees that the background gauge field \hat{A} is the source of the BRST-anti-BRST variation $s\bar{s}A$ of the gauge field A . Notice however that \hat{c} cannot be interpreted as a background for the ghost c , since it has ghost number -1 ; it is also clear that it is not a background for the antighost field, as a shift of the latter field would lead to totally different couplings w.r.t. the ones that are generated for the source \hat{c} .

Thus one arrives at the somewhat surprising conclusion that *requiring the invariance of a $SU(N)$ Yang-Mills action gauge fixed in an R_ξ gauge under both BRST as well as anti-BRST symmetry is equivalent to quantizing the theory in the (R_ξ) BFM*:

$$\begin{aligned} \Gamma^{(0)} &= S_I + S_{\text{GF}} + S_{\text{FPG}} \\ &= S_{\text{YM}} + \hat{S}_{\text{GF}} + \hat{S}_{\text{FPG}} + S_{\text{BV}} + \int d^4x (c_a^\# \bar{s}c_a + \hat{c}_a s\bar{s}c_a), \end{aligned} \quad (3.12)$$

where the background gauge-fixing functional and the background Faddeev-Popov terms are

$$\begin{aligned} \hat{S}_{\text{GF}} &= \int d^4x \left[-\frac{\xi}{2} (b^a)^2 + b^a \hat{\mathcal{F}}^a \right]; \\ \hat{S}_{\text{FPG}} &= - \int d^4x \bar{c}^a s \hat{\mathcal{F}}^a. \end{aligned} \quad (3.13)$$

The standard BFM tree-level vertex functional is recovered by setting $c^\# = \hat{c} = 0$ on the r.h.s. of Eq. (3.12). In this

sense the BFM is not fundamental, as it is naturally emerging from the requirement of anti-BRST invariance.

It is interesting to study the case in which (complex) scalars and/or fermions are added to the theory. Let us start from the former fields, where one has

$$\begin{aligned} s\phi &= ic^a t^a \phi; & \bar{s}\phi &= i\bar{c}^a t^a \phi; \\ s\phi^\dagger &= -ic^a \phi^\dagger t^a; & \bar{s}\phi^\dagger &= -i\bar{c}^a \phi^\dagger t^a, \end{aligned} \quad (3.14)$$

with t^a the generators of the SU(N) representation chosen for ϕ . The corresponding BRST-anti-BRST transformation reads

$$\begin{aligned} s\bar{s}\phi &= ib^a t^a \phi + \bar{c}^a c^b t^a t^b \phi; \\ s\bar{s}\phi^\dagger &= -ib^a \phi^\dagger t^a + \bar{c}^a c^b \phi^\dagger t^b t^a, \end{aligned} \quad (3.15)$$

from which it is immediate to infer that $\{s, \bar{s}\}\phi = 0$.

The extra sources one needs to add to render the action BRST and anti-BRST invariant in the presence of the scalar field ϕ are then¹

$$\begin{aligned} \int d^4x [&\phi^{*\dagger} s\phi + s\phi^\dagger \phi^* + \phi^{\#\dagger} \bar{s}\phi + (\bar{s}\phi^\dagger) \phi^\# \\ &+ \hat{\phi}^\dagger s\bar{s}\phi + (s\bar{s}\phi^\dagger) \hat{\phi}]. \end{aligned} \quad (3.16)$$

Again by identifying $\hat{\phi}$ and $\hat{\phi}^\dagger$ with the background for the scalars ϕ and ϕ^\dagger , respectively, as well as $\phi^\#, \phi^{\#\dagger}$ with their corresponding BRST doublet partners [i.e., $s\phi = \phi^\#, s\phi^\dagger = \phi^{\#\dagger}$, as prescribed by Eq. (3.6)], one recovers the background 't Hooft gauge after the background field $\hat{\phi}$ has acquired an expectation value v .

For fermions ψ and $\bar{\psi}$ the analysis proceeds in the same way as in the scalar case, since Eqs. (3.14) and (3.15) still hold once t^a is identified with the generator of the representation of the fermionic matter field and ϕ replaced by ψ , i.e.,

$$\begin{aligned} s\psi &= ic^a t^a \psi; & \bar{s}\psi &= i\bar{c}^a t^a \psi; \\ s\bar{\psi} &= -ic^a \bar{\psi} t^a; & \bar{s}\bar{\psi} &= -i\bar{c}^a \bar{\psi} t^a. \end{aligned} \quad (3.17)$$

Notice that the requirement of anti-BRST invariance generates unavoidably the sources $\hat{\psi}$ and $\hat{\bar{\psi}}$ which correspond to background fields for the fermions,² as the action will be rendered BRST-anti-BRST invariant through the addition of the term

$$\begin{aligned} \int d^4x [&\bar{\psi}^* s\psi - (s\bar{\psi}) \psi^* + \bar{\psi}^\# \bar{s}\psi \\ &- (\bar{s}\bar{\psi}) \psi^\# - \hat{\psi} s\bar{s}\psi + (s\bar{s}\bar{\psi}) \hat{\psi}], \end{aligned} \quad (3.18)$$

¹We assume that a suitable (gauge invariant) action term S_ϕ (and S_ψ when adding fermions) is added to the classical action (2.1); its concrete form is however irrelevant for the following analysis.

²Fermionic backgrounds have been considered, e.g., in [41]; their physical relevance is however unclear to us at the moment.

where

$$\begin{aligned} s\bar{s}\psi &= ib^a t^a \psi + \bar{c}^a c^b t^a t^b \psi; \\ s\bar{s}\bar{\psi} &= -ib^a \bar{\psi} t^a + \bar{c}^a c^b \bar{\psi} t^b t^a. \end{aligned} \quad (3.19)$$

IV. LOCAL ANTIGHOST EQUATION

The presence of the anti-BRST symmetry leads, as we will explicitly show below, to the existence of a local antighost equation. It should be noticed that for a SU(N) Yang-Mills theory with a conventional R_ξ gauge fixing only an *integrated* antighost equation can be derived, while in the BFM case the existence of a local version of this equation was established in the background Landau gauge in [31] and believed to be valid only for that specific gauge-fixing choice.³

On the other hand, the correspondence just found between BRST and anti-BRST invariance and the BFM shows that there should not be anything special neither when formulating the theory in the BFM nor when choosing $\xi = 0$. Indeed as the existence of the anti-BRST symmetry puts the ghost and antighost fields on the same footing, and given that a local ghost equation (sometimes also referred to as Faddeev-Popov equation) is known to hold, we would expect a local antighost equation to hold as well.

To show that this is indeed the case, let us start by setting to zero the scalar and fermionic matter sector (the complete case will be discussed in the Appendix); then the tree-level action (3.5) can be cast in the form

$$\Gamma^{(0)} = S_{\text{YM}} + sX = S_{\text{YM}} + \bar{s}Y, \quad (4.1)$$

where

$$\begin{aligned} X &= \int d^4x \left[\sum ((-1)^{\text{gh}(\Phi^*)} \Phi^* \Phi + \hat{\Phi} \bar{s}\Phi) + \bar{c}^a \mathcal{F}^a - \frac{\xi}{2} \bar{c}^a b^a \right], \\ Y &= \int d^4x \left[\sum ((-1)^{\text{gh}(\Phi^\#)} \Phi^\# \Phi + \hat{\Phi} s\Phi) - c^a \mathcal{F}^a + \frac{\xi}{2} c^a b^a \right], \end{aligned} \quad (4.2)$$

and the sum is intended, as familiar by now, over all nonzero sources.

To derive the local antighost equation the fastest route turns out to be to calculate the anticommutator between the derivative w.r.t. the ghost field and the anti-BRST operator. Since

$$\bar{s} = \sum \int d^4x \bar{s}\varphi(x) \frac{\delta}{\delta\varphi(x)}; \quad \varphi = \Phi, \Phi^*, \Phi^\#, \hat{\Phi}, \quad (4.3)$$

³Notice that in the case of a ghost-antighost symmetric gauge, the existence of such a local equation is a direct consequence of the Sp[2] symmetry (see, e.g., [42]).

one finds that for any functional $F = F[\varphi]$ with zero ghost charge

$$\left\{ \frac{\delta}{\delta c^a}, \bar{s} \right\} F = \sum \int d^4x \left[\frac{\delta}{\delta c^a} \bar{s} \varphi(x) \right] \frac{\delta F}{\delta \varphi(x)}, \quad (4.4)$$

and therefore

$$\frac{\delta \Gamma}{\delta c^a} = \frac{\delta}{\delta c^a} (\bar{s} Y) = \sum \int d^4x \left[\frac{\delta}{\delta c^a} \bar{s} \varphi(x) \right] \frac{\delta Y}{\delta \varphi(x)} - \bar{s} \frac{\delta Y}{\delta c^a}. \quad (4.5)$$

Then through a lengthy but relatively straightforward calculation, we arrive at the local antighost equation

$$\begin{aligned} \bar{\mathcal{G}}_a \Gamma &\equiv \frac{\delta \Gamma}{\delta c^a} + f^{abc} \frac{\delta \Gamma}{\delta b^b} \bar{c}^c + \xi \frac{\delta \Gamma}{\delta b_a^\#} - \hat{\mathcal{D}}_\mu^{ab} \frac{\delta \Gamma}{\delta A_\mu^{*b}} \\ &\quad - f^{abc} \hat{c}_b \frac{\delta \Gamma}{\delta c_c^\#} - f^{abc} b_b^\# \frac{\delta \Gamma}{\delta \bar{c}_c^\#} \\ &= \mathcal{D}_\mu^{ab} A_b^{*\mu} + f^{abc} c_b^* c^c, \end{aligned} \quad (4.6)$$

where $\hat{\mathcal{D}}_\mu^{ab} = \partial_\mu \delta^{ab} + f^{acb} \hat{A}_\mu^c$. Notice that all the (possibly present) trilinear terms in the ghost and antighost fields have canceled out.

In the case of the local ghost equation one computes the anticommutator of the derivative w.r.t. the antighost field and the BRST operator s :

$$\frac{\delta \Gamma}{\delta \bar{c}^a} = \frac{\delta}{\delta \bar{c}^a} (sX) = \sum \int d^4x \left[\frac{\delta}{\delta \bar{c}^a} s \varphi(x) \right] \frac{\delta X}{\delta \varphi(x)} - s \frac{\delta X}{\delta \bar{c}^a}. \quad (4.7)$$

One then has

$$\begin{aligned} \mathcal{G}_a \Gamma &\equiv \frac{\delta \Gamma}{\delta \bar{c}^a} + \hat{\mathcal{D}}_\mu^{ab} \frac{\delta \Gamma}{\delta A_\mu^{*b}} + f^{abc} \hat{c}_b \frac{\delta \Gamma}{\delta c_c^*} \\ &= \mathcal{D}_\mu^{ab} A_b^{*\mu} + f^{abc} c_b^\# c^c + f^{abc} \bar{c}_b^\# \bar{c}^c - f^{abc} b_b^\# b^c. \end{aligned} \quad (4.8)$$

Finally, the b equation assumes the form

$$\frac{\delta \Gamma}{\delta b^a} = \hat{\mathcal{D}}_\mu^{ab} (A_b^\mu - \hat{A}_b^\mu) - \xi b^a - f^{abc} b_b^\# \bar{c}^c - c_a^\# - f^{abc} \hat{c}_b c^c, \quad (4.9)$$

while the ST and anti-ST identities read respectively⁴

$$\begin{aligned} \int d^4x \left[\frac{\delta \Gamma}{\delta A_\mu^{*a}} \frac{\delta \Gamma}{\delta A_\mu^a} + \frac{\delta \Gamma}{\delta c_a^\#} \frac{\delta \Gamma}{\delta c^a} + A_\mu^{*a} \frac{\delta \Gamma}{\delta \hat{A}_\mu^a} \right. \\ \left. + c_a^\# \frac{\delta \Gamma}{\delta \hat{c}_a} + \bar{c}_a^\# \frac{\delta \Gamma}{\delta b_a^\#} + b^a \frac{\delta \Gamma}{\delta \bar{c}^a} \right] = 0, \end{aligned} \quad (4.10)$$

⁴From here we see that an alternative way of deriving the local antighost equation is to take the derivative w.r.t. b of the anti-ST identity (4.11) and next use the b equation (4.9) to replace the various terms involving the functional derivative w.r.t. b .

$$\begin{aligned} \int d^4x \left[\frac{\delta \Gamma}{\delta A_\mu^{*a}} \frac{\delta \Gamma}{\delta A_\mu^a} + \frac{\delta \Gamma}{\delta c_a^\#} \frac{\delta \Gamma}{\delta c^a} + \frac{\delta \Gamma}{\delta \bar{c}_a^\#} \frac{\delta \Gamma}{\delta \bar{c}^a} \right. \\ \left. + \frac{\delta \Gamma}{\delta b_a^\#} \frac{\delta \Gamma}{\delta b^a} - A_\mu^{*a} \frac{\delta \Gamma}{\delta \hat{A}_\mu^a} - c_a^\# \frac{\delta \Gamma}{\delta \hat{c}_a} \right] = 0. \end{aligned} \quad (4.11)$$

The background Ward identity follows as a consequence of the local antighost equation and the ST identity, since

$$0 = \mathcal{S}_\Gamma (\bar{\mathcal{G}}_a \Gamma - \mathcal{D}_\mu^{ab} A_b^{*\mu} - f^{abc} c_b^* c^c) + \mathcal{G}_a \mathcal{S}(\Gamma) = \mathcal{W}_a \Gamma. \quad (4.12)$$

In the above equation \mathcal{S}_Γ is the linearized ST operator

$$\begin{aligned} \mathcal{S}_\Gamma \equiv \int d^4x \left[\frac{\delta \Gamma}{\delta A_\mu^{*a}} \frac{\delta}{\delta A_\mu^a} + \frac{\delta \Gamma}{\delta A_\mu^a} \frac{\delta}{\delta A_\mu^{*a}} + \frac{\delta \Gamma}{\delta c_a^\#} \frac{\delta}{\delta c^a} \right. \\ \left. + \frac{\delta \Gamma}{\delta c_a^\#} \frac{\delta}{\delta c_a^*} + A_\mu^{*a} \frac{\delta}{\delta \hat{A}_\mu^a} + c_a^\# \frac{\delta}{\delta \hat{c}_a} + \bar{c}_a^\# \frac{\delta}{\delta b_a^\#} + b^a \frac{\delta}{\delta \bar{c}^a} \right], \end{aligned} \quad (4.13)$$

while the background Ward operator reads

$$\begin{aligned} \mathcal{W}_a \equiv -\mathcal{D}_\mu^{ab} \frac{\delta}{\delta A_{b\mu}} - \hat{\mathcal{D}}_\mu^{ab} \frac{\delta}{\delta \hat{A}_{b\mu}} + f^{abc} c_c \frac{\delta}{\delta c_b} \\ + f^{abc} \bar{c}_c \frac{\delta}{\delta \bar{c}_b} + f^{abc} b_c \frac{\delta}{\delta b_b} + f^{abc} c_c^* \frac{\delta}{\delta c_b^*} \\ + f^{abc} c_c^\# \frac{\delta}{\delta c_b^\#} + f^{abc} \bar{c}_c^\# \frac{\delta}{\delta \bar{c}_b^\#} + f^{abc} A_{\mu c}^* \frac{\delta}{\delta A_{\mu b}^*} \\ + f^{abc} A_{\mu c}^\# \frac{\delta}{\delta A_{\mu b}^\#} + f^{abc} \hat{c}_c \frac{\delta}{\delta \hat{c}_b} + f^{abc} b_c^\# \frac{\delta}{\delta b_b^\#}. \end{aligned} \quad (4.14)$$

In a similar fashion, the background Ward identity can also be obtained by taking the anticommutator between the linearized anti-ST operator and the local ghost equation operator \mathcal{G} and then using the anti-ST identity and the local ghost equation.

V. BACKGROUND EFFECTIVE ACTION

The requirement of BRST and anti-BRST invariance in the presence of scalar and fermionic matter leads to the generalization of the ST and anti-ST identities of Eqs. (4.10) and (4.11) shown in Eqs. (A1) and (A2) of the Appendix. The sources of the anti-BRST variations for the gauge, scalar, and fermionic matter fields are to be identified with the corresponding background fields.

In order to further elucidate the physical content of the ST and anti-ST identities, it is convenient to construct an effective action $\tilde{\Gamma}$ for the background configurations by integrating out completely the quantum fields. That is, one is interested in keeping only connected diagrams with external background legs.

The functional $\tilde{\Gamma}$, which is therefore one-particle reducible w.r.t. the quantum fields, can be formally obtained as

follows. The connected generating functional W is obtained by taking a Legendre transform w.r.t. Φ :

$$W = \Gamma + \int d^4x J_\Phi \Phi; \quad J_\Phi = -(-1)^{\epsilon(\Phi)} \frac{\delta \Gamma}{\delta \Phi};$$

$$\Phi = \frac{\delta W}{\delta J_\Phi}; \quad \frac{\delta W}{\delta \zeta} = \frac{\delta \Gamma}{\delta \zeta}; \quad \zeta \in \{\hat{\Phi}, \Phi^*, \Phi^\#\}, \quad (5.1)$$

where we use a collective notation, with J_Φ denoting the source of the quantum field Φ and $\epsilon(\Phi)$ the statistics of the field Φ (1 for anticommuting variables, 0 for commuting ones).

Then one sets

$$\tilde{\Gamma}[\hat{\Phi}, \Phi^*, \Phi^\#] \equiv W[J_\Phi, \hat{\Phi}, \Phi^*, \Phi^\#]|_{J_\Phi=0}. \quad (5.2)$$

Equation (A1) yields the following identity for the connected functional W :

$$\int d^4x \left[-\frac{\delta W}{\delta A_\mu^{*a}} J_{A_\mu^a} + \frac{\delta W}{\delta c_a^*} J_{c^a} + \frac{\delta W}{\delta J_{b^a}} J_{\bar{c}^a} - \frac{\delta W}{\delta \phi^{*\dagger}} J_\phi \right. \\ \left. + \frac{\delta W}{\delta \phi^*} J_{\phi^\dagger} + \frac{\delta W}{\delta \psi^*} J_\psi - \frac{\delta W}{\delta \psi^{*\dagger}} J_{\bar{\psi}} + A_\mu^\# \frac{\delta W}{\delta \hat{A}_{a\mu}} + c_a^\# \frac{\delta W}{\delta \hat{c}_a} \right. \\ \left. + \bar{c}_a^\# \frac{\delta W}{\delta \hat{b}_a} - \phi^\# \frac{\delta W}{\delta \hat{\phi}} + \phi^{*\dagger} \frac{\delta W}{\delta \hat{\phi}^\dagger} - \psi^\# \frac{\delta W}{\delta \hat{\psi}} + \bar{\psi}^\# \frac{\delta W}{\delta \hat{\psi}} \right] = 0. \quad (5.3)$$

By taking a derivative of the above equation w.r.t. any of the anti-BRST sources $\Phi^\#$ and then setting all the sources J_Φ and $\Phi^\#$ to zero, one finds that

$$\frac{\delta \tilde{\Gamma}}{\delta \hat{\Phi}} = \frac{\delta W}{\delta \hat{\Phi}} \Big|_{J_\Phi=\Phi^\#=0} = 0. \quad (5.4)$$

This means that the background field configurations $\hat{\Phi}$ constitute a stationary point for the background effective action $\tilde{\Gamma}$.

Notice that the same result is obtained if one starts from the anti-ST identity for the connected generating functional W , takes one derivative w.r.t. the BRST source Φ^* , and then sets all the sources J_Φ and Φ^* to zero.

As a physical example, one can consider the effective field theory of the color glass condensate [33,34], which describes the physics of high gluon densities and gluon saturation in the small x regime (x denoting the longitudinal momentum fraction of the parton in the collision). In this framework, the fulfillment of the ST identity for $\tilde{\Gamma}$ is crucial for guaranteeing the consistency of the approximations used, as it shows that the background field configuration \hat{A} is still a stationary point of the background effective action, even in the presence of radiative corrections induced by the integration of certain quantum modes of the gluon field.

A. Two- and three-point background gauge functions

In order to illustrate the combinatorics behind the stationary condition (5.4), let us consider the case of the two- and three-point background gauge functions.

The graphs contributing to the two-point function $\tilde{\Gamma}_{\hat{A}\hat{A}}$ are depicted in Fig. 1. From Eq. (5.1) one sees that

$$\int d^4z \Gamma_{A_\mu^a A_\nu^c}(x, z) W_{J_{A_\rho^c} J_{A_\rho^b}}(z, y) = -\delta_{ab} g^{\mu\nu} \delta^4(x-y). \quad (5.5)$$

If one replaces the 1-PI functions $\Gamma_{\hat{A}\hat{A}}$ in the second of the diagrams in Fig. 1(a) by exploiting the background-quantum identity

$$\Gamma_{\hat{A}_\mu^a A_\nu^b} = -\Gamma_{\Omega_\mu^a A_\rho^{*c}} \Gamma_{A_\rho^b A_\rho^c}, \quad (5.6)$$

the gauge propagator cancels against one of the 1-PI 2-point gauge functions by Eq. (5.5). The identity

$$\tilde{\Gamma}_{\hat{A}_{\mu_1}^{a_1} \hat{A}_{\mu_2}^{a_2}} = 0 \quad (5.7)$$

then boils down to the usual background-quantum relation

$$\Gamma_{\hat{A}_{\mu_1}^{a_1} \hat{A}_{\mu_2}^{a_2}}(x_1, x_2) = \int d^4z_1 \int d^4z_2 \Gamma_{\Omega_{\mu_1}^{a_1} A_{c_1\rho_1}^*}(x_1, z_1) \\ \times \Gamma_{\Omega_{\mu_2}^{a_2} A_{c_2\rho_2}^*}(x_2, z_2) \Gamma_{A_{c_1\rho_1} A_{c_2\rho_2}}(z_1, z_2). \quad (5.8)$$

The identity for the three-point function

$$\tilde{\Gamma}_{\hat{A}_{\mu_1}^{a_1} \hat{A}_{\mu_2}^{a_2} \hat{A}_{\mu_3}^{a_3}}(x_1, x_2, x_3) = 0 \quad (5.9)$$

is more involved. The diagrams contributing to Eq. (5.9) are depicted in Fig. 1(b). They are arranged according to the number of \hat{A} insertions in the 1-PI vertex bubbles. In order to establish the connection with the 1-PI background-quantum identities, we notice that the legs involving a gauge propagator and a mixed background-quantum amplitude $\Gamma_{\hat{A}\hat{A}}$ can be reduced with the help of Eq. (5.6) as follows:

$$\Gamma_{\hat{A}_\mu^a A_\nu^b} W_{J_{A_\rho^b} J_{A_\rho^c}} = -\Gamma_{\Omega_\mu^a A_\sigma^{*d}} \Gamma_{A_\sigma^d A_\nu^b} W_{J_{A_\rho^b} J_{A_\rho^c}} = \Gamma_{\Omega_\mu^a A_\rho^{*c}}, \quad (5.10)$$

where Eq. (5.5) has been used.

After the replacement in Eq. (5.10) has been carried out, one obtains from Eq. (5.9) a representation for $\tilde{\Gamma}_{\hat{A}\hat{A}\hat{A}}$ which only involves 1-PI amplitudes. It is uniquely determined by the requirement that the ST identity holds. It can therefore be obtained by applying the method of canonical transformations presented in [27]. Specifically, by taking the derivative of the ST identity w.r.t. Ω and setting Ω to zero afterwards, one obtains quite generally an identity of the form

$$\frac{\delta \Gamma}{\delta \hat{A}_\mu^a} \Big|_{\Omega=0} = -\left\{ \frac{\delta \Gamma}{\delta \Omega_\mu^a}, \Gamma \right\} \Big|_{\Omega=0}, \quad (5.11)$$

where $\{ \cdot, \cdot \}$ denotes the BV bracket associated with the ST identity. Equation (5.11) states that the dependence on the




FIG. 1. The cancellations encoded in the stationary condition (5.4) in the case of the two- and three-point functions of the background gauge field. Small circles attached at the end of lines indicate the background gluons.

background is generated via a canonical transformation with respect to the BV bracket, induced by the generating functional $\frac{\delta\Gamma}{\delta\Omega}$. The latter in general is \hat{A} dependent. As a consequence, the solution cannot be written by simple exponentiation of the BV bracket w.r.t. $\frac{\delta\Gamma}{\delta\Omega}$, but requires the introduction of a Lie series of a suitable functional differential operator [27].

For our purposes it is sufficient to consider the reduced bracket

$$\left\{ \frac{\delta\Gamma}{\delta\Omega_\mu^a(x)}, \cdot \right\} \equiv \int d^4z \left[-\Gamma_{\Omega_\mu^a(x)A_\nu^{*b}}(x, z) \frac{\delta}{\delta A_\nu^b(z)} + \Gamma_{\Omega_\mu^a A_\nu^*}(x, z) \frac{\delta}{\delta A_{b\nu}^*(z)} \right]. \quad (5.12)$$

Then the Lie series generating the background field dependence is obtained by exponentiating the operator

$$\Delta_{\Gamma_{\Omega_\mu^a}(x)} \equiv \left\{ \frac{\delta\Gamma}{\delta\Omega_\mu^a}, \cdot \right\} + \frac{\delta}{\delta \hat{A}_\mu^a}, \quad (5.13)$$

that is, one has

$$\Gamma = E_{\Gamma_\Omega}(\Gamma_0) + \dots, \quad (5.14)$$

where the dots denote amplitudes involving at least one external leg different than A , \hat{A} and the mapping E_{Γ_Ω} is defined according to

$$E_{\Gamma_\Omega}(\Gamma_0) \equiv \sum_{n \geq 0} \frac{1}{n!} \int_1 \dots \int_n \hat{A}_1 \dots \hat{A}_n [\Delta_{\Gamma_{\Omega_1}} \dots \Delta_{\Gamma_{\Omega_n}} \Gamma_0]_{\hat{A}=0}. \quad (5.15)$$

In the above equation Γ_0 denotes the 1-PI vertex functional where fields and sources have been set to zero, with the

only exception of A, A^* ; finally, the shorthand notations $\int_i = \int d^4 z_i$, $\hat{A}_i = \hat{A}_{\mu_i}^{a_i}(z_i)$, and $\Gamma_{\Omega_i} = \Gamma_{\Omega_{\mu_i}^{a_i}}(z_i)$ have been used.

We stress that Eq. (5.14) reproduces the correct dependence of the 1-PI vertex functional on \hat{A} in the gauge sector only. The background dependence of amplitudes involving scalar and fermionic matter fields as well as other external sources can be recovered by making use of the full canonical transformation generated by the functional $\frac{\delta\Gamma}{\delta\Omega}$.

The first two terms of Eq. (5.14) yield (notice that in the following equations \hat{A} is set equal to zero, while further differentiations w.r.t. A are possible)

$$\Gamma_{\hat{A}_{\mu_1}^{a_1}}(x_1) = - \int d^4 z_1 \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1}}(x_1, z_1) \Gamma_{A_{\beta_1}^{b_1}}(z_1), \quad (5.16)$$

and

$$\begin{aligned} \Gamma_{\hat{A}_{\mu_1}^{a_1} \hat{A}_{\mu_2}^{a_2}}(x_1, x_2) &= \frac{1}{2} \int d^4 z_1 \int d^4 z_2 [\Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2}}(x_2, z_2) \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1} A_{\beta_2}^{b_2}}(x_1, z_1, z_2) \Gamma_{A_{\beta_1}^{b_1}}(z_1) + \Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2}}(x_2, z_2) \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1}}(x_1, z_1) \\ &\quad \times \Gamma_{A_{\beta_2}^{b_2} A_{\beta_1}^{b_1}}(z_2, z_1) - \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1} \hat{A}_{\mu_2}^{a_2}}(x_1, z_1, x_2) \Gamma_{A_{\beta_1}^{b_1}}(z_1)] + (a_1 \mu_1 \leftrightarrow a_2 \mu_2), \end{aligned} \quad (5.17)$$

while the third term gives for the three point function $\Gamma_{\hat{A}\hat{A}\hat{A}}$ (we suppress the space-time arguments)

$$\begin{aligned} \Gamma_{\hat{A}_{\mu_1}^{a_1} \hat{A}_{\mu_2}^{a_2} \hat{A}_{\mu_3}^{a_3}} &= \frac{1}{3!} \int d^4 z_1 \int d^4 z_2 \int d^4 z_3 [-\Gamma_{\Omega_{\mu_3}^{a_3} A_{\beta_3}^{*b_3}} \Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2}} \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1} A_{\beta_2}^{b_2}} \Gamma_{A_{\beta_3}^{b_3} A_{\beta_1}^{b_1}} - \Gamma_{\Omega_{\mu_3}^{a_3} A_{\beta_3}^{*b_3}} \Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2} A_{\beta_3}^{b_3}} \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1}} \Gamma_{A_{\beta_2}^{b_2} A_{\beta_1}^{b_1}} \\ &\quad - \Gamma_{\Omega_{\mu_3}^{a_3} A_{\beta_3}^{*b_3}} \Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2}} \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1} A_{\beta_3}^{b_3}} \Gamma_{A_{\beta_2}^{b_2} A_{\beta_1}^{b_1}} - \Gamma_{\Omega_{\mu_3}^{a_3} A_{\beta_3}^{*b_3}} \Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2}} \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1}} \Gamma_{A_{\beta_3}^{b_3} A_{\beta_2}^{b_2} A_{\beta_1}^{b_1}} + \Gamma_{\Omega_{\mu_3}^{a_3} A_{\beta_3}^{*b_3}} \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1} \hat{A}_{\mu_2}^{a_2}} \Gamma_{A_{\mu_3}^{a_3} A_{\beta_1}^{b_1}} \\ &\quad + \Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2} \hat{A}_{\mu_3}^{a_3}} \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1}} \Gamma_{A_{\beta_2}^{b_2} A_{\beta_1}^{b_1}} + \Gamma_{\Omega_{\mu_2}^{a_2} A_{\beta_2}^{*b_2}} \Gamma_{\Omega_{\mu_1}^{a_1} A_{\beta_1}^{*b_1} \hat{A}_{\mu_3}^{a_3}} \Gamma_{A_{\beta_2}^{b_2} A_{\beta_1}^{b_1}}] + \text{symm}, \end{aligned} \quad (5.18)$$

where complete symmetrization of the r.h.s. of the above equation w.r.t. the a_i, μ_i indices is understood. Compatibility of the diagrammatic identity in Fig. 1(b) with Eq. (5.18) follows by taking the appropriate derivatives w.r.t. the quantum fields A of Eqs. (5.16) and (5.17) in order to eliminate recursively the background insertions in the amplitudes of the second and third lines of Fig. 1(b).

Notice in particular the presence of the amplitudes $\Gamma_{\Omega_{A^*} \hat{A}}$. The latter arise due to the dependence of the generating functional Γ_{Ω} on the background \hat{A} . The amplitudes $\Gamma_{\Omega_{A^*} \hat{A}}$ can be fully fixed either by the ST or the anti-ST identities. Their \hat{A} dependence is the cause of the failure of the simple exponentiation in order to derive the solution to Eq. (5.11), which in turn can be overcome by using the appropriate Lie series in Eq. (5.15).

VI. PRACTICAL CONSEQUENCES

A. Two-point ghost sector

The presence of the antighost equation allows one to fully constrain the ghost two-point sector in any gauge. In this sector there are four superficially divergent Green's functions, namely, $\Gamma_{c\bar{c}b}, \Gamma_{\bar{c}^a \Omega_{\mu}^b}, \Gamma_{c^a A_{\mu}^{*b}}$, and, finally, $\Gamma_{\Omega_{\mu}^a A_{\nu}^{*b}}$ (in the following we prefer to switch back to the familiar notation of Ω rather than using its anti-BRST source name $A^{\#}$). The first two functions are constrained by the ghost equation (we factor out the trivial color structure δ^{ab})

$$\begin{aligned} \Gamma_{c\bar{c}}(q) &= -iq^{\mu} \Gamma_{cA_{\mu}^*}(q), \\ \Gamma_{\Omega_{\mu}^a \bar{c}}(q) &= iq_{\mu} - iq^{\nu} \Gamma_{\Omega_{\mu}^a A_{\nu}^*}(q). \end{aligned} \quad (6.1)$$

On the other hand, differentiating the antighost equation (4.6) with respect to a gluon antifield and an antighost, one gets the deformed identities

$$\begin{aligned} \Gamma_{A_{\mu}^* c}(q) &= iq_{\mu} + iq^{\nu} \Gamma_{A_{\mu}^* \Omega_{\nu}^*}(q) - \xi \Gamma_{A_{\mu}^* b^{\#}}(q), \\ \Gamma_{\bar{c} c}(q) &= iq^{\mu} \Gamma_{\bar{c} \Omega_{\mu}^*}(q) - \xi \Gamma_{\bar{c} b^{\#}}(q), \end{aligned} \quad (6.2)$$

and the functions $\Gamma_{b^{\#} A_{\mu}^*}$ and $\Gamma_{b^{\#} \bar{c}}$ related through the identity

$$\Gamma_{b^{\#} \bar{c}}(q) = -iq^{\mu} \Gamma_{b^{\#} A_{\mu}^*}(q). \quad (6.3)$$

Contracting the first equation in (6.2) with q^{μ} and next using the first of the identities (6.1) as well as Eq. (6.3), we find the relation

$$\Gamma_{c\bar{c}}(q) = q^2 - q^{\mu} q^{\nu} \Gamma_{\Omega_{\mu}^a A_{\nu}^*}(q) - \xi \Gamma_{b^{\#} \bar{c}}(q), \quad (6.4)$$

which shows the appearance of the extra function $\Gamma_{b^{\#} \bar{c}}$ with respect to the Landau gauge, where the ghost sector is entirely determined by $\Gamma_{\Omega_{A^*}}$ alone.

Then, observing that

$$\Gamma_{c\bar{c}}(q) = q^2 F^{-1}(q^2), \quad (6.5)$$

where F is the ghost dressing function related to the ghost propagator D through $D(q^2) = F(q^2)/q^2$, and introducing the Lorentz decompositions

$$\begin{aligned} \Gamma_{cA_{\mu}^*}(q) &= iq_{\mu} C(q^2); & \Gamma_{\bar{c} \Omega_{\mu}^*}(q) &= iq_{\mu} E(q^2); \\ \Gamma_{b^{\#} \bar{c}}(q) &= -q^2 K(q^2); \\ \Gamma_{\Omega_{\mu}^a A_{\nu}^*}(q) &= -g_{\mu\nu} G(q^2) - \frac{q_{\mu} q_{\nu}}{q^2} L(q^2), \end{aligned} \quad (6.6)$$

$$Z_A^{-2} = [1 + G(0)]^2 Z_Q^{-2}, \quad (6.16)$$

where we have introduced the background field renormalization constant $\hat{A} = Z_{\hat{A}} \hat{A}_0$. We now take advantage of the residual background gauge invariance which implies the QED-like relation

$$Z_{\hat{A}}^{-1} = Z_g, \quad (6.17)$$

to get finally the desired relation (6.11).

When originally derived in [43] this relation was discussed in the context of the so-called Kugo-Ojima confinement criterion [48], which predicts that, in the Landau gauge, a sufficient condition for color confinement is $1 + G(0) = 0$ (which would in turn imply an IR divergent ghost dressing function). It turns out, however, that lattice data (see [49] for the most recent lattice analysis of the Yang-Mills ghost sector) in conjunction with Schwinger-Dyson techniques [44], show that $1 + G(0) \neq 0$ and thus that there is nothing special about the ratio (6.11) (apart obviously from the fact that it constitutes a universal, albeit gauge-dependent, quantity).

VII. CONCLUSIONS

In this paper we have shown that the (R_ξ) BFM naturally emerges once the requirement of BRST and anti-BRST invariance of the action is fulfilled: indeed, background fields are unequivocally identified as the sources associated to the operator $s\bar{s}$. Correspondingly the existence of the anti-BRST symmetry implies background gauge invariance (and, consequently, a background Ward identity) as well as a (new) local antighost equation, which, when used in conjunction with the local ghost equation enforced by the BRST symmetry, completely determines the algebraic structure of the ghost sector of the theory for any gauge-fixing parameter. In addition, the background fields have been shown to be an extremum of the background effective action obtained by integrating out the quantum fields.

In hindsight, the correspondence

$$\text{BRST} + \text{anti-BRST} \equiv \text{BFM} \quad (7.1)$$

might not appear all that unexpected, as in the BFM background sector the ghost trilinear vertex is proportional to the sum of the ghost and antighost momentum while a quartic vertex involving two background fields and a ghost and an antighost (proportional to the metric tensor) is also generated. Thus the ghost and antighost are treated in a symmetric fashion, exactly as required by the BRST and anti-BRST invariance. It should be stressed however that Eq. (7.1) works only in the R_ξ gauges (which is anyway the only practically relevant case); choosing, e.g., a noncovariant background gauge breaks irremediably the anti-BRST symmetry of the theory.

In Chapter 15 of the second volume of his “Quantum Theory of Fields” [50], Weinberg noticed: “*The discovery of invariance under an ‘anti-BRST’ symmetry showed that,*

*despite appearances, there is a similarity between the roles of [the ghost field] ω^A and [the antighost field] ω^{*A} which remains somewhat mysterious.*”

We hope that this paper helps to shed some light on the mystery.

ACKNOWLEDGMENTS

One of us (A. Q.) thanks the European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT*) where part of this work has been carried out.

APPENDIX: FUNCTIONAL IDENTITIES IN THE PRESENCE OF SCALAR AND FERMIONIC MATTER FIELDS

We give here the relevant functional identities in the presence of scalar and fermionic matter fields. The ST identity takes the form

$$\begin{aligned} \int d^4x \left[\frac{\delta\Gamma}{\delta A_\mu^{*a}} \frac{\delta\Gamma}{\delta A_a^\mu} + \frac{\delta\Gamma}{\delta c_a^*} \frac{\delta\Gamma}{\delta c^a} + A_\mu^{*a} \frac{\delta\Gamma}{\delta \hat{A}_\mu^a} + c_a^\# \frac{\delta\Gamma}{\delta \hat{c}_a} \right. \\ \left. + \bar{c}_a^\# \frac{\delta\Gamma}{\delta b_a^\#} + b^a \frac{\delta\Gamma}{\delta \bar{c}^a} + \frac{\delta\Gamma}{\delta \phi^{*\dagger}} \frac{\delta\Gamma}{\delta \phi} - \frac{\delta\Gamma}{\delta \phi^*} \frac{\delta\Gamma}{\delta \phi^\dagger} \right. \\ \left. + \frac{\delta\Gamma}{\delta \bar{\psi}^*} \frac{\delta\Gamma}{\delta \psi} - \frac{\delta\Gamma}{\delta \psi^*} \frac{\delta\Gamma}{\delta \bar{\psi}} - \phi^\# \frac{\delta\Gamma}{\delta \hat{\phi}} + \phi^{*\dagger} \frac{\delta\Gamma}{\delta \hat{\phi}^{*\dagger}} \right. \\ \left. - \psi^\# \frac{\delta\Gamma}{\delta \hat{\psi}} + \bar{\psi}^\# \frac{\delta\Gamma}{\delta \hat{\psi}^\#} \right] = 0. \end{aligned} \quad (A1)$$

The anti-ST identity is

$$\begin{aligned} \int d^4x \left[\frac{\delta\Gamma}{\delta A_\mu^{*a}} \frac{\delta\Gamma}{\delta A_a^\mu} + \frac{\delta\Gamma}{\delta c_a^*} \frac{\delta\Gamma}{\delta c^a} + \frac{\delta\Gamma}{\delta \bar{c}_a^\#} \frac{\delta\Gamma}{\delta \bar{c}^a} + \frac{\delta\Gamma}{\delta b_a^\#} \frac{\delta\Gamma}{\delta b^a} \right. \\ \left. - A_\mu^{*a} \frac{\delta\Gamma}{\delta \hat{A}_\mu^a} - c_a^* \frac{\delta\Gamma}{\delta \hat{c}_a} + \frac{\delta\Gamma}{\delta \phi^{*\dagger}} \frac{\delta\Gamma}{\delta \phi} - \frac{\delta\Gamma}{\delta \phi^\#} \frac{\delta\Gamma}{\delta \phi^\dagger} \right. \\ \left. + \frac{\delta\Gamma}{\delta \bar{\psi}^*} \frac{\delta\Gamma}{\delta \psi} - \frac{\delta\Gamma}{\delta \psi^\#} \frac{\delta\Gamma}{\delta \bar{\psi}} + \phi^* \frac{\delta\Gamma}{\delta \hat{\phi}} - \phi^{*\dagger} \frac{\delta\Gamma}{\delta \hat{\phi}^{*\dagger}} \right. \\ \left. + \psi^* \frac{\delta\Gamma}{\delta \hat{\psi}} - \bar{\psi}^* \frac{\delta\Gamma}{\delta \hat{\psi}^*} \right] = 0. \end{aligned} \quad (A2)$$

Proceeding in the same way as for the derivation of Eqs. (4.6) and (4.8), we obtain the local antighost equation

$$\begin{aligned} \frac{\delta\Gamma}{\delta c^a} = -f^{abc} \frac{\delta\Gamma}{\delta b^b} \bar{c}^c - \xi \frac{\delta\Gamma}{\delta b_a^\#} + \hat{D}_\mu^{ab} \frac{\delta\Gamma}{\delta A_\mu^{*b}} + \mathcal{D}_\mu^{ab} A_b^{*\mu} \\ + f^{abc} c_b^* c^c + f^{abc} \hat{c}_b \frac{\delta\Gamma}{\delta c_c^\#} + f^{abc} b_b^\# \frac{\delta\Gamma}{\delta \bar{c}_c^\#} \\ + i \frac{\delta\Gamma}{\delta \phi^\#} t^a \hat{\phi} + i \hat{\phi}^{*\dagger} t^a \frac{\delta\Gamma}{\delta \phi^{*\dagger}} + i \frac{\delta\Gamma}{\delta \psi^\#} t^a \hat{\psi} \\ + i \hat{\psi} t^a \frac{\delta\Gamma}{\delta \bar{\psi}^\#} - i \phi^\dagger t^a \phi^* - i \phi^{*\dagger} t^a \phi \\ + i \bar{\psi} t^a \psi^* + i \bar{\psi}^* t^a \psi, \end{aligned} \quad (A3)$$

and the ghost equation

$$\begin{aligned} \frac{\delta\Gamma}{\delta\bar{c}^a} &= \mathcal{D}_\mu^{ab} A_b^{\#\mu} - \hat{\mathcal{D}}_\mu^{ab} \frac{\delta\Gamma}{\delta A_\mu^{*b}} - f^{abc} \hat{c}_b \frac{\delta\Gamma}{\delta c_c^*} + f^{abc} c_b^\# c^c \\ &+ f^{abc} \bar{c}_b^\# \bar{c}^c - f^{abc} b_b^\# b^c - i \frac{\delta\Gamma}{\delta\phi^*} t^a \hat{\phi} \\ &- i \hat{\phi}^{\#\dagger} t^a \frac{\delta\Gamma}{\delta\phi^{*\dagger}} - i \frac{\delta\Gamma}{\delta\psi^*} t^a \hat{\psi} - i \hat{\psi} t^a \frac{\delta\Gamma}{\delta\bar{\psi}^*} \\ &- i \phi^\dagger t^a \phi^\# - i \phi^{\#\dagger} t^a \phi + i \bar{\psi} t^a \psi^\# + i \bar{\psi}^\# t^a \psi. \end{aligned} \quad (\text{A4})$$

Finally the b equation becomes

$$\begin{aligned} \frac{\delta\Gamma}{\delta b^a} &= \hat{\mathcal{D}}_\mu^{ab} (A_b^\mu - \hat{A}_b^\mu) - \xi b^a - f^{abc} b_b^\# \bar{c}^c - c_a^\# - f^{abc} \hat{c}_b c^c, \\ &+ i \hat{\phi}^\dagger t^a \phi - i \hat{\phi} t^a \phi^\dagger + i \hat{\psi} t^a \psi - i \hat{\psi} t^a \bar{\psi}, \end{aligned} \quad (\text{A5})$$

while the background Ward identity yields

$$\begin{aligned} \mathcal{W}_a \Gamma &= -\mathcal{D}_\mu^{ab} \frac{\delta\Gamma}{\delta A_{b\mu}} - \hat{\mathcal{D}}_\mu^{ab} \frac{\delta\Gamma}{\delta \hat{A}_{b\mu}} + f^{abc} c_c \frac{\delta\Gamma}{\delta c_b} \\ &+ f^{abc} \bar{c}_c \frac{\delta\Gamma}{\delta \bar{c}_b} + f^{abc} b_c \frac{\delta\Gamma}{\delta b_b} + f^{abc} c_c^* \frac{\delta\Gamma}{\delta c_b^*} \\ &+ f^{abc} c_c^\# \frac{\delta\Gamma}{\delta c_b^\#} + f^{abc} \bar{c}_c^\# \frac{\delta\Gamma}{\delta \bar{c}_b^\#} + f^{abc} A_{\mu c}^* \frac{\delta\Gamma}{\delta A_{\mu b}^*} \\ &+ f^{abc} A_{\mu c}^\# \frac{\delta\Gamma}{\delta A_{\mu b}^\#} + f^{abc} \hat{c}_c \frac{\delta\Gamma}{\delta \hat{c}_b} + f^{abc} b_c^\# \frac{\delta\Gamma}{\delta b_b^\#} \\ &+ i t^a \phi \frac{\delta\Gamma}{\delta\phi} - i \phi^\dagger t^a \frac{\delta\Gamma}{\delta\phi^\dagger} + i t^a \psi \frac{\delta\Gamma}{\delta\psi} - i \bar{\psi} t^a \frac{\delta\Gamma}{\delta\bar{\psi}} = 0. \end{aligned} \quad (\text{A6})$$

-
- [1] B. S. DeWitt, *Phys. Rev.* **162**, 1195 (1967).
[2] J. Honerkamp, *Nucl. Phys.* **B48**, 269 (1972).
[3] R. E. Kallosh, *Nucl. Phys.* **B78**, 293 (1974).
[4] H. Kluberg-Stern and J. B. Zuber, *Phys. Rev. D* **12**, 482 (1975).
[5] I. Y. Arefeva, L. D. Faddeev, and A. A. Slavnov, *Theor. Math. Phys.* **21**, 1165 (1974).
[6] G. 't Hooft, in *Karpacz 1975, Proceedings, Acta Universitatis Wratislaviensis No. 368*, Vol. 1 (Wroclaw, 1976), pp. 345–369.
[7] S. Weinberg, *Phys. Lett.* **91B**, 51 (1980).
[8] G. M. Shore, *Ann. Phys. (N.Y.)* **137**, 262 (1981).
[9] L. F. Abbott, M. T. Grisaru, and R. K. Schaefer, *Nucl. Phys.* **B229**, 372 (1983).
[10] C. F. Hart, *Phys. Rev. D* **28**, 1993 (1983).
[11] L. F. Abbott, *Nucl. Phys.* **B185**, 189 (1981).
[12] L. F. Abbott, *Acta Phys. Pol. B* **13**, 33 (1982).
[13] S. Ichinose and M. Omote, *Nucl. Phys.* **B203**, 221 (1982).
[14] A. Denner, G. Weiglein, and S. Dittmaier, *Nucl. Phys.* **B440**, 95 (1995).
[15] S. Gates, M. T. Grisaru, M. Rocek, and W. Siegel, *Front. Phys.* **58**, 1 (1983).
[16] A. C. Aguilar and J. Papavassiliou, *J. High Energy Phys.* **12** (2006) 012.
[17] D. Binosi and J. Papavassiliou, *Phys. Rev. D* **77**, 061702 (2008).
[18] D. Binosi and J. Papavassiliou, *J. High Energy Phys.* **11** (2008) 063.
[19] D. Binosi and J. Papavassiliou, *Phys. Rep.* **479**, 1 (2009).
[20] P. A. Grassi, *Nucl. Phys.* **B462**, 524 (1996).
[21] P. A. Grassi, *Nucl. Phys.* **B537**, 527 (1999).
[22] P. A. Grassi, *Nucl. Phys.* **B560**, 499 (1999).
[23] C. Becchi and R. Collina, *Nucl. Phys.* **B562**, 412 (1999).
[24] R. Ferrari, M. Picariello, and A. Quadri, *Ann. Phys. (N.Y.)* **294**, 165 (2001).
[25] D. Binosi and A. Quadri, *Phys. Rev. D* **84**, 065017 (2011).
[26] D. Binosi and A. Quadri, *Phys. Rev. D* **85**, 085020 (2012).
[27] D. Binosi and A. Quadri, *Phys. Rev. D* **85**, 121702 (2012).
[28] G. Curci and R. Ferrari, *Nuovo Cimento A* **32**, 151 (1976).
[29] I. Ojima, *Prog. Theor. Phys.* **64**, 625 (1980).
[30] L. Baulieu and J. Thierry-Mieg, *Nucl. Phys.* **B197**, 477 (1982).
[31] P. A. Grassi, T. Hurth, and A. Quadri, *Phys. Rev. D* **70**, 105014 (2004).
[32] Y. Hatta, E. Iancu, L. McLerran, A. Stasto, and D. Triantafyllopoulos, *Nucl. Phys.* **A764**, 423 (2006).
[33] E. Iancu, A. Leonidov, and L. D. McLerran, *Nucl. Phys.* **A692**, 583 (2001).
[34] E. Ferreiro, E. Iancu, A. Leonidov, and L. McLerran, *Nucl. Phys.* **A703**, 489 (2002).
[35] G. Barnich, F. Brandt, and M. Henneaux, *Phys. Rep.* **338**, 439 (2000).
[36] G. Barnich, F. Brandt, and M. Henneaux, *Commun. Math. Phys.* **174**, 93 (1995).
[37] G. Barnich, F. Brandt, and M. Henneaux, *Commun. Math. Phys.* **174**, 57 (1995).
[38] J. Gomis, J. Paris, and S. Samuel, *Phys. Rep.* **259**, 1 (1995).
[39] A. Quadri, *J. High Energy Phys.* **05** (2002) 051.
[40] N. Nakanishi, *Prog. Theor. Phys.* **60**, 284 (1978).
[41] I. Jack and H. Osborn, *Nucl. Phys.* **B249**, 472 (1985).
[42] M. Tissier and N. Wschebor, *Phys. Rev. D* **79**, 065008 (2009).
[43] T. Kugo, [arXiv:hep-th/9511033](https://arxiv.org/abs/hep-th/9511033).
[44] A. Aguilar, D. Binosi, and J. Papavassiliou, *J. High Energy Phys.* **11** (2009) 066.

- [45] V.M.M. Schaden, D. Zwanziger, and R. Alkofer, [arXiv:1309.0497](#) [Phys. Rev. D (to be published)].
- [46] P.A. Grassi, T. Hurth, and M. Steinhauser, *Ann. Phys. (N.Y.)* **288**, 197 (2001).
- [47] D. Binosi and J. Papavassiliou, *Phys. Rev. D* **66**, 025024 (2002).
- [48] T. Kugo and I. Ojima, *Prog. Theor. Phys. Suppl.* **66**, 1 (1979).
- [49] A. Cucchieri and T. Mendes, [arXiv:1308.1283](#) [Phys. Rev. D (to be published)].
- [50] S. Weinberg, *The Quantum Theory of Fields. Vol 2: Modern Applications* (Cambridge University Press, Cambridge, England, 1996), p. 489.