

Hydrodynamic approach to QGP instabilities

E. Calzetta* and J. Peralta-Ramos†

Departamento de Física, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires and IFIBA, CONICET, Ciudad Universitaria, Buenos Aires 1428, Argentina
(Received 25 September 2013; published 13 November 2013)

We show that the usual linear analysis of quark-gluon plasma Weibel instabilities based on the Maxwell–Boltzmann equation may be reproduced in a purely hydrodynamic model. The latter is derived by the entropy production variational method from a transport equation including collisions and can describe highly nonequilibrium flow. We find that, as expected, collisions slow down the growth of Weibel instabilities. Finally, we discuss the strong momentum anisotropy limit.

DOI: [10.1103/PhysRevD.88.095010](https://doi.org/10.1103/PhysRevD.88.095010)

PACS numbers: 52.27.Ny, 52.35.-g, 47.75.+f, 25.75.-q

I. INTRODUCTION

The hot and dense fireball of nuclear matter created in heavy ion collisions at the Relativistic Heavy Ion Collider and the Large Hadron Collider behaves as an almost perfect fluid, with a viscosity-to-entropy ratio not far from the lower bound $\eta/s = 1/(4\pi)$ derived from the anti-de Sitter/conformal field theory correspondence [1]. This implies that, in spite of the rapid longitudinal expansion, which limits the effectiveness of particle collision in equilibrating the system, the fireball isotropizes extremely fast with characteristic times $\lesssim 1$ fm/c.

Mrówczyński was the first to show that there exists an instability to chromomagnetic fluctuations in the quark-gluon plasma (QGP) for wave vectors transverse to the chromomagnetic field [2,3] (see also Refs. [4–6]). These are the analog of the Weibel instabilities in a electromagnetic (EM) plasma [7] and occur if the one-particle distribution is anisotropic in momentum space (see also Refs. [8–10]). This type of distribution function is relevant in heavy ion collisions because the longitudinal expansion of the fireball causes the system to become much colder in the longitudinal direction than in the transverse ones, implying $\langle p_L^2 \rangle \ll \langle p_T^2 \rangle$ in the local rest frame, where $p_{L,T}$ is the particle momentum in the longitudinal and transverse direction.

The transverse instabilities are found to have a strong impact on the fireball’s evolution due to the large momentum anisotropy present at early times, speeding up the isotropization and equilibration process. Basically, the instability causes the soft sector of the magnetic field to become rapidly amplified, which leads to large-angle scattering of hard particles, thus leading to faster isotropization and thermalization. Therefore, the thermalization process in the QGP is not controlled solely by particle collisions, but the role of collective effects (such as Weibel instabilities) must be taken into account as well.

The chromo-Weibel instability has been extensively studied [2–6,11–24], both analytically in linear response

and numerically in the fully nonlinear case, either within the hard-thermal loop framework or directly from the nonlinear Vlasov equations for anisotropic plasmas (which go beyond the hard-thermal loop approximation; see, e.g., Ref. [25] and references therein). One of the most important results that comes out from these studies is that the nonlinear gauge self-interactions slow down the growth rate of the Weibel instability, an effect that could be studied with a suitable generalization to the non-Abelian case of the formalism presented in this work.

More directly connected to our study is the work by Schenke and coworkers [11], in which the authors study stable and unstable modes as obtained from the Boltzmann–Vlasov equation for non-Abelian fields (but in the effective Abelian approximation) with a Bhatnagar–Gross–Krook collision operator. We shall compare our results to those of Ref. [11] along the way.

The usual kinetic theory approach to study plasma instabilities combines a collisionless transport equation for a one-particle distribution function with the Maxwell equations for the EM field; see, e.g., Refs. [26,27]. In the background, the distribution function is position independent, and the Maxwell field vanishes (see, however, Ref. [12], where initial fluctuations of the currents are taken into account). The linearized equations are a simple transport equation for the perturbation of the distribution function with a source which depends on the linearized Maxwell field. This equation may be solved exactly, leading to an integrodifferential equation for the Maxwell field. Since the background is homogeneous, the perturbations may be Fourier analyzed, whereby a dispersion relation follows.

In this paper we show that the ordinary analysis of QGP instabilities based on the Maxwell–Boltzmann equation may be reproduced in a purely (viscous) hydrodynamic model. To our knowledge, this is the first attempt of this kind that deals with a relativistic plasma and includes the effect of collisions. Manuel and Mrówczyński have derived ideal hydrodynamiclike equations that are applicable to short-time scale color phenomena in the QGP and applied them to study the collective modes in a two-stream system [28]. An alternative set of hydrodynamic equations for a

*calzetta@df.uba.ar

†jperalta@df.uba.ar

non-Abelian plasma is derived in Ref. [29]. Using the chromohydrodynamic formalism developed in Ref. [28], Mannarelli and Manuel investigated jet-induced stream instabilities in Ref. [30] and compared the results with those obtained with kinetic theory in Ref. [31]. Based on a particular closure and neglecting collisions, Basu has studied the Weibel instabilities in a nonrelativistic plasma within a hydrodynamic model and has been able to reproduce, in the strong anisotropy limit, the results from kinetic theory [32].

For simplicity, we shall restrict ourselves to the instability of a homogeneous configuration of an EM plasma (i.e., the background EM field and the background one-particle distribution function are independent of spatial coordinates). We shall reproduce the kinetic theory analysis with a hydrodynamical effective theory derived from kinetic theory by use of the so-called entropy production variational method (EPVM)—see Refs. [33,34] for a detailed account of this method and Refs. [35,36] for its connection to nonequilibrium statistical mechanics. We note that similar approaches to the EPVM have been applied to diverse transport phenomena in Refs. [37–41].

As indicated above, we shall deal with an Abelian plasma. The extension to a non-Abelian plasma is left for the future, but it is worth mentioning that the developments presented here are relevant to the non-Abelian case under the Abelian dominance approximation valid for weak gauge fields $A \ll p_h$, where A and p_h are the vector potential amplitude and the characteristic momentum of hard particles (see, e.g., Refs. [11,12,22]).

This paper is organized as follows. In Sec. II we give a brief overview of the kinetic theory and hydrodynamics of a charged fluid and describe the closure obtained from the EPVM that we use to go from kinetic theory to the hydrodynamic effective theory on which our developments are based. In Sec. III we analyze the Weibel instability as obtained from the effective theory, study the dependence of its growth rate on the collision time, and compare our results to those obtained from kinetic theory. Finally, in Sec. IV we present our conclusions and outlook.

II. THEORETICAL SETUP

A. Kinetic theory and hydrodynamics of a charged fluid

Let us begin by briefly reviewing the kinetic theory and hydrodynamics of a charged fluid.

The fluid has two conserved quantities, the energy-momentum tensor $T^{\mu\nu}$ and current J^μ . They obey conservation laws and the Maxwell equations

$$T_{,\nu}^{\mu\nu} = F^{\mu\rho} J_\rho \quad J_{,\mu}^\mu = 0 \quad F_{,\nu}^{\mu\nu} = J^\mu, \quad (1)$$

where $F^{\mu\nu}$ is the strength tensor

$$F^{\mu\nu} = \begin{pmatrix} 0 & E^1 & E^2 & E^3 \\ -E^1 & 0 & B^3 & -B^2 \\ -E^2 & -B^3 & 0 & B^1 \\ -E^3 & B^2 & -B^1 & 0 \end{pmatrix}. \quad (2)$$

For an ideal fluid (which is not the case studied here), we would have

$$T^{\mu\nu} = (\rho + p)u^\mu u^\nu + pg^{\mu\nu} \quad J^\mu = qu^\mu, \quad (3)$$

where ρ is the energy density, p is the pressure, and q is the charge density.

In the kinetic theory description, the transport equation reads

$$p^\mu \left[\partial_\mu f - eF_{\mu\nu} \frac{\partial f}{\partial p_\nu} \right] = \frac{-1}{\tau} \text{sign}(p^0) I_{\text{col}}, \quad (4)$$

where $f = f(x^\mu, p^\mu, t)$ is the one-particle distribution function, e and p^μ are the particles' charges and momenta, I_{col} is the collision operator, and τ is the collision time. We shall specify I_{col} later on. The current is

$$J^\mu = e \int Dp p^\mu f, \quad (5)$$

and the energy-momentum tensor is

$$T^{\mu\nu} = \int Dp p^\mu p^\nu f, \quad (6)$$

where the measure is given by

$$Dp = \frac{2d^4 p \delta(p^2)}{(2\pi)^3} = \frac{d^4 p}{(2\pi)^3 p} (\delta(p^0 - p) + \delta(p^0 + p)), \quad (7)$$

For simplicity we assume massless particles. To enforce the conservation laws, we require

$$\int Dp \text{sign}(p^0) I_{\text{col}} = \int Dp \text{sign}(p^0) p^\mu I_{\text{col}} = 0. \quad (8)$$

We parametrize the distribution function as follows:

$$f = f_B(1 + Z), \quad (9)$$

with $Z = 0$ at equilibrium. f_B is the background solution. Following Weibel [7], we assume f_B is independent of spatial coordinates. Later on we shall specify f_B , but at the moment we leave this choice open.

Our developments will be based on the EPVM, so we shall make use of the expression giving the entropy production. We assume a simple Boltzmann-type relative entropy flux [42–44]

$$S^\mu = - \int Dp (\text{sign}(p^0)) p^\mu \left[f \ln \left(\frac{f}{f_B} \right) - (f - f_B) \right], \quad (10)$$

so we get the relative (with respect to f_B) entropy production

$$S_{,\mu}^{\mu} = \frac{1}{\tau} \int Dp I_{\text{col}} \ln(1 + Z). \quad (11)$$

To continue we must make an ansatz regarding the collision operator. We shall assume a linear collision term of the form

$$I_{\text{col}}(p) = \int Dp' K[p, p'] Z(p'), \quad (12)$$

where K is a symmetric operator. The idea behind Eq. (12) is that the antisymmetric part of the kernel of the linear collision operator does not affect the entropy production to lowest order. Retaining only the symmetric kernel in the linear collision term, we focus on the part of the dynamics which is directly related to the relaxation of the system toward homogeneity, since this relaxation entails an entropy increase. The conservation laws then mean that $\text{sign}(p^0)$ and $\text{sign}(p^0)p^{\mu}$ are homogeneous solutions. This suggests writing

$$I_{\text{col}}(p) = F(p) f_B(p) \{ Z(p) - \text{sign}(p^0) [A[Z] + p^{\mu} B_{\mu}[Z]] \}, \quad (13)$$

where F is some even function of p^i ($i = 1, 2, 3$), and the quantities $A[Z]$ and $B_{\mu}[Z]$ are functionals of Z introduced to enforce the constraints, namely,

$$\begin{aligned} \int Dp (A + B_{\nu} p^{\nu}) F f_B &= \int Dp \text{sign}(p^0) F f_B Z \\ \int Dp p^{\mu} (A + B_{\nu} p^{\nu}) F f_B &= \int Dp \text{sign}(p^0) p^{\mu} F f_B Z. \end{aligned} \quad (14)$$

We shall not need the explicit expressions for $A[Z]$ and $B_{\mu}[Z]$, but they can be calculated following Ref. [29]. For the moment, we need not specify the function $F(p)$, but we note that $F = p$ corresponds to Anderson–Witting’s ansatz for the collision term [45,46], which we shall use in Sec. III B.

B. Closure from the EPVM

To go from kinetic theory to a fluid description, we need to express Z (the deviation from equilibrium) in terms of hydrodynamic variables, i.e., to provide a closure. The most well-known approaches are the Chapman–Enskog expansion and Grad’s quadratic ansatz. As mentioned in the introduction, in this paper we will use the closure that is obtained from the EPVM (a review can be found in Ref. [33]). We will now briefly review this formalism, focusing on the derivation of the hydrodynamic effective theory that will serve as our basis. A detailed account of this derivation can be found in Refs. [29,47]; see also Refs. [48,49] for concrete applications to heavy ion collisions.

Traditional fluid dynamics derived from kinetic theory by the Chapman–Enskog expansion or Grad’s ansatz has

two important limitations. First, it relies on an expansion in gradients of hydrodynamic variables, which necessarily implies that the system is sufficiently close to equilibrium so that these gradients are small. In turn, this means that the system is also very close to being isotropic in momentum space. Second, it breaks down at large shear viscosity η . The formalism obtained from the EPVM does not suffer from these drawbacks and can successfully track the evolution as given by kinetic theory even for highly nonequilibrium flow. By direct comparison to solutions to Boltzmann’s equation, we have shown in Ref. [48] that for the boost-invariant one-dimensional expansion of matter created in heavy ion collisions the model obtained from the EPVM can reproduce the kinetic theory results, even for highly nonequilibrium situations and/or large values of the shear viscosity of the QGP. Instead, it is well known that the full Israel–Stewart formalism [50–52] (based on Grad’s ansatz) fails in these cases because the pressure becomes negative due to the large values of the shear tensor (see, e.g., Ref. [53]).

If deviations from equilibrium are small, the EPVM closure reduces to Grad’s quadratic ansatz [47]. The effective theory includes nonhydrodynamic variables in addition to the usual hydrodynamic ones (we shall call them $\zeta_{\lambda\sigma}$ and ζ_{λ} in what follows). These variables model the backreaction of f —which may describe a highly nonequilibrium situation—on the hydrodynamic modes which relax much more slowly. In other words, on time scales short with respect to τ , the fluid relaxes to a steady nonequilibrium state characterized by a nonvanishing viscous energy momentum tensor; the relaxation to true equilibrium is a much slower process.

In the EPVM, $\zeta_{\lambda\sigma}$ and ζ_{λ} are identified with the Lagrange multipliers of the variational problem for which the solution gives the f that extremizes the entropy production given fixed values of $T^{\mu\nu}$ and J^{μ} . Physically, one can think of the EPVM as selecting the dynamics of these nonhydrodynamic variables in such a way as to extremize the production of entropy during the evolution of the system [33,34,47].

The EPVM leads, to first order in Lagrange multipliers, to the equation (see Refs. [29,47] for details)

$$2I_{\text{col}}[Z] = \tau f_B [\zeta_{\lambda\sigma} p^{\lambda} p^{\sigma} + \zeta_{\lambda} p^{\lambda}]. \quad (15)$$

Consistency requires

$$\begin{aligned} \int Dp f_B \text{sign}(p^0) [\zeta_{\lambda\sigma} p^{\lambda} p^{\sigma} + \zeta_{\lambda} p^{\lambda}] &= 0 \\ \int Dp f_B \text{sign}(p^0) p^{\mu} [\zeta_{\lambda\sigma} p^{\lambda} p^{\sigma} + \zeta_{\lambda} p^{\lambda}] &= 0. \end{aligned} \quad (16)$$

We now get

$$Z = \frac{\tau}{2F} [\zeta_{\lambda\sigma} p^{\lambda} p^{\sigma} + \zeta_{\lambda} p^{\lambda}] + Z_{\text{hom}}, \quad (17)$$

where

$$Z_{\text{hom}} = \text{sign}(p^0)[\alpha + \beta_\lambda p^\lambda]. \quad (18)$$

Inserting this into the linearized transport equation, we get

$$\begin{aligned} p^\mu \left[f_B \partial_\mu \left\{ \frac{\tau}{2F} [\zeta_{\lambda\sigma} p^\lambda p^\sigma + \zeta_\lambda p^\lambda] + \text{sign}(p^0)[\alpha + \beta_\lambda p^\lambda] \right\} \right. \\ \left. - e F_{\mu\nu} \frac{\partial f_B}{\partial p_\nu} \right] \\ = \frac{-1}{2} \text{sign}(p^0) f_B [\zeta_{\lambda\sigma} p^\lambda p^\sigma + \zeta_\lambda p^\lambda]. \end{aligned} \quad (19)$$

The idea is to get equations for $\zeta_{\lambda\sigma}$, ζ_λ , β_λ , and α by taking moments of this equation [29,54,55]. We shall do this at linear order in the next section.

III. WEIBEL INSTABILITY

To focus on the physics of unstable modes, we assume $f_B = f(\vec{p})\theta(p^0)$, where f is even but anisotropic (as in Weibel's seminal paper [7]). More concretely, $f = C(\beta, b)f_{\beta b}(p)$, where

$$f_{\beta b} = e^{-\beta p} e^{-b^2 p_z^2} \quad (20)$$

as advocated in Ref. [8]. $C(\beta, b)$ is a normalization factor to be fixed later on, and b^2 is a measure of the momentum anisotropy of the background configuration.

Note that considering the collisionless transport equation in the usual approach corresponds to the $\tau \rightarrow \infty$ limit. In the absence of a Maxwell field, in this limit only terms containing derivatives of the hydrodynamic fields remain in the equations of motion. In other words, any space-time independent configuration of fields α , β^μ , ζ^μ , and $\zeta^{\mu\nu}$ will be a solution of the hydrodynamic equations, provided the EM current vanishes (since we want to avoid having a background EM field). To achieve this, we assume that there is a compensating static background. On the other hand, it makes no sense to linearize in τ (this rules out the possibility of using the closure provided by Grad's ansatz or the Chapman–Enskog expansion). For this reason, we shall not write down the full equations of motion but derive the linearized equations directly from kinetic theory instead.

To be even more concrete and to simplify matters, we seek solutions with ($a = 1, 2$) $\zeta_{00} = \zeta_{ab} = \zeta_{0a} = \zeta_{03} = \zeta_{33} = 0$, $\zeta_\lambda = \alpha = 0$, $\beta_0 = \beta_3 = 0$. We further assume a plane wave solution propagating in the z direction; namely, the space-time dependence of all factors is of the form $\exp\{i(kz - \omega t)\}$. The consistency conditions hold, and the transport equation becomes

$$\begin{aligned} i f_{\beta b} (k p^3 - \omega p) \left\{ \frac{\tau}{F} \zeta_{b3} p^b p^3 + \beta_b p^b \right\} - e p^\mu F_{\mu\nu} \frac{\partial f_{\beta b}}{\partial p_\nu} \\ = -f_{\beta b} \zeta_{b3} p^b p^3. \end{aligned} \quad (21)$$

Observe that the normalization drops out.

To compute the moments of this equation, observe that

$$\begin{aligned} \int Dp p^\mu F_{\mu\nu} \frac{\partial f_{\beta b}}{\partial p_\nu} &= 0 \\ \int Dp p_\lambda p^\mu F_{\mu\nu} \frac{\partial f_{\beta b}}{\partial p_\nu} &= - \int Dp p^\mu F_{\mu\lambda} f_{\beta b} \\ \int Dp p_\lambda p_\rho p^\mu F_{\mu\nu} \frac{\partial f_{\beta b}}{\partial p_\nu} &= - \int Dp p^\mu \\ &\quad \times [F_{\mu\lambda} p_\rho + F_{\mu\rho} p_\lambda] f_{\beta b}. \end{aligned} \quad (22)$$

We introduce the notation

$$\langle A \rangle = \int \frac{d^3 p}{(2\pi)^3 p} f_{\beta b} A. \quad (23)$$

The nontrivial first order moments of the kinetic equation are

$$\begin{aligned} ik\tau \zeta_{b3} \left\langle \frac{1}{F} p^a p^b p_3^2 \right\rangle - i\omega \beta_b \langle p p^a p^b \rangle + e F_{0a} \langle p \rangle &= 0 \\ F_{03} \langle p \rangle &= 0. \end{aligned} \quad (24)$$

The nontrivial second order moments are

$$\begin{aligned} ik\tau \zeta_{b3} \left\langle \frac{1}{F} p p^a p^b p_3^2 \right\rangle - i\omega \beta_b \langle p^2 p^a p^b \rangle + e [F_{b0} \langle p^a p^b \rangle \\ + F_{0a} \langle p^2 \rangle] &= 0 \\ ik\beta_b \langle p^a p^b p_3^2 \rangle - i\omega \tau \zeta_{b3} \left\langle \frac{1}{F} p p^a p^b p_3^2 \right\rangle \\ + e [F_{3a} \langle p_3^2 \rangle + F_{b3} \langle p^a p^b \rangle] &= -\zeta_{b3} \langle p^a p^b p_3^2 \rangle. \end{aligned} \quad (25)$$

We shall only use the second of these equations. Finally the current is

$$J^\mu = e C(\beta, b) \left\langle p^\mu \left[\frac{\tau}{F} \zeta_{b3} p^b p^3 + \beta_b p^b \right] \right\rangle. \quad (26)$$

The only nonzero component of the current is

$$J^a = e C(\beta, b) \beta_b \langle p^a p^b \rangle \quad (27)$$

So, the nontrivial Maxwell equations are

$$i\omega F_{a0} + ik F_{a3} = J_a = e C(\beta, b) \beta_b \langle p_a p^b \rangle \quad (28)$$

and

$$-i\omega F_{a3} - ik F_{a0} = 0. \quad (29)$$

To write the final forms of the equations, we observe that

$$\langle p^a p^b A(p, p^3) \rangle = \frac{1}{2} \delta^{ab} \langle (p^2 - p_3^2) A(p, p^3) \rangle. \quad (30)$$

Let us call

$$\begin{aligned} \langle p \rangle &= \frac{n}{C} \quad \frac{1}{2} C \langle (p^2 - p_3^2) \rangle = p_x \\ \langle p^2 - 3p_3^2 \rangle &= \frac{2}{C} (p_x - p_z) \end{aligned} \quad (31)$$

and

$$\begin{aligned} \langle p(p^2 - p_3^2) \rangle &= A_3 & \langle (p^2 - p_3^2)p_3^2 \rangle &= A_4 \\ \left\langle \frac{1}{F}(p^2 - p_3^2)p_3^2 \right\rangle &= A_{4F} & \left\langle \frac{1}{F}p(p^2 - p_3^2)p_3^2 \right\rangle &= A_{5F}. \end{aligned} \quad (32)$$

Therefore,

$$\begin{pmatrix} \omega & k & iep_x & 0 \\ k & \omega & 0 & 0 \\ -i2e\frac{n}{C} & 0 & \omega A_3 & -k\tau A_{4F} \\ 0 & \frac{2ie}{C}(p_x - p_z) & -kA_4 & iA_4 + \omega\tau A_{5F} \end{pmatrix} \times \begin{pmatrix} F_{a0} \\ F_{a3} \\ \beta_a \\ \zeta_{a3} \end{pmatrix} = 0. \quad (33)$$

If $k = 0$, the dispersion relation is

$$[iA_4 + \omega\tau A_{5F}]\omega \left[A_3\omega^2 - 2e^2\frac{n}{C}p_x \right] = 0. \quad (34)$$

Thus, we identify the plasma frequency

$$\omega_p^2 = \frac{2e^2n}{CA_3}p_x. \quad (35)$$

In general, the dispersion relation is

$$\begin{aligned} [iA_4 + \omega\tau A_{5F}]A_3\omega[\omega^2 - k^2 - \omega_p^2] \\ - k^2\tau A_{4F} \left[A_4(\omega^2 - k^2) + \frac{2e^2p_x}{C}(p_x - p_z) \right] = 0. \end{aligned} \quad (36)$$

Let us call

$$k_{st}^2 = \frac{2e^2p_x}{CA_4}(p_x - p_z). \quad (37)$$

Note that k_{st} is a measure of the anisotropy of the background distribution function f_B .

Defining

$$\tau_0 = \frac{A_4}{A_{5F}} \quad (38)$$

$$\lambda = \frac{A_{4F}A_4}{A_3A_{5F}} \quad (39)$$

and putting

$$\omega = i\sigma, \quad (40)$$

the dispersion relation becomes

$$\begin{aligned} P[\sigma] &= \sigma^4 + \frac{\tau_0}{\tau}\sigma^3 + \{k^2 + \omega_p^2 + \lambda k^2\}\sigma^2 \\ &+ \frac{\tau_0}{\tau}[k^2 + \omega_p^2]\sigma - \lambda k^2\{k_{st}^2 - k^2\} = 0. \end{aligned} \quad (41)$$

It is clear that $P[\sigma]$ increases with σ when σ is real and positive. Therefore, for $0 < k^2 < k_{st}^2$ there is one (and only one) real positive root. This root corresponds to the Weibel instability, on which we focus in what follows.

A. Collision time dependence

To investigate how the growth rate of the Weibel instability depends on τ , observe that if $\sigma(\tau)$ denotes the root for a given τ , then

$$P'[\sigma(\tau)]\frac{d\sigma}{d\tau} - \frac{\tau_0}{\tau^2}P_2[\sigma(\tau)] = 0, \quad (42)$$

where

$$P_2[\sigma] = \sigma[\sigma^3 + k^2 + \omega_p^2]. \quad (43)$$

Since P' and P_2 are positive for positive σ , we must have $d\sigma/d\tau \geq 0$. Also write

$$P[\sigma] = P_1[\sigma] + \frac{\tau_0}{\tau}P_2[\sigma] \quad (44)$$

$$P_1[\sigma] = \sigma^4 + \{k^2 + \omega_p^2 + \lambda k^2\}\sigma^2 - \lambda k^2\{k_{st}^2 - k^2\} = 0. \quad (45)$$

Then we must have

$$P_1[\sigma(\tau)] \leq 0, \quad (46)$$

which implies $\sigma(\tau) \leq \sigma_{\max}$, where

$$\begin{aligned} \sigma_{\max}^2 &= \frac{1}{2} \left[\sqrt{4\lambda k^2\{k_{st}^2 - k^2\} + \{k^2 + \omega_p^2 + \lambda k^2\}^2} \right. \\ &\quad \left. - \{k^2 + \omega_p^2 + \lambda k^2\} \right]. \end{aligned} \quad (47)$$

Since $\sigma_{\max}^2 = 0$ both for $k^2 = 0$ and $k^2 = k_{st}^2$, it must have a maximum in k . The maximum is achieved when

$$(1 + \lambda)\sigma_{\max}^2 - \lambda k_{st}^2 + 2\lambda k^2 = 0. \quad (48)$$

We have $\sigma(\tau) \rightarrow \sigma_{\max}$ for $\tau \rightarrow \infty$, which corresponds to the collisionless limit usually assumed to study the Weibel instability. This means that, as expected on physical grounds, collisions slow down the growth of the unstable mode. The particles with momentum orthogonal to the magnetic field, which are responsible for the Weibel instability, can scatter with other particles and avoid getting trapped in the background magnetic field, so a smaller current and therefore a smaller induced magnetic field are generated. This has been verified directly from kinetic theory with a Bhatnagar-Gross-Krook collision term [11].

For $\tau \rightarrow 0$, corresponding to the ideal fluid limit, we get instead

$$\sigma(\tau) = \frac{\tau}{\tau_0} \frac{\lambda k^2}{k^2 + \omega_p^2} (k_{\text{st}}^2 - k^2). \quad (49)$$

In this regime, the maximum growth rate is achieved at

$$k^2 = \omega_p^2 \left[\sqrt{1 + \frac{k_{\text{st}}^2}{\omega_p^2}} - 1 \right] \quad (50)$$

and takes the value

$$\sigma(\tau) = \frac{\tau}{\tau_0} \lambda \omega_p^2 \left[\sqrt{1 + \frac{k_{\text{st}}^2}{\omega_p^2}} - 1 \right]^2. \quad (51)$$

This shows that the instability disappears for all practical purposes when τ is small enough because there is a maximum time scale over which the theory makes sense given by the time scale on which the background solution relaxes to equilibrium. This feature is also present in the kinetic theory approach of Ref. [11].

B. Anisotropy dependence

We wish to discuss now how the instability growth rate depends on the parameters β and b . To obtain analytic results, we shall restrict our discussion to the strong anisotropy limit $b \rightarrow \infty$. We use the Anderson–Witting collision term [45,46], which corresponds to $F = p$.

Let us define

$$\begin{aligned} \langle 1 \rangle &= \int \frac{d^3 p}{(2\pi)^3} p^{-1} e^{-\beta p} e^{-b p^2} \\ &= \frac{1}{(2\pi)^2} \int_0^\infty p dp \int_0^\pi \sin \theta d\theta e^{-\beta p} e^{-b^2 p^2 \cos^2 \theta}. \end{aligned} \quad (52)$$

The idea is that all the necessary expectation values may be obtained as derivatives of $\langle 1 \rangle$ according to the rule

$$\langle p^A (p_z^2)^B \rangle = (-1)^{A+B} \frac{\partial^{A+B}}{\partial \beta^A \partial (b^2)^B} \langle 1 \rangle. \quad (53)$$

Note, however, that for the Anderson–Witting collision term [45,46], A_{4F} cannot be obtained this way but rather from the solution to the equation

$$-\frac{\partial}{\partial \beta} A_{4F} = A_4. \quad (54)$$

Call $q = b/\beta$

$$\langle 1 \rangle = \frac{1}{2\pi^2 \beta^2} \int_0^\infty t dt \int_0^1 dx e^{-t} e^{-q^2 t^2 x^2}. \quad (55)$$

Following Ref. [56], we define the error function

$$\Phi[z] = \text{erf}[z] = \frac{2}{\sqrt{\pi}} \int_0^z dt e^{-t^2}, \quad (56)$$

so we get

$$\langle 1 \rangle = \frac{1}{4\pi^{3/2} \beta b} \left[1 - \Phi \left[\frac{1}{2q} \right] \right] e^{1/4q^2}. \quad (57)$$

Note that no approximation regarding the value of b has been done to get Eq. (57).

In the strong anisotropy limit $b \rightarrow \infty$, Eq. (57) reduces to

$$\langle 1 \rangle_\infty = \frac{1}{4\pi^{3/2} \beta (b^2)^{1/2}}. \quad (58)$$

In this regime we obtain (in all cases to leading order in b^{-1})

$$\begin{aligned} \langle p \rangle &= \frac{n}{C} = \frac{1}{4\pi^{3/2}} \frac{1}{\beta^2 b} \\ \langle p^2 \rangle &= \frac{2p_x}{C} = \frac{1}{4\pi^{3/2}} \frac{2}{\beta^3 b} \\ \langle p^3 \rangle &= A_3 = \frac{1}{4\pi^{3/2}} \frac{6}{\beta^4 b} \\ \langle p^2 p_z^2 \rangle &= A_4 = A_{5F} = \frac{1}{4\pi^{3/2}} \frac{1}{\beta^3 b^3} \\ \langle p p_z^2 \rangle &= A_{4F} = \frac{1}{4\pi^{3/2}} \frac{1}{2\beta^2 b^3}. \end{aligned} \quad (59)$$

We wish to consider configurations with different anisotropy parameter $\xi = b^2/\beta^2$ but the same β and particle density $n = n_0$. Thus, we define

$$C = \frac{n_0}{\langle p \rangle} = 4\pi^{3/2} n_0 \beta^2 b. \quad (60)$$

In this case

$$\omega_p^2 = \frac{1}{3} e^2 n_0 \beta, \quad (61)$$

and $\tau_0 = 1$. The quantity k_{st}^2 is unbounded,

$$k_{\text{st}}^2 = 2(n_0 \beta^3) \frac{b^2}{\beta^4}. \quad (62)$$

Instead λ goes to zero,

$$\lambda = \frac{\beta^2}{12b^2}. \quad (63)$$

Note that the product $\lambda k_{\text{st}}^2 = n_0 \beta / 6$ remains bounded. If τ is small [but not so small as to make the Weibel instability disappear—see the discussion after Eq. (51)], we have from Eq. (49) that

$$\sigma(\tau) = \tau [\lambda k_{\text{st}}^2] \frac{k^2}{k^2 + \omega_p^2}. \quad (64)$$

The behavior of σ with increasing k that we obtain agrees well with the kinetic theory results of Ref. [11] (see Fig. 11 of that reference). The growth rate increases with increasing k as given by Eq. (64), and for large k it saturates at $\tau[\lambda k_{\text{st}}^2]$.

IV. CONCLUSIONS

We have studied the Weibel instability within the framework of a hydrodynamic effective theory derived by the entropy production variational method from the relativistic Maxwell–Boltzmann equation. This method provides a fluid closure that yields an effective theory capable of describing highly nonequilibrium flows.

We have found that the usual linear analysis of QGP Weibel instabilities based on the Maxwell–Boltzmann equation may be reproduced in this particular hydrodynamic model. We have analyzed the dependence of the growth rate of the Weibel instability on the collision time and found that, as expected, the effect of collisions is to slow down the growth of the magnetic field’s amplitude. We have also shown that if the collision time is too short, corresponding to a nearly perfect fluid, the Weibel instability disappears. Our results agree with those obtained from kinetic theory with a BGK collision term [11].

We believe that this study opens up the possibility of investigating the effect of nonlinear self-interactions of the gauge fields on the chromo-Weibel instability relevant to QGP dynamics, within a framework that is simpler than the kinetic theory of non-Abelian plasmas including particle collisions. This may be useful to investigate some issues which are hard to attack from kinetic theory, such as the backreaction of Weibel instabilities on the dynamics of the rapidly expanding QGP [16–18,22]. To this end, we must rely on the extension to non-Abelian plasmas of the effective hydrodynamic formalism presented here, which we carried out in Ref. [29], and study the development of the chromo-Weibel instability in this setting. Work is in progress along this line.

ACKNOWLEDGMENTS

This work has been supported in part by ANPCyT, CONICET and UBA under Project No. UBACYT X032 (Argentina).

-
- [1] U. W. Heinz, in *Relativistic Heavy Ion Physics*, edited by R. Stock, Landolt-Boernstein, New Series, Group I Vol. 23 (Springer Verlag, New York, 2010), Chap. 5.
- [2] S. Mrówczyński, *Phys. Lett. B* **314**, 118 (1993).
- [3] S. Mrówczyński, *Phys. Rev. C* **49**, 2191 (1994).
- [4] S. Mrówczyński, *Nucl. Phys.* **A820**, 95c (2009).
- [5] S. Mrówczyński, *Phys. Rev. D* **77**, 105022 (2008).
- [6] S. Mrówczyński, *Nucl. Phys.* **A774**, 149 (2006).
- [7] E. S. Weibel, *Phys. Rev. Lett.* **2**, 83 (1959).
- [8] R. Schlickeiser, *Phys. Plasmas* **11**, 5532 (2004).
- [9] A. Achterberg and J. Wiersma, *Astron. Astrophys.* **475**, 1 (2007).
- [10] A. Achterberg, J. Wiersma, and C. A. Norman, *Astron. Astrophys.* **475**, 19 (2007).
- [11] B. Schenke, M. Strickland, C. Greiner, and M. H. Thoma, *Phys. Rev. D* **73**, 125004 (2006).
- [12] A. Rebhan and D. Steineder, *Phys. Rev. D* **81**, 085044 (2010).
- [13] A. Rebhan, P. Romatschke, and M. Strickland, *Phys. Rev. Lett.* **94**, 102303 (2005).
- [14] A. Dumitru, Y. Nara, and M. Strickland, *Phys. Rev. D* **75**, 025016 (2007).
- [15] A. Dumitru, Y. Nara, B. Schenke, and M. Strickland, *Phys. Rev. C* **78**, 024909 (2008).
- [16] A. Rebhan, M. Strickland, and M. Attems, *Phys. Rev. D* **78**, 045023 (2008).
- [17] M. Attems, A. Rebhan, and M. Strickland, *Phys. Rev. D* **87**, 025010 (2013).
- [18] P. Romatschke and R. Venugopalan, *Phys. Rev. D* **74**, 045011 (2006).
- [19] P. Arnold, G. D. Moore, and L. G. Yaffe, *Phys. Rev. D* **72**, 054003 (2005).
- [20] P. Romatschke and M. Strickland, *Phys. Rev. D* **68**, 036004 (2003).
- [21] P. Romatschke and M. Strickland, *Phys. Rev. D* **70**, 116006 (2004).
- [22] P. Romatschke and A. Rebhan, *Phys. Rev. Lett.* **97**, 252301 (2006).
- [23] A. Ipp, A. Rebhan, and M. Strickland, *Phys. Rev. D* **84**, 056003 (2011).
- [24] P. Arnold and G. D. Moore, *Phys. Rev. D* **73**, 025013 (2006); **73**, 025006 (2006).
- [25] D. F. Litim and C. Manuel, *Phys. Rep.* **364**, 451 (2002).
- [26] S. Ichimaru, *Basic Principles*, Statistical Plasma Physics Vol. I (Addison-Wesley, Reading, MA, 1992).
- [27] N. A. Krall and A. W. Trivelpiece, *Principles of Plasma Physics* (McGraw-Hill, New York, 1973).
- [28] C. Manuel and S. Mrówczyński, *Phys. Rev. D* **74**, 105003 (2006).
- [29] J. Peralta-Ramos and E. Calzetta, *Phys. Rev. D* **86**, 125024 (2012).
- [30] M. Mannarelli and C. Manuel, *Phys. Rev. D* **76**, 094007 (2007).
- [31] M. Mannarelli and C. Manuel, *Phys. Rev. D* **77**, 054018 (2008).
- [32] B. Basu, *Phys. Plasmas* **9**, 5131 (2002).
- [33] L. M. Martyushev and V. D. Seleznev, *Phys. Rep.* **426**, 1 (2006).
- [34] D. Kondepudi and I. Prigogine, *Modern Thermodynamics: From Heat Engines to Dissipative Structures* (Wiley, New York, 1998).
- [35] G. Gallavotti and E. G. D. Cohen, *J. Stat. Phys.* **80**, 931 (1995).
- [36] D. Ruelle, *J. Stat. Phys.* **95**, 393 (1999).
- [37] M. Trovato and P. Falsaperla, *Phys. Rev. B* **57**, 4456 (1998).
- [38] A. M. Anile and M. Trovato, *Phys. Lett. A* **230**, 387 (1997).

- [39] A. M. Anile and O. Muscato, *Phys. Rev. B* **51**, 16728 (1995).
- [40] T. Christen, *Europhys. Lett.* **89**, 57007 (2010).
- [41] T. Christen and F. Kassubek, *J. Quant. Spectrosc. Radiat. Transfer* **110**, 452 (2009).
- [42] V. Vedral, *Rev. Mod. Phys.* **74**, 197 (2002).
- [43] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer, Amsterdam, 2002).
- [44] T. Sagawa, *Second Law-Like Inequalities with Quantum Relative Entropy: An Introduction*, edited by M. Nakahara and S. Tanaka, Lectures on Quantum Computing, Thermodynamics and Statistical Physics (World Scientific, Singapore, 2013).
- [45] J.L. Anderson and H.R. Witting, *Physica (Amsterdam)* **74**, 466 (1974); **74**, 489 (1974).
- [46] M. Takamoto and S.-I. Inutsuka, *Physica (Amsterdam)* **389**, 4580 (2010).
- [47] E. Calzetta and J. Peralta-Ramos, *Phys. Rev. D* **82**, 106003 (2010).
- [48] J. Peralta-Ramos and E. Calzetta, *Phys. Rev. D* **87**, 034003 (2013).
- [49] J. Peralta-Ramos and E. Calzetta, *Phys. Rev. C* **82**, 054905 (2010).
- [50] R. Baier, P. Romatschke, D.T. Son, A.O. Starinets, and M. A. Stephanov, *J. High Energy Phys.* **04** (2008) 100.
- [51] S. Bhattacharyya, V.E. Hubeny, S. Minwalla, and M. Rangamani, *J. High Energy Phys.* **02** (2008) 045.
- [52] M. Natsuume and T. Okamura, *Phys. Rev. D* **77**, 066014 (2008); **78**, 089902(R) (2008).
- [53] P. Huovinen and D. Molnar, *Phys. Rev. C* **79**, 014906 (2009).
- [54] G.S. Denicol, T. Koide, and D.H. Rischke, *Phys. Rev. Lett.* **105**, 162501 (2010).
- [55] G.S. Denicol, E. Molnár, H. Niemi, and D.H. Rischke, *Eur. Phys. J. A* **48**, 170 (2012).
- [56] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products* (Elsevier, New York, 2007).