

Three-loop matching of the vector currentPeter Marquard,^{1,2} Jan H. Piclum,^{3,4} Dirk Seidel,⁵ and Matthias Steinhauser¹¹*Institut für Theoretische Teilchenphysik, Karlsruhe Institute of Technology (KIT),
D-76128 Karlsruhe, Germany*²*Deutsches Elektronen Synchrotron DESY, Platanenallee 6, D-15738 Zeuthen, Germany*³*Institut für Theoretische Teilchenphysik und Kosmologie, RWTH Aachen, D-52056 Aachen, Germany*⁴*Physik-Department T31, Technische Universität München, D-85748 Garching, Germany*⁵*Theoretische Physik 1, Universität Siegen, D-57068 Siegen, Germany*

(Received 17 January 2014; published 25 February 2014)

We evaluate the three-loop corrections to the matching coefficient of the vector current between quantum chromodynamics (QCD) and nonrelativistic QCD. The result is presented in the $\overline{\text{MS}}$ scheme where large perturbative corrections are observed. The implications on the threshold production of top-quark pairs are briefly discussed.

DOI: [10.1103/PhysRevD.89.034027](https://doi.org/10.1103/PhysRevD.89.034027)

PACS numbers: 12.38.Bx, 14.65.Fy, 14.65.Ha

I. INTRODUCTION

In recent years effective field theories constructed from quantum chromodynamics (QCD) have been enormously successful at describing phenomena where masses and momenta follow certain limits. Among them is nonrelativistic QCD (NRQCD) [1,2] which is applicable to a system of two heavy quarks moving with small relative velocity. Next to properties of the ψ and Υ families also the threshold production of top-quark pairs is among the prominent examples (see, e.g., Ref. [3] for a review).

The common method to construct an effective theory is based on the so-called matching procedure: appropriately chosen Green's functions are computed in the full and effective theory and equality is required up to power-suppressed terms. In this way the couplings of the effective operators (i.e. the matching coefficients) are determined which completely specifies the effective theory.

A crucial operator both in QCD and NRQCD is the vector current of a heavy quark-antiquark pair. The corresponding matching coefficient enters as a building block in a variety of physical observables, for example the bottom-quark mass from Υ sum rules (see, e.g., Refs. [4,5] for recent analyses) and top-quark threshold production at a future electron positron linear collider [6]. The latter process allows for an extraction of the top-quark mass with an accuracy below 100 MeV [7–9]—an improvement of about an order of magnitude as compared to the current results from the Fermilab Tevatron or the CERN Large Hadron Collider [10].

Several quantities are needed in order to perform a third-order analysis of a heavy quark-antiquark system at threshold. Ultrasoft effects have been considered in Refs. [11,12], the three-loop static potential has been computed in Refs. [13–15] and in Refs. [16,17] a preliminary analysis of the top-quark threshold production cross section has been presented including also third-order potential effects. Details on the potential contributions can be found in Refs. [18,19]. In this paper we compute

the three-loop matching coefficient between the vector current in QCD and NRQCD. Thus all ingredients are available to obtain the complete next-to-next-to-next-to-leading order (NNNLO) QCD prediction of the cross section $e^+e^- \rightarrow t\bar{t}$ close to threshold or the decay width of the $\Upsilon(1S)$ meson to leptons. The results for the latter are presented in an accompanying paper [20] where all building blocks are combined to a phenomenological analysis.

II. VECTOR CURRENTS IN QCD AND NRQCD

The vector current in the full theory (QCD) is given by

$$j_v^\mu = \bar{Q}\gamma^\mu Q, \quad (1)$$

where Q denotes a generic heavy quark with mass m_Q . On the other hand in the effective theory (NRQCD) the current is represented by an expansion in $1/m_Q$ where at each order effective operators have to be considered which are multiplied by coefficient functions. The leading contribution involves one operator given by

$$\tilde{j}^k = \phi^\dagger \sigma^k \chi, \quad (2)$$

where ϕ and χ are two-component Pauli spinors for the quark and antiquark, respectively, and σ^k ($k = 1, 2, 3$) are the Pauli matrices. Hence, the matching coefficient of the vector current is defined through

$$j_v^k = c_v(\mu)\tilde{j}^k + \mathcal{O}\left(\frac{1}{m_Q^2}\right). \quad (3)$$

Note that the 0-component of j_v^μ is only relevant for the power-suppressed contributions.

The purpose of this paper is the evaluation of the purely gluonic three-loop corrections to c_v . The fermionic contributions have already been considered in Refs. [21,22].

In order to compute c_v it is convenient to consider on-shell vertex corrections involving the currents j_v^k and \tilde{j}^k . After taking into account the wave function renormalization one obtains (see also Ref. [18])

$$Z_2 \Gamma_v = c_v \tilde{Z}_2 \tilde{Z}_v^{-1} \tilde{\Gamma}_v + \dots, \quad (4)$$

where the quantities with a tilde are defined in the effective theory and the ellipsis represents terms suppressed by the heavy-quark mass. \tilde{Z}_v^{-1} is the renormalization constant of the current \tilde{j}^k which is used to subtract the remaining poles after renormalization. These poles are due to the separation of long and short distance contributions in the effective theory. In order to evaluate physical quantities it is important that the same subtraction scheme is also adopted in the contributions originating from the effective theory [11,12]. It is well known that in the full theory the renormalization constant of the vector current is equal to 1.

In Eq. (4) Z_2 is the on-shell wave function renormalization constant which has been computed up to three-loop accuracy in Refs. [23–25]. Γ_v denotes the one-particle irreducible vertex diagrams with on-shell quarks carrying momenta q_1 and q_2 . It incorporates all one-particle irreducible vertex graphs and the corresponding counterterms for m_Q and α_s . Sample Feynman diagrams are shown in Fig. 1.

The counterparts of Γ_v and Z_2 in the effective theory can be found on the right-hand side of Eq. (4). It is convenient to apply the threshold expansion [26,27] to Eq. (4). This requires the identification of the hard, soft, potential and ultrasoft momentum regions in the integrals contributing to Γ_v and $\tilde{\Gamma}_v$. Since NRQCD is obtained from QCD by integrating out the hard modes one has by construction that the soft, potential and ultrasoft modes agree in Γ_v and $\tilde{\Gamma}_v$ and thus drop out from Eq. (4). As a consequence Γ_v is evaluated for $q^2 = (q_1 + q_2)^2 = 4m_Q^2$, which corresponds to the leading term of the hard integration region, and $\tilde{\Gamma}_v = 1$. Furthermore, we have $\tilde{Z}_2 = 1$.

There are several technical difficulties which one has to overcome in order to compute the vertex corrections. Among them are the large number of diagrams which leads to several thousand Feynman integrals to be evaluated in the first place, their reduction to a small set of about 100 basis




FIG. 1. Feynman diagrams contributing to Γ_v . Straight and curly lines denote heavy quarks with mass m_Q and gluons, respectively.

integrals, so-called master integrals, and the evaluation of the latter in an expansion in $\epsilon = (4 - D)/2$, where D is the space-time dimension. The last two tasks become more complicated by the additional condition $q^2 = 4m_Q^2$ on the external momentum. An automated setup for the calculation has been described in Ref. [22] and applied to the fermionic contributions. Its core parts are a powerful implementation of Laporta's algorithm in the program CRUSHER [28], and FIESTA [29–31] which is based on sector decomposition and is used for the numerical integration of the master integrals. The main differences compared to the gluonic part considered in this paper are the larger number of diagrams and the increased complexity of the integrals which have to be reduced to master integrals. Furthermore, the master integrals are more numerous and more involved.

III. MATCHING COEFFICIENT TO ORDER α_s^3

Before discussing the matching coefficient it is instructive to consider the renormalization constant \tilde{Z}_v . The analytical results can be extracted from Refs. [11,21,32,33]:

$$\begin{aligned} \tilde{Z}_v = & 1 + \left(\frac{\alpha_s^{(n_l)}(\mu)}{\pi} \right)^2 \frac{C_F \pi^2}{\epsilon} \left(\frac{1}{12} C_F + \frac{1}{8} C_A \right) + \left(\frac{\alpha_s^{(n_l)}(\mu)}{\pi} \right)^3 C_F \pi^2 \left\{ C_F^2 \left[\frac{5}{144 \epsilon^2} + \left(\frac{43}{144} - \frac{1}{2} \ln 2 + \frac{5}{48} L_\mu \right) \frac{1}{\epsilon} \right] \right. \\ & + C_F C_A \left[\frac{1}{864 \epsilon^2} + \left(\frac{113}{324} + \frac{1}{4} \ln 2 + \frac{5}{32} L_\mu \right) \frac{1}{\epsilon} \right] + C_A^2 \left[-\frac{1}{16 \epsilon^2} + \left(\frac{2}{27} + \frac{1}{4} \ln 2 + \frac{1}{24} L_\mu \right) \frac{1}{\epsilon} \right] \\ & \left. + T n_l \left[C_F \left(\frac{1}{54 \epsilon^2} - \frac{25}{324 \epsilon} \right) + C_A \left(\frac{1}{36 \epsilon^2} - \frac{37}{432 \epsilon} \right) \right] + C_F T n_h \frac{1}{60 \epsilon} \right\} + \mathcal{O}(\alpha_s^4), \end{aligned} \quad (5)$$

where $C_A = N_c$, $C_F = (N_c^2 - 1)/(2N_c)$ and $T = 1/2$ for a $SU(N_c)$ gauge group and $L_\mu = \ln(\mu^2/m_Q^2)$. Note that the strong coupling is defined in the effective theory with n_l active quarks where $n_l + n_h$ is the total number of quark flavors. In our case we have $n_h = 1$; however, we keep n_h in the formulas for convenience.

From our calculation we can extract the renormalization constant \tilde{Z}_v and compare with Eq. (5). The central values of our numerical coefficients agree at the percent level with the analytical result of Eq. (5) which constitutes a first nontrivial check and provides quite some confidence in the overall setup of our calculation.

We write the perturbative expansion of the matching coefficient in the form

$$c_v = 1 + \frac{\alpha_s^{(n_l)}(\mu)}{\pi} c_v^{(1)} + \left(\frac{\alpha_s^{(n_l)}(\mu)}{\pi} \right)^2 c_v^{(2)} + \left(\frac{\alpha_s^{(n_l)}(\mu)}{\pi} \right)^3 c_v^{(3)} + \mathcal{O}(\alpha_s^4), \quad (6)$$

and decompose $c_v^{(3)}$ according to the color structures as

$$c_v^{(3)} = C_F [C_F^2 c_{FFF} + C_F C_A c_{FFA} + C_A^2 c_{FAA} + T n_l (C_F c_{FFL} + C_A c_{FAL} + T n_h c_{FHL} + T n_l c_{FLL}) + T n_h (C_F c_{FFH} + C_A c_{FAH} + T n_h c_{FHH})] + \text{singlet terms}. \quad (7)$$

Note that all color factors of the nonsinglet part can be expressed in terms of C_F , C_A , and T . In this paper we do not consider the singlet contribution where the external current

does not couple to the fermion line in the final state. At two-loop order such contributions have been computed [34] for axial-vector, scalar and pseudoscalar currents in Ref. [35]. Their numerical effect in those cases is below 3% as compared to the nonsinglet contributions and thus quite small.

The one- and two-loop corrections are known since more than ten years and have been computed in Refs. [36] and [32,35,37], respectively. On the other hand, the fermionic three-loop term became available only a few years ago [21,22]. The so-called renormalon contribution, consisting of the one-loop diagram with arbitrarily many massless quark loop insertions in the gluon propagator, has been computed in Ref. [38]. Supersymmetric one-loop corrections to c_v have been computed in Ref. [39].

In the following we present the results for the individual coefficients in Eq. (7) parametrized in terms of $\alpha_s^{(n_l)}(m_Q)$. The reconstruction of the full dependence on the renormalization scale is straightforward; the corresponding expressions can be obtained from [40]. Our results read [41]

$$\begin{aligned} c_v^{(1)} &= -2C_F, \\ c_v^{(2)} &= \left(-\frac{151}{72} + \frac{89}{144}\pi^2 - \frac{5}{6}\pi^2 \ln 2 - \frac{13}{4}\zeta(3) \right) C_A C_F + \left(\frac{23}{8} - \frac{79}{36}\pi^2 + \pi^2 \ln 2 - \frac{1}{2}\zeta(3) \right) C_F^2 \\ &\quad + \left(\frac{22}{9} - \frac{2}{9}\pi^2 \right) C_F T n_h + \frac{11}{18} C_F T n_l - \frac{1}{2}\pi^2 \left(\frac{1}{2} C_A + \frac{1}{3} C_F \right) C_F L_\mu, \\ c_{FFF} &= 36.55(0.53) + \left(-\frac{9}{16} + \frac{3}{2} \ln 2 \right) \pi^2 L_\mu - \frac{5}{32} \pi^2 L_\mu^2, \\ c_{FFA} &= -188.10(0.83) + \left(-\frac{59}{108} - \frac{3}{4} \ln 2 \right) \pi^2 L_\mu - \frac{47}{576} \pi^2 L_\mu^2, \\ c_{FAA} &= -97.81(0.38) + \left(-\frac{2}{9} - \frac{3}{4} \ln 2 \right) \pi^2 L_\mu + \frac{1}{6} \pi^2 L_\mu^2, \\ c_{FFL} &= 46.691(0.006) + \frac{25}{108} \pi^2 L_\mu - \frac{1}{18} \pi^2 L_\mu^2, \quad c_{FAL} = 39.624(0.005) + \frac{37}{144} \pi^2 L_\mu - \frac{1}{12} \pi^2 L_\mu^2, \\ c_{FHL} &= -\frac{557}{162} + \frac{26}{81} \pi^2, \quad c_{FLL} = -\frac{163}{162} - \frac{4}{27} \pi^2, \quad c_{FFH} = -0.846(0.006) - \frac{1}{20} \pi^2 L_\mu, \\ c_{FAH} &= -0.098(0.051), \quad c_{FHH} = -\frac{427}{162} + \frac{158}{2835} \pi^2 + \frac{16}{9} \zeta(3). \end{aligned} \quad (8)$$

All uncertainties originating from the individual master integrals are added quadratically. In order to obtain a conservative error estimate we interpret the uncertainty of the numerical integration as 1 standard deviation from a Gaussian distribution and multiply it by a factor of 5 which is accounted for in Eq. (8) [42]. The coefficients of L_μ could be obtained in analytic form since all renormalization constants are known analytically.

In most applications it is sufficient to know the result for the matching coefficient with numerically evaluated color

factors. Setting $C_F = 4/3$, $C_A = 3$, $T = 1/2$ and $n_h = 1$ before inserting the master integrals and combining the numerical uncertainties we get

$$\begin{aligned} c_v &\approx 1 - 2.667 \frac{\alpha_s^{(n_l)}}{\pi} + \left(\frac{\alpha_s^{(n_l)}}{\pi} \right)^2 [-44.551 + 0.407 n_l] \\ &\quad + \left(\frac{\alpha_s^{(n_l)}}{\pi} \right)^3 [-2091(2) + 120.66(0.01) n_l - 0.823 n_l^2] \\ &\quad + \text{singlet terms}, \end{aligned} \quad (9)$$

where $\mu = m_Q$ has been chosen. Note that the n_l -independent three-loop term contains the contribution with a closed massive quark loop which amounts to $c_v^{(3)}|_{n_h=0} \approx -0.93(8)$ [22]. One observes that for $n_l = 3, 4$ and 5 all coefficients in Eq. (9) have the same sign and that they grow quite rapidly when going from NLO to NNNLO. At NNLO and NNNLO the fermionic corrections screen the nonfermionic ones, but even for $n_l = 5$ only a reduction of at most 30% is obtained. A first glance at Eq. (9) would suggest that for the quantity c_v perturbation theory breaks down even though the momentum scale involved in the problem, m_Q , is quite large. However, as already mentioned above, c_v itself does not represent a physical quantity. It has to be combined with contributions originating from soft, potential and ultrasoft momentum regions which can compensate the large coefficients in Eq. (9). Further discussions on this topic can be found in Ref. [20]. It might very well be that the $\overline{\text{MS}}$ scheme adopted in our calculation is not well suited for separating the divergences occurring in the different regimes. In fact, also the ultrasoft contribution studied in Refs. [11,12] shows large numerical effects.

We have performed several checks on the correctness of our result which we want to mention in the following. In our calculation we allowed for a general gauge parameter ξ which manifests as a polynomial dependence of the individual diagrams. After summing the three-loop results for Z_2 and Γ_v (taking into account the corresponding quark mass counterterm contribution) we concentrated on the coefficient of the linear ξ dependence and have verified that it vanishes. As a further check we recomputed the n_l contribution [21] using our automated setup. In this context we want to mention that in Ref. [21] all occurring master integrals have been computed either analytically or using a numerical method different from the one used in the present paper. As already mentioned above, with our calculation we could also reproduce the renormalization constant in Eq. (5) with high accuracy which checks all but the highest ϵ coefficients of the master integrals. We note in passing that we have a similar accuracy for the cancellation of the spurious poles up to seventh order occurring due to our reduction procedure.

At this point it is instructive to show a result for a typical master integral contributing to c_v . For the Feynman diagram in Fig. 2, which we need up to order ϵ , we obtain with the help of FIESTA [29–31]

$$M = \frac{e^{3\epsilon\gamma_E}}{m_Q^4} \left(\frac{\mu^2}{m_Q^2} \right)^{3\epsilon} \left(+ \frac{0.411236(3)}{\epsilon^2} + \frac{3.4860(1)}{\epsilon} + 34.520(2) + 339.68(4)\epsilon + \mathcal{O}(\epsilon^2) \right). \quad (10)$$

A very powerful check on the correctness of our result is provided by the change of basis for the master integrals. We employ the integral tables generated during the reduction procedure in order to reexpress the master integrals, which are not known analytically, through different, in general




FIG. 2. Typical master integral appearing in our calculation. The solid lines and dashed lines represent massive and massless lines, respectively. For the external momenta we have the conditions $q_1^2 = q_2^2 = m_Q^2$ and $(q_1 + q_2)^2 = 4m_Q^2$.

more complicated ones. This transformation is done analytically for general space-time dimension D . In a next step the new master integrals are again evaluated with FIESTA and inserted in the new expression for c_v . In Table I we compare the results for the purely gluonic coefficients and the complete result for $c_v^{(3)}$ obtained in the two bases. We observe an excellent agreement within the uncertainties. In the case of the “alternative basis” one has to keep in mind that the integrals to be evaluated numerically are significantly more complicated which explains the larger uncertainties for the coefficients in Table I.

As a further check on the numerical evaluation of the master integrals we have used a different momentum assignment in the input for FIESTA. As a consequence different expressions are generated in intermediate steps leading to different numerical integrations. The final results are in complete agreement with Eq. (9).

We are now in the position to have a first look at the phenomenological consequences of our result for c_v . We consider the residue of the two-point function of the vector currents

$$(-q^2 g_{\mu\nu} + q_\mu q_\nu) \Pi(q^2) = i \int dx e^{iqx} \langle 0 | T j_\mu(x) j_\nu^\dagger(0) | 0 \rangle, \quad (11)$$

TABLE I. Comparison of the purely gluonic coefficients of Eq. (8) and $c_v^{(3)}$ with $n_l = 4$ and $n_l = 5$ for two different choices of the master integral basis. For convenience $\mu = m_Q$ has been adopted. The given uncertainties are obtained by combining the numerical uncertainties of each master integral contribution in quadrature. In contrast to Eqs. (8) and (9) no factor 5 has been introduced for this comparison.

	Default basis [cf. Eq. (8)]	Alternative basis
c_{FFF}	36.55(0.11)	36.61(2.93)
c_{FFA}	-188.10(0.17)	-188.04(2.91)
c_{FAA}	-97.81(0.08)	-97.76(2.05)
$c_v^{(3)}(n_l = 4)$	-1621.7(0.4)	-1621(23)
$c_v^{(3)}(n_l = 5)$	-1508.4(0.4)	-1507(23)




FIG. 3. Residue for the top-quark system normalized to $Z_t^{LO}(\mu_s)$ as a function of the renormalization scale μ . Dotted, dash-dotted, short-dashed and solid lines correspond to LO, NLO, NNLO and NNNLO prediction. In the long-dashed curve only the fermion contributions to $c_v^{(3)}$ are taken into account [NNNLO (ferm.)].

which is obtained by considering $\Pi(q^2)$ close to the $Q\bar{Q}$ threshold. In this limit $\Pi(q^2)$ is dominated by pole contributions originating from bound-state effects

$$\Pi(q^2) \stackrel{E \rightarrow E_n}{\approx} \frac{N_c}{2m_Q^2} \frac{Z_n}{E_n - (E + i0)} + \dots, \quad (12)$$

where the ellipsis denotes contributions from the continuum. Z_n and E_n are the residue and energy of the n th resonance which determine the height and position of the threshold cross section, respectively. In the following we consider the residue of the 1S (pseudo)bound state of top quarks,

$$Z_t = \left[c_v^2 - \frac{E_1}{m_t} c_v \left(c_v + \frac{d_v}{6} \right) \right] |\psi_1(0)|^2, \quad (13)$$

where $\psi_1(0)$ denotes the wave function of the 1S state at the origin and d_v is the matching coefficient of the $1/m_Q^2$ suppressed term in Eq. (3). Note that terms which contribute only to fourth and higher orders have been neglected in Eq. (13). We extend the considerations of Ref. [16,43] by including the nonfermionic contribution of $c_v^{(3)}$ and the $\mathcal{O}(\epsilon)$ term of the $1/m_Q$ potential [20]. Choosing the potential subtracted scheme [44] with $\mu_f = 20$ GeV to define the top-quark mass we obtain $m_t^{\text{PS}} = 171.4$ GeV which leads to

$$\begin{aligned} Z_t &= \frac{(C_F m_t^{\text{PS}} \alpha_s)^3}{8\pi} [1 + (-2.131 + 3.661L)\alpha_s + (8.38 + 1.27x_f - 7.26 \ln \alpha_s - 13.40L + 8.93L^2)\alpha_s^2 \\ &\quad + (5.46 + (-2.23 + 0.78L_f)x_f + 2.21a_3 + 21.48b_{2\epsilon} + 37.53c_f - 134.8(0.1)c_g \\ &\quad + (-9.79 - 44.27L) \ln \alpha_s - 16.35 \ln^2 \alpha_s + (53.17 + 4.66x_f)L - 48.18L^2 + 18.17L^3)\alpha_s^3 + \mathcal{O}(\alpha_s^4)] \\ &= \frac{(C_F m_t^{\text{PS}} \alpha_s)^3}{8\pi} [1 - 2.13\alpha_s + 23.66\alpha_s^2 - 113.0(0.1)\alpha_s^3 + \mathcal{O}(\alpha_s^4)], \end{aligned} \quad (14)$$

where $x_f = \mu_f/(m_t^{\text{PS}} \alpha_s)$, $L = \ln(\mu/(m_t^{\text{PS}} C_F \alpha_s))$, and $L_f = \ln(\mu^2/\mu_f^2)$. We have used $\alpha_s(M_Z) = 0.1184$ to compute $\alpha_s = \alpha_s(\mu_s) \approx 0.141$ where the soft scale $\mu_s = m_Q C_F \alpha_s(\mu_s) \approx 32.16$ GeV has been adopted after the second equality sign. In order to get an impression about the importance of the individual contributions we mark the μ -independent coefficients from the three-loop static potential (a_3), from the two-loop $\mathcal{O}(\epsilon)$ term of the $1/(m_Q r^2)$ potential ($b_{2\epsilon}$), and from the three-loop fermion (c_f) and purely gluonic (c_g) contribution to $c_v^{(3)}$ separately. For this choice of μ one observes quite big NNNLO contributions which are dominated by c_g . Thus, it is instructive to investigate the μ dependence of Z_t which is shown in Fig. 3. Around the soft scale no convergence is observed. Allowing, however, for higher scales one finds a quite flat behavior of the NNNLO curve. Furthermore, the NNNLO corrections become quite small. E.g., considering the top-quark system for $\mu \approx 80$ GeV, the NLO terms amount to about +15% and the NNLO to roughly +20%. The third-order contribution is practically zero. Similar

observations also hold for the bottom-quark case; see Ref. [20].

IV. CONCLUSIONS

The third-order contribution to the matching coefficient of the vector current between QCD and NRQCD has been computed. An automated setup has been developed where even the occurring master integrals are identified automatically, processed with the help of the computer program FIESTA, and prepared for the insertion into the analytic reduction of $c_v^{(3)}$.

In the $\overline{\text{MS}}$ scheme the numerical impact of $c_v^{(3)}$ is quite big as can be seen from Eq. (9) which constitutes the main result of this paper. In a dedicated analysis one has to investigate the consequences for the bottom-quark mass extracted from Υ sum rules and the top-quark threshold production cross section at a future linear collider.

An analysis of the residue of the 1S state indicates that at energy scales around two to three times the soft scale good convergence of the perturbative series is observed.

ACKNOWLEDGMENTS

We would like to thank Martin Beneke and Alexander Penin for many useful discussions and communications and Alexander Smirnov for continuous support on FIESTA.

This work is supported by DFG through Grant No. SFB/TR 9 “Computational Particle Physics.” The Feynman diagrams were drawn with the help of Axodraw [45] and JaxoDraw [46].

-
- [1] W. E. Caswell and G. P. Lepage, *Phys. Lett.* **167B**, 437 (1986).
 [2] G. T. Bodwin, E. Braaten, and G. P. Lepage, *Phys. Rev. D* **51**, 1125 (1995); **55**, 5853(E) (1997).
 [3] N. Brambilla *et al.*, [arXiv:hep-ph/0412158](https://arxiv.org/abs/hep-ph/0412158).
 [4] A. Pineda and A. Signer, *Phys. Rev. D* **73**, 111501 (2006).
 [5] A. Hoang, P. Ruiz-Femenia, and M. Stahlhofen, *J. High Energy Phys.* **10** (2012) 188.
 [6] A. H. Hoang *et al.*, *Eur. Phys. J. direct C* **2**, 1 (2000).
 [7] M. Martinez and R. Miquel, *Eur. Phys. J. C* **27**, 49 (2003).
 [8] K. Seidel, F. Simon, M. Tesar, and S. Poss, *Eur. Phys. J. C* **73**, 2530 (2013).
 [9] T. Horiguchi, A. Ishikawa, T. Suehara, K. Fujii, Y. Sumino, Y. Kiyo, and H. Yamamoto, [arXiv:1310.0563](https://arxiv.org/abs/1310.0563).
 [10] J. Beringer *et al.* (Particle Data Group Collaboration), *Phys. Rev. D* **86**, 010001 (2012).
 [11] M. Beneke, Y. Kiyo, and A. A. Penin, *Phys. Lett. B* **653**, 53 (2007).
 [12] M. Beneke and Y. Kiyo, *Phys. Lett. B* **668**, 143 (2008).
 [13] A. V. Smirnov, V. A. Smirnov, and M. Steinhauser, *Phys. Lett. B* **668**, 293 (2008).
 [14] A. V. Smirnov, V. A. Smirnov, and M. Steinhauser, *Phys. Rev. Lett.* **104**, 112002 (2010).
 [15] C. Anzai, Y. Kiyo, and Y. Sumino, *Phys. Rev. Lett.* **104**, 112003 (2010).
 [16] M. Beneke, Y. Kiyo, A. Penin, and K. Schuller, in *Proceedings of the International Linear Collider Workshop (LCWS07 and ILC07)*, edited by A. Frey and S. Riemann (DESY, Hamburg, 2008).
 [17] M. Beneke, Y. Kiyo, and K. Schuller, *Proc. Sci.*, RADCOR (2007) 051.
 [18] M. Beneke, Y. Kiyo, and K. Schuller, [arXiv:1312.4791](https://arxiv.org/abs/1312.4791).
 [19] M. Beneke, Y. Kiyo, and K. Schuller (to be published).
 [20] M. Beneke, Y. Kiyo, P. Marquard, A. Penin, J. Piclum, D. Seidel, and M. Steinhauser, [arXiv:1401.3005](https://arxiv.org/abs/1401.3005).
 [21] P. Marquard, J. H. Piclum, D. Seidel, and M. Steinhauser, *Nucl. Phys.* **B758**, 144 (2006).
 [22] P. Marquard, J. H. Piclum, D. Seidel, and M. Steinhauser, *Phys. Lett. B* **678**, 269 (2009).
 [23] D. J. Broadhurst, N. Gray, and K. Schilcher, *Z. Phys. C* **52**, 111 (1991).
 [24] K. Melnikov and T. van Ritbergen, *Nucl. Phys.* **B591**, 515 (2000).
 [25] P. Marquard, L. Mihaila, J. H. Piclum, and M. Steinhauser, *Nucl. Phys.* **B773**, 1 (2007).
 [26] M. Beneke and V. A. Smirnov, *Nucl. Phys.* **B522**, 321 (1998).
 [27] V. A. Smirnov, *Springer Tracts Mod. Phys.* **250**, 1 (2012).
 [28] P. Marquard and D. Seidel (unpublished).
 [29] A. V. Smirnov and M. N. Tentyukov, *Comput. Phys. Commun.* **180**, 735 (2009).
 [30] A. V. Smirnov, V. A. Smirnov, and M. Tentyukov, *Comput. Phys. Commun.* **182**, 790 (2011).
 [31] A. V. Smirnov, [arXiv:1312.3186](https://arxiv.org/abs/1312.3186).
 [32] M. Beneke, A. Signer, and V. A. Smirnov, *Phys. Rev. Lett.* **80**, 2535 (1998).
 [33] B. A. Kniehl, A. A. Penin, M. Steinhauser, and V. A. Smirnov, *Phys. Rev. Lett.* **90**, 212001 (2003); **91**, 139903(E) (2003).
 [34] Due to Furry’s theorem the two-loop singlet contribution is zero for the vector current.
 [35] B. A. Kniehl, A. Onishchenko, J. H. Piclum, and M. Steinhauser, *Phys. Lett. B* **638**, 209 (2006).
 [36] G. Källén and A. Sabry, *K. Dan. Vidensk. Selsk., Mat.-Fys. Medd.* **29**, 1 (1955).
 [37] A. Czarnecki and K. Melnikov, *Phys. Rev. Lett.* **80**, 2531 (1998).
 [38] E. Braaten and Y.-Q. Chen, *Phys. Rev. D* **57**, 4236 (1998); **59**, 079901(E) (1999).
 [39] Y. Kiyo, M. Steinhauser, and N. Zerf, *Phys. Rev. D* **80**, 075005 (2009).
 [40] <https://www.ttp.kit.edu/Progdata/ttp14/ttp14-002/>.
 [41] Note that in Ref. [21] the result has been expressed in terms of the coupling defined in the full theory whereas here we use the effective one denoted by $\alpha_s^{(n_f)}$. This explains the difference in the logarithmic part of the coefficient c_{FHL} .
 [42] Note that in Ref. [22], where the fermionic contributions are given, only a factor 2 has been chosen which explains the slight increase of the uncertainty of c_{FHL} in Eq. (8).
 [43] Note that there is a misprint in Eq. (5) of Ref. [16]: the term $E(1 - d_v/3)/m$ should read $E(c_v - d_v/3)/m$.
 [44] M. Beneke, *Phys. Lett. B* **434**, 115 (1998).
 [45] J. A. M. Vermaseren, *Comput. Phys. Commun.* **83**, 45 (1994).
 [46] D. Binosi and L. Theussl, *Comput. Phys. Commun.* **161**, 76 (2004).