

Inflationary steps in the Planck dataVinícius Miranda^{1,2} and Wayne Hu^{3,1}¹*Department of Astronomy & Astrophysics, University of Chicago, Chicago, Illinois 60637, USA*²*The Capes Foundation, Ministry of Education of Brazil, Brasília DF 70359-970, Brazil*³*Kavli Institute for Cosmological Physics, Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

(Received 12 December 2013; published 21 April 2014)

We extend and improve the modeling and analysis of large-amplitude, sharp inflationary steps for second order corrections required by the precision of the Planck cosmic microwave background power spectrum and for arbitrary Dirac-Born-Infeld sound speed. With two parameters, the amplitude and frequency of the resulting oscillations, step models improve the fit by $\Delta\chi^2 = -11.4$. Evidence for oscillations damping before the Planck beam scale is weak: damping only improves the fit to $\Delta\chi^2 = -14.0$ for one extra parameter, if step and cosmological parameters are jointly fit, in contrast to analyses which fix the latter. Likewise, further including the sound speed as a parameter only marginally improves the fit to $\Delta\chi^2 = -15.2$ but has interesting implications for the lowest multipole temperature and polarization anisotropy. Since chance features in the noise can mimic these oscillatory features, we discuss tests from polarization power spectra, lensing reconstruction and squeezed and equilateral bispectra that should soon verify or falsify their primordial origin.

DOI: [10.1103/PhysRevD.89.083529](https://doi.org/10.1103/PhysRevD.89.083529)

PACS numbers: 98.80.Cq, 98.70.Vc

I. INTRODUCTION

Intriguingly, the cosmic microwave background (CMB) seems to favor rapid oscillations in the curvature power spectrum over the smooth power law spectrum given by slow-roll inflation at a level of $\Delta\chi^2 \sim 10$ –20. Such oscillations, first seen in the WMAP data [1–4], persist in the recent Planck data [5–8]. While the significance of this improvement is debatable given the ability of statistical fluctuations from instrument noise or cosmic variance to mimic the signal, its implications for inflationary physics are sufficiently dramatic to merit careful consideration.

Rapidly oscillating power spectra can be generated during inflation if the inflaton rolls over features in much less than an efold, for example oscillations in the potential [9,10], a step in the potential [11] or warp in the Dirac-Born-Infeld (DBI) model [12]. In this paper we consider the less well-explored step feature cases.

On the model side, we extend previous analyses [2,13] by analytically treating large amplitude sharp steps in both the potential and warp at arbitrary sound speeds including new second order corrections that are required by the enhanced precision of the Planck data. Having an analytic model for the inflationary power spectrum greatly enhances the efficiency of the analysis while varying the sound speed provides interesting phenomenology for the lowest multipoles.

On the analysis side, we jointly fit for step and cosmological parameters, unlike the Planck collaboration analysis [5]. Because the presence of step oscillations also changes the broadband average power in the spectrum, joint variation is crucial for interpreting constraints on step

parameters. Although more recent analyses have also jointly varied parameters [6,8], they did so in a different context where the oscillations persist out to arbitrarily high multipoles. We show that joint variation is particularly important for finite width steps and misleading constraints arise when cosmological parameters are fixed.

The outline of the paper is as follows. In Sec. II we describe the improvements and extensions to the modeling of the curvature power spectrum from steps in the potential and warp. These are derived in Appendix A and are shown to be sufficiently accurate for Planck data in Appendix B. The best fit step models at low and high sound speed, found from jointly maximizing the likelihood over step and cosmological parameters in Appendix C, are presented in Sec. III. In Sec. IV we provide falsifiable predictions of these models. We discuss these results in Sec. V.

II. STEP POWER SPECTRA

In this section, we summarize the description of the curvature power spectrum for sharp potential and warp steps in DBI inflation derived in Appendix A. This analytic treatment generalizes previous ones [2,13] to large amplitude, arbitrary sound speed models and employs second order corrections to ensure sufficient accuracy for comparison to the Planck data in the following sections.

We consider models with step features in the DBI Lagrangian,

$$\mathcal{L} = [1 - \sqrt{1 - 2X/T(\phi)}]T(\phi) - V(\phi), \quad (1)$$

where the kinetic term $2X = -\nabla^\mu\phi\nabla_\mu\phi$. We choose units where $M_{\text{pl}} = (8\pi G)^{-1/2} = c = \hbar = 1$ throughout. In braneworld theories that motivate the DBI Lagrangian, ϕ determines the position of the brane, $T(\phi)$ gives the warped brane tension, and $V(\phi)$ is the interaction potential. Note that for $X/T \ll 1$, the sound speed

$$c_s(\phi, X) = \sqrt{1 - 2X/T(\phi)} \quad (2)$$

goes to 1 and this Lagrangian becomes that of a scalar with a canonical kinetic term.

We allow steps to appear in either the warp or the potential,

$$\begin{aligned} T(\phi) &= \frac{\phi^4}{\lambda_B} [1 + b_T F(\phi)], \\ V(\phi) &= V_0 \left(1 - \frac{1}{6}\beta\phi^2\right) [1 + b_V F(\phi)]. \end{aligned} \quad (3)$$

Here, λ_B , V_0 , β parametrize the smooth model and are determined by the tilt and amplitude of the power spectrum as well as the end point for DBI inflation, whereas b_T , b_V give the height of a tanh step,

$$F(\phi) = \tanh\left(\frac{\phi - \phi_s}{d}\right) - 1, \quad (4)$$

at field location ϕ_s , with field width d . Unlike previous treatments [2,13] we allow for the possibility of potential steps at arbitrary sound speed but for simplicity do not consider simultaneous steps in both the warp and the potential.

We show in Appendix A that steps in the warp or potential, over which the inflaton rolls in much less than an efold, generate oscillations in the power spectrum of the following form:

$$\ln \Delta_{\mathcal{R}}^2 = \ln A_s \left(\frac{k}{k_0}\right)^{n_s-1} + I_0(k) + \ln[1 + I_1^2(k)], \quad (5)$$

where we take the normalization scale $k_0 = 0.08 \text{ Mpc}^{-1}$, which is closer to the best constrained scale for the Planck data than the conventional choice of 0.05 Mpc^{-1} . The leading order contribution from the step is

$$I_0(k) = [C_1 W(ks_s) + C_2 W'(ks_s) + C_3 Y(ks_s)] \mathcal{D}\left(\frac{ks_s}{x_d}\right), \quad (6)$$

and the second order contribution is

$$\begin{aligned} \sqrt{2}I_1(k) &= \frac{\pi}{2}(1 - n_s) + [C_1 X(ks_s) + C_2 X'(ks_s) \\ &+ C_3 Z(ks_s)] \mathcal{D}\left(\frac{ks_s}{x_d}\right), \end{aligned} \quad (7)$$

where the windows

$$\begin{aligned} W(x) &= \frac{3 \sin(2x)}{2x^3} - \frac{3 \cos(2x)}{x^2} - \frac{3 \sin(2x)}{2x}, \\ X(x) &= \frac{3}{x^3} (\sin x - x \cos x)^2, \\ Y(x) &= \frac{6x \cos(2x) + (4x^2 - 3) \sin(2x)}{x^3}, \\ Z(x) &= -\frac{3 + 2x^2 - (3 - 4x^2) \cos(2x) - 6x \sin(2x)}{x^3}, \end{aligned} \quad (8)$$

and $' = d/d \ln x$. The sound horizon when the inflaton crosses the step s_s controls the frequency of the oscillations, whereas the finite width $x_d \propto d^{-1}$ determines their damping via

$$\mathcal{D}(y) = \frac{y}{\sinh(y)}. \quad (9)$$

We give the correspondence between these phenomenological parameters and the fundamental ones in Appendix B. There, we also test the accuracy of the analytic model in Eq. (5) against exact calculations. We show that the precision of the Planck data set necessitates the inclusion of the second order I_1 correction whose analytic form is entirely new to this work (cf. [13], [5]). Note that the second order term is determined by exactly the same parameters as the leading order term as a consequence of the generalized slow-roll construction [14].

The constants C_i can be related to fractional changes in c_s and the slow-roll parameter $\epsilon_H = -d \ln H / d \ln a$ induced by the step

$$c_j \equiv \frac{c_{sj}}{c_{sa}}, \quad e_j \equiv \frac{\epsilon_{Hj}}{\epsilon_{Ha}}, \quad (10)$$

where ‘‘a’’ denotes their values on the attractor after the step, ‘‘b’’ is for the same before the step, and ‘‘i’’ is for immediately after the step off of the attractor. More specifically,

$$\begin{aligned} C_1 &= -\ln c_b e_b, \\ C_2 &= -\frac{2c_i - c_b}{3c_i + c_b} + \frac{2e_i - e_b}{3e_i + e_b}, \\ C_3 &= 2 \frac{(1 - c_b) + (c_i - 1)/4}{c_i + c_b}. \end{aligned} \quad (11)$$

For warp steps, the attractor solutions before and after the step and energy conservation at the step gives ($b_V = 0$)

$$\begin{aligned}
c_b &= e_b = \sqrt{\frac{1 - 2b_T}{1 - 2b_T c_{sa}^2}}, \\
c_i &= \frac{c_b}{1 - 2b_T(1 - c_{sa}c_b)}, \\
e_i &= \frac{1 - c_{sa}^2 c_i^2}{c_i(1 - c_{sa}^2)}. \tag{12}
\end{aligned}$$

This generalizes the results of Ref. [13] to large amplitude steps as $c_{sa} \rightarrow 1$. Note that in this limit, arbitrarily large fractional steps in the warp $b_T \rightarrow -\infty$ still only cause infinitesimal changes in the slow-roll parameters $c_j = e_j = 1$ or $C_i \rightarrow 0$. Consequently, there are sound speeds near unity for which a step in the warp cannot explain finite amplitude oscillations in the data.

For potential steps ($b_T = 0$),

$$\begin{aligned}
c_b &= e_b = 1, \\
c_i &= 1 - \frac{3b_V(1 - c_{sa}^2)}{3b_V(1 - c_{sa}^2) - \epsilon_{Ha}}, \\
e_i &= 1 - \frac{3b_V[-3b_V(1 - c_{sa}^2) + (1 + c_{sa}^2)\epsilon_{Ha}]}{\epsilon_{Ha}[-3b_V(1 - c_{sa}^2) + \epsilon_{Ha}]}. \tag{13}
\end{aligned}$$

This generalizes the results of Ref. [2] for potential steps to arbitrary sound speeds. Note that for potential steps $C_1 = 0$. We test the accuracy of these approximations in Appendix B.

In summary, a model is parametrized by five numbers, $\{C_1, C_2, C_3, s_s, x_d\}$. To the leading order, the C_1 term represents a step in the power spectrum at $ks_s \sim 1$, the C_2 term represents a constant amplitude oscillation out to the damping scale $ks_s = x_d$, and the C_3 term represents a change to the shape of the first few oscillations.

In a given model, not all of these parameters are independent as they are determined by the background parameters. Specifically, the three C_i parameters are controlled by the amplitude of the step and mainly the sound speed after the step. For a potential step $C_1 = 0$ and $C_3 \rightarrow 0$ for $c_s \rightarrow 1$. For a warp step, all three C_i are comparable but $C_i \rightarrow 0$ for $c_s \rightarrow 1$, even for arbitrarily large steps.

III. PLANCK DATA ANALYSIS

In this section we analyze the Planck data for the presence of sharp inflationary steps which create high frequency oscillations in the power spectrum. We begin in Sec. III A with potential steps in canonical sound speed models. In Sec. III B we extend the analysis to arbitrary sound speed models where both warp and potential steps can produce the oscillatory phenomenology favored by the Planck data. In Appendix C we discuss details of the analysis that enhance the efficiency of the model search.

A. Canonical sound speed models

We begin with the simplest case of $c_s = 1$ models. Here, warp steps have no effect and potential steps give $C_1 = C_3 = 0$. Step models are thus described by three parameters, $\{s_s, C_2, x_d\}$, the oscillation frequency, amplitude, and damping scale, respectively. The underlying smooth cosmology is taken to be the flat Λ CDM model as defined by $\{A_s, n_s, \theta_A, \Omega_c h^2, \Omega_b h^2, \tau\}$, where θ_A is the angular acoustic scale at recombination, $\Omega_c h^2$ parametrizes the cold dark matter density, $\Omega_b h^2$ parametrizes the baryon density, and τ parametrizes the Thomson optical depth to reionization. We calculate CMB power spectra using a modified version of CAMB [15,16]. In addition, the Planck data are modeled by foreground parameters which we hold fixed throughout to the best fit smooth model (see Appendix C and Tables I and II).

We are interested in the question of whether the step parameters significantly improve the fit to the Planck data rather than marginalized constraints on the parameters themselves. Since the Monte Carlo Markov chain technique is highly inefficient for these purposes, we instead directly maximize the likelihood or minimize the effective $\chi^2 = -2 \ln \mathcal{L}$ in the step and cosmological parameter space jointly. For these oscillatory spectra, the likelihood is a rapidly varying function of frequency s_s , with many local minima. Fortunately, previous works have shown that $s_s \approx 3700$ Mpc is the frequency range that contains the global minimum [5,8]. We therefore search only around this global minimum region. Even so, for efficiency in the minimization it is important to choose combinations of the parameters that are close to the principal components of the curvature or covariance matrix. We discuss such choices in Appendix C.

Unlike the Planck collaboration analysis [5], we simultaneously vary the cosmological and step parameters in the minimization. This step is crucial, as discussed in Appendix B, since the presence of rapid oscillations also changes both the amplitude and shape of the broadband power in multipole space. If the cosmological parameters were not readjusted, the Planck data would falsely suggest that the oscillations could not continue into the $\ell \sim 10^3$ regime where the data is most constraining. For this reason, the minimum found in Ref. [5] is not the global minimum nor is there strong evidence for the damping of the oscillations at high multipoles. The minimum χ^2 as a function of the damping scale x_d is shown in Fig. 1 and is nearly flat for $x_d > 10^2$.

We find that the global minimum is given by

$$\begin{aligned}
C_2 &= 0.075, & s_s &= 3696.9 \text{ Mpc}, \\
x_d &= 105.0, & \Delta\chi^2 &= -14.0, \tag{14}
\end{aligned}$$

where the χ^2 improvement is measured against the best fit smooth model. In contrast, the best fit model of Ref. [5] had




FIG. 1. Minimum χ^2 relative to the best fit smooth (no step) model as a function of the oscillation damping scale x_d for the $c_s = 1$ potential step model (top panel). The minimization is performed jointly over cosmological and step parameters with step position s_s and amplitude of oscillations C_2 shown in the middle and bottom panels respectively.

$x_d = 87$ and a similar amplitude and frequency with a $\Delta\chi^2 = -11.7$.¹ As shown in Table III, this difference is not due to the inclusion of second order corrections from I_1 in Eq. (5), though their omission would bias cosmological parameters such as A_s and n_s . The Planck data thus favor oscillations at the few percent level in C_ℓ , with a peak-to-peak spacing of $\Delta\ell \sim 12$. Note that at $ks_s = x_d$, the damping suppression $\mathcal{D} = 0.85$ and for this model corresponds to $\ell_d \approx 400$. Thus, the oscillations persist at least out to the second acoustic peak (see Fig. 2).

The complete list of cosmological parameters for the best fit step model is given in Table III. The oscillations add broadband power and generally require a lower normalization. Because in this model they damp near the well-constrained third peak, this also requires a higher tilt to keep the total power fixed at this best constrained region. As a result, the model has slightly smaller broadband power at low multipoles, reaching $\sim -5\%$ at the quadrupole.

Furthermore, the Planck data are also compatible with oscillations that persist out to the highest multipole measured in the data, $\ell = 2500$. Taking $x_d \approx 2000$, which is indistinguishable from infinity for Planck,

¹The original version of the Planck collaboration analysis [5] erroneously conflated C_2 with A_c (see Eq. (C2) and correcting this definition also brings the amplitude into agreement with that found for WMAP [2].




FIG. 2 (color online). Best fit models for a potential step with $c_{sa} = 1.0$ (red) and a warp step at $c_{sa} = 0.7$ (blue). Both models have the same few percent oscillations around the best fit smooth model at the first and second peaks (lower panel). The warp step is a marginally better fit of $\Delta\chi^2 = -15.2$ versus the potential step with $\Delta\chi^2 = -14.0$, due to suppression of the lowest multipoles, but introduces the sound speed as an additional parameter.

$$C_2 = 0.043, \quad s_s = 3704.7 \text{ Mpc}, \quad \Delta\chi^2 = -11.4. \quad (15)$$

This fit only differs in χ^2 by 2.6 from that of the global minimum and is comparable to the best fit found in Ref. [5]. Note that even with no damping scale in curvature fluctuations, oscillations in C_ℓ decline with ℓ due to projection and lensing effects (see [2] and Eq. (C2)). Had we fixed cosmological parameters to the best fit smooth model then this x_d would be falsely penalized by $\Delta\chi^2 = 9$. The cosmological parameters for this model are also given in Table III. Notably, with no damping scale, the tilt no longer requires significant adjustment. The change in s_s mainly reflects slightly different cosmological parameters that produce correspondingly different distances to recombination rather than a change in the angular scale.

In summary, the Planck data favor percent level oscillations in C_ℓ produced by potential step features by $\Delta\chi^2 = -11.4$ with two parameters that control the oscillation frequency and amplitude. Minimizing the χ^2 for damping of the oscillations confines the oscillations to roughly the first and second peaks and marginally improves the fit with an additional $\Delta\chi^2 = -2.6$ for a total of -14.0 with one additional parameter for a total of three.

B. Low sound speed models

Low sound speed DBI models allow for two different classes of steps with two different phenomenologies that

impact low multipoles in the Planck data. Both steps in the potential $V(\phi)$ and warp $T(\phi)$ produce the same high multipole oscillations driven by the amplitude parameter C_2 . Given a C_2 that minimizes the Planck χ^2 at high multipoles, the remaining freedom is in choosing a sound speed after the step c_{sa} . In both the potential and warp scenarios, this uniquely fixes the two remaining step parameters C_1 and C_3 . Recall that C_1 controls the step in the power spectrum around the first oscillation and C_3 controls the shape of the first few oscillations. Since the fit is driven by the C_2 oscillations with only small impact from C_1 and C_3 , we fix all the other parameters to the global minimum of Eq. (14) when examining the impact of c_{sa} .

For potential steps, $C_1 = 0$ and $-3/8 < C_3/C_2 < 0$. Even for the maximal case of $-3/8$ and $c_{sa} \rightarrow 0$, there is very little impact on the CMB power spectrum. Consequently, as shown in Fig. 3, the χ^2 surface is essentially flat across c_{sa} .

For warp steps, both C_1/C_2 and C_3/C_2 can be greater than unity and the sound speed has a larger fractional effect on C_ℓ . However, their impact is still limited to the first few oscillations and, given the preference for a horizon scale s_s , severely cosmic variance limited. Raising c_{sa} mainly enhances the step in the power spectrum relative to the oscillations, thus lowering the first few multipoles. Both C_1 and C_3 are important in establishing the shape due to a cancellation in their effects at the first oscillation.

Since warp steps do not produce oscillatory features, as $c_s \rightarrow 1$, there is a maximum $c_{sa} \sim 0.7$ for which they can explain the oscillations (see Fig. 3). The best fit has the maximal possible sound speed

$$c_{sa} = 0.70, \quad \Delta\chi^2 = -15.2 \text{ (warp)}, \quad (16)$$




FIG. 3 (color online). χ^2 improvement as a function of the sound speed after the step c_{sa} for the warp (blue+) and potential (red \times) steps. Other parameters are fixed to their minimum χ^2 values in the $c_s = 1$ potential step model including the oscillation amplitude C_2 . Note that warp steps with $c_{sa} \gtrsim 0.7$ cannot generate the required C_2 (see Eq. (A44)), as marked by the vertical line.

which implies $C_1 = -0.70$, $C_3 = -0.37$ given the fixed parameters in Eq. (14). While the step in the curvature power spectrum is approximately 50% (see Fig. 9) in C_ℓ , this and the changes in the cosmological parameters translates into a $\sim 20\%$ suppression of power at the quadrupole relative to the smooth model (see Fig. 2). Note that the drop between $2 \leq \ell \leq 5$ is particularly sharp for warp steps due to a local maximum in the curvature spectrum oscillations. Nonetheless, with cosmic variance these changes have only a small impact on the fit. As a consequence, while warp steps have interesting phenomenology that may ameliorate low multipole anomalies, there is no statistically significant preference for $c_s < 1$.

IV. FUTURE TESTS

While an improvement of $\Delta\chi^2 \approx -11$ for two parameters and up to -15 in the full step parameter space may sound significant, it has been shown that for more flexible oscillatory models, where not only the amplitude and frequency but also the phase of the oscillation is fit, realizations of smooth models with noise often recover this level of improvement, albeit typically with a smaller oscillation amplitude [4] (see also [6]). Furthermore, the improvement in the WMAP likelihood [2,4] is comparable to that of the Planck likelihood despite the higher precision of Planck, whereas one would have expected the latter to increase for a true signal. For these reasons, it is important to have more definitive tests for the origin of these improvements. In this section we discuss predictions of the best fit models identified above that may be used to verify or falsify the hypothesis of their primordial origin.

As emphasized by Ref. [17], the most incisive consistency test for inflationary features is the E -mode polarization power spectrum and cross spectrum. In Fig. 4 (left panel), we show the predicted E -mode power spectrum of the models in Fig. 2. Consistency with inflationary oscillations demands that oscillations appear at the same frequency while modulated by the acoustic transfer to have nodes that are out of phase with the temperature. Furthermore, due to projection effects, the polarization oscillations are twice as prominent in polarization. In principle, the low ℓ . polarization can also more than double the distinguishing power between the warp and potential fits, albeit this is limited in practice by galactic foregrounds and uncertainties in the reionization model.

Finally, the temperature polarization cross spectrum must also exhibit consistent oscillations, as shown in Fig. 4 (right panel). These predictions should be tested in the next release of the Planck data. More generally they can be tested in any CMB polarization data set that has sufficient amounts of sky to distinguish modes separated by $\Delta\ell \approx 12$ and oscillations in power of 3–10%.

The best fit oscillatory models also predict different CMB lensing effects. High frequency features act in a similar fashion as the acoustic peaks in providing a signal




FIG. 4 (color online). Polarization (left) and temperature polarization cross (right) power spectra for the best fit models of Fig. 2. Step oscillations provide falsifiable predictions for the polarization which would not be mimicked by chance features in the noise.

for lensing. The $\Delta\ell \approx 12$ fineness of the features compared with the acoustic spacing of $\Delta\ell \approx 300$ offsets the smallness of the amplitude. In Fig. 5, we quantify this expectation by showing, for the smooth and best fit step models,

$$\frac{d \ln \ell^2 C_\ell}{d \ln \ell}, \quad (17)$$

which controls lensing and squeezed bispectrum effects [18]. Note that what was a small effect for the power spectrum is an order unity effect for certain lensing effects.




FIG. 5 (color online). Temperature power spectrum derivatives for the best fit models of Fig. 2. Low amplitude, high frequency oscillations produce new signals for lensing reconstruction and squeezed bispectra.

There are two related ways in which the oscillations impact lensing observables. First, if lensing reconstruction were performed with the oscillatory models to construct the filters in the optimal quadratic estimator rather than the smooth models, the noise power in the reconstruction should decrease if the oscillatory features are real. To see this note that in the flat-sky approximation the sample-variance limited reconstruction noise power N_L of the lensing potential is given by [18,19]

$$N_L^{-1} = \int \frac{d^2 \ell_1}{(2\pi)^2} \frac{(\mathbf{L} \cdot \ell_1 C_{\ell_1} + \mathbf{L} \cdot \ell_2 C_{\ell_2})^2}{2C_{\ell_1} C_{\ell_2}}, \quad (18)$$

where $\ell_1 + \ell_2 = \mathbf{L}$. For $L \ll \ell_1$, $\ell_1 \approx -\ell_2$ and for $L \ll \Delta\ell$, C_{ℓ_2} can be Taylor expanded around ℓ_1 . The numerator then scales as the derivative in Eq. (17) squared. For the best fit models, this changes the noise for $L \lesssim 12$.

Relatedly, lensing by long-wavelength modes modulates the angular scale of features in the power spectrum which itself is correlated with the CMB temperature anisotropy through the integrated Sachs-Wolfe (ISW) effect. Thus, the presence of fine scale oscillations changes the squeezed reduced bispectrum of temperature fluctuations [20],

$$b_{L\ell_1\ell_2} \approx \frac{L(L+1) - \ell_1(\ell_1+1) + \ell_2(\ell_2+1)}{2} C_L^{T\phi} C_{\ell_2} + 5\text{perm}, \quad (19)$$

where $C_L^{T\phi}$ is the correlation of the ISW temperature and lensing potential fields. The permutation of $\ell_1 \leftrightarrow \ell_2$ again makes the result scale as the derivative of C_ℓ and is

enhanced by the oscillation. Both of these lensing effects are in principle detectable, though for the best fit frequency with $\Delta\ell \sim 12$ the impact will be limited by cosmic variance. A more detailed study is required to determine their effects on the existing Planck data set.

Features during inflation also produce primordial non-Gaussianity in mainly the equilateral configuration [9,21–23]. For the best fit step models, these should also be observable in Planck [2,24]. Extracting these signals, though, will require using specific templates that include these rapid oscillations [25]. Since the equilateral bispectrum amplitude scales as x_d^2 , the lack of a strong bound on the damping scale implies that the bispectrum signal could be very large at high multipoles, though these models would be beyond the regime of validity of the effective field theory that underlies their calculation [26].

Thus, if the oscillatory fits really reflect inflationary features, there is a battery of consistency tests that the CMB temperature and polarization anisotropy must satisfy.

V. DISCUSSION

In this paper, we have extended and improved the modeling and analysis of sharp inflationary steps for the Planck CMB power spectrum. We find that for the two parameters of the amplitude and frequency of the oscillations, step models improve the fit by $\Delta\chi^2 = -11.4$, whereas additional parameters such as the finite width of the step and sound speed of the inflaton marginally improve the fit to $\Delta\chi^2 = -14.0$ and -15.2 , respectively. In particular, sound speed effects for warp steps lower than the quadrupole power by $\sim 20\%$.

We have shown that it is critical to jointly fit step and cosmological parameters simultaneously. If cosmological parameters, especially the amplitude and tilt, are held fixed then one would falsely infer that the oscillations must damp away at high multipoles due to their excess average power. We have shown that on the contrary there is only marginal preference for a finite damping scale. The improvement in modeling to second order terms in the generalized slow-roll approximation developed here is also required by the increased precision of Planck at high multipoles but their omission would mainly bias the cosmological parameters rather than degrade the fit itself.

Given that chance features in the noise can masquerade as oscillatory step features [4], we have also provided a suite of consistency tests that can verify or falsify the primordial origin of these improved fits. The polarization power and cross spectra should reveal a matching and larger set of oscillations modulated by an out of phase acoustic transfer. The oscillations, if primordial, also provide an extra signal for CMB lensing reconstruction and squeezed bispectra from the lensing-ISW correlation. Finally, the primordial non-Gaussianity in equilateral bispectrum configurations should also be observable.

These predictions may soon be tested in the next release of Planck data.

ACKNOWLEDGMENTS

We thank Peter Adshead, Aurelien Benoit-Levy, Douglas H. Rudd, and Shi Chun Su for useful discussions. We are also grateful for the support of the University of Chicago Research Computing Center for assistance with the calculations carried out in this work. W.H. was supported by the Kavli Institute for Cosmological Physics at the University of Chicago through Grants No. NSF PHY-0114422 and No. NSF PHY-0551142 and an endowment from the Kavli Foundation and its founder Fred Kavli. V. M. and W. H. were additionally supported by U.S. Department of Energy Contract No. DE-FG02-90ER-40560, W. H. by the David and Lucile Packard Foundation, and V.M. by the Brazilian Research Agency CAPES Foundation and by the U.S. Fulbright Organization.

APPENDIX A: ANALYTIC STEP SPECTRUM

In this Appendix, we derive and test the analytic model for the power spectrum used in the main paper, extending previous treatments [2,13] for large, sharp steps in the warp and potential in the DBI context. These necessitate second order corrections to achieve the precision required for the Planck data. We begin with a brief review of the generalized slow-roll (GSR) approximation [27,28] for large power spectrum features in Sec. A 1 [29]. In Sec. A 2, we use exact energy conservation and the attractor solutions before and after the step to derive the general analytic model for the power spectrum in the GSR approximation. In Sec. A 3 and Sec. A 4 we apply this model to steps in the warp and potential.

1. Generalized slow roll

In a general $P(X, \phi)$ model for inflation, the comoving curvature power spectrum,

$$\Delta_{\mathcal{R}}^2 \equiv \frac{k^3 P_{\mathcal{R}}}{2\pi^2} = \lim_{ks \rightarrow 0} \left| \frac{ksy}{f} \right|^2, \quad (\text{A1})$$

is evaluated by solving the field or mode function equation in a spatially flat gauge [28,30],

$$\frac{d^2 y}{ds^2} + \left(k^2 - \frac{2}{s^2} \right) y = \left(\frac{f'' - 3f'}{f} \right) \frac{y}{s^2}. \quad (\text{A2})$$

Here, deviations from de Sitter space are characterized by

$$f^2 = 8\pi^2 \frac{\epsilon_{HCs}}{H^2} \left(\frac{aHs}{c_s} \right)^2, \quad (\text{A3})$$

where

$$\epsilon_H = -\frac{d \ln H}{dN}, \quad (\text{A4})$$

with N as e-folds, $N = 0$ as the end of inflation, c_s denoting the sound speed of field fluctuations, and

$$s(N) = \int_N^0 d\tilde{N} \frac{c_s}{aH} \quad (\text{A5})$$

denoting the sound horizon. Here, and throughout, $' \equiv d/d \ln s$.

Eq. (A2) can be formally solved with the Green function technique by taking its right-hand side as an external source given by the de Sitter mode function with Bunch-Davies initial conditions,

$$y \approx y_0 = \left(1 + \frac{i}{ks}\right) e^{iks}, \quad (\text{A6})$$

and iteratively improving the solution for the presence of the deviations introduced by f . Including the leading second order correction for large features, the power spectrum is given by [29,31]

$$\begin{aligned} \ln \Delta_{\mathcal{R}}^2 \approx & G(\ln s_{\min}) + \int_{s_{\min}}^{\infty} \frac{ds}{s} W(ks) G'(\ln s) \\ & + \ln [1 + I_1^2(k)], \end{aligned} \quad (\text{A7})$$

where the source function

$$G = -2 \ln f + \frac{2}{3} (\ln f)', \quad (\text{A8})$$

and recall W is given by Eq. (8). The validity of the approximation relies on the deviations in the mode functions or the curvature being small rather than the G' deviations from slow roll. It is monitored by the second order corrections

$$I_1(k) = \frac{1}{\sqrt{2}} \int_0^{\infty} \frac{ds}{s} G'(\ln s) X(ks), \quad (\text{A9})$$

with $u = ks$ and X given by Eq. (8). The GSR approximation itself will begin to break down unless [32]

$$I_1 \lesssim \frac{1}{\sqrt{2}}. \quad (\text{A10})$$

Finally, G' carries both the smooth tilt type deviations from scale invariance as well as any impact of sharp features. For sharp features, it can be approximated by [13]

$$\begin{aligned} G' \approx & (1 - n_s) - \frac{1}{3} \sigma_2 + \frac{2}{3} \delta_2 - \frac{5}{3} \sigma_1 - 2\eta_H \\ & + \frac{8}{3} \left(\frac{aHs}{c_s} - 1 \right), \end{aligned} \quad (\text{A11})$$

where the additional slow-roll parameters are defined by

$$\begin{aligned} \eta_H \equiv & \epsilon_H - \frac{1}{2} \frac{d \ln \epsilon_H}{dN}, & \delta_2 \equiv & \epsilon_H \eta_H + \eta_H^2 - \frac{d\eta_H}{dN}, \\ \sigma_1 \equiv & \frac{d \ln c_s}{dN}, & \sigma_2 \equiv & \frac{d\sigma_1}{dN}, \end{aligned} \quad (\text{A12})$$

and we reabsorb the slow-roll deviations in these parameters into the $(1 - n_s)$ factor. Thus, modeling a sharp feature amounts to determining its impact on c_s and ϵ_H .

2. Step sources

We consider sharp steps in the warp $T(\phi)$ and potential $V(\phi)$ of the DBI Lagrangian (1) which generate analogous changes in c_s and ϵ_H . To keep our treatment general, we first parametrize the evolution of these quantities relying on energy conservation and the attractor solution to define their functional form. In the following subsections we give the correspondence of this parametrization to specific step parameters.

The energy density of the inflaton,

$$\rho = \left(\frac{1}{c_s} - 1 \right) T + V, \quad (\text{A13})$$

is conserved as long as the inflaton rolls across the step in much less than an e-fold. This conservation then gives the relationship between the sound speed before (“ b ”; c_{sb}) and immediately after (“ i ”; c_{si}) the step. The acceleration equation

$$\epsilon_H = \frac{3\rho + p}{2\rho} \approx \frac{3}{2V} T \left(\frac{1}{c_s} - c_s \right) \quad (\text{A14})$$

then gives the corresponding change in ϵ_H . After the step, the rolling of the inflaton,

$$\phi_N^2 \equiv \left(\frac{d\phi}{dN} \right)^2 = 2c_s \epsilon_H, \quad (\text{A15})$$

differs from the friction dominated attractor solution,

$$\phi_N = -\frac{c_s}{3} \frac{V_\phi}{H^2} \approx -c_s \frac{V_\phi}{V}, \quad (\text{A16})$$

to which it must decay on the expansion time scale or well after the inflaton has crossed the step (“ a ”). These relations hold for arbitrarily large steps so long as $\epsilon_H \ll 1$.

Together, they imply that the functional form of c_s and ϵ_H is generically given by [13]

$$\begin{aligned}\frac{c_s}{c_{sa}} &= 1 + \frac{1-c_b}{2}F + \frac{c_i-1}{2}(F+2)e^{3(N_s-N)}, \\ \frac{\epsilon_H}{\epsilon_{Ha}} &= 1 + \frac{1-e_b}{2}F + \frac{e_i-1}{2}(F+2)e^{3(N_s-N)},\end{aligned}\quad (\text{A17})$$

where for convenience we have scaled the quantities to their values on the attractor after the step

$$\begin{aligned}c_b &= \frac{c_{sb}}{c_{sa}}, & c_i &= \frac{c_{si}}{c_{sa}}, \\ e_b &= \frac{\epsilon_{Hb}}{\epsilon_{Ha}}, & e_i &= \frac{\epsilon_{Hi}}{\epsilon_{Ha}}.\end{aligned}\quad (\text{A18})$$

Here, F represents a step of infinitesimal width at $N = N_s$ normalized to -2 before the step and 0 after. We discuss the impact of the finite width below.

Following Ref. [13], it is straightforward to derive the source function G' in the approximation that changes to c_s and ϵ_H are small by taking their derivatives and integrals to form the quantities in Eq. (A11). Note that this limit does not necessarily require the steps in the warp itself to be small. In the limit of a large warp factor $\phi_N^2/T \ll 1$, the sound speed approaches unity regardless of the form of T and hence the change in the sound speed is small even for a large fractional change in T .

Integrals over G' are then simply evaluated by recalling that $dF/d \ln s$ is a delta function of amplitude 2. The result of integrating the source by parts is

$$\begin{aligned}\ln \Delta_{\mathcal{R}}^2 &\approx \ln A_s \left(\frac{k}{k_0}\right)^{n_s-1} + C_1 W(k s_s) + C_2 W'(k s_s) \\ &+ C_3 Y(k s_s),\end{aligned}\quad (\text{A19})$$

where

$$\begin{aligned}C_1 &= -(c_b - 1) - (e_b - 1), \\ C_2 &= -\frac{1}{3}(c_i - c_b) + \frac{1}{3}(e_i - e_b), \\ C_3 &= (1 - c_b) + \frac{1}{4}(c_i - 1),\end{aligned}\quad (\text{A20})$$

for the leading order GSR contribution in Eq. (A7) which, once corrected for the finite width of the step below, we shall call GSR0. Here, we have replaced the parameter $G(\ln s_{\min})$ in Eq. (A7) with the power spectrum normalization A_s at k_0 . Note that

$$Y(x) \equiv -\frac{8}{3}x \int d \ln \tilde{x} \frac{W'(\tilde{x})}{\tilde{x}},\quad (\text{A21})$$

which is given in closed form in Eq. (8).

Given that

$$\begin{aligned}\lim_{x \ll 1} W(x) &= 1, & \lim_{x \gg 1} W(x) &= 0, \\ \lim_{x \ll 1} W'(x) &= 0, & \lim_{x \gg 1} W'(x) &= -3 \cos(2x), \\ \lim_{x \ll 1} Y(x) &= 0, & \lim_{x \gg 1} Y(x) &= 0,\end{aligned}\quad (\text{A22})$$

we can further interpret the meaning of the C_i coefficients. C_1 represents a step in the power spectrum and its amplitude is determined by the fact that the inflaton is on the attractor solution before and well after the step. C_2 provides a constant amplitude oscillation whose value is determined by the sharpest part of the feature: the fractional changes in c_s and ϵ_H right at the step. Finally C_3 modifies the shape of the first few oscillations due to the $aHs/c_s - 1$ source.

Likewise, the first order corrections are given by

$$\sqrt{2}I_1 = \frac{\pi}{2}(1 - n_s) + C_1 X(k s_s) + C_2 X'(k s_s) + C_3 Z(k s_s),\quad (\text{A23})$$

where

$$Z(x) = -\frac{8}{3}x \int d \ln \tilde{x} \frac{X'(\tilde{x})}{\tilde{x}},\quad (\text{A24})$$

which is given in closed form in Eq. (8).

We call the analytic model with the I_1 correction GSR1. Using Eq. (A10), we thus expect the GSR expansion itself to be under control for $k s_s \gg 1$ so long as

$$|C_2| < 1/3.\quad (\text{A25})$$

At $k s_s \sim 1$, the exact requirement is a model dependent restriction on a combination of C_1, C_2, C_3 but in the warp and potential step examples this gives roughly the same criteria for the step height.

In principle, this domain of validity includes fractional deviations in c_s and ϵ_H that approach unity, including the region of interest for Planck. However, although the GSR expansion itself remains under control, Eq. (A20) is derived by assuming small fractional deviations and requires correction. Just as we extended the validity of the step approximation to nonlinearities in the step amplitude above, we can also approximately correct for weak non-linearity in the slow-roll parameters by rescaling the C_i coefficients. Here, we extend and generalize the approach of Ref. [2] and [13] for arbitrary sound speeds.

The C_1 amplitude gives the step in power and hence the slow-roll attractor $\Delta_{\mathcal{R}}^2 \propto (c_s \epsilon_H)^{-1}$ determines it as

$$C_1 = -\ln c_b e_b.\quad (\text{A26})$$

The changes in c_s and ϵ_H at the step are already determined nonlinearly and so the only further correction to C_2 comes

from the conversion to slow-roll parameters, e.g. $\sigma_1 = c_s^{-1} dc_s/dN$. Following Ref. [2], we evaluate c_s and ϵ_H at the midpoint of the step and hence

$$C_2 = -\frac{2c_i - c_b}{3c_i + c_b} + \frac{2e_i - e_b}{3e_i + e_b}. \quad (\text{A27})$$

Finally, for C_3 , while there is no direct nonlinear constraint to determine its amplitude, by also renormalizing to the midpoint of the step we approximately preserve the relative relationship between the coefficients that determines the shape of their combined contributions. This is especially important for warp steps where cancellations between C_1 and C_3 occur around the first oscillation. Thus, we take

$$C_3 = 2 \frac{(1 - c_b) + (c_i - 1)/4}{c_i + c_b}. \quad (\text{A28})$$

Finally, we can account for the finite width of the step. If we replace the step function with a tanh function,

$$F(\phi) = \tanh\left(\frac{\phi - \phi_s}{d}\right) - 1, \quad (\text{A29})$$

the integrals over G' will not contribute if the windows in Eq. (8) oscillate many times over the width of the step $k \gg s_s d / \phi_N$. This causes a damping such that the C_i coefficients in Eqs. (A19) and (A23) are replaced by [2]

$$C_i \rightarrow C_i \mathcal{D}\left(\frac{ks_s}{x_d}\right), \quad (\text{A30})$$

where the damping function

$$\mathcal{D}(y) = \frac{y}{\sinh(y)}, \quad (\text{A31})$$

with the damping scale

$$x_d = \frac{1}{\pi d} \frac{d\phi}{d \ln s}. \quad (\text{A32})$$

The derivation of Eq. (5) for the functional form for the analytic model of the step power spectrum is thus complete. We now turn to specific forms for warp and potential steps.

3. Warp steps

For steps in the warp T , we have before and after the step

$$T_b = T_a(1 - 2b_T), \quad V_b = V_a, \quad (\text{A33})$$

and we can then use energy conservation and the attractor solution to give the relevant c and e parameters of the model in Eq. (A18). Our convention is to quote these in terms of b_T and the sound speed on the attractor after the step c_{sa} . The attractor solution tells us that

$$c_b = \sqrt{\frac{1 - 2b_T}{1 - 2b_T c_{sa}^2}}, \quad (\text{A34})$$

and, using Eq. (A14) for ϵ_H , we obtain

$$e_b = c_b. \quad (\text{A35})$$

Now let us consider the sharp changes immediately after the step. Energy conservation tells us

$$\frac{1}{c_{si}} - 1 = \left(\frac{1}{c_{sb}} - 1\right)(1 - 2b_T), \quad (\text{A36})$$

or

$$c_i = \frac{c_b}{1 - 2b_T(1 - c_{sa}c_b)}, \quad (\text{A37})$$

and, with Eq. (A14),

$$e_i = \frac{1 - c_{sa}^2 c_i^2}{c_i(1 - c_{sa}^2)}. \quad (\text{A38})$$

We show an example of the evolution of c_s and ϵ_H in Fig. 6. Since the analytic model only captures the evolution of the parameters around the step and not the evolution on the slow-roll attractor, we plot




FIG. 6 (color online). Evolution of c_s (upper) and ϵ_H (lower) across a warp step. The step in both parameters and transient behavior right after the step is modeled by Eqs. (A17) and (A39) to excellent approximation. Model parameter choices are given in Appendix B.

$$c_s(N - N_s) = c_s^{\text{an}}(N - N_s) \frac{c_s^{\text{at}}(N - N_s)}{c_s^{\text{at}}(0^\pm)}, \quad (\text{A39})$$

where c_s^{an} is the analytic model of Eq. (A17), c_s^{at} is the attractor on either side of the step, and $c_s^{\text{at}}(0^\pm)$ is evaluated approaching the step from either side with $c_{sa} = c_s^{\text{at}}(0^+)$ approached from the side after the step. We likewise account for the slow-roll evolution of ϵ_H . In practice, rather than iterating the attractor solution of Eq. (A16) in the equations of motion to the required accuracy, we numerically solve the equivalent smooth model before and after the step to determine c_s^{at} .

We can now use the general description of Eqs. (A26), (A27), and (A28) to give the C_i coefficients of the analytic power spectrum form. Note that in the small step limit,

$$\begin{aligned} \lim_{b_T \rightarrow 0} c_b &= 1 - (1 - c_{sa}^2)b_T, \\ \lim_{b_T \rightarrow 0} c_i &= 1 + (1 - c_{sa})^2 b_T, \\ \lim_{b_T \rightarrow 0} e_i &= 1 - \frac{(1 - c_{sa})(1 + c_{sa}^2)b_T}{1 + c_{sa}}, \end{aligned} \quad (\text{A40})$$

and so

$$\begin{aligned} \lim_{b_T \rightarrow 0} C_1 &= 2(1 - c_{sa}^2)b_T, \\ \lim_{b_T \rightarrow 0} C_2 &= -\frac{2}{3} \frac{1 - c_{sa}}{1 + c_{sa}} b_T, \\ \lim_{b_T \rightarrow 0} C_3 &= \frac{1}{4}(5 - 2c_{sa} - 3c_{sa}^2)b_T, \end{aligned} \quad (\text{A41})$$

in agreement with Ref. [13]. Our generalized expression lets us explore the $b_T \rightarrow -\infty$ limit,

$$\begin{aligned} \lim_{b_T \rightarrow -\infty} c_b &= \frac{1}{c_{sa}}, \\ \lim_{b_T \rightarrow -\infty} c_i &= \frac{2c_{sa}}{1 + c_{sa}^2}, \\ \lim_{b_T \rightarrow -\infty} e_i &= \frac{1}{2c_{sa}} \frac{1 + 3c_{sa}^2}{1 + c_{sa}^2}. \end{aligned} \quad (\text{A42})$$

Note that for finite c_{sa} these limits are all finite and so the maximal C_i amplitudes are also bounded:

$$\begin{aligned} \lim_{b_T \rightarrow -\infty} C_1 &= 2 \ln c_{sa}, \\ \lim_{b_T \rightarrow -\infty} C_2 &= 4 \frac{1 - c_{sa}^4}{9 + 42c_{sa}^2 + 45c_{sa}^4}, \\ \lim_{b_T \rightarrow -\infty} C_3 &= -\frac{1}{2} \frac{4 - 3c_{sa} + 2c_{sa}^2 - 3c_{sa}^3}{1 + 3c_{sa}^2}. \end{aligned} \quad (\text{A43})$$

Thus, for a fixed observed oscillation amplitude $C_2 > 0$, there is always a maximum c_s for which a warp step cannot explain the data,

$$c_{sa}^2|_{\text{max}} = -\frac{21C_2}{4 + 45C_2} + \frac{2\sqrt{4 + 36C_2 + 9C_2^2}}{4 + 45C_2}. \quad (\text{A44})$$

For example, if $C_2 = 1/15$, $c_{sa}^2|_{\text{max}} \approx 0.724$.

4. Potential steps

For potential steps,

$$\begin{aligned} V_b &= V_a(1 - 2b_V), \\ T_b &= T_a. \end{aligned} \quad (\text{A45})$$

The attractor solution says that to the leading order in ϵ_H , there is no net change in c_s or ϵ_H , only a transient deviation at the step. Thus, $\epsilon_{Hb} = \epsilon_{Ha}$ and $c_{sa} = c_{sb}$ ($c_b = e_b = 1$) or

$$C_1 = 0. \quad (\text{A46})$$

We can also use the attractor to eliminate T/V_a using

$$\epsilon_{Ha} = \frac{3}{2} \frac{T}{V_a} \left(\frac{1}{c_{sa}} - c_{sa} \right). \quad (\text{A47})$$

Energy conservation then gives the transient change as

$$c_i = 1 - \frac{3b_V(1 - c_{sa}^2)}{3b_V(1 - c_{sa}^2) - \epsilon_{Ha}}, \quad (\text{A48})$$

and, using Eq. (A14),




FIG. 7 (color online). Maximum oscillation amplitude C_2 for a warp step as a function of c_{sa} , the sound speed after the step. The blue dashed line corresponds to $C_2 = 1/3$, where the GSR approximation breaks down in the oscillatory regime. Note that for $c_s \gtrsim 1/4$ the oscillation amplitude is limited by physicality rather than the GSR approximation.

$$e_i = 1 - \frac{3b_V[-3b_V(1 - c_{sa}^2) + (1 + c_{sa}^2)\epsilon_{Ha}]}{\epsilon_{Ha}[-3b_V(1 - c_{sa}^2) + \epsilon_{Ha}]} \quad (\text{A49})$$

The general description of Eqs. (A26), (A27), and (A28) then gives the C_i coefficients of the analytic power spectrum form. We show an example of the evolution of c_s and ϵ_H in Fig. 8. Again, to capture the slow-roll evolution of the smooth model, we plot the analytic model corrected, as in Eq. (A39).

In the limit of a small potential step,

$$\lim_{b_V \rightarrow 0} c_i = 1 + 3 \frac{1 - c_{sa}^2}{\epsilon_{Ha}} b_V, \quad \lim_{b_V \rightarrow 0} e_i = 1 - 3 \frac{1 + c_{sa}^2}{\epsilon_{Ha}} b_V, \quad (\text{A50})$$

and so

$$\lim_{b_V \rightarrow 0} C_2 = -\frac{2}{\epsilon_{Ha}} b_V, \quad \lim_{b_V \rightarrow 0} C_3 = \frac{3}{4} \frac{1 - c_{sa}^2}{\epsilon_{Ha}} b_V, \quad (\text{A51})$$

which generalizes the results of Ref. [2] to arbitrary sound speed. The sound speed experiences a transient dip for downward steps $b_V < 0$.

Note that, in the opposite limit,

$$\lim_{b_V \rightarrow -\infty} c_i = \frac{1}{3b_V} \frac{\epsilon_{Ha}}{1 - c_{sa}^2}, \quad \lim_{b_V \rightarrow -\infty} e_i = \frac{3b_V}{\epsilon_{Ha}}, \quad (\text{A52})$$

and so




FIG. 8 (color online). Evolution of c_s (upper) and ϵ_H (lower) across a potential step. The transient change in both parameters is modeled by Eqs. (A17) and (A39) to excellent approximation. Model parameter choices are given in Appendix B.

$$\lim_{b_V \rightarrow -\infty} C_2 = \frac{4}{3}, \quad \lim_{b_V \rightarrow -\infty} C_3 = -\frac{1}{2}. \quad (\text{A53})$$

Since these amplitudes are beyond the limits of the GSR approximation itself according to Eq. (A25), there is effectively no relevant bound on the oscillation amplitude set by energy conservation and the attractor solution unlike the warp step case. Likewise, for a given $0 < C_2 \ll 4/3$, there is no bound on the required sound speed.

APPENDIX B: POWER SPECTRUM ACCURACY

In this section, we test the accuracy of the leading order GSR0 approximation used in previous analyses [2] and the first order GSR1 corrections discussed in Appendix A against an exact computation of the power spectrum from the DBI Lagrangian of Eq. (1). Although GSR0 was previously demonstrated to be sufficiently accurate for WMAP data [2,13], we show here that the increase in precision to the 10^{-3} level in Planck requires second order corrections.

The exact computation of the power spectrum follows from solving Eq. (A2) for a DBI step model that is parametrized by $\{V_0, \beta, \lambda_b, \phi_{\text{end}}\}$, defining the broadband amplitude and slope of the power spectrum, and the step parameters $\{\phi_s, b_T, b_V, d\}$ defining the step position, height parameters, and width (see [13] for computational details). For testing purposes, we choose

$$V_0 = 7.1038 \times 10^{-26}, \quad \beta = 5.5895 \times 10^{-2}, \\ \lambda_b = 2.1771 \times 10^{14}, \quad \phi_{\text{end}} = 8.2506 \times 10^{-8}, \quad (\text{B1})$$

and

$$\phi_s = 3.8311 \times 10^{-8}, \quad d = 9.3835 \times 10^{-13}. \quad (\text{B2})$$

For the warp step, we choose

$$b_T = -3.364, \quad b_V = 0 \text{ (warp)}, \quad (\text{B3})$$

and, for the potential step,

$$b_T = 0, \quad b_V = -6.543 \times 10^{-21} \text{ (potential)}. \quad (\text{B4})$$

These parameters are in fact chosen to be close to the Planck maximum likelihood solution for the amplitude and frequency of warp step oscillations by inverting the steps in this test. Notice that in that case, $|b_T|$, the fractional change in the warp T , exceeds unity. The width d is set so that damping occurs in the $\ell \sim 10^3$ region that Planck is most sensitive to so as to yield the most stringent test of accuracy. The cosmological parameters for the test are given in Table I and coincide with the best fit model without a step.

The analytic models are specified by the conversion of the fundamental parameters into the amplitude parameters

$\{C_1, C_2, C_3\}$, the sound horizon at the step s_s , the effective number of oscillations before damping x_d , as well as the broadband amplitude and tilt parameters A_s and n_s .

Given a solution to the background equations without the step, we set the parameters,

$$c_{sa} = 0.67, \quad \epsilon_{Ha} = 1.70 \times 10^{-19}, \quad (\text{B5})$$

according to their values at $N = N_s$. The C_i amplitude parameters are then determined by Eqs. (A26)–(A28) such that

$$C_1 = -0.65, \quad C_2 = 0.071, \quad C_3 = -0.34 (\text{warp}), \quad (\text{B6})$$

and

$$C_1 = 0, \quad C_2 = 0.071, \quad C_3 = -0.015 (\text{potential}). \quad (\text{B7})$$

Next, the physical scale associated with the step has to be set very precisely in order not to have a phase error after many oscillations. We follow Ref. [13] in defining it numerically to be the sound horizon at which the deviation in the GSR source function due to the step is appropriately centered to a small fraction of the step width,

$$G'(\ln s_s, b_{T,V}) - G'(\ln s_s, 0) = 0. \quad (\text{B8})$$

Using this definition, we obtain

$$s_s = \begin{cases} 3699 \text{ Mpc} & (\text{warp}) \\ 3708 \text{ Mpc} & (\text{potential}) \end{cases}. \quad (\text{B9})$$

Note that although the step is at the same position in field space in both cases, the sound horizon differs slightly due to the change in c_s .

For the damping parameter, we likewise convert the field width d to a physical width s_s/x_d with the numerical solution for $\phi(\ln s)$ through

$$\left. \frac{d\phi}{d \ln s} \frac{1}{\pi d} \right|_{s=s_s} = x_d. \quad (\text{B10})$$

For the test cases, we obtain

$$x_d = \begin{cases} 170.0 & (\text{warp}) \\ 169.9 & (\text{potential}) \end{cases}.$$

Finally, there are the broadband power parameters n_s and A_s . For the tilt parameter, which is slowly varying and essentially independent of the step, we take the slope at $k = k_0 = 0.08 \text{ Mpc}^{-1}$ of the model with $b_{T,V} = 0$. We have chosen parameters in Eq. (B1) so that n_s coincides with the value given in Table I. On the other hand, the effective amplitude A_s depends on the presence of the step as well as the order of the GSR approximation used. In Eq. (A19), the broadband power gains a contribution from the average of the oscillations

$$\begin{aligned} \langle e^{C_1 W + C_2 W' + C_3 Y} \rangle &\approx \mathcal{I}_0 \left[3C_2 \mathcal{D} \left(\frac{ks_s}{x_d} \right) \right] \\ &\approx 1 + \left[\frac{3}{2} C_2 \mathcal{D} \left(\frac{ks_s}{x_d} \right) \right]^2 + \mathcal{O}(C_2^4), \end{aligned} \quad (\text{B11})$$




FIG. 9 (color online). GSR approximations versus exact solutions for the curvature power spectrum (top panel) and the fractional error of the GSR0 and GSR1 analytical solutions (bottom panel). The models are warp step (left) and potential step (right).

where \mathcal{I}_0 here is the modified Bessel function, not to be confused with the GSR integral I_0 . This nonzero average is the fundamental reason why cosmological parameters must be varied jointly with the step parameters when analyzing the Planck data. In the first order correction Eq. (A7), there is the analogous averaging effect

$$\langle I_1^2 \rangle \approx \frac{\pi^2}{8} (1 - n_s)^2 + \left[\frac{3}{2} C_2 \mathcal{D} \left(\frac{k s_s}{x_d} \right) \right]^2, \quad (\text{B12})$$

which is also $\mathcal{O}(C_2^2)$ despite being higher order in the GSR approximation. Moreover, around the damping scale set by x_d , the broadband average of the oscillation changes with k in Eqs. (B11)–(B12) and is not purely an amplitude shift. Note that the error induced by this average term scales as $\delta C_l / C_l \propto C_2^2$ and so rapidly increases with the amplitude of the oscillations.

Since the best choice for A_s depends on both the method and the data set considered, we choose A_s as the amplitude which gives the best agreement between the exact computation and the given GSR computation for the Planck data set. We therefore use the Planck likelihood itself to define A_s for each method. In order to remove the ambiguity caused by the exact model not possessing the maximum likelihood normalization, we in practice maximize both the Planck likelihood over A_s to obtain A'_s and a rescaling of the amplitude of the exact model by R to obtain its best normalization. We then set $A_s = A'_s / R$ to remove the rescaling.

In Fig. 10 we show the residual errors in the GSR0 and GSR1 after the normalization has been set in this way. Note that the residuals $\delta C_l / C_l$ cross zero at $\ell \sim 10^3$, reflecting the pivot or best constrained portion of the Planck spectrum. For scales much smaller or much larger than the

damping scale of the oscillation, the difference between GSR0, GSR1, and exact is nearly constant and can be absorbed into the normalization. However, in our test case, which we have chosen to be the worst case scenario, the damping falls exactly at the pivot. The result is that even with the best fit normalization, GSR0 produces $\sim 1\%$ errors that pivot around $\ell \sim 10^3$. While the error in GSR0 can mainly be absorbed by adjusting cosmological parameters such as the tilt, they are large enough to bias such parameters' non-negligibly. These residuals are reduced to the $\sim 0.1\%$ level with the GSR1 approximation.

More quantitatively, for these specific test cases the residuals produce a change in the Planck likelihood versus exact of

$$\Delta\chi^2 = \begin{cases} -8.6 & \text{GSR0} \\ -0.97 & \text{GSR1} \end{cases} (\text{warp}), \quad (\text{B13})$$

and

$$\Delta\chi^2 = \begin{cases} -7.4 & \text{GSR0} \\ -0.33 & \text{GSR1} \end{cases} (\text{potential}). \quad (\text{B14})$$

Thus, the GSR1 approximation is sufficiently accurate for the Planck analysis. In fact, the χ^2 errors would be even smaller at its global minimum.

The error in these approximations also depends on the step parameter model. For reference, if $x_d \rightarrow \infty$ the error in the GSR0 approximation becomes $\Delta\chi^2 = 0.5$ for the warp step and $\Delta\chi^2 = 0.7$ for the potential step. Here, the error is significantly lower since it takes the form of a constant amplitude rescaling which can be absorbed into A_s .




FIG. 10 (color online). GSR approximations versus exact solutions for the temperature power spectrum (top panel) and the fractional error of the GSR0 and GSR1 analytical solutions (bottom panel). The models are warp step (left) and potential step (right).

In our examples the result of the normalization procedure is to set

$$A_s = \begin{cases} 2.1432 \times 10^{-9} & \text{GSR0} \\ 2.1295 \times 10^{-9} & \text{GSR1} \end{cases} \text{ (warp)} \quad (\text{B15})$$

for the warp step and

$$A_s = \begin{cases} 2.1425 \times 10^{-9} & \text{GSR0} \\ 2.1288 \times 10^{-9} & \text{GSR1} \end{cases} \text{ (potential)} \quad (\text{B16})$$

for the potential step. In the absence of a step ($b_T = b_V = 0$), the same procedure would yield $A_s = 2.1554 \times 10^{-9}$, which is 1% different from the GSR1 value. These changes reflect the broadband power introduced by the oscillations in each case. Given the 0.1% precision of the Planck data, these differences are significant and A_s cannot be held fixed when fitting to step models.

For the minimization procedure in Appendix C, it is nonetheless useful to have an approximate prescription for the renormalization of A_s in the presence of oscillations. Given the average of the oscillatory pieces in Eqs. (B11)–(B12), the normalization parameter that the data should hold approximately fixed is

$$\tilde{A}_s = A_s e^{-2\tau} (1 + \bar{O}), \quad (\text{B17})$$

where \bar{O} contains the average of the oscillatory pieces in each approximation,

$$\bar{O} = \begin{cases} \frac{9}{4} C_2^2 \mathcal{D}^2 \left(\frac{k_0 s_s}{x_d} \right) & \text{GSR0} \\ \frac{\pi^2}{8} (1 - n_s)^2 + \frac{9}{2} C_2^2 \mathcal{D}^2 \left(\frac{k_0 s_s}{x_d} \right) & \text{GSR1} \end{cases}. \quad (\text{B18})$$

Here, the $e^{-2\tau}$ factor accounts for the change in the heights of the acoustic peaks due to anisotropy suppression by scattering during reionization. In our test cases,

$$\tilde{A}_s = \begin{cases} 1.8000 \times 10^{-9} & \text{GSR0} \\ 1.7999 \times 10^{-9} & \text{GSR1} \end{cases} \text{ (warp)} \quad (\text{B19})$$

for the warp step,

$$\tilde{A}_s = \begin{cases} 1.7993 \times 10^{-9} & \text{GSR0} \\ 1.7993 \times 10^{-9} & \text{GSR1} \end{cases} \text{ (potential)} \quad (\text{B20})$$

for the potential step, and $\tilde{A}_s = 1.8021 \times 10^{-9}$ without a step. Note that \tilde{A}_s absorbs most of the changes in the A_s normalization given in Eqs. (B15)–(B16) from the presence of the step.

APPENDIX C: MINIMIZATION

In this Appendix we provide details of the effective χ^2 minimization for the various models presented in Sec. III. In each case, we use the MIGRAD variable metric minimizer from the CERN Minuit2 code [33].

We begin with the smooth Λ CDM cosmology specified by the cosmological parameters $\{\tilde{A}_s, n_s, \theta_A, \Omega_c h^2, \Omega_b h^2, \tau\}$ and 14 foreground parameters defined in the Planck likelihood [34]. We include the Planck low- ℓ spectrum (Commander, $l < 50$), the high- ℓ spectrum (CAMspec, $50 < l < 2500$), and WMAP9 polarization (low-like) likelihoods in our analysis [34,35]. \tilde{A}_s is the effective normalization defined in Eq. (B17); in the absence of a step $\tilde{A}_s = A_s e^{-2\tau}$. In the standard Λ CDM model, the effective number and sum of the masses of neutrinos are held fixed to $N_{\text{eff}} = 3.046$ and $\sum m_\nu = 0.06$ eV, respectively, with the helium fraction $Y_P = 0.2477$. The best fit model is given in Tables I and II and its χ^2 is the baseline from which we quantify any improvements due to the step parameters. We fix the foreground parameters to these values for the following analysis but have spot checked that reoptimization over the foreground parameters does not substantially change our results.

TABLE I. Best fit flat Λ CDM cosmological model without a step with 6 varied parameters (top) and derived parameters (bottom). This model provides the baseline χ^2 for the smooth model but its parameters require adjustment in the presence of a step. \tilde{A}_s is an effective normalization parameter defined in Eq. (B17).

$10^9 \tilde{A}_s$	1.8027
n_s	0.9607
$100\theta_A$	1.04144
$10 \Omega_c h^2$	1.1995
$100 \Omega_b h^2$	2.2039
100τ	8.952
H_0	67.22
$10^9 A_s$	2.1562
D_* (Mpc)	13893.1
χ^2	9802.8

TABLE II. Foreground model. These parameters are jointly minimized with those of Table I in the smooth model and held fixed for the step analysis.

γ^{CIB}	0.538	A_{217}^{PS}	112.4
$r_{143 \times 217}^{\text{PS}}$	0.906	A_{143}^{CIB}	6.18
A_{217}^{CIB}	27.5	c_{100}	1.000580
A_{143}^{tSZ}	6.71	c_{217}	0.9963
$\xi^{\text{tSZ-CIB}}$	0.2	β_1^1	0.55
A_{100}^{PS}	152	A^{KSZ}	3
A_{143}^{PS}	50.8	$r_{143 \times 217}^{\text{CIB}}$	0.365

For the step analysis, the starting point is the canonical $c_s = 1$ potential step where $C_1 = C_3 = 0$. As the oscillatory features from C_2 dominate the fit to the Planck data, the other cases are built from this model. In this case the step is described by $\{C_2, s_s, x_d\}$. While we could directly minimize the χ^2 in this joint cosmological and step parameter space, the efficiency of the search is greatly improved by choosing parameters that are better aligned to the principal axes of the χ^2 surface.

The angular frequency of the oscillation changes with cosmological parameters at fixed s_s . It is thus advantageous to replace s_s with

$$\theta_s = \frac{s_s}{D_*}, \quad (\text{C1})$$

where D_* is the distance to recombination. Note that the oscillations in C_ℓ are then described by sinusoids such as $\sin(2\ell\theta_s)$. Next, due to projection effects, a fixed amplitude C_2 produces an oscillation in C_ℓ that decays as $C_2(\ell\theta_s)^{-1/2}$. In Ref. [2] this scaling was approximately accounted for in the curvature power spectrum description by introducing the amplitude parameter

$$A_c = 3C_2 \left[\sqrt{\frac{s_s}{1 \text{ Gpc}}} \right]^{-1}; \quad (\text{C2})$$

we adopt this convention rather than the more orthogonal angular approach in order to compare with the previous literature. Note that the original version of the Planck collaboration analysis erroneously conflated this parameter with C_2 [5].

Finally, given that the oscillations produce excess broadband power, we use the normalization parameter \tilde{A}_s as defined in Eq. (B17). For the best fit models, this parameter rather than A_s itself is nearly constant. The optimized parameters are therefore $\{\tilde{A}_s, n_s, \theta_A, \Omega_c h^2, \Omega_b h^2, \tau\}$ for the smooth cosmology and $\{A_c, \theta_s, x_d\}$ for the step. The minimum χ^2 potential step model with $c_s = 1$ is given in Table III (GSR1 column) and represents an improvement of $\Delta\chi^2 = -14.0$ over the smooth model of Table I. For reference, we also show here the best fit model at $x_d = 2000$, where the oscillations are undamped all the way to the maximum of $\ell = 2500$ for Planck. Note that most of the improvement due to the step remains. We also repeat the minimization for the GSR0 approximation used in previous treatments for comparison. Note that after adjusting cosmological parameters, steps in either approximation fit equally well but the recovery of such parameters would be biased by using the less accurate GSR0 approximation.

For the arbitrary sound speed warp and potential step models, C_1 and C_3 are set consistently with the step

TABLE III. Best fit potential step model with $c_s = 1$ showing the 9 parameters jointly varied (top) and derived parameters (bottom) using the GSR1 approximation of the main paper versus the less accurate GSR0 approximation. The global minimum is at $x_d = 105$ but $x_d = 2000$ where there is no damping of oscillations for the Planck data gives a comparable fit, albeit with lower a oscillation amplitude A_c . \tilde{A}_s is an effective normalization parameter defined in Eq. (B17) that determines the broadband observed CMB power in the presence of τ and a step.

	GSR1		GSR0	
	105	2000	105	2000
x_d	105	2000	105	2000
$10\theta_s$	2.665	2.667	2.665	2.666
$10A_c$	1.17	0.663	1.11	0.707
$10^9 \tilde{A}_s$	1.8021	1.8024	1.8020	1.8021
n_s	0.9690	0.9608	0.9640	0.9606
$100\theta_A$	1.04140	1.04145	1.04136	1.04140
$10 \Omega_c h^2$	1.2091	1.1995	1.2035	1.1993
$100 \Omega_b h^2$	2.1974	2.2039	2.2053	2.2039
100τ	9.421	9.117	9.361	9.205
H_0	66.82	67.23	67.07	67.22
$10^9 A_s$	2.1669	2.1420	2.1701	2.1565
D_* (Mpc)	13874.7	13893.2	13882.9	13893.9
s_s (Mpc)	3696.9	3704.7	3699.2	3704.5
C_2	0.075	0.043	0.071	0.045
$\Delta\chi^2$	-14.0	-11.4	-13.8	-11.3

amplitude $\{b_T, b_V\}$ and slow-roll parameters after the step $\{c_{sa}, \epsilon_{Ha}\}$ through Eq. (11). These parameters mainly change the power spectrum around $\ell \sim 1/\theta_s$ and hence produce only small changes in the χ^2 due to the limitations of cosmic variance.

We therefore keep the other parameters fixed to the values of the $x_d = 105$ model listed in Table III when considering the additional freedom in these models. Given a fixed C_2 , which fixes the amplitude of the step, this freedom is parametrized by c_{sa} , the sound speed after the step. For warp step, the best fit is given by

$$\begin{aligned} c_{sa} &= 0.70, & C_1 &= -0.70, \\ C_3 &= -0.37, & \Delta\chi^2 &= -15.2 \text{ (warp)}, \end{aligned} \quad (\text{C3})$$

and this corresponds to a $\Delta\chi^2 = -1.2$ improvement over the potential step model at $c_{sa} = 1$. For low sound speed potential step models, the $c_{sa} \rightarrow 0$ limit provides the best fit

$$\begin{aligned} c_{sa} &\rightarrow 0, & C_1 &= 0, \\ C_3 &= -0.03, & \Delta\chi^2 &= -14.1 \text{ (potential)}. \end{aligned} \quad (\text{C4})$$

Given the additional parameter c_{sa} , neither improvement is statistically significant.

- [1] R. Flauger, L. McAllister, E. Pajer, A. Westphal, and G. Xu, *J. Cosmol. Astropart. Phys.* **06** (2010) 009.
- [2] P. Adshead, C. Dvorkin, W. Hu, and E. A. Lim, *Phys. Rev. D* **85**, 023531 (2012).
- [3] H. Peiris, R. Easther, and R. Flauger, *J. Cosmol. Astropart. Phys.* **09** (2013) 018.
- [4] P. D. Meerburg, D. N. Spergel, and B. D. Wandelt, [arXiv:1308.3704](https://arxiv.org/abs/1308.3704).
- [5] P. Ade *et al.* (Planck Collaboration), [arXiv:1303.5082](https://arxiv.org/abs/1303.5082).
- [6] R. Easther and R. Flauger, *J. Cosmol. Astropart. Phys.* **02** (2014) 037.
- [7] M. Benetti, *Phys. Rev. D* **88**, 087302 (2013).
- [8] P. D. Meerburg and D. N. Spergel, [arXiv:1308.3705](https://arxiv.org/abs/1308.3705).
- [9] X. Chen, R. Easther, and E. A. Lim, *J. Cosmol. Astropart. Phys.* **06** (2007) 023.
- [10] E. Silverstein and D. Tong, *Phys. Rev. D* **70**, 103505 (2004).
- [11] J. A. Adams, B. Cresswell, and R. Easther, *Phys. Rev. D* **64**, 123514 (2001).
- [12] R. Bean, X. Chen, G. Hailu, S.-H. Tye, and J. Xu, *J. Cosmol. Astropart. Phys.* **03** (2008) 026.
- [13] V. Miranda, W. Hu, and P. Adshead, *Phys. Rev. D* **86**, 063529 (2012).
- [14] C. Dvorkin and W. Hu, *Phys. Rev. D* **82**, 043513 (2010).
- [15] A. Lewis, A. Challinor, and A. Lasenby, *Astrophys. J.* **538**, 473 (2000).
- [16] C. Howlett, A. Lewis, A. Hall, and A. Challinor, *J. Cosmol. Astropart. Phys.* **04** (2012) 027.
- [17] M. J. Mortonson, C. Dvorkin, H. V. Peiris, and W. Hu, *Phys. Rev. D* **79**, 103519 (2009).
- [18] A. Lewis, A. Challinor, and D. Hanson, *J. Cosmol. Astropart. Phys.* **03** (2011) 018.
- [19] W. Hu, *Astrophys. J.* **557**, L79 (2001).
- [20] D. M. Goldberg and D. N. Spergel, *Phys. Rev. D* **59**, 103002 (1999).
- [21] P. Adshead, W. Hu, C. Dvorkin, and H. V. Peiris, *Phys. Rev. D* **84**, 043519 (2011).
- [22] P. Adshead and W. Hu, *Phys. Rev. D* **85**, 103531 (2012).
- [23] A. Achucarro, J.-O. Gong, G. A. Palma, and S. P. Patil, *Phys. Rev. D* **87**, 121301 (2013).
- [24] P. Adshead, W. Hu, and V. Miranda, *Phys. Rev. D* **88**, 023507 (2013).
- [25] J. Fergusson, M. Liguori, and E. Shellard, *J. Cosmol. Astropart. Phys.* **12** (2012) 032.
- [26] D. Baumann and D. Green, *J. Cosmol. Astropart. Phys.* **09** (2011) 014.
- [27] E. D. Stewart, *Phys. Rev. D* **65**, 103508 (2002).
- [28] W. Hu, *Phys. Rev. D* **84**, 027303 (2011).
- [29] C. Dvorkin and W. Hu, *Phys. Rev. D* **81**, 023518 (2010).
- [30] J. Garriga and V. F. Mukhanov, *Phys. Lett. B* **458**, 219 (1999).
- [31] J. Choe, J.-O. Gong, and E. D. Stewart, *J. Cosmol. Astropart. Phys.* **07** (2004) 012.
- [32] C. Dvorkin and W. Hu, *Phys. Rev. D* **84**, 063515 (2011).
- [33] F. James and M. Roos, *Comput. Phys. Commun.* **10**, 343 (1975).
- [34] P. Ade *et al.* (Planck collaboration), [arXiv:1303.5075](https://arxiv.org/abs/1303.5075).
- [35] C. Bennett, D. Larson, J. Weiland, N. Jarosik, G. Hinshaw *et al.*, *Astrophys. J. Suppl. Ser.* **208**, 20 (2013).