

Exceptional field theory. III. $E_{8(8)}$

Olaf Hohm^{1*} and Henning Samtleben^{2†}

¹*Center for Theoretical Physics, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139, USA*

²*Université de Lyon, Laboratoire de Physique, UMR 5672, CNRS École Normale Supérieure de Lyon 46,
allée d'Italie, F-69364 Lyon cedex 07, France*

(Received 18 July 2014; published 4 September 2014)

We develop exceptional field theory for $E_{8(8)}$, defined on a $(3 + 248)$ -dimensional generalized spacetime with extended coordinates in the adjoint representation of $E_{8(8)}$. The fields transform under $E_{8(8)}$ generalized diffeomorphisms and are subject to covariant section constraints. The bosonic fields include an “internal” dreibein and an $E_{8(8)}$ -valued “zweihundertachtundvierzigbein” (248-bein). Crucially, the theory also features gauge vectors for the $E_{8(8)}$ E bracket governing the generalized diffeomorphism algebra and covariantly constrained gauge vectors for a separate but constrained $E_{8(8)}$ gauge symmetry. The complete bosonic theory, with a novel Chern-Simons term for the gauge vectors, is uniquely determined by gauge invariance under internal and external generalized diffeomorphisms. The theory consistently comprises components of the dual graviton encoded in the 248-bein. Upon picking particular solutions of the constraints the theory reduces to $D = 11$ or type IIB supergravity, for which the dual graviton becomes pure gauge. This resolves the dual graviton problem, as we discuss in detail.

DOI: 10.1103/PhysRevD.90.066002

PACS numbers: 11.25.Yb, 04.50.-h, 04.65.+e

I. INTRODUCTION

In this paper we present the details of the recently announced “exceptional field theory” (EFT) [1] for the group $E_{8(8)}$, complementing the construction for $E_{6(6)}$ and $E_{7(7)}$ given in [2] and [3], respectively. The approach is a generalization of double field theory (DFT) [4–9],¹ with the goal to render the dynamics of the complete $D = 11$ supergravity [11], and that of type IIB [12,13], covariant under the exceptional groups that are known to appear under dimensional reduction [14]. We refer to the Introduction of [2] for a more detailed outline of the general ideas, previous approaches, and extensive references. Here we will mainly present and discuss the novel aspects relevant for the group $E_{8(8)}$ which brings in some distinctive new features as compared to the formulations for the smaller exceptional groups.

The $E_{8(8)}$ EFT is based on a generalized $(3 + 248)$ -dimensional spacetime, with the “external” spacetime coordinates x^μ and “internal” coordinates Y^M in the adjoint representation 248 of $E_{8(8)}$, with dual derivatives ∂_M .² The dependence of all fields on the extended 248 coordinates Y^M is restricted by $E_{8(8)}$ -covariant section constraints [16,17] that project out subrepresentations in the tensor product $248 \otimes 248$,

$$(\mathbb{P}_{1+248+3875})_{MN}{}^{KL} \partial_K \otimes \partial_L = 0. \quad (1.1)$$

As in double field theory, this constraint is meant to hold on any fields, parameters, and their products. This constraint has nontrivial solutions, which break $E_{8(8)}$ to $GL(8)$ or $GL(7) \times SL(2)$, for which the EFT reduces to $D = 11$ supergravity or type IIB, respectively, for appropriate reformulations of these theories, as pioneered in [18,19] for $E_{8(8)}$.

The bosonic field content of the $E_{8(8)}$ EFT is given by

$$\{e_\mu{}^a, \mathcal{V}_M{}^M, A_\mu{}^M, B_{\mu M}\}. \quad (1.2)$$

It incorporates an external frame field (“dreibein”) $e_\mu{}^a$, $\mu = 0, 1, 2$, and an internal generalized frame field (“zweihundertachtundvierzigbein”) $\mathcal{V}_M{}^M$, $M = 1, \dots, 248$, parametrizing the coset space $E_{8(8)}/SO(16)$. From the latter, we may construct the “generalized metric” as $\mathcal{M}_{MN} = (\mathcal{V}\mathcal{V}^T)_{MN}$. Crucially, the theory also requires the presence of generalized gauge connections $A_\mu{}^M$ and $B_{\mu M}$, in order to consistently describe the complete degrees of freedom and dynamics of $D = 11$ supergravity (necessarily including also some of the dual fields). The theory is invariant under gauge symmetries with parameters Λ^M, Σ_M acting as

$$\begin{aligned} \mathbb{L}_{(\Lambda, \Sigma)} V^M &\equiv \Lambda^K \partial_K V^M - 60 \mathbb{P}^M{}_N{}^K{}_L \partial_K \Lambda^L V^N \\ &\quad + \lambda \partial_N \Lambda^N V^M - \Sigma_L f^{LM}{}_N V^N, \end{aligned} \quad (1.3)$$

on a vector V^M of weight λ . The Λ^M transformations generate the generalized diffeomorphisms on the 248-dimensional

*ohohm@mit.edu

†henning.samtleben@ens-lyon.fr

¹See [10] for a review and further references.

²Such generalized spacetimes also appear in the proposal of [15].

space, following the definition for the smaller exceptional groups [16] with \mathbb{P} denoting the projector onto the adjoint representation. The Σ_M gauge symmetry is a new feature of the $E_{8(8)}$ EFT and describes a separate $E_{8(8)}$ gauge symmetry, however, with parameters Σ_M that are “covariantly constrained.” This means that they obey the same algebraic constraints as the derivatives in (1.1), for instance $\mathbb{P}_{MN}{}^{KL}\Sigma_K \otimes \partial_L = 0$, etc. As a result, most of the components vanish after explicitly solving the section constraints, and the $E_{8(8)}$ gauge symmetry is much smaller than is apparent from (1.3) [as it should be, for otherwise all fields encoded in the $E_{8(8)}/SO(16)$ coset space would be pure gauge]. This additional gauge symmetry is necessary for consistency. For instance, the generalized diffeomorphisms in (1.3) with parameter Λ^M do not close into themselves which has been recognized as an obstacle in [16,17]. They do however close in presence of the additional covariantly constrained gauge symmetry that constitutes a separate invariance of the theory. In other words, invariance of an action under generalized diffeomorphisms Λ^M implies its invariance under further Σ_M gauge transformations, as we shall explicitly confirm. This type of gauge structure has first been revealed in the baby example of an $SL(2)$ covariant formulation of four-dimensional Einstein gravity [20].

The constraints on the gauge parameter Σ_M imply that also the associated connection $B_{\mu M}$ is covariantly constrained in the same sense, i.e. it satisfies $\mathbb{P}_{MN}{}^{KL}B_{\mu K} \otimes \partial_L = 0$, etc. Such covariantly constrained compensating gauge fields are a generic feature of the exceptional field theories and show up among the $(D-2)$ -forms (with D counting the number of external dimensions). Therefore in $D=5$, these fields do not even enter the Lagrangian [2], in $D=4$ they appear among the 2-forms with Stückelberg coupling to the Yang-Mills field strengths [3], while in $D=3$ they feature among the vector fields and thus directly affect the algebra of gauge transformations (1.3). In all cases, these constrained gauge fields are related to the appearance of the dual gravitational degrees of freedom as we discuss shortly.

The full $E_{8(8)}$ -covariant action is given by

$$S = \int d^3x d^{248}Y e \left(\hat{R} + e^{-1} \mathcal{L}_{CS} + \frac{1}{240} g^{\mu\nu} D_\mu \mathcal{M}^{MN} D_\nu \mathcal{M}_{MN} - V(\mathcal{M}, g) \right), \quad (1.4)$$

and closely resembles the structure of three-dimensional gauged supergravities [21]. The various terms comprise a (covariantized) Einstein-Hilbert term, a Chern-Simons-type term for the gauge vectors, a covariantized kinetic term for the $E_{8(8)}/SO(16)$ coset fields, and a “potential” V . The Chern-Simons term is a topological term that is needed to ensure the proper on-shell duality relations between

“scalars” and “vectors.” The potential depends only on internal derivatives ∂_M and can be written in a manifestly $E_{8(8)}$ -covariant form as follows:

$$V(\mathcal{M}, g) = -\frac{1}{240} \mathcal{M}^{MN} \partial_M \mathcal{M}^{KL} \partial_N \mathcal{M}_{KL} + \frac{1}{2} \mathcal{M}^{MN} \partial_M \mathcal{M}^{KL} \partial_L \mathcal{M}_{NK} + \frac{1}{7200} f^{NQ}{}_P f^{MS}{}_R \mathcal{M}^{PK} \partial_M \mathcal{M}_{QK} \mathcal{M}^{RL} \partial_N \mathcal{M}_{SL} - \frac{1}{2} g^{-1} \partial_M g \partial_N \mathcal{M}^{MN} - \frac{1}{4} \mathcal{M}^{MN} g^{-1} \partial_M g g^{-1} \partial_N g - \frac{1}{4} \mathcal{M}^{MN} \partial_M g^{\mu\nu} \partial_N g_{\mu\nu}. \quad (1.5)$$

Its form is determined such that it leads to a gauge-invariant action both with respect to the Λ^M and Σ_M gauge transformations of (1.3). Previous attempts to construct an $E_{8(8)}$ -covariant formulation (of truncations of $D=11$ supergravity) missed the third line of (1.5) involving the explicit $E_{8(8)}$ structure constants $f^{MN}{}_K$ [22]. This term is indispensable for gauge invariance of the potential V and for the match with $D=11$ supergravity as we shall explain. All four terms in the action (1.4) are separately gauge invariant with respect to Λ and Σ , but the theory is also invariant under nonmanifest external diffeomorphisms of the x^μ generated by a parameter $\xi^\mu(x, Y)$. This symmetry fixes all the relative coefficients in (1.4), such that this is the unique two-derivative action with all the required symmetries.

We close the Introduction by a discussion of how the above EFT resolves what is often referred to as the “dual graviton problem.” This problem comes about because the $E_{8(8)}$ coset representative \mathcal{M}_{MN} depends on components φ_m , $m=1, \dots, 8$, that in three dimensions are dual to the Kaluza-Klein vectors $A_\mu{}^m$. As the latter originate from components of the $D=11$ metric, this amounts to including in the theory components of a “dual graviton” [23–26] at the full nonlinear level, something that is considered impossible on the grounds of the no-go theorems in [27,28]. In EFT this problem is resolved due to the presence of the extra $E_{8(8)}$ gauge symmetry from (1.3). Solving the section constraints (1.1) such that the theory reduces to $D=11$ supergravity, this covariantly constrained gauge symmetry reduces to a Stückelberg shift symmetry with eight parameters, which can be used to gauge away all the dual graviton components φ_m . Consequently, in the gauge-invariant potential (1.5) all components φ_m drop out upon solving the section constraint, which is necessary for the theory to match $D=11$ supergravity. The same conclusions hold for the solution corresponding to type IIB. Let us finally note that although the dual graviton components φ_m are pure gauge for the $D=11$ and $D=10$ solutions, once we consider strict dimensional reduction to $D=3$, the φ_m are propagating fields among the scalars of the

$E_{8(8)}/\text{SO}(16)$ coset space. Indeed, in this case the Chern-Simons term implies that $B_{\mu m}$ is pure gauge, so the extra gauge symmetry can be fixed by gauging $B_{\mu m}$ away.

This paper is organized as follows. In Sec. II we introduce the $E_{8(8)}$ generalized Lie derivatives and the covariantly constrained $E_{8(8)}$ gauge symmetry. Next, we introduce gauge vectors for these symmetries and define covariant derivatives and field strengths. In Sec. III we present the various terms in the action and prove its gauge invariance under internal and external diffeomorphisms. In particular, we fix all relative coefficients in the action (1.4) by requiring invariance under external diffeomorphisms. Finally, in Sec. IV, we discuss the match with $D = 11$ supergravity and type IIB. Specifically, we discuss how the dual graviton (problem) disappears. We conclude in Sec. V. Some details on the proof of closure of the $E_{8(8)}$ generalized Lie derivatives are presented in the Appendix.

II. $E_{8(8)}$ GAUGE STRUCTURE

In this section we introduce $E_{8(8)}$ -covariant generalized Lie derivatives, which close according to an E bracket, up to a separate covariantly constrained $E_{8(8)}$ gauge symmetry. This means that the $E_{8(8)}$ gauge parameter is subject to the same section constraints as the extended derivatives. Then we introduce gauge fields A_{μ}^M for the E bracket and covariantly constrained gauge fields $B_{\mu M}$ for $E_{8(8)}$.

A. $E_{8(8)}$ generalized lie derivatives

We start by recalling a few generalities of $E_{8(8)}$. Its Lie algebra is 248 dimensional, and the adjoint representation is the smallest fundamental representation. We denote the generators by $(t^M)^N{}_K = -f^{MN}{}_K$, with structure constants $f^{MN}{}_K$, and adjoint indices $M, N = 1, \dots, 248$. The maximal compact subgroup is $\text{SO}(16)$, under which $E_{8(8)}$ decomposes as $\mathbf{248} \rightarrow \mathbf{120} \oplus \mathbf{128}$. There is an invariant symmetric tensor η_{MN} , the Cartan-Killing form, which we normalize by

$$\eta^{MN} = \frac{1}{60} \text{tr}(t^M t^N) = \frac{1}{60} f^{MK}{}_L f^{NL}{}_K, \quad (2.1)$$

and which we freely use to raise and lower adjoint indices. Given this invariant metric, the tensor product of the adjoint with the coadjoint representation is equivalent to $\mathbf{248} \otimes \mathbf{248}$ and decomposes as follows:

$$\mathbf{248} \otimes \mathbf{248} \rightarrow \mathbf{1} \oplus \mathbf{248} \oplus \mathbf{3875} \oplus \mathbf{27000} \oplus \mathbf{30380}. \quad (2.2)$$

In particular, it contains the adjoint representation, and in the following we need the corresponding projector:

$$\begin{aligned} \mathbb{P}^M{}_N{}^K{}_L &= \frac{1}{60} f^M{}_{NP} f^{PK}{}_L \\ &= \frac{1}{30} \delta^M_{(N} \delta^K_{L)} - \frac{7}{30} (\mathbb{P}_{3875})^{MK}{}_{NL} \\ &\quad - \frac{1}{240} \eta^{MK} \eta_{NL} + \frac{1}{120} f^{MK}{}_P f^P{}_{NL}. \end{aligned} \quad (2.3)$$

Here we used Eqs. (2.15) in [29], and the projector onto the 3875 which is given by

$$(\mathbb{P}_{3875})^{MK}{}_{NL} = \frac{1}{7} \delta^M_{(N} \delta^K_{L)} - \frac{1}{56} \eta^{MK} \eta_{NL} - \frac{1}{14} f^P{}_N{}^{(M} f_{PL}{}^{K)}. \quad (2.4)$$

We refer to [19,29] for other useful $E_{8(8)}$ identities.

Let us now discuss the generalized spacetime and geometry based on $E_{8(8)}$. We introduce 248 coordinates Y^M in the adjoint representation, but we subject all functions (i.e. including all fields and gauge parameters and all their products) to the covariant section constraints (1.1). These are necessary in order for the symmetries of the theory to close into an algebra. These symmetries comprise generalized diffeomorphisms on the 248-dimensional space, together with a covariantly constrained $E_{8(8)}$ gauge symmetry. Specifically, denoting by Λ^M and Σ_M the parameters for generalized diffeomorphisms and constrained $E_{8(8)}$, respectively, we define the generalized Lie derivative on a vector by

$$\begin{aligned} \delta V^M &= \mathbb{L}_{(\Lambda, \Sigma)} V^M \\ &\equiv \Lambda^K \partial_K V^M - 60 \mathbb{P}^M{}_N{}^K{}_L \partial_K \Lambda^L V^N + \lambda(V) \partial_N \Lambda^N V^M \\ &\quad - \Sigma_L f^{LM}{}_N V^N. \end{aligned} \quad (2.5)$$

Analogously, one may define the generalized Lie derivative acting on tensors with an arbitrary number of adjoint $E_{8(8)}$ indices. The second line of (2.5) defines the generalized Lie derivative with respect to Λ^M , in accordance with the definition for the smaller exceptional groups [16,17], where we also allowed for a general density weight λ . The third line is a novel feature of the $E_{8(8)}$ EFT. It defines the covariantly constrained $E_{8(8)}$ action, i.e. describes an $E_{8(8)}$ rotation with a parameter Σ_M which itself satisfies the same algebraic conditions (1.1) as the partial derivatives. Concretely, we require that

$$\begin{aligned} (\mathbb{P}_{1+248+3875})_{MN}{}^{KL} C_K \otimes C'_L &= 0, \\ \text{for } C_M, C'_M &\in \{\partial_M, B_{\mu M}, \Sigma_M\}, \end{aligned} \quad (2.6)$$

where $B_{\mu M}$ denotes the gauge connection associated to the Σ_M symmetry of (2.5). This means that for any expression containing two objects, C_M and C'_N , from the list above, the part in the tensor product that is projected out by this

constraint can be consistently set to zero. Explicitly, we have for the individual irreducible representations,

$$\begin{aligned} \eta^{MN} C_M \otimes C'_N = 0, \quad f^{MKN} C_N \otimes C'_K = 0, \\ (\mathbb{P}_{3875})_{MN}{}^{KL} C_K \otimes C'_L = 0. \end{aligned} \quad (2.7)$$

This implies in particular $\eta^{MN} \partial_M \partial_N A = \partial^M \partial_M A = 0$, but also $\partial^M A \partial_M B = 0$, for arbitrary functions A, B , and relations like $f^{MKN} B_{\mu N} \partial_K A = 0$ involving the covariantly constrained gauge field $B_{\mu M}$. These relations imply that for any solution of the section constraint only a subset of coordinates among the Y^M survives, while also only the ‘‘corresponding’’ components of $B_{\mu M}$ are present, as we will explain in more detail below.

Before determining the gauge algebra satisfied by (2.5) we briefly discuss that the above gauge transformations (2.5) possess ‘‘trivial’’ gauge parameters. For these parameters the action of the associated generalized Lie derivative on any field vanishes by virtue of the section constraints (2.7). The following parameters are trivial in this sense,

$$\begin{aligned} \Lambda^M \equiv \eta^{MN} \Omega_N, \quad \Omega_N \text{ covariantly constrained à la (2.6),} \\ \Lambda^M \equiv (\mathbb{P}_{3875})^{MK}{}_{NL} \partial_K \chi^{NL}. \end{aligned} \quad (2.8)$$

Here, in the first line, Ω_N is covariantly constrained in the sense that it satisfies the same constraints as the C_N in (2.6), (2.7). E.g. choosing $\Omega_N = \partial_N \chi$ we infer that $\Lambda^M = \partial^M \chi$ is a trivial parameter, in analogy to DFT. For the first parameter in (2.8) it is straightforward to see with (2.4) and the constraints (2.7) that the generalized Lie derivative (2.5) is zero on fields. As an illustration for the use of constraints, we prove explicitly the triviality of the second parameter in (2.8). We first note that in this case the transport term and density term (i.e. the first and third terms) in (2.5) immediately vanish as a consequence of the third constraint in (2.7). Thus, the action of the generalized Lie derivative reads

$$\begin{aligned} \mathbb{L}_\Lambda V^M &= -60 \mathbb{P}_{3875}^M{}_N{}^{(P} \mathbb{Q}^{(P} (\mathbb{P}_{3875})^{R)Q}{}_{ST} \partial_P \partial_R \chi^{ST} V^N \\ &= -f^M{}_{NX} (f^{X(P} \mathbb{Q}^{(P} (\mathbb{P}_{3875})^{R)Q}{}_{ST}) \partial_P \partial_R \chi^{ST} V^N. \end{aligned} \quad (2.9)$$

Next, we use that \mathbb{P}_{3875} is an invariant tensor under the adjoint action of $E_{8(8)}$, as is manifest from its definition (2.4). This means

$$f^{X(P} \mathbb{Q}^{(P} (\mathbb{P}_{3875})^{R)Q}{}_{ST} - f^{XQ} (\mathbb{P}_{3875})^{PR}{}_{T)Q} = 0. \quad (2.10)$$

Thus, we can replace the structure in (2.9) by the second term in here. Being contracted with $\partial_P \partial_R$ it then follows from the third constraint in (2.7) that this vanishes, completing the proof that the associated generalized Lie derivative acts trivially.

Next, we discuss a novel phenomenon for the $E_{8(8)}$ case: there are combinations of parameters Λ and Σ whose combined action is trivial on all the fields. Specifically, the generalized Lie derivative (2.5) with parameters

$$\begin{aligned} \Lambda^M &= f^{MN}{}_K \Omega_N{}^K, \\ \Omega_N{}^K &\text{ covariantly constrained in first index,} \\ \Sigma_M &= \partial_M \Omega_N{}^N + \partial_N \Omega_M{}^N, \end{aligned} \quad (2.11)$$

acts trivially for a general tensor $\Omega_N{}^K$ that is covariantly constrained in the first index in the sense of (2.6), (2.7). An example is given by $\Omega_M{}^N = \partial_M \chi^N$ with arbitrary χ^N , so we conclude as a special case of (2.11) that

$$\Lambda^M \equiv f^{MN}{}_K \partial_N \chi^K, \quad \Sigma_M \equiv 2 \partial_M \partial_N \chi^N \quad (2.12)$$

has trivial action on all the fields. In order to verify the triviality of (2.11) let us first prove the following useful lemma:

$$f^P{}_M{}^K f_{PN}{}^L C_K \otimes C'_L = C_M \otimes C'_N + C_N \otimes C'_M, \quad (2.13)$$

for any covariantly constrained objects C_M, C'_M . To prove this we compute

$$\begin{aligned} f^P{}_M{}^K f_{PN}{}^L C_K \otimes C'_L &= (f^P{}_M{}^{[K} f_{PN}{}^{L]} + f^P{}_M{}^{(K} f_{PN}{}^{L)}) C_K \otimes C'_L \\ &= \left(-\frac{1}{2} f^{PKL} f_{PNM} + 2 \delta_{(M}^K \delta_{N)}^L \right) \\ &\quad - \frac{1}{4} \eta_{MN} \eta^{KL} - 14 (\mathbb{P}_{3875})^{KL}{}_{MN} \Big) C_K \otimes C'_L \\ &= C_M \otimes C'_N + C_N \otimes C'_M. \end{aligned} \quad (2.14)$$

In the second line we used the Jacobi identity and rewrote the symmetrized ff term in terms of the 3875 projector (2.4). In the final step we used the section constraints (2.7). This completes the proof of (2.13). It is now straightforward to verify the triviality of (2.11). First, the transport and density terms vanish immediately as a consequence of the second constraint in (2.7). The remaining projector term, in the first form of the projector in (2.4), can then be simplified by (2.13) to show that this cancels the Σ terms from (2.11). Another immediate consequence of (2.13) is that for a generalized vector Ω_M (of weight 0) that is covariantly constrained, the generalized Lie derivative reduces to

$$\delta_\Lambda \Omega_M = \Lambda^N \partial_N \Omega_M + \partial_N \Lambda^N \Omega_M + \partial_M \Lambda^N \Omega_N, \quad (2.15)$$

which will be used below.

We close this section by discussing closure of the gauge transformations. In contrast to the analogous structures for $E_{n(n)}$ with $n \leq 7$, the generalized Lie derivatives do not

close by themselves, but only up to (constrained) local $E_{8(8)}$ gauge transformations. Specifically, one finds closure

$$\begin{aligned} [\delta_{(\Lambda_1, \Sigma_1)}, \delta_{(\Lambda_2, \Sigma_2)}] &= \delta_{[(\Lambda_2, \Sigma_2), (\Lambda_1, \Sigma_1)]_E}, \\ [(\Lambda_2, \Sigma_2), (\Lambda_1, \Sigma_1)]_E &\equiv (\Lambda_{12}, \Sigma_{12}), \end{aligned} \quad (2.16)$$

with the effective parameters

$$\begin{aligned} \Lambda_{12}^M &\equiv 2\Lambda_{[2}^N \partial_N \Lambda_{1]}^M - 14(\mathbb{P}_{3875})^{MK}{}_{NL} \Lambda_{[2}^N \partial_K \Lambda_{1]}^L \\ &\quad - \frac{1}{4} \eta^{MK} \eta_{NL} \Lambda_{[2}^N \partial_K \Lambda_{1]}^L + \frac{1}{4} f^{MN}{}_P \partial_N (f^P{}_{KL} \Lambda_2^K \Lambda_1^L), \\ \Sigma_{12M} &\equiv -2\Sigma_{[2M} \partial_N \Lambda_{1]}^N + 2\Lambda_{[2}^N \partial_N \Sigma_{1]M} \\ &\quad - 2\Sigma_{[2}^N \partial_M \Lambda_{1]N} + f^N{}_{KL} \Lambda_{[2}^K \partial_M \partial_N \Lambda_{1]}^L. \end{aligned} \quad (2.17)$$

Note that here is an ambiguity in the form of the effective gauge parameters, because they can be redefined by trivial gauge parameters, (2.8) or (2.11), without spoiling closure. In particular, the term in the second line of Λ_{12} could have been dropped, using (2.11), at the cost of extra terms in Σ_{12} . The form here has been chosen for later convenience. We stress again that closure only holds because of the separate (covariantly constrained) $E_{8(8)}$ gauge symmetry. Note that this is a rather nontrivial statement, because the effective Σ_{12} parameter needs to be compatible with the covariant section constraints (2.7). The compatibility is manifest from the form in (2.17), because in each term the free index M is carried by a constrained object, Σ_M or ∂_M . As this interplay between generalized diffeomorphisms and a separate but constrained gauge symmetry is somewhat unconventional we prove gauge closure (2.16), (2.17) explicitly in the Appendix. We finally note that the gauge

algebra of Σ transformations with themselves is Abelian, for the effective parameter $\Sigma_{12}^M = f^{MNK} \Sigma_{2N} \Sigma_{1K}$ is actually zero by the section constraints (2.7).

B. Gauge fields for $E_{8(8)}$ E-bracket

We now introduce gauge fields for the local symmetries generated by Λ^M and Σ_M . Specifically, these parameters are functions of x^μ and Y^M , requiring in particular covariant derivatives D_μ for the external coordinates. Denoting the gauge fields for the Λ^M symmetries by $A_\mu{}^M$ and those for the Σ_M symmetries by $B_{\mu M}$, the covariant derivative on any tensor with an arbitrary number of adjoint $E_{8(8)}$ indices is defined by

$$D_\mu \equiv \partial_\mu - \mathbb{L}_{(A_\mu, B_\mu)}, \quad (2.18)$$

where the generalized Lie derivative \mathbb{L} acts according to the representation the tensor field lives in. For instance, using (2.5) one finds its action on a vector of zero weight

$$\begin{aligned} D_\mu V^M &= \partial_\mu V^M - A_\mu{}^K \partial_K V^M + 60 \mathbb{P}^M{}_{NKL} \partial_K A_\mu{}^L V^N \\ &\quad + B_\mu{}^L f^M{}_{NL} V^N. \end{aligned} \quad (2.19)$$

The transformation rules for A and B are determined by the requirement that the covariant derivatives (2.18) transform covariantly. In general, their gauge transformations can be computed from

$$(\delta_{(\Lambda, \Sigma)} A, \delta_{(\Lambda, \Sigma)} B) \equiv (\partial \Lambda, \partial \Sigma) + [(\Lambda, \Sigma), (A, B)]_E, \quad (2.20)$$

with the E bracket defined by (2.16). Using (2.17) one computes for the components

$$\begin{aligned} \delta_{(\Lambda, \Sigma)} A_\mu{}^M &= D_\mu \Lambda^M - \partial_N A_\mu{}^N \Lambda^M + 7 \mathbb{P}_{3875}{}^{MN}{}_{KL} (\Lambda^K \partial_N A_\mu{}^L + A_\mu{}^K \partial_N \Lambda^L) \\ &\quad - B_\mu{}^L f^M{}_{NL} \Lambda^N + \frac{1}{8} \eta^{MN} \eta_{KL} (\Lambda^K \partial_N A_\mu{}^L + A_\mu{}^K \partial_N \Lambda^L) + \frac{1}{4} f^{MN}{}_P f^P{}_{KL} (\partial_N \Lambda^K A_\mu{}^L - \Lambda^K \partial_N A_\mu{}^L), \\ \delta_{(\Lambda, \Sigma)} B_{\mu M} &= D_\mu \Sigma_M + \partial_N (B_{\mu M} \Lambda^N) + B_\mu{}^N \partial_M \Lambda_N + \frac{1}{2} f^N{}_{KL} (\Lambda^K \partial_M \partial_N A_\mu{}^L - A_\mu{}^K \partial_M \partial_N \Lambda^L). \end{aligned} \quad (2.21)$$

Note in particular, that the gauge field A and its parameter Λ carries weight 1 [and we have explicitly spelled out the weight term in (2.21)], whereas B and Σ carry weight 0.

Because of the existence of trivial gauge parameters, cf. (2.8) and (2.11) discussed in the previous subsection, the gauge transformations of A and B are determined from the covariance of (2.18) only up to redefinitions by trivial parameters. Specifically, the covariant derivatives (2.18) are invariant under the following vector shift transformations

$$\begin{aligned} \delta A_\mu{}^M &= \partial^M \Xi_\mu + (\mathbb{P}_{3875})^{MK}{}_{NL} \partial_K \Xi_\mu{}^{NL} + f^{MN}{}_K \Xi_{\mu N}{}^K, \\ \delta B_{\mu M} &= \partial_M \Xi_{\mu N} + \partial_N \Xi_{\mu M}, \end{aligned} \quad (2.22)$$

with $\Xi_{\mu M}{}^N$ constrained in the first index. We now redefine the gauge transformations of A and B by adding trivial gauge transformations of this form, with parameters

$$\begin{aligned} \Xi_{\mu 3875}{}^{KL} &= -7(\mathbb{P}_{3875})^{KL}{}_{PQ} A_\mu{}^P \Lambda^Q, \\ \Xi_\mu &= -\frac{1}{8} A_\mu{}^K \Lambda_K, \\ \Xi_{\mu N}{}^K &= -B_{\mu N} \Lambda^K + \frac{1}{4} f^K{}_{PQ} \Lambda^P \partial_N A_\mu{}^Q \\ &\quad - \frac{1}{4} f^K{}_{PQ} \partial_N \Lambda^P A_\mu{}^Q. \end{aligned} \quad (2.23)$$

The gauge transformations (2.21) then take the more compact form

$$\begin{aligned}\delta A_\mu{}^M &= D_\mu^{(1)} \Lambda^M, \\ \delta B_{\mu M} &= D_\mu^{(0)} \Sigma_M - \Lambda^N \partial_M B_{\mu N} + f^N{}_{KL} \Lambda^K \partial_M \partial_N A_\mu{}^L, \end{aligned} \quad (2.24)$$

where we have indicated the respective weights by the superscripts $D_\mu^{(\lambda)}$. This is the final form of the gauge transformations that we use in the following.

Let us now turn to the definition of gauge covariant curvatures or field strengths. Part of these curvatures can be read off from the commutator of covariant derivatives,

$$[D_\mu, D_\nu] V^M = -\mathbb{L}_{(F_{\mu\nu}, G_{\mu\nu})} V^M. \quad (2.25)$$

More precisely, this determines the field strengths up to trivial terms that drop out of the generalized Lie derivatives, for which we find

$$\begin{aligned} F_{\mu\nu}{}^M &= 2\partial_{[\mu} A_{\nu]}{}^M - 2A_{[\mu}{}^N \partial_N A_{\nu]}{}^M \\ &\quad + 14(\mathbb{P}_{3875})^{MN}{}_{KL} A_{[\mu}{}^K \partial_N A_{\nu]}{}^L + \frac{1}{4} A_{[\mu}{}^N \partial^M A_{\nu]}{}^N \\ &\quad - \frac{1}{2} f^{MN}{}_{PQ} f^P{}_{KL} A_{[\mu}{}^K \partial_N A_{\nu]}{}^L, \\ G_{\mu\nu M} &= 2D_{[\mu} B_{\nu]M} - f^N{}_{KL} A_{[\mu}{}^K \partial_M \partial_N A_{\nu]}{}^L. \end{aligned} \quad (2.26)$$

These field strengths do not transform covariantly, but the failure of covariance is of a trivial form that can be compensated by adding 2-form couplings and assigning to them appropriate gauge transformations in the general spirit of the p -form tensor hierarchy [30]. We thus introduce the fully covariant curvatures

$$\begin{aligned} \mathcal{F}_{\mu\nu}{}^M &\equiv F_{\mu\nu}{}^M + 14(\mathbb{P}_{3875})^{MN}{}_{KL} \partial_N C_{\mu\nu}{}^{KL} \\ &\quad + \frac{1}{4} \partial^M C_{\mu\nu} + 2f^{MN}{}_{K} C_{\mu\nu N}{}^K, \\ \mathcal{G}_{\mu\nu M} &= G_{\mu\nu M} + 2\partial_N C_{\mu\nu M}{}^N + 2\partial_M C_{\mu\nu N}{}^N, \end{aligned} \quad (2.27)$$

with 2-form fields $C_{\mu\nu}{}^{KL}$, $C_{\mu\nu}$, and $C_{\mu\nu M}{}^N$, where as in (2.11) the 2-form $C_{\mu\nu M}{}^N$ is covariantly constrained in the first index. The general variation of these curvatures takes a covariant form,

$$\begin{aligned} \delta \mathcal{F}_{\mu\nu}{}^M &= 2D_{[\mu}^{(1)} \delta A_{\nu]}{}^M + 14(\mathbb{P}_{3875})^{MN}{}_{KL} \partial_N \Delta C_{\mu\nu}{}^{KL} \\ &\quad + \frac{1}{4} \partial^M \Delta C_{\mu\nu} + 2f^{MN}{}_{K} \Delta C_{\mu\nu N}{}^K, \\ \delta \mathcal{G}_{\mu\nu M} &= 2D_{[\mu}^{(0)} \delta B_{\nu]M} - 2\partial_M B_{[\mu}{}^N \delta A_{\nu]}{}^N \\ &\quad - 2f^N{}_{KL} \delta A_{[\mu}{}^K \partial_M \partial_N A_{\nu]}{}^L + 2\partial_N \Delta C_{\mu\nu M}{}^N \\ &\quad + 2\partial_M \Delta C_{\mu\nu N}{}^N, \end{aligned} \quad (2.28)$$

where we defined the covariant variations

$$\begin{aligned} \Delta C_{\mu\nu}{}^{KL} &\equiv \delta C_{\mu\nu}{}^{KL} + A_{[\mu}{}^K \delta A_{\nu]}{}^L, \\ \Delta C_{\mu\nu} &\equiv \delta C_{\mu\nu} + A_{[\mu}{}^K \delta A_{\nu]}{}^K, \\ \Delta C_{\mu\nu N}{}^K &\equiv \delta C_{\mu\nu N}{}^K + B_{[\mu N} \delta A_{\nu]}{}^K - \frac{1}{4} f^K{}_{PQ} (A_{[\mu}{}^P \partial_N \delta A_{\nu]}{}^Q \\ &\quad - \partial_N A_{[\mu}{}^P \delta A_{\nu]}{}^Q). \end{aligned} \quad (2.29)$$

We stress that although we had to introduce the additional 2-forms in order to define gauge covariant curvatures, all of them will eventually drop out from the action and the transformation rules. They can be viewed as a convenient tool that allows us to define the Lagrangian in a rather compact form in terms of manifestly covariant quantities whereas we could also have defined the Lagrangian directly in terms of the original fields and confirmed its gauge invariance by an explicit computation. The 2-forms $C_{\mu\nu}$ and $C_{\mu\nu}{}^{KL}$ already show up in the dimensionally reduced theory upon extending on-shell the supersymmetry algebra and first order duality equations beyond the fields present in the Lagrangian [30].

We now specialize to the transformation of the curvatures under Λ and Σ gauge transformations (2.21). The field strength $\mathcal{F}_{\mu\nu}{}^M$ transforms covariantly in that

$$\delta_{\Lambda, \Sigma} \mathcal{F}_{\mu\nu}{}^M = \mathbb{L}_{(\Lambda, \Sigma)} \mathcal{F}_{\mu\nu}{}^M, \quad (2.30)$$

with weight $\lambda = 1$, provided the 2-forms $C_{\mu\nu}$ transform as

$$\begin{aligned} \Delta C_{\mu\nu}{}^{KL} &= \mathcal{F}_{\mu\nu}{}^{(K} \Lambda^{L)}, \\ \Delta C_{\mu\nu} &= \mathcal{F}_{\mu\nu}{}^M \Lambda_M, \\ \Delta C_{\mu\nu N}{}^K &= \frac{1}{4} f^K{}_{PQ} (\partial_N \mathcal{F}_{\mu\nu}{}^P \Lambda^Q - \partial_N \Lambda^Q \mathcal{F}_{\mu\nu}{}^P) \\ &\quad + \frac{1}{2} \mathcal{G}_{\mu\nu N} \Lambda^K + \frac{1}{2} \Sigma_N \mathcal{F}_{\mu\nu}{}^K. \end{aligned} \quad (2.31)$$

On the other hand, the field strength $\mathcal{G}_{\mu\nu M}$ transforms as

$$\begin{aligned} \delta_{\Lambda, \Sigma} \mathcal{G}_{\mu\nu M} &= \mathbb{L}_{(\Lambda, \Sigma)} \mathcal{G}_{\mu\nu M} - f^N{}_{KL} \mathcal{F}_{\mu\nu}{}^K \partial_M \partial_N \Lambda^L \\ &\quad + \partial_M \Sigma_N \mathcal{F}_{\mu\nu}{}^N, \end{aligned} \quad (2.32)$$

where the generalized Lie derivative acts on a tensor of weight 0. These turn out to be the proper transformation rules in order to define a gauge-invariant Chern-Simons term below. To this end we will furthermore derive a set of generalized Bianchi identities (3.14) satisfied by the curvatures $\mathcal{F}_{\mu\nu}{}^M$ and $\mathcal{G}_{\mu\nu M}$.

III. THE ACTION

With the structures set up in the previous section we are now in position to define the various terms in the action (1.4)

$$S = \int d^3x d^{248}Y (\mathcal{L}_{\text{EH}} + \mathcal{L}_{\text{CS}} + \mathcal{L}_{\text{kin}} - eV(\mathcal{M}, g)). \quad (3.1)$$

We describe them one by one. We then verify that the action is invariant under generalized internal and properly defined external diffeomorphisms, which in turn fixes all the relative coupling constants.

A. Einstein-Hilbert and kinetic term

As in [1,31], the Einstein-Hilbert term in (3.1) reads

$$\mathcal{L}_{\text{EH}} = e\hat{R} \equiv e g^{\mu\nu} \hat{R}_{\mu\nu}, \quad (3.2)$$

and is constructed from contraction of the improved Riemann tensor

$$\hat{R}_{\mu\nu}{}^{ab} \equiv R_{\mu\nu}{}^{ab}[\omega] + \mathcal{F}_{\mu\nu}{}^M e^{\rho[a} \partial_M e_{\rho}{}^{b]}, \quad (3.3)$$

where $R_{\mu\nu}{}^{ab}[\omega]$ denotes the covariantized curvature of the spin connection $\omega_{\mu}{}^{ab}$, which in turn is defined by the covariantized vanishing torsion condition

$$0 = D_{[\mu} e_{\nu]}{}^a \equiv \partial_{[\mu} e_{\nu]}{}^a - A_{[\mu}{}^K \partial_K e_{\nu]}{}^a - \partial_K A_{[\mu}{}^K e_{\nu]}{}^a + \omega_{[\mu}{}^{ab} e_{\nu]b}. \quad (3.4)$$

In particular, the dreibein $e_{\mu}{}^a$ is an $E_{8(8)}$ scalar density of weight $\lambda = 1$. Note from the second form in (3.2) that with this weight the Einstein-Hilbert term has a total weight of 1, as needed for local Λ^M gauge invariance. The second term in (3.3) ensures covariance of the Riemann tensor under local Lorentz transformations. As a result, the Einstein-Hilbert term \mathcal{L}_{EH} is invariant under local Lorentz transformations and internal generalized diffeomorphisms. We note that the term is also invariant under the vector shift symmetries (2.22), notably all 2-form contributions in $\mathcal{F}_{\mu\nu}{}^M$ drop out from (3.3).

The matter sector of the theory comprises 128 scalar fields which as in the three-dimensional maximal theory [32] parametrize the coset space $E_{8(8)}/\text{SO}(16)$. In terms of the symmetric group-valued 248×248 matrix \mathcal{M}_{MN} (and its inverse \mathcal{M}^{MN}), the kinetic term in (3.1) takes the form

$$\mathcal{L}_{\text{kin}} = \frac{1}{240} e g^{\mu\nu} D_{\mu} \mathcal{M}_{MN} D_{\nu} \mathcal{M}^{MN} = -\frac{1}{4} e g^{\mu\nu} j_{\mu}{}^M j_{\nu M}, \quad (3.5)$$

in terms of the current $j_{\mu}{}^M$ defined by

$$\begin{aligned} \mathcal{M}^{KN} D_{\mu} \mathcal{M}_{NL} &\equiv j_{\mu}{}^N f_{NL}{}^K, & \text{and satisfying} \\ \mathcal{M}_{MN} j_{\mu}{}^N &= \eta_{MN} j_{\mu}{}^N. \end{aligned} \quad (3.6)$$

All derivatives D_{μ} here are covariantized with respect to generalized internal diffeomorphisms according to (2.18), with the matrix \mathcal{M}_{MN} carrying weight $\lambda = 0$. The second

equation in (3.6) can be verified with (2.1) and the relation,³

$$\mathcal{M}^{PM} \mathcal{M}^{QN} f_{PQ}{}^K = -f^{MN}{}_L \mathcal{M}^{LK}. \quad (3.7)$$

B. Chern-Simons term

The vector fields $A_{\mu}{}^M$ and $B_{\mu M}$ do not carry propagating degrees of freedom, but describe on-shell duals to the scalar fields. Consequently their dynamics in (3.1) is not described by a Yang-Mills coupling but rather by a topological Chern-Simons term which is explicitly given by

$$\begin{aligned} \mathcal{L}_{\text{CS}} = 2\kappa \epsilon^{\mu\nu\rho} &\left(F_{\mu\nu}{}^M B_{\rho M} - f_{KL}{}^N \partial_{\mu} A_{\nu}{}^K \partial_N A_{\rho}{}^L \right. \\ &- \frac{2}{3} f^N{}_{KL} \partial_M \partial_N A_{\mu}{}^K A_{\nu}{}^M A_{\rho}{}^L \\ &\left. - \frac{1}{3} f_{MKL} f^{KP}{}_Q f^{LR}{}_S A_{\mu}{}^M \partial_P A_{\nu}{}^Q \partial_R A_{\rho}{}^S \right), \end{aligned} \quad (3.8)$$

with coupling constant κ that we will determine below. The structure and covariance of the Chern-Simons term become more transparent by calculating its general variation which is given by

$$\begin{aligned} \delta \mathcal{L}_{\text{CS}} &= 2\kappa \epsilon^{\mu\nu\rho} (F_{\mu\nu}{}^M \delta B_{\rho M} + (G_{\mu\nu M} - f_{MN}{}^K \partial_K F_{\mu\nu}{}^N) \delta A_{\rho}{}^M) \\ &= 2\kappa \epsilon^{\mu\nu\rho} (\mathcal{F}_{\mu\nu}{}^M \delta B_{\rho M} + (\mathcal{G}_{\mu\nu M} - f_{MN}{}^K \partial_K \mathcal{F}_{\mu\nu}{}^N) \delta A_{\rho}{}^M). \end{aligned} \quad (3.9)$$

Indeed it follows directly with the section constraints (2.8) and (2.11) that all extra 2-form contributions proportional to $C_{\mu\nu}$ from (2.28) cancel in the second line of (3.9), such that the variation may be expressed entirely in terms of the covariant quantities. Similarly, one confirms with (3.9) that the Chern-Simons term is invariant under the vector shift transformations (2.22). With a little more calculation we may furthermore verify invariance of the Chern-Simons term under generalized internal diffeomorphisms that act as gauge transformations (2.24) on the vector fields. Specifically, after partial integration, the variation (2.24) yields

$$\begin{aligned} \delta \mathcal{L}_{\text{CS}} &= 2\kappa \epsilon^{\mu\nu\rho} \Lambda^K (\mathcal{F}_{\mu\nu}{}^M (-\partial_M B_{\rho K} + f^N{}_{KL} \partial_M \partial_N A_{\rho}{}^L) \\ &- D_{\rho}^{(0)} (\mathcal{G}_{\mu\nu K} - f_{KM}{}^N \partial_N \mathcal{F}_{\mu\nu}{}^M)) \\ &- 2\kappa \epsilon^{\mu\nu\rho} \Sigma_M D_{\rho}^{(1)} \mathcal{F}_{\mu\nu}{}^M. \end{aligned} \quad (3.10)$$

The vanishing of the rhs of this variation corresponds to establishing some generalized Bianchi identities for the curvatures (2.28). This is most conveniently achieved by evaluating three covariant derivatives $\epsilon^{\mu\nu\rho} D_{\mu} D_{\nu} D_{\rho} V^M$ on a vector V^M of weight 0, from which we deduce the identity

³Note the sign, which is due to the fact that unlike η_{MN} the matrix \mathcal{M}_{MN} is not a group invariant tensor, but commutes with the involution which defines the maximal compact subgroup $\text{SO}(16) \subset E_{8(8)}$.

$$\varepsilon^{\mu\nu\rho}\mathbb{L}_{(\mathcal{F}_{\mu\nu},\mathcal{G}_{\mu\nu})}D_\rho V^M = \varepsilon^{\mu\nu\rho}D_\mu(\mathbb{L}_{(\mathcal{F}_{\nu\rho},\mathcal{G}_{\nu\rho})}V^M). \quad (3.11)$$

Its rhs takes the explicit form

$$\varepsilon^{\mu\nu\rho}D_\rho(\mathbb{L}_{(\mathcal{F}_{\mu\nu},\mathcal{G}_{\mu\nu})}V^M) = \varepsilon^{\mu\nu\rho}D_\rho(\mathcal{F}_{\mu\nu}{}^N\partial_N V^M - (\mathcal{G}_{\mu\nu L} - f_{LP}{}^K\partial_K\mathcal{F}_{\mu\nu}{}^P)f^{LM}{}_N V^N),$$

and upon using that

$$D_\rho\partial_N V^M = \partial_N D_\rho V^M - f^{LM}{}_P V^P(\partial_N B_{\rho L} - f_{LQ}{}^K\partial_N\partial_K A_\rho{}^Q), \quad (3.12)$$

for a vector V^M of weight 0, the rhs of (3.11) may be further rewritten as

$$\begin{aligned} \varepsilon^{\mu\nu\rho}D_\rho(\mathbb{L}_{(\mathcal{F}_{\mu\nu},\mathcal{G}_{\mu\nu})}V^M) &= \varepsilon^{\mu\nu\rho}(\mathcal{F}_{\mu\nu}{}^N\partial_N D_\rho V^M - (\mathcal{G}_{\mu\nu L} - f_{LP}{}^K\partial_K\mathcal{F}_{\mu\nu}{}^P)f^{LM}{}_N D_\rho V^N) \\ &\quad + \varepsilon^{\mu\nu\rho}(D_\rho\mathcal{F}_{\mu\nu}{}^N\partial_N V^M - D_\rho(\mathcal{G}_{\mu\nu L} - f_{LP}{}^K\partial_K\mathcal{F}_{\mu\nu}{}^P)f^{LM}{}_N V^N) \\ &\quad - \varepsilon^{\mu\nu\rho}f^{LM}{}_N V^N\mathcal{F}_{\mu\nu}{}^P(\partial_P B_{\rho L} - f_{LQ}{}^K\partial_P\partial_K A_\rho{}^Q). \end{aligned} \quad (3.13)$$

Now the first line in (3.13) reproduces the lhs of (3.11), such that together we obtain the generalized Bianchi identities

$$\begin{aligned} 0 &= \varepsilon^{\mu\nu\rho}D_\rho^{(1)}\mathcal{F}_{\mu\nu}{}^N \otimes \partial_N, \\ 0 &= \varepsilon^{\mu\nu\rho}(D_\rho^{(0)}(\mathcal{G}_{\mu\nu K} - f_{KM}{}^N\partial_N\mathcal{F}_{\mu\nu}{}^M) + \mathcal{F}_{\mu\nu}{}^M(\partial_M B_{\rho K} - f^N{}_{KL}\partial_M\partial_N A_\rho{}^L)). \end{aligned} \quad (3.14)$$

These are sufficient to show that (3.10) vanishes, confirming that the Chern-Simons term is invariant under generalized internal diffeomorphisms. Let us finally note that a more compact presentation of the Chern-Simons term (3.8) can be given as the boundary contribution of a gauge-invariant exact form in four dimensions as

$$S_{\text{CS}} \propto \int_{\Sigma_4} d^4x \int d^{248}Y \left(\mathcal{F}^M \wedge \mathcal{G}_M - \frac{1}{2} f_{MN}{}^K \mathcal{F}^M \wedge \partial_K \mathcal{F}^N \right), \quad (3.15)$$

where again all 2-form contributions $C_{\mu\nu}$ can be checked to drop out from the action. Gauge invariance of (3.15) follows from the transformation behavior of the field strengths under gauge transformations

$$\begin{aligned} \delta_{\Lambda,\Sigma}\mathcal{F}^M &= \mathbb{L}_{(\Lambda,\Sigma)}\mathcal{F}^M, \\ \delta_{\Lambda,\Sigma}(\mathcal{F}^M f_{MN}{}^K \partial_N \mathcal{F}^N) &= \mathbb{L}_{(\Lambda,\Sigma)}(\mathcal{F}^M f_{MN}{}^K \partial_N \mathcal{F}^N) + 2\mathcal{F}^M \partial_M \partial_N \Lambda^K f^N{}_{LK} \mathcal{F}^L - 2\mathcal{F}^M \partial_M \Sigma_N \mathcal{F}^N, \\ \delta_{\Lambda,\Sigma}\mathcal{G}_M &= \mathbb{L}_{(\Lambda,\Sigma)}\mathcal{G}_M - f^N{}_{KL} \mathcal{F}^K \partial_M \partial_N \Lambda^L + \partial_M \Sigma_N \mathcal{F}^N. \end{aligned} \quad (3.16)$$

C. Scalar potential

The last term in the action (3.1) is the scalar potential V which can be given as a function of the external metric $g_{\mu\nu}$ and the internal metric \mathcal{M}_{MN}

$$\begin{aligned} V &= -\frac{1}{240}\mathcal{M}^{MN}\partial_M\mathcal{M}^{KL}\partial_N\mathcal{M}_{KL} + \frac{1}{2}\mathcal{M}^{MN}\partial_M\mathcal{M}^{KL}\partial_L\mathcal{M}_{NK} + \frac{1}{7200}f^{NQ}{}_P f^{MS}{}_R \mathcal{M}^{PK}\partial_M\mathcal{M}_{QK}\mathcal{M}^{RL}\partial_N\mathcal{M}_{SL} \\ &\quad - \frac{1}{2}g^{-1}\partial_M g \partial_N \mathcal{M}^{MN} - \frac{1}{4}\mathcal{M}^{MN}g^{-1}\partial_M g g^{-1}\partial_N g - \frac{1}{4}\mathcal{M}^{MN}\partial_M g^{\mu\nu}\partial_N g_{\mu\nu}. \end{aligned} \quad (3.17)$$

The relative coefficients in this potential are determined by Λ^M and Σ^M gauge invariance by a computation similar to the one presented for the $E_{6(6)}$, $E_{7(7)}$ potentials in [2,3], that we briefly sketch in the following. For the calculation it turns out to be convenient to rewrite the potential as

$$\begin{aligned} V &= \frac{1}{4}j^R{}_M j^S{}_N (\mathcal{M}^{MN}\eta_{RS} - 2\mathcal{M}^{KL}f_{RL}{}^N f_{SK}{}^M + 2\delta_R{}^N \delta_S{}^M) - \frac{1}{2}g^{-1}\partial_M g \mathcal{M}^{MN} f_{NK}{}^P j_P{}^K \\ &\quad - \frac{1}{4}\mathcal{M}^{MN}g^{-1}\partial_M g g^{-1}\partial_N g - \frac{1}{4}\mathcal{M}^{MN}\partial_M g^{\mu\nu}\partial_N g_{\mu\nu}, \end{aligned} \quad (3.18)$$

in terms of the current j_M^P defined in analogy to (3.6) as

$$\mathcal{M}^{KP} \partial_M \mathcal{M}_{PL} \equiv j_M^P f_{PL}^K. \quad (3.19)$$

A short calculation shows that the noncovariant variation of the current j_M^N under generalized diffeomorphisms (2.5) is given by

$$\Delta^{\text{nc}} j_M^N = (\mathcal{M}^{NK} + \eta^{NK}) \partial_M (f_{KQ}^P \partial_P \Lambda^Q - \Sigma_K), \quad (3.20)$$

where we have used the invariance property (3.7) of the structure constants. It is then straightforward to verify that the noncovariant contributions from the variation of the various terms in (3.18) precisely cancel. In particular, we find that under Λ transformations the first line of (3.18) transforms according to

$$\begin{aligned} \frac{1}{4} \Delta_\Lambda^{\text{nc}} (j_M^R j_N^S (\mathcal{M}^{MN} \eta_{RS} - 2\mathcal{M}^{KL} f_{RL}^N f_{SK}^M + 2\delta_R^N \delta_S^M)) \\ = 2\partial_M \partial_P \Lambda^P \partial_N \mathcal{M}^{MN} + \partial_M \partial_L \Lambda^N \partial_N \mathcal{M}^{ML}, \end{aligned} \quad (3.21)$$

whereas the second line of (3.18) transforms into

$$\begin{aligned} \Delta_\Lambda^{\text{nc}} ((3.18), \text{second line}) \\ = -3\partial_M \partial_K \Lambda^K \partial_P \mathcal{M}^{PM} - e^{-1} \partial_M e \mathcal{M}^{MP} \partial_P \partial_R \Lambda^R \\ + e^{-1} \partial_M e \mathcal{M}^{SP} \partial_P \partial_S \Lambda^M. \end{aligned} \quad (3.22)$$

Together, this shows that the scalar potential term (eV) in the Lagrangian is invariant up to total derivatives.

Comparing the expression of (3.17) to other results in the literature [22] shows that the third term of (3.17) has been missed in previous constructions. Here, this term is essential for Λ^M and Σ_M invariance of the scalar potential. Absence of this term is the reason for the observed discrepancy of the scalar potential of [22] with $D = 11$ supergravity as we discuss in more detail in the last section.

D. External diffeomorphism invariance

The various terms of the EFT action (3.1) have been determined by invariance under generalized internal Λ^M , Σ_M diffeomorphisms. In contrast, the relative coefficients between the four terms are determined by invariance of the full action under the remaining gauge symmetries, which are a covariantized version of the external $(2+1)$ -dimensional diffeomorphisms with parameters $\xi^\mu(x, Y)$. For the Y -independent parameter, external diffeomorphism invariance is manifest. On the other hand, gauge invariance for general $\xi^\mu(x, Y)$ determines all relative coefficients, as we shall demonstrate in the following. The computation closely follows the analogous discussion for the $\text{SL}(2, \mathbb{R})$ -covariant formulation of four-dimensional Einstein gravity [20].⁴

⁴We note that here we use a field basis for A and B that is related to the $\text{SL}(2, \mathbb{R})$ treatment of [20] by a field redefinition.

Under general external diffeomorphisms, the external and internal metric transform in the standard (but covariantized) way

$$\delta_\xi \mathcal{M}_{MN} = \xi^\mu D_\mu \mathcal{M}_{MN}, \quad \delta_\xi e_\mu^a = \xi^\rho D_\rho e_\mu^a + D_\mu \xi^\rho e_\rho^a, \quad (3.23)$$

where we recall that the dreibein is an $E_{8(8)}$ scalar density of weight $\lambda = 1$. The transformation behavior of the gauge vectors is more complicated. Inspired by the $\text{SL}(2, \mathbb{R})$ case [20], for these fields we start from the ansatz

$$\begin{aligned} \delta_\xi^{(0)} A_\mu^M &= \xi^\nu F_{\nu\mu}^M + \mathcal{M}^{MN} g_{\mu\nu} \partial_N \xi^\nu, \\ \delta_\xi^{(0)} B_{\mu M} &= \xi^\nu G_{\nu\mu M} - j_M^K g_{\mu\nu} \partial_K \xi^\nu \\ &\quad + \frac{1}{4\kappa} e \varepsilon_{\mu\nu\lambda} g^{\lambda\rho} D^\nu (g_{\rho\sigma} \partial_M \xi^\sigma), \end{aligned} \quad (3.24)$$

where the noncovariant contributions will be required for particular cancellations in the variation of the Lagrangian. The full variation of these fields will be determined as we go along. Note that the form of the variation $\delta_\xi^{(0)} B_{\mu M}$ is manifestly compatible with the constraints (2.7) which this field is required to satisfy, because in the extra noncovariant terms the external index is carried by a derivative.

Let us now compute the variation of the Lagrangian (3.1) under (3.23), (3.24). To start with, let us work out the general variation of the Lagrangian (3.1) with respect to the vector fields which takes the form

$$\delta \mathcal{L} = \varepsilon^{\mu\nu\rho} (\mathcal{E}_{\nu\rho}^{(A)M} \delta B_{\mu M} + \mathcal{E}_{\nu\rho M}^{(B)} \delta A_\mu^M), \quad (3.25)$$

with

$$\begin{aligned} \mathcal{E}_{\mu\nu}^{(A)M} &\equiv 2\kappa F_{\mu\nu}^M + \frac{1}{2} e \varepsilon_{\mu\nu\rho} j^{\rho M}, \\ \mathcal{E}_{\mu\nu M}^{(B)} &\equiv 2\kappa \mathcal{G}_{\mu\nu M} - f_{MN}^K \partial_K \mathcal{E}_{\mu\nu}^{(A)N} \\ &\quad - \frac{1}{4} e \varepsilon_{\mu\nu\rho} (j_M^K j^\rho_K + 2\hat{J}^\rho_M), \end{aligned} \quad (3.26)$$

with the currents j_μ^N , j_M^N from (3.6) and (3.19), respectively, and the current \hat{J}^μ_M describing the contribution from the covariantized Einstein-Hilbert term,

$$\begin{aligned} \delta_A \mathcal{L}_{\text{EH}} &\equiv e \hat{J}^\mu_M \delta A_\mu^M \\ &= -2e e_a^\mu e_b^\nu (\partial_M \omega_\nu^{ab} - D_\nu (e^{\rho[a} \partial_M e_\rho^{b]}) \delta A_\mu^M. \end{aligned} \quad (3.27)$$

Note that not all components of $\mathcal{E}_{\mu\nu}^{(A)M}$ in (3.26) correspond to real equations of motion of the theory, as the field $B_{\mu M}$ is constrained by means of (2.7).

Next we consider the noncovariant variation of the covariantized Einstein-Hilbert term, which is given by [20]

$$\delta_{\xi}^{(0)}(e\hat{R}) = eF^{\mu\nu N}D_{\mu}(\partial_N\xi^{\rho}g_{\rho\nu}) + e\mathcal{M}^{MN}\hat{J}_{\mu M}\partial_N\xi^{\mu}, \quad (3.28)$$

where the second term comes from the noncovariant transformation (3.24) of the vector field A_{μ}^M via (3.27). The noncovariant variation of the Chern-Simons term follows from (3.9) and yields

$$\begin{aligned} \delta_{\xi}^{(0)}\mathcal{L}_{\text{CS}} &= -eF^{\mu\nu M}D_{\mu}(g_{\nu\sigma}\partial_M\xi^{\sigma}) - 2\kappa\varepsilon^{\mu\nu\rho}F_{\mu\nu}^M j_M^K g_{\rho\lambda}\partial_K\xi^{\lambda} - 2\kappa\varepsilon^{\mu\nu\rho}f_{MN}^K\partial_K F_{\mu\nu}^N \mathcal{M}^{ML}g_{\rho\sigma}\partial_L\xi^{\sigma} \\ &\quad + 2\kappa\varepsilon^{\mu\nu\rho}G_{\mu\nu M}\mathcal{M}^{MN}g_{\rho\sigma}\partial_N\xi^{\sigma} + \kappa\varepsilon^{\mu\nu\rho}\partial_K\xi^{\sigma}f_{MN}^K F_{\mu\nu}^M F_{\rho\sigma}^N, \end{aligned} \quad (3.29)$$

up to total derivatives. The first term cancels against the contribution from (3.28). Let us further rewrite the last term of (3.29) in terms of (3.26) as

$$\kappa\varepsilon^{\mu\nu\rho}\partial_K\xi^{\sigma}f_{MN}^K F_{\mu\nu}^M F_{\rho\sigma}^N = \frac{1}{4\kappa}\varepsilon^{\mu\nu\rho}\partial_K\xi^{\sigma}f_{MN}^K \mathcal{E}_{\mu\nu}^{(A)M}\mathcal{E}_{\rho\sigma}^{(A)N} - \partial_K\xi^{\mu}f_{MN}^K j^{\nu M}F_{\mu\nu}^N - \frac{1}{8\kappa}\varepsilon_{\mu\nu\rho}\partial_K\xi^{\mu}f_{MN}^K j^{\nu M}j^{\rho N}. \quad (3.30)$$

For the variation of the scalar kinetic term, we start from the variation

$$\delta^{(0)}(D_{\mu}\mathcal{M}_{MN}) = \mathcal{L}_{\xi}(D_{\mu}\mathcal{M}_{MN}) + \xi^{\nu}[D_{\mu}, D_{\nu}]\mathcal{M}_{MN} - \mathbb{L}_{(\delta_{\xi}^{(0)}A_{\mu}^M, \delta_{\xi}^{(0)}B_{\mu}^M)}\mathcal{M}_{MN}, \quad (3.31)$$

which induces the following variation of the kinetic term (3.5):

$$\begin{aligned} \delta_{\xi}^{(0)}\mathcal{L}_{\text{kin}} &= \frac{1}{2}e\mathcal{M}^{KL}j_K^N j_{\nu N}\partial_L\xi^{\nu} + ef^{MK}{}_L j^{\mu}{}_M F_{\mu\nu}^L \partial_K\xi^{\nu} + ej_L^K j_{\mu}^L \partial_K\xi^{\mu} \\ &\quad - ef^{MK}{}_L j^{\mu}{}_M \partial_K(\mathcal{M}^{LN}g_{\mu\nu}\partial_N\xi^{\nu}) - \frac{1}{4\kappa}\varepsilon^{\mu\nu\rho}j_{\mu}^L D_{\nu}(g_{\rho\sigma}\partial_L\xi^{\sigma}). \end{aligned} \quad (3.32)$$

Upon integration by parts, the last term gives rise to

$$-\frac{1}{4\kappa}\varepsilon^{\mu\nu\rho}g_{\rho\sigma}\partial_L\xi^{\sigma}D_{\mu}j_{\nu}^L = \frac{1}{480\kappa}\varepsilon^{\mu\nu\rho}g_{\rho\sigma}\partial_L\xi^{\sigma}f^{KL}{}_M \mathcal{M}_{KN}[D_{\mu}, D_{\nu}]\mathcal{M}^{MN} + \frac{1}{8\kappa}\varepsilon^{\mu\nu\rho}g_{\rho\sigma}\partial_K\xi^{\sigma}f_{MN}^K j_{\mu}^M j_{\nu}^N, \quad (3.33)$$

and evaluating the commutator of covariant derivatives yields terms that precisely cancel the three terms linear in $F_{\mu\nu}^M$ and $G_{\mu\nu M}$ from (3.29), provided we choose

$$\kappa \equiv \frac{1}{4}, \quad (3.34)$$

for the coupling constant of the CS term. Putting everything together, for the variation of the first three terms of the Lagrangian (3.1) we find up to total derivatives

$$\begin{aligned} \delta_{\xi}^{(0)}(\mathcal{L}_{\text{EH}} + \mathcal{L}_{\text{CS}} + \mathcal{L}_{\text{kin}}) &= \frac{1}{2}e(\mathcal{M}^{KL}\eta_{RS} + 2\delta_R^K\delta_S^L - 2\mathcal{M}^{MN}f^K{}_{MR}f^L{}_{NS})j_{\mu}^S j_L^R \partial_K\xi^{\mu} \\ &\quad + e\mathcal{M}^{MN}(\hat{J}_{\mu M}\partial_N\xi^{\mu} - f^{KL}{}_M j^{\mu}{}_K \partial_L(g_{\mu\nu}\partial_N\xi^{\nu})) + \varepsilon^{\mu\nu\rho}\partial_K\xi^{\sigma}f_{MN}^K \mathcal{E}_{\mu\nu}^{(A)M}\mathcal{E}_{\rho\sigma}^{(A)N}. \end{aligned} \quad (3.35)$$

It remains to compare this variation to the noncovariant variation of the scalar potential (3.17) under (3.23). Noting that

$$\begin{aligned} \delta_{\xi}(\partial_K\mathcal{M}_{MN}) &= \xi^{\mu}D_{\mu}(\partial_K\mathcal{M}_{MN}) + \partial_K\xi^{\mu}D_{\mu}\mathcal{M}_{MN}, \\ \delta_{\xi}(\partial_M g_{\mu\nu}) &= \mathcal{L}_{\xi}(\partial_M g_{\mu\nu}) + (\partial_M\xi^{\rho})D_{\rho}g_{\mu\nu} + 2\partial_M D_{(\mu}\xi^{\rho}g_{\nu)\rho}, \end{aligned} \quad (3.36)$$

it is straightforward to see from (3.18) that the noncovariant variation of the potential due to $\delta_{\xi}^{\text{nc}}(\partial_K\mathcal{M}_{MN})$ precisely cancels the first line of (3.35). Upon further calculation, the remaining contributions from variation of the potential combine with (3.35) into

$$\begin{aligned} \delta_{\xi}^{(0)}\mathcal{L} &= \left(e\hat{J}_{\mu M} - 2eD_{\mu}(e^{-1}\partial_M e) - D_{\nu}(eg_{\mu\rho}\partial_M g^{\nu\rho}) + \frac{1}{2}eD_{\mu}g_{\nu\rho}\partial_M g^{\nu\rho} \right) \mathcal{M}^{MN}\partial_N\xi^{\mu} \\ &\quad + \varepsilon^{\mu\nu\rho}\partial_K\xi^{\sigma}f_{MN}^K \mathcal{E}_{\mu\nu}^{(A)M}\mathcal{E}_{\rho\sigma}^{(A)N}. \end{aligned} \quad (3.37)$$

Using the definite expression (3.27) for $\hat{J}_{\mu M}$, an explicit calculation shows that the first line of (3.37) vanishes identically. We have thus shown that under external diffeomorphisms (3.23), (3.24), the variation of the Lagrangian (3.1) takes the compact form

$$\delta_{\xi}^{(0)} \mathcal{L} = \epsilon^{\mu\nu\rho} \partial_K \xi^\sigma f_{MN}{}^K \mathcal{E}_{\mu\nu}^{(A)M} \mathcal{E}_{\rho\sigma}^{(A)N}. \quad (3.38)$$

Just as in the $SL(2, \mathbb{R})$ case we conclude that invariance of the Lagrangian can be achieved by a further modification of the vector field transformation rules according to [20]

$$\begin{aligned} \delta_{\xi} A_{\mu}{}^M &= \delta_{\xi}^{(0)} A_{\mu}{}^M + 2\xi^{\nu} \mathcal{E}_{\mu\nu}^{(A)M}, \\ \delta_{\xi} B_{\mu M} &= \delta_{\xi}^{(0)} B_{\mu M} + 2\xi^{\nu} (\mathcal{E}_{\mu\nu}^{(B)} + f_{MN}{}^K \partial_K \mathcal{E}_{\mu\nu}^{(A)N}). \end{aligned} \quad (3.39)$$

It is straightforward to see that the new contributions due to the respective terms in $\xi^{\nu} \mathcal{E}_{\mu\nu}^{(A)M}$ and $\xi^{\nu} \mathcal{E}_{\mu\nu}^{(B)}$ take the form of an equation of motion symmetry and mutually cancel. The last term in (3.39) precisely cancels the variation of (3.38). Moreover, we note that the new variation $\delta_{\xi} B_{\mu M}$ continues to be consistent with the constraints (2.7) that this field is required to satisfy.

We may summarize the result of this subsection as follows: the action (3.1) is invariant under external diffeomorphisms parametrized by ξ^{μ} that on the internal and external metric act according to (3.23), while their action on the gauge fields follows from combining (3.24) and (3.39),

$$\begin{aligned} \delta_{\xi} A_{\mu}{}^M &= e \epsilon_{\mu\nu\rho} \xi^{\nu} j^{\rho M} + \mathcal{M}^{MN} g_{\mu\nu} \partial_N \xi^{\nu}, \\ \delta_{\xi} B_{\mu M} &= e \epsilon_{\mu\nu\rho} \left(g^{\rho\lambda} D^{\nu} (g_{\lambda\sigma} \partial_M \xi^{\sigma}) - \frac{1}{2} \xi^{\nu} j_M{}^K j^{\rho}{}_K - \xi^{\nu} \hat{j}^{\rho}{}_M \right) \\ &\quad - j_M{}^K g_{\mu\nu} \partial_K \xi^{\nu}. \end{aligned} \quad (3.40)$$

We have shown that invariance under external diffeomorphisms fixes all the relative coefficients in (3.1); the action is thus uniquely determined by combining internal and external generalized diffeomorphism invariance.

IV. EMBEDDING OF $D = 11$ SUPERGRAVITY

In the previous sections, we have constructed the unique $E_{8(8)}$ -covariant two-derivative action for the fields (1.2), that is invariant under generalized internal and external diffeomorphisms. It remains to establish its relation to $D = 11$ supergravity. Evaluating the field equations descending from (3.1) for an explicit appropriate solution of the section constraints (2.7), one may recover the full dynamics of $D = 11$ supergravity after rearranging the 11-dimensional fields according to a $3 + 8$ Kaluza-Klein split of the coordinates, but retaining the full dependence on all 11 coordinates as first explored in [18,19]. We have done this analysis in all detail in the $E_{6(6)}$ -covariant construction [2] and reproduced the full and untruncated action of 11-dimensional

supergravity from the $E_{6(6)}$ EFT after various redefinitions and redualizations of fields. Here, we keep the discussion brief, sketching the essential steps for the embedding of $D = 11$ supergravity and concentrating on the novel features of the $E_{8(8)}$ case. The complete analysis is left for future work.

The relevant solution of the section condition (1.1) is related to the splitting of coordinates according to the decomposition of the adjoint representation of $E_{8(8)}$ under its maximal $GL(8)$ subgroup:

$$\begin{aligned} \mathbf{248} &\longrightarrow 8_{+3} \oplus 28'_{+2} \oplus 56_{+1} \oplus (1 \oplus 63)_0 \oplus 56'_{-1} \oplus 28_{-2} \oplus 8'_{-3}, \\ \{Y^M\} &\longrightarrow \{y^m, y_{mn}, y^{kmn}, y_m{}^n, y_{kmn}, y^{mn}, y_m\}, \end{aligned} \quad (4.1)$$

with the subscripts referring to the grading with respect to the $GL(1) \subset GL(8)$ generator t_0 . The section constraints (2.7) are solved by truncating the coordinate dependence of all fields and gauge parameters to the coordinates in the 8_{+3} :

$$\Phi(x^\mu, Y^M) \longrightarrow \Phi(x^\mu, y^m). \quad (4.2)$$

In order to see that this truncation provides a solution for the section constraints (2.7), it is sufficient to observe that in the decomposition of the $\mathbf{3875}$ analogous to (4.1), the space of highest grading is an 8_{+5} , which shows that

$$(\mathbb{P}_{3875})_{MN}{}^{mn} = 0. \quad (4.3)$$

Accordingly, for the compensating gauge field constrained by (2.7) we set all but the associated eight components $B_{\mu m}$ to zero,

$$\begin{aligned} B_{\mu}{}^m &\rightarrow 0, & B_{\mu}{}_{mn} &\rightarrow 0, & B_{\mu}{}^{mnk} &\rightarrow 0, \\ B_{\mu m}{}^n &\rightarrow 0, & B_{\mu}{}_{mnk} &\rightarrow 0, & B_{\mu}{}^{mn} &\rightarrow 0, \end{aligned} \quad (4.4)$$

In order to recover the fields of $D = 11$ supergravity, we first express the scalar matrix $\mathcal{M}_{MN} = (\mathcal{V}\mathcal{V}^T)_{MN}$ in terms of a coset-valued vielbein $\mathcal{V} \in E_{8(8)}/SO(16)$ parametrized in triangular gauge associated to the grading of (4.1) as [33]

$$\begin{aligned} \mathcal{V} &\equiv \exp[\phi t_{(0)}] \mathcal{V}_8 \exp[c_{kmn} t_{(+1)}^{kmn}] \\ &\quad \times \exp[e^{klmnpqrs} c_{klmnpq} t_{(+2)rs}] \exp[\varphi_m t_{(+3)}^m]. \end{aligned} \quad (4.5)$$

Here, $t_{(0)}$ is the $E_{8(8)}$ generator associated to the $GL(1)$ grading of (4.1), \mathcal{V}_8 denotes a general element of the $SL(8) \subset GL(8)$ subgroup, whereas the $t_{(+n)}$ refer to the $E_{8(8)}$ generators of positive grading in (4.1).⁵ The scalar fields $c_{mnk} = c_{[mnk]}$ and $c_{mnlpq} = c_{[mnlpq]}$ have an obvious origin in the internal components of the 11-dimensional 3-form and 6-form. The scalar fields on the

⁵Explicit expressions for the matrix exponential (4.5) have been worked out in [22].

other hand represent the degree of freedom dual to the Kaluza-Klein vector fields $A_\mu{}^m$ in the standard decomposition of the 11-dimensional metric. Hence, formally they carry the degrees of the freedom of the dual graviton [23–26] which can be written in more suggestive form by defining

$$c_{m,n_1\dots n_8} \equiv \epsilon_{n_1\dots n_8} \varphi_m. \quad (4.6)$$

Similarly, the gauge field $A_\mu{}^M$ is split according to the decomposition (4.1) into

$$\{A_\mu{}^M\} \longrightarrow \{A_\mu{}^m, A_{\mu mn}, A_{\mu kmnpq}, A_{\mu m}{}^n, A_{\mu}{}^{kmnpq}, A_{\mu}{}^{mn}, A_{\mu m}\}. \quad (4.7)$$

Together with the surviving eight components from (4.4) we count 256 vector fields which appear to largely exceed the number of fields with possible 11-dimensional origin. Rather, from 11 dimensions we expect only the Kaluza-Klein vector fields $A_\mu{}^m$ together with gauge fields $A_{\mu mn}$ and $A_{\mu kmnpq}$ from the 3- and the 6-form, respectively. Fortunately, many of the fields in (4.7) do not in fact enter the Lagrangian (3.1). They are pure gauge as a consequence of the invariance of the action under the vector shift symmetry (2.22). Indeed, closer inspection of the covariant derivatives (2.18) and the Chern-Simons couplings (3.8) shows that out of (4.7) only the components $\{A_\mu{}^m, A_{\mu mn}, A_{\mu kmnpq}, A_{\mu m}{}^n\}$ enter the Lagrangian. More precisely, the covariant derivatives on the scalar fields evaluated in the parametrization of (4.5) are of the schematic form

$$\begin{aligned} D_\mu c_{kmn} &= D_\mu c_{kmn} + \partial_{[k} A_{|\mu|mn]}, \\ D_\mu c_{klmnpq} &= D_\mu c_{klmnpq} + \partial_{[k} A_{|\mu|lmnpq]} + \partial_{[k} A_{|\mu|lm} c_{npq]}, \\ D_\mu \varphi_m &= D_\mu \varphi_m + \dots + \partial_n A_{\mu m}{}^n + B_{\mu m}, \end{aligned} \quad (4.8)$$

where we have denoted by D_μ the derivative covariantized with the Kaluza-Klein vector field $A_\mu{}^m$ with respect to eight-dimensional internal diffeomorphisms. The unspecified terms in (4.8) refer to nonlinear couplings involving the scalar fields c_{kmn} and c_{klmnpq} . Integrating out the gauge field $B_{\mu m}$ thus not only eliminates all the dual graviton components φ_m but simultaneously eliminates all vector fields $A_{\mu m}{}^n$ from the Lagrangian. In this process, it is important that the scalar potential (3.17) does not depend on the scalar fields φ_m . Indeed, invariance of the Lagrangian under the shift $\varphi_m \rightarrow \varphi_m + c_m$ is a direct consequence of the invariance under generalized diffeomorphisms (2.5) with parameter Σ_m . This illustrates once more the role played by the additional covariantly constrained gauge symmetries Σ_M . Their presence and associated gauge connection $B_{\mu M}$ allow us to establish a covariant duality relation involving the degrees of freedom from the 11-dimensional metric and subsequently to

eliminate the dual graviton degrees of freedom φ_m from the Lagrangian.

In turn, this procedure of integrating out $B_{\mu m}$ induces a Yang-Mills-type coupling for the vector fields $A_\mu{}^m$ in a standard mechanism of three-dimensional supergravities [34]. To see this, note that the first line of the field equations (3.26) precisely relates the Yang-Mills field strength $F_{\mu\nu}{}^m$ to the scalar current as

$$F_{\mu\nu}{}^m = -e \epsilon_{\mu\nu\rho} j^{\rho m} = -e \epsilon_{\mu\nu\rho} M^{mn} D^\rho \varphi_n + \dots, \quad (4.9)$$

with $M^{mn} \equiv (\mathcal{V}_8 \mathcal{V}_8^T)^{mn}$.

The resulting Lagrangian then only depends on the fields

$$\{g_{\mu\nu}, \mathcal{V}_8, c_{kmn}, c_{klmnpq}, A_\mu{}^m, A_{\mu mn}, A_{\mu kmnpq}\} \quad (4.10)$$

corresponding to the various components of the 11-dimensional metric, 3-form and 6-form. Its field equations are proper combinations of the 11-dimensional field equations and the duality equation relating the 3-form and the 6-form. As an example, consider the field equations (3.26). With the first line corresponding to (4.9), we observe that the (mn) component of the second line gives rise to

$$f_{mn,N}{}^K \partial_K \mathcal{E}_{\mu\nu}^{(A)N} = 0 \Rightarrow \partial_{[k} (F_{|\mu\nu|mn]} + e j^{\rho mn} \epsilon_{\mu\nu\rho}) = 0, \quad (4.11)$$

which can be integrated to the duality equation

$$F_{\mu\nu mn} + e \epsilon_{\mu\nu\rho} j^{\rho mn} = \partial_{[m} B_{|\mu\nu|n]}, \quad (4.12)$$

with an undetermined 2-form $B_{\mu\nu}$. The latter can be identified with the corresponding component of the 11-dimensional 3-form. Indeed, further derivation $e^{\mu\nu\rho} \partial_\rho$ of (4.12) shows that it is compatible with the component

$$F_{\mu\nu\rho m} = e \epsilon_{\mu\nu\rho} \epsilon_{mn_1\dots n_7} F^{n_1\dots n_7} + \dots, \quad (4.13)$$

of the 11-dimensional duality equation (3-form \leftrightarrow 6-form) relating the field strength of $B_{\mu\nu m}$ on the lhs to the 7-form field strength $F_{n_1\dots n_7} = 7 \partial_{[n_1} c_{c_2\dots c_7]} + \dots$, whose internal derivative $\partial_{n_1} F^{n_1\dots n_7}$ appears as a source in the field equation for $\partial_\mu j^\mu{}_{mn}$. Equations (4.12) and (4.13) can further be used to eliminate all components c_{klmnpq} , $A_{\mu kmnpq}$ from the 11-dimensional 6-form from the equations, and the resulting equations of motion coincide with those coming from $D = 11$ supergravity with its standard field content.

On the level of the action, we get further confirmation from inspecting the scalar potential (3.17). After parametrization (4.5) of the 248-bein, evaluation of (4.2), and truncation of the external metric $g_{\mu\nu}$ to a warped Minkowski₃ geometry, the potential reduces to the schematic form

$$V_{\text{trunc}} \sim R(g) + F_{(4)}^2 + F_{(7)}^2 \quad (4.14)$$

reproducing the contributions from the $D = 11$ kinetic terms and Einstein-Hilbert term in the internal directions in terms of the fields from (4.10). This can be directly inferred from the analysis of [22] which obtains for the first line of (3.17) the expression (4.14) up to a term $F_{\text{dual grav}}^2$ resembling a kinetic term for the dual graviton components (4.6). The role of the third term in the full potential (3.17) (absent in [22]) is precisely to cancel this unwanted contribution. Indeed, the form of (3.18) shows that after imposing (4.2), the extra term is of the form

$$j_M^N j_N^M \rightarrow j_m^n j_n^m = M^{ml} M^{nk} (\partial_m \varphi_n) (\partial_k \varphi_l) + \dots = F_{\text{dual grav}}^2. \quad (4.15)$$

Moreover, since the full potential (3.17) by construction does not depend on φ_m , this confirms the result (4.14).

In view of the duality equation (4.13), the last two terms of the potential (4.14) both correspond to contributions F_{klmn}^2 and $F_{\mu\nu\rho m}^2$ from the original $D = 11$ 3-form kinetic term. This shows the necessity of the $F_{(7)}^2$ term in (4.14), carrying the contribution of the 2-forms $B_{\mu\nu m}$ which are not among the EFT fields in (4.10). The situation is different for the graviton. The $D = 11$ metric gives rise to the external and internal metric and the Kaluza-Klein vector fields, all of which are already encoded in the $E_{8(8)}$ EFT and show up in (4.10). Thus, there is no room for the inclusion of a dual graviton for this would double the number of metric degrees of freedom. Consequently, the match with $D = 11$ supergravity requires that the dual graviton term is absent in (4.14), as observed here. We conclude that there is no dual graviton problem. Summarizing, after rearranging all fields and coordinates of the $E_{8(8)}$ EFT, putting the appropriate solution of the section constraint, the action may eventually be matched to the one obtained by properly parametrizing 11-dimensional supergravity in the standard $3 + 8$ Kaluza-Klein split.

Let us finally mention that also IIB supergravity can be embedded into the $E_{8(8)}$ EFT (3.1). Just as for the $E_{6(6)}$ and $E_{7(7)}$ EFT [1–3], there is another inequivalent solution to the section conditions (2.7) that describes the embedding of the full ten-dimensional IIB theory [12,13] into the $E_{8(8)}$ EFT,⁶ generalizing the situation of type II double field theory [37,38]. For $E_{8(8)}$ the embedding of the IIB theory goes along similar lines as the $D = 11$ embedding described above, with the relevant decomposition $E_{8(8)} \rightarrow \text{GL}(7) \times \text{SL}(2)$ given by

⁶An analogous solution of the $\text{SL}(5)$ covariant section condition, corresponding to some three-dimensional truncation of type IIB, was discussed in the truncation of the theory to its potential term [35]. For a more general discussion of section constraints and type IIB solutions see [36].

$$\begin{aligned} 248 \longrightarrow & (7, 1)_{+4} \oplus (7', 2)_{+3} \oplus (35', 1)_{+2} \oplus (21, 1)_{+1} \\ & \oplus ((48, 1) \oplus (1, 3) \oplus (1, 1))_0 \oplus (21', 1)_{-1} \\ & \oplus (35, 1)_{-2} \oplus (7, 2)_{-3} \oplus (7', 1)_{-4}. \end{aligned} \quad (4.16)$$

The section constraint is then solved by having all fields depend on only the coordinates y^m in the $(7, 1)_{+4}$ and setting to zero all components of $B_{\mu M}$ other than the $B_{\mu m}$ in the $(7', 1)_{-4}$.

V. Summary and Outlook

In this paper we have given the details of the $E_{8(8)}$ exceptional field theory. As discussed in detail in the main text, the novel feature of this case is that the $E_{8(8)}$ valued generalized metric \mathcal{M}_{MN} encodes components of the dual graviton but nevertheless allows for a consistent (in particular gauge-invariant) dynamics thanks to the mechanism of constrained compensator fields introduced in [20] (that in turn is a duality-covariant extension of the proposal in [39]). This mechanism requires the presence of covariantly constrained gauge fields, which in the $D = 3$ case feature among the gauge vectors entering the covariant derivatives. These fields are unconventional, but seem to be indispensable for a gauge and duality invariant formulation. They are a generic feature of the exceptional field theories, corresponding in each case to a subset of the $(D - 2)$ -forms with D denoting the number of external dimensions [2,3].

Studying the truncations of these theories to the internal sector (i.e. neglecting all external coordinate dependence, external metric, and p -form fields), it has been a puzzle for a while how $E_{8(8)}$ generalized diffeomorphisms might be implemented as a consistent structure, given that their transformations do not close into an algebra [16,17]. In the full EFT the resolution is remarkably simple. Also in this case there is a gauge-invariant action (3.1) and nonclosure of generalized diffeomorphisms simply indicates an additional symmetry: the covariantly constrained Σ_M gauge transformations of (2.5). The associated gauge connection $B_{\mu M}$ then takes care of the dynamics of the dual graviton degrees of freedom, just as the analogous $(D - 2)$ -forms do in higher dimensions.

We have restricted the analysis to the bosonic sector of the theory, where generalized diffeomorphism invariance has proved sufficient to uniquely determine the action. We are confident that the extension to include fermions and the construction of a supersymmetric action is straightforward along the lines of the supersymmetric $D = 3$ gauged supergravity [21]. The fermions will transform as scalar densities under generalized diffeomorphisms (2.5) and in the spinor representations of the local ‘‘Lorentz group’’ $\text{SO}(1, 2) \times \text{SO}(16)$, as in [18,32]. For the $E_{7(7)}$ EFT [3] the full supersymmetric completion has recently been constructed in [40].

After completing the detailed construction of exceptional field theory for $E_{d(d)}$, $d = 6, 7, 8$, the question arises

whether one can go even further, perhaps starting with the affine Kac-Moody group $E_{9(9)}$. The pattern of compensating gauge fields in this case would suggest a new set of ‘‘covariantly constrained scalars’’ on top of the infinite hierarchy of fields parametrizing the coset space $E_{9(9)}/K(E_{9(9)})$. We refrain from further speculations.

ACKNOWLEDGMENTS

The work of O. H. is supported by the U.S. Department of Energy (DOE) under the cooperative research Agreement No. DE-FG02-05ER41360 and a DFG Heisenberg fellowship. We would like to thank Hadi and Mahdi Godazgar, Hermann Nicolai, and Barton Zwiebach for useful comments and discussions.

APPENDIX : CLOSURE OF $E_{8(8)}$ GENERALIZED LIE DERIVATIVES

Before proving closure of the gauge transformations, it is convenient to first derive the following lemma:

$$\begin{aligned} & f^R_{UV} f^{UP}{}_K f^{VQ}{}_L \partial_{(P} \otimes \partial_{Q)} \\ & \equiv f^R_{UV} f^{UP}{}_{[K} f^{VQ}{}_{L]} \partial_P \otimes \partial_Q \\ & = -(2\delta^P_{[K} f^{QR}{}_{L]} + \eta^{RP} f^Q{}_{KL}) \partial_{(P} \otimes \partial_{Q)}. \end{aligned} \quad (\text{A1})$$

In order to verify this, we compute by repeated use of the Jacobi identity

$$\begin{aligned} f^R_{UV} f^{UP}{}_{[K} f^{VQ}{}_{L]} & = -f^R_{U^P} f^U{}_{[K|V|} f^{VQ}{}_{L]} \\ & \quad - f^R_{U[K} f^U{}_{|V|}{}^P f^{VQ}{}_{L]} \\ & = -\frac{1}{2} f^R_{U^P} f^{UQ}{}_V f^V{}_{KL} \\ & \quad - f^R_{U[K} f^U{}_{|V|}{}^{UP} f^V{}_{L]}{}^Q. \end{aligned} \quad (\text{A2})$$

Inserting this form back into (A.1) we can apply in each term the Lemma (2.13), which then yields the right-hand side of (A.1). This completes the proof.

Next, we verify closure of the gauge transformations on a vector of weight 0,

$$[\delta_1, \delta_2]V^M = (\delta_{\Lambda_{12}} + \delta_{\Sigma_{12}})V^M, \quad (\text{A3})$$

according to the effective parameters (2.17). We compute for the left-hand side, first including only the Λ transformations,

$$\begin{aligned} [\delta_1, \delta_2]V^M & = \Lambda_2^K \partial_K (\Lambda_1^L \partial_L V^M - f^M{}_{NT} f^{TP}{}_Q \partial_P \Lambda_1^Q V^N) \\ & \quad - f^M{}_{NT} f^{TK}{}_L \partial_K \Lambda_2^L (\Lambda_1^P \partial_P V^N \\ & \quad - f^N{}_{PU} f^{UR}{}_S \partial_R \Lambda_1^S V^P) - (1 \leftrightarrow 2). \end{aligned} \quad (\text{A4})$$

Some terms cancel directly under the $(1 \leftrightarrow 2)$ antisymmetrization, and one finds

$$\begin{aligned} [\delta_1, \delta_2]V^M & = [\Lambda_2, \Lambda_1]^L \partial_L V^M - f^M{}_{NT} f^{TP}{}_Q \Lambda_2^K \partial_K \partial_P \Lambda_1^Q V^N \\ & \quad + f^M{}_{NT} f^{TK}{}_L f^N{}_{PU} f^{UR}{}_S \partial_K \Lambda_2^L \partial_R \Lambda_1^S V^P \\ & \quad - (1 \leftrightarrow 2). \end{aligned} \quad (\text{A5})$$

Here, we denoted by $[\cdot, \cdot]$ the conventional Lie bracket. It turns out, however, that the extra terms in the E bracket (2.17), as compared to the Lie bracket, vanish in the transport term due to the section constraints, so that the transport term already has the desired form. We find it convenient to work for now with a different but equivalent effective parameter,

$$\begin{aligned} \Lambda_{12}^M & = \Lambda_2^N \partial_N \Lambda_1^M - 7(\mathbb{P}_{3875})^{MK}{}_{NL} \Lambda_2^N \partial_K \Lambda_1^L \\ & \quad - \frac{1}{8} \eta^{MK} \eta_{NL} \Lambda_2^N \partial_K \Lambda_1^L - (1 \leftrightarrow 2). \end{aligned} \quad (\text{A6})$$

Comparing then with the form of the gauge transformation with respect to this Λ_{12} we read off for the remaining terms

$$\begin{aligned} [\delta_1, \delta_2]V^M & = \Lambda_{12}^L \partial_L V^M - f^M{}_{NT} f^{TP}{}_Q \partial_P \Lambda_{12}^Q V^N \\ & \quad + f^M{}_{NT} f^{TP}{}_Q \partial_P \Lambda_2^L \partial_L \Lambda_1^Q V^N \\ & \quad - 7 f^M{}_{NT} f^{TP}{}_Q \mathbb{P}^{QK}{}_{RS} \partial_P (\Lambda_2^R \partial_K \Lambda_1^S) V^N \\ & \quad + f^M{}_{NT} f^{TK}{}_L f^N{}_{PU} f^{UR}{}_S \partial_K \Lambda_2^L \partial_R \Lambda_1^S V^P \\ & \quad - (1 \leftrightarrow 2), \end{aligned} \quad (\text{A7})$$

where here and in the following we omit the representation label on the **3875** projector \mathbb{P} , as it can always be distinguished from its index structure. The terms in the first line are the ones desired for closure, while the remaining terms are extra. We next have to show that these are zero or else can be brought to the form of Σ_M gauge transformations.

We investigate terms with $\partial\Lambda\partial\Lambda$ and $\Lambda\partial\partial\Lambda$ separately. The latter originate from the term in the third line. Inserting the projector (2.4) we compute

$$\begin{aligned} & -7 f^M{}_{NT} f^{TP}{}_Q \mathbb{P}^{QK}{}_{RS} \partial_P (\Lambda_2^R \partial_K \Lambda_1^S) V^N \\ & = -\frac{1}{2} f^M{}_{NT} f^{TP}{}_Q [2\delta_R^{(Q} \delta_S^{K)} - f^U{}_R ({}^Q f_{US}{}^K)] \Lambda_2^R \partial_P \partial_K \Lambda_1^S. \end{aligned} \quad (\text{A8})$$

Writing this out yields four terms, two with ff and two with $ffff$. Using the Lemma (A.1) we can then reduce the $ffff$ terms to ff terms. After some algebra, one finds that all ff terms cancel, proving that the $\Lambda\partial\partial\Lambda$ structures in (A.7) actually drop out. Next, we turn to the $\partial\Lambda\partial\Lambda$ structures. The strategy here is to implement the antisymmetry in $(1 \leftrightarrow 2)$ by decomposing the terms into structures of the form $\partial_{(P} \Lambda_2^{[R} \partial_K) \Lambda_1^S]$ and $\partial_{[P} \Lambda_2^{(R} \partial_K) \Lambda_1^S]$. In the former, $ffff$ terms can then be reduced to ff terms by means

of (A.1). After some algebra, one then finds for the terms in the second line of (A.7)

$$\begin{aligned} & -7f^M_{NT}f^{TP}{}_Q\mathbb{P}^{QK}{}_{RS}\partial_P\Lambda_2^R\partial_K\Lambda_1^S - (1\leftrightarrow 2) \\ & = -2f^M_{NT}f^{TP}{}_Q\partial_{[P}\Lambda_2^{(Q}\partial_{K]}\Lambda_1^{K)} \\ & \quad - f^M_{NT}f_S{}^{UQ}f_U{}^K{}_Rf_Q{}^{PT}\partial_{[P}\Lambda_2^{(R}\partial_{K]}\Lambda_1^S). \end{aligned} \quad (\text{A9})$$

Combing with the first term in the second line of (A.7) one obtains

$$\begin{aligned} & f^M_{NT}f^{TP}{}_Q\partial_P\Lambda_2^L\partial_L\Lambda_1^Q - 7f^M_{NT}f^{TP}{}_Q\mathbb{P}^{QK}{}_{RS}\partial_P\Lambda_2^R\partial_K\Lambda_1^S \\ & \quad - (1\leftrightarrow 2) \\ & = 2f^M_{NT}f^{TP}{}_Q\partial_{(P}\Lambda_2^{[K}\partial_{K]}\Lambda_1^{Q]} \\ & \quad - f^M_{NT}f_S{}^{UQ}f_U{}^K{}_Rf_Q{}^{PT}\partial_{[P}\Lambda_2^{(R}\partial_{K]}\Lambda_1^S). \end{aligned} \quad (\text{A10})$$

Next, we have to simplify the terms in the fourth line of (A.7). We first note that the antisymmetrization in $(1\leftrightarrow 2)$ imposes an antisymmetrization of the T, U indices in $f^{TK}{}_L f^{UR}{}_S$. This structure can thus be written as

$$\begin{aligned} & 2f^M_{N[T}f^{TK}{}_L f^N{}_{P|U]}f^{UR}{}_S\partial_K\Lambda_2^L\partial_R\Lambda_1^S \\ & = -f^M_{NP}f^N{}_{UT}f^{TK}{}_L f^{UR}{}_S\partial_K\Lambda_2^L\partial_R\Lambda_1^S, \end{aligned} \quad (\text{A11})$$

where we used the Jacobi identity for the contraction of the first and third structure constants. In this form the antisymmetry in $(1\leftrightarrow 2)$ is manifest. Next, we can decompose the index pair in $\partial_K\Lambda_2\partial_R\Lambda_1$ into its symmetric and antisymmetric part. Applying then for the symmetric part the Lemma (A.1), one finds after some straightforward algebra that these terms equal

$$\begin{aligned} (\text{A.11}) & = f^M_{NP}f^N{}_{TU}f^{TK}{}_L f^{UR}{}_S\partial_{[K}\Lambda_2^L\partial_{R]}\Lambda_1^S \\ & \quad - f^M_{NP}f^{RN}{}_S\partial_{(K}\Lambda_2^K\partial_{R)}\Lambda_1^S \\ & \quad + f^M_{NP}f^{RN}{}_L\partial_{(K}\Lambda_2^L\partial_{R)}\Lambda_1^K \\ & \quad - f^{MK}{}_P f^R{}_{LS}\partial_{(K}\Lambda_2^L\partial_{R)}\Lambda_1^S. \end{aligned} \quad (\text{A12})$$

Combining now (A.10) and (A.12) in the gauge algebra (A.7) one finds that the $ffff$ terms, after relabeling of indices, combine into

$$-f^M_{NT}f^{UK}{}_R(f_{SU}{}^Q f_Q{}^{PT} - f_S{}^{PQ} f_{QU}{}^T)\partial_{[P}\Lambda_2^{(R}\partial_{K]}\Lambda_1^S)V^N. \quad (\text{A13})$$

The terms in parentheses can now be combined by the Jacobi identity to give the structure $f_S{}^{TQ} f_{QU}{}^P$, after which it follows with the Lemma (2.13) that this term vanishes. The ff terms combine to give the following final result

$$[\delta_1, \delta_2]V^M = \delta_{\Lambda_{12}}V^M - f^{MK}{}_P(f^R{}_{LS}\partial_{(K}\Lambda_2^L\partial_{R)}\Lambda_1^S)V^P. \quad (\text{A14})$$

As required, the extra term can be interpreted as a Σ_M gauge transformation, so that we established in total

$$\begin{aligned} [\delta_1, \delta_2]V^M & = \delta_{\Lambda_{12}}V^M + \delta_{\Sigma_{12}}V^M, \\ \Sigma_{12M} & = -f^N{}_{PQ}\partial_{(M}\Lambda_2^P\partial_{N)}\Lambda_1^Q. \end{aligned} \quad (\text{A15})$$

Note that the Σ_M gauge parameter is manifestly covariantly constrained in that its free index is always carried by a derivative. This completes the proof of closure. Finally, we may redefine these gauge parameters by trivial parameters of the form (2.12),

$$\chi^K = \frac{1}{4}f^K{}_{PQ}\Lambda_2^P\Lambda_1^Q. \quad (\text{A16})$$

This brings the gauge algebra into the equivalent form (2.17) that we used in the main text. We finally note that the closure of Σ and Λ transformations as indicated in (2.17) follows by a straightforward computation that uses the Lemma (2.13) for the constrained parameters Σ_M . This concludes our proof of closure of the $E_{8(8)}$ generalized Lie derivatives.

[1] O. Hohm and H. Samtleben, *Phys. Rev. Lett.* **111**, 231601 (2013).
 [2] O. Hohm and H. Samtleben, *Phys. Rev. D* **89**, 066016 (2014).
 [3] O. Hohm and H. Samtleben, *Phys. Rev. D* **89**, 066017 (2014).
 [4] W. Siegel, *Phys. Rev. D* **48**, 2826 (1993).
 [5] C. Hull and B. Zwiebach, *J. High Energy Phys.* **09** (2009) 099.

[6] C. Hull and B. Zwiebach, *J. High Energy Phys.* **09** (2009) 090.
 [7] O. Hohm, C. Hull, and B. Zwiebach, *J. High Energy Phys.* **07** (2010) 016.
 [8] O. Hohm, C. Hull, and B. Zwiebach, *J. High Energy Phys.* **08** (2010) 008.
 [9] O. Hohm and S. K. Kwak, *J. Phys. A* **44**, 085404 (2011).
 [10] O. Hohm, D. Lüst, and B. Zwiebach, *Fortschr. Phys.* **61**, 926 (2013).

- [11] E. Cremmer, B. Julia, and J. Scherk, *Phys. Lett. B* **76**, 409 (1978).
- [12] J. H. Schwarz and P. C. West, *Phys. Lett. B* **126**, 301 (1983).
- [13] P. S. Howe and P. C. West, *Nucl. Phys.* **B238**, 181 (1984).
- [14] E. Cremmer and B. Julia, *Nucl. Phys.* **B159**, 141 (1979).
- [15] P. C. West, *Phys. Lett. B* **575**, 333 (2003).
- [16] A. Coimbra, C. Strickland-Constable, and D. Waldram, *J. High Energy Phys.* **02** (2014) 054.
- [17] D. S. Berman, M. Cederwall, A. Kleinschmidt, and D. C. Thompson, *J. High Energy Phys.* **01** (2013) 064.
- [18] H. Nicolai, *Phys. Lett. B* **187**, 316 (1987).
- [19] K. Koepsell, H. Nicolai, and H. Samtleben, *Classical Quantum Gravity* **17**, 3689 (2000).
- [20] O. Hohm and H. Samtleben, *J. High Energy Phys.* **09** (2013) 080.
- [21] H. Nicolai and H. Samtleben, *Phys. Rev. Lett.* **86**, 1686 (2001).
- [22] H. Godazgar, M. Godazgar, and M. J. Perry, *J. High Energy Phys.* **06** (2013) 044.
- [23] T. Curtright, *Phys. Lett. B* **165**, 304 (1985).
- [24] C. Hull, *Nucl. Phys.* **B583**, 237 (2000).
- [25] P. C. West, *Classical Quantum Gravity* **18**, 4443 (2001).
- [26] C. Hull, *J. High Energy Phys.* **09** (2001) 027.
- [27] X. Bekaert, N. Boulanger, and M. Henneaux, *Phys. Rev. D* **67**, 044010 (2003).
- [28] X. Bekaert, N. Boulanger, and S. Cnockaert, *J. Math. Phys. (N.Y.)* **46**, 012303 (2005).
- [29] K. Koepsell, H. Nicolai, and H. Samtleben, *J. High Energy Phys.* **04** (1999) 023.
- [30] B. de Wit, H. Nicolai, and H. Samtleben, *J. High Energy Phys.* **02** (2008) 044.
- [31] O. Hohm and H. Samtleben, *Phys. Rev. D* **88**, 085005 (2013).
- [32] N. Marcus and J. H. Schwarz, *Nucl. Phys.* **B228**, 145 (1983).
- [33] E. Cremmer, B. Julia, H. Lu, and C. N. Pope, *Nucl. Phys.* **B523**, 73 (1998).
- [34] H. Nicolai and H. Samtleben, *Nucl. Phys.* **B668**, 167 (2003).
- [35] C. D. Blair, E. Malek, and J.-H. Park, *J. High Energy Phys.* **01** (2014) 172.
- [36] C. Strickland-Constable, [arXiv:1310.4196](https://arxiv.org/abs/1310.4196).
- [37] O. Hohm, S. K. Kwak, and B. Zwiebach, *Phys. Rev. Lett.* **107**, 171603 (2011).
- [38] O. Hohm, S. K. Kwak, and B. Zwiebach, *J. High Energy Phys.* **09** (2011) 013.
- [39] N. Boulanger and O. Hohm, *Phys. Rev. D* **78**, 064027 (2008).
- [40] H. Godazgar, M. Godazgar, O. Hohm, H. Nicolai, and H. Samtleben, [arXiv:1406.3235](https://arxiv.org/abs/1406.3235) [*J. High Energy Phys.* (to be published)].