

Measurement of the antineutrino neutral-current elastic differential cross section

A. A. Aguilar-Arevalo,¹² B. C. Brown,⁶ L. Bugel,¹¹ G. Cheng,⁵ E. D. Church,¹⁵ J. M. Conrad,¹¹ R. Dharmapalan,¹ Z. Djurcic,² D. A. Finley,⁶ R. Ford,⁶ F. G. Garcia,⁶ G. T. Garvey,⁹ J. Grange,⁷ W. Huelsnitz,⁹ C. Ignarra,¹¹ R. Imlay,¹⁰ R. A. Johnson,³ G. Karagiorgi,⁵ T. Katori,^{11,*} T. Kobilarcik,⁶ W. C. Louis,⁹ C. Mariani,¹⁶ W. Marsh,⁶ G. B. Mills,⁹ J. Mirabal,⁹ C. D. Moore,⁶ J. Mousseau,⁷ P. Nienaber,¹⁴ B. Osmanov,⁷ Z. Pavlovic,⁹ D. Perevalov,⁶ C. C. Polly,⁶ H. Ray,⁷ B. P. Roe,¹³ A. D. Russell,⁶ M. H. Shaevitz,⁵ J. Spitz,¹¹ I. Stancu,¹ R. Tayloe,⁸ R. G. Van de Water,⁹ M. O. Wascko,¹⁷ D. H. White,⁹ D. A. Wickremasinghe,³ G. P. Zeller,⁶ and E. D. Zimmerman⁴

(MiniBooNE Collaboration)

¹University of Alabama, Tuscaloosa, Alabama 35487, USA

²Argonne National Laboratory, Argonne, Illinois 60439, USA

³University of Cincinnati, Cincinnati, Ohio 45221, USA

⁴University of Colorado, Boulder, Colorado 80309, USA

⁵Columbia University, New York, New York 10027, USA

⁶Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

⁷University of Florida, Gainesville, Florida 32611, USA

⁸Indiana University, Bloomington, Indiana 47405, USA

⁹Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

¹⁰Louisiana State University, Baton Rouge, Louisiana 70803, USA

¹¹Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

¹²Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, D.F. 04510, México

¹³University of Michigan, Ann Arbor, Michigan 48109, USA

¹⁴Saint Mary's University of Minnesota, Winona, Minnesota 55987, USA

¹⁵Yale University, New Haven, Connecticut 06520, USA

¹⁶Center for Neutrino Physics, Virginia Tech, Blacksburg, Virginia 24061, USA

¹⁷Imperial College London, London SW7 2AZ, United Kingdom

(Received 30 September 2013; published 8 January 2015)

We report the measurement of the flux-averaged antineutrino neutral current elastic scattering cross section ($d\sigma_{\bar{\nu}N \rightarrow \bar{\nu}N}/dQ^2$) on CH_2 by the MiniBooNE experiment using the largest sample of antineutrino neutral current elastic candidate events ever collected. The ratio of the antineutrino to neutrino neutral current elastic scattering cross sections and a ratio of the antineutrino neutral current elastic to antineutrino charged current quasielastic cross sections are also presented.

DOI: 10.1103/PhysRevD.91.012004

PACS numbers: 13.15.+g, 12.15.Mm, 13.85.Dz, 14.20.Dh

I. INTRODUCTION

One of the simplest weak neutral current interactions is the elastic scattering of a neutrino from a nucleon (NCE). This process is sensitive to both isoscalar and isovector weak currents carried by the nucleon, whereas charge current quasielastic (CCQE) scattering is sensitive to only the isovector current. Both NCE and CCQE neutrino interactions are important for accelerator-based neutrino oscillation experiments, and to date, very few measurements in the GeV regime have been made, particularly with antineutrinos [1]. With regards to the definition of a NCE interaction, it should be noted that in the antineutrino-nucleon NCE scattering reported here the antineutrino scatters off of both free and bound nucleons as we report in Appendix A.

Recent measurements of neutrino-nucleus CCQE scattering on ^{12}C show an enhanced cross section [2] relative to the prediction from impulse approximation calculations, such as the relativistic Fermi gas (RFG) model of the nucleus [3]. The enhancement is likely to arise from nucleon-nucleon correlations absent in the RFG model, and NCE scattering provides a complementary channel to further examine the nuclear effects common to both CCQE and NCE neutrino-nucleon scattering. Electron-nucleus scattering can also provide clues to understand neutrino-nucleus scattering and have been studied recently [4–6].

The Mini Booster Neutrino Experiment (MiniBooNE) has previously reported high-statistics measurements of neutrino CCQE (νCCQE) and neutrino NCE (νNCE) scattering cross sections [2,7] on carbon. Recently, a measurement of the neutrino content of the antineutrino mode flux was carried out [8], and an antineutrino CCQE

*Present address: Queen Mary University of London, London E1 4NS, United Kingdom.

($\bar{\nu}$ CCQE) cross section measurement was published [9]. Sizable nuclear effects are observed in both the ν CCQE and $\bar{\nu}$ CCQE data, suggesting contributions from nucleon-nucleon correlations and two-body exchange currents [10–15]. The ν NCE cross section has been studied to quantify these nuclear effects [16–19]. As the RFG model does not incorporate the nuclear effects, agreement between the measured cross sections (ν CCQE, ν NCE, and $\bar{\nu}$ CCQE) and the cross section model was achieved by assigning a higher value ($\sim 30\%$) to the axial mass (M_A) parameter in the axial-vector form factor. For a detailed discussion of neutrino cross section measurements and model predictions, see Ref. [1].

The antineutrino-nucleus NCE ($\bar{\nu}$ NCE) scattering measurement reported here is part of a series of measurements providing an understanding of the neutrino flux and cross sections in the energy regime accessible to MiniBooNE [2,7,9,20–22]. The data set corresponds to 10.09×10^{20} protons on the neutrino production target. The experimental signature is the same as the ν NCE scattering [7]—the scintillation light produced by the recoil nucleons. The sample size for the $\bar{\nu}$ NCE scattering cross section—60,605 events with 40% sample purity—is the largest collected to date for this type of interaction. Also reported here are the first experimental measurements of the antineutrino to neutrino NCE cross section and $\bar{\nu}$ NCE to $\bar{\nu}$ CCQE cross section ratios.

Previously, a few experiments have measured neutrino-nucleon NCE scattering [23,24], most notably the E734 experiment [25] at Brookhaven National Laboratory (BNL) which reported both neutrino-proton and antineutrino-proton NCE scattering measurements as a function of four-momentum transfer squared (Q^2) with 1,686 and 1,821 candidate events respectively.

II. MINIBOONE EXPERIMENT

A. Experimental setup

The MiniBooNE experiment, located at Fermi National Accelerator Laboratory, proposed to test the short baseline neutrino oscillations reported by the LSND experiment [26,27]. In addition, the experiment is well suited to measure a variety of high-statistic, neutrino cross sections [2,7,20–22]. It is situated in the Booster Neutrino Beamline (BNB) that produces the neutrino beam via the decay of mesons produced in a proton-beryllium interaction. The primary proton beam with a momentum of 8.89 GeV/c is extracted from the Fermilab Booster in 1.6 μ s pulses with $\sim 4 \times 10^{12}$ protons in each beam pulse. They impinge on a beryllium target placed in a magnetic focusing horn. The p-Be interactions produce a secondary beam of mesons that can be selectively focused or defocused by the magnetic horn. In antineutrino mode, the magnetic horn focuses negatively charged particles and defocuses positively charged particles. The mesons then

decay in an air-filled decay pipe producing a beam of (anti) neutrinos. Magnetic horn focusing also increases the desired neutrino flux reaching the MiniBooNE detector by a factor of ~ 6 . The average energy of the antineutrino beam is about 650 MeV. Further details on the BNB can be found in Ref. [28].

The MiniBooNE detector, situated 545 m from the Be target, is a spherical steel tank with a radius of 610 cm filled with 800 tons of mineral oil. The mineral oil [29] serves both as a target for the neutrino beam and the medium in which the resultant particles from neutrino interactions propagate. The detector is divided into two optically isolated regions separated by a spherical shell of radius 575 cm. The inner sphere, referred to as the *signal* region, is lined with 1,280 inward-pointing 8-in. photomultiplier tubes (PMTs) [30]. The outer shell is the *veto* region with 240 PMTs arranged in back-to-back pairs pointed along the circumference of the detector. Charged particles produced in the neutrino interaction emit Cherenkov and scintillation light that is collected by the PMTs. Six steel legs support the detector, situated in a vault along with the detector electronics and data acquisition systems. The entire assembly is buried under approximately 3 m of earth overburden to reduce cosmic ray backgrounds. Further details about the MiniBooNE detector can be found in Ref. [31].

B. Flux prediction, cross section model, and detector simulation

A GEANT4-based Monte Carlo (MC) beam simulation [32] is used to calculate the neutrino and antineutrino flux at the detector. The simulation accepts as input the shape, location, and material of the components of the BNB, the MiniBooNE target hall, and the meson decay volume through which the primary protons, the secondary mesons, and tertiary neutrinos propagate. The various components of simulation depend on the specific processes in the beamline and arise from a combination of constraints which include other particle production software, external measurements by MiniBooNE or other experiments in a similar energy regime, theoretical predictions, and extrapolation of external measurements to MiniBooNE energies.

Most of the neutrinos seen by the detector come from the decay of primary π^+ and π^- produced in the p-Be interaction as well as their subsequent μ^+ and μ^- decays. The π^+ and π^- production tables used in the MC simulation come from a parametrization of the HARP experiment [33] which measured pion production on a replica Be target at 8.89 GeV/c. The resulting neutrino flux prediction for antineutrino mode running is shown in Fig. 1 (the flux tables are available in Ref. [34]). The neutrino contamination in the antineutrino mode beam is higher ($\sim 16\%$) as compared to the corresponding antineutrino contamination of the neutrino mode beam ($\sim 6\%$). For details on the MiniBooNE flux prediction in both modes, see Ref. [28]. The neutrino contamination in the antineutrino mode beam




FIG. 1 (color online). The predicted antineutrino mode flux at the MiniBooNE detector for different types of neutrinos as a function of their energy as reported in Refs. [28] and [34].

was measured by the MiniBooNE collaboration [8] and the results applied to the flux estimate used in the present measurement.

Neutrino interaction rates, products, and their kinematics in the MiniBooNE detector are predicted using the NUANCE [35] neutrino event generator that has been customized to the MiniBooNE experiment. It has as input the neutrino flux prediction described above, as well as the detector target material and geometry. The mineral oil target is CH_2 with a density of 0.845 g/cm^3 . The (anti)neutrino NCE scattering off of free protons is modeled using the Llewellyn–Smith formalism [36], while for bound nucleons the RFG model of Smith and Moniz [3] is used. In NUANCE pion production is assumed to occur via delta production as per Rein and Sehgal’s prescription [37].

The various parameters in NUANCE are tuned as follows: All the nucleon’s vector form factors are assumed to retain their conventional values while the mass in the axial vector M_A is assigned a value of 1.23 GeV for the nucleons bound in carbon (as per MiniBooNE νCCQE measurement [38]) and 1.13 GeV for free nucleons (which is the average between the values measured by the deuterium based scattering experiments and MiniBooNE). In the low Q^2 regime, the Pauli blocking parameter was scaled up by a scaling factor, $\kappa = 1.022$, in order to match the observed MiniBooNE νCCQE data—see Ref. [39] for details. Note that these values are used for the CCQE/NCE channels that are background to this measurement, and the changes as indicated in more recent analyses [2,7] are covered by systematic errors on these parameters. In the case of neutrino induced resonant pion production, the form factors are assumed to be identical to those used in NCE and CCQE interaction, with the exception of the axial vector mass, where we take $M_A = 1.1 \text{ GeV}$ —see Ref. [21]. A 20% probability is assigned to the possibility that the outgoing pion is absorbed within the nucleus through final state interactions (FSIs). In this case the final product of the interaction is just the nucleon, similar to a neutrino NCE

interaction. Lastly, the strange quark contribution to the vector and axial vector form factor is taken to be zero with an uncertainty of 0.1.

The neutrino-generated final states output by the MiniBooNE neutrino event generator (NUANCE) are passed on to the MiniBooNE detector simulation. A GEANT3 simulation software [40] in conjunction with a customized optical model is used to simulate particle propagation, the resulting light emission and propagation, and the PMT response in the MiniBooNE detector. Some modifications to the standard GEANT3 routines include an improved model for Dalitz decay ($\pi^0 \rightarrow e^+e^-\gamma$), muon decay ($\mu \rightarrow e\nu\nu$), and the possibility of μ^- capture by carbon. The default GFLUKA [41] package is used to model hadron interactions. The MiniBooNE optical model has 12 components with a total of 35 adjustable parameters which have been tuned using external measurements and calibration data. The various components of the optical model include the index of refraction of the oil, light extinction length, the propagation and detection efficiency of Cherenkov light, scintillation and fluorescence yields of the different fluors present in the oil, scattering and reflections in the detector, and relative and angular efficiencies of the PMTs. The scintillation photons were modelled as per Birk’s law [42], and its coefficients are additional parameters in the optical model. The charge and time response of the PMTs were modelled by parametrization of data collected by the PMT studies using a pulsed laser source and calibration light sources in the detector [30]. Finally, the detector simulation includes modelling of signal digitization of the PMT outputs and the data acquisition.

III. NEUTRAL-CURRENT ELASTIC ANALYSIS

A. Event reconstruction

In the case of NCE event reconstruction, each event is assumed to be due to a proton of which the Cherenkov and scintillation light profiles are determined from the MC simulation. NCE scattering resulting in outgoing neutrons is only seen through their subsequent strong interactions resulting in protons; hence, NCE neutrons are indistinguishable from NCE protons. Most NCE protons are below Cherenkov threshold (350 MeV) and are reconstructed primarily via the scintillation light yields. Figure 2 (top) shows the MC prediction of reconstructed energy spectrum for NCE protons and neutrons. We see that most of the scattered nucleons are below threshold and that NCE neutrons and NCE protons have a similar energy profile.

The charge and time information from the PMTs is used to determine the position, time, direction, and energy of an event by employing a log-likelihood minimization method. Outgoing protons from NCE scattering have a characteristic light emission profile which readily allows for their particle identification. The ability to differentiate protons from beam unrelated events (mostly electrons) is illustrated




FIG. 2 (color online). $\bar{\nu}$ NCE reconstruction in MiniBooNE. The top plot shows the MC predicted energy spectrum for $\bar{\nu}$ NCE protons and neutrons. We see that most of the scattered nucleons are below the Cherenkov threshold for protons in the MiniBooNE medium (350 MeV—shown by the dotted red line) and that both neutrons and protons have a similar spectrum. The bottom plot shows the fraction of late (scintillation) light for $\bar{\nu}$ NCE data and MC, as a function of reconstructed nucleon energy. The errors on data represent the uncertainty in the modelling of the optical photons resulting from the scintillation light. We see that there is good agreement between the prediction and data from 50 to 350 MeV and also the Cherenkov threshold transition matches at 350 MeV (shown by the dotted red line).

in Fig. 3 where the fraction of prompt light emitted is plotted vs the number of tank PMT hits. Prompt light is defined as the fraction of PMT hits with corrected time between -5 and 5 ns, where the corrected time is the time difference between the PMT hit time and the reconstructed event time with light propagation time from the reconstructed vertex to the PMT taken into account.

The MiniBooNE detector position resolution for NCE protons is ~ 0.75 and ~ 1.35 m for neutrons. The energy resolution is $\sim 20\%$ for protons and $\sim 30\%$ for neutrons. Energy scaling for NCE protons was checked by plotting the fraction of scintillation or late light as a function of reconstructed energy, as shown in Fig. 2 (bottom). There is agreement (within errors) between data and MC both in the energy regime of interest (50 to 350 MeV) and the




FIG. 3 (color online). Fraction of prompt photons compared to the total number for beam unrelated events and NCE MC events reconstructed under an electron hypothesis. The events out of time with the beam are due to Michel decay, while the events in time with the beam are due to scintillation light from recoiling nucleons. The error bars correspond to the rms of the distributions.

Cherenkov threshold transition at 350 MeV. For details on reconstruction methods used in MiniBooNE, see Ref. [43]; for the NCE event reconstruction in particular, see Ref. [44]. The energy calibration of NCE protons is discussed in Appendix C of Ref. [44].

B. Event selection

To isolate a sample of $\bar{\nu}$ NCE events, a series of analysis cuts based on the physics processes and Monte Carlo studies was applied. The cuts (listed below) are various restrictions on the experimental variables, like PMT charge, time, or reconstructed energy, which differentiate the NCE events from other events:

- (1) Only one *subevent* to ensure selection of NC events with no decaying particles. A subevent is a cluster of at least ten tank hits with no more than ten ns between any two consecutive hits. A typical NCE interaction has only one subevent associated with the primary neutrino interaction.
- (2) Number of veto PMT hits less than 6. This cut excludes events that are entering or exiting the detector and register activity in the veto region. Cosmic rays and neutrino interactions in the material surrounding the detector with the outgoing nucleon entering the detector account for most of the events constrained by this selection cut. The veto cut removes almost all (99.9%) of the cosmic ray background. CCQE interactions in which the muon exits the detector before decaying are also excluded by this cut.
- (3) The reconstructed event time must occur within the neutrino beam time window.
- (4) Number of PMT tank hits greater than 12 to ensure that the event can be reliably reconstructed.




FIG. 4 (color online). Log-likelihood ratio between electron and proton event hypotheses for MC-generated NCE scattering events and beam unrelated data. Both histograms are normalized to unit area. Events with $\ln(\mathcal{L}_e/\mathcal{L}_p) < 0.42$ are selected for the analysis.

- (5) Reconstructed proton energy less than 650 MeV, above which the signal-to-background ratio decreases significantly.
- (6) A cut on the log-likelihood ratio between events reconstructed with a proton hypothesis and an electron hypothesis: $\ln(\mathcal{L}_e/\mathcal{L}_p) < 0.42$. This cut removes beam-unrelated (Michel) electrons from cosmic ray muon decays. Figure 4 shows the likelihood difference between events reconstructed under an electron and a proton hypothesis, for both Monte Carlo $\bar{\nu}$ NCE scattering events and beam-unrelated backgrounds (data).
- (7) Finally, a fiducial volume cut of 5 m. This cut ensures that the events in the sample are well reconstructed and well contained. It also reduces the neutrino events resulting from interaction with the dirt surrounding the detector.

A total of 60,605 events passes the analysis cuts, representing the largest $\bar{\nu}$ NCE candidate sample ever collected to date. Table I shows the results of a MC study to determine the efficiency and purity of the $\bar{\nu}$ NCE sample for each selection cut applied in the order shown. Figure 5 shows the reconstructed nucleon energy spectrum for NCE

TABLE I. Results from a MC study for $\bar{\nu}$ NCE efficiency and purity as a function of the selection cuts.

Selection cut	Efficiency (%)	Purity (%)
No cuts	100	0.2
1 subevent & veto hits < 6	59	2
Event in beam window	57	15
Tank hits > 12	55	16
Energy < 650 MeV	45	16
$\ln(\mathcal{L}_e/\mathcal{L}_p) < 0.42$	42	34
Fiducial cut $R < 5$ m	32	40




FIG. 5 (color online). Reconstructed nucleon kinetic energy spectra for the data and MC after the NCE event selection and a uniform fiducial volume cut of $R < 5$ m are applied. All MC distributions are normalized to the number of protons on target (POT).

events (data) along with the MC prediction of the sample composition, after subtracting beam-unrelated events and estimation of backgrounds (next section). After removing beam-unrelated events, the predicted fraction of $\bar{\nu}$ NCE scattering events in the sample is 48%. The remaining 52% of events is various backgrounds to this measurement. Neutrino induced interactions constitute 19% of the background. The next largest source of background is the so called “dirt events”(17%). These are neutrino interactions happening in the dirt just outside the detector with the recoil nucleon entering the detector without firing enough veto PMTs. Finally, there is a contribution from NCE-like events (14%), which are NC-pion producing events where the pion is absorbed in the target nucleus resulting in an event with a nucleon mimicking the neutrino NCE scattering signal.

C. Estimation of backgrounds

The various backgrounds to the MiniBooNE $\bar{\nu}$ mode NCE measurement are similar to those in the ν mode NCE measurement [7] with one notable exception: the neutrino induced events in the antineutrino mode beam. Data-driven methods are used to constrain all of the backgrounds as explained below.

As previously mentioned, the neutrino contamination in the antineutrino flux ($\sim 16\%$) is significantly larger than the corresponding antineutrino contamination in neutrino mode ($\sim 4\%$). HARP [33] did not cover all the phase space for π^+ production necessary to specify the neutrino background in the antineutrino beam. The neutrino contamination in the antineutrino mode beam was measured by MiniBooNE to constrain the flux outside of the region where data from the HARP experiment are available. Three independent and complimentary techniques are employed to measure the neutrino background utilizing the high-statistic neutrino mode cross section measurements made in

an almost pure neutrino beam. Briefly, the first technique exploits the difference in the angular distribution of outgoing muons in $\bar{\nu}$ and ν CCQE interactions to tag neutrino induced events. The second technique inferred the rate of ν induced events from a study of the charged current (CC) single pion production channel. The neutrino CC single pion interaction leads to a π^+ of which the decay into a muon is seen in the detector; however, the corresponding antineutrino CC single pion interaction produces a π^- that is absorbed in the detector medium most of the time [45]. The third technique to constrain the ν component of the beam exploits an external measurement of the rate of μ^- nuclear capture in ν CC interactions [46]. The results from the three techniques are consistent. For details on the first two techniques, see Ref. [8], and for the third technique, see Appendix A in Ref. [9]. Accordingly, a correction factor of 0.78 was applied to the MC prediction of the original neutrino flux (in the antineutrino mode) based on the HARP measurement. The antineutrino flux prediction was unchanged in this procedure. The cross section of neutrino induced NCE events in the antineutrino mode is inferred from the high-statistic neutrino mode ν NCE cross section measurement [7]. The total uncertainty in the estimation of the ν NCE background events in the sample is 14%.

The next major background is neutrino/antineutrino interactions occurring in the dirt surrounding the detector resulting in nucleons (mostly neutrons) which penetrate the detector without firing veto PMTs. The so-called dirt events are a significant fraction of the total background, particularly at low (below 300 MeV) energies. They are difficult to model as they result from interactions with various media outside the detector (the soil, detector support structures, etc.), of which the exact composition is not known. However, they have distinct kinematics and spatial distributions that can be used to constrain their contribution. Dirt events are mostly low in energy, preferentially reconstructed in the upstream part (the side facing the beam), and close to the edge of the detector. We use this kinematic information to select a sample of “dirt-enriched” events in both data and MC. A chi-square minimization is then employed to fit the MC prediction with a single scale factor to the observed data. The resulting fits show agreement across the three variables which were chosen: the reconstructed Z variable (axis along the beam direction), reconstructed R variable (radius), and reconstructed energy. Figure 6 shows a representative energy bin in a sample enriched in dirt events produced with the reconstructed Z variable. For details on the dirt measurement method used in MiniBooNE NCE analyses, see Appendix A in Ref. [7]. This background was separately measured in this $\bar{\nu}$ NCE analysis and ν NCE analysis [7], as the dirt background composition could be different in the two cases. The resultant dirt scaling factor of 0.62 was applied to the MC dirt prediction with an uncertainty of 10%. This may be compared to the scaling factor of 0.68 for the neutrino mode [7].




FIG. 6 (color online). Fit to the data using MC templates for the in-tank and dirt events in reconstructed Z variable for energy between 40 and 110 MeV. The “total MC before the fit” is a sum of the in-tank and dirt templates, which are absolutely (POT) normalized.

The final background is the $\bar{\nu}$ NCE-like events which are NC-pion events where the pion is absorbed within the nucleus, resulting in a final state identical to a $\bar{\nu}$ NCE scattering signal event. We rely on the MC prediction (together with the MiniBooNE measured neutral current π^0 cross section [47]) to estimate this background. The NUANCE cross section model assigns an error of 30% for pion absorption. Note that the $\bar{\nu}$ NCE-like background is the only background not directly measured in this analysis, and hence we report the calculated contribution which was subtracted to obtain our final antineutrino-nucleon NCE scattering cross section measurement (Fig. 7). Table II lists




FIG. 7 (color online). MiniBooNE $\bar{\nu}N \rightarrow \bar{\nu}N$ flux-integrated differential cross section on CH_2 . The uncertainty includes all errors—systematic and statistical. Also shown is the NCE-like background which has been subtracted from the reported cross section. The Monte Carlo predictions for the cross section with two different values of M_A —MiniBooNE ν CCQE measured value, $M_A = 1.35$ GeV (black) and $M_A = 1.02$ GeV, global average from deuterium based scattering experiments (dashed blue)—are shown for comparison.

TABLE II. The MiniBooNE $\bar{\nu}$ NCE analysis sample composition after backgrounds estimation.

Sample composition	Fraction (%)
$\bar{\nu}$ NCE (signal)	49
ν NCE (background)	19
Dirt events	17
$\bar{\nu}$ NCE-like	13
Others	2

the MiniBooNE $\bar{\nu}$ NCE analysis sample composition after estimation of backgrounds.

D. Antineutrino neutral-current elastic flux-averaged cross section

We report the $\bar{\nu}$ NCE scattering differential cross section as a function of the *measured* quasielastic momentum transfer ($Q_{QE(N)}^2$) from the total kinetic energy of scattered nucleons. This is to be distinguished from the *true* neutrino-nucleon four-momentum transferred (Q^2), a model-dependent quantity. This $Q_{QE(N)}^2$ is a well-defined experimental variable and was used for the ν NCE measurement [7]. It corresponds to the true Q^2 in the simple case of a static target nucleon (no Fermi motion) and no FSI.

From our studies of FSI, the total kinetic energy of all outgoing nucleons is fairly independent of the underlying nuclear model and is a better variable to report than the energy of the leading nucleon, for example. In addition, the total energy is a better experimental variable to report with the calorimetric measurement that MiniBooNE detector makes for this channel. With the qualifications explained above, the expression for measured neutrino-nucleon four-momentum transferred is

$$Q_{QE(N)}^2 := 2m_N T = 2m_N \sum_i T_i, \quad (1)$$

where m_N is the nucleon mass and T is the sum of the kinetic energies of the final state nucleons. The sum T_i is used in the definition due to the calorimetric nature of the MiniBooNE measurement and is more inclusive with respect to possible nuclear effects as compared to track-based reconstruction used in the SciBooNE experiment [48] or the BNL E734 experiment [25].

The constrained backgrounds—the beam unrelated background, the dirt background, and the neutrino induced backgrounds—are subtracted from the reconstructed energy spectrum for data. The $\bar{\nu}$ NCE-like background is removed by doing a bin-by-bin multiplication of the data spectrum by the signal fraction, i.e. the ratio of the number of $\bar{\nu}$ NCE events to the total number of antineutrino induced in-tank events, based on the MC prediction. Finally, a Bayesian unfolding procedure [49,50] is used to correct the background subtracted data for limited detector resolution, mis-reconstruction, and sources of detector inefficiency.

The flux-averaged $\bar{\nu}$ NCE scattering differential cross section is extracted as per the formula

$$\frac{d\sigma_i^{\bar{\nu}NCE}}{dQ_{QE(N)}^2} = \frac{\sum_j U_{ij}(d_j - D_j - V_j - N_j) \frac{S_j}{S_j + B_j}}{\epsilon_i \cdot (2M_N \Delta T) \cdot N^{tar} \cdot N^{POT} \cdot \Phi_{\bar{\nu}}}, \quad (2)$$

where U_{ij} is the unfolding matrix, the index j labels the reconstructed energy bin, and i labels the unfolded true energy bin (as per the MC prediction). In the above equation, d_j represents data; D_j , V_j , and N_j are the data-driven corrected backgrounds of dirt events, neutrino induced events, and beam unrelated events respectively; S_j is the MC predicted number of $\bar{\nu}$ NCE scattering events; B_j is the rest of the backgrounds which mostly consist of the $\bar{\nu}$ NCE-like events; ϵ is the efficiency; ΔT is the bin width; N^{tar} is the number of nucleons in the detector; N^{POT} is the number of protons on target corresponding to the data set; and $\Phi_{\bar{\nu}}$ is the total integrated antineutrino flux within the energy range 0 to 10 GeV (both $\bar{\nu}_\mu$ and $\bar{\nu}_e$).

The unfolding matrix U_{ij} is calculated from the predicted correlation between reconstructed nucleon energy and true nucleon kinetic energy (the sum of the kinetic energies of all nucleons in the final state) resulting in a well-behaved, but biased, solution. The error due to the bias in the unfolding procedure is estimated by employing an iterative method where each successive unfolded spectrum is used as the true energy spectrum for the next iteration. The resulting spread in the cross section measurement, from the first iteration to the last one when it converges, is the error in the unfolding procedure. The details of the unfolding procedure and the estimation of the associated error with it can be found in Refs. [44,51].

The resulting flux-averaged $\bar{\nu}$ NCE scattering differential cross section is shown in Fig. 7. Also shown is the $\bar{\nu}$ NCE-like background which was subtracted from the total $\bar{\nu}$ NCE-like cross section. Though the systematic uncertainties are higher in the lowest energy bins, as $Q_{QE(N)}^2$ approaches 0, the “rollover” associated with the binding energy of the carbon nucleus is clearly seen for the first time.

Since the MiniBooNE target is mineral oil (CH_2), the measured $\bar{\nu}$ NCE scattering is a sum of three different processes: scattering on free protons in hydrogen, bound protons in carbon, and bound neutrons in carbon. The contribution of these individual processes to the total cross section is discussed in Appendix A. Integrating over 0.033 to 1.655 GeV² Q_{QE}^2 bins, the total $\bar{\nu}$ NCE scattering cross section per nucleon is $(5.06 \pm 0.990) \times 10^{-40}$ cm².

The various uncertainties in the $\bar{\nu}$ NCE cross section measurement are listed in Table III. The flux error encompasses the uncertainties in the pion propagation and decay in the BNB. The cross section error includes the uncertainties in the cross section model of the various background processes. The error associated with the detector electronics, the PMT response, and the uncertainty in

TABLE III. The total integrated normalization error in the MiniBooNE $\bar{\nu}$ NCE scattering cross section measurement along with the key individual error contributions.

Error	Value (%)
Statistical	4.5
Flux uncertainty	5.8
Cross section uncertainty (background processes)	3.0
Detector effects	14.5
Estimation of ν induced events	3.6
Estimation of dirt events	1.7
Unfolding error	6.8
<i>Total</i>	19.5

modeling the production and propagation of optical photons within the detector medium contribute to the detector error. The next two errors are from the uncertainty in the measurement of neutrino induced NCE events and dirt events. The final error is due to the unfolding procedure.

For each systematic uncertainty, there is an associated error matrix that encompasses the information about the parameters describing the particular physical process, the uncertainties in the parameters, and any correlation among them. The error matrices are added in quadrature to obtain the total error matrix, and Table III lists the normalization error (sum of the diagonal elements of the error matrix) for both the individual errors and the total error.

E. Ratio of antineutrino neutral-current elastic to neutrino neutral-current elastic cross-section measurement

Both the neutrino-nucleus NCE scattering cross section [7] and the antineutrino-nucleus NCE cross section reported here represent the largest sample of such events ever collected to date. Since both measurements were made in the same beamline and with the same detector, we expect a bin-by-bin ratio of the two cross section measurements would cancel the common systematic errors. The resulting cross section ratio plot encompasses information from both the neutrino and antineutrino NCE scattering cross sections while minimizing the errors. However, it should be noted that $Q_{QE(N)}^2$ is sensitive to the neutrino flux and the two measurements are made in the same beamline but with opposite horn polarities, resulting in nonidentical flux spectra. One of the main motivations for measurement of this cross section is to better understand and model neutrino nucleus interactions. We believe that such a ratio measurement where the errors are carefully accounted for would aid the theoretical physics community to test various models.

The data set for the ratio measurement consists of the entire neutrino mode and antineutrino mode NCE scattering cross section data from MiniBooNE. This consists of 94,531 ν NCE candidate events and 60,605 $\bar{\nu}$ NCE candidate events that pass selection cuts.




FIG. 8 (color online). Ratio of the antineutrino to neutrino NCE scattering cross section in MiniBooNE with total error. Also plotted are the predicted ratios from MC simulations with $M_A = 1.02$ GeV and $M_A = 1.35$ GeV—determined by the MiniBooNE ν CCQE measurement ($M_A^{\nu\text{CCQE}}$).

The systematic error for the ratio measurement was evaluated by dividing the errors into two types: *correlated* errors and *uncorrelated* errors. The correlated systematic errors are common to both ν NCE and $\bar{\nu}$ NCE scattering measurements. Since both measurements are made using the same detector and have the same observed final state, the detector systematic errors—the uncertainty in the optical photon production and propagation, the error associated with the detector electronics, and the error associated with the PMT response—are categorized as correlated errors. The uncorrelated errors include the error associated with the measurement of the dirt background, the error in the measurement of the neutrino component of the antineutrino beam, and the error accrued due to the bias in the unfolding procedure implemented. The resulting $\bar{\nu}$ NCE to ν NCE cross section ratio measurement is shown in Fig 8. The error bars represent the total normalization error due to both systematic and statistical errors. The total uncertainty in the $\bar{\nu}$ NCE to ν NCE cross section ratio measurement is about 20%.

F. Ratio of antineutrino neutral-current elastic to antineutrino charged-current quasielastic cross-section measurement

We also report a $\bar{\nu}$ NCE-to- $\bar{\nu}$ CCQE scattering ratio measurement as a function of $Q_{QE(N/\mu^+)}^2$. MiniBooNE has previously reported this ratio in neutrino mode (ν NCE-to- ν CCQE ratio) in Ref. [7]. Although there are significant differences between the extraction of Q_{QE}^2 for the individual cross sections, this measurement is a ratio of two well-defined experimental quantities. As was the case for NCE, the expression used for $Q_{QE(\mu^+)}^2$ [2] assumes that the muon is recoiling from a stationary nucleus. Corrections for that are model dependent and not provided here. Instead, the various models can be compared to this




FIG. 9 (color online). MiniBooNE $\bar{\nu}$ NCE/ $\bar{\nu}$ CCQE cross section ratio on CH_2 as a function of $Q^2_{QE(N/\mu^+)}$. Also shown are the MC prediction for the ratio with $M_A = 1.02$ GeV as well as that determined by the MiniBooNE ν CCQE measurement ($M_A^{\nu\text{CCQE}} = 1.35$ GeV). The individual $\bar{\nu}$ NCE and $\bar{\nu}$ CCQE cross sections are per target nucleon—there are 14/8 times more target nucleons in the numerator than in the denominator. The error bars include both statistical and systematic errors (except the flux errors) taken in quadrature. The corresponding ν NCE/ ν CCQE cross section ratio has been reported in Ref. [7].

experimentally defined ratio with reduced correlated experimental errors, for a sensitive test of the underlying nuclear model.

In the $\bar{\nu}$ NCE/ $\bar{\nu}$ CCQE scattering cross section ratio (Fig. 9), the uncertainties in the flux estimation are assumed to cancel out whereas other errors have been added in quadrature.

IV. SUMMARY

To summarize, using a high-statistics sample of $\bar{\nu}$ NCE scattering interactions collected by the MiniBooNE experiment, the $\bar{\nu}$ NCE ($\bar{\nu}N \rightarrow \bar{\nu}N$) flux-averaged differential cross section, $d\sigma/dQ^2$, on CH_2 was measured. The $\bar{\nu}$ NCE cross section (Fig. 7) shows good agreement to a simple RFG model with $M_A = 1.35$ GeV—determined by the MiniBooNE ν CCQE measurement [2]. This is interesting as it shows that a simple tuning of M_A , presumably to effectively handle more complex nuclear effects, provides a reasonable description.

For the first time, an antineutrino-to-neutrino NCE scattering cross section ratio has been reported, that accounts for all the systematic errors common to both measurements. Finally, the $\bar{\nu}$ NCE-to- $\bar{\nu}$ CCQE cross section ratio is provided. The corresponding neutrino mode ratio (ν NCE to ν CCQE) was reported in Ref. [7], facilitating a comparison between the two modes.

Any other models designed to explain the MiniBooNE ν CCQE data [10–15] need to consider the entire MiniBooNE data set (ν CCQE, $\bar{\nu}$ CCQE, ν NCE, and $\bar{\nu}$ NCE) to be considered complete.

ACKNOWLEDGMENTS

The authors would like to acknowledge the support of Fermilab, the Department of Energy, and the National Science Foundation in the construction, operation, and data analysis of the Mini Booster Neutrino Experiment.

APPENDIX A: MINIBOOONE ANTINEUTRINO NEUTRAL-CURRENT ELASTIC CROSS-SECTION DISCUSSION

The antineutrino-nucleon NCE scattering cross section reported here is in terms of $Q^2_{QE(N)}$ which in MiniBooNE is proportional to the total kinetic energies of all the final state nucleons that are produced in the interaction. Also in MiniBooNE, NCE scattering on protons is indistinguishable from NCE scattering on neutrons as the neutrons are seen only via their subsequent strong interaction with protons. The MiniBooNE target is mineral oil (CH_2), hence the scattering is off of both bound nucleons (in carbon) and free nucleons (in hydrogen). In fact, the cross section is a sum of three different processes: the antineutrino scattering off free protons in the hydrogen atom, the bound protons in the carbon atom, and the bound neutrons in the carbon atom. Each of the individual processes have different efficiencies in the MiniBooNE detector. Figure 10 shows the efficiency correction functions $C_{\bar{\nu}p,H}$, $C_{\bar{\nu}p,C}$, and $C_{\bar{\nu}n,C}$ for the three processes. The efficiency correction is defined as the ratio of the efficiency for a particular type of $\bar{\nu}$ NCE event to the average efficiency for all $\bar{\nu}$ NCE events as a function of $Q^2_{QE(N)}$. Therefore, the flux-averaged $\bar{\nu}$ NCE differential cross section on CH_2 shown in Fig. 7 can be expressed as

$$\frac{d\sigma_{\bar{\nu}N \rightarrow \bar{\nu}N}}{dQ^2} = \frac{1}{7} C_{\bar{\nu}p,H}(Q^2_{QE(N)}) \frac{d\sigma_{\bar{\nu}p \rightarrow \bar{\nu}p,H}}{dQ^2} + \frac{3}{7} C_{\bar{\nu}p,C}(Q^2_{QE(N)}) \times \frac{d\sigma_{\bar{\nu}p \rightarrow \bar{\nu}p,C}}{dQ^2} + \frac{3}{7} C_{\bar{\nu}n,C}(Q^2_{QE(N)}) \frac{d\sigma_{\bar{\nu}n \rightarrow \bar{\nu}n,C}}{dQ^2},$$




FIG. 10 (color online). The efficiency corrections for the different processes (as labeled) contributing to the MiniBooNE $\bar{\nu}$ NCE scattering cross section, as a function of $Q^2_{QE(N)}$.

where $d\sigma_{\bar{\nu}p \rightarrow \bar{\nu}p,H}/dQ^2$ is the $\bar{\nu}$ NCE cross section on free protons (per free proton), $d\sigma_{\bar{\nu}p \rightarrow \bar{\nu}p,C}/dQ^2$ is the $\bar{\nu}$ NCE cross section on bound protons (per bound proton), and $d\sigma_{\bar{\nu}n \rightarrow \bar{\nu}n,C}/dQ^2$ is the $\bar{\nu}$ NCE cross section on bound neutrons (per bound neutron). The efficiency corrections should be applied to the predicted cross sections of the individual processes in order to compare with the MiniBooNE $\bar{\nu}$ NCE scattering cross section result.

APPENDIX B: TABLES

Here we tabulate the results presented in this paper. Table VI lists the $\bar{\nu}$ NCE differential cross section, the $\bar{\nu}$ NCE-like background (as shown in Fig. 7), and the correction coefficients, in bins of $Q_{QE(N)}^2$. A data release page is also available in Ref. [52] where the experimentally reconstructed nucleon scattering energy spectrum, the

TABLE IV. MiniBooNE measured $\bar{\nu}$ NCE/ $\bar{\nu}$ CCQE cross section ratio as a function of $Q_{QE(N/\mu^+)}^2$.

$Q_{QE(N/\mu^+)}^2$ (GeV ²)distribution	$\frac{\sigma_{\bar{\nu}NCE}^j}{\sigma_{\bar{\nu}CCQE}^j}$
0.100–0.150	0.245 ± 0.013
0.150–0.200	0.248 ± 0.014
0.200–0.250	0.249 ± 0.017
0.250–0.300	0.264 ± 0.020
0.300–0.350	0.275 ± 0.024
0.350–0.400	0.272 ± 0.027
0.400–0.450	0.277 ± 0.033
0.450–0.500	0.275 ± 0.037
0.500–0.600	0.269 ± 0.045
0.600–0.700	0.281 ± 0.059
0.700–0.800	0.284 ± 0.072
0.800–1.000	0.294 ± 0.086
1.000–1.200	0.391 ± 0.139
1.200–1.500	0.535 ± 0.276

TABLE V. MiniBooNE $\bar{\nu}$ NCE/ ν NCE scattering cross section ratio measured as a function of $Q_{QE(N)}^2 = 2m_N \sum_i T_i$.

$Q_{QE(N)}^2$ (GeV ²)distribution	$\frac{\sigma_{\bar{\nu}NCE}^j}{\sigma_{\nu NCE}^j}$
0.067–0.135	0.555 ± 0.0371
0.135–0.202	0.473 ± 0.0304
0.202–0.270	0.393 ± 0.0257
0.270–0.337	0.344 ± 0.0230
0.337–0.405	0.300 ± 0.0232
0.405–0.472	0.265 ± 0.0210
0.472–0.540	0.228 ± 0.0183
0.540–0.608	0.202 ± 0.0189
0.608–0.675	0.184 ± 0.0193
0.675–0.743	0.170 ± 0.0210
0.743–0.810	0.160 ± 0.0225
0.810–0.878	0.155 ± 0.0254
0.878–0.945	0.148 ± 0.0269
0.945–1.013	0.149 ± 0.0335
1.013–1.080	0.159 ± 0.0310
1.080–1.148	0.157 ± 0.0370
1.148–1.216	0.151 ± 0.0385
1.216–1.283	0.144 ± 0.0329
1.283–1.351	0.138 ± 0.0321
1.351–1.418	0.139 ± 0.0329
1.418–1.486	0.132 ± 0.0325
1.486–1.553	0.132 ± 0.0342
1.553–1.621	0.141 ± 0.0406
1.621–1.689	0.136 ± 0.0434

TABLE VI. MiniBooNE measured $\bar{\nu}$ NCE differential cross section, predicted $\bar{\nu}$ NCE-like background, and predicted correction coefficients for the three different $\bar{\nu}$ NCE scattering contributions as a function of $Q_{E(N)}^2 = 2m_N \sum_i T_i$.

$Q_{E(N)}^2$ (GeV ²)distribution	$\bar{\nu}$ NCE cross section, (cm ² /GeV ²)	$\bar{\nu}$ NCE-like background, (cm ² /GeV ²)	$C_{\bar{\nu}p,H}$	$C_{\bar{\nu}p,C}$	$C_{\bar{\nu}n,C}$
0.033–0.100	$(9.700 \pm 0.791) \times 10^{-40}$	1.048×10^{-41}	0.484	1.017	1.229
0.100–0.168	$(1.646 \pm 0.119) \times 10^{-39}$	2.931×10^{-41}	1.134	1.128	0.868
0.168–0.235	$(1.225 \pm 0.084) \times 10^{-39}$	3.094×10^{-41}	1.055	1.050	0.944
0.235–0.303	$(9.039 \pm 0.724) \times 10^{-40}$	3.365×10^{-41}	1.028	1.018	0.979
0.303–0.370	$(6.749 \pm 0.661) \times 10^{-40}$	4.288×10^{-41}	0.998	0.997	1.002
0.370–0.438	$(5.053 \pm 0.584) \times 10^{-40}$	5.623×10^{-41}	0.972	0.984	1.018
0.438–0.506	$(3.863 \pm 0.461) \times 10^{-40}$	7.192×10^{-41}	1.001	0.981	1.019
0.506–0.573	$(2.963 \pm 0.416) \times 10^{-40}$	8.520×10^{-41}	0.979	0.983	1.020
0.573–0.641	$(2.292 \pm 0.361) \times 10^{-40}$	9.154×10^{-41}	0.986	0.980	1.022
0.641–0.708	$(1.799 \pm 0.314) \times 10^{-40}$	9.245×10^{-41}	0.981	0.985	1.018
0.708–0.776	$(1.409 \pm 0.274) \times 10^{-40}$	8.970×10^{-41}	0.997	0.985	1.016
0.776–0.844	$(1.111 \pm 0.235) \times 10^{-40}$	8.260×10^{-41}	0.969	0.980	1.025
0.844–0.911	$(8.747 \pm 2.088) \times 10^{-41}$	7.278×10^{-41}	0.966	0.969	1.038
0.911–0.979	$(7.061 \pm 1.681) \times 10^{-41}$	6.289×10^{-41}	0.936	0.973	1.036
0.979–1.046	$(5.833 \pm 1.079) \times 10^{-41}$	5.392×10^{-41}	0.979	0.983	1.022
1.046–1.114	$(4.711 \pm 0.989) \times 10^{-41}$	4.316×10^{-41}	0.977	0.980	1.025
1.114–1.181	$(3.883 \pm 1.139) \times 10^{-41}$	3.543×10^{-41}	0.904	0.976	1.035
1.181–1.249	$(3.101 \pm 0.726) \times 10^{-41}$	2.846×10^{-41}	0.736	0.893	1.152
1.249–1.317	$(2.513 \pm 0.482) \times 10^{-41}$	2.366×10^{-41}	0.685	0.865	1.192
1.317–1.384	$(2.060 \pm 0.384) \times 10^{-41}$	1.976×10^{-41}	0.464	0.867	1.202
1.384–1.452	$(1.720 \pm 0.294) \times 10^{-41}$	1.657×10^{-41}	0.386	0.851	1.241
1.452–1.519	$(1.391 \pm 0.279) \times 10^{-41}$	1.386×10^{-41}	0.323	0.811	1.286
1.519–1.587	$(1.165 \pm 0.315) \times 10^{-41}$	1.145×10^{-41}	0.526	0.840	1.233
1.587–1.655	$(1.024 \pm 0.378) \times 10^{-41}$	9.840×10^{-42}	0.240	0.854	1.222

migration matrices from reconstructed to true energy (for the bound nucleons in carbon, free protons in hydrogen, and the $\bar{\nu}$ NCE-like background), the detector efficiency, and the total error matrix is reported.

Table V quantifies the antineutrino NCE-to-neutrino NCE scattering cross section ratio measurement shown in Fig. 8. And Table IV lists the $\bar{\nu}$ NCE-to- $\bar{\nu}$ CCQE differential cross section ratio measurement shown in Fig. 9.

-
- [1] J. A. Formaggio and G. P. Zeller, *Rev. Mod. Phys.* **84**, 1307 (2012).
- [2] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **81**, 092005 (2010).
- [3] R. A. Smith and E. J. Moniz, *Nucl. Phys.* **B43**, 605 (1972); **101**, 547(E) (1975).
- [4] J. Carlson, J. Jourdan, R. Schiavilla, and I. Sick, *Phys. Rev. C* **65**, 024002 (2002).
- [5] O. Benhar, N. Farina, H. Nakamura, M. Sakuda, and R. Seki, *Phys. Rev. D* **72**, 053005 (2005).
- [6] J. E. Amaro, M. B. Barbaro, J. A. Caballero, T. W. Donnelly, and C. Maieron, *Phys. Rev. C* **71**, 065501 (2005).
- [7] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **82**, 092005 (2010).
- [8] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **84**, 072005 (2011).
- [9] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **88**, 032001 (2013).
- [10] J. E. Amaro, M. B. Barbaro, J. A. Caballero, T. W. Donnelly, and J. M. Udías, *Phys. Rev. D* **84**, 033004 (2011).
- [11] A. Bodek, H. S. Budd, and M. E. Christy, *Eur. Phys. J. C* **71**, 1726 (2011).
- [12] C. Giusti and A. Meucci, arXiv:1110.4005.
- [13] M. Martini *et al.*, *Phys. Rev. C* **84**, 055502 (2011).
- [14] J. Nieves, I. R. Simo, and M. J. V. Vacas, *J. Phys. Conf. Ser.* **408**, 012040 (2013).
- [15] J. T. Sobczyk, *Eur. Phys. J. C* **72**, 1850 (2012).
- [16] J. E. Amaro, M. Barbaro, J. Caballero, and T. Donnelly, *Phys. Rev. C* **73**, 035503 (2006).
- [17] O. Benhar and G. Veneziano, *Phys. Lett. B* **702**, 433 (2011).
- [18] A. Meucci, C. Giusti, and F. D. Pacati, *Phys. Rev. D* **84**, 113003 (2011).
- [19] A. V. Butkevich and D. Perevalov, *Phys. Rev. C* **84**, 015501 (2011).
- [20] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **103**, 081801 (2009).

- [21] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **83**, 052009 (2011).
- [22] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **81**, 013005 (2010).
- [23] D. Cline, A. Entenberg, W. Kozanecki, A. Mann, D. Reeder, C. Rubbia, J. Strait, L. Sulak, and H. Williams, *Phys. Rev. Lett.* **37**, 252 (1976).
- [24] A. Entenberg, J. Horstkotte, W. Kozanecki, A. Mann, C. Rubbia, J. Strait, L. Sulak, P. Wanderer, and H. Williams, *Phys. Rev. Lett.* **42**, 1198 (1979).
- [25] L. Ahrens *et al.*, *Phys. Rev. D* **35**, 785 (1987).
- [26] C. Athanassopoulos *et al.* (LSND), *Phys. Rev. Lett.* **81**, 1774 (1998).
- [27] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **110**, 161801 (2013).
- [28] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **79**, 072002 (2009).
- [29] J. L. Raaf, A. O. Bazarko, G. T. Garvey, E. A. Hawker, R. A. Johnson, G. B. Mills, A. Pla-Dalmau, and R. Tayloe, *IEEE Trans. Nucl. Sci.* **49**, 957 (2002).
- [30] S. Brice *et al.* (MiniBooNE Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **562**, 97 (2006).
- [31] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **599**, 28 (2009).
- [32] S. Agostinelli *et al.* (GEANT4), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [33] M. G. Catanesi *et al.*, *Eur. Phys. J. C* **52**, 29 (2007).
- [34] MiniBooNE flux data release, http://www-boone.fnal.gov/for_physicists/data_release/flux/.
- [35] D. Casper, *Nucl. Phys. B, Proc. Suppl.* **112**, 161 (2002).
- [36] C. Llewellyn Smith, *Phys. Rep.* **3**, 261 (1972).
- [37] D. Rein and L. M. Sehgal, *Nucl. Phys.* **B223**, 29 (1983).
- [38] A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. Lett.* **100**, 032301 (2008).
- [39] T. Katori, Report No. FERMILAB-THESIS-2008-64, 2008.
- [40] R. Brun *et al.*, “GEANT 3”, CERN DD/EE/84-1 (1987).
- [41] A. Ferrari *et al.*, Report No. CERN-2005-010, 2005.
- [42] J. D. Jackson, *Classical Electrodynamics Third Edition*, 3rd ed. (Wiley, New York, 1998).
- [43] R. B. Patterson, E. M. Laird, Y. Liu, P. D. Meyers, I. Stancu, and H. A. Tanaka, *Nucl. Instrum. Methods Phys. Res., Sect. A* **608**, 206 (2009).
- [44] D. Perevalov, Report No. FERMILAB-THESIS-2009-47, 2009.
- [45] A. Shinohara, T. Muroyama, J. Shintai, J. Kurachi, M. Furukawa, T. Miura, Y. Yoshimura, T. Saito, T. Ohdaira, and N. Imanishi, *Phys. Rev. A* **53**, 130 (1996).
- [46] T. Suzuki, D. Measday, and J. Roalsvig, *Phys. Rev. C* **35**, 2212 (1987).
- [47] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), *Phys. Rev. D* **81**, 013005 (2010).
- [48] H. Takei, Report No. FERMILAB-THESIS-2009-19, 2009.
- [49] G. Cowan, *Statistical data analysis* (Oxford University Press, Oxford, 1998).
- [50] G. D’Agostini, *Nucl. Instrum. Methods Phys. Res., Sect. A* **362**, 487 (1995).
- [51] R. Dharmapalan, Report No. FERMILAB-THESIS-2012-29, 2012.
- [52] MiniBooNE antineutrino NCE data release, http://www-boone.fnal.gov/for_physicists/data_release/nce_l_nubar.