

**Consistent use of effective potentials**Anders Andreassen,<sup>\*</sup> William Frost,<sup>†</sup> and Matthew D. Schwartz<sup>‡</sup>*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*

(Received 28 August 2014; published 28 January 2015)

It is well known that effective potentials can be gauge dependent while their values at extrema should be gauge invariant. Unfortunately, establishing this invariance in perturbation theory is not straightforward, since contributions from arbitrarily high-order loops can be of the same size. We show in massless scalar QED that an infinite class of loops can be summed (and must be summed) to give a gauge-invariant value for the potential at its minimum. In addition, we show that the exact potential depends on both the scale at which it is calculated and the normalization of the fields, but the vacuum energy does not. Using these insights, we propose a method to extract some physical quantities from effective potentials which is self-consistent order by order in perturbation theory, including improvement with the renormalization group.

DOI: 10.1103/PhysRevD.91.016009

PACS numbers: 11.30.Qc, 12.15.Lk

**I. INTRODUCTION**

Effective actions provide a powerful organizing framework for quantum field theory. Often one computes an effective action by integrating out a massive particle, usually at fixed order in perturbation theory as in the four-Fermi theory, but sometimes to all orders in couplings but leading order in  $\frac{1}{m^2}$ , as in the Euler-Heisenberg Lagrangian [1,2]. In these cases, the effective action is an action just like the original one, but with a subset of its fields. Thus it can be used for the same calculations, such as  $S$ -matrix elements, in the same way. An advantage of the effective action approach is that it is often easier to expand in  $\frac{1}{m^2}$  and then calculate the physical quantity than to perform the calculation before the expansion.

A more subtle use of effective actions occurs when one integrates out *all* the particles. Integrating out everything of course just produces a number (usually  $\infty$ ). To get functional dependence one can instead integrate out all the particles in the presence of classical sources  $J$  for each to get a functional  $W[J]$  of those sources. The Legendre transform of  $W[J]$  gives an effective action  $\Gamma[\phi] = W[J] - J\phi$ , which we call the one-particle irreducible (1PI) effective action. In the 1PI effective action, the fields are classical background fields. One cannot perform loops using  $\Gamma[\phi]$  since all the loops have already been performed in integrating out the particles.  $\Gamma[\phi]$  nevertheless encodes all the classical and quantum physics of the original theory. This  $\Gamma[\phi]$  is therefore an extremely powerful object. It is also extremely difficult to compute.

These two types of effective action are of course closely related; for example, the 1PI effective action for QED reduces to the Euler-Heisenberg Lagrangian when restricted to a single electron loop and constant

electromagnetic fields. It is nevertheless important to understand that the 1PI effective action does *not* function like the kind of integral over a local Lagrangian one usually uses in quantum field theory.  $\Gamma[\phi]$  describes fluctuations in the presence of background sources, which are normally absent. The fields  $\phi$  on which  $\Gamma$  depends are classical fields—functions of space-time, not quantum operators acting on a Hilbert space. The value of the action at generic classical field values  $\phi$  has no clear physical meaning. Indeed, its value depends on the way the calculation is performed: what gauge is chosen, the normalization of the fields, and even the scale at which the action is computed (see Sec. II below). On-shell quantities, such as  $S$ -matrix elements computed from the effective action do not depend on these conventions. An important property of the effective action, or the effective potential to which it reduces for constant fields, is that its minimum gives the energy of the true ground state of a theory. For this reason, effective actions provide unrivaled insight into spontaneous symmetry breaking [3,4].

Despite the unphysical nature of the effective action at generic field values, its gauge dependence has elicited a surprising amount of consternation. Since the gauge dependence was first observed 40 years ago [5], papers are regularly written claiming that the effective potential should be defined in some other way, often through a field redefinition, so that it is manifestly gauge invariant. The easiest way to make the potential gauge invariant is simply to pick a gauge: in the mid 1970s, Dolan and Jackiw argued that only unitary gauge ( $\xi = \infty$ ) makes sense [6] while S. Weinberg insisted upon Landau gauge [7] ( $\xi = 0$ ). Around the same time, Frere and Nicoletopoulos (based on observations of Fischler and Brout [8]) argued that one should use dressed fields which have no field-strength renormalization [9], which is similar to Yennie gauge ( $\xi = 3$ ). In 1998, Tye and Vtorov-Karevsky argued that one should use an exact nonlinear field redefinition, replacing a scalar doublet  $\phi_1 + i\phi_2$  by a linear sigma model  $\sigma e^{i\pi}$  [10]. Others

<sup>\*</sup>anders@physics.harvard.edu<sup>†</sup>wfrost@physics.harvard.edu<sup>‡</sup>schwartz@physics.harvard.edu

have tried using composite fields, such as  $\phi^*\phi$  [11], using a displacement operator formalism [12,13], working in a Hamiltonian formulation [14] or the Vikovisky-DeWitt [15] formalism, where the introduction of gauge invariance (and therefore gauge noninvariance) is sidestepped completely. In 2014, Nielsen [16] proposed a nonlinear, but perturbative, field redefinition determined by solving a differential equation with a boundary condition at  $\xi = \xi_0$  to remove the dependence on  $\xi$ .

The periodic frustration with the gauge dependence of the effective potential seems to us somewhat misguided. Gauge invariance of unphysical quantities is commonplace. One is not normally bothered by the  $\xi$  dependence of the photon two-point function or the anomalous dimension of a charged scalar field. Similarly, there is no reason to be dismayed about the gauge dependence of the effective action. While field redefinitions can treat a symptom of the unphysical nature of the potential, they cannot make it any more physical. In fact, none of the papers cited above seem to address the incongruity of both using the freedom in quantum field theory to perform field redefinitions and claiming that a particular field redefinition is preferred. Of course, physical quantities, such as on-shell  $S$ -matrix elements must be gauge invariant. Furthermore, gauge invariance provides a very useful theoretical consistency check. Not only does it help verify that you did the relevant loop calculations correctly, but it can also verify that the quantity you are calculating could possibly be physical. Thus, for an object as unintuitive as the effective potential, gauge dependence is in many ways actually desirable.

With these observations in mind, let us consider how the gauge invariance of some physical quantities calculated from the effective potential have been established in perturbation theory and beyond. Our focus in this paper is on scalar QED at zero temperature. With a negative mass squared for the scalar this model is called the Abelian Higgs model. With zero mass we call it the Coleman-Weinberg model [7]. General nonperturbative demonstrations that certain quantities calculated from the effective potential in these models should be gauge invariant were provided by Nielsen [17] and independently by Fukuda and Kugo [18]. Quantities which are expected to be gauge invariant in these models include the value of the potential at its extrema and the pole masses of the scalar and vector bosons in the spontaneously broken phase. We review some of the main results from these papers in Secs. II and III.

Although all of our new results pertain to the Coleman-Weinberg model, we devote Sec. III to the Abelian Higgs model since it illustrates a number of aspects of gauge invariance not present when  $m = 0$ . We try to provide a useful summary of this model since the literature is scattered and somewhat inconsistent. Also, issues which are not present in the Coleman-Weinberg model, such as an unphysical gauge-dependent minimum of the effective potential and the presence of an additional dimensionful

scale, provide a broader picture of some subtleties associated with gauge invariance.

The main result of this paper concerns the extrema of the Coleman-Weinberg effective potential. We demonstrate that the minimum of the renormalization-group-improved potential in this model is gauge invariant order by order in perturbation theory. To show this requires first an understanding of the appropriate perturbation expansion. Perturbation theory must respect the relation  $\lambda \sim \hbar e^4$  which determines the minimum at one loop (see Sec. IV). We calculate all the two-loop contributions in  $R_\xi$  gauges proportional to  $\hbar^2 e^6$  (Sec. V), as well as all the terms like  $\hbar^{n+2} e^6 \frac{e^{4n}}{\lambda^n}$  coming from three-loop and higher-order graphs (Sec. VI) which also contribute at order  $\hbar^2 e^6$  when  $\lambda \sim \hbar e^4$ . Demonstrating gauge invariance in perturbation theory also requires careful consideration of what dimensionful scale the potential at the minimum should be expressed in terms of. The most obvious dimensionful scale  $v = \langle \phi \rangle$  is a questionable candidate since it is gauge dependent itself. These issues are discussed in Sec. VII, where we show explicitly that the instability scale  $\Lambda_I$  is defined so that  $V(\Lambda_I) = 0$  depends on  $\xi$ .

Next we augment our perturbative calculation with a proposal for how perturbation theory in which gauge invariance holds order by order can be used in a resummed effective potential. The challenge is that, first, resummation breaks the strict  $\lambda \sim \hbar e^4$  power counting required for gauge invariance and, second, that the renormalization-group kernel for the field-strength renormalization,  $\Gamma = \int \gamma d \ln \mu$  has gauge dependence to all orders in any expansion. Our solution exploits the invariance of physical quantities in the effective potential to the scale where the potential is calculated: we simply run the couplings towards the minimum before calculating the potential in perturbation theory. Then the field-strength renormalization plays no direct role.

Although the potential at the minimum is gauge invariant, field values are gauge dependent. For example, the expectation value of the field,  $v = \langle \phi \rangle$ , depends on  $\xi$  [cf. Eq. (3.18)] as does the *instability scale*  $\Lambda_I$ , defined as the field value where  $V(\Lambda_I) = 0$  [cf. Eq. (7.10)]. Although the gauge dependence of field values is often emphasized [17,19–21] there are still a surprising number of papers which try to ascribe physical significance to values of  $\phi$ . Sometimes the physical question one wants to answer is akin to when higher-dimension operators, for example from Planck-scale physics, can significantly affect the potential. We present one gauge-invariant way to deal with these higher-dimension operators in Sec. IX. Our conclusions and implications of our work are discussed in Sec. X.

## II. EFFECTIVE ACTIONS

We begin with a rapid introduction to effective actions and their gauge dependence. In this section,  $\phi$  generically

denotes all the fields in a given theory (in the rest of the paper  $\phi$  will denote a specific field in scalar QED).

A very useful object in quantum field theory is the 1PI effective action  $\Gamma[\phi]$ . When  $\phi$  is independent of  $x$ , the effective action reduces to the effective potential

$$\Gamma[\phi] = - \int d^4x V(\phi) + \text{derivative and nonlocal terms.} \quad (2.1)$$

The interactions in the effective action embody a sum over all of the off-shell 1PI graphs in the quantum theory. Since any graph can be produced by sewing together 1PI graphs, the effective action when used at tree level can reproduce all the loops in the original theory. Thus the (quantum) effective action, when used classically, reproduces the full quantum physics of a given classical theory. In compensation for this enormous power, one must sacrifice manifest locality: in general,  $\Gamma[\phi] \neq \int d^4x \mathcal{L}[\phi(x)]$  for any  $\mathcal{L}$ . Knowing the effective action amounts to solving a theory. As this is generally impossible, one must resort to an approximation to the effective action and argue that this approximation is valid for the computation of interest.

Although one could compute the 1PI effective action by computing all 1PI graphs in a theory, this is generally inefficient. Instead, one often computes the effective action by relating it to the Legendre transform of the classical action and doing a background field expansion [5]. We first define the functional

$$W[J] \equiv -i \ln Z[J] = -i \ln \left[ \int \mathcal{D}\phi e^{iS[\phi] + i \int d^4x \phi J} \right] \quad (2.2)$$

where  $J(x)$  is an external current. The effective action is the Legendre transform of  $W[J]$ :

$$\Gamma[\phi] = W[J_\phi] - \int d^4x J_\phi \phi. \quad (2.3)$$

Here  $J_\phi$  denotes the value of the external current which would make a given  $\phi$  solve the equations of motion. Equivalently, in the presence of a current  $J_\phi$ , the expectation value of the field is  $\phi_J$ :

$$\phi_J = \langle J_\phi | \phi(x) | J_\phi \rangle = -i \frac{1}{Z} \frac{\partial Z[J]}{\partial J(x)} \Big|_{J=J_\phi} = \frac{\partial W[J]}{\partial J(x)} \Big|_{J=J_\phi}. \quad (2.4)$$

In this equation  $J_\phi$  is the independent variable which determines  $\phi_J$  as the expectation value of the field in its presence. For example, when  $J = 0$ , so that no current is turned on, then  $\phi_0 = \langle \phi \rangle$  is the true vacuum in the theory. The conjugate relation is

$$\frac{\partial \Gamma[\phi]}{\partial \phi} \Big|_{\phi=\phi_J} = -J_\phi. \quad (2.5)$$

Here  $\phi_J$  is the independent variable to be chosen and used to compute  $J_\phi$ . Importantly, this equation implies that the true vacuum of the theory,  $\phi_0$ , is an extremum of the effective action. There is an assumed one-to-one correspondence between  $\phi_J$  and  $J_\phi$ , which is why the effective potential should always be convex.<sup>1</sup>

The effective action is to be used classically, so that fields are always on-shell, satisfying their equations of motion. When  $\phi$  is not an extremum of  $\Gamma$ , it represents the expectation value of the field in the presence of a nonzero background current; it is the classical solution to a different system. In other words,  $\Gamma[\phi]$  encodes how the system responds to external currents for which Eq. (2.4) is satisfied. That this Legendre transform produces the object whose vertices are 1PI graphs in the original theory is not obvious. It is also not obvious that  $\Gamma[\phi]$  can be computed by shifting the fields  $\phi \rightarrow \phi + \phi_q$  and computing 1PI graphs involving  $\phi_q$  with  $\phi$  held fixed. These three representations of  $\Gamma[\phi]$  are explained in more detail in Refs. [4,24,25].

### A. Gauge dependence

The main issue of concern in this paper is the gauge dependence of the effective action. The gauge dependence arises because when the source  $J$  is nonzero, the system has a nondynamical background charge density. Since there is no corresponding coupling of this charge density to the photon, the Ward identity will consequently be violated [5]. In the special case when  $J = 0$ , there is no charged background. Then  $\phi_0 = \langle 0 | \phi(x) | 0 \rangle$  is extremal and we have simply

$$\Gamma[\phi_0] = W[0] = -i \ln Z[0] = -i \ln \int \mathcal{D}\phi e^{iS[\phi]}. \quad (2.6)$$

This expression is gauge invariant, since gauge-fixing modifies  $Z[0]$  only by an infinite constant, which drops out of  $\ln Z[0]$  (see Ref. [18] for a longer discussion with a careful treatment of the divergences in  $Z[0]$ ). Thus when  $\frac{\partial \Gamma[\phi]}{\partial \phi} = 0$ , then the effective action is gauge invariant by Eq. (2.5).

An alternative way to show gauge invariance at the extremum is by exploring how and when Ward identities are violated in the presence of a background current [17,18]. In particular, Ref. [17] demonstrated that in Fermi gauges, where the gauge-fixing term is  $-\frac{1}{2\xi} (\partial_\mu A_\mu)^2$ , the effective potential satisfies

<sup>1</sup>The convexity issue is a thorny one [22,23], as the effective potential in the Standard Model appears to be perturbatively nonconvex. Conveniently, all the effective potentials we consider in this paper are convex.

$$\left[ \xi \frac{\partial}{\partial \xi} + C(\phi, \xi) \frac{\partial}{\partial \phi} \right] V(\phi, \xi) = 0 \quad (2.7)$$

where  $C(\phi, \xi)$  is some precisely well-defined object, independently calculable order by order in perturbation theory. What this equation implies is that the gauge dependence of the effective potential can be compensated for by a rescaling of  $\phi$ . It also apparently implies that at an extremum, where  $\frac{\partial}{\partial \phi} V(\phi, \xi) = 0$ , the effective potential is automatically gauge invariant,  $\frac{\partial}{\partial \xi} V(\phi, \xi) = 0$ , consistent with the simple argument in the previous paragraph.

There are a number of unsettling features of the Nielsen identity. First of all, it is somewhat trivial, in the sense that one can always define a function  $C(\phi, \xi)$  so that Eq. (2.7) holds; one simply defines  $C(\phi, \xi)$  by Eq. (2.7). Indeed, this is a shortcut to calculating  $C(\phi, \xi)$ . This shortcut makes it clear that the Nielsen identity only guarantees that  $\frac{\partial}{\partial \xi} V(\phi, \xi) = 0$  at an extremum if  $C(\phi, \xi)$  is finite at the extremum. A second unsettling feature of the Nielsen identity is that  $C(\phi, \xi)$  is sometimes infinite [17]. When this happens, one cannot conclude that the potential at the extremum is gauge invariant, or that it is not. It is also possible for  $C(\phi, \xi)$  to be infinite at the extremum to all orders in perturbation theory but finite nonperturbatively.

For a concrete example, as we will see in Sec. VII, in the Coleman-Weinberg model  $C(\phi, \xi)$  is logarithmically divergent as  $\phi$  approaches an extremum when the potential is expressed in terms of the expectation value  $v = \langle \phi \rangle$ . The origin of this divergence is that  $v$  itself is gauge dependent. If instead one expresses  $V$  in terms of the modified minimal subtraction ( $\overline{\text{MS}}$ ) scale  $\mu$ , then  $C(\phi, \xi)$  is finite at the minimum. Thus, as we show in this paper, the value of the potential at the minimum is gauge invariant order by order only in a carefully considered perturbation expansion.

### B. Rescaling and calculation scale invariance

Another feature of the potential at the minimum is that it is independent of field rescaling:  $\phi \rightarrow \kappa \phi$  for any  $\kappa$ . This is not a deep observation, as the value of any function at a minimum is invariant under a rescaling of the dependent variable (see Fig. 1). More generally, the value of the potential at an extremum is independent of any field redefinition,  $\phi \rightarrow \phi'(\phi)$ . These are intuitive features of quantum field theories, since the path integral is also invariant under field redefinitions, and the effective action in the true vacuum is defined as a path integral in Eq. (2.6). Indeed, the gauge invariance of the action at extrema is a special case of general field-redefinition invariance, since one can view gauge transformations as field redefinitions. In doing so, however, one must allow for the possibility that if with one definition an extremum is at  $A_\mu = 0$ , with another definition it may be at a nonzero and  $x$ -dependent expectation value for  $A_\mu$ .




FIG. 1 (color online). Under the rescaling of the dependent variable, a function changes but its values at extrema do not [21]. This elementary mathematical fact explains why the effective potential can depend on the field normalization, but  $V_{\min}$  does not.

A corollary of the above argument is that the potential away from its extrema *does* depend on how the field is normalized and defined. This is also obvious from Fig. 1. Away from an extremum, the action describes the system in the presence of a current  $J$ . When one rescales the field  $\phi$ , the  $J\phi$  term with  $J$  fixed breaks the invariance of the path integral under rescaling. Equivalently, from Eq. (2.5), we see that  $J_{\kappa\phi} = \frac{1}{\kappa} J_\phi$  so that when a field is rescaled,  $\Gamma[\kappa\phi]$  gives the least action in the presence of a rescaled current.

A number of authors have proposed that the gauge dependence of the effective potential can be removed through a field redefinition [8–11,16]. For example, Tye and Vtorov-Karevsky argue that one should replace  $\phi_1 + i\phi_2 \rightarrow \sigma \exp(i\pi)$  [10]. Then  $\sigma$  is a U(1) singlet and so its source  $J$  is neutral and the interaction  $J\sigma$  in the Lagrangian does not cause the Ward identity to be violated. Although there is nothing wrong with this argument, physical quantities, such as the value of the potential at its minimum, should be independent of field redefinitions. A field redefinition is in a sense similar to a gauge choice. It does not make the potential away from the minimum any more physical. Moreover, with this nonlinear field redefinition, a renormalizable theory becomes nonrenormalizable and nominally straightforward calculations can become drastically more complicated (try computing  $\beta_\lambda$  at one loop in this theory). The point is that physics should be independent of field redefinitions, so one should choose a basis which simplifies the calculations, not one that makes unphysical quantities more comforting.

An interesting and underappreciated fact about the effective potential is that its form depends on the scale where it is computed. The effective potential satisfies a renormalization group equation (RGE):

$$\left( \mu \frac{\partial}{\partial \mu} - \gamma_i \phi_i \frac{\partial}{\partial \phi_i} + \beta_i \frac{\partial}{\partial \lambda_i} \right) V = 0. \quad (2.8)$$

This equation says that the explicit dependence on the RG scale  $\mu$  in the effective potential can be compensated by redefining the couplings  $\lambda_i$ , through their  $\beta$  functions, and rescaling the fields  $\phi_i$  based on their anomalous dimensions  $\gamma_i$ . One can compute the potential to fixed order at a scale  $\mu_0$ , in terms of  $\lambda_i(\mu_0)$  and  $\mu_0$  and then evolve to some other scale  $\mu$  by solving this equation simultaneously with the RGEs for the coupling constants; we call this method 1. Alternatively, one could evolve the couplings to  $\mu$  and then compute the effective potential there; we call this method 2. The potentials at  $\mu$  computed in these two different ways will not agree: they differ because in method 1, the rescaling of  $\phi_i$  from  $\mu_0$  to  $\mu$  is included, while in method 2, it is not. We show through an explicit example in Sec. VIII A that in the Coleman-Weinberg model, the two methods give potentials which indeed differ by  $\ln \frac{\mu}{\mu_0}$  terms at next-to-leading order.

One can easily write a differential equation similar to Eq. (2.7) for the scale dependence:

$$\left[ \mu_0 \frac{\partial}{\partial \mu_0} - \gamma_i \phi_i \frac{\partial}{\partial \phi_i} \right] V(\phi, \mu_0, \mu) = 0 \quad (2.9)$$

where  $\mu_0$  is the scale at which the potential is calculated before it is run to the scale  $\mu$  by solving Eq. (2.8). In a sense, this equation although trivial to derive is more useful than the Nielsen identity, since  $\gamma_i$  cannot be infinite [unlike  $C(\phi, \xi)$  in Eq. (2.7)]. It immediately indicates that at an extremum, the effective potential is  $\mu_0$  independent. We use this calculation-scale invariance in Sec. VIII C to show how a resummed effective potential can be computed so that it is gauge invariant order by order in a particular perturbation expansion.

For another perspective, recall that the vertices of the effective action encode (amputated) 1PI Green's functions. General amputated Green's functions satisfy an RGE identical to Eq. (2.8), including the  $\gamma$  term. Thus they also depend on the scale where they are calculated. In converting such Green's functions to  $S$ -matrix elements, the Lehmann-Symanzik-Zimmermann reduction theorem instructs us to multiply by  $\prod_i \sqrt{Z_i}$ , where  $\sqrt{Z_i}$  is the on-shell wave-function renormalization factor for the external leg  $i$ . In  $\overline{\text{MS}}$ , these  $Z$  factors are not 1 but are gauge and scale dependent. Indeed, their  $\mu$  dependence explicitly cancels the  $\gamma$  term in Eq. (2.8) so that all scale dependence in  $S$ -matrix elements is compensated for by rescaling the couplings according to their  $\beta$  functions. As the  $\beta$  functions are gauge invariant in  $\overline{\text{MS}}$ , the  $S$  matrix can then be gauge invariant at any scale.

In summary, the effective potential at general field values depends on an arbitrary calculation scale  $\mu_0$ . This is different from the usual dependence on the renormalization scale  $\mu$  arising from a truncation to fixed order in perturbation theory. The effective potential has that dependence as well, but it depends on  $\mu_0$  even nonperturbatively. The value of the potential at extrema is  $\mu_0$  independent.

That the  $\mu_0$  dependence drops out at extrema does not require field redefinitions or multiplication by the wave-function renormalization factors included in going from Green's functions to  $S$ -matrix elements.

### III. ABELIAN HIGGS MODEL

This paper focuses on scalar QED, a renormalizable theory with two scalar fields  $\phi_1$  and  $\phi_2$ , a photon,  $A_\mu$  and an  $O(2)$  symmetry. If the scalar doublet is given a negative mass squared in the classical potential, the model is called the Abelian Higgs model. This section is a review of results in Refs. [6,19,26,27] with some details filled in and a few added comments. Establishing gauge invariance in this model is simpler than in the Coleman-Weinberg model for the essential reason that there is spontaneous symmetry breaking already in the classical potential. Consequently, a traditional loop expansion is justified and the first nontrivial check occurs at one loop. Nevertheless, the Abelian Higgs model presents interesting features and subtleties that are complementary to those in the Coleman-Weinberg model, which is why we discuss it here.

We write the Lagrangian as

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu}^2 + \frac{1}{2} (\partial_\mu \phi_1 - e A_\mu \phi_2)^2 + \frac{1}{2} (\partial_\mu \phi_2 + e A_\mu \phi_1)^2 - V_0 + \mathcal{L}_{\text{GF}} + \mathcal{L}_{\text{ghost}} \quad (3.1)$$

with the classical potential

$$V_0(\phi) = -\frac{1}{2} m^2 \phi^2 + \frac{\lambda}{24} \phi^4 \quad (3.2)$$

where

$$\phi^2 \equiv \phi_1^2 + \phi_2^2. \quad (3.3)$$

Since the theory has a global  $O(2)$  symmetry rotating  $\phi_1$  and  $\phi_2$ , the effective potential should only depend on  $\phi$ . When  $m^2 > 0$ , the classical potential is minimized when

$$v_{\text{cl}} \equiv \langle \phi_1 \rangle_{\text{classical}} = \sqrt{\frac{6}{\lambda}} m. \quad (3.4)$$

We have used the global  $O(2)$  symmetry to put the vacuum expectation value (VEV) in  $\phi_1$  with  $\langle \phi_2 \rangle = 0$ . Since we have normalized the vacuum energy so that  $V_0(0) = 0$ , the value of the classical potential at the minimum is  $V_0(v_{\text{cl}}) = -\frac{3}{2\lambda} m^4 = -\frac{\lambda}{24} v_{\text{cl}}^4$ .

To compute quantum corrections to the vacuum energy, we need to fix a gauge. In gauge fixing, as always, one looks for a deformation of the theory with the fewest nasty features. To appreciate the relevant tradeoffs, it is helpful to use a three-parameter family ( $\xi$ ,  $\Upsilon_1$  and  $\Upsilon_2$ ) of gauges [6]. We write

$$\mathcal{L}_{\text{GF}} = -\frac{1}{2\xi}(\partial_\mu A^\mu + e\xi\Upsilon_1\phi_1 + e\xi\Upsilon_2\phi_2)^2. \quad (3.5)$$

The associated Fadeev-Popov ghost Lagrangian is

$$\mathcal{L}_{\text{ghost}} = (\partial_\mu \bar{c})(\partial_\mu c) - e^2\xi\Upsilon_1\phi_2\bar{c}c + e^2\xi\Upsilon_2\phi_1\bar{c}c \quad (3.6)$$

where the ghosts  $c$  and antighosts  $\bar{c}$  are Grassmann-valued scalar fields.

For  $\Upsilon_i = 0$ , these are the *Fermi gauges*. Fermi gauges are appealing because they do not explicitly break the global  $O(2)$  symmetry rotating  $\phi_1$  and  $\phi_2$  and because the ghosts decouple. A drawback of Fermi gauges is that there is uncanceled kinetic mixing between the gauge field and the scalars. A more serious problem with Fermi gauges is that they can lead to singular intermediate results, as we will see, which make some calculations difficult.

The  $R_\xi$  gauges [28] are  $\Upsilon_1 = v_{\text{cl}}$  and  $\Upsilon_2 = 0$ . Note that even in  $R_\xi$  gauges, there is still kinetic mixing when  $\phi \neq v_{\text{cl}}$  and, except for Landau gauge,  $\xi = 0$ , the ghosts are not free. In the  $R_\xi$  gauges, the global  $O(2)$  symmetry is explicitly broken. This breaking is not dangerous, since gauge-independent quantities must be independent of the gauge fixing and should respect the  $O(2)$  symmetry. A more troubling fact about  $\Upsilon_1 = v_{\text{cl}}$  is that the *classical* minimum  $v_{\text{cl}}$  is then hardcoded into the Lagrangian, when presumably only the true quantum minimum has any physical significance. It would be even more troubling to allow  $\Upsilon_i$  to be dynamical, for example with  $\Upsilon_1 = \phi$ ; such gauges led Weinberg to declare  $R_\xi$  gauges problematic in all but Landau gauge,  $\xi = 0$  [7] (see Ref. [18] for a careful refutation of the argument in Ref. [7]). In the three-parameter family we consider here,  $\Upsilon_i$  are arbitrary, so  $v_{\text{cl}}$  does not appear.

Another feature of the Abelian Higgs model with this gauge fixing is that it has a spurious unphysical minimum. The gauge fixing induces a coupling between  $\phi_1$  and  $\phi_2$  in the classical potential. Thus there is a solution for constant fields with  $A_\mu = c = \bar{c} = 0$  and

$$\phi_i = e\Upsilon_i \sqrt{\frac{6m^2}{\lambda e^2(\Upsilon_1^2 + \Upsilon_2^2)} + \frac{6\xi}{\lambda}}. \quad (3.7)$$

The potential in this solution has the value

$$V_{\text{min}} = -\frac{3}{2\lambda}[m^2 + e^2\xi(\Upsilon_1^2 + \Upsilon_2^2)]^2 \quad (3.8)$$

which is gauge dependent and unphysical. This solution, found by Dolan and Jackiw [6], led them to argue for unitary gauge,  $\xi = \infty$ . As explained in Ref. [18] (see also Ref. [27]), the trouble with this solution is not the gauge dependence, but that, despite satisfying the Euler-Lagrange equations, it does not extremize the action. An extremal solution related to this one has a nonvanishing and position-dependent expectation value for  $A_\mu$ ; the gauge-fixing term contributes at spatial infinity so the boundary terms cannot be dropped in deriving the Euler-Lagrange equations. In this sense the three-parameter gauge fixing provides what Ref. [18] called a *bad gauge*. Note that there is nothing inconsistent about bad gauges; they are just rather inconvenient for doing calculations because of the position-dependent extrema. If we stick to solutions close to the gauge-independent classical one, with  $\phi_1 = v_{\text{cl}}$  and  $\phi_2 = 0$ , the position-dependent solutions can be ignored. Another criterion for choosing one gauge over another was discussed in Ref. [13].

We can always use the  $O(2)$  symmetry to keep  $\langle\phi_2\rangle = 0$  to all orders. If we do so, then we do not even need to turn on a background field for  $\phi_2$ , since it will not contribute to the extremal solutions for  $\phi_1$ . To avoid the bad gauges discussed above, we should correspondingly take  $\Upsilon_1 = 0$ . A more general  $O(2)$ -invariant condition is to only turn on the linear combination which has  $\Upsilon_i \cdot \langle\phi_i\rangle = 0$ . This is essentially what was done in Ref. [26]. Then, the renormalized one-loop effective potential in  $\overline{\text{MS}}$  with  $m \neq 0$  is

$$V_1(\phi) = \frac{\hbar}{16\pi^2} \left[ \frac{3}{4}M_A^4 \left( \ln \frac{M_A^2}{\mu^2} - \frac{5}{6} \right) + \frac{1}{4}M_B^4 \left( \ln \frac{M_B^2}{\mu^2} - \frac{3}{2} \right) - \frac{1}{2}M_G^4 \left( \ln \frac{M_G^2}{\mu^2} - \frac{3}{2} \right) + \frac{1}{4}M_+^4 \left( \ln \frac{M_+^2}{\mu^2} - \frac{3}{2} \right) + \frac{1}{4}M_-^4 \left( \ln \frac{M_-^2}{\mu^2} - \frac{3}{2} \right) \right] \quad (3.9)$$

where

$$M_A^2 = e^2\phi^2, \quad (3.10)$$

$$M_B^2 = \frac{\lambda}{2}\phi^2 - m^2, \quad (3.11)$$

$$M_G^2 = \xi e^2(\Upsilon_1\phi_2 - \Upsilon_2\phi_1) \quad (3.12)$$

are the contributions from  $A_\mu$ ,  $\phi$  and ghosts, respectively and

$$M_{\pm}^2 = \frac{\lambda}{12}(\phi^2 - v_{\text{cl}}^2) + \xi e^2(\Upsilon_1\phi_2 - \Upsilon_2\phi_1) \pm \frac{1}{2}\sqrt{\frac{\lambda}{6}[\phi^2 - v_{\text{cl}}^2] \left[ \frac{\lambda}{6}(\phi^2 - v_{\text{cl}}^2) - 4\xi e^2\phi^2 + 4\xi e^2(\Upsilon_1\phi_2 - \Upsilon_2\phi_1) \right]} \quad (3.13)$$

come from the kinetic mixing.

Then, assuming that the VEV of  $\phi_1$  gets a perturbative correction  $v = \langle\phi_1\rangle = v_{\text{cl}} + v_1 + \dots$  for some  $v_1$  of order  $\hbar$ , we can evaluate the potential at the minimum perturbatively:

$$V_{\text{min}} = V_0(v) + V_1(v) + \dots \quad (3.14)$$

$$= V_0(v_{\text{cl}}) + v_1 V_0'(v_{\text{cl}}) + V_1(v_{\text{cl}}) + \dots \quad (3.15)$$

$$= V_0(v_{\text{cl}}) + V_1(v_{\text{cl}}) + \dots + \mathcal{O}(\hbar^2) \quad (3.16)$$

where the omitted terms are  $\mathcal{O}(\hbar^2)$  and  $V_0'(v_{\text{cl}}) = 0$  has been used. The importance of doing an analytic expansion in  $\hbar$  with an appropriate truncation before evaluating  $V_{\text{min}}$  or other physical quantities was emphasized by Patel and Ramsey-Musolf in Ref. [21] (see also Ref. [19]). Since  $M_+^2 = M_-^2 = M_G^2 = -\xi e^2 \Upsilon_2 v_{\text{cl}}$  when  $\phi_1 = v_{\text{cl}}$  and  $\phi_2 = 0$ , we find that  $V(v)$  is gauge invariant to order  $\hbar$ . Explicitly,

$$V_{\text{min}} = v_{\text{cl}}^4 \left[ -\frac{\lambda}{24} + \frac{\hbar}{16\pi^2} \left( -\frac{5}{8}e^4 - \frac{1}{24}\lambda^2 + \frac{1}{36}\lambda^2 \ln \frac{\lambda v_{\text{cl}}^2}{3\mu^2} + \frac{3}{4}e^4 \ln \frac{e^2 v_{\text{cl}}^2}{\mu^2} \right) + \mathcal{O}(\hbar^2) \right]. \quad (3.17)$$

Thus the vacuum energy is manifestly gauge independent.

To determine  $v_1$ , we note that the minimum condition is

$$0 = V_0'(v) + V_1'(v) + \dots = V_0'(v_{\text{cl}}) + v_1 V_0''(v_{\text{cl}}) + V_1'(v_{\text{cl}}) + \dots \quad (3.18)$$

Using again that  $V_0'(v_{\text{cl}}) = 0$ , we get  $v_1 = -\frac{V_1'(v_{\text{cl}})}{V_0''(v_{\text{cl}})}$  which gives

$$v_1 = v_{\text{cl}} \frac{\hbar}{16\pi^2} \left[ \frac{3e^4}{\lambda} + \frac{\lambda}{2} - \frac{9e^4}{\lambda} \ln \frac{e^2 v_{\text{cl}}^2}{\mu^2} - \frac{\lambda}{2} \ln \frac{v_{\text{cl}}^2 \lambda}{3\mu^2} + \frac{e^2 \xi}{2} \ln \frac{-e^2 v_{\text{cl}} \xi \Upsilon_2}{\mu^2} \right] + \Upsilon_2 e^2 \xi \frac{\hbar}{16\pi^2} \left[ -\frac{1}{2} + \ln \frac{-e^2 \xi v_{\text{cl}} \Upsilon_2}{\mu^2} \right]. \quad (3.19)$$

This expression manifestly depends on both  $\xi$  and  $\Upsilon_2$ . The dangers of assigning physical significance to  $\langle\phi\rangle$  have been emphasized in Refs. [19,21] and elsewhere.

Note that  $v_1$  is singular in Fermi gauges,  $\Upsilon_i \rightarrow 0$ , but not in Landau gauge  $\xi \rightarrow 0$ . The singularity is an infrared divergence associated with the ghosts becoming massless. As pointed out in Refs. [17] and [26], the function  $C(\phi, \xi)$  in the Nielsen identity also has an infrared divergence in Fermi gauge. As discussed in Sec. II A, the fact that  $C(\phi, \xi)$  can be infinite calls into question the usefulness of the Nielsen identity; as we have seen, the potential at the minimum is gauge invariant for any choice of  $\Upsilon_2$  and  $\xi$ .

At this point, we can conclude that everything works swimmingly: the vacuum energy is gauge invariant but the VEV is not, exactly as foretold by the Nielsen identity. At the risk of being too cautious, it is perhaps worth adding that in the Abelian Higgs model,  $V_{\text{min}}$  is not actually observable; the true renormalized vacuum energy has an arbitrary subtraction associated with the cosmological constant. Even if the vacuum energy can be measured (for example, by weakly coupling this model to gravity), one still needs a scheme to measure the difference  $\Delta V = V_{\text{min}} - V(0)$  between this energy and the energy at  $\phi = 0$ . This difference, as a function of  $v_{\text{cl}}$ ,  $e$  and  $\lambda$  is what we have calculated.

Even then, although  $\Delta V$  is independent of gauge, it is expressed in terms of  $v_{\text{cl}} = \sqrt{\frac{e}{\lambda}} m$  which is itself unphysical. Indeed, even  $m$  is not physical: it is not the mass of the scalar in the vacuum. That mass, which we denote as  $m_S$ , itself gets corrections of order  $\hbar$  from self-energy graphs. The gauge-dependent part of these corrections at order  $e^2 \lambda$  has been calculated in Ref. [26], with the result that  $\frac{d}{d\xi} m_S = 0$ . Therefore  $\Delta V/m_S^4$  is gauge independent and only a function of the  $\overline{\text{MS}}$  couplings  $e$  and  $\lambda$ . This ratio is in principle observable, if  $\Delta V$  is, and depends only on other observables,  $e$  and  $\lambda$ . With this long-winded disclaimer, we can now conclude that quantum corrections in the effective potential can make (in principle) testable experimental predictions.

In summary, in the Abelian Higgs model both  $\Delta V = V_{\text{min}} - V(0)$  and  $m_S$  are separately independent of the gauge parameters  $\xi$  and  $\Upsilon_2$  at order  $\hbar$  when expressed in terms of the classical expectation value of  $\phi$ ,  $v_{\text{cl}}$ . That  $\Delta V$  and  $m_S$  are separately gauge invariant is not strictly necessary. All that is required is that the gauge dependence cancels in the ratio  $\Delta V/m_S^4$  which is a calculable function of the observables  $e$  and  $\lambda$ . As we will see, in the Coleman-Weinberg model, where the classical theory is scaleless, such considerations are critical. We also found that the expectation value of the

field  $v = \langle \phi \rangle$  is *not* gauge invariant at order  $\hbar$ . In the Fermi gauges,  $\Upsilon_i = 0$ ,  $v$  even has an infrared divergence.

#### IV. COLEMAN-WEINBERG MODEL

Now let us turn to the main subject of this paper, i.e., scalar QED with a massless scalar, also known as the Coleman-Weinberg model. The Lagrangian is as in Eq. (3.1), with

$$V_0 = \frac{\lambda}{24} \phi^4 \quad (4.1)$$

and the gauge-fixing term is

$$V_1(\phi) = \phi^4 \frac{\hbar}{16\pi^2} \left[ \frac{3}{4} e^4 \left( \ln \frac{e^2 \phi^2}{\mu^2} - \frac{5}{6} \right) + \frac{\lambda^2}{16} \left( \ln \frac{\lambda \phi^2}{2\mu^2} - \frac{3}{2} \right) + \left( \frac{\lambda^2}{144} - \frac{1}{12} e^2 \lambda \xi \right) \left( \ln \frac{\phi^2}{\mu^2} - \frac{3}{2} \right) + \frac{1}{4} K_+^4 \ln K_+^2 + \frac{1}{4} K_-^4 \ln K_-^2 \right] \quad (4.3)$$

with

$$K_{\pm}^2 = \frac{1}{12} (\lambda \pm \sqrt{\lambda^2 - 24\lambda e^2 \xi}). \quad (4.4)$$

The relation

$$K_+^4 + K_-^4 = \frac{\lambda^2}{36} - \frac{1}{3} e^2 \lambda \xi \quad (4.5)$$

has been used to simplify the one-loop potential.

The tree-level potential in this model has only a single minimum, at  $\phi = 0$ , where the  $O(2)$  symmetry is unbroken. For there to be a minimum at one loop, the corrections must be large enough to turn over the potential. For  $\lambda$  small, so that the theory is perturbative, this is only possible if the  $\frac{\hbar}{16\pi^2} e^4$  term is as large as the tree-level  $\frac{\lambda}{24}$  piece. So let us assume  $\lambda \sim \frac{\hbar}{16\pi^2} e^4$  and that there is a minimum at some scale  $v$ . Then the condition for the minimum,  $V'(v) = 0$  provides a precise relationship between  $\lambda$  and  $e$ :

$$\lambda = \frac{\hbar}{16\pi^2} e^4 \left( 6 - 36 \ln \frac{ev}{\mu} \right) + \mathcal{O}(e^6). \quad (4.6)$$

As discussed in Ref. [4], to understand this equation one must appreciate dimensional transmutation. In scalar QED, the only scale is the scale  $\mu$  at which the couplings are defined.  $\mu$  is arbitrary, so we may as well take  $\mu = v$ . Then Eq. (4.6) reduces to

$$\lambda = \frac{\hbar}{16\pi^2} e^4 (6 - 36 \ln e) + \mathcal{O}(e^6). \quad (4.7)$$

This equation should be thought of as a condition on  $\lambda = \lambda(\mu)$  and  $e = e(\mu)$ : the minimum occurs at the scale  $v = \mu$

$$\mathcal{L}_{\text{GF}} = -\frac{1}{2\xi} (\partial_\mu A^\mu)^2. \quad (4.2)$$

These can be thought of as  $R_\xi$  gauges (since  $v_{\text{cl}} = 0$  in this theory) or Fermi gauges. They correspond to  $\Upsilon_1 = \Upsilon_2 = 0$  in the three-parameter gauge family discussed above, and so the ghosts decouple and can be ignored. There is unavoidable kinetic mixing between  $\phi$  and  $\partial_\mu A_\mu$  in this theory, but as in the Abelian Higgs model, this is an inconvenient but not insurmountable complication.

The renormalized one-loop effective potential in  $\overline{\text{MS}}$  in this theory is the  $m \rightarrow 0$  limit of Eq. (3.9):

where Eq. (4.7) holds. Of course,  $v$  will get corrections and, as we will see, is gauge dependent (unlike  $\mu$ ). But at least at one loop, this is an acceptable way to think about the minimum in the effective potential in scalar QED.

Since  $\lambda$  and  $e$  can be anything, it is natural to wonder whether Eq. (4.7) requires some kind of finite tuning. As explained in Ref. [4] it does not. The evolution of  $e$  and  $\lambda$  are determined by the  $\beta$  functions:

$$\mu \frac{d}{d\mu} e = \beta_e, \quad \mu \frac{d}{d\mu} \lambda = \beta_\lambda, \quad (4.8)$$

where, at one loop,

$$\beta_e = \frac{\hbar}{16\pi^2} \left( \frac{e^3}{3} \right) + \dots, \quad (4.9)$$

$$\beta_\lambda = \frac{\hbar}{16\pi^2} \left( 36e^4 - 12e^2\lambda + \frac{10\lambda^2}{3} \right). \quad (4.10)$$

The key feature of these equations is that  $e$  is multiplicatively renormalized ( $e = 0$  is a fixed point of the RG flow) while  $\lambda$  can get an additive correction even at  $\lambda = 0$ . What this means is that if  $\lambda$  and  $e$  start off small,  $e$  runs logarithmically, but  $\lambda$  will grow at an ever increasing rate until it hits a Landau pole. Indeed, the exact solutions to the one-loop RGEs are [4]

$$e^2(\mu) = \frac{e^2(\mu_0)}{1 - \frac{e^2(\mu_0)}{24\pi^2} \ln \frac{\mu}{\mu_0}} \quad (4.11)$$

which expresses  $e(\mu)$  in terms of  $e$  at some reference scale  $\mu_0$  and



$$\lambda(\mu) = \frac{e^2(\mu)}{10} \left[ 19 + \sqrt{719} \tan \left( \frac{\sqrt{719}}{2} \ln \frac{e(\mu)^2}{C} \right) \right] \quad (4.12)$$

where  $C$  is an integration constant which can be traded for  $\lambda(\mu_0)$ . The tangent in  $\lambda(\mu)$  implies that as  $e^2$  changes by a factor of  $\approx 1.2$ ,  $\lambda$  will go from  $-\infty$  to  $\infty$ . In particular, if  $\lambda$  and  $e$  are small, there will *always* be a point where Eq. (4.7) is satisfied.

At the minimum, we find

$$V_{\min} = v^4 \frac{\hbar}{16\pi^2} e^4 \left( -\frac{3}{8} \right) + \mathcal{O}(e^6). \quad (4.13)$$

This is gauge invariant, simply because the  $e^4$  terms in  $V_1(\phi)$  are gauge invariant.

The first nontrivial check on the gauge invariance of  $V(v)$  requires the terms in the effective potential of order  $\hbar^2 e^6$ , with  $\lambda$  counting as order  $\hbar e^4$ , and  $\ln e$  and  $\ln \lambda$  counting as order  $e^0$ . In scalar QED, each loop comes with a factor of  $\hbar e^2$  or  $\hbar \lambda$  from the vertices, so  $\hbar e^2 \lambda$  terms come from one-loop graphs and  $\hbar^2 e^6$  terms from two-loop graphs. Thus we need at least the two-loop Coleman-Weinberg potential. In addition, effective-potential calculations involve an infinite number of background-field insertions, which can be conveniently resummed into dressed propagators with  $e$ - and  $\lambda$ -dependent masses. These propagators allow for graphs to have extra factors of  $e^2/\lambda$  in them. For example, a three-loop graph gives a term proportional to  $\hbar^3 e^{10}/\lambda$  which also scales like  $\hbar^2 e^6$

when  $\lambda \sim \hbar e^4$ . These higher-loop terms are discussed in Sec. VI.

## V. THE TWO-LOOP COLEMAN-WEINBERG POTENTIAL

The complete two-loop potential in scalar QED with gauge dependence does not appear in the literature, as far as we are aware. The two-loop potential in Landau gauge is known in the complete Standard Model [29] and for general renormalizable theories [30]. Kang computed the terms in the two-loop potential at order  $e^6$  that depend on  $\ln \frac{\phi^2}{\mu^2}$ , but with a subtraction scheme slightly different from  $\overline{\text{MS}}$  [31]. For our analysis, we also require the  $\mu$ -independent part of the effective potential at order  $e^6$ . We will also need an infinite series of other graphs, the *daisy* graphs, which are discussed in Sec. VI, but let us start with the two-loop computation. To compute the effective potential at this order, there is a counterterm contribution and a contribution from two-loop graphs. We work in  $d = 4 - 2\epsilon$  dimensions throughout, and the scale  $\mu$  is the  $\overline{\text{MS}}$  renormalization scale which implicitly contains the  $\ln 4\pi$  and  $\gamma_E$  factors.

### A. Counterterm contribution

The counterterm contribution can be extracted from the  $\mathcal{O}(\epsilon)$  part of the one-loop potential by multiplying the  $\frac{1}{\epsilon}$  poles in the renormalization factors. In  $d = 4 - 2\epsilon$  dimensions, the one-loop potential can be written as

$$V_1 = \phi^4 [(3 - 2\epsilon)f(e^2) + f\left(\frac{\lambda}{2}\right) + f(K_+^2) + f(K_-^2)] + \mathcal{O}(\epsilon^2) \quad (5.1)$$

where

$$f(x) = -\frac{x^2}{64\pi^2} \left[ \frac{1}{\epsilon} + \frac{3}{2} - \ln \frac{x\phi^2}{\mu^2} + \frac{\epsilon}{2} \left( \ln \frac{x\phi^2}{\mu^2} - \frac{3}{2} \right)^2 + \epsilon \left( \frac{5}{8} + \frac{\pi^2}{12} \right) \right] \quad (5.2)$$

with  $K_{\pm}$  defined in Eq. (4.4). The one-loop renormalization factors are

$$Z_{\phi} = 1 + \frac{\hbar}{16\pi^2} \frac{1}{\epsilon} (3 - \xi)e^2 + \dots, \quad Z_e = 1 + \frac{\hbar}{16\pi^2} \frac{1}{\epsilon} e^2 + \dots, \quad \lambda Z_{\lambda} = \lambda + \frac{\hbar}{16\pi^2} \frac{1}{\epsilon} \left( 18e^4 - 6e^2\lambda + \frac{5}{3}\lambda^2 \right) + \dots \quad (5.3)$$

Replacing  $e \rightarrow Z_e e$ ,  $\lambda \rightarrow Z_{\lambda} \lambda$  and  $\phi \rightarrow \sqrt{Z_{\phi}} \phi$  and expanding to order  $\hbar^2 e^0$  gives the counterterm contribution to the two-loop potential:

$$V_2^{\text{ct}} = \left( \frac{\hbar}{16\pi^2} \right)^2 \phi^4 e^6 \left[ (-10 + 6\xi) \ln^2 \frac{e\phi}{\mu} + \left( \frac{10}{3} - 4\xi + \frac{3}{2}\xi \ln \frac{\lambda\xi}{6e^2} \right) \ln \frac{e\phi}{\mu} - \frac{5}{3} - \frac{5\pi^2}{12} + 2\xi + \frac{\pi^2}{16}\xi - \frac{3}{4}\xi \ln \frac{\lambda\xi}{6e^2} + \frac{3}{16}\xi \ln^2 \frac{\lambda\xi}{6e^2} \right] + \mathcal{O}(e^4\lambda, e^2\lambda^2, \lambda^3). \quad (5.4)$$

**B. Two-loop contribution**

The two-loop contribution to the next-to-next-to-leading-order (NNLO) effective potential can be computed from Feynman diagrams with dressed propagators (called *prototype diagrams* in Ref. [4]). The most straightforward way to compute the effective potential is to turn on a background field for  $\phi_1$  by replacing  $\phi_1 \rightarrow \phi_1 + \phi$  in the classical Lagrangian. After

substituting this into the classical Lagrangian and dropping the terms linear in the propagating fields  $\phi_1$ ,  $\phi_2$  and  $A_\mu$ , we compute the propagators by diagonalizing the kinetic terms, treating the background field  $\phi$  as constant. The resulting propagators are

$$1 \text{ ————— } 1 = D_{11}(k) = \frac{i}{k^2 - \frac{\lambda}{2}\phi^2} \quad (5.5)$$

$$2 \text{ ————— } 2 = D_{22}(k) = i \frac{k^2 - e^2\xi\phi^2}{(k^2 - \phi^2 K_+)(k^2 - \phi^2 K_-)} \quad (5.6)$$

$$\mu \text{ ~~~~~ } \nu = \Delta_{\mu\nu}(k) = -i \left[ \frac{1}{k^2 - e^2\phi^2} \left( g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) + \frac{\xi(k^2 - \frac{1}{6}\lambda\phi^2)}{(k^2 - \phi^2 K_+)(k^2 - \phi^2 K_-)} \frac{k_\mu k_\nu}{k^2} \right], \quad (5.7)$$

with  $K_\pm$  given in Eq. (4.4) and

$$\mu \text{ ~~~~~ } 2 = T_{\mu 2}(k) = \frac{\xi e\phi}{(k^2 - \phi^2 K_+)(k^2 - \phi^2 K_-)} k_\mu. \quad (5.8)$$

This last propagator comes from the kinetic mixing between  $\phi_2$  and  $A_\mu$  in the presence of a background field for  $\phi_1$ . The asymmetry between  $D_{11}$  and  $D_{22}$  arises because we are only turning on a background for  $\phi_1$  for simplicity. The full effective potential of course also depends on background values for  $\phi_2$ . However, since the effective potential has a global  $O(2)$  symmetry, it is enough to turn

on a  $\phi_1$  background for the questions we address in this paper.

The propagators in Eqs. (5.5)–(5.8) are effective propagators, where all of the flipping between scalars and longitudinal vector bosons that can occur between interactions has already been resummed. In this way, an infinite set of diagrams is replaced by a single one:

$$\text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3} + \text{Diagram 4} + \dots \rightarrow \text{Diagram 5}. \quad (5.9)$$

Diagrams where there is one flip between vertices are still allowed (for example, see Fig. 2 below). The hair on the left-hand diagrams indicates background field insertions which are implicit in prototype diagrams, even without kinetic mixing.

There are five types of interactions:

$$\begin{array}{c} i \\ \diagdown \\ j \end{array} \begin{array}{c} l \\ \diagup \\ k \end{array} = -i\frac{\lambda}{3}(\delta_{ij}\delta_{kl} + \delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk}), \quad \begin{array}{c} i \\ \diagdown \\ k \end{array} \begin{array}{c} j \\ \diagup \\ k \end{array} = -i\frac{\lambda}{3}\phi(\delta_{ij}\delta_{k1} + \delta_{jk}\delta_{i1} + \delta_{ki}\delta_{j1}), \quad (5.10)$$

$$\begin{array}{c} i \\ \diagdown \\ j \end{array} \begin{array}{c} \mu \\ \diagup \\ \nu \end{array} = 2ie^2\delta_{ij}g_{\mu\nu}, \quad \begin{array}{c} i \\ \diagdown \\ j \end{array} \begin{array}{c} \mu \\ \diagup \\ \nu \end{array} = e\epsilon_{ij}(k_i^\mu + k_j^\mu), \quad \begin{array}{c} \mu \\ \diagdown \\ \nu \end{array} \begin{array}{c} j \\ \diagup \\ \nu \end{array} = 2ie^2\phi\delta_{j1}g_{\mu\nu} \quad (5.11)$$




FIG. 2. There are 12 prototype diagrams which contribute to the two-loop Coleman-Weinberg potential in  $R_\xi$  gauges. Only the four diagrams in the top row contribute at next-to-leading order (NLO) (order  $\hbar^2$ ) when  $\lambda \sim \hbar$ .

where the solid lines can be either  $\phi_1$  or  $\phi_2$ . These Feynman rules can also be found in Ref. [31] after rescaling  $\lambda \rightarrow \frac{4}{3}$  to match our normalization, with an extended discussion of their derivation in Appendix A of that reference. Diagrams constructed with these Feynman rules were dubbed *prototype diagrams* in Ref. [4] to emphasize that they represent an infinite number of diagrams with arbitrary numbers of background-field insertions.

The effective potential is computed with these Feynman rules by evaluating vacuum energy contributions, i.e., diagrams with no external propagating fields. Each such diagram corresponds to an infinite number of background-field insertions. One can also compute terms in the effective action which have derivatives, such as corrections to the kinetic term  $(\partial_\mu \phi)(\partial_\mu \phi)$ . To do so, one inserts a finite momentum  $p^\mu$  into a single background field line. The procedure is described in Ref. [32]. The one-loop effective potential is a special case: the vacuum diagrams are quadratically divergent. A trick to computing it is to add external legs with  $p^\mu = 0$ . Inserting one such leg produces the first derivative of the potential,  $V'(\phi)$ . Two legs gives the second derivative,  $V''(\phi)$  and so in. Conversely, the second derivative of the potential  $V''(\phi)$  determines the scalar propagator at zero momentum, which in turn can be used to calculate the mass of the scalar at one loop. The path of least resistance is often to compute the one-loop potential using functional determinants [5] and the potential at two loops and beyond with prototype Feynman diagrams.

At two loops, no additional tricks are needed and we can compute the effective potential directly from the vacuum diagrams with dressed propagators. All of the two-loop diagrams can be computed using two master integrals. The first is a standard one-loop integral

$$I_1(\Delta) = \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2 - \Delta + i\epsilon} = \frac{-i}{(4\pi)^{d/2}} \frac{1}{\Delta^{1-\frac{d}{2}}} \Gamma\left(\frac{2-d}{2}\right). \quad (5.12)$$

The second is the two-loop scalar vacuum sunset graph with three masses [29,30,33]:

$$I_2(A, B, C) = \int \frac{d^d k_1}{(2\pi)^d} \int \frac{d^d k_2}{(2\pi)^d} \frac{1}{(k_1^2 - A + i\epsilon)(k_2^2 - B + i\epsilon)((k_1 + k_2)^2 - C + i\epsilon)}. \quad (5.13)$$

This integral is fully symmetric in  $A$ ,  $B$  and  $C$ . The full result in  $d$  dimensions can be found in Ref. [33] along with its  $\epsilon$  expansion. Writing  $d = 4 - 2\epsilon$ , the expansion is

$$\begin{aligned} I_2(A, B, C) = & \frac{1}{(4\pi)^{4-2\epsilon}} \frac{\Gamma(1+\epsilon)^2}{(1-\epsilon)(1-2\epsilon)} \frac{1}{2C^{2\epsilon}} \\ & \times \left\{ -\frac{1}{\epsilon^2} (A+B+C) + \frac{2}{\epsilon} \left( A \ln \frac{A}{C} + B \ln \frac{B}{C} \right) - A \ln^2 \frac{A}{C} - B \ln^2 \frac{B}{C} + (C-A-B+C\lambda) \ln \frac{A}{C} \ln \frac{B}{C} \right. \\ & - \frac{1}{3} C \lambda \pi^2 - 2C\lambda \ln \frac{A-B+C-C\lambda}{2C} \ln \frac{B-A+C-C\lambda}{2C} + 2C\lambda \text{Li}_2 \left( \frac{B-A+C-C\lambda}{2C} \right) \\ & \left. + 2C\lambda \text{Li}_2 \left( \frac{A-B+C-C\lambda}{2C} \right) \right\} \end{aligned} \quad (5.14)$$

with

$$\lambda = \sqrt{\frac{A^2 + (B - C)^2 - 2A(B + C)}{C^2}}. \quad (5.15)$$

This expansion holds if  $\lambda^2 \geq 0$  and  $\sqrt{A} + \sqrt{B} \leq \sqrt{C}$ . The expansion in other regions can be obtained by permuting  $A$ ,  $B$  and  $C$ .

There are four diagrams that contribute at order  $\hbar^2 e^6$ :

$$\begin{aligned} \mathcal{M}_A &= \text{Diagram A} = -\frac{\hbar^2 e^2}{2} \int \frac{d^d k_1}{(2\pi)^d} \int \frac{d^d k_2}{(2\pi)^d} D_{22}(k_1) \Delta_{\mu\mu}(k_2) \\ &= \frac{\hbar^2 \phi^4 e^6}{(16\pi^2)^2} \xi \left[ -12 \ln^2 \frac{e\phi}{\mu} + \left( 8 - 3 \ln \frac{\lambda\xi}{6e^2} \right) \ln \frac{e\phi}{\mu} - \frac{5}{2} - \frac{\pi^2}{16} - \frac{3}{16} \ln^2 \frac{\lambda\xi}{6e^2} + \ln \frac{\lambda\xi}{6e^2} \right] + \dots, \end{aligned} \quad (5.16)$$

$$\begin{aligned} \mathcal{M}_B &= \text{Diagram B} = \frac{i}{2} e^2 \int \frac{d^d k_1}{(2\pi)^d} \int \frac{d^d k_2}{(2\pi)^d} (k_2^\mu - k_1^\mu)(k_1^\nu - k_2^\nu) D_{11}(k_1) D_{22}(k_2) \Delta_{\mu\nu}(k_1 + k_2) \\ &= \frac{\hbar^2 \phi^4 e^6}{(16\pi^2)^2} \left[ (2 + 6\xi) \ln^2 \frac{e\phi}{\mu} - (3 + 7\xi) \ln \frac{e\phi}{\mu} + \frac{7}{4} + \frac{\pi^2}{8} + \frac{15}{4} \xi + \frac{3\pi^2}{8} \xi \right] + \dots, \end{aligned} \quad (5.17)$$

$$\begin{aligned} \mathcal{M}_C &= \text{Diagram C} = -i\hbar^2 e^4 \phi^2 \int \frac{d^d k_1}{(2\pi)^d} \int \frac{d^d k_2}{(2\pi)^d} \Delta_{\mu\nu}(k_1) \Delta_{\mu\nu}(k_2) D_{11}(k_1 + k_2) \\ &= \frac{\hbar^2 \phi^4 e^6}{(16\pi^2)^2} \left[ (18 + 6\xi) \ln^2 \frac{e\phi}{\mu} - (21 + 7\xi) \ln \frac{e\phi}{\mu} + \frac{47}{4} + \frac{7\pi^2}{24} + \frac{15}{4} \xi + \frac{3\pi^2}{8} \xi \right] + \dots, \end{aligned} \quad (5.18)$$

$$\begin{aligned} \mathcal{M}_D &= \text{Diagram D} = -2\hbar^2 e^3 \phi \int \frac{d^d k_1}{(2\pi)^d} \int \frac{d^d k_2}{(2\pi)^d} (k_1^\mu - k_2^\mu) \Delta_{\mu\nu}(k_1 + k_2) T_{\nu 2}(k_2) D_{11}(k_1) \\ &= \frac{\hbar^2 \phi^4 e^6}{(16\pi^2)^2} \xi \left[ -12 \ln^2 \frac{e\phi}{\mu} + 14 \ln \frac{e\phi}{\mu} - \frac{15}{2} - \frac{3\pi^2}{4} \right] + \dots \end{aligned} \quad (5.19)$$

with the  $\dots$  vanishing as  $\lambda \rightarrow 0$ . Kang computed some of the logarithmic terms in these amplitudes in Ref. [31], and we agree with his results. We also confirm the observation of Ref. [34] that the sum of graphs  $B$ ,  $C$  and  $D$  are gauge invariant on their own. We have supplemented the integrand-level observation of Ref. [34] with explicit calculations, which are required for the full two-loop potential.

Adding the contribution of these graphs to the counterterm contribution in Eq. (5.4) gives the  $\hbar^2$  contributions to the effective potential:

$$V_2 = \left( \frac{\hbar}{16\pi^2} \right)^2 e^6 \phi^4 \left[ (10 - 6\xi) \ln^2 \frac{e\phi}{\mu} + \left( -\frac{62}{3} + 4\xi - \frac{3}{2} \xi \ln \frac{\lambda\xi}{6e^2} \right) \ln \frac{e\phi}{\mu} + \xi \left( -\frac{1}{2} + \frac{1}{4} \ln \frac{\lambda\xi}{6e^2} \right) + \frac{71}{6} \right] + \dots \quad (5.20)$$

where again the  $\dots$  vanish as  $\lambda \rightarrow 0$ .

### C. RGE cross-check

As a cross-check, it is easy to verify that  $V = V_0 + V_1 + V_2$  satisfies the renormalization group equation

$$\left( \mu \frac{\partial}{\partial \mu} - \gamma \phi \frac{\partial}{\partial \phi} + \beta_e \frac{\partial}{\partial e} + \beta_\lambda \frac{\partial}{\partial \lambda} \right) V = 0 \quad (5.21)$$

up to order  $\hbar^2$ . The anomalous dimensions and  $\beta$ -function coefficients in scalar QED up to two-loop order can be extracted from Refs. [35,36]. We find

$$\begin{aligned} \gamma &= \frac{\hbar}{16\pi^2} e^2 (\xi - 3) + \frac{\hbar^2}{(16\pi^2)^2} \left( \frac{10}{3} e^4 + \frac{1}{9} \lambda^2 \right) + \dots, \\ \beta_\lambda &= \frac{\hbar}{16\pi^2} \left( 36e^4 - 12e^2\lambda + \frac{10\lambda^2}{3} \right) \\ &\quad + \frac{\hbar^2}{(16\pi^2)^2} \left( -416e^6 + \frac{316e^4\lambda}{3} + \frac{56e^2\lambda^2}{3} - \frac{20\lambda^3}{3} \right) + \dots, \\ \beta_e &= \frac{\hbar}{16\pi^2} \frac{e^3}{3} + \dots. \end{aligned} \quad (5.22)$$

Using these expansions, our two-loop effective potential indeed satisfies the RGE. A similar cross-check was done in Landau gauge for the two-loop Standard Model potential in Ref. [29] and for general renormalizable theories in Ref. [30].

In fact, the RGE could have been used to bootstrap all of the  $\ln \frac{\phi}{\mu}$  dependence in the two-loop effective potential. Note however that the  $\ln \lambda$  and  $\ln e$  terms in the one-loop effective potential are critical to determining the  $\ln \frac{\phi}{\mu}$  dependence at two loops, and these terms are not fixed by RG invariance. In other words, we needed the exact one-loop potential to determine the two-loop  $\ln \frac{\phi}{\mu}$  dependence. Coleman and Weinberg argue (in Appendix A2 of Ref. [4]) that one can ignore logarithms of couplings when working in a subtraction scheme with  $\lambda \equiv \frac{d^4 V}{d\phi^4} \Big|_\mu$  for a fixed scale  $\mu$ . This may be true, but since the fourth derivative of the potential is not gauge invariant (even at an extremum), this renormalization condition induces unnecessary complications when studying gauge dependence. Moreover, in  $\overline{\text{MS}}$ , logarithms of couplings play an important role and must be kept.

While we are on the subject of RG invariance, note that the  $\overline{\text{MS}}$   $\beta$ -function coefficients for  $e$  and  $\lambda$  are gauge invariant. This is true to all orders, since the  $\beta$  functions describe the evolution of couplings which can in principle be measured from scattering experiments. The anomalous dimension for  $\phi$  is gauge dependent; it describes the evolution of the field strength  $Z_\phi$  for  $\phi$ , which is not itself observable. We will return to the RG when considering the resummed potential in Sec. VIII.

### VI. DAISY RESUMMATION

As pointed out by Nielsen [17,37], there can be contributions to the effective potential proportional to  $\lambda^{-1} e^{10}$  coming from *daisy diagrams*, like those in Fig. 3. When  $\lambda \sim e^4$ , these terms are of the same order as the  $e^2 \lambda$  terms in the one-loop potential and the  $e^6$  terms in the two-loop potential. Thus they should be relevant to showing the gauge invariance of physical quantities at order  $e^6$  in this power counting. Nielsen argued that the reason the scalar mass Kang calculated does not satisfy his identity was due to the absence of these terms [17]. Some time later, Johnston [38] showed how these terms can be summed into a dressed propagator for the scalar fields, suggesting that these terms could be computed and the Nielsen identity restored. Shortly afterwards Bazeia [39] showed that even at one loop, where daisies are irrelevant, the vacuum energy in the Coleman-Weinberg model has gauge dependence. Part of the contribution of daisy resummation to the vector and scalar masses in Landau and unitary gauges was evaluated in Ref. [40]. In this section, we compute all of the daisy graphs relevant at  $e^6$  in a general  $R_\xi$  gauge. In the next section we demonstrate that, after carefully keeping track of the independent variables, the effective potential at its minimum is indeed gauge invariant.

Before beginning, it is worth remarking that the daisy resummation we perform here is related to, but not identical to, daisy resummation in finite-temperature field theory [41–43]. At finite temperature, the resummation of daisy diagrams is necessary to calculate a critical temperature because, in the limit  $m \ll T$ , new infrared divergences arise. Daisy resummation as a solution to infrared problems associated with massless Goldstone bosons has recently been proposed in Refs. [44,45]. The relevance of daisy resummation to solving gauge-dependence problems at finite temperature has also been discussed [21,46,47]. It is therefore not surprising that daisy resummation is relevant to the gauge-dependence problem in the Coleman-Weinberg model. Nevertheless, the relevant calculations have never been done, to our knowledge, which is why we resum the relevant daisy graphs here.

Normally, one does not get inverse powers of coupling constants from Feynman diagrams. Indeed, at any fixed order in perturbation theory, there are always positive powers of couplings, even in effective-potential calculations. However, the effective potential always involves





FIG. 3. Example daisy graphs which contribute inverse powers of  $\lambda$ .

summing an infinite number of graphs, namely those with an arbitrary number of background-field insertions. It is this infinite sum which gives the  $\ln \phi$  dependence in the effective potential and which can generate inverse powers of couplings. As discussed in Sec. V, we simplify the infinite sums by using dressed propagators. For example, we see from Eq. (5.5) that the  $\phi_1$  propagator is  $D_{11} = \frac{i}{k^2 - \frac{\lambda}{2}\phi^2}$

which has an effective mass  $m^2 = \frac{\lambda}{2}\phi^2$ . In the daisy graphs, each photon loop (the petals) gives a factor proportional to  $e^2\phi^2$ , each vertex gives a factor of  $e^2$ , and the loop integral over the scalar propagators can give inverse powers of the effective mass. For example, a four-loop three-petal daisy will give



$$\propto (e^2)^3 (e^2\phi^2)^3 \int \frac{d^4k}{2\pi^4} \left( \frac{i}{k^2 - \frac{\lambda}{2}\phi^2} \right)^3 \propto \phi^4 \frac{e^{12}}{\lambda}. \quad (6.1)$$

Here, we have simply done the integral by dimensional analysis, since it is UV and IR finite. This graph therefore contributes at order  $e^8$  in the  $\lambda \sim e^4$  power counting. It is therefore beyond the order we need for the first nontrivial gauge-invariance check of  $V_{\min}$  (not to mention that this particular loop is itself  $\xi$  independent).

It is not hard to see, using dimensional analysis, that the only graphs which could contribute at order  $e^6$  (with the  $\lambda \sim e^4$  power counting) must involve  $\phi_2$  propagators and have petal-type photon loops. The petals factorize off from

the central disk, and from each other. Thus a daisy graph with  $n$  petals has the form

$$I_n = i^{n+1} \frac{e^{2n}}{2n} A_{n\phi_2} (A_\gamma)^n \quad (6.2)$$

where  $A_\gamma$  includes the photon loop and its counterterm,  $A_{n\phi_2}$  is the scalar integral with  $n$   $\phi_2$  propagators, and the factor  $\frac{1}{2n} e^{2n}$  comes from the vertex Feynman rule and the symmetry factor for the circle.


Each photon loop gives

$$\begin{aligned} A_\gamma^{\text{loop}} &= \text{[diagram of a photon loop with a counterterm dot]} = \int \frac{d^d k}{(2\pi)^d} \Delta_{\mu\mu} \\ &= \frac{\hbar}{16\pi^2} e^2 \phi^2 \left[ \frac{3}{\varepsilon} + \left( 1 - 6 \ln \frac{\phi e}{\mu} \right) + \varepsilon \left( 1 + \frac{\pi^2}{4} - 2 \ln \frac{\phi e}{\mu} + 6 \ln^2 \frac{\phi e}{\mu} \right) + \dots \right]. \end{aligned} \quad (6.3)$$

The UV pole in this loop is removed by the one-loop counterterm:

$$A_\gamma^{\text{c.t.}} = \frac{\hbar}{16\pi^2} e^2 \phi^2 \left[ -\frac{3}{\varepsilon} \right]. \quad (6.4)$$

The  $\mathcal{O}(\varepsilon)$  in  $A_\gamma^{\text{loop}}$  can be important if the scalar loop it multiplies is UV divergent. This only happens for two petals ( $n = 2$ ). For  $n = 2$ , the scalar loop gives



$$= \int \frac{d^d k}{(2\pi)^d} D_{22}^2 = -i \frac{\hbar}{16\pi^2} \left[ \frac{1}{\varepsilon} + \frac{3e^2\xi}{\lambda} + \dots \right]. \quad (6.5)$$

Since the  $\frac{1}{\varepsilon}$  term in  $A_{2\phi_2}$  has no  $\frac{1}{\lambda}$  piece, the cross term between it and the  $\mathcal{O}(\varepsilon)$  part of  $A_\gamma^{\text{loop}}$  will not contribute at order  $e^6$  (with  $\lambda \sim e^4$ ). Thus, for all the daisies, we can drop the  $\mathcal{O}(\varepsilon)$  terms in  $A_\gamma^{\text{loop}}$  and take

$$A_\gamma = A_\gamma^{\text{loop}} + A_\gamma^{\text{c.t.}} = \frac{\hbar}{16\pi^2} e^2 \phi^2 \left( 1 - 6 \ln \frac{e\phi}{\mu} \right). \quad (6.6)$$

For  $n > 3$ , the scalar loop is UV and IR finite. We find

$$A_{n\phi_2} = \text{Diagram} = \int \frac{d^4 k}{(2\pi)^4} D_{22}^n = \frac{i\hbar}{16\pi^2} \phi^4 \frac{e^2 \lambda \xi}{12(n-1)} \left( \frac{-6i}{\lambda \phi^2} \right)^n \quad (6.7)$$

so that

$$I_n = \frac{\hbar}{16\pi^2} \left( -\frac{1}{24} e^2 \lambda \xi \phi^4 \right) \frac{1}{n(n-1)} \left[ \frac{\hbar e^4}{(16\pi^2) \lambda} \left( 6 - 36 \ln \frac{e\phi}{\mu} \right) \right]^n. \quad (6.8)$$

Each term in this series contributes at order  $e^6$  when  $\lambda \sim e^4$ . Thus they are all equally important for checking gauge invariance and we must sum the series. Summing the series is easy enough to do using

$$\sum_{n=2}^{\infty} \frac{1}{n(n-1)} x^n = x + (1-x) \ln(1-x) \quad (6.9)$$

which gives

$$V^{e^6 \text{ daisies}} = \phi^4 \frac{\hbar}{16\pi^2} \left( -\frac{e^2 \lambda \xi}{24} \right) \times \left[ \frac{\hat{\lambda}(\phi)}{\lambda} + \left( 1 - \frac{\hat{\lambda}(\phi)}{\lambda} \right) \ln \left( 1 - \frac{\hat{\lambda}(\phi)}{\lambda} \right) \right] \quad (6.10)$$

where

$$\hat{\lambda}(\phi) \equiv \frac{\hbar e^4}{16\pi^2} \left( 6 - 36 \ln \frac{e\phi}{\mu} \right). \quad (6.11)$$

We have defined  $\hat{\lambda}(\phi)$  so that according to Eq. (4.6)  $\hat{\lambda}(v) = \lambda$  at the scale  $v$  where the one-loop potential has its minimum. Remarkably, while each daisy graph with  $n > 1$  is individually power divergent as  $\lambda \rightarrow 0$  with  $e$  fixed, the sum of all daisies scales only like  $\ln \lambda$ . An alternative approach to calculating Eq. (6.10) is outlined at the end of Ref. [37].

One might also be concerned about graphs where petals attach to the  $\phi_2$  legs in the two-loop diagrams  $\mathcal{M}_B$ ,  $\mathcal{M}_C$  and  $\mathcal{M}_D$  in Eqs. (5.17)–(5.19) (petals attached to diagram  $\mathcal{M}_A$  are the same daisy chain we have already summed). Since petals are not suppressed, it seems like these should be of the same order. However, these two-loop daisies are in fact subleading. A quick way to see that they are subleading is to note that to order  $e^6$ ,  $\mathcal{M}_B$ ,  $\mathcal{M}_C$  and  $\mathcal{M}_D$  are all independent of  $\lambda$ . Thus they are independent of the scalar

masses. Since the daisy resummation can be thought of as correcting the propagators and shifting the masses (see next paragraph), they have no effect on expressions that are independent of the masses, such as  $\mathcal{M}_B$ ,  $\mathcal{M}_C$  and  $\mathcal{M}_D$ . Similarly, it is not hard to show that all self-energy corrections other than the one we included contribute first at order  $e^8$ . Thus we have included all the graphs relevant at order  $e^6$ .

Before moving on, it is worth pointing out that daisy resummation is important even in Landau gauge,  $\xi = 0$ . In Landau gauge, there is no kinetic mixing and the scalar propagators are

$$D_{11} = \frac{i}{k^2 - \frac{\lambda}{2} \phi^2}, \quad D_{22} = \frac{i}{k^2 - \frac{\lambda}{6} \phi^2}. \quad (6.12)$$

These propagators still have  $\lambda$ -dependent masses and can produce  $\frac{1}{\lambda}$  dependence from daisy graphs. Since there is no kinetic mixing, and since the scalar propagators are of the conventional form with single poles, the daisy resummation can be done by summing the geometric series of self-energy graphs. That is, the daisy summation along a  $\phi_1$  line amounts to replacing

$$D_{11} \rightarrow \hat{D}_{11} = D_{11} \sum_{n=0}^{\infty} \left( \frac{ie^2}{2} D_{11} A_\gamma \right)^n = \frac{i}{k^2 - \frac{\lambda}{2} \phi^2 + \frac{\hat{\lambda}}{6} \phi^2}. \quad (6.13)$$

Thus the new effective mass squared for  $\phi_1$  is  $m_1^2 = \frac{3\lambda - \hat{\lambda}}{6} \phi^2$ . Similarly, the new effective mass squared for  $\phi_2$  is  $m_2^2 = \frac{\lambda - \hat{\lambda}}{6} \phi^2$ . Note that at the scale where the one-loop potential has its minimum,  $\lambda = \hat{\lambda}$ , and the Goldstone mode  $\phi_2$  becomes massless, as observed in Ref. [40]. The net result is that the sum of all  $\phi_1$  or  $\phi_2$  daisy graphs (including the one-loop zero-daisy graph) is simply of the general one-loop form, Eq. (3.9), with shifted masses:

$$V_{\xi=0}^{\text{daisies}} = \frac{\phi^4}{64\pi^2} \left\{ \left( \frac{3\lambda - \hat{\lambda}}{6} \right)^2 \left[ \ln \frac{(3\lambda - \hat{\lambda})\phi^2}{6\mu^2} - \frac{3}{2} \right] + \left( \frac{\lambda - \hat{\lambda}}{6} \right)^2 \left[ \ln \frac{(\lambda - \hat{\lambda})\phi^2}{6\mu^2} - \frac{3}{2} \right] \right\}. \quad (6.14)$$

Since  $\lambda \sim e^4$ , these terms have effects comparable to terms in the three-loop Coleman-Weinberg potential. Thus, while the infinite sum of daisy graphs is always important, an advantage of Landau gauge is that it postpones the relevance of daisy resummation by one loop. Landau gauge does not however let us ignore the daisy graphs completely.

Two recent papers also observed that the resummation of certain graphs to all orders is necessary starting at three loops in Landau gauge [44,45]. These two papers were concerned with resolving an infrared-divergence problem associated with massless Goldstone bosons starting at three loops. While these two papers discussed diagrams similar to the ones here, the problem they solved was different (infrared divergences, not gauge dependence) and their results are not directly transferable. However, these two papers, along with the earlier work in Refs. [38,40,48], do explain in a more systematic way how daisy and other

relevant diagrams can be resummed through a modification of the effective propagators.

In summary, the full Coleman-Weinberg potential up to order  $e^6$  with  $\lambda \sim e^4$  is the sum of Eqs. (4.1), (4.3), (5.20) and (6.10). It is helpful to write the result as

$$V = V^{\text{LO}} + V^{\text{NLO}} + \dots \quad (6.15)$$

where the leading-order (LO) potential

$$V^{\text{LO}} = \frac{\lambda}{24} \phi^4 + \frac{\hbar e^4}{16\pi^2} \phi^4 \left( -\frac{5}{8} + \frac{3}{2} \ln \frac{e\phi}{\mu} \right) \quad (6.16)$$

scales as  $\mathcal{O}(\hbar)$  when  $\lambda \sim \hbar e^4$  and the NLO potential, scaling like  $\mathcal{O}(\hbar^2)$ , is

$$\begin{aligned} V^{\text{NLO}} = & \frac{\hbar e^2 \lambda}{16\pi^2} \phi^4 \left( \frac{\xi}{8} - \frac{\xi}{24} \ln \frac{e^2 \lambda \xi \phi^4}{6\mu^4} \right) \\ & + \frac{\hbar^2 e^6}{(16\pi^2)^2} \phi^4 \left[ (10 - 6\xi) \ln^2 \frac{e\phi}{\mu} + \left( -\frac{62}{3} + 4\xi - \frac{3}{2} \xi \ln \frac{\lambda \xi}{6e^2} \right) \ln \frac{e\phi}{\mu} + \xi \left( -\frac{1}{2} + \frac{1}{4} \ln \frac{\lambda \xi}{6e^2} \right) + \frac{71}{6} \right] \\ & + \phi^4 \frac{\hbar e^2 \lambda}{16\pi^2} \left( -\frac{\xi}{24} \right) \left[ \frac{\hat{\lambda}(\phi)}{\lambda} + \left( 1 - \frac{\hat{\lambda}(\phi)}{\lambda} \right) \ln \left( 1 - \frac{\hat{\lambda}(\phi)}{\lambda} \right) \right] \end{aligned} \quad (6.17)$$

where  $\hat{\lambda}(\phi)$  is defined in Eq. (6.11). Note that there are tree and one-loop contributions to the LO potential and that the NLO potential get contributions from one, two and all higher-order loops.

## VII. GAUGE INVARIANCE OF $V_{\min}$

With the two-loop Coleman-Weinberg potential in hand and the contribution of daisy graphs understood, we can now demonstrate the gauge invariance of the potential at its minimum.

Recall from Eq. (4.13) that at LO,  $V_{\min}^{\text{LO}} = -v^4 e^4 \frac{3}{128\pi^2}$ . One might naturally expect that the NLO contribution could be written as  $V_{\min}^{\text{NLO}} = v^4 e^6 C$  for some  $\xi$ -independent constant  $C$ . Unfortunately, we cannot expect  $V_{\min}$  to be explicitly  $\xi$  independent when written in terms of  $v$ . The problem is that  $v = \langle \phi \rangle$  is a field value and therefore gauge dependent, so that it infects all dimensionful quantities expressed in terms of it. One alternative is to calculate not  $V_{\min}$  but rather the ratio of  $V_{\min}$  to some other dimension-four quantity, such as  $m_S^4$ , where  $m_S$  is the scalar mass. Indeed, Coleman and Weinberg [4] and

later Kang [31] discussed the gauge invariance of the dimensionless ratio of the scalar-to-vector masses. However, we would really like to see whether  $V_{\min}$  is physical on its own. But then, if not  $v$ , what are we to express  $V_{\min}$  in terms of?

An alternative to expressing  $V_{\min}$  in terms of  $v$  is to express it in terms of the renormalization group scale  $\mu$ . This scale is as physical as the  $\overline{\text{MS}}$  couplings; the two are intrinsically connected. To be concrete, let us define the scale  $\mu_X$  as the scale where Eq. (4.7) is satisfied exactly. That is,  $\mu_X$  is defined by the exact relation

$$\lambda(\mu_X) \equiv \frac{\hbar}{16\pi^2} e^4(\mu_X) \{6 - 36 \ln[e(\mu_X)]\}. \quad (7.1)$$

Since this relation is exact, we no longer can or need to solve for  $e^6$  terms in the relation between  $\lambda$  and  $e$ , as was done in Ref. [31]. One can instead now solve for corrections to  $v$

$$v = \langle \phi \rangle = \mu_X + v_1 + v_2 + \dots \quad (7.2)$$



with  $v_1 \sim e(\mu_X)^2$ ,  $v_2 \sim e(\mu_X)^4$ , etc., and  $\mu_X$  is defined by Eq. (7.1). Then one can consistently expand  $V_{\min}$  at  $\mu = \mu_X$ :

$$\begin{aligned}
 V_{\min} = & V^{\text{LO}}(\mu_X) + V^{\text{NLO}}(\mu_X) + \overbrace{v_1 \frac{d}{d\phi} V^{\text{LO}}|_{\phi=\mu_X}}^{=0} \\
 & + V^{\text{NNLO}}(\mu_X) + v_1 \frac{d}{d\phi} V^{\text{NLO}}|_{\phi=\mu_X} \\
 & + \underbrace{v_2 \frac{d}{d\phi} V^{\text{LO}}|_{\phi=\mu_X}}_{=0} + \frac{1}{2} v_1^2 \frac{d^2}{d\phi^2} V^{\text{LO}}|_{\phi=\mu_X} + \dots
 \end{aligned} \tag{7.3}$$

where  $V^{\text{LO}}$  is the effective potential truncated to order  $\hbar$  with  $\lambda \sim \hbar$ ,  $V^{\text{NLO}}$  is truncated to  $\hbar^2$  and so on, as in Eqs. (6.16) and (6.17).

A convenient feature of setting  $\phi = \mu_X$  is that the daisy contribution simplifies. Indeed, from Eq. (6.11) we see that when  $\phi = \mu = \mu_X$  then  $\hat{\lambda}(\mu_X) = \lambda(\mu_X)$ . Then the entire daisy contribution in the last line of Eq. (6.17) reduces to simply

$$V^{\text{NLO,daisies}} = -\frac{\xi}{24} \phi^4 \frac{\hbar e^2 \lambda}{16\pi^2} = -\frac{\xi}{24} \frac{\hbar^2 e^6}{(16\pi^2)^2} (6 - 36 \ln e). \tag{7.4}$$

Also, as indicated in Eq. (7.3), we can use the fact that  $V^{\text{LO}'}(\mu_X) = 0$ , which was the defining equation for  $\mu_X$ . This simplifies  $V_{\min}$  to

$$V_{\min} = V^{\text{LO}}(\mu_X) + V^{\text{NLO}}(\mu_X) + \dots \tag{7.5}$$

$$\begin{aligned}
 \Lambda_I = \mu_I - \frac{V^{\text{NLO}}(\mu_I)}{V^{\text{LO}'}(\mu_I)} + \dots = \mu_I + \mu_I \frac{\hbar e^2}{16\pi^2} \left\{ -\frac{71}{9} - \frac{11\xi}{12} + \left( \frac{5\xi}{2} + \frac{124}{9} \right) \ln e - \frac{20}{3} \ln^2 e + \frac{1}{4} \xi \ln \left[ \frac{h\xi}{32\pi^2} (5 - 12 \ln e) \right] \right. \\
 \left. - \frac{\xi}{12} \left( -2 + 12 \ln e - 3 \ln \frac{3}{5 - 12 \ln e} \right) \right\} + \dots
 \end{aligned} \tag{7.10}$$

This instability scale is linearly dependent on the gauge parameter  $\xi$ , and therefore should not be used to draw physical conclusions. The  $\xi$  dependence of other field values can be computed in a similar way, confirming that they are indeed unphysical.

### VIII. RENORMALIZATION GROUP IMPROVEMENT

At this point we have shown that if the effective potential in scalar QED is expressed in terms of the  $\overline{\text{MS}}$  couplings  $e$

Plugging this into Eqs. (6.16) and (6.17) and using Eq. (7.1) we then find

$$V_{\min} = -\frac{3\hbar e^4}{128\pi^2} \mu_X^4 + \frac{e^6 \hbar^2}{(16\pi^2)^2} \mu_X^4 \left( \frac{71}{6} - \frac{62}{3} \ln e + 10 \ln^2 e \right) \tag{7.6}$$

which is manifestly gauge invariant! The daisies have exactly canceled the  $\xi$  dependence of the NLO one-loop and two-loop contributions.

Next, let us look at a field value expressed in terms of the  $\overline{\text{MS}}$  scale  $\mu$ , to double-check that somehow all of its gauge dependence is not miraculously absent. Consider the value of the field where the potential is zero,  $\Lambda_I$ , which in the Standard Model is sometimes given an interpretation as an instability scale [49]. Setting  $V(\Lambda_I) = 0$  gives a different relation between  $\lambda$  and  $e$  than  $V'(v) = 0$  did. The condition on the running couplings so that  $V^{\text{LO}} = 0$  at  $\phi = \mu = \mu_I$  is

$$\lambda(\mu_I) = \frac{\hbar}{16\pi^2} e(\mu_I)^4 \{15 - 36 \ln[e(\mu_I)]\}. \tag{7.7}$$

At NLO, we then find

$$\Lambda_I = \mu_I - \frac{V^{\text{NLO}}(\mu_I)}{V^{\text{LO}'}(\mu_I)} + \dots = \mu_I - \frac{32\pi^2}{3e^4 \hbar \mu_I^3} V^{\text{NLO}}(\mu_I). \tag{7.8}$$

To evaluate the daisy contribution, we can no longer use  $\hat{\lambda} = \lambda$ . Instead we now find

$$\frac{\hat{\lambda}}{\lambda} = \frac{\hat{\lambda}(\phi = \mu = \mu_I)}{\lambda(\mu = \mu_I)} = 1 - \frac{3}{5 - 12 \ln e}. \tag{7.9}$$

Up to NLO, we then find

and  $\lambda$  and the scale  $\mu_X$  where they satisfy Eq. (7.1), then the value of the potential at the quantum minimum  $V_{\min}$  is gauge invariant. We checked this to the first nonleading order, which required the two-loop potential and the summation of the leading daisy diagrams. A natural question is whether we *must* express  $V_{\min}$  in terms of  $\mu_X$ . Clearly, we should be able to write any physical quantity in terms of any other scale. More importantly, the check required the truncation of the effective potential to order  $e^6$  with  $\lambda(\mu_X) \sim e(\mu_X)^4$ . We would like to be able

to use effective potentials in other contexts, where the couplings are defined at some other arbitrary scale  $\mu_Y$  and the renormalization group is used to evolve the potential to a scale near its minimum. So how are we to use this unusual truncation in an RG-improved effective potential, where some  $\lambda$  and  $e$  dependence is necessarily included to all orders in  $\hbar$ ? In this section, we show how to answer both of these questions, and that the answers are related.

### A. Calculation-scale invariance

First, let us consider how the calculation of  $V_{\min}$  would change if we had calculated the effective potential with the couplings defined at a scale  $\mu_Y$  instead of  $\mu_X$ . At  $\mu_Y$ , the potential obviously has the same form, Eq. (6.17), but with  $e$  and  $\lambda$  interpreted as  $e_Y \equiv e(\mu_Y)$  and  $\lambda_Y \equiv \lambda(\mu_Y)$  rather than  $e_X \equiv e(\mu_X)$  and  $\lambda_X \equiv \lambda(\mu_X)$ . That is,

$$V = \frac{\lambda_Y}{24} \phi^4 + \frac{\hbar}{16\pi^2} \phi^4 \left[ e_Y^4 \left( -\frac{5}{8} + \frac{3}{2} \ln \frac{e_Y \phi}{\mu_Y} \right) + e_Y^2 \lambda_Y \xi \left( \frac{1}{8} - \frac{1}{24} \ln \frac{e_Y^2 \lambda_Y \xi \phi^4}{6\mu_Y^4} \right) \right] + \dots \quad (8.1)$$

For simplicity, let us assume for now that  $\mu_Y$  is close enough to  $\mu_X$  so that  $\lambda_Y$  is still much smaller than  $e_Y^2$ . For the daisy contribution evaluated at  $\mu_Y$ , we can then still use  $\hat{\lambda} \sim \lambda$ , since the difference is of higher order. This lets us continue to drop most of Eq. (6.10).

The easiest way to study  $V$  with couplings at  $\mu_Y$  is simply to use the renormalization group to move the couplings back to  $\mu_X$ . Then we can recycle the previous analysis. Expanding Eq. (4.11) gives

$$e_Y = e_X + \frac{\hbar}{48\pi^2} e_X^3 \ln \frac{\mu_Y}{\mu_X} + \dots \quad (8.2)$$

Expanding Eq. (4.12) gives

$$\lambda_Y = \lambda_X + \frac{\hbar}{16\pi^2} [36e_X^4 - 12e_X^2 \lambda_X] \ln \frac{\mu_Y}{\mu_X} + \frac{\hbar^2}{(16\pi^2)^2} e_X^6 \left[ -416 \ln \frac{\mu_Y}{\mu_X} - 192 \ln^2 \frac{\mu_Y}{\mu_X} \right] + \dots \quad (8.3)$$

By plugging these into Eq. (8.1) we find

$$V = \frac{\lambda_X}{24} \phi^4 + \frac{\hbar}{16\pi^2} \phi^4 \left[ e_X^4 \left( -\frac{5}{8} + \frac{3}{2} \ln \frac{e_X \phi}{\mu_X} \right) + e_X^2 \lambda_X \left( \frac{\xi}{8} - \frac{\xi}{24} \ln \frac{e_X^2 \lambda_X \xi \phi^4}{6\mu_X^4} + \frac{\xi - 3}{6} \ln \frac{\mu_Y}{\mu_X} \right) \right] + \dots \quad (8.4)$$

We have not even shown the two-loop terms, but already one can see that this potential is not identical to what we got from computing  $V$  starting at  $\mu = \mu_X$ : there is an uncancelled  $\ln \frac{\mu_Y}{\mu_X}$  term. Despite this modification of the potential, we still find that

$$V_{\min} = \frac{e^4 \hbar}{16\pi^2} \mu_X^4 \left( -\frac{3}{8} \right) + \frac{e^6 \hbar^2}{(16\pi^2)^2} \mu_X^4 \left( \frac{71}{6} - \frac{62}{3} \ln e + 10 \ln^2 e \right) + \dots \quad (8.5)$$

exactly as in Eq. (7.6). That is, the new  $\ln \frac{\mu_Y}{\mu_X}$  term has no effect on the value of the potential at the minimum.

That the  $\mu_Y$  dependence would drop out of  $V_{\min}$  was anticipated in Sec. II B, with  $\mu_0$  replaced by  $\mu_Y$  and  $\mu$  replaced by  $\mu_X$ . We next review and extend that discussion in the context of this concrete example.

### B. Discussion

There is a quick way to see why computing the potential at a different scale affected  $V(\phi)$  but not  $V_{\min}$ . The effective potential satisfies the RGE in Eq. (5.21):

$$\left( \mu \frac{\partial}{\partial \mu} - \gamma \phi \frac{\partial}{\partial \phi} + \beta_e \frac{\partial}{\partial e} + \beta_\lambda \frac{\partial}{\partial \lambda} \right) V = 0. \quad (8.6)$$

This equation says that the explicit  $\mu$  dependence in the effective potential is exactly compensated for by a rescaling of the coupling constants, according to their  $\beta$  functions, and a rescaling of the field  $\phi$ , according to its anomalous dimension. In writing the potential computed at  $\mu_Y$  in terms of couplings at  $\mu_X$  we only rescaled the couplings, not the field. Indeed, the extra  $\frac{\xi-3}{6} \ln \frac{\mu_Y}{\mu_X}$  term in Eq. (8.4) would be exactly removed if we rescaled the field as  $\phi \rightarrow \kappa \phi$  with

$$\kappa = 1 + \frac{e^2}{16\pi^2} (3 - \xi) \ln \frac{\mu_Y}{\mu_X} + \dots \quad (8.7)$$

Because the two potentials (calculated at  $\mu_X$  or  $\mu_Y$ ) differ by terms which can be compensated by a field rescaling, the value of  $V_{\min}$  will be unchanged. To see this, simply note that the extrema of any function  $f(x)$  are invariant under a rescaling of the dependent variable: this rescaling just

scales the  $x$  axis (see Fig. 1). Thus the invariance of  $V_{\min}$  follows as a special case.

This property of the effective potential, that it depends on the scale where it is calculated, may seem unsettling. One might imagine that the  $\ln \frac{\mu_Y}{\mu_X}$  terms can be removed by invoking canonical normalization, perhaps by demanding that the kinetic term in the effective action be reset to  $\frac{1}{2}(\partial_\mu \phi)^2$  after including all the quantum corrections. One cannot just demand this, however. First of all, in the effective action even the two-derivative kinetic terms have a complicated field dependence

$$\begin{aligned} \Gamma[\phi] = & \dots + \frac{1}{2} Z_1[\phi] (\partial_\mu \phi_i) (\partial_\mu \phi_i) \\ & + \frac{1}{2} Z_2[\phi] \frac{\phi_i \phi_j}{\phi^2} (\partial_\mu \phi_i) (\partial_\mu \phi_j) \\ & + Y_1[\phi] \frac{1}{\phi^4} ((\partial_\mu \phi_i) (\partial_\mu \phi_i))^2 + \dots \end{aligned} \quad (8.8)$$

The functionals  $Z_1$ ,  $Z_2$ ,  $Y_1$  etc. can be calculated from prototype diagrams as a power expansion in the external momentum  $p^\mu$  [32]. Explicit results at one loop for  $Z_1$  and  $Z_2$  can be found in Ref. [10] and for  $Y_1$  in  $\phi^4$  theory in Ref. [32]. So it is not clear how one would go about canonically normalizing  $\phi$  apart from some byzantine implicit nonlinear field redefinition.

A more elementary objection to renormalizing the fields is that we have already renormalized them. In  $\overline{\text{MS}}$  the field-strength renormalization  $Z_1$  is unambiguously fixed, so there is no remaining freedom. One could have chosen a different normalization convention, for example, an on-shell renormalization for which the residue of the propagator at the physical pole is set to 1 exactly. However, since these are conventions, physical quantities must be independent of them. Thus we can work in  $\overline{\text{MS}}$ . Indeed, as we have seen  $V_{\min}$  is independent of the normalization condition.

### C. RG-improved effective potential

We have shown that the effective potential depends on the scale  $\mu_Y$  where it is calculated. We have also shown that when  $\mu_Y$  is close to the scale  $\mu_X$  where the couplings satisfy Eq. (7.1), then the value of  $V_{\min}$  is independent of  $\mu_Y$ . Now suppose  $\mu_Y$  is not close to  $\mu_X$ , so that  $\ln \frac{\mu_Y}{\mu_X}$  is not small. Then we cannot work in fixed-order perturbation theory. For example, in the Standard Model, one normally extracts the  $\overline{\text{MS}}$  couplings through perturbative threshold calculations at a scale  $\mu_Y \sim 100$  GeV near the weak scale. The equivalent of  $\mu_X$  is a scale near where  $\lambda \sim 0$  which is many orders of magnitude higher in energy; so,  $\mu_X \gg \mu_Y$ . The minimum of the effective potential is found by solving the RGE for the effective potential to evolve it from  $\mu_Y$  to  $\mu_X$ . Thus the potential at  $\mu = \mu_X$  includes terms to all orders in perturbation theory whether it is written in terms of  $g(\mu_Y)$  or

$g(\mu_X)$ . This seems to limit the use of fixed-order relations like Eq. (7.1).

In order to combine the  $\lambda \sim e^4$  scaling, which is required for gauge invariance, with resummation, we can exploit the observation from Sec. VIII B that  $V_{\min}$  is independent of the rescaling  $\phi$ . In particular, we see that we do not have to evolve the entire effective potential from  $\mu_Y$  to  $\mu_X$ . Instead, we can just evolve the couplings, and then compute the fixed-order potential directly at  $\mu_X$ . The scale  $\mu_Y$  only enters for determining the boundary conditions of the renormalization equations. Since the  $\beta$  functions are all gauge invariant, the gauge invariance of the potential at the minimum then automatically follows from the arguments in Sec. VII.

To be concrete, the resummed effective potential is often written as

$$\begin{aligned} V[\phi] = & \frac{1}{4!} e^{4\Gamma(\mu=\phi)} \\ & \times [\lambda_{\text{eff}}^{(0)}(\mu=\phi) + \lambda_{\text{eff}}^{(1)}(\mu=\phi) + \lambda_{\text{eff}}^{(2)}(\mu=\phi) + \dots] \phi^4 \end{aligned} \quad (8.9)$$

with  $\Gamma(\mu) = \int_{\mu_Y}^{\mu} \gamma(\mu') \frac{d\mu'}{\mu}$  and  $\frac{1}{4!} \lambda_{\text{eff}}^{(j)}(\mu) \phi^4$  is the order- $\hbar^j$  term in the fixed-order calculation of the effective potential. We are proposing instead that one should use Eq. (7.3):

$$\begin{aligned} V_{\min} = & V^{\text{LO}}(\mu_X) + V^{\text{NLO}}(\mu_X) + V^{\text{NNLO}}(\mu_X) \\ & + v_1 \frac{d}{d\phi} V^{\text{NLO}}|_{\phi=\mu_X} + \frac{1}{2} v_1^2 \frac{d^2}{d\phi^2} V^{\text{LO}}|_{\phi=\mu_X} + \dots \end{aligned} \quad (8.10)$$

where  $V^{\text{LO}} \sim \hbar$ ,  $V^{\text{NLO}} \sim \hbar^2$ ,  $v_1 \sim \hbar$  and so on, using  $\lambda \sim \hbar$ . Using Eq. (8.10), the value of  $V_{\min}$  will be gauge invariant order by order using this expansion, while  $V_{\min}$  computed using Eq. (8.9) will not be.

Note that this consistent prescription is not equivalent to using dressed fields where the field-strength renormalization is absorbed into the field. Such a redefinition was advocated in Ref. [9] (based on Ref. [8]) as removing all of the gauge dependence in the effective potential. First of all, that redefinition does not work as advertised: even after absorbing the field-strength renormalization, the effective potential is still gauge dependent [17,18]. Secondly, we are still using the same fields and  $\overline{\text{MS}}$  normalizations relevant for  $S$ -matrix calculations, which greatly facilitates applications to the Standard Model. Furthermore, since  $V^{\text{NLO}}(\mu_X)$  is known in the Standard Model [29] (and in general renormalizable theories [30]) and since the daisy graphs begin at NNLO in Landau gauge, Eq. (7.3) can immediately be implemented for many theories of direct phenomenological interest. The effect of using Eq. (8.10) instead of Eq. (8.9) for the Standard Model effective potential is discussed in Ref. [50].

## IX. HIGHER-DIMENSION OPERATORS

As we have seen,  $V_{\min}$  is gauge invariant but the VEV at the minimum,  $v = \langle \phi \rangle$  is not. One reason one would like  $v$  to be physical is that it gives a criteria for when higher-dimension operators can have a significant effect. For example, one might expect an operator  $\Delta\mathcal{L} = -\frac{1}{\Lambda^2}\phi^6$  to be relevant at field values where this term contributes at the same order as the leading one:  $\Lambda \sim \frac{1}{\sqrt{\lambda(v)}}v$ . Since  $v$  is  $\xi$  dependent, this criterion is not meaningful.

The correct way to evaluate the influence of a gauge-invariant higher-dimension operator is to add it to the classical potential and see how  $V_{\min}$  is affected. So let us add  $-\frac{1}{\Lambda^2}\phi^6$  to the Lagrangian (that is, we add  $+\frac{1}{\Lambda^2}\phi^6$  to the classical potential). First, consider the case when  $\Lambda$  is very small (so this new term is large). Then the sensitive cancellation between  $\lambda$  and  $e^4$  will be severely disrupted and the new quantum minimum will disappear. With the sign given, the only minimum would be at  $\phi = 0$  with  $V_{\min} = 0$ . If we flip the sign, then the potential will be unbounded from below and again there will be no quantum minimum, with  $V = 0$  still the only extremum and clearly gauge invariant.

Now suppose this new term has a very small effect. To be precise, recall that the quantum minimum in the absence of this term occurs at leading order at the scale  $\phi = \mu_X$  where Eq. (7.1) is satisfied. Thus if this term is very small, at LO, it shifts the minimum to

$$V_{\min} = \mu_X^4 \left[ -\frac{3}{8} \frac{e^4 \hbar}{16\pi^2} + \frac{\mu_X^2}{\Lambda^2} + \mathcal{O}\left(\frac{\mu_X^4}{\Lambda^4}\right) \right]. \quad (9.1)$$

This is gauge invariant, but trivially so.

The key observation is that to compute subleading corrections when the effect is small, we cannot simply plug the next-to-leading-order VEV into  $\frac{1}{\Lambda^2}\phi^6$ . Doing so would give a gauge-dependent correction. Instead, we must carefully reevaluate our perturbation expansion in the presence of this new term.

If we are careful we must commit to whether this new term is important or not. If we say it is important, then minimally, the new term in Eq. (9.1) must be comparable to the original one. That means that  $\Lambda^2 \sim \mu_X^2 \frac{16\pi^2}{e^4 \hbar}$ . This power counting means that the one-loop minimum is not at the scale  $\mu_X$  but rather at a new scale  $\mu_Z$  where

$$\lambda(\mu_Z) \equiv \frac{\hbar}{16\pi^2} e^4(\mu_Z) [6 - 36 \ln[e(\mu_Z)]] - 36 \frac{\mu_Z^2}{\Lambda^2} \quad (9.2)$$

which gives

$$V_{\min} = \mu_Z^4 \left[ -\frac{3}{8} \frac{e^4 \hbar}{16\pi^2} - \frac{1}{2} \frac{\mu_Z^2}{\Lambda^2} + \mathcal{O}(e^6) \right]. \quad (9.3)$$

As before, the first nontrivial check will be at order  $e^6$ . To perform this check, one would need to compute the

two-loop graphs with this modification as well as the daisy resummation in the  $R_\xi$  gauges.

The main implication of the discussion in this section is that arguments about the relevance of higher-dimension operators must be refined to have a consistent perturbative expansion. For example, one might traditionally calculate  $\phi_{\min}$  defined as the field value at the minimum in Landau gauge, and compare it to  $M_{\text{Pl}}$  to see gravitational effects on the effective potential. This comparison is not gauge invariant. Instead, one must actually insert the new operators and see how physical quantities determined by the effective potential change. An example implementation of this procedure for the Standard Model was discussed in Ref. [50].

## X. CONCLUSIONS

In this paper we have resolved a long-standing puzzle about gauge dependence in the Coleman-Weinberg model. We have shown that the potential at the minimum is gauge invariant at the first nontrivial order in perturbation theory. The invariance is only manifest if an appropriate power counting is used and if  $V_{\min}$  is expressed in terms of another gauge-invariant scale, such as a value of the  $\overline{\text{MS}}$  parameter  $\mu$  where the couplings satisfy some relation. To establish gauge invariance required a calculation of the full two-loop Coleman-Weinberg potential in a general gauge to order  $e^6$  as well as an infinite number of daisy diagrams which are also relevant at next-to-leading order.

Besides providing the first explicit check that a potential is gauge invariant at a loop-generated minimum, our work has implications for how effective potentials should be used consistently. First of all, we showed that one cannot simply truncate to fixed order in the usual loop expansion. Some terms at a given order must be dropped, and other terms from diagrams of arbitrarily high-loop order must be included. Only then will the potential have the desired gauge-invariance properties. At leading order in the appropriate power counting, only a finite number of diagrams contribute. At next-to-leading order, the first infinite class of diagrams becomes relevant in a general gauge and must be resummed. A propitious feature of Landau gauge,  $\xi = 0$ , is that this resummation is postponed to next-to-next-to-leading order, competitive with three-loop graphs in the normal loop expansion. Nevertheless, even in Landau gauge at NLO, one *must* truncate the effective potential appropriately to have self-consistent results. This result is consistent with recent observations in Refs. [44,45].

We also discussed how the consistent use we advocate is compatible with an RG-improved effective potential. The key is to run the couplings first to a scale where the one-loop potential, appropriately truncated, has its minimum. This scale is gauge invariant. Then one can add corrections to the effective potential around that scale order by order in perturbation theory using couplings at that scale. This essentially amounts to dropping the field-strength

renormalization contribution to the resummed effective potential. However, we are not performing any field redefinitions. Thus the same threshold corrections can be used to convert observables to  $\overline{\text{MS}}$  parameters at the low scale, and then run to the scale near the quantum minimum. The effective potential computed this way is still gauge dependent. Moreover, even in Landau gauge (or any other gauge), the potential is different if we first run the couplings and then calculate the potential or if we calculate the potential and then run it by solving its RGE. The difference is not an artifact of the truncation in perturbation theory: the exact potentials computed in these two ways would differ. However, this difference, like the difference between potentials computed in different gauges, should not affect physical observables. We have shown that indeed it does not affect the value of the potential at the minimum.

Our work definitively resolves the question of how one physical quantity, the true vacuum energy in a theory, can be computed consistently in perturbation theory. We also know that  $S$ -matrix elements are gauge invariant. There are of course many other quantities that effective actions are used to calculate, and it would be very interesting to understand definitively how they can be computed consistently as well. To compute the scalar and vector masses, one would need to resum the daisy contributions to the scalar and vector self-energy graphs at small momentum, supplementing the results of Ref. [31]. Part of this resummation was performed in Ref. [40] in Landau and unitarity gauges. It would be interesting to complete the calculation, and also to understand why Kang's contributions were gauge invariant on their own. Another interesting quantity is the tunneling rate between two extrema. A traditional tunneling-rate calculation depends on unphysical regions of the potential [51,52], away from the minima. However, any rate calculation should also depend on the kinetic terms in the effective action, so there are new opportunities for cancellation. Although there has been some progress in understanding how tunneling rates may be consistent [23,34], there are, to our knowledge, no explicit demonstrations.

At finite temperature, the critical temperature for a phase transition should be physical. The spectrum of gravity waves as the Universe cools through this transition should

also be physical. While some progress has been made in showing that these quantities can be computed in a gauge-invariant way in perturbation theory [14,21,46,53] the question does not seem to be completely resolved. Other apparently gauge-dependent quantities include the compactification radius in certain extra-dimensional models [37,54] and the inflation scale in Higgs inflation models [55].

Often the gauge dependence in a quantity extracted from the effective potential originates in a dependence on field values. For example, the instability scale in the Standard Model  $\Lambda_I$ , defined as the value of  $\phi$  where the potential goes negative [49], is gauge dependent [56]. The scale where the Standard Model effective potential is maximal,  $\Lambda_{\text{max}}$ , is also sometimes used to make physical arguments [57], despite its gauge dependence. A common criterion for stability in the Standard Model is that the value  $v$  of the field at the minimum is less than  $M_{\text{Pl}}$ . Since  $v$  is gauge dependent, so is this criterion [58]. Even if one asks only that the potential at the minimum,  $V_{\text{min}}$  be less than 0, which is a gauge-invariant criterion, one would presumably still want to know when this bound is affected by Planck-suppressed or other higher-dimension operators. To this end, we have sketched one possible solution in Sec. IX: one should add these new operators to the theory and see how  $V_{\text{min}}$  changes. Since the new operators affect the perturbation expansion, this procedure is not equivalent to replacing  $\phi \rightarrow v$  in these operators and seeing how the vacuum energy changes. With these examples, we are optimistic that the apparent gauge invariance of many physical quantities can be eliminated by the self-consistent use of an appropriate perturbation expansion.

## ACKNOWLEDGMENTS

We thank M. Reece, M. Ramsey-Musolf, M. Strassler and A. Strumia for helpful discussions. The authors are supported in part by Grant No. DE-SC003916 from the Department of Energy. A. A. is supported in part by the Stolt-Nielsen Fund for Education of the American-Scandinavian Foundation and the Norway-America Association. W.F. is supported in part by the Lord Rutherford Memorial Research Fellowship.

- 
- [1] W. Heisenberg and H. Euler, Consequences of dirac's theory of positrons, *Z. Phys.* **98**, 714 (1936).
  - [2] J. S. Schwinger, On gauge invariance and vacuum polarization, *Phys. Rev.* **82**, 664 (1951).
  - [3] G. Jona-Lasinio, Relativistic field theories with symmetry breaking solutions, *Nuovo Cimento* **34**, 1790 (1964).

- [4] S. R. Coleman and E. J. Weinberg, Radiative corrections as the origin of spontaneous symmetry breaking, *Phys. Rev. D* **7**, 1888 (1973).
- [5] R. Jackiw, Functional evaluation of the effective potential, *Phys. Rev. D* **9**, 1686 (1974).
- [6] L. Dolan and R. Jackiw, Gauge invariant signal for gauge symmetry breaking, *Phys. Rev. D* **9**, 2904 (1974).

- [7] S. Weinberg, Perturbative calculations of symmetry breaking, *Phys. Rev. D* **7**, 2887 (1973).
- [8] W. Fischler and R. Brout, Gauge invariance in spontaneously broken symmetry, *Phys. Rev. D* **11**, 905 (1975).
- [9] J.-M. Frere and P. Nicoletopoulos, Gauge invariant content of the effective potential, *Phys. Rev. D* **11**, 2332 (1975).
- [10] S.H. Tye and Y. Vtorov-Karevsky, Effective action of spontaneously broken gauge theories, *Int. J. Mod. Phys. A* **13**, 95 (1998).
- [11] W. Buchmuller, Z. Fodor, and A. Hebecker, Gauge invariant treatment of the electroweak phase transition, *Phys. Lett. B* **331**, 131 (1994).
- [12] D. Binosi, J. Papavassiliou, and A. Pilaftsis, Displacement operator formalism for renormalization and gauge dependence to all orders, *Phys. Rev. D* **71**, 085007 (2005).
- [13] L.P. Alexander and A. Pilaftsis, The one-loop effective potential in non-linear gauges, *J. Phys. G* **36**, 045006 (2009).
- [14] D. Boyanovsky, D. Brahm, R. Holman, and D. Lee, The gauge invariant effective potential: Equilibrium and non-equilibrium aspects, *Phys. Rev. D* **54**, 1763 (1996).
- [15] G.-L. Lin and T.-K. Chyi, Gauge independent effective potential and the Higgs mass bound, *Phys. Rev. D* **60**, 016002 (1999).
- [16] N. Nielsen, Removing the gauge parameter dependence of the effective potential by a field redefinition, *Phys. Rev. D* **90**, 036008 (2014).
- [17] N. Nielsen, On the gauge dependence of spontaneous symmetry breaking in gauge theories, *Nucl. Phys.* **B101**, 173 (1975).
- [18] R. Fukuda and T. Kugo, Gauge invariance in the effective action and potential, *Phys. Rev. D* **13**, 3469 (1976).
- [19] W. Loinaz and R. Willey, Gauge dependence of lower bounds on the Higgs mass derived from electroweak vacuum stability constraints, *Phys. Rev. D* **56**, 7416 (1997).
- [20] M. Gonderinger, H. Lim, and M. J. Ramsey-Musolf, Complex scalar singlet dark matter: Vacuum stability and phenomenology, *Phys. Rev. D* **86**, 043511 (2012).
- [21] H. H. Patel and M. J. Ramsey-Musolf, Baryon washout, electroweak phase transition, and perturbation theory, *J. High Energy Phys.* **07** (2011) 029.
- [22] E. J. Weinberg and A.-q. Wu, Understanding complex perturbative effective potentials, *Phys. Rev. D* **36**, 2474 (1987).
- [23] A. Strumia and N. Tetradis, A consistent calculation of bubble nucleation rates, *Nucl. Phys.* **B542**, 719 (1999).
- [24] L. Abbott, Introduction to the background field method, *Acta Phys. Pol. B* **13**, 33 (1982).
- [25] M. D. Schwartz, *Quantum Field Theory and the Standard Model* (Cambridge University Press, Cambridge, England, 2014).
- [26] I. Aitchison and C. Fraser, Gauge invariance and the effective potential, *Ann. Phys. (N.Y.)* **156**, 1 (1984).
- [27] J. Do Nascimento and D. Bazeia, Gauge invariance of the effective potential, *Phys. Rev. D* **35**, 2490 (1987).
- [28] K. Fujikawa, B. W. Lee, and A. I. Sanda, Generalized renormalizable gauge formulation of spontaneously broken gauge theories, *Phys. Rev. D* **6**, 2923 (1972).
- [29] C. Ford, I. Jack, and D. Jones, The standard model effective potential at two loops, *Nucl. Phys.* **B387**, 373 (1992).
- [30] S. P. Martin, Two loop effective potential for a general renormalizable theory and softly broken supersymmetry, *Phys. Rev. D* **65**, 116003 (2002).
- [31] J. Kang, Gauge invariance of the scalar-vector mass ratio in the Coleman-Weinberg model, *Phys. Rev. D* **10**, 3455 (1974).
- [32] C. Fraser, Calculation of higher derivative terms in the one loop effective lagrangian, *Z. Phys. C* **28**, 101 (1985).
- [33] A. I. Davydychev and J. Tausk, Two loop self-energy diagrams with different masses and the momentum expansion, *Nucl. Phys.* **B397**, 123 (1993).
- [34] D. Metaxas and E. J. Weinberg, Gauge independence of the bubble nucleation rate in theories with radiative symmetry breaking, *Phys. Rev. D* **53**, 836 (1996).
- [35] M. E. Machacek and M. T. Vaughn, Two loop renormalization group equations in a general quantum field theory: 1. wave function renormalization, *Nucl. Phys.* **B222**, 83 (1983).
- [36] M. E. Machacek and M. T. Vaughn, Two loop renormalization group equations in a general quantum field theory: 3. scalar quartic couplings, *Nucl. Phys.* **B249**, 70 (1985).
- [37] N. Nielsen, Gauge dependence of effective potential and self-consistent dimensional reduction, *Z. Phys. C* **33**, 579 (1987).
- [38] D. Johnston, Coleman-Weinberg, Nielsen and Daisies, *Phys. Lett. B* **186**, 185 (1987).
- [39] D. Bazeia, Gauge invariance and daisies, *Phys. Lett. B* **207**, 53 (1988).
- [40] A. Larsen and N. Nielsen, Vector tadpole graphs in the Coleman-Weinberg model, *Z. Phys. C* **44**, 689 (1989).
- [41] D. Kirzhnits and A. D. Linde, Macroscopic consequences of the Weinberg model, *Phys. Lett.* **42B**, 471 (1972).
- [42] S. Weinberg, Gauge and global symmetries at high temperature, *Phys. Rev. D* **9**, 3357 (1974).
- [43] M. Carrington, The effective potential at finite temperature in the standard model, *Phys. Rev. D* **45**, 2933 (1992).
- [44] S. P. Martin, Taming the goldstone contributions to the effective potential, *Phys. Rev. D* **90**, 016013 (2014).
- [45] J. Elias-Miro, J. Espinosa, and T. Konstandin, Taming infrared divergences in the effective potential, *J. High Energy Phys.* **08** (2014) 034.
- [46] C. Wainwright, S. Profumo, and M. J. Ramsey-Musolf, Gravity waves from a cosmological phase transition: Gauge artifacts and daisy resummations, *Phys. Rev. D* **84**, 023521 (2011).
- [47] M. Garny and T. Konstandin, On the gauge dependence of vacuum transitions at finite temperature, *J. High Energy Phys.* **07** (2012) 189.
- [48] M. Quiros, On daisy and superdaisy resummation of the effective potential at finite temperature, [arXiv:hep-ph/9304284](https://arxiv.org/abs/hep-ph/9304284).
- [49] D. Buttazzo, G. Degrassi, P. Paolo Giardino, G. F. Giudice, F. Sala, A. Salvio, and A. Strumia, Investigating the near-criticality of the Higgs boson, *J. High Energy Phys.* **12** (2013) 089.
- [50] A. Andreasen, W. Frost, and M. D. Schwartz, Consistent Use of the Standard Model Effective Potential, *Phys. Rev. Lett.* **113**, 241801 (2014).

- [51] J. Espinosa and M. Quiros, Improved metastability bounds on the standard model Higgs mass, *Phys. Lett. B* **353**, 257 (1995).
- [52] G. Isidori, G. Ridolfi, and A. Strumia, On the metastability of the standard model vacuum, *Nucl. Phys.* **B609**, 387 (2001).
- [53] D. E. Morrissey and M. J. Ramsey-Musolf, Electroweak baryogenesis, *New J. Phys.* **14**, 125003 (2012).
- [54] G. Kunstatter and H. Leivo, On the gauge dependence of the one loop effective potential in self-consistent dimensional reduction, *Phys. Lett.* **166B**, 321 (1986).
- [55] J. L. Cook, L. M. Krauss, A. J. Long, and S. Sabharwal, Is Higgs inflation ruled out?, *Phys. Rev. D* **89**, 103525 (2014).
- [56] L. Di Luzio and L. Mihaila, On the gauge dependence of the standard model vacuum instability scale, *J. High Energy Phys.* **06** (2014) 079.
- [57] A. Hook, J. Kearney, B. Shakya, and K. M. Zurek, Probable or improbable universe? Correlating electroweak vacuum instability with the scale of inflation, [arXiv:1404.5953](https://arxiv.org/abs/1404.5953).
- [58] W. Loinaz and R. Willey, Monte Carlo simulation calculation of critical coupling constant for two-dimensional continuum  $\varphi^4$  theory, *Phys. Rev. D* **58**, 076003 (1998).