

Baryon spin-flavor structure from an analysis of lattice QCD results of the baryon spectrum

I. P. Fernando^{*} and J. L. Goity[†]

Department of Physics, Hampton University, Hampton, Virginia 23668, USA
Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA
 (Received 6 October 2014; published 19 February 2015)

The excited baryon masses are analyzed in the framework of the $1/N_c$ expansion using the available physical masses and also the masses obtained in lattice QCD for different quark masses. The baryon states are organized into irreducible representations of $SU(6) \times O(3)$, where the $[56, \ell^P = 0^+]$ ground-state and excited baryons, and the $[56, 2^+]$ and $[70, 1^-]$ excited states are analyzed. The analyses are carried out to $\mathcal{O}(1/N_c)$ and first order in the quark masses. The issue of state identifications is discussed. Numerous parameter-independent mass relations result at those orders, among them the well known Gell-Mann-Okubo and equal-spacing relations, as well as additional relations involving baryons with different spins. It is observed that such relations are satisfied at the expected level of precision. From the quark-mass dependence of the coefficients in the baryon mass formulas an increasingly simpler picture of the spin-flavor composition of the baryons is observed with increasing pion mass (equivalently, increasing $m_{u,d}$ masses), as measured by the number of significant mass operators.

DOI: [10.1103/PhysRevD.91.036005](https://doi.org/10.1103/PhysRevD.91.036005)

PACS numbers: 11.15.Pg, 12.38.Gc, 14.20.Dh, 14.20.Gk

I. INTRODUCTION

One of the most important present objectives in lattice QCD (LQCD) is the calculation of the light baryon spectrum, where in recent years substantive progress has been made. The implementation of optimized baryon source operators [1–4] has enabled improved signals for excited baryons, leading to remarkable progress in identifying states by their quantum numbers and in the determination of their masses. In calculations performed with quark masses corresponding to $390 \text{ MeV} \leq M_\pi \leq 702 \text{ MeV}$, the spectrum of nonstrange baryons [3] and also of strange baryons [4] were obtained. The case of $M_\pi = 702 \text{ MeV}$ corresponds to the limit of exact $SU(3)$ flavor symmetry. These calculations were performed on anisotropic lattices $16^3 \times 128$ with a gluon Symanzik-improved action with tree-level tadpole-improved coefficients and an anisotropic clover fermion action as explained in Ref. [5]. Although the effects of finite widths of the baryons are not yet implemented in these calculations, the results are very significant. The extraction of the baryonic resonance parameters (mass and width) by means of finite-volume effects on the two-body spectrum (e.g., πN) as it has been carried out for the ρ meson [6], in baryons is still to be fully implemented in a LQCD calculation. A nice example of the latter was shown in a continuum chiral perturbation theory study of those effects for extracting the Δ resonance [7]. The results used in this work pertain to the use of quasilocal baryon source/sink operators, which are not entirely sufficient for extracting the resonance parameters, and therefore

the quoted masses will probably be (slightly) shifted in the more complete framework employing the finite-volume effects. In fact, for the LQCD states to be analyzed here, the available phase space for the two-body decay of the excited baryons is increasingly suppressed with increasing M_π , which for the considered range of quark masses result in state widths which are significantly smaller than in the physical case. An estimate using the available phase space and the phenomenological widths gives widths $\sim 50 \text{ MeV}$ or smaller for the S -wave decays and even smaller for P - and D -waves. Thus, the present results of the LQCD baryon masses are expected to be very close to those one would obtain with the more complete method.

Although other recent works on baryon LQCD spectroscopy have been carried out in Refs. [1,2,8,9–11], the present work will use the results obtained by the Jefferson Lab Lattice QCD Collaboration in Refs. [3,4]. The study can similarly be applied to other results, in particular those of the BGR Collaboration [8] where the masses of the states analyzed here have been calculated.

A key observation from the analysis carried out in Refs. [3,4] is that source/sink operators which, in the continuum limit, are in irreducible representations of the spin-flavor and quark orbital angular momentum groups $SU(2N_f) \times O(3)$ are very close to being at the optimum for the selective overlap with the baryon states. This is a strong indication that the baryon mass eigenstates themselves must be approximately organized into multiplets of that group, a fact that is well known to hold phenomenologically. This has been tested explicitly in the LQCD calculations by measuring the coupling strengths of sources in different representations to each of the baryon levels studied. The state admixture of different $SU(2N_f) \times O(3)$

^{*}ishara@jlab.org
[†]goity@jlab.org

irreducible representations, known as *configuration mixing*, cannot however be directly inferred from those strengths, as they depend on details of the operators. Since in the exact symmetry limit the couplings would be “diagonal,” it is expected that the existence of small off-diagonal couplings necessarily translates into small state-configuration mixings. In the present work, configuration mixings will be altogether neglected, and thus all claims are restricted to the approximate validity of that assumption. The states studied in this work are the ones corresponding to the $SU(6) \times O(3)^P$ $[56, 0^+]$ or Roper multiplet, the $[56, 2^+]$ and the $[70, 1^-]$. These are of particular interest because they have been previously analyzed phenomenologically in the framework of the $1/N_c$ expansion employed here [12], where the assumption of no configuration mixing works very well up to the degree of accuracy that the input masses and other observables permit.

The existence of a spin-flavor symmetry in baryons can be rigorously justified in the large- N_c limit of QCD. The symmetry is the result of a consistency requirement imposed by unitarity on pion-baryon scattering in that limit [13–15], and spin-flavor symmetry is thus broken by corrections which can be organized in powers of $1/N_c$. Under the assumption that the real world with $N_c = 3$ baryons can be analyzed using a $1/N_c$ expansion, starting at the lowest order with an exact spin-flavor symmetry, many analyses of baryon masses and other properties have been carried out. In particular, excited baryon masses have been analyzed in numerous works for the cases considered in this work [16–24] as well as for other multiplets [25–28]. Although spin-flavor symmetry is justified in the large- N_c limit, the larger $SU(2N_f) \times O(3)$ is not. The latter can be broken due to spin-orbit effects at $\mathcal{O}(N_c^0)$, as is the case in the $[70, 1^-]$ baryons, and it is in principle not such a good symmetry. However, phenomenologically it has been known for a long time that spin-orbit effects in baryons are small, and actually smaller than the hyperfine (HF) effects which are subleading in $1/N_c$. In addition, configuration mixings which are not suppressed in the large- N_c limit turn out to be driven by operators of the spin-orbit type [29,30], and seem to be small as well. As mentioned earlier, these observations also apply to the LQCD baryons.

Particular predictions result when configuration mixings are disregarded. They have the form of parameter-independent mass relations which hold up to higher-order corrections in the $1/N_c$ or $SU(3)$ -breaking expansions. Among those relations are the well known Gell-Mann-Okubo (GMO) and equal-spacing (EQS) relations, which are valid in general, and additional ones involving different spin-flavor states such as relations in the 56 -plets that follow from the Gürsey-Radicati mass formula, and other relations in the 70 -plet [21]. As it will be shown in the present analysis, LQCD baryon masses fulfill to the expected accuracy those relations.

The objective of this work is to analyze the LQCD results for baryon masses using the $1/N_c$ expansion to $\mathcal{O}(1/N_c)$ and to first order in $SU(3)$ symmetry breaking. Although the LQCD results, as mentioned above, are at larger than physical quark masses and do not have a complete implementation of the effects due to the finite baryon decay widths, they provide complete sets of states, i.e., states that complete the experimentally partially filled multiplets, which is a very useful addition for more accurate analyses as the ones carried out here. In addition, since the $1/N_c$ expansion of QCD applies even in cases where such approximations are made (e.g., quenched QCD, larger quark masses, etc.), the present study also serves as a test of the $1/N_c$ expansion itself.

In the phenomenological analyses, the excited baryon masses used as inputs are those provided by the Particle Data Group (PDG) [31]. For two flavors and the multiplets considered here all states are established, but for three flavors there is a significant number of missing strange baryon states. For example, in the $[70, 1^-]$ multiplet there are 30 theoretical masses and only 17 are currently experimentally known. Although those 17 masses are sufficient for the purpose of the $1/N_c$ analysis, they are not sufficient for a thorough test of the mass relations. On the other hand, the LQCD results provide complete multiplets, enabling a complete test of mass relations. In the particular case of the $[70, 1^-]$, the issue of state mixing can be sorted out in the phenomenological case thanks to the simultaneous analysis of partial decay widths and photocouplings, as shown most recently in Ref. [24] for the nonstrange baryons. These inputs are however not possible for the LQCD baryons, and therefore the state mixing relies very strongly on the criterion for identifying states. In this regard, level-crossing effects are possible as the quark masses are varied in the LQCD calculations [8,10]. This is a present topic of interest in LQCD, which is still in its early stages.

This work is organized as follows. In Sec. II a brief description of the $1/N_c$ expansion framework is given. Section III contains the results and their analysis. Section IV gives the summary and conclusions. Appendix A displays the bases of operators and the respective matrix elements needed in this work, and Appendix B gives the baryon masses, both from the PDG [31] and LQCD [3,4], which are the inputs for the fits.

II. THE $1/N_c$ EXPANSION AND SPIN-FLAVOR SYMMETRY IN BARYONS

Consistency of baryons in the ordinary large- N_c limit as defined by ’t Hooft [32] requires that baryons form multiplets of a contracted spin-flavor group $SU(2N_f)$ [13–15]. The generators of that symmetry are denoted by S^i (spin), T^a (flavor) and $X^{ia} = G^{ia}/N_c$ (spin-flavor). In the case of excited baryons, the observation that quark spin and orbital angular momentum are in general weakly

coupled in baryons has lead to a phenomenologically successful scheme of organizing the states in multiplets of $SU(2N_f) \times O(3)$. For finite N_c it is possible to work with the ordinary rather than the contracted spin-flavor group for the purposes of building the operator bases [33]. Any static baryonic observable can be expressed by an effective operator which is decomposed in a basis of operators ordered in powers of $1/N_c$ and which can be expressed as appropriate tensor products of the symmetry generators. In the present case of baryon masses, the bases of operators are well known. The details for obtaining those bases can be found in Refs. [16,18–21,34,35].

The excited states considered here will be either in the totally symmetric or in the mixed symmetric irreducible representations of $SU(6)$. Following the large- N_c Hartree picture of a baryon, without a loss of generality and for the purpose of dealing with the group theory of the spin-flavor and orbital degrees of freedom, one can describe a low-excitation baryon as a spin-flavor symmetric core with $N_c - 1$ quarks and one excited quark. In this way it becomes straightforward to obtain the matrix elements of bases operators. Appendix A gives the mass operator bases to the needed order and the corresponding matrix elements.

The mass operator bases are organized in powers of $1/N_c$ and involve $SU(3)$ singlet and octet operators, the latter for symmetry breaking by the parameter $\epsilon \equiv m_s - \hat{m}$, where $\hat{m} = (m_u + m_d)/2$. One may consider the expansion to $\mathcal{O}(\epsilon^0/N_c)$ and $\mathcal{O}(\epsilon)$. It turns out that contributions $\mathcal{O}(\epsilon/N_c)$ are almost insignificant in most cases as shown later.

The multiplets to be analyzed have the following state contents: (i) $[56, 0^+]$, one $SU(3)$ **8** with $S = 1/2$ and one **10** with $S = 3/2$; (ii) $[56, 2^+]$, one **8** for each $S = 3/2$ and $5/2$, and one **10** for each $S = 1/2$ through $7/2$; (iii) $[70, 1^-]$, one **1** Λ baryon for each $S = 1/2$ and $3/2$, two **8**'s for each $S = 1/2$ and $3/2$, one **8** for $S = 5/2$ and one **10** for each $S = 1/2$ and $3/2$.

For each case, the mass operators to the order needed here are as follows,

$[56, 0^+]$: In this case the mass operator is the famous Gürsey-Radicati (GR) mass formula, which, explicitly displaying the $1/N_c$ power counting, reads as

$$M_{[56,0^+]} = c_1 N_c + \frac{c_2}{N_c} S(S+1) + b_1 N_s + \frac{b_2}{2\sqrt{12}N_c} \left(3I(I+1) - S(S+1) - \frac{3}{4} N_s(N_s+2) \right) + \mathcal{O}(1/N_c^2), \quad (1)$$

where S is the baryon spin operator, I is the isospin, and N_s is the number of strange quarks, and the c_i and b_i are coefficients determined by the QCD dynamics, which are obtained by fitting to the masses. The mass operators are defined such that all coefficients are $\mathcal{O}(N_c^0)$. The $SU(3)$ -breaking parameter ϵ is here included in the coefficients b_i

and b_2 . For all mass formulas, the quark-mass dependence is implicitly absorbed into the coefficients.

$[56, 2^+]$: In this case the basis has three $SU(3)$ -symmetric and three breaking operators,

$$M_{[56,2^+]} = \sum_{i=1}^3 c_i O_i + \sum_{i=1}^3 b_i \bar{B}_i. \quad (2)$$

The basis of operators along with the matrix elements are given in Appendix A 1, Table XVI.

$[70, 1^-]$: In the case of nonstrange baryons, where the states belong to the mixed symmetric $20'$ -plet of $SU(4)$, the mass formula reads [19]

$$M_{[20',1^-]} = \sum_{i=1}^8 c_i O_i, \quad (3)$$

where the eight basis operators up to and including $\mathcal{O}(1/N_c)$ are given in Table XVII of Appendix A 2. For three flavors the mass formula reads [20,21]

$$M_{[70,1^-]} = \sum_{i=1}^{11} c_i O_i + \sum_{i=1}^4 b_i \bar{B}_i, \quad (4)$$

where the basis operators up to and including $\mathcal{O}(1/N_c)$ or $\mathcal{O}(\epsilon)$ are given in Table XVIII and Table XIX of Appendix A 3. In order that the $SU(3)$ -breaking operators do not contribute to the nonstrange baryon masses, they have been redefined according to $\bar{B}_1 = t_8 - \frac{1}{2\sqrt{3}N_c} O_1$, $\bar{B}_2 = T_8 - \frac{N_c-1}{2\sqrt{3}N_c} O_1$, $\bar{B}_3 = \frac{10}{N_c} d_{8ab} g_{ia} G_{ib} + \frac{5(N_c^2-9)}{8\sqrt{3}N_c^2(N_c-1)} O_1 + \frac{5}{2\sqrt{3}(N_c-1)} O_6 + \frac{5}{6\sqrt{3}} O_7$, $\bar{B}_4 = 3\ell_i g_{i8} - \frac{\sqrt{3}}{2} O_2$.

Since in general the number of states is larger than the number of coefficients of the fit, and the masses are linear in the coefficients, there must be linear mass relations which are independent of the coefficients. Such mass relations have been derived in previous works, and will be tested here with the LQCD results. Many of the mass relations involve $SU(3)$ -breaking mass differences, and are thus identically satisfied in the limit of $SU(3)$ symmetry. There are however some mass relations which test exclusively the breaking of the spin symmetry at $\mathcal{O}(1/N_c)$; these occur in the $[56, 2^+]$ multiplet. The mass relations will be presented in the analysis of each case below, and they are depicted in Tables III, IV, VII, XIII, and XIV.

III. FITS TO THE LQCD RESULTS

In this section the fits to the LQCD masses are performed. The LQCD results used here are as follows: for two flavors the results are those of Ref. [3], of which only the results for the negative-parity baryon masses will be analyzed, and for three flavors the results of Ref. [4] are used. For two flavors the quark masses used correspond to

TABLE I. Coefficients of the GR mass formula for the ground-state baryons. The case $M_\pi = 702$ MeV corresponds to exact SU(3) symmetry. χ^2_{dof} is the χ^2 per degree of freedom.

Coefficients [MeV]	M_π [MeV]			
	PDG	391	524	702
c_1	293 ± 6	377 ± 3	420 ± 2	474 ± 1
c_2	247 ± 12	296 ± 5	251 ± 3	200 ± 2
b_1	189 ± 12	75 ± 6	45 ± 3	0
b_2	94 ± 26	43 ± 11	14 ± 7	0
χ^2_{dof}	0.19	0.15	1.43	0

TABLE II. Fit to the $[56, 0^+]$ Roper baryons. It is found that the SU(3)-breaking effects on the HF interactions can be neglected, and thus $b_2 = 0$ throughout.

Coefficients [MeV]	M_π [MeV]			
	PDG	391	524	702
c_1	469 ± 8	714 ± 6	760 ± 5	770 ± 3
c_2	175 ± 44	165 ± 12	124 ± 9	115 ± 20
b_1	204 ± 18	48 ± 12	15 ± 12	0
χ^2_{dof}	0.16	0.53	0.76	0

$M_\pi = 396$ and 524 MeV, and for three flavors m_s has been kept fixed, and $M_\pi = 391, 524$ and 702 MeV. For each of the multiplets it is necessary to identify the states with the LQCD mass levels. This procedure is not unique and thus it requires some analysis, as shown below. In the following the notation used to designate the states will be as follows: B_S or B'_S for states with baryon spin S which belong predominantly to octets, and B''_S for baryons which belong predominantly in singlet or decuplet. For the case of the Δ and Ω baryons which can only belong in a decuplet when isospin is exact, no primes are used, and the same for the $[56, 0^+]$ baryons where $S = 1/2(3/2)$ states necessarily belong to **8** (**10**).

A. $[56, 0^+]$ baryons

Here, the ground-state and excited $[56, 0^+]$ (Roper multiplet) baryon masses are fitted using the GR mass formula Eq. (1). The input masses used here are depicted in Appendix B, Table XX. In all **56**-plet masses the flavor singlet breaking of $SU(6) \times O(3)$ is $\mathcal{O}(1/N_c)$, i.e., suppressed by a factor $1/N_c^2$ with respect to the leading symmetric mass. Thus, under the assumption of no configuration mixing, $SU(6) \times O(3)$ must be particularly good. The possible significance of the SU(3)-breaking effects on the HF terms, controlled by the coefficient b_2 , is considered. Table I gives the results of the fits for the ground-state baryons, and Table II for the Roper baryons.

The analysis of LQCD ground-state baryon masses including higher-order terms in the SU(3) breaking has been carried out in Ref. [36], for LQCD calculations other than the present ones. It is noted that the HF mass splittings have the behavior observed also in other LQCD calculations, where it increases with M_π up to $M_\pi \sim 400$ MeV, and decreases for higher M_π (for a current summary see Ref. [37]). On the other hand for the excited baryons the HF splittings are almost always monotonously decreasing with increasing M_π , both in the **56**- and **70**-plets.

In the Roper baryons, the identification of the $\mathbf{8}_{1/2}$ is obvious, being the lightest positive-parity excited states above the ground state, but for the $\mathbf{10}_{3/2}$ one needs to distinguish between two possible excited multiplets: one will be a Roper and the other will be in the $[56, 2^+]$. One of the choices, namely that in which the Roper $\mathbf{10}_{3/2}$ is the lightest one, does not seem to be consistent with the magnitude of the HF splittings observed throughout the spectrum. One is therefore lead to conclude that the $\mathbf{10}_{3/2}$ belonging to the $[56, 2^+]$ are the lowest-lying excitations, followed by the Roper ones.

In Fig. 1 the dependencies on M_π of the coefficients are displayed. The well-known dramatic downturn with decreasing M_π of the Roper baryon masses is clearly driven by the spin-flavor singlet component of the masses, given by the coefficient c_1 . The HF effects determined by c_2 have a smooth behavior in M_π but significantly different strength in the GS and in the Roper states, being reduced in




FIG. 1 (color online). Evolution of the coefficients with M_π for the ground-state baryons (left panel) and the Roper baryons (right panel).

TABLE III. Mass relations for the ground-state octet and decuplet. The relations are valid up to corrections $\mathcal{O}(e^{\frac{3d}{2}}/N_c)$ in the case of the GMO and EQS relations which stem from the one-loop chiral corrections [36], and up to $\mathcal{O}(1/N_c^2)$ for the rest of the relations.

Relation	M_π [MeV]		
	PDG	391	524
$2(N + \Xi) - (3\Lambda + \Sigma) = 0$	30.2 ± 0.4	38 ± 75	32 ± 32
$\Sigma'' - \Delta = \Xi'' - \Sigma'' = \Omega'' - \Xi''$	155 ± 2	64 ± 25	40 ± 11
	149.0 ± 0.5	55 ± 19	33 ± 13
	140.7 ± 0.5	54 ± 17	40 ± 14
$\frac{1}{3}(\Sigma + 2\Sigma'') - \Lambda - \frac{2}{3}(\Delta - N) = 0$	9 ± 1	1 ± 28	14 ± 12
$\Sigma'' - \Sigma - (\Xi'' - \Xi) = 0$	23.5 ± 0.5	12 ± 25	12 ± 15
$3\Lambda + \Sigma - 2(N + \Xi) + (\Omega - \Xi'' - \Sigma'' + \Delta) = 0$	16 ± 2	29 ± 81	32 ± 36
$\Sigma'' - \Delta + \Omega - \Xi'' - 2(\Xi'' - \Sigma'') = 0$	2.5 ± 2.4	8 ± 51	14 ± 37

TABLE IV. Mass relations for the Roper multiplet. The relations hold at the same orders as in the case of the ground-state baryons.

Relation	M_π [MeV]	
	391	524
$2(N + \Xi) - (3\Lambda + \Sigma) = 0$	179 ± 180	106 ± 155
$\Sigma'' - \Delta = \Xi'' - \Sigma'' = \Omega'' - \Xi''$	13 ± 45	-27 ± 26
	84 ± 40	41 ± 49
	48 ± 42	41 ± 57
$\frac{1}{3}(\Sigma + 2\Sigma'') - \Lambda - \frac{2}{3}(\Delta - N) = 0$	51 ± 65	29 ± 41
$\Sigma'' - \Sigma - (\Xi'' - \Xi) = 0$	58 ± 63	77 ± 80
$3\Lambda + \Sigma - 2(N + \Xi) + (\Omega'' - \Xi'' - \Sigma'' + \Delta) = 0$	144 ± 189	174 ± 170
$\Sigma'' - \Delta + \Omega'' - \Xi'' - 2(\Xi'' - \Sigma'') = 0$	107 ± 110	67 ± 147

the latter. Unlike the GS baryons, no significant SU(3) breaking in the HF interaction is observed in the Roper baryons, so the coefficient b_2 is consistent with zero for the LQCD masses.

The mass relations are given in Tables III and IV, which show that they are satisfied within errors for the LQCD results. In the physical case, the knowledge of the Roper states is rather incomplete. Based on the mass relations the predictions shown in Table V are made. As shown below, the listed PDG candidate states may also match predictions from the $[56, 2^+]$ multiplet, as discussed later.

B. $[56, 2^+]$ baryons

Here the lowest excited baryons that can fit into the $[56, 2^+]$ are considered. The input masses used here are depicted in Appendix B, Table XXI. The first step is the

identification of the states in the LQCD results. With the exception of the $\mathbf{10}_{3/2}$, all the states are in spin-flavor states which appear for the first time, and thus the lightest states with given spin and flavor are the ones of interest. In the case of the $\mathbf{10}_{3/2}$, as discussed earlier, there are two possible excited levels to consider, one of which will belong to the excited $[56, 0^+]$, where the arguments for the identification were already given. For Σ and Ξ baryons, the LQCD analysis [4] has assigned the dominant SU(3) multiplet to which they belong, $\mathbf{8}$ or $\mathbf{10}$. Therefore, there is no ambiguity about the identification of states in the present multiplet.

There is mixing between states in the octet and decuplet, namely the Σ and the Ξ pairs of states with $S = 3/2$ and with $S = 5/2$, namely $(\Sigma_S^{(8)}, \Sigma_S^{(10)})$ and $(\Xi_S^{(8)}, \Xi_S^{(10)})$. These mixings obviously result from SU(3) breaking, and the physical states are defined as follows:

TABLE V. Predictions of physically unknown states in the Roper multiplet. These predictions agree with the ones in Ref. [38]. (*) fit with all basis operators.

Baryon	Predicted mass [MeV]	Fitted Mass (*) [MeV]	PDG candidate and mass [MeV]
$\Sigma''_{3/2}$	1790 ± 131	1800 ± 26	$\Sigma(1840)(3/2^+)^*$ with mass ~ 1840
$\Xi''_{1/2}$	1825 ± 108	1815 ± 31	...
$\Xi''_{3/2}$	1955 ± 171	1975 ± 35	$\Xi(1950)(?)^{***}$ with mass $\sim 1950 \pm 15$
$\Omega''_{3/2}$	2120 ± 219	2150 ± 46	...

TABLE VI. First four fits include all operators and the next three fits with only O_1 , O_2 , O_3 and \bar{B}_1 operators. The fit with the minimal set of operators does not describe well the physical baryons.

Coefficients [MeV]	M_π [MeV]						
	PDG	391	524	702	391	524	702
c_1	540 ± 11	704 ± 2	718 ± 1	754 ± 1	710 ± 2	724 ± 1	753 ± 1
c_2	18 ± 5	48 ± 6	28 ± 3	-6 ± 5	59 ± 6	21 ± 3	0
c_3	244 ± 4	169 ± 5	166 ± 3	104 ± 4	151 ± 5	148 ± 3	106 ± 4
b_1	217 ± 4	75 ± 3	54 ± 1	0	56 ± 3	36 ± 1	0
b_2	95 ± 14	-23 ± 11	13 ± 5	0	0	0	0
b_3	268 ± 16	59 ± 9	55 ± 4	0	0	0	0
Mixing angles [rad]							
$\theta_{\Sigma_{3/2}}$	-0.16 ± 0.02	0.06 ± 0.03	-0.03 ± 0.01	0	0	0	0
$\theta_{\Xi_{3/2}}$	-0.26 ± 0.04	0.07 ± 0.03	-0.03 ± 0.01	0	0	0	0
$\theta_{\Sigma_{5/2}}$	-0.22 ± 0.03	0.05 ± 0.03	-0.03 ± 0.01	0	0	0	0
$\theta_{\Xi_{5/2}}$	-0.20 ± 0.02	0.08 ± 0.04	-0.03 ± 0.01	0	0	0	0
χ^2_{dof}	0.84	0.60	0.47	0.92	0.63	0.53	0.80

$$\begin{pmatrix} \Sigma_S \\ \Sigma'_S \end{pmatrix} = \begin{pmatrix} \cos \theta_{\Sigma_S} & \sin \theta_{\Sigma_S} \\ -\sin \theta_{\Sigma_S} & \cos \theta_{\Sigma_S} \end{pmatrix} \begin{pmatrix} \Sigma_S^{(8)} \\ \Sigma_S^{(10)} \end{pmatrix},$$

$$\begin{pmatrix} \Xi_S \\ \Xi'_S \end{pmatrix} = \begin{pmatrix} \cos \theta_{\Xi_S} & \sin \theta_{\Xi_S} \\ -\sin \theta_{\Xi_S} & \cos \theta_{\Xi_S} \end{pmatrix} \begin{pmatrix} \Xi_S^{(8)} \\ \Xi_S^{(10)} \end{pmatrix}. \quad (5)$$

Two different fits are carried out: one includes all the SU(3)-breaking operators, and a second one only includes the one-body operator giving the spin-independent breaking effects. Since the symmetry breaking by the operator \bar{B}_1 does not produce mixing between **8** and **10**, the mixing angles are actually $\propto \epsilon/N_c$, and thus naturally very small. The results are shown in Table VI. It is checked that the present fit fully agrees with a previous one for the physical case [22]. One important observation is that based on the quality of the fits the mixings cannot be definitely established for the LQCD results.

The evolution with M_π of the coefficients is shown in Fig. 2. It is interesting to notice that the coefficient c_1 has qualitatively similar but less dramatic behavior than in the case of the Roper baryons, which must be an indication of a

similar mechanism as the one which drives down the Roper masses with decreasing M_π . The HF interaction given by c_3 behaves smoothly with M_π , decreasing slowly as M_π increases, and it has similar strength as in the Roper baryons. Although the operators \bar{B}_2 and \bar{B}_3 are significant in the physical case, their contributions are negligible in the LQCD cases, as shown by the second fit in Table VI. The latter observation implies that the mixing between octet and decuplet states, which are driven by those operators, are very small as confirmed by the small mixing angles in the first fit in Table VI.

The mass relations for the $[56, 2^+]$ are depicted in Table VII [35]. In addition to the GMO and EQS relations, there are several relations which relate SU(3) mass splittings in multiplets with different baryon spin, as well as relations among the masses of baryons with the same strangeness but different baryon spin. Almost all relations are satisfied by the LQCD results, with the exception of the results at $M_\pi = 702$ MeV, where the deviations are however within the expected magnitude of higher order corrections.

The predictions of the fit and the mass relations for the experimentally unknown or poorly known states are shown


FIG. 2 (color online). Evolution of the operator coefficients with M_π for the two fits in Table VI.

TABLE VII. Mass relations for the $[56, 2^+]$ multiplet. The relations hold at the same orders as in the case of the ground-state baryons. The last column corresponds to the SU(3) symmetric limit.

Relation	M_π [MeV]		
	391	524	702
$2(N_{3/2} + \Xi_{3/2}) - (3\Lambda_{3/2} + \Sigma_{3/2}) = 0$	98 ± 126	49 ± 173	0
$2(N_{5/2} + \Xi_{5/2}) - (3\Lambda_{5/2} + \Sigma_{5/2}) = 0$	40 ± 98	55 ± 65	0
$\Sigma''_{1/2} - \Delta_{1/2} = \Xi''_{1/2} - \Sigma''_{1/2} = \Omega_{1/2} - \Xi''_{1/2}$	-13 ± 110	36 ± 33	0
	23 ± 44	43 ± 22	0
	85 ± 54	35 ± 19	0
$\Sigma''_{3/2} - \Delta_{3/2} = \Xi''_{3/2} - \Sigma''_{3/2} = \Omega_{3/2} - \Xi''_{1/2}$	48 ± 46	36 ± 23	0
	56 ± 29	30 ± 16	0
	45 ± 31	41 ± 15	0
$\Sigma''_{5/2} - \Delta_{5/2} = \Xi''_{5/2} - \Sigma''_{5/2} = \Omega_{5/2} - \Xi''_{5/2}$	35 ± 40	34 ± 26	0
	62 ± 31	26 ± 23	0
	57 ± 34	52 ± 18	0
$\Sigma''_{7/2} - \Delta_{7/2} = \Xi''_{7/2} - \Sigma''_{7/2} = \Omega_{7/2} - \Xi''_{7/2}$	38 ± 38	35 ± 25	0
	67 ± 31	36 ± 20	0
	59 ± 31	22 ± 18	0
$\Delta_{5/2} - \Delta_{3/2} - (N_{5/2} - N_{3/2}) = 0$	70 ± 68	4 ± 68	44 ± 33
$(\Delta_{7/2} - \Delta_{5/2}) - \frac{7}{5}(N_{5/2} - N_{3/2}) = 0$	68 ± 78	2.5 ± 92	75 ± 41
$\Delta_{7/2} - \Delta_{1/2} - 3(N_{5/2} - N_{3/2}) = 0$	129 ± 175	13 ± 192	133 ± 74
$\frac{8}{15}(\Lambda_{3/2} - N_{3/2}) + \frac{22}{15}(\Lambda_{5/2} - N_{5/2}) - (\Sigma_{5/2} - \Lambda_{5/2}) - 2(\Sigma_{7/2}'' - \Delta_{7/2}) = 0$	91 ± 100	29 ± 75	0
$\Lambda_{5/2} - \Lambda_{3/2} + 3(\Sigma_{5/2} - \Sigma_{3/2}) - 4(N_{5/2} - N_{3/2}) = 0$	10 ± 207	10 ± 272	0
$\Lambda_{5/2} - \Lambda_{3/2} + \Sigma_{5/2} - \Sigma_{3/2} - 2(\Sigma_{5/2}'' - \Sigma_{3/2}'') = 0$	111 ± 81	12 ± 72	87 ± 59
$7(\Sigma''_{3/2} - \Sigma''_{7/2}) - 12(\Sigma''_{5/2} - \Sigma''_{7/2}) = 0$	44 ± 319	39 ± 268	67 ± 266
$4(\Sigma''_{1/2} - \Sigma''_{7/2}) - 5(\Sigma''_{3/2} - \Sigma''_{7/2}) = 0$	83 ± 170	87 ± 104	58 ± 161

in Table VIII. The PDG candidate state $\Sigma(1840)(3/2^+)^*$ could be identified with the $\Sigma_{3/2}(1889)$ state in Table VIII, but as discussed earlier it can also be identified with the Roper $\Sigma_{3/2}$. The PDG candidate state $\Xi(2120)^*(?)$ is consistent with both $\Xi_{3/2}$ and $\Xi''_{7/2}$ in Table VIII, so its parity could be predicted to be positive.

C. $[70, 1^-]$ baryons

In the case of two flavors, there are two mixing angles for the pairs of nucleon states with $S = 1/2$ and $S = 3/2$. Denoting by $^{(2s+1)}N_s$ the nucleon state with spin S and quark spin s , the physical states are given by

TABLE VIII. Predictions of physically unknown states in the $[56, 2^+]$ multiplet, and suggested identifications with PDG listed states. The first two GMO relations and the 12th equation in Table VII, which is a large- N_c parameter-independent mass relation, were used to predict the above masses.

Missing states	Fitted mass [MeV]	Mass listed in PDG [MeV]	Mass from mass relations [MeV]
$\Sigma_{3/2}$	1889 ± 30	$\Sigma(1840)(3/2^+)^*$ with mass ~ 1840	1920 ± 70
$\Xi_{3/2}$	2074 ± 24	$\Xi(2120)^*(?)$: 2130 ± 7	2080 ± 75
$\Xi_{5/2}$	2000 ± 31	$\Xi(2030)^{***}$ ($S \geq 5/2^+$) with 2025 ± 5	2006 ± 14
$\Sigma''_{1/2}$	2060 ± 6	...	2127 ± 120
$\Xi''_{1/2}$	2221 ± 6	$\Xi(2250)^{**}(?)$: 2214 ± 5	...
$\Omega_{1/2}$	2382 ± 7
$\Sigma''_{3/2}$	2059 ± 29	$\Sigma(2080)^{**}(3/2^+)$: 2120 ± 40	2109 ± 96
$\Xi''_{3/2}$	2212 ± 24
$\Omega_{3/2}$	2350 ± 6
$\Sigma''_{5/2}$	2053 ± 23	$\Sigma(2070)^*(5/2^+)$: 2070 ± 10	2077 ± 56
$\Xi''_{5/2}$	2178 ± 31
$\Omega_{5/2}$	2297 ± 6
$\Xi''_{7/2}$	2129 ± 6	$\Xi(2120)^*(?)$: 2130 ± 7	...
$\Omega_{7/2}$	2222 ± 6

$$\begin{pmatrix} N_S \\ N'_S \end{pmatrix} = \begin{pmatrix} \cos \theta_{2S} & \sin \theta_{2S} \\ -\sin \theta_{2S} & \cos \theta_{2S} \end{pmatrix} \begin{pmatrix} {}^2N_S \\ {}^4N_S \end{pmatrix} \quad (6)$$

Understanding these mixings is very important, as the decays and photo-couplings are sensitive to them. These mixings are predicted at the leading level of breaking of spin-flavor symmetry [23]. Indeed, if the $\mathcal{O}(N_c^0)$ spin-orbit operators $O_{2,3,4}$ would have contributions of natural size, the mixing angles would be $\theta_1 = \cos^{-1}(-\sqrt{2/3}) = 2.526$ and $\theta_3 = \cos^{-1}(-\sqrt{5/6}) = 2.721$ up to $1/N_c$ corrections. However, it is known phenomenologically that the contributions of those operators are weak, and thus the mixing angles are significantly affected by the subleading in $1/N_c$ operators, in particular the hyperfine operator O_6 . The determination of the mixing angles requires in principle more information than just the masses, as there are seven masses, and nine mass operators up to the order $1/N_c$, which means that the angles cannot be predicted. A biased prediction is obtained by neglecting the three-body operators, which gives one angle as a function of the other one according to the relation [24]

$$\begin{aligned} & 3(M_{N_{\frac{1}{2}}} + M_{N'_{\frac{1}{2}}} - 4M_{N_{\frac{3}{2}}} - 4M_{N'_{\frac{3}{2}}} + 6M_{N_{\frac{5}{2}}} + 8M_{\Delta_{\frac{1}{2}}} - 8M_{\Delta_{\frac{3}{2}}}) \\ &= (13 \cos 2\theta_1 + \sqrt{32} \sin 2\theta_1)(M_{N'_{\frac{1}{2}}} - M_{N_{\frac{1}{2}}}) \\ & \quad - 4(\cos 2\theta_3 - \sqrt{20} \sin 2\theta_3)(M_{N'_{\frac{3}{2}}} - M_{N_{\frac{3}{2}}}). \end{aligned} \quad (7)$$

However a determination of the angles in a more rigorous way requires the input of additional observables, namely the partial decay widths and/or photo-couplings. The details of that analysis are provided in Ref. [24].

In the case of three flavors there are two-state and also three-state mixings. For the nucleons one has the same case as for two flavors, while for Σ , Λ and Ξ baryons there is three-state mixing. The physical states are given in terms of the quark spin and SU(3) eigenstates by

$$\begin{pmatrix} \mathbf{10}_S \quad \text{or} \quad \mathbf{1}_S \\ \mathbf{8}_S \\ \mathbf{8}'_S \end{pmatrix} = \Theta \begin{pmatrix} {}^2\mathbf{10}_S \quad \text{or} \quad {}^2\mathbf{1}_S \\ {}^2\mathbf{8}_S \\ {}^4\mathbf{8}_S \end{pmatrix}, \quad (8)$$

where the physical states are indicated by the dominant SU(3) content, and the Euler mixing matrix is given by

$$\Theta = \begin{pmatrix} c\phi c\psi - c\theta s\phi s\psi & c\psi s\phi + c\theta c\phi s\psi & s\theta s\psi \\ -c\theta c\psi s\phi - c\phi s\psi & c\theta c\phi c\psi - s\phi s\psi & c\psi s\theta \\ s\theta s\phi & -c\phi s\theta & c\theta \end{pmatrix}, \quad (9)$$

$c\theta \equiv \cos \theta$, $s\theta \equiv \sin \theta$, etc.

The angles θ can always be taken in the interval $[0, \pi)$. The mixing angles ϕ and ψ vanish in the limit of exact SU(3) symmetry, and are thus proportional to the parameter ϵ . The

SU(3)-symmetric limit becomes similar to the nonstrange case except that there are two additional masses, namely the ones of the singlet Λ baryons. The determination of the mixing angles would be similar to the nonstrange case. In the absence of additional information to that of the masses, the angles can be determined only through the exclusion of some operators. For instance, one strategy would be to exclude the three-body operators, which seem in general to have particularly weak contributions to masses.

For the states which are subjected to mixing it is necessary to make the identification of the physical states. As mentioned in the Introduction, for the physical case the identification has been clear for a long time, thanks to the simultaneous use of strong decay partial widths and helicity amplitudes [21,24,39,40], but that information is not available for the LQCD baryons. The identifications of the LQCD states were analyzed separately (a total of 256 possibilities). It turns out that most assignments pass the tests of χ^2 , mass relations and the naturalness of the coefficients. Thus on a general rigorous ground the problem of state assignment is not completely resolved. However, if one requires that the coefficients flow reasonably smoothly towards the physical ones which are known, then only one assignment becomes possible, namely the one discussed here.

Since the mixing angles would be an indicator of level-crossing effects as M_π is varied, understanding them is an important task. In fact, recent studies of lower-lying negative-parity states in Ref. [8,10] identified the two lowest-lying $N_{1/2}^-$ masses and may give the first evidence of such a level crossing as M_π varies.

For two flavors, and following the global analysis of Ref. [24], the two mixing angles are given as input, and in this way it is possible to fit with the complete basis of operators up to two-body. If the additional information provided by partial decay widths and/or photo-couplings is not available, as is the case for the LQCD results, one possibility is to neglect some of the basis operators, which allows one to predict the mixing angles solely using the masses. Some guidance regarding what operators to exclude is given by the rather clear hierarchy of the importance the different operators have, as measured by the magnitude of their coefficients. In fact, it becomes clear that the mixing angles are mostly controlled by the operators O_2 , O_6 and to a lesser extend O_4 and O_5 .

For two flavors, the LQCD results are those in Ref. [3], and the corresponding fits are shown in Table IX. The physical case is in good agreement with previous works [18,19]. If one considers only the seven known masses as inputs to the fit, one operator must be eliminated: the operator O_8 is thus dismissed as it is always virtually irrelevant. A second fit where only the three dominant operators are kept turns out to be consistent for the LQCD results, but gives a poor fit to the physical case. In that case, the M_π evolution of the coefficients is shown in Fig. 3.

TABLE IX. Fits to the nonstrange $[20', 1^-]$ baryon masses. Unless the mixing angles are inputs to the fit, the operator O_8 is not necessary due to linear dependence as there are only seven mass inputs to fit. For the physical case with the seven-parameter fit, the mixing angles from the global analysis ($\theta_{N_{1/2}} = 0.49 \pm 0.29$, $\theta_{N_{3/2}} = 3.01 \pm 0.17$) were used as inputs. For the minimal fit with c_1 , c_5 , c_6 , the mixing angles in the physical case are not inputs.

Coefficients [MeV]	M_π [MeV]					
	PDG	396	524	PDG	396	524
c_1	463 ± 2	543 ± 5	598 ± 3	459 ± 2	533 ± 5	579 ± 3
c_2	-36 ± 12	39 ± 35	13 ± 14	0	0	0
c_3	313 ± 69	-83 ± 215	-96 ± 74	0	0	0
c_4	65 ± 31	-70 ± 71	-95 ± 30	0	0	0
c_5	71 ± 18	99 ± 48	107 ± 24	16 ± 18	122 ± 46	106 ± 23
c_6	443 ± 10	446 ± 25	307 ± 13	443 ± 10	502 ± 25	414 ± 13
c_7	-20 ± 31	-0.37 ± 62.89	-66 ± 34	0	0	0
$\theta_{N_{1/2}}$ [rad]	0.52 ± 0.13	2.94 ± 0.21	2.76 ± 0.06	3.13 ± 0.01	3.04 ± 0.05	3.03 ± 0.03
$\theta_{N_{3/2}}$ [rad]	3.02 ± 0.09	2.88 ± 0.42	2.38 ± 0.11	3.12 ± 0.02	2.98 ± 0.08	2.97 ± 0.05
χ^2_{dof}	0.05	0	0	0.68	0.52	1.0

A comparison of the physical case shows that it is consistent with earlier work [18,19], but differs significantly for the coefficients c_3 and c_6 with respect to the recent global analysis carried out in Ref. [24]. Since all those fits are consistent in terms of the χ^2 , it is an indication of the ambiguity that results when only the masses are fitted. This means that for the LQCD results one should also expect several consistent fits in terms of the value of the χ^2 , which will have significantly different values for some of the parameters.

Now the fits to the three-flavor case are presented. The identification of the states has been performed as described earlier. Such identification is displayed in Table XXII of Appendix B. For the sake of brevity, only those operators which have effects of any significance have been included here: after an initial analysis, several operators whose coefficients are consistent with zero have been eliminated. The fits for three flavors are given in Tables X and XII for the corresponding subsets of operators. Because of the different definitions of the basis operators for the different multiplets, in order to compare contributions which are of

common nature across multiplets such as the spin-flavor singlet contributions, the HF and the SU(3) breaking, the following identification of coefficients should be done: $c_{1_{56}} \leftrightarrow (c_1 + (b_1 + b_2)/\sqrt{3})_{70}$, $c_{2_{[56,0^+]}} \leftrightarrow c_{3_{[56,2^+]}} \leftrightarrow \frac{1}{3}c_{6_{70}}$, $b_{1_{56}} \leftrightarrow -((b_1 + b_2)\sqrt{3}/2)_{70}$.

The fits in the physical case are checked to be consistent with previous analyses [20,21]. The evolution of the fit coefficients with M_π is depicted in Fig. 4. It is interesting to observe the evolution of the mixing angles θ with M_π , as they can give a clue to the possible level crossing as M_π evolves. Similar to what is observed in the nonstrange baryons discussed above, in the $S = 3/2$ baryons the mixing angles remain approximately continuous from the physical case to $M_\pi = 702$ MeV, while in the $S = 1/2$ baryons there is a change by more than $\pi/2$, indicating a level crossing along the way. This qualitatively agrees with the LQCD results in Refs. [8,10]. It is interesting to

TABLE X. Fit to the $[70, 1^-]$ masses using a subset of operators chosen as a minimal subset such that the χ^2_{dof} is acceptable for all input sets. For the physical case the mixing angles from the global analysis [24] ($\theta_{N_{1/2}} = 0.49 \pm 0.29$, $\theta_{N_{3/2}} = 3.01 \pm 0.17$) were used as inputs.

Coefficients [MeV]	M_π [MeV]			
	PDG	391	524	702
c_1	444.3 ± 0.3	572 ± 2	585 ± 1	636 ± 1
c_2	84 ± 2	68 ± 12	-7 ± 6	-16 ± 4
c_3	117 ± 13	59 ± 22	-40 ± 18	2 ± 8
c_4	115 ± 5	-12 ± 12	-28 ± 9	-13 ± 4
c_5	84 ± 10	134 ± 17	132 ± 14	84 ± 7
c_6	538 ± 5	327 ± 10	350 ± 6	262 ± 4
c_7	-159 ± 13	49 ± 27	-59 ± 17	13 ± 11
b_1	-214 ± 5	-100 ± 13	-43 ± 9	0
b_2	-188 ± 2	-62 ± 6	-46 ± 4	0
b_3	-92 ± 2	-41 ± 10	-6 ± 7	0
χ^2_{dof}	0.74	0.65	0.14	0.09




FIG. 3 (color online). Evolution with M_π of the minimum set of operator coefficients for the $[20, 1^-]$ baryon masses in $SU(4) \times O(3)$.

TABLE XI. Mixing angles in the $[70, 1^-]$ predicted from the fit in Table X.

Mixing angles [rad]	M_π [MeV]			
	PDG	391	524	702
$\theta_{N_{1/2}}$	0.76 ± 0.03	0.61 ± 0.12	2.77 ± 0.06	2.98 ± 0.05
$\theta_{N_{3/2}}$	3.09 ± 0.40	0.10 ± 0.81	2.70 ± 0.10	2.84 ± 0.03
$\phi_{\Lambda_{1/2}}$	-0.15 ± 0.01	-0.15 ± 0.01	-0.14 ± 0.01	0
$\theta_{\Lambda_{1/2}}$	0.83 ± 0.01	0.70 ± 0.01	2.76 ± 0.01	2.98 ± 0.05
$\psi_{\Lambda_{1/2}}$	0.05 ± 0.01	0.11 ± 0.01	-0.18 ± 0.02	0
$\phi_{\Lambda_{3/2}}$	-0.21 ± 0.03	-0.16 ± 0.04	-0.12 ± 0.02	0
$\theta_{\Lambda_{3/2}}$	3.08 ± 0.01	0.13 ± 0.01	2.69 ± 0.02	2.84 ± 0.03
$\psi_{\Lambda_{3/2}}$	-0.18 ± 0.01	0.07 ± 0.03	-0.03 ± 0.01	0
$\phi_{\Sigma_{1/2}}$	-0.25 ± 0.02	0.03 ± 0.01	-0.05 ± 0.04	0
$\theta_{\Sigma_{1/2}}$	1.01 ± 0.01	0.75 ± 0.01	2.75 ± 0.01	2.98 ± 0.05
$\psi_{\Sigma_{1/2}}$	-0.10 ± 0.01	0.01 ± 0.07	0.03 ± 0.04	0
$\phi_{\Sigma_{3/2}}$	-0.08 ± 0.06	0.06 ± 0.04	-0.02 ± 0.04	0
$\theta_{\Sigma_{3/2}}$	3.05 ± 0.01	0.16 ± 0.02	2.66 ± 0.01	2.84 ± 0.03
$\psi_{\Sigma_{3/2}}$	0.04 ± 0.02	0.03 ± 0.02	0.005 ± 0.001	0
$\phi_{\Xi_{1/2}}$	-0.30 ± 0.03	0.03 ± 0.01	-0.05 ± 0.06	0
$\theta_{\Xi_{1/2}}$	0.94 ± 0.01	0.78 ± 0.01	2.77 ± 0.04	2.98 ± 0.05
$\psi_{\Xi_{1/2}}$	-0.14 ± 0.02	0.01 ± 0.07	0.03 ± 0.06	0
$\phi_{\Xi_{3/2}}$	-0.09 ± 0.07	0.05 ± 0.03	-0.02 ± 0.04	0
$\theta_{\Xi_{3/2}}$	3.07 ± 0.01	0.19 ± 0.03	2.69 ± 0.02	2.84 ± 0.03
$\psi_{\Xi_{3/2}}$	0.05 ± 0.03	0.02 ± 0.01	0.006 ± 0.001	0

observe that for $M_\pi = 702$ MeV all baryons are stable, and almost all are still stable for $M_\pi = 524$ MeV, while below $M_\pi = 391$ MeV they are unstable. Since the $S = 1/2$ baryons have S -wave decays, they are the ones that are sensitive to the opening of the decay. These observations suggest a synchronization between the mixing angle and the stability of the baryon. In fact, the change in θ_1 shown in Table XI in going from $M_\pi = 391$ to 524 MeV is approximately $\pi/2$, as expected for a level crossing. Is this an explanation for the observed level crossings? Perhaps, but it is not clear at this point, and it deserves further study.




FIG. 4 (color online). Evolution with respect to M_π of the coefficients of the basis operators used to fit both the physical and the LQCD $[70, 1^-]$ masses.

Consistent fits to only LQCD results can be achieved by a minimal set of significant operators. It is found that the relevant SU(3) singlet operators are the spin-flavor singlet O_1 , the HF O_6 and the two spin-orbit ones O_2 and O_5 and the first two SU(3)-breaking operators. These results are illustrated in Fig. 5. Note that all the SU(3)-breaking operators are relevant for fitting the physical case. The operator O_3 is found to be important for the physical masses, but irrelevant for the LQCD masses, where the operator O_5 is instead significant. It is interesting to observe that in models with pion exchange between quarks,

TABLE XII. Fit results with minimal set of mass operators for the $[70, 1^-]$. Only masses are used as inputs.

Coefficients	M_π [MeV]			
	PDG	391	524	702
c_1	462 ± 0.3	582 ± 2	587 ± 1	637 ± 1
c_2	83 ± 2	92 ± 10	13 ± 8	-11 ± 4
c_5	-67 ± 11	136 ± 17	127 ± 13	96 ± 7
c_6	420 ± 4	270 ± 9	344 ± 6	257 ± 4
c_7	-78 ± 14	4 ± 31	-47 ± 16	21 ± 11
b_1	-92 ± 4	-53 ± 13	-34 ± 9	0
b_2	-179 ± 2	-58 ± 6	-48 ± 4	0
$\theta_{N_{1/2}}$	0.33 ± 0.02	0.79 ± 0.21	2.95 ± 0.05	2.94 ± 0.02
$\theta_{N_{3/2}}$	0.45 ± 0.02	0.79 ± 0.13	2.86 ± 0.07	2.84 ± 0.03
χ^2_{dof}	6.7	0.86	0.46	0.13




FIG. 5 (color online). Evolution with respect to M_π of the coefficients in Table XII.

such as certain versions of the chiral quark model, O_3 is naturally important, and should fade as M_π increases.

The mass relations are depicted in Tables XIII and XIV. All are well satisfied, except for the EQS relation for $M_\pi = 391$ MeV involving $\Sigma''_{3/2}$. A shift of its mass by $\sim +30$ MeV leads to consistency. The mass predictions

are given in Table XV. Since, the PDG candidate state $\Xi(1950)^{***}(\?)$ is consistent with $\Xi'_{3/2}$, $\Xi_{5/2}$ and $\Xi''_{1/2}$ in Table XV, its parity could be predicted to be negative.

IV. COMMENTS AND CONCLUSIONS

From the study presented here of recent LQCD results for the low-lying baryon excitations, it can be concluded that a picture of the spin-flavor composition of excited baryons is derived from their masses calculated in LQCD and the $1/N_c$ expansion. The results obtained entirely support the picture seen from the lattice QCD analysis of the mass eigenstate couplings to source/sink operators. A similar, and even simpler picture than the physical case emerges at increasing quark masses, where with very few dominant operators the LQCD masses can be described. The expected narrowness of the states analyzed for the quark masses in the LQCD results suggests that those results are very realistic. For higher excited baryons, which will be broader, the present LQCD results may be a poorer approximation. Nonetheless, they should be interesting to study.

TABLE XIII. GMO and EQS relations for the $[70, 1^-]$ multiplet. Due to the insufficient number of known states with three or more stars, the mass relations cannot be checked for the physical case.

Relation	M_π [MeV]		
	PDG	391	524
$2(N_{1/2} + \Xi_{1/2}) - (3\Lambda_{1/2} + \Sigma_{1/2}) = 0$...	59 ± 156	17 ± 125
$2(N_{3/2} + \Xi_{3/2}) - (3\Lambda_{3/2} + \Sigma_{3/2}) = 0$...	31 ± 121	13 ± 74
$2(N_{5/2} + \Xi_{5/2}) - (3\Lambda_{5/2} + \Sigma_{5/2}) = 0$...	46 ± 91	6 ± 64
$\Sigma''_{1/2} - \Delta_{1/2} = \Xi''_{1/2} - \Sigma''_{1/2} = \Omega_{1/2} - \Xi''_{1/2}$...	67 ± 47	35 ± 56
...	...	34 ± 36	40 ± 41
...	...	24 ± 49	22 ± 26
$\Sigma''_{3/2} - \Delta_{3/2} = \Xi''_{3/2} - \Sigma''_{3/2} = \Omega_{3/2} - \Xi''_{3/2}$...	2 ± 49	39 ± 23
...	...	82 ± 47	37 ± 21
...	...	61 ± 43	31 ± 21

TABLE XIV. Octet-decuplet mass relations for the $[70, 1^-]$ multiplet. S_B is the mass splitting between the state B and the nonstrange states in the SU(3) multiplet to which it belongs. The results shown correspond to the relation divided by the sum of the positive coefficients in the relation (e.g., 163 for the first relation).

Relation	M_π [MeV]	
	391	524
$14(S_{\Lambda_{3/2}} + S_{\Lambda'_{3/2}}) + 63S_{\Lambda_{5/2}} + 36(S_{\Sigma_{1/2}} + S_{\Sigma'_{1/2}}) - 68(S_{\Lambda_{1/2}} + S_{\Lambda'_{1/2}}) - 27S_{\Sigma_{5/2}} = 0$	9.4 ± 40	0.96 ± 34
$14(S_{\Sigma_{3/2}} + S_{\Sigma'_{3/2}}) + 21S_{\Lambda_{5/2}} - 9S_{\Sigma_{5/2}} - 18(S_{\Lambda_{1/2}} + S_{\Lambda'_{1/2}}) - 2(S_{\Sigma_{1/2}} + S_{\Sigma'_{1/2}}) = 0$	37 ± 45	5.4 ± 38
$14S_{\Sigma'_{1/2}} + 49S_{\Lambda_{5/2}} + 23(S_{\Sigma_{1/2}} + S_{\Sigma'_{1/2}}) - 45(S_{\Lambda_{1/2}} + S_{\Lambda'_{1/2}}) - 19S_{\Sigma_{5/2}} = 0$	9.4 ± 40	0.7 ± 34
$14S_{\Sigma'_{3/2}} + 28S_{\Lambda_{5/2}} + 11(S_{\Sigma_{1/2}} + S_{\Sigma'_{1/2}}) - 27(S_{\Lambda_{1/2}} + S_{\Lambda'_{1/2}}) - 10S_{\Sigma_{5/2}} = 0$	0.8 ± 40	0.1 ± 33

TABLE XV. Predictions of physically unknown states in the $[70, 1^-]$ multiplet from the fit in Table X.

Missing state	Fitted mass with union set of operators [MeV]	PDG[MeV]
$\Sigma_{1/2}$	1645 ± 11	$\Sigma(1620)1/2^{***} = 1620 \pm 10$
$\Xi_{1/2}$	1801 ± 11	...
$\Xi_{1/2}'$	1930 ± 8	...
$\Lambda_{3/2}'$	1825 ± 6	...
$\Sigma_{3/2}'$	1780 ± 7	...
$\Xi_{3/2}'$	1944 ± 7	$\Xi(1950)(?)^{***} = 1950 \pm 15$
$\Xi_{5/2}$	1939 ± 6	$\Xi(1950)(?)^{***} = 1950 \pm 15$
$\Sigma_{1/2}''$	1828 ± 7	...
$\Xi_{1/2}''$	1969 ± 8	$\Xi(1950)(?)^{***} = 1950 \pm 15$
$\Omega_{1/2}$	2107 ± 10	...
$\Sigma_{3/2}''$	1916 ± 6	$\Sigma(1940)3/2^{***} = 1950 \pm 30$
$\Xi_{3/2}''$	2057 ± 7	...
$\Omega_{3/2}$	2198 ± 9	...

A strong conclusion is that the LQCD masses are even closer to an approximate $SU(6) \times O(3)$ symmetry limit than the physical ones. This is most likely due to the fact that the composition of baryons becomes increasingly closer to a constituent quark model picture as the quark masses increase, emphasizing the mass operators which are naturally large in those models and suppressing the rest.

For the quark masses employed in the LQCD calculations used here, the dramatic downturn in c_1 for the Roper baryons is not manifest. This is an effect where probably chiral symmetry plays an important role as it becomes restored towards lighter M_π . In recent LQCD work on nucleon resonances [8,10] a first evidence of that downturn is observed. It remains to be determined what precise mechanism drives that effect, and perhaps some clever strategy in LQCD calculations could be used for that purpose. While in the Roper multiplet the c_1 coefficient should have a large negative curvature as a function of M_π to match the physical masses, it lies along an almost perfect straight line for the ground-state baryons, and it has a moderate negative curvature in the other cases.

Identifying the HF coefficients as mentioned earlier, one finds that for the LQCD results the strength of the HF in the ground-state baryons is almost twice as large as in the excited baryons, which is significantly different from the physical case, where it is only about 25% larger. One should point out that in a picture of large N_c baryons with

very heavy quark masses the hyperfine interaction will scale as m_Q/N_c , implying that in LQCD calculations at even larger quark masses than the ones used so far the hyperfine interaction coefficient should eventually scale approximately as m_Q .

The spin-orbit contributions are all smaller than the natural size. In the $[56, 2^+]$ it is an effect $\mathcal{O}(1/N_c)$, and the coefficient slowly decreases with increasing M_π . In the $[70, 1^-]$ the $\mathcal{O}(N_c^0)$ contribution is determined by c_2 which decreases with increasing M_π , and the $\mathcal{O}(1/N_c)$ contribution determined by c_5 remains roughly independent of M_π .

An interesting open problem is how to relate the $SU(6) \times O(3)$ decomposition of the physical baryons determined via the $1/N_c$ expansion as presented here, with the information on the coupling strengths of the mass eigenstates to the different source/sink operators obtained in the LQCD calculations.

ACKNOWLEDGMENTS

The authors thank Jozef Dudek and Robert Edwards for useful discussions. This work was supported in part by DOE Contract No. DE-AC05-06OR23177 under which JSA operates the Thomas Jefferson National Accelerator Facility (J. L. G.), and by the National Science Foundation through Grants No. PHY-0855789 and PHY-1307413 (I. P. F. and J. L. G.).

APPENDIX A: BASES OF MASS OPERATORS

This Appendix gives the bases of mass operators with the respective normalization factors used in this work.

1. Operator basis and matrix elements for the $[56,2^+]$ multiplet

TABLE XVI. Matrix elements of SU(3) singlet operators (top) and SU(3)-breaking operators (bottom). Here, $a_S = 1, -2/3$ for $S = 3/2, 5/2$, respectively and $b_S = 1, 2/3, 1/9, -2/3$ for $S = 1/2, 3/2, 5/2, 7/2$, respectively

	O_1	O_2	O_3
	$N_c \mathbf{1}$	$\frac{1}{N_c} \ell_i S_i$	$\frac{1}{N_c} S_i S_i$
$8_{3/2}$	N_c	$-\frac{3}{2N_c}$	$\frac{3}{4N_c}$
$8_{5/2}$	N_c	$\frac{1}{N_c}$	$\frac{3}{4N_c}$
$10_{1/2}$	N_c	$-\frac{9}{2N_c}$	$\frac{15}{4N_c}$
$10_{3/2}$	N_c	$-\frac{3}{N_c}$	$\frac{15}{4N_c}$
$10_{5/2}$	N_c	$-\frac{1}{2N_c}$	$\frac{15}{4N_c}$
$10_{7/2}$	N_c	$\frac{3}{N_c}$	$\frac{15}{4N_c}$
	\bar{B}_1	\bar{B}_2	\bar{B}_3
	N_S	$\frac{1}{N_c} \ell_i G_{i8} - \frac{1}{2\sqrt{3}} O_2$	$\frac{1}{N_c} S_i G_{i8} - \frac{1}{2\sqrt{3}} O_3$
N_S	0	0	0
Λ_S	1	$\frac{3\sqrt{3}a_S}{4N_c}$	$-\frac{3\sqrt{3}}{8N_c}$
Σ_S	1	$-\frac{\sqrt{3}a_S}{4N_c}$	$\frac{\sqrt{3}}{8N_c}$
Ξ_S	2	$\frac{\sqrt{3}a_S}{N_c}$	$-\frac{\sqrt{3}}{2N_c}$
Δ_S	0	0	0
Σ''_S	1	$\frac{3\sqrt{3}b_S}{4N_c}$	$-\frac{5\sqrt{3}}{8N_c}$
Ξ''_S	2	$\frac{3\sqrt{3}b_S}{2N_c}$	$-\frac{5\sqrt{3}}{4N_c}$
Ω_S	3	$\frac{9\sqrt{3}b_S}{4N_c}$	$-\frac{15\sqrt{3}}{8N_c}$
$\Sigma_{3/2} - \Sigma''_{3/2}$	0	$\frac{\sqrt{3}}{2N_c}$	0
$\Sigma_{5/2} - \Sigma''_{5/2}$	0	$\frac{\sqrt{3}}{2N_c}$	0
$\Xi_{3/2} - \Xi''_{3/2}$	0	$\frac{\sqrt{42}}{6N_c}$	0
$\Xi_{5/2} - \Xi''_{5/2}$	0	$\frac{\sqrt{42}}{6N_c}$	0

2. Operator basis and matrix elements for the $[20,1^-]$ multipletTABLE XVII. Mass operator basis and matrix elements for the $[20', 1^-]$ multiplet [19].

	O_1	O_2	O_3	O_4
	$N_c \mathbf{1}$	$\ell_i s_i$	$\frac{3}{N_c} \ell_{ij}^{(2)} g_{ia} G_{ja}^c$	$\ell_i s_i + \frac{4}{N_c+1} \ell_i t_a G_{ia}^c$
$N_{1/2}$	N_c	$-\frac{(2N_c-3)}{3N_c}$	0	$\frac{2}{N_c+1}$
$N'_{1/2}$	N_c	$-\frac{5}{6}$	$-\frac{5(N_c+1)}{48N_c}$	0
$N'_{1/2} - N_{1/2}$	0	$-\frac{1}{3} \sqrt{\frac{N_c+3}{2N_c}}$	$-\frac{5}{48N_c} \sqrt{\frac{(N_c+3)(2N_c-1)^2}{2N_c}}$	$-\sqrt{\frac{N_c+3}{2N_c(N_c+1)^2}}$
$N_{3/2}$	N_c	$\frac{(2N_c-3)}{6N_c}$	0	$-\frac{1}{N_c+1}$
$N'_{3/2}$	N_c	$-\frac{1}{3}$	$\frac{1}{12N_c} (N_c + 1)$	0
$N'_{3/2} - N_{3/2}$	0	$-\frac{1}{6} \sqrt{\frac{5(N_c+3)}{N_c}}$	$\frac{1}{96N_c} \sqrt{\frac{5(N_c+3)(2N_c-1)^2}{N_c}}$	$-\sqrt{\frac{5(N_c+3)}{4N_c(N_c+1)^2}}$
$N'_{5/2}$	N_c	$+\frac{1}{2}$	$-\frac{1}{48N_c} (N_c + 1)$	0
$\Delta_{1/2}$	N_c	$+\frac{1}{3}$	0	0
$\Delta_{3/2}$	N_c	$-\frac{1}{6}$	0	0
	O_5	O_6	O_7	O_8
	$\frac{1}{N_c} \ell_i S_i^c$	$\frac{1}{N_c} S_i^c S_i^c$	$\frac{1}{N_c} s_i S_i^c$	$\frac{2}{N_c} \ell_{ij}^{(2)} s_i S_j^c$
$N_{1/2}$	$-\frac{(N_c+3)}{3N_c^2}$	$\frac{(N_c+3)}{2N_c^2}$	$-\frac{(N_c+3)}{4N_c^2}$	0
$N'_{1/2}$	$-\frac{5}{3N_c}$	$\frac{2}{N_c}$	$\frac{1}{2N_c}$	$\frac{5}{6N_c}$
$N'_{1/2} - N_{1/2}$	$\sqrt{\frac{N_c+3}{18N_c^2}}$	0	0	$\frac{5}{12N_c} \sqrt{\frac{N_c+3}{2N_c}}$
$N_{3/2}$	$\frac{(N_c+3)}{6N_c^2}$	$\frac{(N_c+3)}{2N_c^2}$	$-\frac{(N_c+3)}{4N_c^2}$	0
$N'_{3/2}$	$-\frac{2}{3N_c}$	$\frac{2}{N_c}$	$\frac{1}{2N_c}$	$-\frac{2}{3N_c}$
$N'_{3/2} - N_{3/2}$	$\sqrt{\frac{5(N_c+3)}{36N_c^3}}$	0	0	$-\frac{1}{24} \sqrt{\frac{5(N_c+3)}{N_c^3}}$
$N'_{5/2}$	$\frac{1}{N_c}$	$\frac{2}{N_c}$	$\frac{1}{2N_c}$	$\frac{1}{6N_c}$
$\Delta_{1/2}$	$-\frac{4}{3N_c}$	$\frac{2}{N_c}$	$-\frac{1}{N_c}$	0
$\Delta_{3/2}$	$\frac{2}{3N_c}$	$\frac{2}{N_c}$	$-\frac{1}{N_c}$	0

3. Operator basis and matrix elements for the $[70, 1^-]$ multipletTABLE XVIII. SU(3) singlet basis of operators for the $[70, 1^-]$ masses [21].

	O_1	O_2	O_3	O_4	O_5	O_6
	$N_c 1$	$\ell_i s_i$	$\frac{3}{N_c} \ell_{ij}^{(2)} g_{ia} G_{ja}^c$	$\frac{4}{N_c+1} \ell_i t_a G_{ia}^c$	$\frac{1}{N_c} \ell_i S_i^c$	$\frac{1}{N_c} S_i^c S_i^c$
${}^2 8_{1/2}$	N_c	$\frac{3-2N_c}{3N_c}$	0	$\frac{2(N_c+3)(3N_c-2)}{9N_c(N_c+1)}$	$-\frac{N_c+3}{3N_c^2}$	$\frac{N_c+3}{2N_c^2}$
${}^4 8_{1/2}$	N_c	$-\frac{5}{6}$	$-\frac{5(3N_c+1)}{48N_c}$	$\frac{5(3N_c+1)}{18(N_c+1)}$	$-\frac{5}{3N_c}$	$\frac{2}{N_c}$
${}^2 8_{1/2} - {}^4 8_{1/2}$	0	$-\sqrt{\frac{(N_c+3)}{18N_c}}$	$-\frac{5}{24} \sqrt{\frac{(N_c+3)(3N_c-2)^2}{2N_c^3}}$	$-\frac{(5-3N_c)}{9(N_c+1)} \sqrt{\frac{N_c+3}{2N_c}}$	$\sqrt{\frac{N_c+3}{18N_c^3}}$	0
${}^2 1_{1/2}$	N_c	-1	0	0	0	0
${}^2 10_{1/2}$	N_c	$\frac{1}{3}$	0	$-\frac{(3N_c+7)}{9(N_c+1)}$	$-\frac{4}{3N_c}$	$\frac{2}{N_c}$
${}^2 8_{3/2}$	N_c	$\frac{2N_c-3}{6N_c}$	0	$-\frac{(N_c+3)(3N_c-2)}{9N_c(N_c+1)}$	$\frac{N_c+3}{6N_c^2}$	$\frac{N_c+3}{2N_c^2}$
${}^4 8_{3/2}$	N_c	$-\frac{1}{3}$	$\frac{3N_c+1}{12N_c}$	$\frac{3N_c+1}{9(N_c+1)}$	$-\frac{2}{3N_c}$	$\frac{2}{N_c}$
${}^2 8_{3/2} - {}^4 8_{3/2}$	0	$-\sqrt{\frac{5(N_c+3)}{36N_c}}$	$-\frac{1}{48} \sqrt{\frac{5(N_c+3)(2-3N_c)^2}{N_c^3}}$	$-\sqrt{\frac{5(N_c+3)(5-3N_c)^2}{324N_c(N_c+1)^2}}$	$\sqrt{\frac{5(N_c+3)}{36N_c^3}}$	0
${}^2 1_{3/2}$	N_c	$\frac{1}{2}$	0	0	0	0
${}^2 10_{3/2}$	N_c	$-\frac{1}{6}$	0	$\frac{3N_c+7}{18(N_c+1)}$	$\frac{2}{3N_c}$	$\frac{2}{N_c}$
${}^4 8_{5/2}$	N_c	$\frac{1}{2}$	$-\frac{3N_c+1}{48N_c}$	$-\frac{3N_c+1}{6(N_c+1)}$	$\frac{1}{N_c}$	$\frac{2}{N_c}$
	O_7	O_8	O_9	O_{10}	O_{11}	
	$\frac{1}{N_c} s_i S_i^c$	$\frac{2}{N_c} \ell_{ij}^{(2)} s_i S_j^c$	$\frac{3}{N_c^2} \ell_i g_{ja} \{S_i^c, G_{ja}^c\}$	$\frac{2}{N_c^2} t_a \{S_i^c, G_{ia}^c\}$	$\frac{3}{N_c^2} \ell_i g_{ia} \{S_j^c, G_{ja}^c\}$	
${}^2 8_{1/2}$	$-\frac{N_c+3}{4N_c^2}$	0	$\frac{(N_c+3)(7-15N_c)}{24N_c^3}$	$-\frac{(N_c+3)(3N_c+1)}{12N_c^3}$	$-\frac{(N_c+3)(3N_c+1)}{24N_c^3}$	
${}^4 8_{1/2}$	$\frac{1}{2N_c}$	$\frac{5}{3N_c}$	$\frac{5(3N_c+1)}{24N_c^2}$	$-\frac{(3N_c+1)}{3N_c^2}$	$\frac{5(3N_c+1)}{12N_c^2}$	
${}^2 8_{1/2} - {}^4 8_{1/2}$	0	$\sqrt{\frac{25(N_c+3)}{72N_c^3}}$	$\sqrt{\frac{(N_c+3)(3N_c-2)^2}{288N_c^3}}$	0	$\sqrt{\frac{(N_c+3)(3N_c+1)^2}{72N_c^3}}$	
${}^2 1_{1/2}$	0	0	0	0	0	
${}^2 10_{1/2}$	$-\frac{1}{N_c}$	0	$\frac{(3N_c+7)}{6N_c^2}$	$\frac{(3N_c+7)}{6N_c^2}$	$\frac{(3N_c+7)}{12N_c^2}$	
${}^2 8_{3/2}$	$-\frac{N_c+3}{4N_c^2}$	0	$\frac{(N_c+3)(15N_c-7)}{48N_c^3}$	$-\frac{(N_c+3)(3N_c+1)}{12N_c^3}$	$\frac{(N_c+3)(3N_c+1)}{48N_c^3}$	
${}^4 8_{3/2}$	$\frac{1}{2N_c}$	$-\frac{4}{3N_c}$	$\frac{(3N_c+1)}{12N_c^2}$	$-\frac{(3N_c+1)}{3N_c^2}$	$\frac{(3N_c+1)}{6N_c^2}$	
${}^2 8_{3/2} - {}^4 8_{3/2}$	0	$-\sqrt{\frac{5(N_c+3)}{144N_c^3}}$	$\sqrt{\frac{5(N_c+3)(3N_c-2)^2}{576N_c^3}}$	0	$\sqrt{\frac{5(N_c+3)(3N_c+1)^2}{144N_c^3}}$	
${}^2 1_{3/2}$	0	0	0	0	0	
${}^2 10_{3/2}$	$-\frac{1}{N_c}$	0	$-\frac{(3N_c+7)}{12N_c^2}$	$\frac{(3N_c+7)}{6N_c^2}$	$-\frac{(3N_c+7)}{24N_c^2}$	
${}^4 8_{5/2}$	$\frac{1}{2N_c}$	$\frac{1}{3N_c}$	$-\frac{(3N_c+1)}{8N_c^2}$	$-\frac{(3N_c+1)}{3N_c^2}$	$-\frac{(3N_c+1)}{4N_c^2}$	

TABLE XIX. SU(3) basis of SU(3)-breaking octet operators for the [70, 1⁻] masses [21].

	B_1	B_2
	t_8	T_8^c
${}^2 8_{1/2}, {}^2 8_{3/2}$	$\frac{N_c^3 - (7N_s - 8I^2)N_c^2 + 3(4N_s - 8I^2 + 1)N_c - 9N_s}{2\sqrt{3}N_c(N_c - 1)(N_c + 3)}$	$\frac{N_c^4 - (3N_s - 1)N_c^3 + (N_s - 8I^2 - 3)N_c^2 - 3(N_s - 8I^2 + 1)N_c + 9N_s}{2\sqrt{3}N_c(N_c - 1)(N_c + 3)}$
${}^4 8_{1/2}, {}^4 8_{3/2}, {}^4 8_{5/2}$	$\frac{N_c - N_s - 4I^2}{2\sqrt{3}(N_c - 1)}$	$\frac{N_c^2 - (3N_s + 2)N_c + 4(I^2 + N_s)}{2\sqrt{3}(N_c - 1)}$
${}^2 8_{1/2} - {}^4 8_{1/2}, {}^2 8_{3/2} - {}^4 8_{3/2}$	0	0
${}^2 1_{1/2}, {}^2 1_{3/2}$	$\frac{(3 - N_c)}{\sqrt{3}(N_c + 3)}$	$\frac{(N_c + 5)(N_c - 3)}{2\sqrt{3}(N_c + 3)}$
${}^2 8_{1/2} - {}^2 1_{1/2}, {}^2 8_{3/2} - {}^2 1_{3/2}$	$-\frac{3(N_c - 1)}{2\sqrt{N_c}(N_c + 3)}$	$-\frac{3(N_c - 1)}{2\sqrt{N_c}(N_c + 3)}$
${}^4 8_{1/2} - {}^2 1_{1/2}, {}^4 8_{3/2} - {}^2 1_{3/2}$	0	0
$10_{1/2}, 10_{3/2}$	$\frac{N_c - 8N_s + 5}{2\sqrt{3}(N_c + 5)}$	$\frac{N_c^2 - (3N_s - 4)N_c - 7N_s - 5}{2\sqrt{3}(N_c + 5)}$
${}^2 8_{1/2} - {}^2 10_{1/2}, {}^2 8_{3/2} - {}^2 10_{3/2}$	$-\sqrt{\frac{2}{3}} \frac{N_c + 3}{N_c(N_c - 1)(N_c + 5)}$	$\sqrt{\frac{2}{3}} \frac{N_c + 3}{N_c(N_c - 1)(N_c + 5)}$
${}^4 8_{1/2} - {}^2 10_{1/2}, {}^4 8_{3/2} - {}^2 10_{3/2}$	0	0
		B_3
		$\frac{10}{N_c} d_{8ab} g_{ia} G_{ib}^c$
${}^2 8_{1/2}, {}^2 8_{3/2}$		$\frac{3N_c^3 - (13N_s - 8I^2 + 3)N_c^2 + (31N_s - 44I^2 - 12)N_c - 6(N_s - 14I^2)}{-\frac{24}{5}\sqrt{3}N_c^2(N_c - 1)}$
${}^4 8_{1/2}, {}^4 8_{3/2}, {}^4 8_{5/2}$		$\frac{3N_c^2 - (7N_s + 4I^2 - 3)N_c + (N_s - 20I^2)}{-\frac{24}{5}\sqrt{3}N_c(N_c - 1)}$
${}^2 8_{1/2} - {}^4 8_{1/2}, {}^2 8_{3/2} - {}^4 8_{3/2}$		0
${}^2 1_{1/2}, {}^2 1_{3/2}$		0
${}^2 8_{1/2} - {}^2 1_{1/2}, {}^2 8_{3/2} - {}^2 1_{3/2}$		$\frac{5(3N_c + 1)}{16N_c\sqrt{N_c}}$
${}^4 8_{1/2} - {}^2 1_{1/2}, {}^4 8_{3/2} - {}^2 1_{3/2}$		0
$10_{1/2}, 10_{3/2}$		$-\frac{3N_c^2 - 14(N_s - 1)N_c - 22N_s - 5}{\frac{24}{5}\sqrt{3}N_c(N_c + 5)}$
${}^2 8_{1/2} - {}^2 10_{1/2}, {}^2 8_{3/2} - {}^2 10_{3/2}$		$\frac{5(N_c + 2)}{6\sqrt{6}N_c} \sqrt{\frac{N_c + 3}{N_c(N_c - 1)(N_c + 5)}}$
${}^4 8_{1/2} - {}^2 10_{1/2}, {}^4 8_{3/2} - {}^2 10_{3/2}$		0
		B_4
		$3\ell_i g_{i8}$
${}^2 8_{1/2}$		$-\frac{N_c^3 - (10N_s - 14I^2 + 3)N_c^2 + 3(7N_s - 8I^2)N_c - 9(N_s - 2I^2)}{\sqrt{3}N_c(N_c - 1)(N_c + 3)}$
${}^4 8_{1/2}$		$-\frac{5(N_c - N_s - 4I^2)}{4\sqrt{3}(N_c - 1)}$
${}^2 8_{1/2} - {}^4 8_{1/2}$		$-\frac{N_c - N_s - 4I^2}{2\sqrt{6}(N_c - 1)} \sqrt{1 + \frac{3}{N_c}}$
${}^2 1_{1/2}$		$\frac{\sqrt{3}(N_c - 3)}{(N_c + 3)}$
${}^2 8_{1/2} - {}^2 1_{1/2}$		$\frac{9(N_c - 1)}{2(N_c + 3)\sqrt{N_c}}$
${}^4 8_{1/2} - {}^2 1_{1/2}$		0
${}^2 10_{1/2}$		$\frac{N_c - 8N_s + 5}{2\sqrt{3}(N_c + 5)}$
${}^2 8_{1/2} - {}^2 10_{1/2}$		$-\sqrt{\frac{2}{3}} \sqrt{\frac{N_c + 3}{N_c(N_c - 1)(N_c + 5)}}$
${}^4 8_{1/2} - {}^2 10_{1/2}$		$\frac{4}{\sqrt{3}} \frac{1}{\sqrt{(N_c - 1)(N_c + 5)}}$
${}^2 8_{3/2}$		$\frac{N_c^3 - (10N_s - 14I^2 + 3)N_c^2 + 3(7N_s - 8I^2)N_c - 9(N_s - 2I^2)}{2\sqrt{3}N_c(N_c - 1)(N_c + 3)}$
${}^4 8_{3/2}$		$-\frac{N_c - N_s - 4I^2}{2\sqrt{3}(N_c - 1)}$
${}^2 8_{3/2} - {}^4 8_{3/2}$		$-\sqrt{\frac{5}{3}} \frac{N_c - N_s - 4I^2}{4(N_c - 1)} \sqrt{1 + \frac{3}{N_c}}$
${}^2 1_{3/2}$		$-\frac{\sqrt{3}(N_c - 3)}{2(N_c + 3)}$

(Table continued)

TABLE XIX. (Continued)

	B_4
	$3\mathcal{L}_{i9i8}$
${}^2 8_{3/2} - {}^2 1_{3/2}$	$-\frac{9(N_c-1)}{4(N_c+3)\sqrt{N_c}}$
${}^4 8_{3/2} - {}^2 1_{3/2}$	0
${}^2 10_{3/2}$	$-\frac{N_c-8N_c+5}{4\sqrt{3}(N_c+5)}$
${}^2 8_{3/2} - {}^2 10_{3/2}$	$\sqrt{\frac{N_c+3}{6N_c(N_c-1)(N_c+5)}}$
${}^4 8_{3/2} - {}^2 10_{3/2}$	$\frac{2\sqrt{10}}{\sqrt{3(N_c-1)(N_c+5)}}$
${}^4 8_{5/2}$	$\frac{\sqrt{3(N_c-N_s-4I^2)}}{4(N_c-1)}$

APPENDIX B: INPUT MASSES

TABLE XX. Ground-state (left), and $[56, 0^+]$ Roper (right) baryon masses in MeV. The inversion in the ordering of the masses of the $\Xi_{1/2}$ and the Δ masses at and above $M_\pi = 391$ MeV is similar to that observed in other LQCD calculations [1].

Baryon	PDG	M_π [MeV]			Baryon	PDG	M_π [MeV]		
		391	524	702			391	524	702
$N_{1/2}$	938 ± 30	1202 ± 15	1309 ± 9	1473 ± 4	$N_{1/2}$	1450 ± 20	2221 ± 52	2300 ± 30	2339 ± 21
$\Lambda_{1/2}$	1116 ± 30	1279 ± 20	1371 ± 7	1473 ± 4	$\Lambda_{1/2}$	1630 ± 70	2189 ± 44	2330 ± 26	2339 ± 21
$\Sigma_{1/2}$	1189 ± 30	1309 ± 13	1375 ± 6	1473 ± 4	$\Sigma_{1/2}$	1660 ± 30	2252 ± 46	2357 ± 52	2339 ± 21
$\Xi_{1/2}$	1315 ± 30	1351 ± 15	1420 ± 9	1473 ± 4	$\Xi_{1/2}$...	2278 ± 22	2321 ± 54	2339 ± 21
$\Delta_{3/2}$	1228 ± 30	1518 ± 20	1582 ± 9	1673 ± 6	$\Delta_{3/2}$	1625 ± 75	2356 ± 33	2450 ± 17	2454 ± 55
$\Sigma_{3/2}$	1383 ± 30	1582 ± 15	1622 ± 6	1673 ± 6	$\Sigma_{3/2}$...	2369 ± 31	2423 ± 19	2454 ± 55
$\Xi_{3/2}$	1532 ± 30	1636 ± 11	1655 ± 11	1673 ± 6	$\Xi_{3/2}$...	2453 ± 26	2463 ± 45	2454 ± 55
$\Omega_{3/2}$	1672 ± 30	1691 ± 13	1694 ± 9	1673 ± 6	$\Omega_{3/2}$...	2501 ± 33	2504 ± 35	2454 ± 55

TABLE XXI. $[56, 2^+]$ masses. The experimental values are those for baryons with a three-star or higher rating by the PDG.

Baryon	PDG	M_π [MeV]			Baryon	PDG	M_π [MeV]		
		391	524	702			391	524	702
$N_{3/2}$	1700 ± 50	2148 ± 33	2178 ± 61	2314 ± 17	$\Delta_{3/2}$	1935 ± 35	2270 ± 37	2344 ± 17	2387 ± 19
$\Lambda_{3/2}$	1800 ± 30	2225 ± 28	2227 ± 39	2314 ± 17	$\Sigma''_{3/2}$...	2318 ± 26	2379 ± 15	2387 ± 19
$\Sigma_{3/2}$...	2243 ± 24	2238 ± 26	2314 ± 17	$\Xi''_{3/2}$...	2374 ± 13	2409 ± 6	2387 ± 19
$\Xi_{3/2}$...	2263 ± 31	2305 ± 15	2314 ± 17	$\Omega_{3/2}$...	2420 ± 28	2450 ± 13	2387 ± 19
$N_{5/2}$	1683 ± 8	2140 ± 31	2198 ± 17	2271 ± 13	$\Delta_{5/2}$	1895 ± 25	2333 ± 35	2359 ± 17	2388 ± 17
$\Lambda_{5/2}$	1820 ± 5	2228 ± 20	2249 ± 15	2271 ± 13	$\Sigma''_{5/2}$...	2368 ± 20	2392 ± 19	2388 ± 17
$\Sigma_{5/2}$	1918 ± 18	2229 ± 22	2253 ± 17	2271 ± 13	$\Xi''_{5/2}$...	2430 ± 24	2418 ± 13	2388 ± 17
$\Xi_{5/2}$...	2296 ± 22	2275 ± 13	2271 ± 13	$\Omega_{5/2}$...	2487 ± 24	2470 ± 13	2388 ± 17
$\Delta_{1/2}$	1895 ± 25	2284 ± 107	2312 ± 28	2398 ± 32	$\Delta_{7/2}$	1950 ± 10	2390 ± 31	2384 ± 19	2403 ± 21
$\Sigma_{1/2}''$...	2270 ± 26	2348 ± 17	2398 ± 32	$\Sigma''_{7/2}$	2033 ± 8	2428 ± 22	2418 ± 15	2403 ± 21
$\Xi''_{1/2}$...	2293 ± 35	2391 ± 13	2398 ± 32	$\Xi''_{7/2}$...	2494 ± 22	2455 ± 13	2403 ± 21
$\Omega_{1/2}$...	2378 ± 42	2426 ± 13	2398 ± 32	$\Omega_{7/2}$...	2553 ± 22	2477 ± 13	2403 ± 21

TABLE XXII. $[70, 1^-]$ masses. The experimental values are those for baryons with a three-star or higher rating by the PDG.

Baryon	PDG	M_π [MeV]			Baryon	PDG	M_π [MeV]		
		391	524	702			391	524	702
$N_{1/2}$	1538 ± 18	1681 ± 51	1797 ± 32	1968 ± 8	$N_{5/2}$	1678 ± 8	2012 ± 26	2033 ± 20	2109 ± 11
$\Lambda_{1/2}$	1670 ± 10	1777 ± 32	1852 ± 27	1968 ± 8	$\Lambda_{5/2}$	1820 ± 10	2057 ± 19	2068 ± 12	2109 ± 11
$\Sigma_{1/2}$...	1783 ± 25	1852 ± 27	1968 ± 8	$\Sigma_{5/2}$	1775 ± 5	2059 ± 21	2066 ± 15	2109 ± 11
$\Xi_{1/2}$...	1846 ± 32	1899 ± 32	1968 ± 8	$\Xi_{5/2}$...	2127 ± 21	2105 ± 15	2109 ± 11
$N_{3/2}$	1523 ± 8	1820 ± 40	1896 ± 17	2000 ± 8	$\Delta_{1/2}$	1645 ± 30	1885 ± 40	1964 ± 42	2023 ± 60
$\Lambda_{3/2}$	1690 ± 5	1904 ± 25	1939 ± 17	2000 ± 8	Σ''_{12}	...	1952 ± 25	1998 ± 37	2023 ± 60
$\Sigma_{3/2}$	1675 ± 10	1905 ± 23	1940 ± 20	2000 ± 8	$\Xi''_{1/2}$...	1987 ± 27	2038 ± 17	2023 ± 60
$\Xi_{3/2}$	1823 ± 5	1974 ± 25	1976 ± 17	2000 ± 8	$\Omega_{1/2}$...	2011 ± 41	2060 ± 20	2023 ± 60
$N'_{1/2}$	1660 ± 20	1892 ± 35	1928 ± 37	2045 ± 11	$\Delta_{3/2}$	1720 ± 50	1955 ± 32	2033 ± 17	2098 ± 11
$\Lambda'_{1/2}$	1785 ± 65	1849 ± 36	1944 ± 37	2045 ± 11	$\Sigma''_{3/2}$...	1958 ± 36	2071 ± 15	2098 ± 11
$\Sigma'_{1/2}$	1765 ± 35	1840 ± 36	1941 ± 37	2045 ± 11	$\Xi''_{3/2}$...	2040 ± 31	2108 ± 15	2098 ± 11
$\Xi'_{1/2}$...	1876 ± 27	2001 ± 22	2045 ± 11	$\Omega_{3/2}$...	2101 ± 30	2139 ± 15	2098 ± 11
$N'_{3/2}$	1700 ± 50	1895 ± 29	1935 ± 37	2077 ± 10	$\Lambda''_{1/2}$	1407 ± 4	1710 ± 32	1796 ± 20	1922 ± 11
$\Lambda'_{3/2}$...	1936 ± 30	1981 ± 27	2077 ± 10	$\Lambda''_{3/2}$	1520 ± 1	1817 ± 21	1816 ± 40	1903 ± 11
$\Sigma'_{3/2}$...	1951 ± 27	1977 ± 25	2077 ± 10					
$\Xi'_{3/2}$...	1998 ± 31	2030 ± 27	2077 ± 10					

- [1] A. Walker-Loud *et al.*, *Phys. Rev. D* **79**, 054502 (2009).
[2] J. M. Bulava *et al.*, *Phys. Rev. D* **79**, 034505 (2009).
[3] R. G. Edwards, J. J. Dudek, D. G. Richards, and S. J. Wallace, *Phys. Rev. D* **84**, 074508 (2011).
[4] R. G. Edwards, N. Mathur, D. G. Richards, and S. J. Wallace (Hadron Spectrum Collaboration), *Phys. Rev. D* **87**, 054506 (2013).
[5] H.-W. Lin *et al.* (Hadron Spectrum Collaboration), *Phys. Rev. D* **79**, 034502 (2009).
[6] J. J. Dudek, R. G. Edwards, and C. E. Thomas, *Phys. Rev. D* **87**, 034505 (2013) and references therein.
[7] V. Bernard, U.-G. Meissner, and A. Rusetsky, *Nucl. Phys.* **B788**, 1 (2008).
[8] G. P. Engel, C. B. Lang, D. Mohler, and A. Schäfer (BGR [Bern-Graz-Regensburg] Collaboration), *Phys. Rev. D* **87**, 074504 (2013).
[9] C. Alexandrou, T. Korzec, G. Koutsou, and T. Leontiou, *Phys. Rev. D* **89**, 034502 (2014).
[10] M. S. Mahbub, W. Kamleh, D. B. Leinweber, P. J. Moran, and A. G. Williams, *Phys. Rev. D* **87**, 094506 (2013).
[11] M. S. Mahbub, W. Kamleh, D. B. Leinweber, P. J. Moran, and A. G. Williams (CSSM Lattice Collaboration), *Phys. Rev. D* **87**, 011501 (2013).
[12] N. Matagne and F. Stancu, [arXiv:1406.1791](https://arxiv.org/abs/1406.1791) [Rev. Mod. Phys. (to be published)] and references therein.
[13] J.-L. Gervais and B. Sakita, *Phys. Rev. Lett.* **52**, 87 (1984).
[14] J.-L. Gervais and B. Sakita, *Phys. Rev. D* **30**, 1795 (1984).
[15] R. F. Dashen and A. V. Manohar, *Phys. Lett. B* **315**, 425 (1993).
[16] J. L. Goity, *Phys. Lett. B* **414**, 140 (1997).
[17] D. Pirjol and T.-M. Yan, *Phys. Rev. D* **57**, 5434 (1998).
[18] C. E. Carlson, C. D. Carone, J. L. Goity, and R. F. Lebed, *Phys. Lett. B* **438**, 327 (1998).
[19] C. E. Carlson, C. D. Carone, J. L. Goity, and R. F. Lebed, *Phys. Rev. D* **59**, 114008 (1999).
[20] C. L. Schat, J. L. Goity, and N. N. Scoccola, *Phys. Rev. Lett.* **88**, 102002 (2002).
[21] J. L. Goity, C. L. Schat, and N. N. Scoccola, *Phys. Rev. D* **66**, 114014 (2002).
[22] J. L. Goity, C. L. Schat, and N. N. Scoccola, *Phys. Lett. B* **564**, 83 (2003).
[23] D. Pirjol and C. Schat, *Phys. Rev. D* **67**, 096009 (2003).
[24] E. Gonzalez de Urreta, J. L. Goity, and N. N. Scoccola, *Phys. Rev. D* **89**, 034024 (2014).
[25] N. Matagne and F. Stancu, *Phys. Rev. D* **71**, 014010 (2005).
[26] N. Matagne and F. Stancu, *Phys. Lett. B* **631**, 7 (2005).
[27] N. Matagne and F. Stancu, *Phys. Rev. D* **74**, 034014 (2006).
[28] N. Matagne and F. Stancu, *Phys. Rev. D* **85**, 116003 (2012).
[29] J. L. Goity, *Phys. At. Nucl.* **68**, 624 (2005).
[30] J. L. Goity, in *Large N_c QCD 2004: Proceedings of the Workshop, Trento, Italy, 2004*, edited by J. L. Goity, R. F. Lebed, A. Pich, C. L. Schat, and N. N. Scoccola (World Scientific, Singapore, 2005), p. 211.
[31] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012).
[32] G. 't Hooft, *Nucl. Phys.* **B72**, 461 (1974).
[33] R. F. Dashen, E. E. Jenkins, and A. V. Manohar, *Phys. Rev. D* **51**, 3697 (1995).
[34] R. F. Dashen, E. E. Jenkins, and A. V. Manohar, *Phys. Rev. D* **49**, 4713 (1994).

- [35] J. L. Goity, C. L. Schat, and N. N. Scoccola, *Phys. Lett. B* **564**, 83 (2003).
- [36] E. E. Jenkins, A. V. Manohar, J. W. Negele, and A. Walker-Loud, *Phys. Rev. D* **81**, 014502 (2010).
- [37] A. Calle Cordon, T. DeGrand, and J. L. Goity, *Phys. Rev. D* **90**, 014505 (2014).
- [38] C. E. Carlson and C. D. Carone, *Phys. Lett. B* **484**, 260 (2000).
- [39] N. Isgur and G. Karl, *Phys. Rev. D* **18**, 4187 (1978).
- [40] C. Jayalath, J. L. Goity, E. Gonzalez de Urreta, and N. N. Scoccola, *Phys. Rev. D* **84**, 074012 (2011).