

**Fermion masses and mixing in general warped extra dimensional models**Mariana Frank,<sup>1,\*</sup> Cherif Hamzaoui,<sup>2,†</sup> Nima Pourtolami,<sup>1,‡</sup> and Manuel Toharia<sup>1,§</sup><sup>1</sup>*Department of Physics, Concordia University 7141 Sherbrooke Street West, Montreal, Quebec H4B 1R6, Canada*<sup>2</sup>*Groupe de Physique Théorique des Particules, Département des Sciences de la Terre et de L'Atmosphère, Université du Québec à Montréal, Case Postale 8888, Succursale Centre-Ville, Montréal, Québec H3C 3P8, Canada*

(Received 17 April 2015; published 3 June 2015)

We analyze fermion masses and mixing in a general warped extra dimensional model, where all the Standard Model (SM) fields, including the Higgs, are allowed to propagate in the bulk. In this context, a slightly broken flavor symmetry imposed universally on all fermion fields, without distinction, can generate the full flavor structure of the SM, including quarks, charged leptons and neutrinos. For quarks and charged leptons, the exponential sensitivity of their wave functions to small flavor breaking effects yield hierarchical masses and mixing as it is usual in warped models with fermions in the bulk. In the neutrino sector, the exponential wave-function factors can be flavor blind and thus insensitive to the small flavor symmetry breaking effects, directly linking their masses and mixing angles to the flavor symmetric structure of the five-dimensional neutrino Yukawa couplings. The Higgs must be localized in the bulk and the model is more successful in generalized warped scenarios where the metric background solution is different than five-dimensional anti-de Sitter (AdS<sub>5</sub>). We study these features in two simple frameworks, flavor complementarity and flavor democracy, which provide specific predictions and correlations between quarks and leptons, testable as more precise data in the neutrino sector becomes available.

DOI: [10.1103/PhysRevD.91.116001](https://doi.org/10.1103/PhysRevD.91.116001)

PACS numbers: 11.10.Kk, 12.15.Ff, 14.60.Pq

**I. INTRODUCTION**

The discovery of a SM-like Higgs boson at the LHC with a mass of 125 GeV was a huge step forward in confirming the validity of the Standard Model (SM) and probing the electroweak symmetry breaking mechanism. But despite its experimental success, the SM still fails to provide an explanation, among other things, for the origin and observed pattern of fermion masses and mixings.

In the quark sector, the masses are extremely hierarchical, with the top much heavier than the rest of the quarks and with a strong ordering. In the up sector, the masses are separated by 3 orders of magnitude, while in the down sector the mass ratios are separated by 2 orders of magnitude. The quarks also exhibit mixing patterns given by three small mixing angles and a large ( $CP$ ) phase. In the lepton sector, the charged lepton masses obey a similar hierarchical pattern as the down-type quarks. On the other hand, though not known exactly, neutrino masses are known to be very small and their square mass differences imply a closer mass pattern,  $\Delta m_{31(32)}^2 : \Delta m_{21}^2 \sim 10^2 : 1$ . Neutrinos appear to mix maximally and this has been long seen as pointing towards a different flavor origin between quarks and leptons, and also to the necessity of introducing

new physics. After the successful measurement of the neutrino mixing angle  $\theta_{13}$  by the Daya Bay [1,2], T2K [3,4], MINOS [5,6], RENO [7] and Double Chooz [8,9] collaborations, the determination of the neutrino mass hierarchy has become a priority for theoretical studies and for future neutrino experiments. A great deal of theoretical work in this area has been trying to provide answers, based on such diverse frameworks as seesaw mechanisms [10–16], Abelian [17–19] and non-Abelian [20–37] symmetries imposed on the leptonic sector (both charged and neutral), and many texture structures for leptonic mass matrices, including modifications of accepted paradigms, such as tribimaximal [38–43], bimaximal [44–49] and democratic [50–52] neutrino mixing matrices. While various attempts to unify the description of quarks and leptons already exist (mostly based on quark-lepton complementarity [53–66]), an attractive possibility would be that quarks and leptons obey the same symmetry at a higher scale, which is then slightly broken at lower scales, yielding different patterns for masses and mixing for the quarks/leptons than for the neutrinos. This is the scenario we plan to investigate here, in the context of warped extra dimensions, where small symmetry breaking terms have very different effects on quarks and leptons due to the geometry of the model.

Introducing a warped extra dimension provides an elegant way to address both the hierarchy problem (to stabilize the Higgs mass against large radiative corrections) and the fermion flavor hierarchy problem. These models

\* [mariana.frank@concordia.ca](mailto:mariana.frank@concordia.ca)† [hamzaoui.cherif@uqam.ca](mailto:hamzaoui.cherif@uqam.ca)‡ [n\\_pour@live.concordia.ca](mailto:n_pour@live.concordia.ca)§ [mtoharia@physics.concordia.ca](mailto:mtoharia@physics.concordia.ca)

were first proposed to deal with the hierarchy problem of the SM [67,68], where introducing an extra dimension produced a five-dimensional anti-de Sitter ( $\text{AdS}_5$ ) geometry bounded by two hard walls (branes) along the extra dimension, referred to as the Planck (or UV) brane and the TeV (or IR) brane. Allowing for exponential modulation from the gravity scale down to the weak scale along this compact extra dimension [67,68] yields the weak-Planck mass hierarchy.

The original Randall-Sundrum (RS) model localized all SM fields on the IR brane, leading to severe flavor violation bounds on the new physics scale. It was later shown that if the fermions were allowed to propagate in the bulk of the extra dimension [69–74], the same model could address the flavor hierarchy problem of the SM as well. The model thus emerged as a geometric theory of flavor. By localizing the Higgs on the IR brane with anarchic order-one couplings to the bulk fermions, the profiles of the fermion zero-modes can be adjusted to reproduce the observed Yukawa couplings in the low energy theory. Since the first and second generation fermions are localized towards the UV brane, they inherit substantial flavor protection from the RS–Glashow-Iliopoulos-Maiani mechanism [75]. However, by allowing the SM fields to propagate in the bulk, from an effective four-dimensional point of view, a tower of Kaluza-Klein (KK) fermions exists for all the SM fields, yielding enhanced contributions to electroweak and flavor observables in the SM [73,75–82]. This effect imposes a stringent bound on the scale of new physics of some  $\sim 10$  TeV [83,84] and hence renders these models completely out of the reach of current experiments. Some solutions were proposed to relieve these restrictions. One way was to extend the gauge symmetry of the model by introducing an  $SU(2)_R$  gauge sector with custodial protection [73,76,85]. Here the basic idea is to align the down-type Yukawa couplings using the additional symmetry, so that the primary sources of intergenerational mixing are the up-type Yukawa couplings. Since the dominant constraints on flavor-changing neutral currents come from the down-type sector, the constraint on KK masses is substantially relaxed. A different approach to address the issue is through a slight modification of the warping factor along the extra dimension, allowing it to deviate slightly from the  $\text{AdS}_5$  metric [86–91]. This deviation is such that the warping is more drastic near the TeV brane, while the background becomes more  $\text{AdS}_5$ -like near the Planck brane. These types of metric solutions can help suppress dangerous contributions to the electroweak and flavor observables by reducing the constraints on new physics down to about  $\sim 1$  TeV.

Warped extra dimensional models were shown to provide new contributions to the Higgs production rate through gluon fusion which could conflict with the collider data [74,92–100].

Interestingly, in the modified five-dimensional (5D) metric scenarios the region of parameter space in which

the dangerous contributions to flavor and electroweak precision observables are small is the same as the region where the new contributions to the Higgs production cross section are also small (and thus safe) [94]. Moreover, this is achievable only when the Higgs field in these models is allowed to leak considerably into the bulk.

In the context of warped extra dimensional models with  $\mathcal{O}(1)$  5D Yukawa couplings and with no *a priori* structure (i.e., flavor anarchy), one can easily generate the hierarchical structure of the quark and charged lepton sectors [101] while, due to large mixing angles, the neutrino sector must be treated differently. In particular, in [102], it was shown that if the Higgs field leaks sufficiently into the bulk it is possible that the (exponentially small) neutrino wave functions become independent on the flavor structure of the 5D neutrino mass parameters ( $c'_\nu$ ), and thus the 4D neutrino flavor structure depends directly on the flavor structure of the 5D neutrino Yukawa couplings.

In a previous work [103], we proposed a unified picture of fermion masses and mixings in the context of a warped extra dimensional model with  $\text{AdS}_5$  background metric, and with all the SM fields in the bulk, including the Higgs. In that picture, the same flavor symmetric structure is imposed on all the fermions of the SM, including neutrinos. Small flavor breaking effects are exponentially enhanced in the quark and charged lepton sectors, thus producing hierarchical masses and mixings. With a sufficiently delocalized Higgs field, the neutrino wave functions are flavor blind and the flavor structure is governed by the 5D neutrino Yukawa flavor structure. Previous analyses on masses in warped space are available in the literature [104–121].

In this work, we revisit this idea in the context of the modified  $\text{AdS}_5$  metric solutions. We show that the SM masses and mixing can be generated successfully, and the mass generation in the neutrino sector appears to be less fine tuned than in the case where a pure  $\text{AdS}_5$  background metric is assumed.

Our work is organized as follows. We summarize the features of the modified  $\text{AdS}_5$  model in Sec. II, with particular emphasis on fermion mass generation, Sec. III. We explore an explicit implementation of the scenario, flavor complementarity, in Sec. IV and another, of a democratic flavor symmetry, in Sec. V. We summarize our results and conclude in Sec. VI. We leave the details of some calculations for the Appendix.

## II. WARPED SPACE MODELS WITH FIELDS IN THE BULK

We consider a 5D warped space with the extra dimension compactified and allow all SM fields to propagate in the following generalized warped space-time metric:

$$ds^2 = e^{-2A(y)} \eta_{\mu\nu} dx^\mu dx^\nu - dy^2, \quad (2.1)$$

$\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$  being the flat metric. The fifth dimension,  $y$ , is bounded by two branes localized at  $y = 0$  and  $y = y_1$  and  $A(y)$  is a model-dependent function. As mentioned in the Introduction, some generalized warped models can be safe from precision electroweak tests and flavor bounds for very low KK masses. Motivated by this, we consider such a model within a modified AdS<sub>5</sub> scenario with the following warp exponent [90,122,123]:

$$A(y) = ky + \frac{1}{\nu^2} \ln \left( 1 - \frac{y}{y_s} \right), \quad (2.2)$$

where  $k \sim M_{Pl}$  is the AdS<sub>5</sub> curvature, expected to be of the order of the Planck mass scale,  $y_s$  is the position of the metric singularity, always chosen to be outside of the physical region,  $y_s > y_1$ , and  $\nu > 0$  is a model parameter taken to be real. The  $\nu$  parameter, alongside  $\Delta = y_s - y_1$ , the distance between the location of the metric singularity and the IR brane, measures the departure of the metric from the pure AdS<sub>5</sub> background. The smaller the values of  $\Delta$  and  $\nu$ , the more modified the metric; intuitively, the singularity has a larger effect on the physics at the IR brane the closer it gets to it. One can calculate the curvature along the fifth dimension and obtain

$$R(y) = 8A''(y) - 20(A'(y))^2. \quad (2.3)$$

The curvature radius,  $L(y) = \sqrt{-20/R}$ , in units of  $k$  along the fifth dimension is then given by

$$kL(y) = \frac{k\Delta}{\sqrt{1 - 2\nu^2/5 + 2\nu^2k\Delta + (\nu^2k\Delta)^2}}. \quad (2.4)$$

One can see that for values of  $\nu > \sqrt{5/2}$ , this function has a minimum before the singularity and therefore the curvature can change sign within the physical region. Following [91], we impose that this minimum is located outside of the physical region and hence the curvature radius is a monotonically decreasing function between the UV and the IR branes.

The more familiar RS metric is recovered by taking the limits  $\nu \rightarrow \infty$  and  $y_s \rightarrow \infty$ , yielding  $A(y) = ky$ , with the curvature radius being constant,  $kL = 1$ . In Fig. 1, we compare the two metrics and plot the warp exponent function  $A(y)$  for the AdS<sub>5</sub> and the modified AdS<sub>5</sub> cases. We can see that the amount of warping near the IR brane at around  $ky = 35$  is larger for the modified AdS<sub>5</sub>. Thus, as the figure indicates, the same amount of warping from the UV brane to the IR brane in the modified scenario requires a slightly smaller length of the fifth dimension and hence an IR brane slightly closer to the UV brane. The curvature radius [Eq. (2.4)] at the UV brane is approximately equal for the pure and modified AdS<sub>5</sub> spaces with,  $ky(y) \approx 1$ . On the contrary, at the IR brane, as  $kL(y)$  is a monotonically decreasing function,  $ky(y)$  assumes its minimal value for the modified AdS<sub>5</sub> space and hence  $kL_1 \equiv kL(y_1)$  is a




FIG. 1 (color online). The modified AdS<sub>5</sub> warp factor  $A(y)$  versus the standard RS warp exponent,  $y$ . The horizontal line corresponds to  $ky = 35$ . For the same amount of warping, the modified scenario requires a shorter length scale along the fifth dimension.

good measure of the amount of deviation from the pure AdS<sub>5</sub> space with constant curvature radius.

The 5D fermion Lagrangian density with Dirac neutrinos is

$$\begin{aligned}
 \mathcal{L}_q = & \mathcal{L}_{\text{kinetic}} + M_{q_i} \bar{Q}_i Q_i + M_{u_i} \bar{U}_i U_i + M_{d_i} \bar{D}_i D_i \\
 & + (Y_{ij}^{u5D} H \bar{Q}_i U_j + \text{H.c.}) + (Y_{ij}^{d5D} H \bar{Q}_i D_j + \text{H.c.}) \\
 & + (Q_i \rightarrow L_i, U_i \rightarrow N_i, D_i \rightarrow E_i), \quad (2.5)
 \end{aligned}$$

where,  $i, j$  are flavor indices and the 5D Yukawa parameters,  $Y_{ij}^{5D}$ , are dimensionful quantities of  $\mathcal{O}(1) \times \sqrt{k}$ .  $Q_i$  ( $L_i$ ) are 5D quark (lepton) fields for  $SU(2)$  doublets while  $U_i$  ( $N_i$ ) and  $D_i$  ( $E_i$ ) are  $SU(2)$  singlet quark (lepton) fields. The bulk mass,  $M_{\psi_i}$ , originating from the momentum along the fifth dimension, can be taken in general to be  $y$  dependent. To be able to compare, we choose it such that it coincides with its usual definition in RS models, and express it in units of the fifth dimension curvature,  $k$ , as  $M_{\psi_i} = c_{\psi}^i k$ , where  $c_{\psi}^i$  are localization parameters, dimensionless quantities of  $\mathcal{O}(1)$ , and  $\psi_i$  runs over all SM quarks and leptons.<sup>2</sup> Dimensional reduction then yields the normalized profile for the fermion and the Higgs fields along the bulk of the extra dimension,  $q_L^{0,i}(y)$ ,  $u_R^{0,i}(y)$  and  $h(y)$ , which are given by

$$q_L^{0,i}(y) = q_0^i e^{(2-c_q^i)A(y)}, \quad (2.6)$$

<sup>1</sup>We use throughout  $c_q$  for the doublets ( $c_q$  and  $c_l$ ) and  $c_u$  for the singlets ( $c_u$ ,  $c_d$ ,  $c_\nu$  and  $c_e$ ).

<sup>2</sup>Alternative fermion and Higgs profiles can be found in [94,122], where different bulk mass conventions are adopted.

$$u_R^{0,i}(y) = u_0^i e^{(c_u^i+2)A(y)}, \quad (2.7)$$

$$h(y) = h_0 e^{aky}, \quad (2.8)$$

with  $q_0^i = f(c_q^i)$ ,  $u_0^i = f(-c_u^i)$  and  $h_0 = e^{-(a-1)ky_s} \mathfrak{h}_0$ , and where  $f(c)$  and  $\mathfrak{h}_0$  are normalization factors which depend on  $c$ ,  $\nu$  and  $y_s$ , given explicitly in the Appendix, along with their limiting expressions for the usual RS (AdS<sub>5</sub>) metric background. From these profiles, one can check that localization of the fields in the bulk of the extra dimension is determined by the values of the  $c_\psi^i$  for the fermion fields, such that a value of  $c_\psi^i > 1/2$  indicates an UV localized field, while a value of  $c_\psi^i < 1/2$  localizes the field near the IR brane.<sup>3</sup> The Higgs field localization along the fifth dimension is given by the parameter  $a$ , the dimension of the Higgs condensate operator. A completely IR brane localized field corresponds to the limit  $a \rightarrow \infty$ , while for a delocalized Higgs field  $a$  is small. However, in order to maintain the original Randall-Sundrum solution to the hierarchy problem without fine-tuning, the Higgs field localization should be such that  $a \gtrsim 2$  (for an AdS<sub>5</sub> metric background). If the 5D Higgs potential is of the form  $V_{\text{bulk}}(H) = M_{5d}^2 H^2$ , with associated brane potentials at each boundary, the Higgs profile has two solutions, one growing towards the IR and another one decaying at the IR brane. This last one is proportional to  $e^{(4-a)ky}$  in the AdS<sub>5</sub> background. In order to maintain the RS solution to the hierarchy problem, the decaying solution has to be subdominant, and this happens for  $a > 2$ . In that situation, the Higgs field profile [see Eq. (2.10)] is localized near the IR brane and mass terms depending on the Higgs profile vacuum expectation value are warped down to the energy scale of the IR brane as

$$m_{\text{Higgs}} \simeq M_{5d} e^{-A(y_1)}. \quad (2.9)$$

This resolution of the hierarchy problem depends on the actual length of the extra dimension, as well as the choice of the 5D metric parameters,  $\nu$  and  $y_s$ , which are usually set such that  $A(y_1) \simeq 35$ .

When  $a < 2$ , and in order to maintain a Higgs profile localization towards the IR brane, some fine-tuning between the parameters of the bulk scalar potential and the brane potentials is necessary in order to suppress the unwanted solution. In the modified AdS<sub>5</sub> scenario the lowest value of  $a$  that does not require fine-tuning depends on the various new metric parameters [82]. In this case, the Higgs profile is given by

$$h(y) = h_0 e^{aky} [1 + (M_0/k - a)[F(y) - F(0)]], \quad (2.10)$$

<sup>3</sup>This convention is for left-handed doublets. For right-handed singlet fields, our convention is such that  $c_u^i > -1/2$  for an UV localized field and  $c_u^i < -1/2$  for an IR localized field.

where  $M_0$  is the brane Higgs mass term [coefficient of the  $|H|^2 \delta(y - y_1)$  term at the IR brane] and the function  $F(y)$  is given by

$$F(y) = e^{-2(a-2)ky_s} k y_s [-2(a-2)k y_s]^{-1+4/\nu^2} \times \Gamma \left[ 1 - \frac{4}{\nu^2}, -2(a-2)k(y_s - y) \right]. \quad (2.11)$$

The decaying term at the IR brane is the second term in Eq. (2.10) and, forcing  $M_0/k \simeq a$  (fine-tuning parameters), the solution can become subdominant. In order to avoid this fine-tuning of parameters, we note that  $F(y)$  is a monotonically increasing function. We can thus introduce  $\delta \equiv |F(y_1)|$  as a measurement of the amount of fine-tuning needed for the RS solution to survive the hierarchy problem. When  $\delta \sim \mathcal{O}(1)$ , no fine-tuning is needed to guarantee that the increasing solution for the Higgs profile dominates. When the parameter  $\delta = F(y_1)$  becomes larger, this solution needs fine-tuning of parameters to suppress its value.

As expressed in the Introduction, scenarios of generalized warped metric backgrounds can yield lower bounds on KK masses coming from precision and flavor constraints. The precision electroweak oblique parameters,  $S$  and  $T$  are proportional to the dimension-6 operators

$$|H^\dagger D_\mu h|^2, \quad H^\dagger W_{\mu\nu} H B^{\mu\nu}.$$

Due to the deviations of the metric from AdS<sub>5</sub> near the IR brane all the KK modes, including the gauge KK modes, become more localized near the IR brane. Intuitively, for a Higgs field that is sufficiently delocalized from the IR brane, couplings to these KK modes will be suppressed, which will result in lower bounds for the KK mode masses in these models (which can be as low as  $m_{KK} \gtrsim 1$  TeV). As the Higgs field becomes more localized towards the IR brane, these couplings become large (even larger than in the pure AdS<sub>5</sub> case) and the bounds can become much worse. Therefore the success of this model is crucially dependent on the localization of the Higgs field along the fifth dimension [86,124].

Figure 2 shows the no-fine-tuning region (above the red solid  $\delta = 1$  curve) in the  $(a, \nu)$  plane, where  $a$  is the Higgs localization parameter and  $\nu$  is the metric parameter of the modified metric solution. The region below is the one fine tuned, requiring an adjustment of Lagrangian parameters with a tuning precision growing exponentially. (The close dashed curve locates the points where  $\delta = 10$ , i.e., where the tuning is already 10%.)

In producing these graphs, for each case we first set the value of the IR brane position,  $ky_1$ , which in turn fixes the value of  $y_s$ , the position of the singularity. Then for each value of the parameter  $\nu$  we solved for  $a$  in  $\delta(a, \nu, y_1, y_s) = 1$ .

To first order, the effective SM Yukawa couplings are obtained from the overlap integral




FIG. 2 (color online). The  $\delta = 1$  plots and the connection with neutrino masses in the  $\nu - a$  plane for (left panel)  $kL_1 = 0.2$  (large deviation) and (right panel)  $kL_1 = 0.99$  (more RS-like). The red (solid) curve locates the no-fine-tuning threshold in which  $\delta = 1$ . Above this curve,  $\delta < 1$ , and below the curve,  $\delta > 1$  (corresponding to the unwanted fine-tuned region). The shaded area corresponds to a heavy neutrino mass ( $m_3$  in *normal* ordering),  $m_\nu \sim 5 \times 10^{-11}$  GeV for different values of the  $c$  parameters, but such that the mass expression has no exponential sensitivity to the  $c$  parameters. Note that in the RS-like case (right panel) it is not possible to obtain neutrino masses without fine-tuning, or quitting the neutrino plateau parameter region (as would be the case in usual RS scenarios). With modified-AdS<sub>5</sub> metric (left panel) it is possible to find nontuned points with neutrino masses in the plateau (corresponding to no exponential  $c$ -dependence).

$$y_{ij}^u = \frac{Y_{ij}^u}{\sqrt{k}} \int_0^{y_1} dy e^{-4A(y)} h(y) q_L^{0,i}(y) u_R^{0,j}(y), \quad (2.12)$$

where the index  $u$  denotes the four types of Yukawa couplings of the SM, i.e.,  $u = u, d, e, N$  and we have defined the *dimensionless* 5D Yukawa couplings as  $Y_{ij}^u = Y_{ij}^{5D}/\sqrt{k} \sim \mathcal{O}(1)$ . Given the profiles (2.6), (2.7) and (2.8) and the metric (2.2), the integral above can be performed analytically and written as

$$y_{ij}^u = \tilde{Y}_{ij}^u h_0 f(c_q^i) f(-c_u^j), \quad (2.13)$$

where the factor  $\tilde{Y}_{ij}^u$ , defined in the Appendix, has very mild  $c^i$  dependence. The function  $f(c)$  is such that depending on the value of  $c$ , the Yukawa couplings can be exponentially sensitive to  $c$ , or only mildly dependent. Finally  $h_0$  is the bulk Higgs normalization factor and does not depend on the fermion mass parameters  $c^i$ . Note that, although throughout the paper we have suppressed the explicit dependence of the fields on the metric parameters,  $\nu, y_s, y_1$  and  $k$ , and in particular all of the factors of Eq. (2.13), are metric dependent. As shown in the Appendix, one can always retrieve the RS limit for these terms by taking the limit  $\nu, y_s \rightarrow \infty$ .

### III. FERMION MASSES IN WARPED SPACE

We now return to a more detailed discussion of fermion masses in this model. These masses are given, to first order, by the eigenvalues of the  $3 \times 3$  Yukawa matrices of the form

$$v y^u = v \begin{pmatrix} y_{11}^u & y_{12}^u & y_{13}^u \\ y_{21}^u & y_{22}^u & y_{23}^u \\ y_{31}^u & y_{32}^u & y_{33}^u \end{pmatrix}. \quad (3.1)$$

In general, in order to get the correct SM masses, no exponential  $c$ -dependence is needed for the top quark, which corresponds to  $c_q^t \lesssim 1/2$  and  $c_u^t > -1/2$  (this region of parameter space corresponds to the top plateau shown in Fig. 3). For the rest of the SM particles  $c_q^i \gtrsim 1/2$  and  $c_u^i < -1/2$ , which implies that the Yukawa couplings will depend exponentially on the values of the  $c^i$ . In the case of neutrinos however, in order to accommodate their tiny masses, one needs localization parameters  $c_\nu < -1$ . It was shown [102] that, for a delocalized Higgs with  $a$  parameter small compared to the localization parameter  $c^i$ , the 4D effective neutrino masses depend exponentially on  $a$  but lose their dependence on the  $c^i$ 's. This region of parameter space corresponds to the neutrino plateau shown in Fig. 3. In other words, the limit of the function in Eq. (2.13) for large  $c_\nu$ 's is given by

$$y_{ij}^u \sim \tilde{Y}_{ij}^u h_0. \quad (3.2)$$

To make this more transparent, in the formula for fermion masses given by [cf. Eq. (2.13)]

$$(M_f)_{ij} = v \tilde{Y}_{ij}^u h_0 f(c_q^i) f(-c_u^j), \quad (3.3)$$

we factor the exponential behaviors and write them in the following two distinct limits:




FIG. 3 (color online). Effective 4D Yukawa couplings for fermions as functions of the fermion bulk mass parameter  $c$  for (a) the RS and (b) general warped space-time metric. The plots on the right-hand side are produced by taking the  $c$  parameters for the doublet and the singlet to be equal. In the left-handed plots, each contour depicts 1 order of magnitude difference with respect to its adjacent contour. A typical location for neutrino masses of order  $m_\nu \sim 10^{-11}$  GeV is also shown. The shaded vertical band shows the region where all fermions of the SM should be located. (a) RS with all SM elds including the Higgs in the bulk,  $a = 2.04$  and  $\delta = 1$  and (b) Modified AdS5 with  $\nu = 0.32$ ,  $kL(y_1) = 0.2$ ,  $A(y_1) = 35$ ,  $a = 4.46$  and  $\delta = 1$ .

$$\begin{aligned}
 (M_f)_{ij} &\sim v Y_{ij}^f \epsilon^{(c_q - \frac{1}{2})} \epsilon^{-(c_u + \frac{1}{2})} \quad \text{for } c_q > 1/2, \quad c_u < -1/2, \\
 (M_\nu)_{ij} &\sim v Y_{ij}^N \epsilon^{-ky_1(a-1)} \quad \text{for } c_q - c_u > a,
 \end{aligned}
 \quad (3.4)$$

where  $\epsilon = e^{-A(y)}$  (see the Appendix for details).<sup>4</sup>

The limits described in Eq. (3.4) imply that, as the quark and charged lepton mass matrix elements are exponentially dependent on the  $c^i$  parameters, any structure in the 5D Yukawa matrix elements  $Y_{ij}^f$  will

<sup>4</sup>For modified AdS<sub>5</sub>,  $\epsilon_{\text{mod}} = e^{-ky_1} (1 - \frac{y_1}{y_s})^{\frac{1}{2}} \sim 10^{-15}$  while for RS  $\epsilon_{\text{RS}} = e^{-ky_1} \sim 10^{-15}$ .

be largely washed out and will always produce generically hierarchical fermion masses as well as small mixing angles. For the neutrinos, on the other hand, since there is no exponential sensitivity on the flavorful  $c^i$  parameters, any structure inherent in the 5D Yukawa matrix elements will survive in the 4D effective theory. This is the region of neutrino parameter space we are interested in, shown in Fig. 3 as the neutrino plateau. In RS models, the actual height of the neutrino plateau is determined exclusively by the value of the  $a$  parameter. For the value of the warp factor required to solve the hierarchy problem, the highest possible neutrino masses in the plateau are too small by 1 to 2 orders of magnitude




FIG. 4 (color online). (Left panel) Neutrino masses as functions of the metric parameter  $\nu$  for different values of the Higgs localization parameter  $a$  and fixed  $kL_1 = 0.2$ . The fermion mass parameters ( $c$ 's) are fixed to a region where there is no exponential dependence on them (the neutrino plateau). The curves end whenever the fine-tuning threshold ( $\delta = 1$ ) is reached (thick overlapping curve). Note that for small values of  $\nu$  the neutrino masses become larger while still remaining in the nontuned regime and in the neutrino plateau. (Right panel) Same plot for different values of  $kL_1$  and fixed  $\delta = 1$ . The graph corresponding to  $kL_1 = 0.9$  remains constant at larger values of  $\nu$  and (approximately) coincides with the RS limit.

(without fine-tuning) to be phenomenologically viable (see the upper panels in Fig. 3). Some tuning, or some enhancement of the 5D Yukawa couplings, and/or trespassing the edge of the plateau would then be required, making the scenario less attractive for our purposes. On the other hand, in modified AdS<sub>5</sub> scenarios, although the level of the plateau is still highly sensitive to the value of  $a$ , the masses could actually be increased by some 2 orders of magnitude and thus allow for phenomenologically acceptable neutrino masses in the desired region of the model (see Fig. 2 and the lower panels in Fig. 3). This feature occurs because the modified AdS<sub>5</sub> metric [see Eq. (2.2)] exhibits a richer parameter space. In the RS metric, in order to produce the correct neutrino masses in the plateau, we need  $a \sim 1.85$ , a value which amounts to about 1/10000 fine-tuning of parameters in the 5D Higgs potential. While for modified AdS<sub>5</sub> scenarios, one can produce a neutrino plateau within the experimental bounds for some range of  $a$  parameter values without fine-tuning. It is interesting that the parameter space for which the neutrino plateau is most favorable coincides with the region where small KK masses satisfy bounds coming from flavor and electroweak precision measurements [125,126] and constraints from Higgs phenomenology [93].

In order to further illustrate this issue, in Fig. 4 we show the resulting neutrino masses in the plateau region as a function of the parameter  $\nu$ , for different values of  $a$  (left panel) and  $kL_1$  (right panel). The area below the curves is the no-fine-tuning region, and one can see

that the largest neutrino masses (in the plateau) occur for  $\nu \sim 0.3$  and  $kL_1 = 0.2$ , whereas in the RS limit, i.e.,  $kL_1 \sim 1$  and  $\nu$  large, the masses are some 2 orders of magnitude lower (slightly too low). This makes the extended metric scenarios a framework for the flavor mechanism investigated here, and adds to the advantages of these scenarios (i.e., a much lower KK scale consistent with electroweak and flavor bounds, and with Higgs production).

Qualitatively the general features of the flavor structure of these modified AdS<sub>5</sub> models are very similar to these features in the pure AdS<sub>5</sub> models for the bulk mass  $c_{\psi}^i$  parameters. Therefore, in order to illustrate our flavor setup, it is useful to consider the simpler case of the RS metric. In this case the fermion mass formulas in Eq. (3.4) can be simplified dramatically. As usual, we consider the mass matrix for the neutrino sector separately from the case of quarks and charged lepton mass matrices. Consider the case with  $c_{q,u}^i > \frac{1}{2}$ . In this case we have,

$$(M_f)_{ij} \simeq v e^{(c_q^i - \frac{1}{2})} e^{-(c_u^j + \frac{1}{2})} \sqrt{2(a-1)|1-2c_q^i|1+2c_u^j} \tilde{Y}_{ij}^f$$

$$(M_\nu)_{ij} \simeq v e^{a-1} \frac{\sqrt{2(a-1)|1-2c_l^i|1+2c_\nu^j}}{\sqrt{e^{(1-2c_l^i)}-1}\sqrt{e^{(1+2c_\nu^j)}-1}} \tilde{Y}_{ij}^\nu, \quad (3.5)$$

where the 5D Yukawa couplings are given by<sup>5</sup>

<sup>5</sup>For exact formulas see the Appendix.

$$\tilde{Y}_{ij}^u \simeq \frac{1}{a - c_q^i + c_u^j} Y_{ij}^u. \quad (3.6)$$

From Eqs. (2.6), (2.7) and (2.12) one can see that there are two sources of flavor structure: the 5D dimensionless Yukawa couplings  $Y_{ij}^u$  and the bulk mass coefficients  $c_\psi^i$ . We are interested in scenarios in which all Yukawa matrices ( $Y_F = Y_{ij}^u, Y_{ij}^d, Y_{ij}^\nu$  and  $Y_{ij}^e$ ) and fermion bulk mass matrices from the 5D Lagrangian ( $\mathbf{c}_f = \mathbf{c}_q, \mathbf{c}_u, \mathbf{c}_d, \mathbf{c}_l, \mathbf{c}_\nu$  and  $\mathbf{c}_e$ ) share the same symmetry structure, which is then slightly broken through some high energy mechanism according to

$$Y_F = Y_F^0 + \delta Y_F, \quad (3.7)$$

$$\mathbf{c}_f = \mathbf{c}_f^0 + \delta \mathbf{c}_f, \quad (3.8)$$

where the matrices  $Y_F^0$  and  $\mathbf{c}_f^0$  are flavor symmetric while the perturbation matrices  $\delta Y_F$  and  $\delta \mathbf{c}_f$  are random. Inserting these perturbations in Eq. (3.5), the fermion masses receive corrections to leading order as follows:

$$\begin{aligned} m_t &= m_t^0 + \delta m_t c_q^3, & c_u^3 &< 1/2, \\ (m_f)_{ij} &= (m_f)_{ij}^0 f(\delta c_q^i) f(\delta c_u^j) \sim (m_f)_{ij}^0 \epsilon^{(\delta c_q^i + \delta c_u^j)} \\ & a > c_q^i + c_u^j, \\ (m_\nu)_{ij} &= (m_\nu)_{ij}^0 + \delta (m_\nu)_{ij} & a &< c_q^i + c_\nu^j. \end{aligned} \quad (3.9)$$

Therefore the same exponential sensitivity on the bulk mass  $c_\psi^i$  parameters,  $\epsilon \sim 10^{-15}$ , responsible for producing the SM hierarchy in the standard RS, is now translated into exponential sensitivity of the symmetry breaking terms. As a consequence, small symmetry breaking terms ( $|\delta c^i| \sim 0.1$ ) can produce mass corrections of order  $10^{-15(\delta c^i + \delta c^j)}$  (i.e., a hierarchy of order  $\sim 10^6$ ) to the quark and charged lepton mass matrices. This is in complete agreement with the observed hierarchy in these sectors. As mentioned before, the neutrinos and the top quark fields live in the two plateaus (see Fig. 3) with mild  $c_\psi^i$  sensitivity.

For the mixing angles, the eigenvector matrix that diagonalizes the neutrino sector should be very close to the eigenvector matrix of the 5D Yukawa matrix. However, in the quark and charged lepton sectors the mixing matrices are generically close to the unit matrix with off-diagonal entries hierarchically small as<sup>6</sup>

<sup>6</sup>Similar expressions are obtained for  $V_L^d$  and  $V_L^q$  (see for example [112]).

$$V_L^u \simeq \begin{pmatrix} 1 & \frac{f_q^1(\tilde{M}_u)_{21}}{f_q^2(\tilde{M}_u)_{11}} & \frac{f_q^1(\tilde{Y}_u)_{13}}{f_q^2(\tilde{Y}_u)_{33}} \\ -\frac{f_q^1(\tilde{M}_u)_{21}}{f_q^2(\tilde{M}_u)_{11}} & 1 & \frac{f_q^2(\tilde{Y}_u)_{23}}{f_q^3(\tilde{Y}_u)_{33}} \\ \frac{f_q^1(\tilde{M}_u)_{31}}{f_q^2(\tilde{M}_u)_{11}} & -\frac{f_q^2(\tilde{Y}_u)_{23}}{f_q^3(\tilde{Y}_u)_{33}} & 1 \end{pmatrix}, \quad (3.10)$$

where  $f_q^i$  is shorthand for the profile functions  $f(c_q^i)$ ,  $(\tilde{M}_u)_{ij}$  is the  $(ij)$  minor of the matrix in parenthesis, and  $(\tilde{Y}_u)_{ij}$  is the  $ij$  element of the Yukawa matrix. We define the Cabibbo-Kobayashi-Maskawa (CKM) and Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrices as the following:

$$V_{\text{CKM}} \equiv V_L^u V_L^{d\dagger} \quad \text{and} \quad V_{\text{PMNS}} \equiv V_L^\nu V_L^{e\dagger}. \quad (3.11)$$

As mentioned after Eq. (3.4), for the quarks and charged leptons, the off-diagonal mixing angles are also exponentially sensitive to the  $c_\psi^i$  parameters and hence to the small symmetry breaking terms. The  $V_L^f$  matrix elements in Eq. (3.10) can then be written as

$$(V_L^{u,d,e})_{ij} \sim (V_L^{u,d,e})_{ij}^0 \epsilon^{(\delta c_q^i - \delta c_q^j)} \quad (\text{for } u, d \text{ and } e), \quad (3.12)$$

where  $(V_L^{u,d,e})_{ij}^0$  are the matrix elements before the flavor symmetry breaking and  $\{i, j\}$  can be  $\{1, 2\}$ ,  $\{1, 3\}$  and  $\{2, 3\}$ . We see again that, due to exponential warping, all original symmetries present in the high energy theory are washed out in the quark and charged lepton sectors. Contrary to this, in the Dirac neutrino sector, the terms in the  $V_L^\nu$  are much less sensitive to the symmetry breaking terms, since their own dependence on the  $c_\psi^i$  is mild in the region of parameters of interest. If we define the eigenvalues of the 5D neutrino Yukawa matrix as

$$Y_\nu^{\text{diag}} = V_{Y_L} Y_\nu V_{Y_R}, \quad (3.13)$$

then the matrix diagonalizing the 4D effective neutrino mass matrix is

$$(V_L^\nu)_{ij} \sim (V_{Y_L})_{ij}. \quad (3.14)$$

So far the framework is general, and we did not specify the type of symmetry imposed. In the next two sections we will consider two concrete implementations, one within flavor complementarity and then a second within flavor democracy. The background metric considered will always be the modified AdS<sub>5</sub> solution, in the most favorable region of parameter space.

#### IV. FLAVOR COMPLEMENTARITY

In the first symmetry implementation, in this section we study the effects of enforcing a strong correlation between 5D quark and 5D lepton parameters. This scenario is



motivated by the observation that charged lepton masses and down-quark masses obey similar patterns. In warped space models, masses are obtained from exponentially small overlap integrals, and having similar mass patterns can indicate that both the 5D Yukawa structure and the 5D bulk fermion masses ( $c$  parameters) are very similar for the down quarks and for the charged leptons. We therefore assume that  $Y_d = Y_e$  and  $\mathbf{c}_q = \mathbf{c}_l$  to a high order of precision, so that the differences in observed masses between down quarks and charged leptons come from deviations in the right-handed bulk parameters  $\mathbf{c}_d$  and  $\mathbf{c}_e$ .

At the same time, inspired by neutrino phenomenology, we implement a popular 2-3 ( $\mu - \tau$ ) symmetry [22,24,127–148] within the general flavor structure of the model. We require that all the 5D Yukawa matrices must be symmetric under two to three permutations and consider, for simplicity, the case where the 5D fermion bulk mass matrices  $\mathbf{c}_f^0$  are diagonal and degenerate in that basis [maybe a remnant of some global  $U(3)$  flavor symmetry broken in the Yukawa sector], i.e.,

$$\mathbf{c}_f^0 = \begin{pmatrix} c_f^0 & 0 & 0 \\ 0 & c_f^0 & 0 \\ 0 & 0 & c_f^0 \end{pmatrix}, \quad (4.1)$$

where  $c_f^0$  are real parameters, and  $f$  denotes both doublets and singlets  $f = q, l, u, d, e, \nu$ . In 4D scenarios, discrete permutation symmetries of the neutrino mass matrix are known to lead to interesting (and phenomenology viable) neutrino masses and mixing patterns (such as the tribimaximal mixing matrix [38–43], or the bimaximal mixing matrix [44–49]).

We will impose the zeroth order Yukawa matrices to be

$$Y_{u,\nu}^0 = y_u \begin{pmatrix} \sqrt{6}d & 0 & 0 \\ d & d+1 & d-1 \\ d & d-1 & d+1 \end{pmatrix} \quad \text{and} \\ Y_{d,e}^0 = y_d \begin{pmatrix} a & 1 & 1 \\ c & g & e \\ c & e & g \end{pmatrix}, \quad (4.2)$$

where the parameters  $y_u, y_d, a, c, d, g$  and  $e$  are complex.<sup>7</sup> As explained above, we impose the down-type Yukawa couplings to match the charged lepton ones, i.e.,  $Y_d = Y_e \equiv Y_{d,e}$ . We further require that the up-type Yukawa couplings be the same as the neutrino Yukawa

couplings, i.e.,  $Y_u = Y_\nu \equiv Y_{u,\nu}$ , in order to reduce free parameters and enhance the degree of correlation between quarks and leptons.

The structure of the down-type Yukawa  $Y_{d,e}^0$  is set only by the 2-3 symmetry, but the structure of neutrino-type Yukawa couplings  $Y_{u,\nu}^0$  in Eq. (4.2) requires more explanation. The number of parameters has been reduced to two complex parameters,  $y_u$  and  $d$ , such that the rotation matrix that diagonalizes  $Y_\nu^0$  is the bimaximal mixing matrix<sup>8</sup> and the eigenvalues of  $Y_{u,\nu}^0$  are simply given by  $3.08d$ ,  $1.59d$  and  $2$  (in units of  $y_u$ ). When  $|d| \sim 1/6$ , we should naively generate a hierarchy between the eigenvalues of the matrix matching the solar and atmospheric neutrino mass hierarchies, for the case of a normal ordering in the neutrino masses. Of course, in warped scenarios, the 5D Yukawa matrices are in general not directly proportional to the effective 4D fermion mass matrices. In the neutrino sector however, since we are near the neutrino mass “plateau” of the  $c$  parameters, the structure of the effective 4D mass matrix should be very similar to the 5D Yukawa matrix. We therefore expect that the effective 4D neutrino structure should be diagonalized by a matrix close to the bimaximal mixing matrix, with a normal ordering in the masses matching the observed mass differences. 5D wave function effects, as well as random, but small, symmetry breaking terms, would perturb this structure, but the main features survive.

Finally, the two texture zeros of  $Y_u^0$  in Eq. (4.2) cause the vanishing of the (21) minor of that matrix, which in turn contributes to the quark-lepton correlations which we want to address here; the vanishing, or the smallness of those entries, is therefore a critical requirement in the setup, as it links the value of the Cabibbo angle to the PMNS element  $V_{13}$  (i.e., to  $\theta_{13}$ ). To see how this occurs, let us write the approximate expression for the Cabibbo angle  $V_{us}$  in warped extra dimensions,

$$V_{us} \simeq \left| \frac{f(c_q^1)}{f(c_q^2)} \right| \left| \frac{(\tilde{M}_d)_{21}}{(\tilde{M}_d)_{11}} - \frac{(\tilde{M}_u)_{21}}{(\tilde{M}_u)_{11}} \right|, \quad (4.3)$$

where  $(\tilde{M}_u)_{ij}$  and  $(\tilde{M}_d)_{ij}$  are the  $(ij)$  minors of the  $Y_u$  and  $Y_d$  matrices. We can see that the texture zeros in the 5D Yukawa matrix  $Y_u$  ensure that  $(\tilde{M}_u)_{12} = 0$ , so that the Cabibbo angle is controlled, to first order, by the down sector Yukawa couplings and by the quark doublet bulk  $c$  parameters, i.e.,

$$V_{us} \simeq \frac{f(c_q^1)}{f(c_q^2)} \frac{(\tilde{M}_d)_{21}}{(\tilde{M}_d)_{11}}. \quad (4.4)$$

<sup>7</sup>We also assume that the symmetry is broken by some small perturbations, i.e.,  $Y_i = Y_i^0 + \delta Y_i$  and  $\mathbf{c}_f = \mathbf{c}_f^0 + \delta \mathbf{c}_f$ , although we require that the constraints  $Y_d = Y_e$  and  $\mathbf{c}_q = \mathbf{c}_l$  survive the flavor symmetry breaking in order to maintain the quark lepton complementarity.

<sup>8</sup>The tribimaximal scheme predicts the  $CP$  phase in the PMNS matrix,  $\delta$ , to be zero, contrary to phenomenological conjectures which favor  $\delta = -\frac{\pi}{2}$ .

Since we imposed  $Y_d^0 = Y_e^0$  and  $\mathbf{c}_q = \mathbf{c}_l$ , the left mixing matrices  $W_L^d$  and  $W_L^e$  for the down sector and charged lepton sector are very similar, and in particular the 12 entries have the exact same first order expansion  $(W_L^d)_{12} \simeq (W_L^e)_{12} \simeq V_{us}$ . This is the origin of the quark-lepton complementarity relations we wish to study here.

In the lepton sector, the effective 4D neutrino mass matrix should be diagonalized by a unitary matrix, almost bimaximal, with its third eigenvector (associated to  $m_\nu^3$ ) close to  $(0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ . The charged lepton Yukawa coupling, on the other hand, will be diagonalized by  $W_L^d$ , since both the Yukawa couplings and the doublet  $c$  parameters are the same in the down quark and charged lepton sectors. The unitary matrices generating the charged flavor part of the PMNS matrix  $V_{\text{PMNS}} = (W_L^e)^\dagger W_L^\nu$  should be close to

$$W_L^e \simeq \begin{pmatrix} 1 & V_{us} & V_{ub} \\ -V_{us}^* & 1 & \frac{V_{ub}}{V_{us}} - \frac{f(c_q^2)}{f(c_q^3)} \\ -\frac{f(c_q^2)}{f(c_q^3)} V_{us}^* & -\frac{V_{ub}^*}{V_{us}^*} + \frac{f(c_q^2)}{f(c_q^3)} & 1 \end{pmatrix} \quad \text{and}$$

$$W_L^\nu \simeq \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{2} & \frac{1}{2} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \end{pmatrix} + \mathcal{O}(\epsilon), \quad (4.5)$$

where we have  $W_L^d \simeq W_L^e$ , and where the  $\mathcal{O}(\epsilon)$  entries represent small corrections coming from wave-function effects and flavor symmetry breaking terms. The PMNS matrix elements  $V_{e3}$ ,  $V_{\mu 3}$  and  $V_{e2}$  are then expected to be

$$V_{e3} \simeq \left(1 - \frac{f(c_q^2)}{f(c_q^3)}\right) \frac{V_{us}}{\sqrt{2}} + \mathcal{O}(\epsilon), \quad (4.6)$$

$$V_{\mu 3} \simeq -\frac{1}{\sqrt{2}} \left[1 + \frac{V_{ub}}{V_{us}} - \frac{f(c_q^2)}{f(c_q^3)}\right] + \mathcal{O}(\epsilon), \quad (4.7)$$

$$V_{e2} \simeq \frac{1}{\sqrt{2}} - \frac{1}{2} \left(1 + \frac{f(c_q^2)}{f(c_q^3)}\right) V_{us} + \mathcal{O}(\epsilon). \quad (4.8)$$

Combining these equations we obtain the relations linking the PMNS and CKM mixing matrix elements:

$$V_{e3} \simeq -V_{\mu 3} V_{us} - \frac{V_{ub}}{\sqrt{2}} + \mathcal{O}(\epsilon), \quad (4.9)$$

$$V_{e2} \simeq \frac{1}{\sqrt{2}} \left(1 - \frac{V_{ub}}{\sqrt{2}}\right) - V_{us} \left(1 + \frac{V_{\mu 3}}{\sqrt{2}}\right) + \mathcal{O}(\epsilon). \quad (4.10)$$

In order to check these relations we perform a random scan of the free parameters in the Yukawa couplings, fixing only the absolute value of  $|d| = 1/7$  and allowing its phase and the rest of the complex parameters of Eq. (4.2),  $a, c, e, g$ , to be random, with absolute values of  $\mathcal{O}(1)$ . Still at the level of the Yukawa couplings, we also implement full random perturbations of order 5% as a small flavor symmetry breaking effect. The quark and charged lepton  $c$  parameters are fixed (without a scan) in order to obtain good first order values of CKM angles and masses, and the precise numerical values used are such that the flavor symmetry is also slightly broken. In the neutrino sector we take random values of  $c_\nu^1, c_\nu^2$  and  $c_\nu^3$ , with the only constraint that the values are in the ‘‘neutrino plateau’’ (see Fig. 3) and that their values remain relatively degenerate. In particular we consider three windows, where we randomly scan, i.e.,  $c_\nu^i \in [-2.5, -2.3]$ ,  $c_\nu^i \in [-2.2, -2.0]$  and  $c_\nu^i \in [-2.0, -1.8]$ . During the scan over random parameters, we keep only points that produce numerically correct CKM angles  $|V_{us}| \simeq 0.22$ ,  $|V_{cb}| \simeq 0.041$  and  $|V_{ub}| \simeq 0.0035$  [149], as well as correct neutrino mass differences. In Fig. 5, we test the correlation between  $|V_{e3}|$  and  $|V_{\mu 3}|$  and find it in very good agreement with Eq. (4.9). The bands in the graph represent the  $3\sigma$  uncertainty around the central values obtained from experimental global fits [150,151]. It is quite remarkable that the theoretical correlation curve as well as most of the points generated for the intermediate window lie within these bounds. Higher values of  $c_\nu^i$  and lower values of  $c_\nu^i$  produce points with too high or too small  $|V_{\mu 3}|$ .<sup>9</sup> Note that the lower values of  $c_\nu^i$  lie at the end of the plateau and the beginning of the exponential sensitivity, and so these points start to show greater deviations from the expected correlation. For higher  $c_\nu^i$  values however, the correlation is expected to be quite robust, although the experimental values might not agree with the obtained results. In order to further check this scenario, we perform the same scan as before but including all the points generated, without checking for correct CKM values (although in general they are quite CKM-like). This means that sometimes, the value of the Cabibbo angle is larger or smaller than expected due to accidental alignments or suppressions coming from the Yukawa couplings (taken randomly). Nevertheless the expectation is that the correlation between  $V_{e3}$  and  $V_{us}$  from Eq. (4.6) should survive, and indeed we see in Fig. 6 that this is the case. The exact values of the parameters used in the scan are obtained by adding small deviations to the zero-order values. In the case of the  $c$  parameters we have thus  $c_f^i = 0.6 \pm \mathcal{O}(0.1)$  for quarks and leptons, and  $c_\nu^i = 2 \pm \mathcal{O}(0.1)$

<sup>9</sup>Breaking further the degeneracy in  $c_\nu^i$  will increase the range of possible results.




FIG. 5 (color online). Scans of  $|V_{e3}|$  versus  $|V_{\mu 3}|$  (upper panels) and  $|V_{e2}|$  versus  $|V_{\mu 3}|$  (lower panels) using the 2-3 symmetric Yukawa couplings of Eq. (4.2), with  $|d| = 1/7$  and random  $a, c, e, g$ , and their phases. Small random general perturbations of order 5% to these terms are also included. The right-hand neutrino  $c$  parameters are also taken randomly, within three different windows, i.e.,  $c_\nu^i \in [-2.5, -2.3]$  (left panels),  $c_\nu^i \in [-2.2, -2.0]$  (center panels) and  $c_\nu^i \in [-2.0, -1.8]$  (right panels). The points show agreement with the nontrivial correlation of Eq. (4.9) (represented by the diagonal line in the upper graphs).




FIG. 6 (color online). Scan of the  $V_{us}$  versus  $V_{e3}$  with random perturbations around 2-3 symmetric Yukawa couplings as in the previous figures, with  $c_\nu^i \in [-2.2, -2.0]$ , and allowing for unphysical CKM entries (although still hierarchical), in order to test the quark-lepton correlation expressed in Eq. (4.6) and shown in the graph with the diagonal line.

for right-handed neutrinos. The Yukawa couplings are those given in Eq. (4.2), with the addition of small random perturbations.

## V. FLAVOR DEMOCRACY

In the second symmetry implementation, in this section we assume a democratic structure [152–156] for all flavor parameters, meaning that in our case the 5D Yukawa couplings,  $Y_F^0$  are invariant under  $S_3 \times S_3$  symmetry, while the 5D fermion bulk mass matrices,  $\mathbf{c}_f^0$  are invariant under  $S_3$  permutations. Explicitly, the democratic 5D Yukawa couplings and 5D fermion bulk mass matrices are given by

$$Y_F^{\text{Dem}} \propto \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \quad \text{and} \quad \mathbf{c}_f^{\text{Dem}} = \begin{pmatrix} a_f & b_f & b_f \\ b_f & a_f & b_f \\ b_f & b_f & a_f \end{pmatrix}. \quad (5.1)$$

Overall, there are four Yukawa matrices,  $Y_u^{\text{Dem}}$ ,  $Y_d^{\text{Dem}}$ ,  $Y_e^{\text{Dem}}$  and  $Y_\nu^{\text{Dem}}$ , corresponding to the up-quark sector, the down-quark sector, the charged leptons and the neutrinos. There are six fermion bulk  $c$  matrices, namely  $\mathbf{c}_q^{\text{Dem}}$ ,  $\mathbf{c}_u^{\text{Dem}}$ ,  $\mathbf{c}_d^{\text{Dem}}$ ,  $\mathbf{c}_l^{\text{Dem}}$ ,  $\mathbf{c}_e^{\text{Dem}}$  and  $\mathbf{c}_\nu^{\text{Dem}}$ .

Both matrices can be simultaneously diagonalized by the same unitary transformation resulting in two zero eigenvalues for the Yukawa matrices, and two degenerate eigenvalues for the bulk mass matrices,  $\mathbf{c}_f^{\text{Dem}}$ . The 5D Yukawa and bulk mass matrices thus become, in their diagonal basis,

$$Y_F^0 = y_F^0 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \mathbf{c}_f^0 = \begin{pmatrix} c_f^{10} & 0 & 0 \\ 0 & c_f^{10} & 0 \\ 0 & 0 & c_f^{30} \end{pmatrix}, \quad (5.2)$$

where  $y_F^0$  are complex Yukawa couplings and the index  $F$  runs over  $u, d, e$  and  $\nu$ . The elements  $c_f^{i0}$  are real and the index  $f_i$  runs over doublets  $q_i, l_i$  as well as singlets  $u_i, d_i, \nu_i$  and  $e_i$ , with  $i$  the flavor index. Note that in this flavor symmetric limit, all fermions except the  $t$  quark,  $b$  quark,  $\tau$  lepton and  $\nu_\tau$  neutrino are massless. The 5D flavor structure of Eq. (5.2) yields the zeroth order CKM and PMNS matrices for this scenario:

$$V_i^0 = \begin{pmatrix} \cos \theta_i^0 & \sin \theta_i^0 & 0 \\ -\sin \theta_i^0 & \cos \theta_i^0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (5.3)$$

where  $i = \text{CKM, PMNS}$ . The angle  $\theta_i^0$ , depends on the detailed structure of the symmetry breaking terms,  $\delta Y_F$  and  $\delta \mathbf{c}_f$  in Eqs. (3.8) and (3.7), and is not fixed by the underlying  $S_3 \times S_3$  symmetry. Addition of generic small perturbations, as in Eqs. (3.7) and (3.8), breaks the flavor symmetry and lifts the degeneracies to produce SM-like masses and mixing angles. In the neutrino sector, the two level degeneracy is lifted by a small amount  $(\delta Y_F)_{ij}$ . This suggests a normal hierarchy ordering with one heavier eigenstate, and with two lighter ones having similar masses. Using Eq. (3.4) and taking the generic size of the perturbations as  $(\delta Y_F)_{ij} \approx \delta Y^\nu$  for simplicity yields the following relations for the neutrino masses:

$$\begin{aligned} m_1 &\sim \delta Y^\nu v e^{-ky_1(a-1)}, \\ m_2 &\sim \delta Y^\nu v e^{-ky_1(a-1)}, \\ m_3 &\sim (1 + \delta Y^\nu) v e^{-ky_1(a-1)}. \end{aligned} \quad (5.4)$$

Neutrino mass data requires  $v e^{-ky_1(a-1)} \approx 0.3$  eV and explaining the hierarchy problem requires  $ky_1 \approx 35$ , which means that the value of the Higgs localization parameter

should be about  $a \approx 1.8$ . As explained in a previous section, this value of  $a$  requires some fine-tuning of parameters in the RS 5D Higgs potential. With modified AdS<sub>5</sub> metrics it is possible to remain in a nontuned region, and in particular we find that the best region is obtained for  $\nu \sim 0.2$  and  $kL_1 \sim 0.3$ , where the parameter  $a$  can have values as large as 4.5.<sup>10</sup> In order to obtain the observed neutrino mass hierarchy ratio  $r$ , defined as  $r = (|m_2|^2 - |m_1|^2)/(|m_3|^2 - |m_1|^2) \approx 0.03$ , the size of the Yukawa coupling perturbations  $\delta Y$  must be fixed to  $\delta Y^\nu \lesssim \sqrt{r} \approx 0.17$ . There are no restrictions on the values of bulk mass parameter  $c_\psi^i$ 's (as long as they are within the bounds  $a < c_\nu + c_l$ ).

Consider the elements  $V_{e2}$ ,  $V_{e3}$  and  $V_{\mu3}$  of the PMNS matrix. As mentioned before, due to the plateau in the neutrino sector, for a small  $a$  parameter, the 5D Yukawa matrix structure is more or less preserved and therefore these elements should be close to the ones shown in Eq. (5.3) plus some perturbations. This matrix predicts very small values for both  $|V_{e3}|$  and  $|V_{\mu3}|$  and so the perturbations should lift them (especially  $|V_{\mu3}|$ ). The value of  $V_{e2}$  on the other hand is highly sensitive to the structure of the neutrino Yukawa flavor violating matrix  $\delta Y_{ij}^\nu$  and can be large or small. It turns out that when the  $c$  parameters are such that  $a < c_l^3 + c_\nu^3$  and  $c_l^3 < 1/2$  (a region not generic in usual warped extra dimension scenarios) the lifting of the zero-order PMNS can be successful and we have

$$V_{e2} \sim \sin \theta_\nu^0, \quad (5.5)$$

$$V_{e3} \propto \delta Y_{13}^\nu f(c_l^3), \quad (5.6)$$

$$V_{\mu3} \propto \delta Y_{23}^\nu f(c_l^3). \quad (5.7)$$

Note in particular that we expect that  $V_{e3}/V_{\mu3} \propto \delta Y_{13}^\nu/\delta Y_{23}^\nu$ . In Fig. 7 we present a scan of the model parameters to verify the validity of Eq. (5.5). In this scan we randomly perturbed the values of the 5D parameters  $Y_\nu, Y_e, c_l$  and  $c_\nu$ , but kept the values of the three relevant parameters  $c_l^3, \delta Y_{13}^\nu$  and  $\delta Y_{23}^\nu$  fixed. We see from the figure that the values obtained for the matrix elements  $V_{\mu3}$  and  $V_{e3}$  can be made to lie within the experimental bounds by fixing only these three parameters with all other terms randomly perturbed. In particular, the formulas show the sensitivity of these two PMNS mixing angles to the flavor structure of the neutrino Yukawa matrix  $\delta Y^\nu$ , but not to the charged lepton Yukawa matrix  $\delta Y^l$  or to the bulk masses  $\delta c_\psi^i$ , except for  $\delta c_l^3$ . Knowing that experimentally  $V_{\mu3}^{\text{exp}} \approx 0.65$  and  $V_{e3}^{\text{exp}} \approx 0.15$ , for the numerical evaluations we took the bulk mass parameter of the third family lepton

<sup>10</sup>This is due to the fact that in the modified AdS<sub>5</sub> case, Eq. (5.4) will be slightly modified and higher values of the  $a$  parameter become acceptable.




FIG. 7 (color online). Scatter plots of  $V_{e3}$  versus  $V_{\mu 3}$  with random values of  $\delta Y_\nu$ ,  $\delta Y_e$ ,  $c_l$  and  $c_\nu$ . We have fixed  $c_l^3 = 0.41$  and  $(\delta Y^\nu)_{23} = 0.13$  and have then taken  $(\delta Y^\nu)_{13} = 0.002$  (left panel)  $(\delta Y^\nu)_{13} = 0.008$  (center panel) and  $(\delta Y^\nu)_{13} = 0.032$  (right panel). The concentration of points in a precise region shows that the mixing angles  $V_{e3}$  and  $V_{\mu 3}$  are sensitive to only these three parameters. The obtained values of  $V_{\mu 3}$  lie on the lower side of the experimental window, whereas  $V_{e3}$  depends on the ratio of Yukawa couplings  $(\delta Y^\nu)_{13}/(\delta Y^\nu)_{23}$ , so that its smallness is due to a slight hierarchy in these.

doublet  $c_l^3 < 1/2$  to obtain larger mixing angles for small  $\delta Y \approx 0.1$ , the same as in the quark sector, where  $c_q^3 \lesssim 1/2$  is needed to obtain a large top quark mass, and thus it could be a hint of a possible additional family symmetry among the  $SU(2)$  doublets of the third family. The particular values used in the scan are  $c_l^3 = 0.41$ ,  $\delta Y_{13}^\nu = 0.008$  and  $\delta Y_{23}^\nu = 0.13$ , which produce PMNS angles consistent with the experiment but with  $|V_{\mu 3}|$  on the lower experimental side. When the experimental uncertainty decreases and if the central value of  $|V_{\mu 3}^{\text{exp}}|$  ends up on the higher end of the current allowed region, this scenario will be under great pressure. We also show how indeed the ratio  $V_{e3}/V_{\mu 3}$  depends on the ratio  $\delta Y_{13}^\nu/\delta Y_{23}^\nu$  by increasing and decreasing the value of  $\delta Y_{13}^\nu$ , resulting in larger or smaller values of  $|V_{e3}|$ .

In the charged lepton sector and the up- and down-quark sectors, massless states are also lifted by the flavor symmetry breaking, and these masses emerge directly proportional to the generic size of the perturbations in the Yukawa matrix,  $\delta Y$ , as described in [103].

TABLE I. A parameter point for democratic flavor symmetry in the localization parameter space (out of many possible points) in the zeroth order 5D fermion  $c$  parameter space, consistent with all the experimental and model constraints. For this point, we have set all the zeroth order Yukawa coefficients to be universal,  $y_u^0 = y_d^0 = y_\nu^0 = y_e^0 = 4.4$ , and the Higgs localization parameter to  $a = 2.1$ . The modified AdS<sub>5</sub> metric parameters are  $\nu = 1.1$ ,  $y_1 = 2.8 \times 10^{-17}$  and  $A(y_1) = 35$ .

$f$	$q$	$l$	$u$	$d$	$e$	$\nu$
$c_f^{10}$	0.55	0.55	0.55	0.65	0.65	5.00
$c_f^{30}$	0.45	0.45	0.45	0.65	0.65	3.00

Finally, in Table I we present a set of  $S_3$  symmetric zeroth order bulk  $c$  parameters (many other points with the same symmetry in the parameter space are possible), which, with a small perturbation of order  $\delta c \approx \delta Y \approx 10\%$ , can lead to the SM in the modified AdS<sub>5</sub> scenario.<sup>11</sup> For this specific point we have taken the value of the warp exponent at the IR brane,  $A(y_1)$ , to be exactly 35. With this assignment, there is only one more free parameter in the model left to completely fix the metric. This parameter can be either  $\nu$  or the position of the singularity,  $y_s$ . Therefore one has the freedom to choose the amount of departure from the pure AdS<sub>5</sub>. For the point presented in Table I, we took  $\nu \approx 0.32$  and the value of the localization of the Higgs field operator  $a$  is 4.46 as in the previous example. With this starting point, the SM is then easily reproduced by breaking the symmetry through adding perturbations  $\delta c^i$ 's and  $\delta Y_{ij}$ 's to lift the degeneracies of the symmetric scenario. We require the  $\delta Y_{ij} \lesssim 0.1$  and  $\delta c_\psi^i \lesssim 0.1$  parameters to produce the SM masses, angles and phases, by systematically using the approximate formulas in this section up to any order of precision consistent with the SM. In general the sensitivity to the precise values of the 5D Yukawa couplings,  $\delta Y_{ij}$ , is minimal and randomly chosen matrices with  $\delta Y_{ij} \lesssim 0.1$  produce all the required SM features.

## VI. CONCLUSION

In this work, we have provided a warped extra dimensional framework in which all of the fermions, including neutrinos, are treated on equal footing, and where the SM fermion flavor structure can still emerge out of a slightly

<sup>11</sup>A similar point for the pure AdS<sub>5</sub> was presented in [103].

broken universal flavor symmetry. Essential for this scenario is that all the matter fields must be in the bulk and, in particular, the Higgs field should be as delocalized as possible from the IR boundary. We have explored a general warped scenario in which the metric is modified from the usual AdS<sub>5</sub> background. This setup has the advantage of allowing lower KK masses ( $\sim 1$  to 2 TeV), while still safe from precision electroweak tests and flavor bounds. For our study, the modified metric presents a further advantage, as the neutrino mass generation in our framework is less fine tuned. This is due to a numerical accident by which the neutrino masses generated in the AdS<sub>5</sub> background are too small (at most  $\sim 10^{-4}$  eV) in the parameter region of interest (the neutrino plateau), while in the modified metric neutrino masses can be up to 2 orders of magnitude larger, in the same qualitative region of parameter space. A way out in AdS<sub>5</sub> would be to further delocalize the Higgs field, although requiring such a delocalized Higgs to generically solve the hierarchy problem, some degree of fine-tuning must be introduced in the Higgs potential. These tensions disappear when we use a modified AdS<sub>5</sub> geometry, where our flavor setup can be successfully implemented without further fine-tuning.

In the parameter region of interest (the neutrino plateau) the effective 4D neutrino masses do not have exponential dependence on the bulk mass parameters  $c^i$ , in contrast with quark and charged lepton masses. Once a universal flavor symmetry is slightly broken, the SM flavor structure emerges due to the inherent features of warped space models, i.e., the wave-function profiles of light quarks and charged leptons are exponentially sensitive to the symmetry violating terms, resulting in masses and mixing controlled by small flavor violating terms. In the neutrino sector, in the plateau region with a highly delocalized Higgs field, the wave functions are not exponentially sensitive to Lagrangian parameters and thus the original flavor symmetry is essentially preserved. Overall results are similar in both RS and modified AdS<sub>5</sub> types of scenarios, indicating that the precise nature of the flavor symmetry or the precise nature of the metric solution is not crucial for the main property of the scenarios.

For illustration, we choose two simple examples of different symmetries which can provide an implementation for this mechanism. In the first scenario, we study quark-lepton complementarity associated with  $\mu$ - $\tau$  symmetry (or 2-3 symmetry), known to lead to phenomenologically viable neutrino masses and mixings. Assuming equal Yukawa matrix couplings in the charged lepton and down-quark sector, and equal Yukawa matrix couplings

in the neutrino and up-quark sector, as well as identical localization for the doublet quark and lepton representations, mass differences emerge entirely from singlet representation localization. The predictions of this implementation are definite connections between PMNS and CKM matrix elements, as given by Eqs. (4.9) and (4.10).

In the second example, flavor democracy, the 5D Yukawa couplings for the fermions are invariant under the  $S_3 \times S_3$  symmetry, while the 5D fermion bulk mass matrices are invariant under  $S_3$  family permutation invariance. Small perturbations of  $\mathcal{O}(10\%)$  are enough to generate the full flavor structure of the SM in both quark and lepton sectors, and the ratio of PMNS matrix elements can be simply expressed in terms of these perturbations. Explicit expressions appear in Eq. (5.5). Localization of the third lepton family follows localization of the third quark family, hinting at an additional quark-lepton symmetry. As distinctive predictions and correlations appear in each implementation, the model studied here would yield a very promising novel laboratory for studying fermion flavor symmetries.

## ACKNOWLEDGMENTS

We thank NSERC for partial financial support under Grant No. SAP105354.

## APPENDIX: EXPLICIT EXPRESSIONS FOR THE FIELD PROFILES

Here we derive the explicit expressions for the fermion and Higgs profiles, as well as for the Yukawa couplings for the modified AdS<sub>5</sub> scenario. The fermion profiles are given by

$$q_L^{0,i}(y) = q_0^i e^{(2-c_q^i)A(y)}, \quad u_R^{0,i} = u_0^i e^{(c_u^i+2)A(y)}, \quad (\text{A1})$$

while the Higgs profile is

$$h(y) = h_0 e^{aky}, \quad (\text{A2})$$

with

$$\begin{aligned} q_0^i &= \sqrt{k} e^{\frac{1}{2}-c_q^i} f(c_q^i) \equiv \sqrt{k} \tilde{f}(c_q^i), \\ u_0^i &= \sqrt{k} e^{\frac{1}{2}+c_u^i} f(-c_u^i) \equiv \sqrt{k} \tilde{f}(-c_u^i), \end{aligned} \quad (\text{A3})$$

and

$$\begin{aligned} h_0 &\equiv \sqrt{k} e^{-(a-1)ky_s} \mathfrak{h}_0 \\ \mathfrak{h}_0 &= \frac{1}{\sqrt{ky_s(2(a-1)ky_s)^{-\frac{2}{v^2}-1} (\Gamma(1 + \frac{2}{v^2}, 2(a-1)k(y_s - y_1)) - \Gamma(1 + \frac{2}{v^2}, 2(a-1)ky_s))}}, \end{aligned} \quad (\text{A4})$$

where

$$\epsilon = e^{-A(y)} = e^{-ky_1} \left(1 - \frac{y_1}{y_s}\right)^{\frac{1}{\nu^2}}. \quad (\text{A5})$$

The reduced profile functions  $\mathfrak{f}(c)$  are defined as  $\mathfrak{f}(c) \equiv \epsilon^{\frac{1}{2}-c} f(c)$ , with

$$f(c) \equiv \frac{\epsilon^{c-\frac{1}{2}}}{\sqrt{ky_s e^{(1-2c)ky_s} ((1-2c)ky_s)^{\frac{1-2c}{\nu^2}-1} (\Gamma(1 - \frac{1-2c}{\nu^2}, (1-2c)k(y_s - y_1)) - \Gamma(1 - \frac{1-2c}{\nu^2}, (1-2c)ky_s))}}. \quad (\text{A6})$$

Note that the Higgs and fermion profiles are defined differently, due to the specific Higgs potential we have considered [157]. We can now write down the most general form for the Yukawa couplings as

$$y_{ij}^u = \tilde{Y}_{ij}^u h_0 f(c_q^i) f(-c_u^j), \quad (\text{A7})$$

where the  $\tilde{Y}_{ij}^u$ 's are related to the 5D Yukawa couplings via the equation

$$\begin{aligned} \tilde{Y}_{ij}^u &\equiv Y_{ij}^u \sqrt{k} \epsilon^{1-c_q+c_u} y_s e^{ky_s(a-c_q^i+c_u^j)} [ky_s(a-c_q^i+c_u^j)]^{\frac{c_u^j-c_q^i}{\nu^2}-1} \\ &\times \left[ \Gamma\left(\frac{c_q^i-c_u^j}{\nu^2} + 1, (a-c_q^i+c_u^j)k(y_s-y_1)\right) - \Gamma\left(\frac{c_q^i-c_u^j}{\nu^2} + 1, (a-c_q^i+c_u^j)ky_s\right) \right]. \end{aligned} \quad (\text{A8})$$

Before switching to the expressions in the RS metric, we use the asymptotic expansion of the incomplete gamma function

$$\Gamma(a, z) \sim z^{a-1} e^{-z} \left(1 + \frac{a-1}{z} + \frac{(a-1)(a-2)}{z^2} + \mathcal{O}(z^{-3})\right). \quad (\text{A9})$$

Then we get the following asymptotic behavior, up to the first order in  $\mathcal{O}(z^{a-1})$ , with  $\epsilon$  defined as in Eq. (A5):

$$f(c) \sim \sqrt{\frac{1-2c}{1-\epsilon^{1-2c}}}. \quad (\text{A10})$$

$$h_0 \sim \sqrt{\frac{2(1-a)}{1-e^{2aky_1}\epsilon^2}}, \quad (\text{A11})$$

$$\tilde{Y}_{ij}^u \sim \frac{e^{aky_1} \epsilon^{c_q^i-c_u^j} - 1}{a-c_q^i+c_u^j} Y_{ij}^u. \quad (\text{A12})$$

Keeping the  $\mathcal{O}(z^{a-2})$  term not only gives a much better approximation for the top and neutrino plateaus, but it also gives a more transparent expression for these functions:

$$f(c) \sim \epsilon^{c-\frac{1}{2}} \sqrt{\frac{(1-2c)ky_s\nu^2}{ky_s\nu^2(\epsilon^{2c-1}-1) - \epsilon^{2c-1} + \left(\frac{y_s}{y_s-y_1}\right)}}. \quad (\text{A13})$$

$$h_0 \sim \epsilon^{-1} e^{-aky_1} \sqrt{\frac{2(1-a)^2 ky_s \nu^2}{ky_s \nu^2 (1-a)(\epsilon^{-2} e^{-2aky_1} - 1) - \epsilon^{-2} e^{-2aky_1} + \left(\frac{y_s}{y_s-y_1}\right)}}. \quad (\text{A14})$$

From these formulas, valid for the general modified AdS<sub>5</sub> metric, one can see that, taking the limits  $\nu \rightarrow \infty$  and  $y_s \rightarrow \infty$  we obtain the expressions for the profiles in the RS metric:

$$f^{\text{RS}}(c) = \sqrt{\frac{1-2c}{1-\epsilon^{1-2c}}} \equiv \epsilon^{c-\frac{1}{2}} \mathfrak{f}^{\text{RS}}(c), \quad h_0^{\text{RS}} = e^{(1-a)ky_1} \sqrt{\frac{2(1-a)}{\epsilon^{2(a-1)} - 1}}, \quad (\text{A15})$$

$$\tilde{Y}_{ij}^{\text{RS},u} \equiv \frac{\epsilon^{-(a-c_q^i+c_u^j)} - 1}{a-c_q^i+c_u^j} Y_{ij}^{\text{RS},u}, \quad y_{ij}^{\text{RS},u} = \tilde{Y}_{ij}^{\text{RS},u} h_0^{\text{RS}} f^{\text{RS}}(c_q^i) f^{\text{RS}}(-c_u^j), \quad (\text{A16})$$

where  $\epsilon \equiv e^{-ky_1}$ .

- [1] F. P. An *et al.* (DAYA-BAY Collaboration), Observation of Electron-Antineutrino Disappearance at Daya Bay, *Phys. Rev. Lett.* **108**, 171803 (2012).
- [2] F. P. An *et al.* (Daya Bay Collaboration), Improved measurement of electron antineutrino disappearance at Daya Bay, *Chin. Phys. C* **37**, 011001 (2013).
- [3] K. Abe *et al.* (T2K Collaboration), Indication of Electron Neutrino Appearance from an Accelerator-Produced Off-Axis Muon Neutrino Beam, *Phys. Rev. Lett.* **107**, 041801 (2011).
- [4] K. Abe *et al.* (T2K Collaboration), Precise Measurement of the Neutrino Mixing Parameter  $\theta_{23}$  from Muon Neutrino Disappearance in an Off-Axis Beam, *Phys. Rev. Lett.* **112**, 181801 (2014).
- [5] P. Adamson *et al.* (MINOS Collaboration), Measurement of the Neutrino Mass Splitting and Flavor Mixing by MINOS, *Phys. Rev. Lett.* **106**, 181801 (2011).
- [6] P. Adamson *et al.* (MINOS Collaboration), Combined Analysis of  $\nu_\mu$  Disappearance and  $\nu_\mu \rightarrow \nu_e$  Appearance in MINOS Using Accelerator and Atmospheric Neutrinos, *Phys. Rev. Lett.* **112**, 191801 (2014).
- [7] J. K. Ahn *et al.* (RENO Collaboration), Observation of Reactor Electron Antineutrino Disappearance in the RENO Experiment, *Phys. Rev. Lett.* **108**, 191802 (2012).
- [8] Y. Abe *et al.* (Double Chooz Collaboration), Reactor electron antineutrino disappearance in the Double Chooz experiment, *Phys. Rev. D* **86**, 052008 (2012).
- [9] Y. Abe *et al.* (Double Chooz Collaboration), Improved measurements of the neutrino mixing angle  $\theta_{13}$  with the Double Chooz detector, *J. High Energy Phys.* **10** (2014) 86.
- [10] P. Minkowski,  $\mu \rightarrow e\gamma$  at a rate of one out of 1-billion muon decays?, *Phys. Lett.* **67B**, 421 (1977).
- [11] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1980), p. 315.
- [12] T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, Japan, 1979), p. 95.
- [13] S. L. Glashow, in *Proceedings of the 1979 Cargese Summer Institute on Quarks and Leptons*, edited by M. Levy *et al.* (Plenum Press, New York, 1980), p. 687.
- [14] R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Violation, *Phys. Rev. Lett.* **44**, 912 (1980).
- [15] J. Schechter and J. W. F. Valle, Neutrino masses in  $SU(2) \times U(1)$  theories, *Phys. Rev. D* **22**, 2227 (1980).
- [16] G. Lazarides, Q. Shafi, and C. Wetterich, Proton lifetime and fermion masses in an  $SO(10)$  model, *Nucl. Phys.* **B181**, 287 (1981).
- [17] P. H. Frampton, S. L. Glashow, and D. Marfatia, Zeroes of the neutrino mass matrix, *Phys. Lett. B* **536**, 79 (2002).
- [18] L. Lavoura, Zeros of the inverted neutrino mass matrix, *Phys. Lett. B* **609**, 317 (2005).
- [19] S. Dev, S. Verma, S. Gupta, and R. R. Gautam, Neutrino mass matrices with a texture zero and a vanishing minor, *Phys. Rev. D* **81**, 053010 (2010).
- [20] C. S. Lam, Determining Horizontal Symmetry from Neutrino Mixing, *Phys. Rev. Lett.* **101**, 121602 (2008).
- [21] C. S. Lam, Unique horizontal symmetry of leptons, *Phys. Rev. D* **78**, 073015 (2008).
- [22] W. Grimus, L. Lavoura, and P. O. Ludl, Is  $S(4)$  the horizontal symmetry of tri-bimaximal lepton mixing?, *J. Phys. G* **36**, 115007 (2009).
- [23] S. F. Ge, D. A. Dicus, and W. W. Repko,  $Z_2$  symmetry prediction for the leptonic Dirac  $CP$  phase, *Phys. Lett. B* **702**, 220 (2011).
- [24] H. J. He and F. R. Yin, Common origin of  $\mu$ - $\tau$  and  $CP$  breaking in neutrino seesaw, baryon asymmetry, and hidden flavor symmetry, *Phys. Rev. D* **84**, 033009 (2011).
- [25] R. d. A. Toorop, F. Feruglio, and C. Hagedorn, Discrete flavour symmetries in light of T2K, *Phys. Lett. B* **703**, 447 (2011).
- [26] S. F. Ge, D. A. Dicus, and W. W. Repko, Residual Symmetries for Neutrino Mixing with a Large  $\theta_{13}$  and Nearly Maximal  $\delta_D$ , *Phys. Rev. Lett.* **108**, 041801 (2012).
- [27] R. de Adelhart Toorop, F. Feruglio, and C. Hagedorn, Finite modular groups and lepton mixing, *Nucl. Phys.* **B858**, 437 (2012).
- [28] H. J. He and X. J. Xu, Octahedral symmetry with geometrical breaking: New prediction for neutrino mixing angle  $\theta_{13}$  and  $CP$  violation, *Phys. Rev. D* **86**, 111301 (2012).
- [29] D. Hernandez and A. Y. Smirnov, Lepton mixing and discrete symmetries, *Phys. Rev. D* **86**, 053014 (2012).
- [30] C. S. Lam, Finite symmetry of leptonic mass matrices, *Phys. Rev. D* **87**, 013001 (2013).
- [31] D. Hernandez and A. Y. Smirnov, Discrete symmetries and model-independent patterns of lepton mixing, *Phys. Rev. D* **87**, 053005 (2013).
- [32] M. Holthausen, K. S. Lim, and M. Lindner, Lepton mixing patterns from a scan of finite discrete groups, *Phys. Lett. B* **721**, 61 (2013).
- [33] B. Hu, Neutrino mixing and discrete symmetries, *Phys. Rev. D* **87**, 033002 (2013).
- [34] D. Hernandez and A. Y. Smirnov, Relating neutrino masses and mixings by discrete symmetries, *Phys. Rev. D* **88**, 093007 (2013).
- [35] S. F. King, T. Neder, and A. J. Stuart, Lepton mixing predictions from  $\Delta(6n^2)$  family symmetry, *Phys. Lett. B* **726**, 312 (2013).
- [36] C. Hagedorn, A. Meroni, and L. Vitale, Mixing patterns from the groups  $\Sigma(n\phi)$ , *J. Phys. A* **47**, 055201 (2014).
- [37] L. Lavoura and P. O. Ludl, Residual  $Z_2 \times Z_2$  symmetries and lepton mixing, *Phys. Lett. B* **731**, 331 (2014).
- [38] P. F. Harrison, D. H. Perkins, and W. G. Scott, A redetermination of the neutrino mass squared difference in trimaximal mixing with terrestrial matter effects, *Phys. Lett. B* **458**, 79 (1999).
- [39] P. F. Harrison, D. H. Perkins, and W. G. Scott, Tri-bimaximal mixing and the neutrino oscillation data, *Phys. Lett. B* **530**, 167 (2002).
- [40] Z. z. Xing, Nearly tri-bimaximal neutrino mixing and  $CP$  violation, *Phys. Lett. B* **533**, 85 (2002).
- [41] P. F. Harrison and W. G. Scott, Symmetries and generalizations of tri-bimaximal neutrino mixing, *Phys. Lett. B* **535**, 163 (2002).
- [42] P. F. Harrison and W. G. Scott, Permutation symmetry, tri-bimaximal neutrino mixing and the  $S_3$  group characters, *Phys. Lett. B* **557**, 76 (2003).



- [43] X. G. He and A. Zee, Some simple mixing and mass matrices for neutrinos, *Phys. Lett. B* **560**, 87 (2003).
- [44] F. Vissani, A study of the scenario with nearly degenerate Majorana neutrinos, [arXiv:hep-ph/9708483](https://arxiv.org/abs/hep-ph/9708483).
- [45] V. D. Barger, S. Pakvasa, T. J. Weiler, and K. Whisnant, Bimaximal mixing of three neutrinos, *Phys. Lett. B* **437**, 107 (1998).
- [46] A. J. Baltz, A. S. Goldhaber, and M. Goldhaber, Solar Neutrino Puzzle: An Oscillation Solution with Maximal Neutrino Mixing, *Phys. Rev. Lett.* **81**, 5730 (1998).
- [47] I. Stancu and D. V. Ahluwalia,  $L/E$  flatness of the electron-like event ratio in super-Kamiokande and a degeneracy in neutrino masses, *Phys. Lett. B* **460**, 431 (1999).
- [48] H. Georgi and S. L. Glashow, Neutrinos on Earth and in the heavens, *Phys. Rev. D* **61**, 097301 (2000).
- [49] N. Li and B. Q. Ma, A new parametrization of the neutrino mixing matrix, *Phys. Lett. B* **600**, 248 (2004).
- [50] H. Fritzsch and Z. Z. Xing, Lepton mass hierarchy and neutrino oscillations, *Phys. Lett. B* **372**, 265 (1996).
- [51] H. Fritzsch and Z. z. Xing, Large leptonic flavor mixing and the mass spectrum of leptons, *Phys. Lett. B* **440**, 313 (1998).
- [52] H. Fritzsch and Z. z. Xing, Maximal neutrino mixing and maximal  $CP$  violation, *Phys. Rev. D* **61**, 073016 (2000).
- [53] Y. Zhang, X. Zhang, and B. Q. Ma, Quark-lepton complementarity and self-complementarity in different schemes, *Phys. Rev. D* **86**, 093019 (2012).
- [54] X. Zhang, Y. j. Zheng, and B. Q. Ma, Quark-lepton complementarity revisited, *Phys. Rev. D* **85**, 097301 (2012).
- [55] G. N. Li, H. H. Lin, and X. G. He, Comment on reparametrization invariance of quark-lepton complementarity, *Phys. Lett. B* **711**, 57 (2012).
- [56] S. K. Kang, Revisiting the quark-lepton complementarity and triminimal parametrization of neutrino mixing matrix, *Phys. Rev. D* **83**, 097301 (2011).
- [57] Y. H. Ahn, H. Y. Cheng, and S. Oh, Quark-lepton complementarity and tribimaximal neutrino mixing from discrete symmetry, *Phys. Rev. D* **83**, 076012 (2011).
- [58] Y. Shimizu and R. Takahashi, Deviations from tribimaximality and quark-lepton complementarity, *Europhys. Lett.* **93**, 61001 (2011).
- [59] K. M. Patel, An  $SO(10) \times S_4$  model of quark-lepton complementarity, *Phys. Lett. B* **695**, 225 (2011).
- [60] J. Barranco, F. Gonzalez Canales, and A. Mondragon, Universal mass texture,  $CP$  violation and quark-lepton complementarity, *Phys. Rev. D* **82**, 073010 (2010).
- [61] Y. j. Zheng, Dependence of the quark-lepton complementarity on parametrizations of the CKM and PMNS matrices, *Phys. Rev. D* **81**, 073009 (2010).
- [62] Z. z. Xing, Right unitarity triangles, stable  $CP$ -violating phases and approximate quark-lepton complementarity, *Phys. Lett. B* **679**, 111 (2009).
- [63] P. H. Frampton and S. Matsuzaki, Strong incompatibility of tribimaximal mixing with quark-lepton complementarity, *Mod. Phys. Lett. A* **24**, 429 (2009).
- [64] W. Rodejohann and K. A. Hochmuth, Lepton flavor violation, leptogenesis and neutrino mixing in quark-lepton complementarity scenarios, *Int. J. Mod. Phys. A* **22**, 5875 (2007).
- [65] F. Plentinger, G. Seidl, and W. Winter, Seesaw mechanism in quark-lepton complementarity, *Phys. Rev. D* **76**, 113003 (2007).
- [66] M. Picariello, B. C. Chauhan, J. Pulido, and E. Torrente-Lujan, Predictions from non-trivial quark-lepton complementarity, *Int. J. Mod. Phys. A* **22**, 5860 (2007).
- [67] L. Randall and R. Sundrum, Large Mass Hierarchy from a Small Extra Dimension, *Phys. Rev. Lett.* **83**, 3370 (1999).
- [68] L. Randall and R. Sundrum, An Alternative to Compactification, *Phys. Rev. Lett.* **83**, 4690 (1999).
- [69] Y. Grossman and M. Neubert, Neutrino masses and mixings in nonfactorizable geometry, *Phys. Lett. B* **474**, 361 (2000).
- [70] T. Gherghetta and A. Pomarol, Bulk fields and supersymmetry in a slice of AdS, *Nucl. Phys.* **B586**, 141 (2000).
- [71] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, Bulk gauge fields in the Randall-Sundrum model, *Phys. Lett. B* **473**, 43 (2000).
- [72] S. Chang, J. Hisano, H. Nakano, N. Okada, and M. Yamaguchi, Bulk standard model in the Randall-Sundrum background, *Phys. Rev. D* **62**, 084025 (2000).
- [73] S. J. Huber, Flavor violation and warped geometry, *Nucl. Phys.* **B666**, 269 (2003).
- [74] S. Casagrande, F. Goertz, U. Haisch, M. Neubert, and T. Pfoh, The custodial Randall-Sundrum model: From precision tests to Higgs physics, *J. High Energy Phys.* **09** (2010) 014.
- [75] K. Agashe, G. Perez, and A. Soni, Flavor structure of warped extra dimension models, *Phys. Rev. D* **71**, 016002 (2005).
- [76] K. Agashe, A. Delgado, M. J. May, and R. Sundrum, RS1, custodial isospin and precision tests, *J. High Energy Phys.* **08** (2003) 050.
- [77] G. Burdman, Constraints on the bulk standard model in the Randall-Sundrum scenario, *Phys. Rev. D* **66**, 076003 (2002).
- [78] G. Burdman, Flavor violation in warped extra dimensions and  $CP$  asymmetries in  $B$  decays, *Phys. Lett. B* **590**, 86 (2004).
- [79] M. S. Carena, A. Delgado, E. Ponton, T. M. P. Tait, and C. E. M. Wagner, Warped fermions and precision tests, *Phys. Rev. D* **71**, 015010 (2005).
- [80] M. S. Carena, A. Delgado, E. Ponton, T. M. P. Tait, and C. E. M. Wagner, Precision electroweak data and unification of couplings in warped extra dimensions, *Phys. Rev. D* **68**, 035010 (2003).
- [81] A. Delgado and A. Falkowski, Electroweak observables in a general 5D background, *J. High Energy Phys.* **05** (2007) 097.
- [82] M. Quiros, Higgs bosons in extra dimensions, *Mod. Phys. Lett. A* **30**, 1540012 (2015).
- [83] C. Csaki, A. Falkowski, and A. Weiler, The flavor of the composite pseudo-Goldstone Higgs, *J. High Energy Phys.* **09** (2008) 008.
- [84] K. Agashe, A. Azatov, and L. Zhu, Flavor violation tests of warped/composite SM in the two-site approach, *Phys. Rev. D* **79**, 056006 (2009).
- [85] K. Agashe, R. Contino, L. Da Rold, and A. Pomarol, A custodial symmetry for  $Zb\bar{b}$ , *Phys. Lett. B* **641**, 62 (2006).

- [86] A. Falkowski and M. Perez-Victoria, Electroweak breaking on a soft wall, *J. High Energy Phys.* **12** (2008) 107.
- [87] B. Batell, T. Gherghetta, and D. Sword, The soft-wall standard model, *Phys. Rev. D* **78**, 116011 (2008).
- [88] S. Mert Aybat and J. Santiago, Bulk fermions in warped models with a soft wall, *Phys. Rev. D* **80**, 035005 (2009).
- [89] J. A. Cabrer, G. von Gersdorff, and M. Quiros, Soft-wall stabilization, *New J. Phys.* **12**, 075012 (2010).
- [90] J. A. Cabrer, G. von Gersdorff, and M. Quiros, Warped electroweak breaking without custodial symmetry, *Phys. Lett. B* **697**, 208 (2011).
- [91] J. A. Cabrer, G. von Gersdorff, and M. Quiros, Suppressing electroweak precision observables in 5D warped models, *J. High Energy Phys.* **05** (2011) 083.
- [92] A. Azatov, M. Toharia, and L. Zhu, Higgs production from gluon fusion in warped extra dimensions, *Phys. Rev. D* **82**, 056004 (2010).
- [93] M. Frank, N. Pourtolami, and M. Toharia, Higgs bosons in warped space, from the bulk to the brane, *Phys. Rev. D* **87**, 096003 (2013).
- [94] M. Frank, N. Pourtolami, and M. Toharia, Higgs production in general 5D warped models, *Phys. Rev. D* **89**, 016012 (2014).
- [95] R. Malm, M. Neubert, K. Novotny, and C. Schmell, 5D perspective on Higgs production at the boundary of a warped extra dimension, *J. High Energy Phys.* **01** (2014) 173.
- [96] M. Carena, S. Casagrande, F. Goertz, U. Haisch, and M. Neubert, Higgs production in a warped extra dimension, *J. High Energy Phys.* **08** (2012) 156.
- [97] P. R. Archer, M. Carena, A. Carmona, and M. Neubert, Higgs production and decay in models of a warped extra dimension with a bulk Higgs, *J. High Energy Phys.* **01** (2015) 060.
- [98] C. Bouchart and G. Moreau, Higgs boson phenomenology and VEV shift in the RS scenario, *Phys. Rev. D* **80**, 095022 (2009).
- [99] A. Djouadi and G. Moreau, Higgs production at the LHC in warped extra-dimensional models, *Phys. Lett. B* **660**, 67 (2008).
- [100] A. Falkowski, Pseudo-goldstone Higgs production via gluon fusion, *Phys. Rev. D* **77**, 055018 (2008).
- [101] C. Csaki, C. Delaunay, C. Grojean, and Y. Grossman, A model of lepton masses from a warped extra dimension, *J. High Energy Phys.* **10** (2008) 055.
- [102] K. Agashe, T. Okui, and R. Sundrum, Common Origin for Neutrino Anarchy and Charged Hierarchies, *Phys. Rev. Lett.* **102**, 101801 (2009).
- [103] M. Frank, C. Hamzaoui, N. Pourtolami, and M. Toharia, Unified flavor symmetry from warped dimensions, *Phys. Lett. B* **742**, 178 (2015).
- [104] K. Agashe, G. Perez, and A. Soni, *B*-Factory Signals for a Warped Extra Dimension, *Phys. Rev. Lett.* **93**, 201804 (2004).
- [105] K. Agashe, A. E. Blechman, and F. Petriello, Probing the Randall-Sundrum geometric origin of flavor with lepton flavor violation, *Phys. Rev. D* **74**, 053011 (2006).
- [106] K. Agashe, G. Perez, and A. Soni, Collider signals of top quark flavor violation from a warped extra dimension, *Phys. Rev. D* **75**, 015002 (2007).
- [107] F. del Aguila and J. Santiago, Universality limits on bulk fermions, *Phys. Lett. B* **493**, 175 (2000).
- [108] F. del Aguila and J. Santiago, Top mixing in effective theories, *Nucl. Phys. B, Proc. Suppl.* **89**, 43 (2000).
- [109] S. Chang, C. S. Kim, and M. Yamaguchi, Hierarchical mass structure of fermions in warped extra dimension, *Phys. Rev. D* **73**, 033002 (2006).
- [110] G. Moreau and J. I. Silva-Marcos, Flavor physics of the RS model with KK masses reachable at LHC, *J. High Energy Phys.* **03** (2006) 090.
- [111] G. Moreau and J. I. Silva-Marcos, Neutrinos in warped extra dimensions, *J. High Energy Phys.* **01** (2006) 048.
- [112] S. Casagrande, F. Goertz, U. Haisch, M. Neubert, and T. Pfoh, Flavor physics in the Randall-Sundrum model: I. Theoretical setup and electroweak precision tests, *J. High Energy Phys.* **10** (2008) 094.
- [113] S. J. Huber and Q. Shafi, Fermion masses, mixings and proton decay in a Randall-Sundrum model, *Phys. Lett. B* **498**, 256 (2001).
- [114] S. J. Huber and Q. Shafi, Neutrino oscillations and rare processes in models with a small extra dimension, *Phys. Lett. B* **512**, 365 (2001).
- [115] S. J. Huber and Q. Shafi, Majorana neutrinos in a warped 5-D standard model, *Phys. Lett. B* **544**, 295 (2002).
- [116] S. J. Huber and Q. Shafi, Seesaw mechanism in warped geometry, *Phys. Lett. B* **583**, 293 (2004).
- [117] G. J. Ding and Y. L. Zhou, Dirac neutrinos with  $S_4$  flavor symmetry in warped extra dimensions, *Nucl. Phys.* **B876**, 418 (2013).
- [118] A. E. Cárcamo Hernández, C. O. Dib, N. A. Neill, and A. R. Zerwekh, Quark masses and mixings in the RS1 model with a condensing 4th generation, *J. High Energy Phys.* **02** (2012) 132.
- [119] S. Gopalakrishna, T. Mandal, S. Mitra, and G. Moreau, LHC signatures of warped-space vectorlike quarks, *J. High Energy Phys.* **08** (2014) 079.
- [120] S. Gopalakrishna, T. Mandal, S. Mitra, and R. Tibrewala, LHC signatures of a vector-like  $b'$ , *Phys. Rev. D* **84**, 055001 (2011).
- [121] A. M. Iyer and S. K. Vempati, Lepton masses and flavor violation in Randall-Sundrum model, *Phys. Rev. D* **86**, 056005 (2012).
- [122] A. Carmona and J. Santiago, The effective Lagrangian for bulk fermions in models with extra dimensions, *J. High Energy Phys.* **01** (2012) 100.
- [123] A. Carmona, E. Ponton, and J. Santiago, Phenomenology of non-custodial warped models, *J. High Energy Phys.* **10** (2011) 137.
- [124] K. Agashe, C. Csaki, C. Grojean, and M. Reece, The  $S$ -parameter in holographic technicolor models, *J. High Energy Phys.* **12** (2007) 003.
- [125] J. A. Cabrer, G. von Gersdorff, and M. Quiros, Flavor phenomenology in general 5D warped spaces, *J. High Energy Phys.* **01** (2012) 033.
- [126] J. de Blas, A. Delgado, B. Ostdiek, and A. de la Puente, LHC signals of noncustodial warped 5D models, *Phys. Rev. D* **86**, 015028 (2012).
- [127] E. Ma and M. Raidal, Neutrino Mass, Muon Anomalous Magnetic Moment, and Lepton Flavor Nonconservation, *Phys. Rev. Lett.* **87**, 011802 (2001); **87**, 159901(E) (2001).

- [128] W. Grimus and L. Lavoura, Softly broken lepton numbers and maximal neutrino mixing, *J. High Energy Phys.* **07** (2001) 045.
- [129] R. N. Mohapatra and S. Nasri, Leptogenesis and  $\mu$ - $\tau$  symmetry, *Phys. Rev. D* **71**, 033001 (2005).
- [130] R. N. Mohapatra, S. Nasri, and H. B. Yu, Leptogenesis,  $\mu$ - $\tau$  symmetry and  $\theta_{13}$ , *Phys. Lett. B* **615**, 231 (2005).
- [131] S. Nasri, Implications of  $\mu \leftrightarrow \tau$  symmetry on neutrinos and leptogenesis, *Int. J. Mod. Phys. A* **20**, 6258 (2005).
- [132] R. N. Mohapatra, S. Nasri, and H. B. Yu, Grand unification of  $\mu$ - $\tau$  symmetry, *Phys. Lett. B* **636**, 114 (2006).
- [133] R. N. Mohapatra, S. Nasri, and H. B. Yu,  $S(3)$  symmetry and tri-bimaximal mixing, *Phys. Lett. B* **639**, 318 (2006).
- [134] Z. z. Xing, H. Zhang, and S. Zhou, Nearly tri-bimaximal neutrino mixing and  $CP$  violation from  $\mu$ - $\tau$  symmetry breaking, *Phys. Lett. B* **641**, 189 (2006).
- [135] Y. H. Ahn, S. K. Kang, C. S. Kim, and J. Lee, Phased breaking of  $\mu$ - $\tau$  symmetry and leptogenesis, *Phys. Rev. D* **73**, 093005 (2006).
- [136] I. Aizawa and M. Yasue, New type of complex neutrino mass texture and  $\mu$ - $\tau$  symmetry, *Phys. Rev. D* **73**, 015002 (2006).
- [137] K. Fuki and M. Yasue, What does  $\mu$ - $\tau$  symmetry imply in neutrino mixings?, *Phys. Rev. D* **73**, 055014 (2006).
- [138] A. S. Joshipura, Universal 2-3 symmetry, *Eur. Phys. J. C* **53**, 77 (2008).
- [139] B. Adhikary, A. Ghosal, and P. Roy,  $\mu$ - $\tau$  symmetry, tribimaximal mixing and four zero neutrino Yukawa textures, *J. High Energy Phys.* **10** (2009) 040.
- [140] Z. z. Xing and Y. L. Zhou, A generic diagonalization of the  $3 \times 3$  neutrino mass matrix and its implications on the  $\mu$ - $\tau$  flavor symmetry and maximal  $CP$  violation, *Phys. Lett. B* **693**, 584 (2010).
- [141] S. F. Ge, H. J. He, and F. R. Yin, Common origin of soft  $\mu$ - $\tau$  and  $CP$  breaking in neutrino seesaw and the origin of matter, *J. Cosmol. Astropart. Phys.* **05** (2010) 017.
- [142] I. de Medeiros Varzielas, R. Gonzalez Felipe, and H. Serodio, Leptonic mixing, family symmetries, and neutrino phenomenology, *Phys. Rev. D* **83**, 033007 (2011).
- [143] S. Gupta, A. S. Joshipura, and K. M. Patel, How good is  $\mu$ - $\tau$  symmetry after results on non-zero  $\theta_{13}$ ?, *J. High Energy Phys.* **09** (2013) 035.
- [144] B. Adhikary, M. Chakraborty, and A. Ghosal, Masses, mixing angles and phases of general Majorana neutrino mass matrix, *J. High Energy Phys.* **10** (2013) 043; **09** (2014) 180(E).
- [145] B. Adhikary, A. Ghosal, and P. Roy,  $\theta_{13}$ ,  $\mu$  $\tau$  symmetry breaking and neutrino Yukawa textures, *Int. J. Mod. Phys. A* **28**, 1350118 (2013).
- [146] J. Liao, D. Marfatia, and K. Whisnant, Perturbations to  $\mu$ - $\tau$  symmetry in neutrino mixing, *Phys. Rev. D* **87**, 013003 (2013).
- [147] C. Hamzaoui, S. Nasri, and M. Toharia, Partial  $\mu$ - $\tau$  textures and leptogenesis, *Phys. Rev. D* **89**, 073019 (2014).
- [148] E. I. Lashin, N. Chamoun, C. Hamzaoui, and S. Nasri, Neutrino mass textures and partial  $\mu$ - $\tau$  symmetry, *Phys. Rev. D* **89**, 093004 (2014).
- [149] K. A. Olive *et al.* (Particle Data Group Collaboration), Review of particle physics, *Chin. Phys. C* **38**, 090001 (2014).
- [150] M. C. Gonzalez-Garcia, M. Maltoni, and T. Schwetz, Updated fit to three neutrino mixing: Status of leptonic  $CP$  violation, *J. High Energy Phys.* **11** (2014) 052.
- [151] M. Maltoni, T. Schwetz, M. A. Tortola, and J. W. F. Valle, Status of global fits to neutrino oscillations, *New J. Phys.* **6**, 122 (2004).
- [152] H. Harari, H. Haut, and J. Weyers, Quark masses and Cabibbo angles, *Phys. Lett.* **78B**, 459 (1978).
- [153] S. Pakvasa and H. Sugawara, Discrete symmetry and Cabibbo angle, *Phys. Lett.* **73B**, 61 (1978).
- [154] S. Pakvasa and H. Sugawara, Mass of the  $t$  quark in  $SU(2) \times U(1)$ , *Phys. Lett.* **82B**, 105 (1979).
- [155] E. Derman and H. S. Tsao,  $[SU_2 \times U_1] \times S_n$  flavor dynamics and a bound on the number of flavors, *Phys. Rev. D* **20**, 1207 (1979).
- [156] Y. Yamanaka, H. Sugawara, and S. Pakvasa, Permutation symmetries and the fermion mass matrix, *Phys. Rev. D* **25**, 1895 (1982); **29**, 2135(E) (1984).
- [157] A. Azatov, M. Toharia, and L. Zhu, Higgs mediated flavor changing neutral currents in warped extra dimensions, *Phys. Rev. D* **80**, 035016 (2009).