

# Hydrodynamics with gauge anomaly: Variational principle and Hamiltonian formulation

Gustavo M. Monteiro,<sup>1</sup> Alexander G. Abanov,<sup>1,2</sup> and V. P. Nair<sup>3</sup>

<sup>1</sup>*Department of Physics and Astronomy, Stony Brook University, Stony Brook, New York 11794, USA*

<sup>2</sup>*Simons Center for Geometry and Physics, Stony Brook University, Stony Brook, New York 11794, USA*

<sup>3</sup>*Physics Department, City College of the CUNY, New York, New York 10031, USA*

(Received 28 February 2015; published 26 June 2015)

We present a variational principle for relativistic hydrodynamics with gauge-anomaly terms for a fluid coupled to an Abelian background gauge field. For this we utilize the Clebsch parametrization of the velocity field. We also set up the Hamiltonian formulation and the canonical framework for the theory. While the equations of motion only involve the density and velocity fields, i.e., the Clebsch potentials only appear in the combination which is the velocity field, the generators of symmetry transformations (including the Hamiltonian) depend explicitly on one of the Clebsch potentials, if the background field is time-dependent. For the special case of time-independent background fields, this feature is absent.

DOI: [10.1103/PhysRevD.91.125033](https://doi.org/10.1103/PhysRevD.91.125033)

PACS numbers: 04.20.Fy, 12.39.Fe, 11.30.Rd

## I. INTRODUCTION

Hydrodynamics is a long-wavelength effective description of interacting systems based on the assumption of local equilibrium. Hydrodynamic equations are essentially local conservation laws supplemented by the constitutive relations between conserved densities. These conservation laws are macroscopic manifestations of symmetries of the system. Constitutive relations are often written phenomenologically and involve unknown “equations of state,” which in principle should be obtainable from the underlying “microscopic” theory such as kinetic theory, many body models or quantum field theory [1].

If the underlying theory is a quantum field theory (QFT) with *quantum anomalies*, the conservation laws corresponding to anomalous symmetries are broken. However, the anomalous symmetry breaking is rather subtle and one might hope for an applicability of a universal hydrodynamic description with additional hydrodynamic terms taking anomalies into account. This possibility was noticed initially in AdS/CFT systems [2,3], and then in genuine relativistic hydrodynamic formulation by Son and Surowka for a particular case of Abelian gauge anomaly [4].

The goal of this work is to find variational and Hamiltonian formulations of the hydrodynamics with gauge anomaly [4]. Variational and Hamiltonian approaches to hydrodynamics have a long history and we refer the reader to Refs. [5,6] for reviews. The Hamiltonian formalism is appropriate to study wavelike excitations and instabilities near the fixed point—through the linear analysis of the eigenmodes—and provides the most appropriate framework to study perturbation theory and symmetries of the system. We aim to understand how the quantum anomaly affects the canonical generators of gauge transformations and diffeomorphisms as well as their semidirect product algebra. Our approach will be entirely 3 + 1 dimensional, providing a

minimal generalization of the standard action principle for fluid dynamics to accommodate anomalies.

Let us start with equations of anomalous hydrodynamics of [4]. The current and energy-momentum conservation laws for anomalous QFT in the background gauge field can be written as:

$$\partial_\lambda j^\lambda = -\frac{C}{8} \epsilon^{\lambda\nu\sigma\tau} F_{\lambda\nu} F_{\sigma\tau}, \quad (1)$$

$$\partial_\lambda T^{\lambda\nu} = F^{\nu\sigma} j_\sigma. \quad (2)$$

The right-hand side of Eq. (2) is the Lorentz force, while the right-hand side of (1) is the gauge anomaly term, fully characterized by a single dimensionless constant  $C$ . Here and in the following we drop the angular brackets denoting expectation values, e.g.,  $\langle j \rangle \rightarrow j$ , so that  $j^\lambda$  and  $T^{\lambda\nu}$  are classical fields representing the current and the energy-momentum tensor.

Assuming local equilibrium and imposing the local form of the second law of thermodynamics, Son and Surowka were able to constrain the form of constitutive relations. In this paper we are interested in the case of zero temperature and absence of dissipation. Thus, we will use a particular form of the constitutive relations found in [4], which is given by:

$$j^\lambda = nu^\lambda + \frac{C}{12} \epsilon^{\lambda\nu\sigma\tau} \mu u_\nu (2\mu \partial_\sigma u_\tau + 3F_{\sigma\tau}), \quad (3)$$

$$T^{\lambda\nu} = n\mu u^\lambda u^\nu + P(\mu) g^{\lambda\nu}. \quad (4)$$

Here we have introduced the equation of state of the fluid  $P(\mu)$  which gives the fluid pressure  $P$  as a function of the chemical potential  $\mu$ . The charge density in the fluid rest frame is given by  $n = P'(\mu)$ . The fluid 4-velocity  $u^\lambda$  satisfies  $u^\lambda u_\lambda = -1$  and, therefore, has only three independent components. In this case, the zeroth component of

Eq. (2)—the energy conservation—is not independent, but can be viewed as a consequence of the other four equations (1), (2). The latter four independent equations fully determine the evolution of  $n$  and three independent components of 4-velocity  $u^\lambda$ .

We notice that Eqs. (1)–(4) constitute the first-order hydrodynamics equations written in Landau frame. Namely, the constitutive relations (3), (4) are first order in derivatives and the ambiguity in the definition of 4-velocity is resolved by defining it as an eigenvector of the energy-momentum tensor. Landau frame was used in [1] and was adopted in [4] to construct the hydrodynamics with gauge anomaly.

The variational problem for hydrodynamics with gauge anomaly in  $1 + 1$  dimensions was successfully developed in [7], however it cannot be trivially generalized to  $3 + 1$  dimensions. The most successful attempt so far in finding an effective action for Eqs. (1)–(4) was given in [8], but the obtained action contained unphysical hydrodynamic excitations propagating in a fourth auxiliary spatial dimension. All these approaches rely on an effective action for the Lagrangian specification of fluid variables [9,10]. On the other hand, the action principle for non-Abelian hydrodynamics was presented in [6], where the authors introduced the idea of coarse graining the coadjoint orbit action. A similar approach to fluid dynamics for spinning particles has been recently developed in [11]. An action that includes anomalies in the standard model of particle physics within the framework of the coadjoint orbit method was given in [12]. The anomaly structure in the standard model is different from what is given in (1)–(4) and so the effective action for anomalies in [12] is not immediately applicable to the present problem.

In this work, we will use the so-called Clebsch potentials to parametrize the Eulerian variables [13] and to write down a variational principle that produces the Son-Surowka equations at zero temperature. We restrict ourselves to the flat Minkowski spacetime, though the generalization to more general geometric backgrounds is straightforward. Unless otherwise specified, we use the Cartesian orthonormal frame, where the pseudometric can be chosen as  $g_{\lambda\nu} = \text{diag}(-1, 1, 1, 1)$ .

The variational principle and the symmetries are analyzed in Secs. II and III. Using the obtained action, we then derive the corresponding Hamiltonian formulation specifying the form of the relativistic Hamiltonian and the Poisson brackets. We emphasize the symmetries of the system and their manifestations in Hamiltonian formalism, pointing out the special feature of one of the Clebsch potentials appearing separately and not via the combination in the dynamic velocity field. This feature is commented on in Sec. VII and we conclude with the discussion of the obtained results and their possible generalizations.

## II. HYDRODYNAMIC ACTION

The variational principle for perfect relativistic fluid dynamics is well known and goes back to [14–16]. The

key point in finding a hydrodynamic action is the introduction of a set of variables appropriate to the canonical framework, the so-called Clebsch potentials. The use of the Clebsch parametrization enlarges the phase space and removes the degeneracy of the Poisson algebra between hydrodynamic variables. The latter degeneracy of the Poisson's bracket makes the writing a symplectic form only in terms of hydrodynamic quantities impossible. The Clebsch potentials are scalar fields which parametrize the hydrodynamic variables, such as momentum and charge densities. Namely, we write the velocity one-form in terms of 3 scalar potentials  $(\theta, \alpha, \beta)$  and the chemical potential  $\mu$ , such that  $u = \mu^{-1}(d\theta + \alpha d\beta)$ , vide [15]. In the following we find an additional term in the hydrodynamic action of [15,16] reproducing the gauge anomaly in hydrodynamic equations.

The field content of the hydrodynamic action is given by 4 components of the 4-current  $J^\lambda$  and 3 scalar Clebsch potentials  $(\theta, \alpha, \beta)$  parametrizing dynamic velocity  $\xi_\lambda$ :

$$\xi_\lambda = \partial_\lambda \theta + \alpha \partial_\lambda \beta. \quad (5)$$

Then one of the main results of this work is that the action generating Eqs. (1)–(4) is given by:

$$S = - \int [J^\lambda (\xi_\lambda - A_\lambda) + \varepsilon(n)] d^4x + \frac{C}{6} \int A \wedge \xi \wedge d(\xi + A). \quad (6)$$

Here  $\varepsilon(n)$  is the proper energy density of the fluid which is assumed to be a known function of the proper charge density  $n$ . The latter is given by an absolute value of the 4-current  $J^\lambda$  as  $n \equiv \sqrt{-g_{\lambda\nu} J^\lambda J^\nu}$ . The second term on the right-hand side of (6) describes the anomaly and is written in the differential form language, that is,  $\xi = d\theta + \alpha d\beta$ . Taking  $C = 0$  in (6) we recover the action for a relativistic perfect fluid without anomaly [15,16].

The full set of variational equations is obtained by varying (6) over  $J^\lambda, \theta, \alpha, \beta$ . We start with:

$$\frac{\delta S}{\delta J^\lambda} = -(\xi_\lambda - A_\lambda) + \varepsilon'(n) \frac{J_\lambda}{n} = 0. \quad (7)$$

It is convenient to introduce a complete parametrization of the 4-current  $J^\lambda$  in terms of its absolute value  $n$  and its direction given by 4-velocity  $u^\lambda$  as:

$$J^\lambda \equiv n u^\lambda, \quad u^\lambda u_\lambda = -1. \quad (8)$$

Then Eq. (7) can be viewed as a relation between the dynamic velocity, density and the 4-velocity<sup>1</sup>:

<sup>1</sup>For the case of irrotational flows, such as superfluids, the dynamic velocity can be fully characterized by  $d\theta$  and Eq. (9) corresponds to the Josephson condition.

$$\xi_\lambda - A_\lambda = \mu u_\lambda, \quad (9)$$

where the chemical potential  $\mu(n)$  is given by the derivative of the energy density as:

$$\mu(n) \equiv \varepsilon'(n). \quad (10)$$

The Clebsch potentials  $\theta, \alpha, \beta$  enter (6) only through  $\xi$  given by (5). The corresponding variations give the following equations of motion:

$$\frac{\delta S}{\delta \theta} = \partial_\lambda \left( \frac{\delta S}{\delta \xi_\lambda} \right) = 0, \quad (11)$$

$$\frac{\delta S}{\delta \alpha} = \frac{\delta S}{\delta \xi_\lambda} \partial_\lambda \beta = 0, \quad (12)$$

$$\frac{\delta S}{\delta \beta} = \partial_\lambda \left( \alpha \frac{\delta S}{\delta \xi_\lambda} \right) = \frac{\delta S}{\delta \xi_\lambda} \partial_\lambda \alpha = 0, \quad (13)$$

with

$$-\frac{\delta S}{\delta \xi_\lambda} = nu^\lambda + \frac{C}{6} \epsilon^{\lambda\nu\eta\sigma} [2A_\nu \partial_\eta \xi_\sigma - (\xi_\nu - A_\nu) \partial_\eta A_\sigma]. \quad (14)$$

Introducing the charge current:

$$j^\lambda = -\frac{\delta S}{\delta \xi_\lambda} + \frac{C}{6} \epsilon^{\lambda\nu\eta\sigma} [3\partial_\nu (A_\eta \xi_\sigma) - 3A_\nu \partial_\eta A_\sigma + \xi_\nu \partial_\eta \xi_\sigma], \quad (15)$$

we obtain (1) from (11) and (5). The relations (15), (14) give the constitutive relation (3).

Defining the energy-momentum tensor by (4), one can derive the conservation law (2) from (9) and (11)–(13) after some tedious but straightforward manipulations.<sup>2</sup> We do not go through this derivation in more detail, since, in the next section, Sec. III, we will derive Eqs. (1)–(4) more straightforwardly from symmetries of the action (6).

In the absence of the gauge field background  $A_\mu = 0$  the action (6) becomes the conventional action for relativistic perfect fluid dynamics [15,16]. The only manifestation of the gauge anomaly in this case is the nonconventional relation between current and 4-velocity. Namely, the relation (3) becomes  $j^\lambda = nu^\lambda + \frac{C}{3} \mu^2 \omega^\lambda$  with relativistic vorticity defined as  $\omega^\lambda = \frac{1}{2} \epsilon^{\lambda\nu\sigma\tau} u_\nu \partial_\sigma u_\tau$ . This current is conserved  $\partial_\lambda j^\lambda = 0$  because both relations  $\partial_\lambda (nu^\lambda) = 0$  and  $\partial_\lambda (\mu^2 \omega^\lambda) = 0$  follow from (6) in the absence of the gauge background<sup>3</sup>—this consequence can be observed directly from [4] by setting the temperature and the external fields to zero. Such “removal” of the anomaly responses by

<sup>2</sup>Technical remark: it is convenient to start this derivation with an obvious equation  $\frac{\delta S}{\delta \xi_\lambda} [\partial_\lambda (\frac{\delta S}{\delta j^\lambda}) - \partial_\nu (\frac{\delta S}{\delta j^\nu})] = 0$ .

<sup>3</sup>One can think of the second relation as a consequence of (1), (2). We notice that the second conserved quantity  $\mu^2 \omega^\lambda$  can be identified as a density of the Casimir (helicity) of the relativistic perfect fluid dynamics.

current redefinition is not possible though when a nontrivial gauge field background is present.

### III. SYMMETRIES

In this section we show explicitly that Eqs. (1), (2) can be obtained as consequences of (anomalous) gauge symmetry and space-time translational symmetry of the action (6), respectively.

We notice that the first line of (6) is symmetric with respect to the gauge transformation with the gauge parameter  $\Lambda(x)$

$$\delta_\Lambda A_\lambda = \partial_\mu \Lambda, \quad \delta_\Lambda \theta = \Lambda. \quad (16)$$

Indeed, from (5), (16) we have  $\delta_\Lambda \xi_\lambda = \partial_\lambda \Lambda$  and see that the combination  $\xi_\lambda - A_\lambda$  entering (6) is gauge invariant.

This gauge invariance, however, is broken by the anomalous (second line) part of the action (6). It is easy to verify that, up to boundary terms, the gauge transformation of the action is given by

$$\delta_\Lambda S = \int \partial_\lambda \Lambda \left( \frac{\delta S}{\delta \xi_\lambda} + \frac{\delta S}{\delta A_\lambda} \right) d^4 x = \frac{C}{6} \int \Lambda dA \wedge dA. \quad (17)$$

Unlike the case of a general breaking of a symmetry, the loss of symmetry due to anomalies is rather special. The gauge variation of the action depends only on the background gauge field and has a very specific form, the latter being determined by the densities of certain topological invariants. It is easy to see that the action can be made fully gauge invariant by supplementing it with the Chern-Simons term  $-\frac{C}{6} \int_{M_5} A \wedge dA \wedge dA$ . The integral in this term is taken over an auxiliary 5-dimensional space  $M_5$  which boundary coincides with the physical space-time. This gives an elegant interpretation of the anomaly of the 4-dimensional theory as being due to the inflow of charge from the fifth dimension, a setup known as *anomaly inflow*; this is standard and well known in QFT with quantum anomalies [17]. With the variation with respect to the Clebsch potential  $\theta$  satisfying Eq. (11), the variation of (17) over  $\Lambda$  gives the charge conservation law modulo the anomaly as

$$\partial_\lambda \left( \frac{\delta S}{\delta A_\lambda} \right) = -\frac{C}{24} \epsilon^{\lambda\nu\sigma\tau} F_{\lambda\nu} F_{\sigma\tau}. \quad (18)$$

The quantity  $\delta S / \delta A_\lambda$  is known as the *consistent current* versus the *covariant current*  $j^\lambda$  defined in (3). A quick calculation shows that

$$j^\lambda = \frac{\delta S}{\delta A_\lambda} - \frac{C}{6} \epsilon^{\lambda\nu\sigma\tau} A_\nu F_{\sigma\tau}. \quad (19)$$

Taking the divergence of (19), we obtain (1).

We now turn to the energy-momentum conservation (4). The standard way of deriving this law is to gauge space-time translational symmetries by introducing the background metric and study the invariance of the action under diffeomorphisms  $x^\lambda \rightarrow x^\lambda + \zeta^\lambda(x)$ .

We consider (6) in an arbitrary background metric by replacing the measure  $d^4x$  by the invariant one  $\sqrt{-g}d^4x$  and by introducing the metric into all scalar products. Notice that  $\xi_\lambda$  is naturally a covariant vector, being derivatives of the scalar Clebsch potentials, and thus  $J^\lambda \xi_\lambda$  being an invariant scalar product does not require additional metric factors. However, a scalar product like  $J^2$  will become  $J^\mu J^\nu g_{\mu\nu}$ . The resulting action is invariant under diffeomorphisms, i.e.,  $\delta_\zeta S = 0$ , and on equations of motion we have

$$\int \left[ (\mathcal{L}_\zeta \mathbf{g})_{\nu\lambda} \frac{\delta S}{\delta g_{\nu\lambda}} + (\mathcal{L}_\zeta A)_\lambda \frac{\delta S}{\delta A_\lambda} \right] d^4x = 0, \quad (20)$$

since the terms corresponding to the variations of the fields vanish by the equations of motion. Here  $\mathcal{L}_\zeta$  denotes the Lie derivative with respect to the vector field  $\zeta$ . Explicitly

$$(\mathcal{L}_\zeta \mathbf{g})_{\nu\lambda} = \partial_\nu \zeta_\lambda + \partial_\lambda \zeta_\nu, \quad (21)$$

$$(\mathcal{L}_\zeta A)_\lambda = \zeta^\nu F_{\nu\lambda} + \partial_\lambda (\zeta^\nu A_\nu). \quad (22)$$

Using these formulas and setting the coefficient of  $\zeta^\nu$  in (20) to zero we obtain<sup>4</sup>

$$\partial_\lambda T^\lambda{}_\nu = F_{\nu\lambda} \frac{\delta S}{\delta A_\lambda} - \frac{C}{6} F_{\nu\lambda} \epsilon^{\lambda\eta\sigma\tau} A_\eta F_{\sigma\tau}, \quad (23)$$

with

$$T^{\lambda\nu} \equiv -\frac{2}{\sqrt{-g}} \frac{\delta S}{\delta g_{\lambda\nu}}. \quad (24)$$

A quick calculation shows that the energy-momentum tensor (24) is the same as (4). This is expected as the last term of (6) is the integral of a 4-form—which is metric-independent—and gives no contribution to the energy-momentum tensor. Therefore, (4) is identical in form to the energy-momentum tensor for conventional perfect fluid dynamics. We see that the metric independence of the anomalous contribution to (6) is an essential feature of the analysis in the hydrodynamic Landau frame where the energy-momentum tensor is not modified by corrections which are of the first order in gradients of the velocity.

Finally, it is easy to see that Eq. (23) with the relation (19) is equivalent to (2). This completes the demonstration that the action (6) does indeed reproduce Eqs. (1)–(4).

<sup>4</sup>The identity  $A_\nu \epsilon^{\lambda\eta\sigma\tau} F_{\lambda\eta} F_{\sigma\tau} = -4F_{\nu\lambda} \epsilon^{\lambda\eta\sigma\tau} A_\eta F_{\sigma\tau}$  can be useful.

#### IV. HAMILTONIAN FORMALISM

In this section we set up the Hamiltonian formulation of Eqs. (1)–(4) starting with the action (6).

We start by reducing the seven independent variational fields of (6) to four given by  $J^0$  and by the Clebsch parameters  $\theta, \alpha, \beta$ . The spatial components of (8), (9) give

$$J_i = \frac{n}{\mu} (\xi_i - A_i) \quad (25)$$

and we can eliminate the spatial components of the current  $J^i$  using (25). Using this relation and the defining relation (8) for  $n$ , namely,  $(J^0)^2 - (J^i)^2 = n^2$ , we find

$$J^0 \equiv \rho = \frac{n}{\mu} \sqrt{\mu^2 + (\xi_i - A_i)^2}. \quad (26)$$

Here and in the following we use  $\rho$  to denote  $J^0$ . We may regard  $J^0 = \rho$  as the independent variable, with  $n$  given implicitly as a function of  $\rho$  by (26).<sup>5</sup>

Substituting (25), (26) into (6) we obtain the action in a form linear in the time-derivatives and depending only on fields  $\rho, \theta, \alpha, \beta$ . After some integrations by parts, it can be brought to the following form:

$$S = \int (\langle \pi_\theta, \dot{\theta} \rangle + \langle \pi_\beta, \dot{\beta} \rangle - H) dt, \quad (27)$$

where  $\langle f, g \rangle \equiv \int f(x)g(x)d^3x$  denotes the  $L^2$ -inner product in the space of real functions,  $H$  is the Hamiltonian,  $\pi_\theta$  and  $\pi_\beta$  are the canonical field momenta conjugate to  $\theta$  and  $\beta$ , respectively. The explicit formulas for the canonical momenta are

$$\pi_\theta = -\left[ \rho + \frac{C}{6} (A_i + \alpha \partial_i \beta) B^i \right], \quad (28)$$

$$\pi_\beta = -\alpha \left[ \rho + \frac{C}{6} (A_i - \partial_i \theta) B^i \right]. \quad (29)$$

The Hamiltonian  $H$  in (27) is given by

$$H = \int \left[ \rho \sqrt{\mu^2 + (\xi_i - A_i)^2} - P(\mu) - \rho A_0 \right] d^3x - \frac{C}{6} \int [\xi_i B^i A_0 + \epsilon^{ijk} (\partial_i \theta - A_i) \xi_j E_k] d^3x. \quad (30)$$

The pressure  $P(\mu)$  is related to the energy density by the Legendre transform  $\varepsilon(n) = n\mu - P(\mu)$ , with  $P'(\mu) = n$  and we have also introduced the magnetic and electric fields  $B^i = \epsilon^{ijk} \partial_j A_k$  and  $E_i = \partial_i A_0 - \partial_0 A_i$  with  $\epsilon^{ijk} \equiv \epsilon^{0ijk}$ .

<sup>5</sup>As  $\mu(n)$  is assumed to be a known function of  $n$  (10) Eq. (26) can in principle be solved to obtain  $n(\rho, \xi_i)$ ,  $\mu(\rho, \xi_i)$ , etc.

Once again, we may note that if the anomaly vanishes, that is, for  $C = 0$ , the Hamiltonian formulation (28)–(30) reduces to the known Hamiltonian formulation for the perfect relativistic fluid [15,16,18,19]. We notice that in this case the Hamiltonian depends on Clebsch potentials only through  $\xi_i$ . This feature is lost in the presence of the anomaly, i.e., when  $C \neq 0$ , although the equations of motion (1)–(4) still do not contain the Clebsch potentials explicitly.

We shall comment on the meaning of this explicit dependence on  $\theta$  in the following sections. Here we just point out that the coefficient of  $E_k$  in the last term of (30) may be interpreted as an intrinsic electric dipole moment of the fluid. It is worth recalling that one of the main predictions of the anomaly for fluids is the chiral magnetic effect which leads to charge separation in a magnetic field. An electric dipole moment obviously suggests a charge separation and we may regard the last term of Eq. (30) as a reflection of this feature in the Hamiltonian framework.

So far we have considered the background gauge field as space- and time-dependent. An interesting special case is when the magnetic field is time-independent. It is then possible to choose a vector potential  $A_i$  which is independent of time as well. Then the last term of (30) can be integrated by parts and the Hamiltonian takes the form

$$H = \int \left[ \rho \sqrt{\mu^2 + (\xi_i - A_i)^2} - P(\mu) - A_0 \rho \right] d^3x - \frac{C}{6} \int A_0 [2\xi_i B^i + \epsilon^{ijk}(\xi_i - A_i) \partial_j \xi_k] d^3x. \quad (31)$$

In this case, the explicit dependence on  $\theta$  has disappeared and the Clebsch potentials only appear in the combination  $\xi_i$ . It is straightforward to observe that, for a time-independent gauge field, the potential term is simply  $\int A_0 j^0 d^3x$ , as one should expect.

In the next section we discuss the effect of the anomaly on the Poisson structure of the Hamiltonian formulation derived in this section.

## V. POISSON BRACKETS

The variational principle (27) which is linear in time-derivatives immediately provides us with the canonically conjugate pairs  $\theta, \pi_\theta$  and  $\beta, \pi_\beta$ . The Poisson brackets of all fields follow then from the canonical ones for the above fields

$$\{\theta, \pi'_\theta\} = \{\beta, \pi'_\beta\} = \delta(\mathbf{x} - \mathbf{x}'), \quad (32)$$

where we have listed only the nonvanishing Poisson brackets. Here and below we use a concise notation omitting the spatial arguments of the fields so that, e.g.,  $\beta$  means  $\beta(\mathbf{x})$ ,  $\pi'_\theta$  means  $\pi_\theta(\mathbf{x}')$ , etc.

The hydrodynamic equations of motion (1)–(4) can be formulated as equations written entirely in terms of  $\rho$  and  $\xi_i$

without an explicit dependence on the Clebsch parameters. Therefore, we shall look for the possible Hamiltonian reduction of (30), (32). The reduction consists of the dynamic reduction, i.e., the Hamiltonian should be expressible only in terms of the density  $\rho$  and dynamic velocity  $\xi_i$ , and the kinematic reduction, i.e., the closure of Poisson brackets of  $\rho$  and  $\xi_i$  without the use of the Clebsch parameters [20].

As we remarked before, with the inclusion of the anomaly, the dynamic reduction is only partially successful. Namely, the Hamiltonian (30) does depend on  $\partial_i \theta$  in the case of general time-dependent gauge field background. In the case of time-independent background the dynamic reduction is complete and the Hamiltonian (31) depends on the Clebsch parameters only through  $\xi_i$ .

Remarkably, the Poisson algebra of  $\rho$  and  $\xi_i$  is closed for any gauge field background so that the kinematic reduction is achieved. Indeed, after some straightforward calculations, we derive from (32) and the definition (5) the following set of Poisson brackets closed with respect to the fields  $\rho$  and  $\xi_i$ ,

$$\{\rho_+, \rho'_+\} = \frac{C}{3} B^i \partial_i \delta(\mathbf{x} - \mathbf{x}'), \quad (33)$$

$$\{\tilde{\xi}_i, \rho'_+\} = \partial_i \delta(\mathbf{x} - \mathbf{x}'), \quad (34)$$

$$\{\tilde{\xi}_i, \tilde{\xi}'_j\} = -\frac{\partial_i \tilde{\xi}_j - \partial_j \tilde{\xi}_i - \epsilon_{jik} B^k}{\rho_-} \delta(\mathbf{x} - \mathbf{x}'). \quad (35)$$

Here, for the sake of brevity, we introduced the following compact notation,

$$\tilde{\xi}_i \equiv \xi_i - A_i, \quad (36)$$

$$\rho_\pm \equiv \rho \pm \frac{C}{6} \tilde{\xi}_i B^i. \quad (37)$$

A comment on the first of these equations, namely, (33), is appropriate at this point. It is well known that the  $[j^0, j^{0i}]$  commutator will be modified by a Schwinger term in the presence of an anomaly for the corresponding symmetry [21]. This can be shown by explicit computation of the corrections to commutators via Feynman diagrams, the triangle diagram leading to the specific form given.<sup>6</sup> It can also be seen from a 2-cocycle constructed in terms of the descent equations which lead to the anomalies [22]. Our action effectively reproduces this in the Poisson brackets. We may also note that an expression analogous to (33) has appeared in [23].

We remark here that the dynamic velocity  $\tilde{\xi}_i$  and the modified densities  $\rho_\pm$  are invariant under the

<sup>6</sup>The computation of modified commutators follows a procedure known as the Bjorken-Johnson-Low method where correlators of currents at slightly unequal times are calculated and a suitable equal-time limit is taken.

transformations (16), therefore, the Poisson algebra (33)–(35) is written in terms of explicitly gauge-invariant quantities. However, as it is well known [22] that the generator of gauge transformations cannot be realized canonically in the presence of anomaly (see Sec. VI).

The algebra (33)–(35) is obtained as a result of Hamiltonian reduction and is degenerate. It admits two Casimirs—the quantities having vanishing Poisson brackets with fields entering Poisson algebra. They are given by

$$C_1 = \int \rho_+ d^3x, \quad (38)$$

$$C_2 = \int \epsilon^{ijk} \tilde{\xi}_i \partial_j (\tilde{\xi}_k + 2A_k) d^3x. \quad (39)$$

The charge density  $j^0$  defined in (3) is given by

$$j^0 = \rho_+ + \frac{C}{6} \epsilon^{ijk} \tilde{\xi}_i \partial_j (\tilde{\xi}_k + 2A_k). \quad (40)$$

It is a combination of densities of two Casimirs of the algebra. It is worthwhile to point out that when the gauge field is time-independent the anomaly term can be written as a total derivative, i.e.,  $E_i B^i = \partial_i (A_0 B^i)$ , what automatically implies that the total charge is indeed conserved.

In the absence of anomaly  $C = 0$ , all expressions (30), (33)–(40) become the known formulas for perfect fluid dynamics [5,6]. Even when the anomaly is present, i.e.,  $C \neq 0$ , if we consider the case of the background gauge field being absent, we obtain again the formulas of anomaly-free hydrodynamics with a single exception. Namely, the definition of the charge density (40) still differs from  $\rho$  by the density of Casimir (39). The latter is known as the helicity of the hydrodynamic flow.

Having Hamiltonian and Poisson brackets one can obtain equations of motion for any quantity  $Q$  as  $\dot{Q} = \partial Q / \partial t + \{H, Q\}$ , where  $\partial Q / \partial t$  denotes the “explicit” time-derivative. In our case this explicit derivative acts only on the time varying external gauge field. The dynamical fields  $\xi_i$  and  $\rho$  do not depend on time explicitly. For example, the equation of motion for  $\xi_i$  will read  $\dot{\xi}_i = -\partial_t A_i + \{H, \xi_i\}$ , etc.

While the Clebsch variables appear in the algebra (33)–(35) only via  $\xi_i$ , we should note that, in the presence of the time-dependent gauge field background, the Hamiltonian (30) contains  $\partial_t \theta$  in addition to the density and the dynamic velocity fields. Thus the algebra (33)–(35) is not adequate for a complete Hamiltonian description, and it should be supplemented by Poisson brackets involving the  $\theta$  field. We list those brackets here for completeness

$$\{\rho_+, \partial_k \theta'\} = \partial_k \delta(\mathbf{x} - \mathbf{x}'), \quad (41)$$

$$\{\tilde{\xi}_i, \partial_k \theta'\} = \frac{\tilde{\xi}_i + A_i - \partial_i \theta}{\rho_-} \partial_k \delta(\mathbf{x} - \mathbf{x}'). \quad (42)$$

## VI. SYMMETRY GENERATORS

The Poisson algebra (33)–(35) is closed and, in the case of the time-independent background, produces the hydrodynamic equations with the use of the Hamiltonian (31). However, the brackets (33)–(35) are nonlinear and therefore do not have the Lie-Poisson form. For the symmetry analysis it is preferable to find an equivalent set of Poisson brackets corresponding to the algebra of symmetry generators of the system.

It is easy to see from (27) that the momentum densities can be defined as:

$$\Theta_{0i} = -\pi_\theta \partial_i \theta - \pi_\beta \partial_i \beta. \quad (43)$$

The momentum densities  $\Theta_{0i}$  satisfy the diffeomorphism algebra and act as local translations in the absence of background field. However, one cannot express (43) only in terms of the density  $\rho$  and the dynamic velocity in the background of nonvanishing magnetic field. More precisely, the canonical energy-momentum tensor acquires an explicit  $\theta$  dependence:

$$\Theta_{0i} = \left( \rho + \frac{C}{6} A_k B^k \right) \xi_i + \frac{C}{6} B^k (\xi_k \partial_i \theta - \xi_i \partial_k \theta). \quad (44)$$

Let us now turn to gauge transformations which can be viewed as shifts in the field  $\theta$ . The naive canonical gauge generator for this symmetry is  $-\pi_\theta$ . Using (28), (5) we can write it as

$$-\pi_\theta = \rho + \frac{C}{6} (A_i + \xi_i - \partial_i \theta) B^i \quad (45)$$

and notice that it also depends explicitly on  $\partial_i \theta$ .

It is straightforward to check that the Poisson structure (33)–(35) can be put in a semidirect product Lie-Poisson algebra [18,19] in terms of (44), (45).

A gauge transformation of an arbitrary functional  $F$  of basic fields generated by  $-\pi_\theta$  is given by:

$$\delta_\Lambda F \equiv \int \left( -\Lambda(\mathbf{x}') \{ \pi'_\theta, F \} + \frac{\delta F}{\delta A_i(\mathbf{x}')} \partial'_i \Lambda \right) d^3x', \quad (46)$$

where the transformation of the gauge potential has also been added.

However, it is easy to see that (46) gives  $\delta_\Lambda \alpha \neq 0$ , as well as  $\delta_\Lambda \rho \neq 0$  in apparent contradiction with gauge invariance of  $\alpha$  and  $\rho$ . In fact, one can show that the gauge symmetry (16) is not canonically realizable.

Let us now consider  $\rho_+$  given by (37) as a generator of gauge transformations instead of  $-\pi_\theta$ . We easily check that  $\delta_\Lambda \alpha = \delta_\Lambda \beta \equiv 0$  and  $\delta_\Lambda \theta \equiv \Lambda$ . Moreover, under the modified gauge transformations generated by  $\rho_+$  the density  $\rho$  transforms as:

$$\delta_{\Lambda}\rho = -\frac{C}{6}B^i\partial_i\Lambda, \quad (47)$$

and there exists the gauge invariant quantity  $\rho + \frac{C}{6}B^iA_i$ .

While  $\rho_+$  can be considered as a modified generator of gauge transformations two subsequent gauge transformations generated by  $\rho_+$  do not commute and the commutative algebra of gauge transformations has acquired a central extension (33). This is, of course, a classical manifestation of a well-known phenomenon in studies of quantum anomalies [22]. At this point it is not clear whether similar modifications can be made for diffeomorphism generators (43).<sup>7</sup>

## VII. CONCLUSION AND DISCUSSION

We have presented a variational principle for hydrodynamic equations with gauge anomaly at zero temperature. From the obtained action, we derived the Poisson structure and the Hamiltonian for the system. The most noteworthy feature of the obtained Hamiltonian formulation is that in the presence of gauge anomaly, the Hamiltonian reduction to the density and velocity fields is not complete and one of the Clebsch potentials becomes physical and is present in the Hamiltonian in the presence of the time-dependent gauge field background.

The case of the time-independent external gauge fields is more natural for the Hamiltonian formulation. In this case one has a complete Hamiltonian reduction with both Hamiltonian and Poisson brackets expressed purely in terms of the charge density  $\rho$  and dynamic velocity  $\xi_i$ .

<sup>7</sup>The variables  $\alpha$  and  $\rho$  do not transform nicely under these generators.

It turns out, however, that the generators of gauge transformations  $\rho_+$  (37) cease to commute and that the generators of spatial translations (44) can be written only with the explicit use the Clebsch potential  $\theta$ . The origin of the explicit appearance of  $\theta$  in the Hamiltonian and in (45) and (43) can be traced to the term  $A \wedge \xi \wedge d\xi = A \wedge d\theta \wedge d\alpha \wedge d\beta$  in the action. This term is needed in the hydrodynamic action to make sure that the anomalous nonconservation of the charge corresponds to the one of the underlying QFT and following from the computation of the triangle diagram. In our variational approach the presence of  $A \wedge \xi \wedge d\xi$  term does not lead to any entropy production and is in agreement with the requirement of positive semidefinite entropy production which was central in Son and Surowka analysis [4]. A connection of the anomalous term with the entropy arguments might become more explicit if the variational principle could be generalized to finite temperature hydrodynamics. A possibility of such a generalization is worthy of further investigation.

## ACKNOWLEDGMENTS

We would like to acknowledge the Simons Center for Geometry and Physics for the support of the program on quantum anomalies in hydrodynamics in Spring 2014 during which this work has been done. We are grateful to V. Cheianov and B. Khesin and other participants of that program for multiple discussions and suggestions related to this work. The work of A. G. A. was supported by the NSF under Grant No. DMR-1206790 and the work of V. P. N. was supported by NSF Grant No. PHY-1213380 and by PSC-CUNY awards.

- 
- [1] L. Landau and E. Lifshitz, *Fluid Mechanics* (Butterworth-Heinemann, Washington, DC, 1987).
  - [2] J. Erdmenger, M. Haack, M. Kaminski, and A. Yarom, Fluid dynamics of R-charged black holes, *J. High Energy Phys.* **01** (2009) 055.
  - [3] N. Banerjee, J. Bhattacharya, S. Bhattacharyya, S. Dutta, R. Loganayagam, and P. Surowka, Hydrodynamics from charged black holes, *J. High Energy Phys.* **01** (2011) 094.
  - [4] D. T. Son and P. Surowka, Hydrodynamics with Triangle Anomalies, *Phys. Rev. Lett.* **103**, 191601 (2009).
  - [5] V. E. Zakharov and E. Kuznetsov, Hamiltonian formalism for nonlinear waves, *Usp. Fiz. Nauk* **167**, 1137 (1997).
  - [6] R. Jackiw, V. P. Nair, S.-Y. Pi, and A. P. Polychronakos, Perfect fluid theory and its extensions, *J. Phys. A* **37**, R327 (2004).
  - [7] S. Dubovsky, L. Hui, and A. Nicolis, Effective field theory for hydrodynamics: Wess-Zumino term and anomalies in two spacetime dimensions, *Phys. Rev. D* **89**, 045016 (2014).
  - [8] F. M. Haehl, R. Loganayagam, and M. Rangamani, Effective actions for anomalous hydrodynamics, *J. High Energy Phys.* **03** (2014) 034.
  - [9] S. Dubovsky, L. Hui, A. Nicolis, and D. T. Son, Effective field theory for hydrodynamics: Thermodynamics and the derivative expansion, *Phys. Rev. D* **85**, 085029 (2012).
  - [10] A. Nicolis and D. T. Son, Hall viscosity from effective field theory, [arXiv:1103.2137](https://arxiv.org/abs/1103.2137).
  - [11] D. Karabali and V. P. Nair, Relativistic particle and relativistic fluids: Magnetic moment and spin-orbit interactions, *Phys. Rev. D* **90**, 105018 (2014).
  - [12] V. P. Nair, R. Ray, and S. Roy, Fluids, anomalies and the chiral magnetic effect: A group-theoretic formulation, *Phys. Rev. D* **86**, 025012 (2012).

- [13] A. Clebsch, Über die integration der hydrodynamischen gleichungen, *J. Reine Angew. Math.* **1859**, 1 (1859).
- [14] C. C. Lin, in *International School of Physics E. Fermi (XXI)*, edited by G. Careri (Academic Press, New York, 1963).
- [15] B. F. Schutz, Perfect fluids in general relativity: Velocity potentials and a variational principle, *Phys. Rev. D* **2**, 2762 (1970).
- [16] B. F. Schutz, Hamiltonian theory of a relativistic perfect fluid, *Phys. Rev. D* **4**, 3559 (1971).
- [17] C. G. Callan Jr. and J. A. Harvey, Anomalies and fermion zero modes on strings and domain walls, *Nucl. Phys.* **B250**, 427 (1985); see also R. Bertlmann, *Anomalies in Quantum Field Theory*, International Series of Monographs on Physics 91 (Clarendon-Oxford University Press, Oxford, England, 1996).
- [18] D. D. Holm and B. A. Kupershmidt, Relativistic fluid dynamics as a Hamiltonian system, *Phys. Lett.* **101A**, 23 (1984).
- [19] J. E. Marsden, T. Ratiu, and A. Weinstein, Semidirect products and reduction in mechanics, *Trans. Am. Math. Soc.* **281**, 147 (1984).
- [20] R. L. Seliger and G. Whitham, Variational principles in continuum mechanics, *Proc. R. Soc. A* **305**, 1480 (1968).
- [21] R. Jackiw and K. Johnson, Anomalies of the axial-vector current, *Phys. Rev.* **182**, 1459 (1969); S. L. Adler and D. G. Boulware, Anomalous commutators and the triangle diagram, *Phys. Rev.* **184**, 1740 (1969); R. A. Bertlmann and T. Sykora, Point-splitting method of the commutator anomaly of Gauss law operators, *Phys. Rev. D* **56**, 2236 (1997) and references therein; see also *Current Algebra and Anomalies*, edited by S. Treiman, R. Jackiw, B. Zumino, and E. Witten (World Scientific, Singapore, 1985).
- [22] L. Faddeev, Operator anomaly for the Gauss law, *Phys. Lett.* **145B**, 81 (1984); J. Mickelsson, Chiral anomalies in even and odd dimensions, *Commun. Math. Phys.* **97**, 361 (1985); L. Faddeev and S. Shatashvili, Algebraic and Hamiltonian methods in the theory of non-Abelian anomalies, *Theor. Math. Phys.* **60**, 770 (1984); see also the book by S. Treiman *et al.* in [21].
- [23] A. Alekseev, V. Cheianov, and J. Frohlich, Universality of Transport Properties in Equilibrium, the Goldstone Theorem and Chiral Anomaly, *Phys. Rev. Lett.* **81**, 3503 (1998).