Nonleptonic decays of charmed mesons into two pseudoscalars

Aritra Biswas and Nita Sinha

The Institute of Mathematical Sciences, C.I.T. Campus, Tharamani, Chennai 600 113, India

Gauhar Abbas

IFIC, Universitat de València-CSIC, Apt. Correus 22085, E-46071 València, Spain

(Received 21 April 2015; published 28 July 2015)

We examine the role of resonant coupled channel final state interactions (FSIs), as well as weak annihilation and exchange contributions, in explaining all the two-body hadronic $D \rightarrow PP$ decay modes. In the un-unitarized amplitudes we include modified Wilson coefficients with nonfactorizable corrections as parameters. For the hadronic form factors, the z-series expansion method is used to get the q^2 dependence. The FSI effects are incorporated via a phenomenological approach with widths of resonances to various channels taken from observations where available, and others as additional parameters to be determined from fits of all the theoretical rates to the measured ones. Our results for the rather hard to explain $D^0 \rightarrow K^+K^-, \pi^+\pi^-$ are in agreement with measured values. We demonstrate that both weak exchange and FSI effects are required to get the correct branching ratio for the $D^0 \rightarrow K^0 \bar{K^0}$ mode. Using our unitarized amplitudes we evaluate the strong phase difference between the amplitudes for $D^0 \rightarrow K^-\pi^+$ and $D^0 \rightarrow K^+\pi^-$ and find it to be in complete agreement with the recent BES III result.

DOI: 10.1103/PhysRevD.92.014032

PACS numbers: 13.25.Ft

I. INTRODUCTION

With the advent of the B factories, along with even the Tevatron having sufficient luminosities to perform excellent heavy flavor studies, and more recently with the data pouring in from LHCb and BESIII, quark flavor physics is changing its role from being the ground for confirming the Kobayashi-Maskawa mechanism for CP violation to providing the observables that can test or constrain physics beyond the Standard Model. With improved precision measurements, the goals of hadronic weak decays will cease to be seeking observables free from hadronic uncertainties and instead become actually precisely understanding the QCD effects.

Along with the plethora of data on semileptonic and hadronic rare decays, CP asymmetries, and mixing in B decays, the discovery of charm mixing and hints of CP violation in the charm sector resulted in nonleptonic charm meson decays being a focus of attention in the last few years [1,2]. Since CP violation in charm, direct, and mixing is expected to be negligible, any hint of CP violation in charmed mesons is expected to be from physics beyond the Standard Model and hence charm may be instrumental in providing insights into new physics [3–11]. In fact, earlier FOCUS [12], CLEO [13], and various other collaborations had produced many interesting results in the charm sector. The 3.2 σ hint [14,15] of a difference of *CP* asymmetries between the singly suppressed K^+K^- and $\pi^+\pi^-$ charmed decay modes resulted in a large volume of work [16-32], mostly using different models of new physics to explain the result. The hint has since then been slowly moving towards zero and currently there seems to be no evidence for any

direct *CP* violation in charm in any mode [33]. Even though this hint for the *CP* asymmetry slowly disappeared, from all the recent work done on the charm decays to the two pseudoscalar modes motivated by this hint, it was clear that it is critical to first understand the observed branching ratios of all the charmed hadronic decay modes well, within the Standard Model, before any observation of an anomalous rate or any new *CP* asymmetry can be claimed as due to the presence of new physics.

However, this is not an easy task. The mass of the charm quark (1.275 GeV) makes it very difficult to come up with a proper theoretical technique for calculation of hadronic charmed meson decays. The charm, unlike the bottom quark, is not sufficiently heavy to allow realization of the infinitely heavy quark limit. Therefore, the well-known theoretical approaches based on QCD, for example, heavy quark effective theory [34,35], QCD factorization [36,37], the perturbative QCD approach [38–41], and the soft-collinear effective theory [42], which lead to very satisfactory predictions for *B* decays, cannot be used to explain data in the case of charmed mesons. Furthermore, the charm quark is also not light enough for a chiral expansion to be applicable.

In the absence of any other reliable and effective theoretical methods, the factorization approach is still one of the most successful ways to study two-body charm meson decays [43,44]. However, it is well known now that in the naive factorization approach, calculation of Wilson coefficients of effective operators faces the problem of γ_{5} - and renormalization scheme dependence. These difficulties can be overcome in the framework of the

"generalized factorization approach" where Wilson coefficients are effective and include important nonfactorizable (NF) corrections [45,46].

In the past there was another attempt to explain hadronic D decays using the so-called large $1/N_c$ (where N_c is number of color degrees of freedom) approach [47]. It was observed that dropping Fierz transformed terms characterized by $1/N_c$ can narrow the gap between predictions and observations up to a satisfactory level. The calculations based on QCD sum rules showed that Fierz terms were certainly compensated by the NF corrections [48–50].

There exists another model-independent, so-called "quark diagram" or "topological diagram" approach in the literature [51–59], where all two-body nonleptonic weak decays of heavy mesons are expressed in terms of distinct quark diagrams, depending on the topologies of weak interactions, including all strong interaction effects. It is based on SU(3) symmetry and allows extraction of the quark diagram amplitudes by fitting against experimental data. However SU(3) breaking effects in charmed meson decays have been shown to be important and need to be carefully incorporated [60–62].

The importance of final state interactions (FSIs) in nonleptonic charm decays had been realized and discussed in several papers [63–66] in the early 1980s, where the authors had been intrigued by the anomalies in the observed branching ratios of the Cabibbo favored (CF) neutral modes versus the charged $K\pi$ modes, and the differing rates of the singly Cabibbo suppressed (SCS) K^+K^- and $\pi^+\pi^-$ modes, followed by measurements of rates of a few other modes that had unexpected suppression/enhancement. Many of these were conjectured to be due to FSI. Surprisingly, even in the last couple of years, in many of the papers that worried about the charm *CP* asymmetry problem, these old puzzles were still considered unresolved.

Even for the case of hadronic B meson decays, the role of FSIs has been examined rather carefully in the last few years [67,68]. The mass of the charmed meson lies right in the heart of the resonance region. Hence, resonant final state rescattering is bound to play a bigger role in the twobody hadronic charm decays and needs to be evaluated. Of course dynamical calculations of these long distance effects are not possible and hence they can only be determined phenomenologically after comparison of the theoretical estimates with experimental data. Unitarity constraints play an important role in providing the theoretical estimates.

Another contribution in hadronic two-body decays that has been debated for a long period is that of the weak annihilation and exchange diagrams. One of Rosen's proposals [69] was that the W-exchange diagrams may be large and since this appears only in D^0 and not in D^+ decays, it could account for the difference in the lifetimes of these two mesons. Bigi and Fukugita [70] then proposed several D and B meson decay modes that could be the smoking gun signals of the W-exchange contributions and yet, when the mode $D^0 \rightarrow \phi \bar{K}^0$ was observed, it was argued [71] that it could have been generated from the decay mode $D^0 \rightarrow K^* \eta$, with this final state rescattering to the $\phi \bar{K}^0$ mode. Annihilation type contributions along with FSIs were incorporated in the hadronic two-body vectorpseudoscalar modes of charmed meson decays in Ref. [72]. Studies using the quark diagram approach of Ref. [73] had also indicated that annihilation type contributions are needed to explain the observed data.

In this paper, we study the role of FSI in the two-body D(*D* here can be any of the D^0 , D^+ , or D_s^+) meson decays. We assume that FSI effects are dominated by resonance states close to the mass of D mesons. In fact, there exist isospin 0, 1, and 1/2 resonances near the D mass, that may contribute to rescatterings among different channels in these respective isospin states and enhance/suppress some of the decay rates. In the next section, we give the formalism for the calculation of the un-unitarized amplitudes, using a modified factorization approach, where the effective Wilson coefficients include NF corrections. This is in analogy with the QCD factorization approach of Beneke-Neubert for hadronic *B* meson decays [74], where however, using the hard scattering approach, the NF corrections are calculable in heavy quark approximation. However, for charm, since this approximation fails, these NF corrections are not calculable and are left as parameters. We also indicate our parametrization of the annihilation contributions and discuss our inputs: the decay constants and the form factors, for which we have used a z-series expansion approach. In Sec. III, we discuss the need to incorporate additional long distance FSI effects and show how this can be done with a K matrix formalism for coupled channels. Using the observed widths, masses, and known decay rates of the resonances to the various channels to evaluate the diagonal elements and leaving the unknown elements of the K matrix as parameters, the unitarized amplitudes are calculated (as discussed in [72]) to estimate the branching ratios of all the SCS, CF, and doubly Cabibbo suppressed (DCS) $D \rightarrow PP$ decay modes. In Sec. IV, we list the isospin decomposition of all the decay modes, the parameters that need to be determined from our fits, and the errors in theoretical inputs used. We list all the branching ratios after our numerical χ^2 fits and the values of the fitted parameters. Finally we conclude in Sec. V.

II. THE UN-UNITARIZED AMPLITUDES

A. Weak Hamiltonian and Wilson coefficients

The study of weak decays of charmed mesons to twobody hadronic modes necessarily requires a careful evaluation of the strong interaction corrections. The weak effective Hamiltonian may be expressed in terms of coefficient functions, which incorporate the strong interaction effects above the scale $\mu \sim m_c$ and the currentcurrent operators as NONLEPTONIC DECAYS OF CHARMED MESONS INTO TWO ...

$$\mathcal{H}_{w} = \frac{G_{F}}{\sqrt{2}} [C_{1}(\mu)O_{1}(\mu) + C_{2}(\mu)O_{2}(\mu)] + \text{H.c.} \quad (1)$$

where G_F is the Fermi coupling constant, C_1 and C_2 are the Wilson coefficients, and the operators are

$$O_1 = (\bar{u}_{\alpha}q_{2\alpha})_{V-A}(\bar{q}_{1\beta}c_{\beta})_{V-A}$$
$$O_2 = (\bar{u}_{\alpha}q_{2\beta})_{V-A}(\bar{q}_{1\beta}c_{\alpha})_{V-A}.$$

 α and β in the above are color indices, while q_1, q_2 can be either the *d* or the *s* quark. The quark diagrams dominantly contributing to the branching ratios of $D \rightarrow P_1P_2$ [59] are the color-favored tree amplitude *T*, the color-suppressed amplitude *C*, the W-exchange amplitude *E*, and the W-annihilation amplitude *A*, shown in Fig. 1.

Penguin contributions in charmed meson decays are highly suppressed as the dominant down type quark contribution to the flavor changing neutral current $c \rightarrow u$ transition is from the *b* quark which is accompanied by the presence of the tiny product, $V_{cb}^*V_{ub}$, of the Cabibbo– Kobayashi–Maskawa (CKM) matrix elements. Hence, the two operators in Eq. (1) are sufficient for calculating the amplitudes and branching ratios of the $D \rightarrow PP$ modes.

In the naive factorization approach, the matrix element of the four-fermion operator in the heavy quark decay is replaced by a product of two currents. The amplitudes for the nonleptonic two-body decay modes are then the product of a transition form factor and a decay constant. However, NF corrections must exist; while such corrections for scales larger than μ are taken into consideration in the effective weak Hamiltonian, those below this scale also need to be carefully incorporated. In the QCD factorization approach for *B* meson decays [36,37,75], these NF corrections are handled using the hard scattering approach, where the vertex corrections and the hard spectator interactions are



FIG. 1 (color online). The dominant quark diagram amplitudes.

added at the next to leading order in α_s and its accuracy is limited only by the corrections to the heavy quark limit. But, in the case of charm decays, where the heavy quark expansion is not a very good approximation, it is best to parametrize these NF corrections and then determine them by fitting the theoretical branching ratios with the experimental data. In the diagrammatic approach of Ref. [59] also, either the Wilson coefficients themselves or the NF corrections appearing in the Wilson coefficients are determined from fits to data.

Hence, we write the scale dependent Wilson coefficients, modified to include the NF corrections which are parametrized by χ_1 and χ_2 with their respective phases ϕ_1 and ϕ_2 as

$$a_1(\mu) = C_1(\mu) + C_2(\mu) \left(\frac{1}{N_c} + \alpha(\mu)\chi_1 e^{i\phi_1}\right)$$
(2)

$$a_{2}(\mu) = C_{2}(\mu) + C_{1}(\mu) \left(\frac{1}{N_{c}} + \alpha(\mu)\chi_{2}e^{i\phi_{2}}\right).$$
(3)

The dominant tree and color amplitudes for $D \rightarrow P_1 P_2$, where P_1 is the final meson which carries the spectator quark, while P_2 represents the meson emitted from the weak vertex (as depicted in Fig. 1), are then written as

$$T(C) = \frac{G_F}{\sqrt{2}} V_{\text{CKM}} a_1(\mu) (a_2(\mu)) f_{P_2}(m_D^2 - m_{P_1}^2) F_0^{DP_1}(m_{P_2}^2),$$
(4)

where f_{P_2} is the P_2 meson decay constant and $F_0^{DP_1}(m_{P_2}^2)$ denotes the transition form factor for $D \to P_1$ evaluated at $m_{P_2}^2$. We follow the prescription of Ref. [76] and choose the scale μ to be the energy release in individual decay processes rather than fixed at m_c . This scale, which is dependent on the final state masses, allows for SU(3) breaking, in addition to that coming from different decay constants and form factors. This scale is taken to be $\mu = \sqrt{\Lambda m_D(1 - r_2^2)}$, where $r_2^2 = m_{P_2}^2/m_D^2$ and Λ is another free parameter. Λ , χ_1 , χ_2 , ϕ_1 , and ϕ_2 are taken to be universal for all the decay modes and are fitted from experimental data.

B. Weak annihilation contributions

For a long time, W-exchange and W-annihilation contributions used to be neglected due to the so-called helicity suppression. However, observation of many decay modes of charmed and bottom mesons, which is possible only via the annihilation or exchange diagrams, has indicated that these contributions could be substantial. These short distance weak annihilation effects were hence included in the diagrammatic approach. In principle these could result from rescattering, even in the absence of annihilation and exchange processes. In fact, weak annihilation

topologies were assumed to be induced by nearby resonances through FSIs in Ref. [73]. The authors in [73] as well as Ref. [77] use SU(3) to relate the couplings of the final state mesons with the resonances. This leads to the result that the long distance W-exchange contribution can be induced by a tree amplitude, while W-annihilation can be induced by a color-suppressed internal W-emission. The resonant FSIs modify the W-exchange and W-annihilation amplitudes but the T and C amplitudes are unaffected. We emphasize that the assumption of SU(3) plays an important role in these results. In a most general coupled channel formalism, all contributions in the various channels will be affected by the resonant FSIs, as will be shown in Sec III. We parametrize the W-exchange and W-annihilation contributions in the amplitudes by $\chi^{E(A)}$ and estimate them from phenomenological fits to data. The exchange (annihilation) amplitudes are hence written as

$$E_{q,s}(A_{q,s}) = \frac{G_F}{\sqrt{2}} V_{\text{CKM}} C_1(\mu) (C_2(\mu)) \chi_{q,s}^{E(A)} \frac{C_F}{N_c^2} f_D f_{P_1} f_{P_2}.$$
(5)

Since the initial charmed meson is annihilated and both the final mesons are produced from the weak vertex, after the production of a quark-antiquark pair from a gluon, these amplitudes are a product of the decay constants of the initial D meson (f_D) and that of $P_1(f_{P_1})$ and $P_2(f_{P_2})$. Apart from this, the strengths of the exchange (annihilation) amplitudes $\chi_{q,s}^{E(A)}$ are assumed to be the same for all modes and the subscripts distinguish the contributions of the pair production of the light quark-antiquark from that of the strange pair. Since the annihilation and exchange contributions are necessarily nonfactorizable, they depend only on $C_{1,2}$ rather than the modified coefficients $a_{1,2}$.

We wish to emphasize that for the case of B mesons decaying to two pseudoscalar mesons, it has been shown [74] that the nonvanishing annihilation contribution at order α_s arises only for gluons emitted from the initial state quarks; on the other hand, the resonant FSIs (to be discussed in Sec. III) will necessarily result from gluons emitted from final state quarks. Hence, assuming that a similar distinction will hold for *D* decays as well, adding FSIs when annihilation contributions are already added should not amount to double counting, as pointed out in Ref. [78].

The scale of the Wilson coefficients for the exchange and annihilation amplitudes must depend on both the mass ratios, $r_{1,2} = m_{P_{1,2}}/m_D$, and is taken to be $\mu = \sqrt{\Lambda m_D (1 - r_1^2)(1 - r_2^2)}$.

C. Nonperturbative inputs: Form factors and decay constants

We start by specifying our convention for the different mesons involved in our analysis:

$$\begin{aligned} \pi^{+} &= -u\bar{d}, & \pi^{-} = d\bar{u}, & \pi^{0} = \frac{u\bar{u} - dd}{\sqrt{2}}, \\ K^{0} &= d\bar{s}, & \bar{K^{0}} = -s\bar{d}, & K^{+} = u\bar{s}, & K^{-} = s\bar{u}, \\ D^{0} &= c\bar{u}, & D^{+} = -c\bar{d}, & D^{+}_{s} = c\bar{s}. \end{aligned}$$

In the $D \rightarrow P_1$ transitions, the matrix element of the vector current is written in terms of the form factors F_+ and F_0 as

$$\begin{split} \langle P_1(p') | \bar{q} \gamma^{\mu} c | D(p) \rangle &\equiv F_+(q^2) \left(p^{\mu} + p'^{\mu} - \frac{m_D^2 - m_{P_1}^2}{q^2} q^{\mu} \right) \\ &+ F_0(q^2) \frac{m_D^2 - m_{P_1}^2}{q^2} q^{\mu}, \end{split} \tag{6}$$

where $q \equiv p - p'$. The matrix element for the production of the second meson P_2 is given by

$$\langle P_2(q)|\bar{q_1}\gamma_\mu q_2|0\rangle = if_{P_2}q_\mu. \tag{7}$$

Hence, in the amplitude of the nonleptonic two pseudoscalar decay modes of charmed mesons involving the product of the two matrix elements specified in Eqs. (6) and (7), only the transition form factor F_0 appears. Transition form factors can in principle be experimentally measured from the semileptonic decays; however, in the massless lepton limit, only the $F_+(q^2)$ contributes to the semileptonic amplitude distributions. However, the semileptonic information is still useful, since at zero momentum transfer the form factors obey the kinematic constraint $F_0(0) = F_+(0)$. The q^2 dependence of the F_0 on the other hand is accessible only with massive leptons in the semileptonic decays or in lattice simulations. Simple and modified pole models have been widely used to parametrize the q^2 dependence of the form factors, but these have poor convergence properties. Recently the zexpansion [79,80] has been introduced as a model independent parametrization of the q^2 dependence of form factors over the entire kinematic range and has been shown to have improved convergence properties. In this approach, based on analyticity and unitarity, the form factors are expressed as a series expansion in powers of z^n , where z is a nonlinear function of q^2 , with an overall multiplicative function accounting for the subthreshold poles and branch cuts,

$$F(t) = \frac{1}{P(t)\phi(t,t_0)} \sum_{k=0}^{\inf} a_k(t_0) z(t,t_0)^k.$$
 (8)

The series coefficients and prefactors can only be determined from fits to lattice or experimental data. In fact, the CLEO Collaboration has determined these coefficients for the $D \rightarrow \pi$, K, η form factors from the semileptonic decays, but in the massless lepton limit. Hence, for $F_0(q^2)$, we use lattice results to determine the first two coefficients.

In the Becirevic-Kaidalov (BK) ansatz [81],

$$F_0(q^2) = \frac{F_0(0)}{1 - \frac{q^2}{\beta m_D^{*2}}},\tag{9}$$

where m_D^* is the mass of the vector meson with flavor $c\bar{d}$ or $c\bar{s}$, depending on the transition being $c \rightarrow d$ or $c \rightarrow s$, respectively.¹ $F_0(0)$ and β are parameters to be fitted to experimental data and in fact, in Ref. [83], the Fermilab and Lattice MILC collaborations have fitted these parameters to CLEO-c data. The normalization f(0) and shape parameter determine the physical observables describing the form factors at large recoil and are given by

$$f(0) \equiv F_{+}(0) = F_{0}(0),$$

$$\frac{1}{\beta} \equiv \frac{(M_{H}^{2} - M_{L}^{2})}{F_{+}(0)} \frac{dF_{0}}{dq^{2}}\Big|_{q^{2} = 0}.$$
 (10)

Using these input parameters we can determine the first two coefficients of the series expansion in Eq. (8) for $F_0(q^2)$.

Few details regarding the z-series form factor expansion can be found in Appendix B. If the series is rapidly converging, even two coefficients may be sufficient to determine the q^2 dependent form factor $F_0(q^2)$. Equating the normalization and slope obtained using Eq. (8) to those obtained from the lattice parameters $(f(0), \beta)$, which in turn had been obtained by fits to experimental data, we can obtain the z-expansion series (up to linear order) for all the form factors. The rst two coefficients of the z-expansion for $D \rightarrow \pi$, $D \rightarrow K$ and $D \rightarrow \eta$ form factors are given in Table II.

In Fig. 2 we show the plots for our results for $D \to \pi$ and $D \to K$ where the lattice input parameters from Ref. [83] are used. We would like to point out that our $D \to \pi$ and $D \to K$ form factor values at $q^2 = 0$ are in very good agreement with those given in Refs. [84,85], which are the most precise published calculations for $D \to \pi l\nu$ and $D \to K l\nu$ form factors, according to Lattice Review [86]. Further, the shape of $F_0(q^2)$ for $D \to \pi, K$ that we obtain after the z-series expansion is consistent with that of Ref. [87]. Moreover, the f(0) values in Table I for $D \to \pi$ and $D \to K$ are also in agreement with the CLEO results of Ref. [88].

Regarding $D \to \eta$ and $D \to \eta'$ form factors, since in these transitions only the nonstrange component η_q is involved, one expects $F_0(q^2)$ for $D \to \eta_q, \eta'_q \sim D \to \pi$. For $D \to \eta$, CLEO has determined $F_+(0)V_{cd} =$ $0.086 \pm 0.006 \pm 0.001$ using the semileptonic decay mode





FIG. 2 (color online). The q^2 dependence of the scalar form factors. The plot on the left displays $F_0(q^2)$ for the $D \to \pi$ transition, while that on the right is for the $D \to K$ transition.

 $D^+ \rightarrow \eta e^+ \nu_e$ [89]. Hence we use this value to estimate $F_0(0)$. However, for η' , while the first observation of the decay mode $D^+ \rightarrow \eta' e^+ \nu_e$ has been reported by CLEO in the same paper, the form factor was not determined in this case, and hence we approximate that the $F_0(0)$ for η' will be the same as that for η . Further, since it has been shown [90] that the form factors and particularly their shape are insensitive to the spectator quark, the shapes for both η and η' are assumed to be the same as those for the $D \rightarrow \pi$ case.

The $D_s \rightarrow \eta, \eta'$ have been estimated with some lattice studies using the $(f(0), \beta)$ values from a recent exploratory paper by Bali *et al.* [91]. However, these have larger uncertainties, since even the lowest pion mass used is still far from the physical mass. Hence, for $D_s \rightarrow \eta, \eta'$ we take the form factors to be similar to that of $D \rightarrow K$ and for $D_s \rightarrow K$ we take them to be similar to $D \rightarrow \pi$. Note that in all these cases the masses of the final mesons for each of the respective decay process are used in obtaining their zexpansion coefficients; the approximations are used only for the input parameters, f(0) and β . Due to this

TABLE I. Best-fit values of BK parameters for the scalar form factors.

Decay	f(0)	β
$\overline{D \to \pi}$	$0.64 \pm 0.03 \pm 0.06$	$1.41 \pm 0.06 \pm 0.07$
$D \to K$	$0.73 \pm 0.03 \pm 0.07$	$1.31 \pm 0.07 \pm 0.13$

TABLE II. z-expansion coefficients obtained after using the BK parameters in Table I.

$0.02 -0.41 \pm 0.05$
$0.01 -0.32 \pm 0.03$
$0.004 -0.27 \pm 0.02$

¹It has been pointed out in Ref. [82] that F_0 , but not F_+ , can be modeled by a single pole.

uncertainty, we have added an additional 3% theoretical error to these form factors.

Turning now to the decay constants, for π and *K* mesons, the $f_{\pi,K}$ are taken from the Particle Data Group (PDG) [92]. For the η and η' , following the method described in [93], it is assumed that the decay constants in the quark flavor basis follow the pattern of particle state mixing. The η and η' are expressed as linear combinations of the orthogonal flavor states,

$$\eta_q = \frac{1}{\sqrt{2}}(u\bar{u} + d\bar{d}), \quad \text{and} \quad \eta_s = s\bar{s}.$$
 (11)

The physical states η and η' are related to these flavor states by

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix},$$

where the $\eta - \eta'$ mixing angle denoted by ϕ represents the sum of the ideal mixing angle and the $\eta - \eta'$ mixing angle (θ) in the octet-singlet basis, $\phi = \theta + \tan^{-1}\sqrt{2}$. Hence the decay constants (form factors) f_q and f_s (F_{0_q} and F_{0_s}), corresponding to those for η_q and η_s ($D \rightarrow \eta_q$ and $D \rightarrow \eta_s$), respectively, are given by

$$\begin{aligned} f_{\eta}^{q} &= f_{q} \cos \phi, & f_{\eta}^{s} = -f_{s} \sin \phi, \\ F_{0_{\eta}}^{q} &= F_{0_{q}} \cos \phi, & F_{0_{\eta}}^{s} = -F_{0_{s}} \sin \phi, \\ f_{\eta'}^{q} &= f_{q} \sin \phi, & f_{\eta'}^{s} = f_{s} \cos \phi, \\ F_{0_{\eta'}}^{q} &= F_{0_{q}} \sin \phi, & F_{0_{\eta'}}^{s} = F_{0_{s}} \cos \phi. \end{aligned}$$

A widely used phenomenological fit for the decay constants as well as the mixing angle ϕ is given in Ref. [93]. This fit was based on various ratios of decay rates having η' in the final state with respect to that with η , for example, $\Gamma(J/\psi \rightarrow \eta'\rho)/\Gamma(J/\psi \rightarrow \eta\rho)$, comparison of cross-sections of scattering processes for $\pi^- p \rightarrow \eta' n$ with that of $\pi^- p \rightarrow \eta n$ etc. Recently, *BABAR*, with more accurate data on two photon widths of light pseudoscalar mesons, did a combined analysis [94] along with CLEO data to yield a mixing angle and decay constants with reduced uncertainties: $\phi = 37.66 \pm 0.70$, $\frac{f_a}{f_x} = 1.078 \pm 0.044$, and $\frac{f_s}{f_z} = 1.246 \pm 0.087$, which are used in this work.

With the above inputs, the un-unitarized amplitudes for all the two-body pseudoscalar-pseudoscalar (*PP*) modes SCS, CF, and DCS may be written and are listed subsequently. For the decay modes involving η and η' , to distinguish the case in which η_q is the P_2 meson of Eq. (4) from that where η_s is, the notation used is

$$\begin{split} C^{f}_{\eta_{q}} &= \frac{G_{F}}{\sqrt{2}} V_{\text{CKM}} a_{2}(\mu) f_{q}(m_{D}^{2} - m_{P_{1}}^{2}) F_{0}^{DP_{1}}(m_{\eta}^{2}), \\ C^{f}_{\eta_{s}} &= \frac{G_{F}}{\sqrt{2}} V_{\text{CKM}} a_{2}(\mu) f_{s}(m_{D}^{2} - m_{P_{1}}^{2}) F_{0}^{DP_{1}}(m_{\eta}^{2}), \\ C^{f}_{\eta_{q}'} &= \frac{G_{F}}{\sqrt{2}} V_{\text{CKM}} a_{2}(\mu) f_{q}(m_{D}^{2} - m_{P_{1}}^{2}) F_{0}^{DP_{1}}(m_{\eta'}^{2}), \\ C^{f}_{\eta_{s}'} &= \frac{G_{F}}{\sqrt{2}} V_{\text{CKM}} a_{2}(\mu) f_{s}(m_{D}^{2} - m_{P_{1}}^{2}) F_{0}^{DP_{1}}(m_{\eta'}^{2}). \end{split}$$

Similarly, to distinguish the cases where η_q or η_s is the P_1 meson, which incidentally appear in both tree (*T*) and color-suppressed (*C*) amplitudes (unlike for the case discussed above), we use the notation

(1) SCS decays

$$\begin{split} A(D^{0} \to \pi^{+}\pi^{-}) &= -V_{cd}V_{ud}(T+E_{q}) \\ A(D^{0} \to \pi^{0}\pi^{0}) &= \frac{V_{cd}V_{ud}}{\sqrt{2}}(-C+E_{q}) \\ A(D^{0} \to \pi^{0}\eta) &= \frac{V_{cd}V_{ud}}{2}(-C_{\eta_{q}}^{F_{0}} + C_{\eta_{q}}^{f})\cos\phi \\ &- V_{cs}V_{us}C_{\eta_{s}}^{f}\frac{\sin\phi}{\sqrt{2}} - V_{cd}V_{ud}E_{q}\cos\phi \\ A(D^{0} \to \pi^{0}\eta') &= \frac{V_{cd}V_{ud}}{2}(-C_{\eta_{s}}^{F_{0}} + C_{\eta_{s}}^{f})\sin\phi \\ &+ V_{cs}V_{us}C_{\eta_{s}}^{f}\frac{\cos\phi}{\sqrt{2}} - V_{cd}V_{ud}E_{q}\sin\phi \\ A(D^{0} \to \eta\eta) &= \frac{V_{cd}V_{ud}}{\sqrt{2}}(C_{\eta_{q}}^{f} + E_{q})\cos^{2}\phi \\ &+ V_{cs}V_{us}\left(-C_{\eta_{s}}^{f}\frac{\sin2\phi}{2\sqrt{2}} + \sqrt{2}E_{s}\sin^{2}\phi\right) \\ A(D^{0} \to \eta\eta') &= V_{cd}V_{ud}\left(E_{q}\frac{\sin2\phi}{2} + (C_{\eta_{q}}^{f} + C_{\eta_{s}}^{f})\frac{\sin2\phi}{\sqrt{2}}\right) \\ &+ V_{cs}V_{us}\left(-C_{\eta_{s}}^{f}\frac{\sin^{2}\phi}{\sqrt{2}} + C_{\eta_{s}}^{f}\frac{\cos^{2}\phi}{\sqrt{2}}\right) \\ A(D^{0} \to \pi^{0}\pi') &= -V_{cs}V_{us}(T+E_{q}) \\ A(D^{0} \to K^{0}\bar{K}^{0}) &= -(V_{cs}V_{us}E_{q} + V_{cd}V_{ud}E_{s}) \\ A(D^{+} \to \pi^{+}\eta) &= \frac{V_{cd}V_{ud}}{\sqrt{2}}(T_{\eta_{q}}^{F_{0}} + C_{\eta_{q}}^{f} + 2A_{q})\cos\phi \\ &- V_{cs}V_{us}C_{\eta_{s}}^{f}\sin\phi \\ A(D^{+} \to \pi^{+}\eta') &= \frac{V_{cd}V_{ud}}{\sqrt{2}}(T_{\eta_{q}}^{F_{0}} + C_{\eta_{q}}^{f} + 2A_{q})\sin\phi \\ &+ V_{cs}V_{us}C_{\eta_{s}}^{f}\cos\phi \\ A(D^{+} \to K^{+}\bar{K}^{0}) &= V_{cd}V_{ud}A_{s} + V_{cs}V_{us}T \end{split}$$

$$\begin{split} A(D_{s}^{+} \to \pi^{+}K^{0}) &= -(V_{cd}V_{ud}T + V_{cs}V_{us}A_{q}) \\ A(D_{s}^{+} \to \pi^{0}K^{+}) &= -\frac{1}{\sqrt{2}}(V_{cd}V_{ud}C - V_{cs}V_{us}A_{q}) \\ A(D_{s}^{+} \to K^{+}\eta) &= (V_{cd}V_{ud}C_{\eta_{q}}^{f} + V_{cs}V_{us}A_{q})\frac{\cos\phi}{\sqrt{2}} \\ &- V_{cs}V_{us}(T_{\eta_{s}}^{F_{0}} + C_{\eta_{s}}^{f} + A_{s})\sin\phi \\ A(D_{s}^{+} \to K^{+}\eta') &= (V_{cd}V_{ud}C_{\eta_{q}}^{f} + V_{cs}V_{us}A_{q})\frac{\sin\phi}{\sqrt{2}} \\ &+ V_{cs}V_{us}(T_{\eta_{s}}^{F_{0}} + C_{\eta_{s}}^{f} + A_{s})\cos\phi \end{split}$$

The $D^0 \rightarrow \pi \pi$ SCS decays obey the following triangular isospin relation:

$$A(D^0 \to \pi^+ \pi^-) + \sqrt{2}A(D^0 \to \pi^0 \pi^0) = \sqrt{2}A(D^+ \to \pi^+ \pi^0).$$

(2) CF decays

$$\begin{split} &(D^0 \to K^- \pi^+) = -V_{cs} V_{ud} (T + E_q) \\ &(D^0 \to \bar{K}^0 \pi^0) = -V_{cs} V_{ud} \frac{(C - E_q)}{\sqrt{2}} \\ &(D^0 \to \bar{K}^0 \eta) = V_{cs} V_{ud} \Big((-C_{\eta_q}^{F_0} - E_q) \frac{\cos \phi}{\sqrt{2}} + E_s \sin \phi \Big) \\ &(D^0 \to \bar{K}^0 \eta') = V_{cs} V_{ud} \Big((-C_{\eta_q'}^{F_0} - E_q) \frac{\sin \phi}{\sqrt{2}} - E_s \cos \phi \Big) \\ &(D^+ \to \bar{K}^0 \pi^+) = -V_{cs} V_{ud} (T + C) \\ &(D_s^+ \to \bar{K}^0 K^+) = -V_{cs} V_{ud} (C + A_s) \\ &(D_s^+ \to \pi^+ \eta) = \frac{V_{cs} V_{ud}}{\sqrt{2}} (T_{\eta_s}^{F_0} \sin \phi - A_q \cos \phi) \\ &(D_s^+ \to \pi^+ \eta') = \frac{V_{cs} V_{ud}}{\sqrt{2}} (-T_{\eta_s'}^{F_0} \cos \phi - A_q \sin \phi) \end{split}$$

The $D^0 \rightarrow K\pi$ CF decays obey the following triangular isospin relation:

$$A(D^0 \to K^- \pi^+) + \sqrt{2}A(D^0 \to \bar{K^0}\pi^0) = A(D^+ \to \bar{K^0}\pi^+).$$

(3) DCS decays

$$\begin{split} &(D^0 \to K^+ \pi^-) = V_{cd} V_{us} (T + E_q) \\ &(D^0 \to K^0 \pi^0) = V_{cd} V_{us} \frac{(C - E_q)}{\sqrt{2}} \\ &(D^0 \to K^0 \eta) = V_{cd} V_{us} \Big((C_{\eta_q}^{F_0} + E_q) \frac{\cos \phi}{\sqrt{2}} - E_s \sin \phi \Big) \\ &(D^0 \to K^0 \eta') = V_{cd} V_{us} \Big((C_{\eta_q'}^{F_0} + E_q) \frac{\sin \phi}{\sqrt{2}} + E_s \cos \phi \Big) \\ &(D^+ \to K^0 \pi^+) = V_{cd} V_{us} (C + A_q) \\ &(D^+ \to K^+ \pi^0) = V_{cd} V_{us} \frac{T - A_q}{\sqrt{2}} \\ &(D^+ \to K^+ \eta) = -V_{cd} V_{us} \Big((T_{\eta_q}^{F_0} + A_q) \frac{\cos \phi}{\sqrt{2}} - A_s \sin \phi \Big) \\ &(D^+ \to K^+ \eta') = -V_{cd} V_{us} \Big((T_{\eta_q'}^{F_0} + A_q) \frac{\sin \phi}{\sqrt{2}} + A_s \cos \phi \Big) \\ &(D_s^+ \to K^+ K^0) = V_{cd} V_{us} (T + C) \end{split}$$

The $D^0 \rightarrow K\pi$ DCS decays obey the following quadrilateral isospin relation:

$$\begin{split} A(D^0 \to K^+ \pi^-) &+ \sqrt{2} A(D^0 \to K^0 \pi^0) \\ &= A(D^+ \to K^0 \pi^+) + \sqrt{2} A(D^+ \to K^+ \pi^0). \end{split}$$

$$\begin{split} C_{\eta_q}^{F_0}(T_{\eta_q}^{F_0}) &= \frac{G_F}{\sqrt{2}} V_{\text{CKM}} a_2(\mu)(a_1(\mu)) \\ &\times f_{P_2}(m_D^2 - m_\eta^2) F_0^{D\eta_q}(m_{P_2}^2), \\ C_{\eta_s}^{F_0}(T_{\eta_s}^{F_0}) &= \frac{G_F}{\sqrt{2}} V_{\text{CKM}} a_2(\mu)(a_1(\mu)) \\ &\times f_{P_2}(m_D^2 - m_\eta^2) F_0^{D\eta_s}(m_{P_2}^2), \\ C_{\eta_q}^{F_0}(T_{\eta_q}^{F_0}) &= \frac{G_F}{\sqrt{2}} V_{\text{CKM}} a_2(\mu)(a_1(\mu)) \\ &\times f_{P_2}(m_D^2 - m_{\eta'}^2) F_0^{D\eta_q'}(m_{P_2}^2), \\ C_{\eta_s'}^{F_0}(T_{\eta_s}^{F_0}) &= \frac{G_F}{\sqrt{2}} V_{\text{CKM}} a_2(\mu)(a_1(\mu)) \\ &\times f_{P_2}(m_D^2 - m_{\eta'}^2) F_0^{D\eta_q'}(m_{P_2}^2). \end{split}$$

III. FINAL STATE INTERACTIONS

Final state interaction effects are incorporated using unitarity relations, where the contribution to any channel is a result of the sum over all possible hadronic intermediate states. Hence for all the $n \ D \rightarrow PP$ decays, the FSI

corrected amplitudes or the "unitarized" amplitudes, A_i^U with i = 1, ...n, are given by²

$$\mathcal{A}_{i}^{U} = \sum_{k=1}^{N} \mathcal{S}_{ik}^{1/2} \mathcal{A}_{k}, \qquad (12)$$

where S is the strong interaction matrix and k = $1, \dots, n, n+1, \dots N$ stands for all possible states that can rescatter into the PP states. In the heavy quark limit the hard rescattering dominates, in which case the sum can be interpreted to be over all intermediate states of partons and the number of these states will hence be very large. Parton hadron duality will then permit this estimation. These corrections are incorporated into the hard scattering contributions in the QCD factorization approach of Ref. [74] for the case of B meson decays. For the case of charmed meson decays, since m_c is not large enough, we include these NF corrections in the modified Wilson coefficients, in terms of parameters χ_1 and χ_2 . However, some residual long distance FSIs may be left which are particularly important for charmed meson decays, due to the nearby resonances. This residual rescattering is considered in the limited set of $D \rightarrow PP$ decays, to which the duality cannot be applied, and therefore these effects may not be incorporated in the NF corrections.

The S matrix in Eq. (12) can be written in terms of a residual matrix (S_{res}) for the rescattering among the $D \rightarrow PP$ states alone and the scattering matrix which accounts for the hard rescattering from all possible hadronic states into these channels, resulting in the factorization amplitudes as

$$S_{ik} = \sum_{j=1}^{n} (S_1)_{ij} (S_2)_{jk}, \text{ where } S_1 = S_{res} \text{ and}$$

$$S_2 = S_1^{-1} S, \text{ with } \mathcal{A}_j^{fac} = \sum_{k=1}^{N} (S_2^{1/2})_{jk} \mathcal{A}_k, \text{ resulting in}$$

$$\mathcal{A}_i^{U} = \sum_{k=1}^{n} (S_{res}^{1/2})_{ij} \mathcal{A}_j^{fac}.$$
(13)

Any S matrix can be written in terms of a real, symmetric K matrix as $S = (1 - iK)^{-1}(1 + iK)$. Hence, the unitarized amplitudes in Eq. (13) may be written as

$$\mathcal{A}_{i}^{U} = \sum_{k=1}^{n} ((1 - iK)^{-1})_{ij} \mathcal{A}_{j}^{fac}.$$
 (14)

The K matrix parametrization has the advantage that the resonances coupling two-body channels are represented by poles in the K matrix. The summation in Eq. (14) corresponds to summing the geometric series, where the final state hadrons are produced from scattering via resonance at

PHYSICAL REVIEW D 92, 014032 (2015)

different orders, starting from zero, i.e., directly from the decaying meson without the resonance contribution, or resonant rescattering occurring once, twice, and so on. While such coupled FSIs have been considered in the past in many papers [31,65,72,77,96–99], most of the papers on charm decays further assume SU(3) to relate the parameters of the coupling matrix. Since SU(3) is broken, we prefer to use the measured decay rates of the resonances to various channels to fix the K matrix parameters as far as possible and the ones not measured are left as parameters to be determined by fits of all the theoretical branching ratios to the observed values.

For each of the SCS, CF, and DCS modes, states with the same isospin are coupled together. In general the *K* matrix coupling three channels will have the form

$$K(s) = \frac{1}{(m_{\text{Res}}^2 - s)} \begin{bmatrix} k_1 \Gamma_{11} & \sqrt{k_1 k_2} \Gamma_{12} & \sqrt{k_1 k_3} \Gamma_{13} \\ \sqrt{k_2 k_1} \Gamma_{21} & k_2 \Gamma_{22} & \sqrt{k_2 k_3} \Gamma_{23} \\ \sqrt{k_3 k_1} \Gamma_{31} & \sqrt{k_3 k_2} \Gamma_{32} & k_3 \Gamma_{33} \end{bmatrix},$$

where m_{Res} denotes the mass of the resonance through which the different channels are coupled; and k_1 , k_2 , and k_3 are the c.m. momenta of the three decay modes. There are six independent parameters Γ_{ij} . To reduce the independent parameters to a manageable number, we impose the requirement that the diagonal cofactors of K(s) vanish [or equivalently, det K(s) = 0]. This leads us to three conditions,

$$\Gamma_{12}^2 = \Gamma_{11}\Gamma_{22}, \quad \Gamma_{13}^2 = \Gamma_{11}\Gamma_{33}, \text{ and } \Gamma_{23}^2 = \Gamma_{22}\Gamma_{33}.$$
 (15)

The Γ_{ii} 's are related to the partial decay width of the resonance to the *i*th channel.

To illustrate this, we consider first the case of isospin zero states of SCS decay modes of the D^0 meson. The isospin zero combination of the $\pi^+\pi^-$ and $\pi^0\pi^0$, K^+K^- and $K^0\bar{K}^0$, and the $\eta\eta$ modes, are coupled via Eq. (14) with the $f_0(1710)$ pole in the K matrix. Hence, for this specific case of coupling of the isospin zero states, in the K matrix, $m_{\text{Res}} = 1.720 \text{ GeV}$, $k_1 = \frac{1}{2}(m_{D^0}^2 - 4m_{\pi^0}^2)$, $k_2 = \frac{1}{2}(m_{D^0}^2 - 4m_{K^0}^2)$, $k_3 = \frac{1}{2}(m_{D^0}^2 - 4m_{\eta}^2)$, and we have

$$\begin{split} \Gamma(f_0 \to \pi\pi) &= \frac{\Gamma_{11}k_1}{m_{\text{Res}}}, \qquad \Gamma(f_0 \to K\bar{K}) = \frac{\Gamma_{22}k_2}{m_{\text{Res}}}, \\ \text{d} \quad \Gamma(f_0 \to \eta\eta) &= \frac{\Gamma_{33}k_3}{m_{\text{Res}}}. \end{split}$$

Experimentally only the two ratios of the decay rates, $\Gamma(f_0 \to K\bar{K})/\Gamma(f_0 \to \pi\pi)$ and $\Gamma(f_0 \to K\bar{K})/\Gamma(f_0 \to \eta\eta)$, have been determined. Hence we keep $g_{\rm pe} \equiv \Gamma(f_0 \to K\bar{K})$ as a parameter, to be determined from fits of our theoretical estimates to the observed branching ratios.

Similarly, for the I = 1 case, we take the $a_0(1450)$ resonance with $m_{\text{Res}} = 1.474$ GeV and $\Gamma_{\text{Res}} = 0.265$ GeV,

an

²For part of the formalism used in this section, we closely follow Refs. [67,95].

to be responsible for the rescattering among the channels $K\bar{K}$, $\pi\eta$, and $\pi\eta'$, to which this resonance decays. Here again, the decay rate $\Gamma(a_0 \rightarrow \pi \eta)$ is not yet accurately measured and is treated as a parameter (h_{pe}) that may be predicted from the fits of all the branching ratios of the $D \rightarrow PP$ modes to experimental data. Note that the $K\bar{K}$, $\pi\eta$, and $\pi\eta'$ states appear as final states not only of SCS D^0 and D^+ decays, but also in the CF decays of the D_s^+ decays. For all three sets of decays, the same K matrix (apart from tiny modifications in the c.m. momenta and the mass squared of the decaying meson) will suffice, and more importantly with the same one unknown parameter, while many additional observables (all the branching ratios of these D^0 , D^+ , and D_s^+) will get added to the χ^2 fit. If this one same parameter, along with the other unknowns in our analysis, can simultaneously explain all the observed data, it would indicate that our naive technique of incorporating the FSI effects is satisfactory.

We also couple the isospin 1/2 states of the $K\pi$, $K\eta$, and $K\eta'$ channels, which are the final states in the SCS decays of D_s^+ , CF decays of D^0 , and DCS decays of D^0 and D^+ mesons. Here we use the $K_0^*(1950)$ resonance with $m_{\text{Res}} = 1.945$ GeV. Only the branching ratio, $\Gamma(K_0^* \to K\pi)/\Gamma_{\text{total}}$, has been measured. We take the other two decay rates, $\Gamma(K_0^* \to K\eta)$ and $\Gamma(K_0^* \to K\eta')$, as parameters $(j_{pe_1} \text{ and } j_{pe_2})$ that can be determined by the overall fits of all the branching ratios to data.

IV. NUMERICAL ANALYSIS AND RESULTS

To estimate all the possible sets of coupled channels, the isospin decompositions of all the SCS, CF, and DCS modes are listed subsequently. Here $A^{(U)}$ denotes the bare or ununitarized (unitarized or FSI corrected) amplitudes, respectively, for each of the decay modes, while $A_i^{\text{mode}(U)}$ denotes the corresponding un-unitarized (unitarized) isospin, I = iamplitudes for those modes. With the absence of resonances, in particular isopin components with the right quantum numbers in the vicinity of the charmed meson masses, some of the isospin components of many modes remain un-unitarized.

With all the unitarized isospin amplitudes, we construct the corresponding unitarized decay amplitudes for all the decay modes. The decay rates for all the $D \rightarrow P_1P_2$ are then calculated as

$$\Gamma(D \to P_1 P_2) = \frac{p_c}{8\pi m_D^2} |A(D \to P_1 P_2)|^2.$$
(16)

Here p_c is the center of mass momentum of the mesons in the final state given by

$$p_c = \frac{\sqrt{(m_D^2 - (m_{P_1} + m_{P_2})^2)(m_D^2 - (m_{P_1} - m_{P_2})^2)}}{2m_D}.$$

TABLE III. Parameter best-fit values.

Name	Values	Name	Values	Name	Values
Λ	0.625645	\dot{j}_{pe_1}	0.0000239368	χ^A_a	132.685
χ_1	-2.68215	j_{pe_2}	0.096456	χ^{A}_{s}	193.447
χ_2	2.23605	χ^{E}_{a}	-334.805	ϕ_1	0.302258
$g_{\rm pe}$	0.0471262	χ^{E}_{s}	-81.3363	ϕ_2	2.87681
h _{pe}	0.118834				

The theoretical branching ratios for each of the decay modes of the D^0 , D^+ , or D_s^+ mesons are then obtained by dividing the corresponding decay rates by the total decay widths of these mesons. We then perform a χ^2 fit of these theoretical branching ratios with the experimentally measured branching fractions, estimating all the unknown parameters from the best fit to data.

The unknown parameters in our study are the four parameters representing the NF corrections, χ_1 , χ_2 ; and the four parameters of their respective phases ϕ_1 , ϕ_2 : $\chi^E_{q,s}$ and

(1) SCS decays

$$\begin{split} &A^{(U)}(D^{0} \to \pi^{+}\pi^{-}) \equiv \sqrt{2}\mathcal{A}_{2}^{\pi\pi} + \sqrt{2}\mathcal{A}_{0}^{\pi\pi(U)} \\ &A^{(U)}(D^{0} \to \pi^{0}\pi^{0}) \equiv 2\mathcal{A}_{2}^{\pi\pi} - \mathcal{A}_{0}^{\pi\pi(U)} \\ &A^{(U)}(D^{0} \to \pi^{0}\eta) \equiv \sqrt{3}\mathcal{A}_{1}^{\pi\eta'(U)} \\ &A^{(U)}(D^{0} \to \pi^{0}\eta) \equiv \sqrt{3}\mathcal{A}_{0}^{\eta\eta'(U)} \\ &A^{(U)}(D^{0} \to \eta\eta) \equiv \sqrt{3}\mathcal{A}_{0}^{\eta\eta'} \\ &A^{(U)}(D^{0} \to \eta\eta') \equiv \sqrt{3}\mathcal{A}_{0}^{\eta\eta'} \\ &A^{(U)}(D^{0} \to K^{+}K^{-}) \equiv \sqrt{\frac{3}{2}} \Big(\mathcal{A}_{1}^{KK(U)} + \mathcal{A}_{0}^{KK(U)}\Big) \\ &A^{(U)}(D^{0} \to K^{0}\bar{K}^{0}) \equiv \sqrt{\frac{3}{2}} \Big(\mathcal{A}_{1}^{KK(U)} - \mathcal{A}_{0}^{KK(U)}\Big) \\ &A^{(U)}(D^{+} \to \pi^{+}\pi^{0}) \equiv 3\mathcal{A}_{2}^{\pi\pi} \\ &A^{(U)}(D^{+} \to \pi^{+}\eta) \equiv \mathcal{A}_{1}^{\pi^{+}\eta(U)} \\ &A^{(U)}(D^{+} \to \pi^{+}\eta') \equiv \mathcal{A}_{1}^{\pi^{+}\eta'(U)} \\ &A^{(U)}(D^{+} \to \pi^{+}\eta') \equiv \mathcal{A}_{1}^{\pi^{+}\eta'(U)} \\ &A^{(U)}(D^{+} \to \pi^{+}K^{0}) \equiv \frac{1}{\sqrt{3}}\mathcal{A}_{\frac{3}{2}}^{\pi K} + \sqrt{\frac{2}{3}}\mathcal{A}_{\frac{1}{2}}^{\pi K(U)} \\ &A^{(U)}(D_{s}^{+} \to \pi^{0}K^{+}) \equiv \sqrt{\frac{2}{3}}\mathcal{A}_{\frac{3}{2}}^{\pi K} - \frac{1}{\sqrt{3}}\mathcal{A}_{\frac{1}{2}}^{\pi K(U)} \\ &A^{(U)}(D_{s}^{+} \to K^{+}\eta) \equiv \mathcal{A}_{\frac{1}{2}}^{K^{+}\eta'(U)} \\ &A^{(U)}(D_{s}^{+} \to K^{+}\eta') \equiv \mathcal{A}_{\frac{1}{2}}^{K^{+}\eta'(U)} \end{split}$$

(2) CF decays

$$\begin{split} A^{(U)}(D^{0} \to K^{-}\pi^{+}) &= \frac{1}{3}\mathcal{A}_{\frac{3}{2}}^{\bar{K}\pi} + \frac{2}{3}\mathcal{A}_{\frac{1}{2}}^{\bar{K}\pi(U)} \\ A^{(U)}(D^{0} \to \bar{K}^{0}\pi^{0}) &= \frac{\sqrt{2}}{3}\left(\mathcal{A}_{\frac{3}{2}}^{\bar{K}\pi} - \mathcal{A}_{\frac{1}{2}}^{\bar{K}\pi(U)}\right) \\ A^{(U)}(D^{0} \to \bar{K}^{0}\eta) &= \sqrt{\frac{2}{3}}\mathcal{A}_{\frac{1}{2}}^{\bar{K}\eta(U)} \\ A^{(U)}(D^{0} \to \bar{K}^{0}\eta') &= \sqrt{\frac{2}{3}}\mathcal{A}_{\frac{1}{2}}^{\bar{K}\eta'(U)} \\ A^{(U)}(D^{+} \to \bar{K}^{0}\pi^{+}) &= \mathcal{A}_{\frac{3}{2}}^{\bar{K}\pi^{+}} \\ A^{(U)}(D^{+}_{s} \to \bar{K}^{0}K^{+}) &= \mathcal{A}_{1}^{\bar{K}\bar{K}(U)} \\ A^{(U)}(D^{+}_{s} \to \pi^{+}\eta) &= \mathcal{A}_{1}^{\pi^{+}\eta'(U)} \\ A^{(U)}(D^{+}_{s} \to \pi^{+}\eta') &= \mathcal{A}_{1}^{\pi^{+}\eta'(U)} \end{split}$$

(3) DCS decays

$$\begin{split} A^{(U)}(D^{0} \to K^{+}\pi^{-}) &= \frac{\sqrt{2}}{3} \mathcal{A}_{\frac{1}{2}}^{K\pi} - \frac{\sqrt{2}}{\sqrt{3}} \mathcal{A}_{\frac{1}{2}}^{K\pi(U)} \\ A^{(U)}(D^{0} \to K^{0}\pi^{0}) &= \frac{2}{3} \mathcal{A}_{\frac{3}{2}}^{K\pi} + \frac{1}{\sqrt{3}} \mathcal{A}_{\frac{1}{2}}^{K\pi(U)} \\ A^{(U)}(D^{0} \to K^{0}\eta) &= \mathcal{A}_{\frac{1}{2}}^{K\eta(U)} \\ A^{(U)}(D^{0} \to K^{0}\eta') &= \mathcal{A}_{\frac{1}{2}}^{K\eta'(U)} \\ A^{(U)}(D^{+} \to K^{0}\pi^{+}) &= \frac{\sqrt{2}}{3} \mathcal{A}_{\frac{3}{2}}^{K\pi^{+}} + \frac{\sqrt{2}}{\sqrt{3}} \mathcal{A}_{\frac{1}{2}}^{K\pi^{+}(U)} \end{split}$$

$$\begin{split} A^{(U)}(D^+ \to K^+ \pi^0) &= \frac{2}{3} \mathcal{A}_{\frac{3}{2}}^{K\pi^+} - \frac{1}{\sqrt{3}} \mathcal{A}_{\frac{1}{2}}^{K\pi^+(U)} \\ A^{(U)}(D^+ \to K^+ \eta) &= A_{\frac{1}{2}}^{K^+ \eta(U)} \\ A^{(U)}(D^+ \to K^+ \eta') &= A_{\frac{1}{2}}^{K^+ \eta'(U)} \\ A^{(U)}(D_s^+ \to K^+ K^0) &= \frac{1}{\sqrt{2}} (\mathcal{A}_1^{KK} + \mathcal{A}_0^{KK}) \end{split}$$

 $\chi^{A}_{q,s}$ depicting the strength of the W-exchange and Wannihilation amplitudes with distinct strengths for $q\bar{q}$ and $s\bar{s}$ pair production; one unknown in each of the isospin zero and isospin one K matrices coupling modes from decays of D^{0} and D^{+} mesons; two parameters in the isospin half K matrix coupling various decay modes of D_{s}^{+} ; and one parameter Λ , representing the momentum of the soft degrees of freedom in the charmed mesons, that is used to define the scale for each of the individual decay modes, making a total of 13 unknown parameters. On the other hand, out of all 33 decay modes considered, 28 have been measured, resulting in sufficient observables to determine all the unknown parameters and give predictions for five of the branching fractions of DCS modes that are not yet measured.

Apart from the experimental errors in the observed branching ratios, the calculated errors in our theoretical branching ratio estimates arise from the errors in the form factors, the η and η' decay constants, the $\eta - \eta'$ mixing angle, and the errors in the measured decay widths of the various resonances into the different channels that are included in our χ^2 fits. Errors due to the other theoretical inputs, like meson masses; decay constants of pion, kaon,

TABLE IV. $D \rightarrow PP$ SCS branching ratios. Columns 2 and 3 show our results with annihilation included, for the cases with and without FSI, respectively, while column 4, which displays results without annihilation, includes FSI.

Modes	With FSI	Without FSI	Without Ann.	Experimental Value
$D^0 o \pi^+ \pi^-$	$(1.44 \pm 0.027) \times 10^{-3}$	$(4.35 \pm 1.67) \times 10^{-3}$	$(4.02 \pm 1.75 \times) \times 10^{-3}$	$(1.402 \pm 0.026) \times 10^{-3}$
$D^0 o \pi^0 \pi^0$	$(1.14 \pm 0.56) \times 10^{-3}$	$(3.66 \pm 1.43) \times 10^{-3}$	$(2.04 \pm 0.79) \times 10^{-3}$	$(8.209 \pm 0.35) \times 10^{-4}$
$D^0 \rightarrow K^+ K^-$	$(4.06 \pm 0.77) \times 10^{-3}$	$(4.27 \pm 2.34) \times 10^{-3}$	$(6.78 \pm 3.08) \times 10^{-3}$	$(3.96 \pm 0.08) \times 10^{-3}$
$D^0 \rightarrow K^0 \bar{K^0}$	$(3.42 \pm 0.52) \times 10^{-4}$	$(5.61 \pm 0.00) \times 10^{-4}$	$(2.80 \pm 0.84) \times 10^{-4}$	$(3.4 \pm 0.8) \times 10^{-4}$
$D^0 \rightarrow \pi^0 \eta$	$(1.47 \pm 0.90) \times 10^{-3}$	$(6.47 \pm 2.98 \times 10^{-3}$	$(3.25 \pm 1.51 \times 10^{-3}$	$(6.8 \pm 0.7) imes 10^{-4}$
$D^0 ightarrow \pi^0 \eta'$	$(2.17 \pm 0.65) imes 10^{-3}$	$(3.81 \pm 1.43) \times 10^{-3}$	$(1.85 \pm 0.79) \times 10^{-3}$	$(9.0 \pm 1.4) \times 10^{-4}$
$D^0 \rightarrow \eta \eta$	$(1.27 \pm 0.27) imes 10^{-3}$	$(1.32 \pm 0.41) \times 10^{-3}$	$(1.34 \pm 0.29) \times 10^{-3}$	$(1.67 \pm 0.20) \times 10^{-3}$
$D^0 o \eta \eta'$	$(9.53 \pm 1.83) \times 10^{-4}$	$(1.04 \pm 0.27) \times 10^{-3}$	$(5.38 \pm 1.63) \times 10^{-4}$	$(1.05 \pm 0.26) \times 10^{-3}$
$D^+ o \pi^+ \pi^0$	$(8.89 \pm 4.51) imes 10^{-4}$	$(8.70 \pm 6.70) \times 10^{-4}$	$(9.73 \pm 3.94) \times 10^{-4}$	$(1.19 \pm 0.06) \times 10^{-3}$
$D^+ \rightarrow K^+ \bar{K^0}$	$(3.75 \pm 0.63) \times 10^{-3}$	$(1.02\pm0.37) imes10^{-2}$	$(1.99 \pm 0.56) \times 10^{-2}$	$(5.66 \pm 0.32) \times 10^{-3}$
$D^+ o \pi^+ \eta$	$(4.72 \pm 0.21) \times 10^{-3}$	$(2.34 \pm 1.26) \times 10^{-2}$	$(1.66 \pm 0.77) \times 10^{-2}$	$(3.53 \pm 0.21) \times 10^{-3}$
$D^+ o \pi^+ \eta'$	$(6.76 \pm 2.19) \times 10^{-3}$	$(3.00 \pm 0.76) \times 10^{-2}$	$(9.78 \pm 3.35) \times 10^{-3}$	$(4.67 \pm 0.29) \times 10^{-3}$
$D_s^+ \to \pi^+ K^0$	$(1.96 \pm 0.90) \times 10^{-3}$	$(1.46 \pm 1.10) \times 10^{-3}$	$(1.32 \pm 1.01) \times 10^{-3}$	$(2.42 \pm 0.12) \times 10^{-3}$
$D_s^+ \to \pi^0 K^+$	$(8.17 \pm 4.64) \times 10^{-4}$	$(1.74 \pm 1.00) \times 10^{-4}$	$(1.01 \pm 0.54) \times 10^{-3}$	$(6.3 \pm 2.1) \times 10^{-4}$
$D_s^+ \to K^+ \eta$	$(1.50 \pm 0.75) \times 10^{-3}$	$(6.40 \pm 4.52) \times 10^{-3}$	$(2.23 \pm 1.82) \times 10^{-3}$	$(1.76 \pm 0.35) \times 10^{-3}$
$D_s^+ \to K^+ \eta'$	$(7.07 \pm 0.49) \times 10^{-4}$	$(2.09\pm 0.87)\times 10^{-3}$	$(0.57\pm 0.47)\times 10^{-4}$	$(1.8 \pm 0.6) \times 10^{-3}$

charmed mesons; and CKM mixing elements (involving the first two generations) are negligibly small.

As discussed in Sec. II C, the z-series expansion has been obtained for the form factors, keeping the first two terms of this series. The coefficients a_0 and a_1 of these two terms are functions of normalization and shape parameters f(0) and β , obtained from lattice results. The errors in these lattice parameters are used to obtain the errors in the expansion coefficient functions and then propagated to get the errors in the form factors. We find that the errors in the form

factors vary from $\approx 5\%$ to $\approx 25\%$. In Tables IV, V, and VI, we list the values of the branching ratios of all the SCS, CF, and DCS $D \rightarrow PP$ modes obtained from our analysis, after incorporating the FSI effects (shown in the second column), as well as in the absence of the FSI (column 3) and absence of annihilation (column 4), along with the corresponding observed experimental branching ratios (column 5) given in PDG [92]. We also predict the branching ratios for a few DCS modes that have not been experimentally measured yet and are given in the second column of Table VI

After incorporating all the errors, the χ^2 minimization results in the best-fit values of all the parameters given in Table III.

PHYSICAL REVIEW D 92, 014032 (2015)

V. CONCLUSIONS

For several decades, various ratios of decay rates of many of the $D \rightarrow PP$ modes remained a puzzle, as these ratios were expected to be close to one in the SU(3) limit, but the measured values exhibited large deviations from unity.

We have evaluated the bare amplitudes of all the $D \rightarrow PP$ modes using factorization; however, we add nonfactorizable corrections, weak annihilation, and exchange contributions as parameters. In the hadron matrix elements, the q^2 dependence of the form factors involved are evaluated using the z-expansion method and, finally, resonant final state interaction effects are incorporated. The parameters of the *K* matrix coupling the various channels are defined using the measured decay widths of the resonances (where available) and those unobserved are left as parameters to be fitted from all the measured 28 $D \rightarrow PP$ branching ratios. Our best fit has a χ^2 /degree of freedom of 2.25, which is an improvement over the previous results in Refs. [59,76].

(i) It may be noted that we have added the NF corrections to the color-suppressed and tree amplitudes, in contrast to Ref. [76] where they have

TABLE V. $D \rightarrow PP$ CF branching ratios. Inclusion of annihilation/FSI in our branching ratio estimates shown in various columns is the same as those specified for Table IV.

Modes	With FSI	Without FSI	Without Ann.	Experimental Value
$\overline{D^0 \to K^- \pi^+}$	$(3.70 \pm 1.33) \times 10^{-2}$	$(8.83 \pm 2.47) \times 10^{-2}$	$(5.63 \pm 1.81) \times 10^{-2}$	$(3.88 \pm 0.05) \times 10^{-2}$
$D^0 \rightarrow \bar{K^0} \pi^0$	$(1.88 \pm 0.99) \times 10^{-2}$	$(1.29 \pm 0.44) \times 10^{-1}$	$(3.30 \pm 1.47) \times 10^{-2}$	$(2.38 \pm 0.08) \times 10^{-2}$
$D^0 \rightarrow \bar{K^0} \eta$	$(1.59\pm0.48) imes10^{-2}$	$(0.97 \pm 0.33) imes 10^{-2}$	$(1.09 \pm 0.34) \times 10^{-2}$	$(0.958\pm0.06) imes10^{-2}$
$D^0 \rightarrow \bar{K^0} \eta'$	$(2.29 \pm 0.43) \times 10^{-2}$	$(2.06\pm 0.30)\times 10^{-2}$	$(2.45\pm 0.47)\times 10^{-2}$	$(1.88 \pm 0.1) \times 10^{-2}$
$D^+ \rightarrow \bar{K^0}\pi^+$	$(3.42 \pm 1.78) \times 10^{-2}$	$(1.35 \pm 1.12) imes 10^{-1}$	$(5.25 \pm 3.34) \times 10^{-2}$	$(2.94\pm0.14) imes10^{-2}$
$D_s^+ \rightarrow \bar{K^0}K^+$	$(5.65 \pm 1.29) \times 10^{-2}$	$(1.70 \pm 0.79) \times 10^{-1}$	$(1.35 \pm 0.53) \times 10^{-1}$	$(2.95\pm0.14) imes10^{-2}$
$D_s^+ o \pi^+ \eta$	$(2.26 \pm 0.82) \times 10^{-2}$	$(0.78\pm0.56) imes10^{-2}$	$(2.14 \pm 0.90) \times 10^{-2}$	$(1.69 \pm 0.10) \times 10^{-2}$
$D_s^+ \to \pi^+ \eta'$	$(2.64 \pm 0.78) \times 10^{-2}$	$(3.73 \pm 1.52) \times 10^{-2}$	$(2.52 \pm 0.85) \times 10^{-2}$	$(3.94 \pm 0.25) \times 10^{-2}$

TABLE VI. $D \rightarrow PP$ DCS branching ratios. Inclusion of annihilation/FSI in our branching ratio estimates shown in various columns is the same as specified for Table IV.

Modes	With FSI	Without FSI	Without Ann.	Experimental Value
$D^0 \rightarrow K^+ \pi^-$	$(1.77 \pm 0.88) \times 10^{-4}$	$(3.71 \pm 1.33) \times 10^{-4}$	$(2.48 \pm 1.07) \times 10^{-4}$	$(1.38 \pm 0.028) \times 10^{-4}$
$D^0 \rightarrow K^0 \pi^0$	$(2.11 \pm 0.26) \times 10^{-4}$	$(3.70 \pm 1.35) \times 10^{-4}$	$(0.68 \pm 0.46) \times 10^{-4}$	••••
$D^0 \to K^0 \eta$	$(0.94 \pm 0.45) imes 10^{-4}$	$(0.28 \pm 0.10) imes 10^{-4}$	$(0.96 \pm 0.32) \times 10^{-4}$	
$D^0 \to K^0 \eta'$	$(8.02 \pm 3.32) \times 10^{-4}$	$(0.59 \pm 0.08) imes 10^{-4}$	$(9.22 \pm 1.61) \times 10^{-4}$	• • •
$D^+ \rightarrow K^0 \pi^+$	$(3.27 \pm 1.86) \times 10^{-4}$	$(1.19 \pm 0.55) \times 10^{-3}$	$(3.51 \pm 2.11) \times 10^{-4}$	• • •
$D^+ \to K^+ \pi^0$	$(3.07 \pm 1.02) \times 10^{-4}$	$(2.15 \pm 1.17) \times 10^{-4}$	$(3.27 \pm 1.39) \times 10^{-4}$	$(1.83 \pm 0.26) imes 10^{-4}$
$D^+ \to K^+ \eta$	$(0.98 \pm 0.26) imes 10^{-4}$	$(1.04 \pm 0.23) \times 10^{-4}$	$(0.89 \pm 0.27) imes 10^{-4}$	$(1.08 \pm 0.17) \times 10^{-4}$
$D^+ \rightarrow K^+ \eta'$	$(1.40 \pm 0.39) \times 10^{-4}$	$(1.82\pm0.18) imes10^{-4}$	$(1.35 \pm 0.39) \times 10^{-4}$	$(1.76 \pm 0.22 \times 10^{-4})$
$D_s^+ \to K^+ K^0$	$(7.84 \pm 2.31) \times 10^{-4}$	$(0.68\pm 0.09)\times 10^{-4}$	$(0.72\pm 0.44)\times 10^{-4}$	

only been added to the color-suppressed amplitudes. Of course NF corrections in the color-suppressed contributions are well motivated; however, vertex corrections, etc., will be present even in the tree contributions and hence these need to be added. Even for *B* decays, Ref. [74] evaluates them for both the color-suppressed and tree amplitudes. For D decays, since the nonfactorizable corrections are expected to be even more important due to the $1/m_c$ corrections being non-negligible, perhaps incorporating them into tree and color-suppressed contributions would result in better fits to data. In fact, such corrections were incorporated into both tree and color-suppressed amplitudes in Ref. [73] and several other papers.

- (ii) We have incorporated resonant FSIs in addition to the NF corrections to the amplitudes. We clarify that this will not result in any double counting. It is true that nonfactorizable contributions include final state interactions. In general these FSIs should include all possible states that can rescatter into the PP states being considered. However, when contributions from all hadronic states at large enough energy (short distance part) are summed over, one should be able to work in terms of the quark and gluon degrees of freedom (using duality). However, in addition there can be soft (long distance) leftover, residual scattering among a limited set of PP states. This final state interaction at the hadronic level is elastic in the sense of probability conservation among the limited subset of states and incorporates only this residual scattering, not all possible FSIs. This scattering among the limited PP modes cannot be treated in the factorization amplitudes by duality and hence there should be no double counting. Moreover, the Wilson coefficients which incorporate the nonfactorizable pieces are expected to be universal. On the other hand, the resonant final state interactions coupling different channels in various isospin states cannot be universal but have to be channel dependent. Hence, this needs to be incorporated separately.
- (iii) We are able to get reasonable fits to almost all the observed branching ratios. In particular, our branching fractions for $D \rightarrow KK$, $D \rightarrow \pi\pi$ modes that have been a longstanding puzzle are in agreement with the corresponding measured values.
- (iv) For constructing the FSI K matrix, if SU(3) couplings are used, then due to the symmetry of these matrices, diagonalization results in one or more eigenvalues being zero. As a result, in this case it turns out that FSI modifies only the annihilation and exchange contributions, while tree and color-suppressed amplitudes remain unaffected. However if SU(3) is not used, but the K matrix is constructed from measured rates of the

resonances to different channels, and one leaves those not observed as parameters (as is done in this work), then it is the net un-unitarized amplitudes in different isospin states that get enhanced or suppressed, and not just the annihilation/exchange contributions alone.

- (v) In Tables IV, V, and VI, we have added columns comparing the theoretical branching ratios for all the *PP* modes considered, for cases with and without the coupled channel FSI. It can be clearly seen that the fits to data are indeed much better with FSI.
- (vi) We have evaluated the cosine of the strong phase difference between the unitarized amplitudes for $D^0 \rightarrow K^-\pi^+$ and $D^0 \rightarrow K^+\pi^-$ and obtain $\cos \delta_{K\pi} = 0.94 \pm 0.027$. This result is consistent with the recently measured BESIII result, $\cos \delta_{K\pi} = 1.02 \pm 0.11 \pm 0.06 \pm 0.01$ [100].
- (vii) The mode $D^0 \to K^0 \bar{K^0}$ does not have any tree or color-suppressed contributions, but can come only from W-exchange. In fact, there are two exchange contributions, one appearing with a $d\bar{d}$ and the other with an $s\bar{s}$, which under exact SU(3) symmetry would cancel each other, resulting in a null amplitude. However, since our parameters for these two contributions are distinct, our bare amplitude for this mode is small but nonvanishing. There have been speculations [73,77] that this mode can arise just from final state interactions, even in the absence of a weak exchange contribution. However, from Table IV it is clear that without the exchange contribution we are unable to generate a large enough rate: both final state interaction and the exchange contribution are necessary for consistency with the measured branching fraction.
- (viii) We have also evaluated the four ratios of amplitudes that had been specified in a recent paper [101]. In the SU(3) limit these are all expected to be unity. Our theoretical estimates for these ratios are given below:

$$\begin{split} R_1 &\equiv \frac{|A(D^0 \to K^+ \pi^-)|}{|A(D^0 \to \pi^+ K^- |\tan^2 \theta_c} = 1.27 \pm 0.32, \\ R_2 &\equiv \frac{|A(D^0 \to K^+ K^-)|}{|A(D^0 \to \pi^+ \pi^-)|} = 1.27 \pm 0.42, \\ R_3 &\equiv \frac{|A(D^0 \to K^+ K^-)| + |A(D^0 \to \pi^+ \pi^-)|}{|A(D^0 \to \pi^+ K^- |\tan \theta_c + |A(D^0 \to K^+ \pi^-)|\tan^{-1} \theta_c} \\ &= 1.19 \pm 0.28, \\ R_4 &\equiv \sqrt{\frac{|A(D^0 \to K^+ K^-)||A(D^0 \to \pi^+ \pi^-)|}{|A(D^0 \to \pi^+ K^-)||A(D^0 \to K^+ \pi^-)|}} \\ &= 1.19 \pm 0.26. \end{split}$$

Furthermore, the following combination of these ratios is expected to be vanishing up to fourth order in U-spin breaking,

$$\Delta R \equiv R_3 - R_4 + \frac{1}{8} [(\sqrt{2R_1 - 1} - 1)^2 - (\sqrt{2R_2 - 1} - 1)^2].$$

Using our unitarized amplitudes, we find the central value of ΔR to be indeed very tiny, however, with a large error:

$$\Delta R = -0.000013 \pm 0.006.$$

Our theoretical errors (in form factors, K-matrix parameters, etc.) are propagated to evaluate the errors in the ratios R_1 , R_2 , R_3 , R_4 and finally in ΔR .

- (ix) We would like to mention that in many modes involving η and η' , we have additional terms in our amplitude due to our distinction of the different form factors, compared to, for example, those that appear in Ref. [59]. A naive look at the color-suppressed diagrams for $D \to \pi^0 \eta(\eta')$ will indicate that the contributions are from the case where the spectator is part of the π^0 , and that where it constitutes the $\eta(\eta')$ must cancel. However, in terms of the specific decay constants and form factors, one is proportional to $-f_{\pi}F_0^{D\eta_q}(m_{\pi}^2)$, while the other is proportional to $f_{\eta_q}F_0^{D\eta_\pi}(m_{\eta}^2)$, which are unequal and hence the net result is non-vanishing.
- (x) While the Particle Data Group [92] does not include a world average for $\Gamma(f_0(1710) \rightarrow K\bar{K})$, it does list two values for the ratio $\frac{\Gamma(f_0(1710) \rightarrow K\bar{K})}{\Gamma_{\text{total}}}$: 0.36 ± 0.12 (Ref. [102]) and 0.38^{+0.09}_{-0.19} (Ref. [103]); our fit value of $g_{\text{pe}}(\equiv \Gamma(f_0(1710) \rightarrow K\bar{K}))$ corresponds to 0.35 for the branching ratio, which is consistent with these values.
- (xi) Our theoretical errors are rather large and could be reduced in the future with more precise form factors available either from measurement of semileptonic D, D_s modes at BES III (where if even the lepton mass could be incorporated, e.g., by looking at modes with muon in the final state, then the q^2 dependent F_0 could be known) or with improved lattice studies, especially for $D \rightarrow \eta', D_s \rightarrow K$, and $D_s \rightarrow \eta, \eta'$.
- (xii) Accurate measurements of the decay widths of the resonances (used for the final state interactions) to many of the coupled channels can reduce the theoretical uncertainties and possibly allow for better fits to data. For example, our fits seem to indicate a rather large value for $h_{\rm pe}$ or the width of $a_0(1450) \rightarrow \pi^0 \eta$. This seems to result in larger branching fractions for many of the isospin one

modes. Future measurement of this width can help reduce this uncertainty and perhaps result in better fits to data for these modes.

(xiii) Out of the 28 observed *PP* modes, we are unable to fit 7 of the modes well. Many of these modes involve η or η' in the final state. Including a gluonium component in the η , η' states may possibly be one way of improving these fits. This, along with improved form factor measurements and observation of decay rates of the resonances (playing a role in final state interactions) to these decay modes, as mentioned in the last two points above, could go a long way toward improving our fits. One glaring misfit is the mode $D^+ \to K^+ \bar{K^0}$. This mode does couple to $\pi^+\eta, \pi^+\eta'$ modes and hence may possibly improve, along with the improvements in the other modes. Finally, perhaps a more sophisticated statistical analysis may also play a role in improving our results, which we hope to carry out in the future [104].

ACKNOWLEDGMENTS

The authors thank L. Gibbons, R. Hill, and J. Libby for correspondence related to form factors. N. S. thanks H.-Y. Cheng for discussions at the early stage of this work and A. Paul, S. Pakvasa, and N. G. Deshpande for comments. She also thanks T. E. Browder for information regarding the BESIII strong phase measurement. A. B. thanks S. Patra for suggestions regarding the numerical analysis. G. A. thanks A. Pich for discussions. G. A.'s work has been supported in part by the Spanish Government and ERDF funds from the EU Commission [Grants No. FPA2011-23778, No. FPA2014-53631-C2-1-P, and No. CSD2007-00042 (Consolider Project CPAN)].

APPENDIX A: THE WILSON COEFFICIENTS

The Wilson coefficients used in the evaluation of all the bare amplitudes have been calculated at the final state hadronic scale. This allows for an additional SU(3) breaking effect and they have been incorporated using the procedure outlined in Ref. [76]. The Wilson coefficients at lower scale are calculated in Ref. [105]. The essential steps are the following.

- (i) In the first step, the Wilson coefficients $C_i(m_W)$ at weak scale are calculated by requiring the equality of the effective theory with five active flavors q = u, d, s, c, b onto the full theory.
- (ii) Next, the coefficients undergo the evolution from the scale m_W to μ through the equation

$$C(\mu) = U_5(\mu, m_W)C(m_W).$$
 (A1)

(iii) In the next step, coefficients are calculated at the scale of the b quark

$$C(m_b) \to Z(m_b).$$
 (A2)

(iv) Now, the Wilson coefficients can be evaluated at the required scale (μ_c or μ_{hadron}) through the equation

$$C(\mu) = U_4(\mu, m_b)Z(m_b).$$
 (A3)

In the above steps, U_5 and U_4 are the 2×2 and 7×7 evolution matrices for five and four active flavors, respectively. The $Z(m_b)$ is given in the Eqs. (A.7) to (A.10) of Ref. [105].

The explicit expressions of the Wilson coefficients obtained after following the above steps are given in the Ref. [76]. They are

$$\begin{split} C_1(\mu) &= -0.2334\alpha^{1.444} + 0.0459\alpha^{0.7778} \\ &\quad + 1.313\alpha^{0.4444} - 0.3041\alpha^{-0.2222}, \\ C_2(\mu) &= 0.2334\alpha^{1.444} + 0.0459\alpha^{0.7778} \\ &\quad - 1.313\alpha^{0.4444} + 0.3041\alpha^{-0.2222}, \end{split}$$

in terms of the running coupling constant α :

$$\alpha = \alpha_s(\mu) = \frac{4\pi}{\beta_0 \ln(\mu^2/\Lambda_{\overline{\text{MS}}}^2)} \left[1 - \frac{\beta_1 \ln \ln(\mu^2/\Lambda_{\overline{\text{MS}}}^2)}{\beta_0^2 \ln(\mu^2/\Lambda_{\overline{\text{MS}}}^2)} \right],$$

with the coefficients

$$\beta_0 = \frac{33 - 2f}{3}, \qquad \beta_1 = 102 - \frac{38}{3}N_f.$$

We take active flavor number $N_f = 3$, and the QCD scale $\Lambda_{\overline{\text{MS}}} = \Lambda_{\overline{\text{MS}}} = 375$ MeV. Again, note that the scale dependent strong coupling constant $\alpha_s(\mu)$ is evaluated at the final state hadronic scales for each individual decay, to take care of the SU(3) breaking.

APPENDIX B: SERIES EXPANSION METHOD FOR FORM FACTORS

Using the analytic properties of $F(q^2)$, a transformation of variables is made which maps the cut on the q^2 plane onto a unit circle |z| < 1, where

$$z(t,t_0) = \frac{\sqrt{t_+ - t} - \sqrt{t_+ - t_0}}{\sqrt{t_+ - t} + \sqrt{t_+ - t_0}}, \qquad t = q^2.$$

where

$$t_{\pm} = (m_D \pm m_{P_1})^2$$
 and $t_0 = t_+ \left(1 - \left(1 - \frac{t_-}{t_+}\right)^{1/2}\right).$

This transformation allows the form factors to be given by an expansion about $q^2 = t_0$, given as

$$F(t) = \frac{1}{P(t)\phi(t,t_0)} \sum_{k=0}^{\inf} a_k(t_0) z(t,t_0)^k,$$

given also as Eq. (8) in the text. The function P(t) in the above is 1 for $D \to \pi$ form factors. For $D_s \to \eta$ and $D \to K$ form factors, $P(t) = z(t, M_{D_{s0}^*}^2)$, where $M_{D_{s0}^*}$ is the nearest 0⁺ resonance mass. The outer function ϕ is given by [106]

$$\begin{split} \phi(t,t_0) &= \sqrt{\frac{3t_+t_-}{32\pi\chi_0}} \bigg(\frac{z(q^2,0)}{-q^2}\bigg)^2 \bigg(\frac{z(q^2,t_0)}{t_0-q^2}\bigg)^{-1/2} \\ &\times \bigg(\frac{z(q^2,t_-)}{t_--q^2}\bigg)^{-1/4} \frac{\sqrt{t_+-q^2}}{(t_+-t_0)^{1/4}}, \end{split}$$

where χ_0 has been calculated [106] using OPE and is given by

$$\chi_0 = \frac{1 + 0.751\alpha_s(m_c)}{8\pi^2}.$$

For simplicity, we ignore the condensate contribution which is of the order $\mathcal{O}(m_c^{-3})$ and $\mathcal{O}(m_c^{-4})$. The strong coupling at charm scale is computed with the package RunDec [107].

- M. Artuso, B. Meadows, and A. A. Petrov, Annu. Rev. Nucl. Part. Sci. 58, 249 (2008).
- [2] A. Ryd and A. A. Petrov, Rev. Mod. Phys. 84, 65 (2012).
- [3] G. Burdman and I. Shipsey, Annu. Rev. Nucl. Part. Sci. 53, 431 (2003).
- [4] G. Burdman, E. Golowich, J. L. Hewett, and S. Pakvasa, Phys. Rev. D 66, 014009 (2002).
- [5] A. A. Petrov, Proc. Sci., BEAUTY2009 (2009) 024 [arXiv:1003.0906].
- [6] G. D'Ambrosio and D. N. Gao, Phys. Lett. B 513, 123 (2001).
- [7] B. D. Yabsley, Int. J. Mod. Phys. A 19, 949 (2004).
- [8] F. Close and H. Lipkin, Phys. Lett. B 551, 337 (2003).
- [9] S. Fajfer, S. Prelovsek, and P. Singer, Phys. Rev. D 64, 114009 (2001).

- [10] Y. Grossman, A. L. Kagan, and Y. Nir, Phys. Rev. D 75, 036008 (2007).
- [11] N. Sinha, R. Sinha, T. E. Browder, S. Pakvasa, and N. G. Deshpande, Phys. Rev. Lett. 99, 262002 (2007).
- [12] J. Link et al. (FOCUS), Phys. Lett. B 485, 62 (2000).
- [13] G. Brandenburg *et al.* (CLEO), Phys. Rev. Lett. **87**, 071802 (2001).
- [14] R. Aaij et al. (LHCb), Phys. Rev. Lett. 108, 111602 (2012).
- [15] T. Aaltonen *et al.* (CDF), Phys. Rev. Lett. **109**, 111801 (2012).
- [16] T. Feldmann, S. Nandi, and A. Soni, J. High Energy Phys. 06 (2012) 007.
- [17] J. Brod, Y. Grossman, A. L. Kagan, and J. Zupan, J. High Energy Phys. 10 (2012) 161.
- [18] J. Brod, A. L. Kagan, and J. Zupan, Phys. Rev. D 86, 014023 (2012).
- [19] H.-Y. Cheng and C.-W. Chiang, Phys. Rev. D 85, 034036 (2012).
- [20] B. Bhattacharya, M. Gronau, and J. L. Rosner, Phys. Rev. D 85, 054014 (2012).
- [21] E. Franco, S. Mishima, and L. Silvestrini, J. High Energy Phys. 05 (2012) 140.
- [22] G. Isidori, J. F. Kamenik, Z. Ligeti, and G. Perez, Phys. Lett. B 711, 46 (2012).
- [23] Y. Grossman and D. J. Robinson, J. High Energy Phys. 04 (2013) 067.
- [24] D. Atwood and A. Soni, Prog. Theor. Exp. Phys. 2013, 093B05 (2013).
- [25] G. F. Giudice, G. Isidori, and P. Paradisi, J. High Energy Phys. 04 (2012) 060.
- [26] G. Hiller, Y. Hochberg, and Y. Nir, Phys. Rev. D 85, 116008 (2012).
- [27] Y. Hochberg and Y. Nir, Phys. Rev. Lett. 108, 261601 (2012).
- [28] W. Altmannshofer, R. Primulando, C.-T. Yu, and F. Yu, J. High Energy Phys. 04 (2012) 049.
- [29] G. Isidori and J. F. Kamenik, Phys. Rev. Lett. 109, 171801 (2012).
- [30] G. Hiller, M. Jung, and S. Schacht, Phys. Rev. D 87, 014024 (2013).
- [31] F. Buccella, M. Lusignoli, A. Pugliese, and P. Santorelli, Phys. Rev. D 88, 074011 (2013).
- [32] S. Muller, U. Nierste, and S. Schacht, arXiv:1503.06759.
- [33] M. Charles, arXiv:1411.4581.
- [34] H. D. Politzer and M. B. Wise, Phys. Lett. B 208, 504 (1988).
- [35] H. Georgi, in Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 91): Perspectives in the Standard Model, River Edge, N.J. (World Scientific, Boulder, Colorado, 1992), 589.
- [36] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, Nucl. Phys. B591, 313 (2000).
- [37] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, Phys. Rev. Lett. 83, 1914 (1999).
- [38] Y.-Y. Keum, H.-n. Li, and A. Sanda, Phys. Lett. B 504, 6 (2001).
- [39] Y.-Y. Keum, H.-n. Li, and A. Sanda, Phys. Rev. D 63, 054008 (2001).
- [40] C.-D. Lu, K. Ukai, and M.-Z. Yang, Phys. Rev. D 63, 074009 (2001).

- [41] C.-D. Lu and M.-Z. Yang, Eur. Phys. J. C 23, 275 (2002).
- [42] C. W. Bauer, D. Pirjol, and I. W. Stewart, Phys. Rev. D 65, 054022 (2002).
- [43] M. Bauer, B. Stech, and M. Wirbel, Z. Phys. C 34, 103 (1987).
- [44] M. Wirbel, B. Stech, and M. Bauer, Z. Phys. C 29, 637 (1985).
- [45] A. Ali, G. Kramer, and C. D. Lu, Phys. Rev. D 59, 014005 (1998).
- [46] A. Ali, G. Kramer, and C. D. Lu, Phys. Rev. D 58, 094009 (1998).
- [47] A. Buras, J. Gerard, and R. Ruckl, Nucl. Phys. B268, 16 (1986).
- [48] B. Y. Blok and M. A. Shifman, Sov. J. Nucl. Phys. 45, 135 (1987).
- [49] B. Y. Blok and M. A. Shifman, Sov. J. Nucl. Phys. 45, 301 (1987).
- [50] B. Y. Blok and M. A. Shifman, Sov. J. Nucl. Phys. 45, 522 (1987).
- [51] L.-L. Chau and H.-Y. Cheng, Phys. Rev. D 36, 137 (1987).
- [52] L.-L. Chau and H.-Y. Cheng, Phys. Lett. B 222, 285 (1989).
- [53] J. L. Rosner, Phys. Rev. D 60, 114026 (1999).
- [54] C.-W. Chiang and J. L. Rosner, Phys. Rev. D 65, 054007 (2002).
- [55] C.-W. Chiang, Z. Luo, and J. L. Rosner, Phys. Rev. D 67, 014001 (2003).
- [56] B. Bhattacharya and J. L. Rosner, Phys. Rev. D 77, 114020 (2008).
- [57] B. Bhattacharya and J. L. Rosner, Phys. Rev. D 81, 014026 (2010).
- [58] B. Bhattacharya and J. L. Rosner, Phys. Rev. D 82, 037502 (2010).
- [59] H.-Y. Cheng and C.-W. Chiang, Phys. Rev. D 81, 074021 (2010).
- [60] L.-L. Chau and H.-Y. Cheng, Phys. Lett. B 333, 514 (1994).
- [61] Y.-L. Wu, M. Zhong, and Y.-F. Zhou, Eur. Phys. J. C 42, 391 (2005).
- [62] D. N. Gao, Phys. Lett. B 645, 59 (2007).
- [63] H. Lipkin, Phys. Rev. Lett. 44, 710 (1980).
- [64] J. F. Donoghue and B. R. Holstein, Phys. Rev. D 21, 1334 (1980).
- [65] C. Sorensen, Phys. Rev. D 23, 2618 (1981).
- [66] A. Kamal and E. Cooper, Z. Phys. C 8, 67 (1981).
- [67] C.-K. Chua and W.-S. Hou, Phys. Rev. D 72, 036002 (2005).
- [68] M. Gronau, D. London, and J. L. Rosner, Phys. Rev. D 87, 036008 (2013).
- [69] S. P. Rosen, Phys. Rev. Lett. 44, 4 (1980).
- [70] I. Bigi and M. Fukugita, Phys. Lett. 91B, 121 (1980).
- [71] J. F. Donoghue, Phys. Rev. D 33, 1516 (1986).
- [72] A. Kamal, N. Sinha, and R. Sinha, Z. Phys. C 41, 207 (1988).
- [73] H.-Y. Cheng, Eur. Phys. J. C 26, 551 (2003).
- [74] M. Beneke and M. Neubert, Nucl. Phys. B675, 333 (2003).
- [75] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, in 30th International Conference on High-Energy Physics (ICHEP 2000), 27 Jul–2 Aug 2000 Osaka, Japan (World)

Scientific, 2001), 882, http://ichep2000.hep.sci.osaka-u .ac.jp/.

- [76] H.-n. Li, C.-D. Lu, and F.-S. Yu, Phys. Rev. D 86, 036012 (2012).
- [77] P. Zenczykowski, Acta Phys. Pol. B 28, 1605 (1997).
- [78] J.-H. Lai and K.-C. Yang, Phys. Rev. D 72, 096001 (2005).
- [79] T. Becher and R. J. Hill, Phys. Lett. B 633, 61 (2006).
- [80] R. J. Hill, in Proceedings of the 4th Conference on Flavor Physics and CP Violation (FPCP), 9-12 Apr 2006, Vancouver, British Columbia, Canada, eConf no.: C06-04-09 (2006), 027.
- [81] D. Becirevic and A. B. Kaidalov, Phys. Lett. B 478, 417 (2000).
- [82] R. J. Hill, Phys. Rev. D 73, 014012 (2006).
- [83] C. Bernard, C. DeTar, M. Di Pierro, A. El Khadra, R. Evans *et al.*, Phys. Rev. D 80, 034026 (2009).
- [84] H. Na, C. T. H. Davies, E. Follana, G. P. Lepage, and J. Shigemitsu (HPQCD Collaboration), Phys. Rev. D 82, 114506 (2010).
- [85] H. Na, C. T. H. Davies, E. Follana, J. Koponen, G. P. Lepage, and J. Shigemitsu (HPQCD Collaboration), Phys. Rev. D 84, 114505 (2011).
- [86] S. Aoki, Y. Aoki, C. Bernard, T. Blum, G. Colangelo *et al.*, Eur. Phys. J. C 74, 2890 (2014).
- [87] H. Na, C. T. H. Davies, E. Follana, J. Koponen, G. P. Lepage, and J. Shigemitsu, Phys. Rev. D 84, 114505 (2011).
- [88] D. Besson *et al.* (CLEO Collaboration), Phys. Rev. D 80, 032005 (2009).
- [89] J. Yelton *et al.* (CLEO Collaboration), Phys. Rev. D 84, 032001 (2011).

- [90] J. Koponen, C. Davies, and G. Donald (HPQCD), arXiv:1208.6242.
- [91] G. S. Bali, S. Collins, S. Durr, and I. Kanamori, Phys. Rev. D 91, 014503 (2015).
- [92] K. Olive *et al.* (Particle Data Group), Chin. Phys. C **38**, 090001 (2014).
- [93] T. Feldmann, P. Kroll, and B. Stech, Phys. Rev. D 58, 114006 (1998).
- [94] F.-G. Cao, Phys. Rev. D 85, 057501 (2012).
- [95] C.-K. Chua, Nucl. Phys. A844, 260c (2010).
- [96] A. Kamal and R. Sinha, Phys. Rev. D 36, 3510 (1987).
- [97] F. Buccella, M. Forte, G. Miele, and G. Ricciardi, Z. Phys. C 48, 47 (1990).
- [98] F. Buccella, M. Lusignoli, G. Miele, and A. Pugliese, Z. Phys. C 55, 243 (1992).
- [99] F. Buccella, M. Lusignoli, G. Miele, A. Pugliese, and P. Santorelli, Phys. Rev. D 51, 3478 (1995).
- [100] M. Ablikim *et al.* (BESIII), Phys. Lett. B **734**, 227 (2014).
- [101] M. Gronau, Phys. Rev. D 91, 076007 (2015).
- [102] M. Albaladejo and J. A. Oller, Phys. Rev. Lett. 101, 252002 (2008).
- [103] R. Longacre, A. Etkin, K. Foley, W. Love, T. Morris *et al.*, Phys. Lett. B **177**, 223 (1986).
- [104] A. Biswas, A. Paul, and N. Sinha (to be published).
- [105] S. Fajfer, P. Singer, and J. Zupan, Eur. Phys. J. C 27, 201 (2003).
- [106] M. C. Arnesen, B. Grinstein, I. Z. Rothstein, and I. W. Stewart, Phys. Rev. Lett. 95, 071802 (2005).
- [107] K. Chetyrkin, J. H. Kuhn, and M. Steinhauser, Comput. Phys. Commun. 133, 43 (2000).