

Anti-de Sitter momentum spaceMichele Arzano,¹ Giulia Gubitosi,² João Magueijo,² and Giovanni Amelino-Camelia¹¹*Dipartimento di Fisica, Università La Sapienza and Sez. Roma1 INFN, P.le A. Moro 2, 00185 Roma, Italia*²*Theoretical Physics, Blackett Laboratory, Imperial College, London SW7 2BZ, United Kingdom*
(Received 8 December 2014; revised manuscript received 22 June 2015; published 16 July 2015)

We investigate the anti-de Sitter (AdS) counterpart to the well-studied de Sitter (dS) model for energy-momentum space, viz “ κ -momentum space” space (with a structure based on the properties of the κ -Poincaré Hopf algebra). On the basis of previous preliminary results one might expect the two models to be complementary: dS exhibiting an invariant maximal spatial momentum but unbounded energy, AdS a maximal energy but unbounded momentum. If that were the case AdS momentum space could be used to implement a principle of maximal Planck-scale energy, just as several studies use dS momentum space to postulate of maximal Planck-scale spatial momentum. However, several unexpected features are uncovered in this paper, which limit the scope of the expected complementarity, and interestingly they take different forms in different coordinatizations of AdS momentum space. “Cosmological” AdS coordinates mimic the dS construction used for κ -momentum space, and produce a Carroll limit in the ultraviolet. However, unlike the κ -momentum space, the boundary of the covered patch breaks Lorentz invariance, thereby introducing a preferred frame. In “horospherical” coordinates we achieve full consistency with frame independence as far as boost transformations are concerned, but find that rotational symmetry is broken, leading to an anisotropic model for the speed of light. Finally, in “static” coordinates we find a way of deforming relativistic transformations that successfully enforces frame invariance and isotropy, and produces a Carroll limit in the ultraviolet. Our results are also relevant for a long-standing debate on whether or not coordinate redefinitions in momentum space lead to physically equivalent theories: our three proposals are evidently physically inequivalent, leading to alternative models of Planck-scale effects. As a corollary we study the UV running of the Hausdorff dimension of momentum space in the first and third model, obtaining different results.

DOI: [10.1103/PhysRevD.92.024028](https://doi.org/10.1103/PhysRevD.92.024028)

PACS numbers: 04.20.-q, 04.60.-m

I. INTRODUCTION

In recent years findings in several areas of quantum-gravity research (see, e.g., Refs. [1,2] and references therein) have motivated the investigation of Planck-scale modified dispersion relations (MDRs), and this has attracted interest in MDRs as a possible avenue for Planck-scale phenomenology associated with astrophysical and cosmological observations [3–7]. It has become clear that some of the key predictions arising from MDRs depend crucially on whether the relevant framework *breaks* or merely *deforms* relativistic symmetries. A preferred-frame scenario is inevitable if the transformation laws between inertial observers remain the standard special-relativistic ones, since they only leave invariant the usual Einsteinian dispersion relation $E^2 - p^2 = m^2$. However, it is possible to introduce a deformation of relativistic symmetries preserving the equivalence of reference frames and leaving the MDR observer-independent [8–10]. Notable examples of such “DSR” (doubly-special, or deformed-special relativity) scenarios include theories based on a maximally-symmetric curved momentum space. This has been investigated in great detail if momentum space has de Sitter geometry. Here we seek to investigate momentum space with anti-de Sitter geometry, a possibility which has so far received very little attention in the literature (see, however, [11,12]). In doing so

we will uncover several significant differences between dS and AdS models of momentum space.

If DSR-relativistic scenarios arise from maximally-symmetric momentum space it is easy to see how one can achieve compatibility between some MDRs and the laws of transformation between inertial observers. One usually introduces ordinary special relativity by taking as the starting point the isometries of Minkowski spacetime, but one could equally well start from the isometries of Minkowski momentum space. In either case one can derive the transformation laws of momenta and spacetime coordinates by consistency [13]. Since the isometries of de Sitter (or anti-de Sitter) space can be seen as a deformation of the isometries of Minkowski space, any set of transformation laws derived from the isometries of de Sitter (or anti-de Sitter) momentum space is as “relativistic” as special relativity (i.e. it abides by the principle of the relativity of inertial frames). However, such a construction entails a deformation of the transformation laws between inertial observers, and these will leave invariant a modified (deformed) dispersion relation.

Constructions based on de Sitter momentum space have been extensively studied in the literature, with many authors registering the expectation that the counterpart AdS model would have properties easily obtainable from those of dS momentum space. Several arguments suggest

that the two models should be complementary, with dS exhibiting an invariant maximal spatial momentum but unbounded energy, and AdS a maximal energy but unbounded spatial momentum. However, as we will show in this paper, many crucial novelties arise in AdS curved momentum space that are not captured by this expected “complementarity.” Whereas previous arguments focused exclusively on local properties of the two momentum spaces, one of the key ingredients of our analysis is the realization of the fact that different coordinates cover different patches of the manifold, and that this leads to different physical statements on what is the free theory. A number of options appear, mimicking—or not—constructions previously considered for dS. We will find that in all of them AdS momentum space is *qualitatively* very different from dS, the main point made in this paper.

An important reference for us is the so-called “ κ -momentum space,” a coordinatization of a certain patch of de Sitter momentum space which has been found to have remarkably good relativistic properties, and can be inspired by the formal structure of the κ -Poincaré Hopf algebra [14–17].¹

As we observed in Ref. [18], κ -momentum space can be viewed as the momentum space equivalent of the “cosmological” representation of dS spacetime. While our focus here is on the AdS momentum space, an important source of intuition for us is provided by these previous results on the dS momentum space, and specifically the κ -momentum-space construction, so we find appropriate to review those results in Sec. II. Then in Sec. III A we find the corresponding construction for AdS. In such “cosmological” coordinates a simple representation for the Casimir invariant, momentum space metric and integration measure is found. However, unlike with dS, the boundary of the covered patch breaks Lorentz invariance. If analyzed only at the level of infinitesimal transformations the model is DSR-relativistic, but a breakdown of Lorentz invariance is noticed when considering finite Lorentz transformations. We know of no previous examples in the literature of such subtle breakdown of relativistic symmetries (see, however [19]).

Another possible approach mimicking the κ -momentum space consists of using “horospherical” coordinates, which cover a patch of AdS. We do this in Sec. III B, only to

¹The main point is that on the portion of de Sitter momentum space covered by such “ κ -coordinates” the action of relativistic symmetry generators on the momentum manifold can be interpreted in terms of the action of the quantum κ -Poincaré algebra. Some nontrivial properties of the κ -Poincaré algebra, namely the antipode and coproduct of the algebra elements, have a role in the definition of the conservation laws in interactions of particles whose momentum space is this “ κ -momentum space.” Here, we will not be concerned with this last aspect, as we are focusing on the single-particle properties of momentum space. For the same reason, we will not look for possible quantum-algebraic structures associate to AdS momentum space. In fact, in order to do this, we would need to define the properties of multiparticle systems as well, which goes beyond the scope of this work.

encounter a similar problem to that found for cosmological coordinates, but this time regarding the rotations. Similarly to what happens for boosts in “cosmological” coordinates, the boundary of the patch covered by horospherical coordinates breaks invariance under rotations, and so the theory is anisotropic. The ensuing formalism is somewhat awkward, and the expression for the Casimir is far more complex. However, we argue that this could be a good model for encoding anisotropic MDRs and speed of light. We should however bear in mind some potential pathologies: the model does not allow one spatial momentum to take arbitrary negative values if we want to preserve invariance under finite boosts.

In view of the symmetry breaking properties of these two models, in Sec. III C we investigate an alternative construction which does not purport to mimic the κ -momentum space. We propose a set of coordinates analogous to “static” coordinates in the spacetime picture. They cover the whole of AdS and do not break Lorentz invariance in any way. They lead to simple expressions for the metric, Casimir and integration measure. As with the first model, we find a Carroll limit in the UV, i.e.: the speed of light goes to zero in the UV.

As an application, in Sec. IV we briefly investigate the issue of running of dimensionality for the first and third model (the matter is far less obvious for the second model, due to its anisotropy). We do this by choosing linearizing coordinates and evaluating the *measure of integration on momentum space* (a procedure described in [18,20], known to match the spectral dimension in the UV limit in all cases studied so far). We find that the two models exhibit running to different dimensions, a particularly transparent indication of the fact that they are physically distinct models, though both based on AdS momentum space.

Given that static coordinates have not been considered for dS, for completeness in Sec. V we present them. We find that they break Lorentz invariance in a fashion similar to that found for AdS in cosmological coordinates. We also examine running of dimensionality in the corresponding model, finding a very suggestive result. In a concluding section we collect the main results of this paper and discuss their implications.

II. DE SITTER MOMENTUM SPACE

As mentioned in the Introduction the action of relativistic symmetries on momenta can be *deformed* if one considers a maximally symmetric curved momentum space. A widely studied example of deformed Poincaré symmetries reflecting such nontrivial geometry of momentum space is the so-called κ -Poincaré algebra [14,15,17]. Indeed, as first shown in [21], in a κ -deformed framework momenta can be seen as coordinates on a portion of de Sitter momentum space defined as a four-dimensional hypersurface:

$$-P_0^2 + P_1^2 + P_2^2 + P_3^2 + P_4^2 = \frac{1}{\ell^2}, \quad (1)$$

embedded in five dimensional Minkowski space, with line element:

$$ds^2 = -dP_0^2 + dP_1^2 + dP_2^2 + dP_3^2 + dP_4^2, \quad (2)$$

selected by the inequality

$$P_0 - P_4 > 0, \quad (3)$$

where the ‘‘cosmological constant’’ is the inverse of the parameter which governs the deformation of the algebraic structures in κ -Poincaré, $\kappa = 1/\ell$. The natural parametrization of this submanifold, inherited by the bicrossproduct basis of the κ -Poincaré algebra [22], is given by *bicrossproduct coordinates*, which correspond in position space to the ‘‘cosmological’’ or ‘‘flat slicing’’ rendition of de Sitter space. They are related to the embedding coordinates via:

$$\begin{aligned} P_0(E, \vec{p}) &= \frac{\sinh(\ell E)}{\ell} + \frac{\ell p^2}{2} e^{\ell E}, \\ P_i(E, \vec{p}) &= -p_i e^{\ell E}, \\ P_4(E, \vec{p}) &= -\frac{\cosh(\ell E)}{\ell} + \frac{\ell p^2}{2} e^{\ell E}, \end{aligned} \quad (4)$$

where $p \equiv |\vec{p}|$. With these coordinates the line element takes the familiar ‘‘cosmological’’ de Sitter metric form:

$$ds^2 = -dE^2 + e^{2\ell E} \sum_{j=1}^3 dp_j^2 \quad (5)$$

from which it is easy to infer the invariant integration measure in momentum space:

$$d\mu(E, \vec{p}) = e^{3\ell E} p^2 dE dp. \quad (6)$$

The deformed mass-shell is given by the intersection of a plane $P_4 = \text{const}$ with the momentum manifold:

$$-P_0^2 + \vec{P}^2 = \frac{1}{\ell^2} - P_4^2 = m^2. \quad (7)$$

In Fig. 1 we show the κ -momentum space and the mass-shells given by the above constraint. In the massless case, using the relations above, it can be shown that the mass-shell condition reads

$$\mathcal{C} \left(1 + \frac{\ell^2 \mathcal{C}}{4} \right) = 0, \quad (8)$$

where \mathcal{C} is the Casimir invariant of the κ -Poincaré algebra in bicrossproduct coordinates:

$$\mathcal{C} = -\frac{4}{\ell^2} \sinh^2(\ell E/2) + e^{\ell E} p^2. \quad (9)$$

Looking at the mass-shell, it is clear that one has to perform a further restriction on the allowed range for the embedding coordinates in order for the theory to be relativistic. In fact, a crucial request is that any mass-shell is completely within the allowed portion of de Sitter momentum space. Failure to meet this condition would result in the possibility for finite boosts to bring outside the allowed region some




FIG. 1 (color online). The portion of (2-dimensional) de Sitter momentum space known as the ‘‘2D κ -momentum space.’’ The blue plane is defined by the condition $P_0 - P_4 = 0$, so the κ -momentum space is on the upper-left side of the plane. The red lines represent the mass-shells, defined by the condition $P_4 = \text{const}$. In order to have the mass-shells completely within the allowed region one has to further restrict to $P_0 > 0$ and $P_4 < 0$.

points that were originally within it. The restriction one has to enforce is given by the conditions

$$P_0 > 0, \quad P_4 < 0. \quad (10)$$

However let us mention that at a field theoretic level the Hopf algebraic structures of the κ -Poincaré algebra ensure that the model is fully consistent without the further restriction above [23].

A. Running of Hausdorff dimension of momentum space for the κ -momentum space scenario

In [18] we showed that the κ -momentum space is characterized by a running of its Hausdorff dimension when going from the IR regime to the UV. We considered a general $D + 1$ de Sitter manifold, and we allowed for the mass-shell to be a generic function of the κ -Poincaré Casimir, parametrized as $m^2 = \mathcal{C}(1 + \ell^{2\gamma} \mathcal{C}^\gamma)$. Here we review the argument found in [18] for UV dimensional running, specializing to the $D = 3$, $\gamma = 1$ case, which is the one discussed in the previous subsection (the exact coefficient of the UV-dominant term in the mass-shell relation does not affect the UV value of the Hausdorff dimension). In doing so, we will add some remarks that will facilitate comparison with the AdS constructions.

The phenomenon of dimensional running can be characterized by choosing a set of ‘‘linearizing coordinates,’’ rendering the dispersion relations trivial in the UV, and examining the dimensionality associated with the integration measure in such coordinates. The linearizing coordinates for \mathcal{C} , are [18]

$$\begin{aligned}\tilde{E} &= \frac{2 \sinh(\ell E/2)}{\ell} \\ \tilde{p}_i &= p_i e^{\ell E/2},\end{aligned}\quad (11)$$

which in the UV limit (defined as $E \rightarrow \infty$ and $p \rightarrow 1/\ell$) become:

$$\begin{aligned}\tilde{E} &\approx \frac{e^{\ell E/2}}{\ell} \\ \tilde{p}_i &\approx \frac{e^{\ell E/2}}{\ell}\end{aligned}\quad (12)$$

(we note that in the UV limit $\tilde{E} \approx \tilde{p}$, even off-shell). In terms of the new coordinates the measure (6) is given by:

$$d\mu = \tilde{E}^2 \tilde{p}^2 d\tilde{E} d\tilde{p}. \quad (13)$$

As explained in [18], for on shell relations which in the UV limit take the form $\mathcal{C}^{1+\gamma}$, one finds

$$d_H = \frac{6}{1+\gamma}, \quad (14)$$

so in the case of interest here ($\gamma = 1$) one finds that the Hausdorff dimension runs to 3 in the UV.

Notice that we could obtain the same result by linearizing directly the on-shell relation that comes out of the κ -momentum space construction, Eq. (7). This amounts to choosing the embedding coordinates themselves as linearizing coordinates. In the UV, their relation to the bicross-product coordinates is

$$\begin{aligned}\tilde{E} = P_0 &\approx \frac{e^{\ell E}}{2\ell} (1 + \ell^2 p^2) \\ \tilde{p}_i = P_i &\approx p_i e^{\ell E}\end{aligned}\quad (15)$$

where the approximate signs refer to the UV limit approximation. We note also here that in the UV limit $\tilde{E} \approx \tilde{p}$, even off-shell. The measure (6) in the new coordinates now reads:

$$d\mu = \frac{\tilde{p}^2}{\tilde{E}} d\tilde{p} d\tilde{E} \quad (16)$$

from which we can directly read $d_H = 3$.² The last description will be useful in establishing a comparison with AdS constructions. It implies that if we take the MDR that comes

²In the more general $D + 1$ -dimensional case, and allowing for redefinitions of the mass-shell with UV limit $m^2 = \ell^{2\gamma} (P_0^2 - \vec{P}^2)^{1+\gamma}$ one would get

$$d_H = \frac{D}{1+\gamma}. \quad (17)$$

Note that this is another example of correspondence between the UV Hausdorff dimension of momentum space and the UV limit of the spectral dimension. The second one was computed in [24] for the 4- and 3- dimensional cases, and the results are in agreement with formula (17).

most naturally out of dS [i.e., Eq. (7)] then we would observe dimensional reduction from $D + 1$ to D . This can be equivalently obtained from \mathcal{C} with $\gamma = 1$.

III. ADS MOMENTUM SPACE

As with dS space, AdS momentum space can be described as a four-dimensional hypersurface embedded in a five-dimensional flat space, this time with signature $-, -, +, +, +$. The submanifold is now defined by:

$$-P_0^2 + P_1^2 + P_2^2 + P_3^2 - P_4^2 = -\frac{1}{\ell^2} \quad (18)$$

and the corresponding line element is

$$ds^2 = -dP_0^2 + dP_1^2 + dP_2^2 + dP_3^2 - dP_4^2. \quad (19)$$

A. Cosmological coordinates for AdS

In analogy with the κ -momentum-space construction over dS we seek coordinates casting a portion of AdS in the form of a cosmological metric (which is no longer a ‘‘flat slicing,’’ as it was for dS). It can be shown (see the Appendix) that the cosmological AdS coordinates are defined by the following relation with the embedding coordinates:

$$\begin{aligned}P_0(E, \vec{p}) &= \frac{1}{\ell} \sin \ell E \\ P_1(E, \vec{p}) &= p_1 \cos(\ell E) \\ P_2(E, \vec{p}) &= p_2 \cos(\ell E) \\ P_3(E, \vec{p}) &= p_3 \cos(\ell E) \\ P_4(E, \vec{p}) &= \frac{1}{\ell} \sqrt{1 + (\ell p)^2} \cos \ell E.\end{aligned}\quad (20)$$

In these coordinates the metric reads:

$$ds^2 = -dE^2 + \cos^2(\ell E) \left(\frac{dp^2}{1 + \ell^2 p^2} + p^2 d\Omega^2 \right). \quad (21)$$

The submanifold covered by these coordinates is defined by the constraint $-1/\ell \leq P_0 \leq 1/\ell$, or, if we require the energy to be positive, $0 \leq P_0 \leq 1/\ell$. Using the line element (21) we easily deduce the invariant integration measure for AdS momentum space in these coordinates:

$$d\mu(E, p) = \frac{\cos^3(\ell E)}{\sqrt{1 + \ell^2 p^2}} p^2 dE dp. \quad (22)$$

In analogy with dS, the mass-shell relation can be inferred by imposing $P_4 = \text{const}$ upon the surface condition:




FIG. 2 (color online). Portion of AdS momentum space covered by cosmological coordinates. The condition for the allowed region is $-1/\ell < P_0 < 1/\ell$, which is the portion of the AdS manifold between the two blue planes. The red lines represent the mass-shells, defined by the condition $P_4 = \text{const}$.

$$-P_0^2 + \vec{P}^2 = -\frac{1}{\ell^2} + P_4^2 = -m^2. \quad (23)$$

From this we see that we must require that $m \leq 1/\ell$. In terms of the cosmological AdS coordinates the mass-shell condition takes the form:

$$-\frac{1}{\ell^2} \sin^2 \ell E + p^2 \cos^2(\ell E) = -m^2. \quad (24)$$

In Fig. 2 we plot the submanifold of AdS covered by cosmological coordinates as well as the mass-shells given by the constraint (23).

1. Maximal energy and speed of light in the UV limit

The mass-shell condition given by Eq. (24) implies the presence of a maximal energy in the theory (just like on the κ -momentum space there is a maximal spatial momentum). Let us consider a massless particle, in this case the dispersion relation is given by

$$\frac{1}{\ell} \tan(\ell E) = p \quad (25)$$

and it is evident that

$$E \leq E_{\text{max}} = \frac{\pi}{2\ell}, \quad (26)$$

whereas there is no maximal spatial momentum. In addition, we see that the speed of light goes to zero as $p \rightarrow \infty$

$$c = \frac{dE}{dp} \rightarrow 0. \quad (27)$$

This is nothing but the Carroll limit [25]. For massive particles the mass shell can be written as:

$$\cos^2 \ell E = \frac{1 - \ell^2 m^2}{1 + \ell^2 p^2} \quad (28)$$

(remember that the constraint $m < 1/\ell$ must be satisfied). As $p \rightarrow \infty$ again we get $E \rightarrow E_{\text{max}}$. Notice that if $m = 1/\ell$ then the MDR does not fix the momentum, and the energy saturates. A plot of the behavior of the MDR is shown in Fig. 3.

2. Violation of Lorentz invariance

Despite the care taken not to introduce a preferred frame, this has in fact sneaked in by virtue of the fact that the boundary of the submanifold is not invariant under the action of the Lorentz group. This is a crucial difference between the analogous constructions for dS (for which the boundary is $P_0 = P_4$) and AdS (where the boundary is $|P_0| = 1/\ell$). It might seem a subtle point, but the implications are obvious if we write down the Lorentz transformations in momentum space.

These can be described as a nonlinear representation of the Lorentz group, inferred from the standard ones as applied to the embedding coordinates through the relations (20).

For a finite boost in the $\hat{1}$ direction, the explicit transformation rules are

$$E' = \frac{1}{\ell} \arcsin \left(\gamma \ell \left(\frac{1}{\ell} \sin(\ell E) - v p_1 \cos(\ell E) \right) \right), \quad (29)$$

$$p'_1 = \frac{\gamma (p_1 \cos(\ell E) - v \frac{1}{\ell} \sin(\ell E))}{\sqrt{1 - \ell^2 \gamma^2 \left(\frac{1}{\ell} \sin(\ell E) - v p_1 \cos(\ell E) \right)^2}}, \quad (30)$$

$$p'_2 = \frac{p_2 \cos(\ell E)}{\sqrt{1 - \ell^2 \gamma^2 \left(\frac{1}{\ell} \sin(\ell E) - v p_1 \cos(\ell E) \right)^2}}, \quad (31)$$




FIG. 3 (color online). Dispersion relation for a massive particle (28) in AdS momentum space with cosmological coordinates, with $\ell = 1$ and $m = 0.1$.

$$p'_3 = \frac{p_3 \cos(\ell E)}{\sqrt{1 - \ell^2 \gamma^2 \left(\frac{1}{\ell} \sin(\ell E) - v p_1 \cos(\ell E)\right)^2}}. \quad (32)$$

These can be shown to be generated by:

$$L_{01} = p_1 \cos(\ell E) \partial_E + \frac{1}{\ell} \tan(\ell E) \partial_{p_1}. \quad (33)$$

It is obvious that there is something pathological with the finite transformations: as the transformation for the energy shows, there is clearly a maximal boost parameter, such that any larger boost would bring the value of energy outside the allowed range.

Another way to see that this framework is not invariant under finite boosts is by noticing that any mass shell that goes through the allowed region of the AdS manifold is not completely included within that region (see Fig. 2), and there is no way to further restrict the allowed region so as to solve the problem. This means that for any value of the mass and energy, there will always be a finite boost pushing the particle outside the allowed range of parameters.

Yet another indicator of the breakdown of relativistic invariance is the fact that not only is the energy E bounded, but also the embedding one:

$$P_0 \leq P_{0\max} = \frac{1}{\ell} \quad (34)$$

and this is clearly incompatible with its standard transformation rules under boosts. This is in sharp contrast with the corresponding situation in de Sitter space, where p is bounded, but not its embedding counterpart.

B. Horospherical coordinates

An AdS coordinate system which mimics more closely the dS properties of the κ -momentum space is given by the so-called *horospherical coordinates* [26,27]. In terms of the embedding coordinates they read

$$\begin{aligned} P_0 &= \frac{1}{\ell} \cosh(\ell k_0) + \frac{\ell}{2} e^{\ell k_0} k_i k^i, \\ P_4 &= e^{\ell k_0} k_1, \\ P_2 &= e^{\ell k_0} k_2, \\ P_3 &= e^{\ell k_0} k_3, \\ P_1 &= \frac{1}{\ell} \sinh(\ell k_0) - \frac{\ell}{2} e^{\ell k_0} k_i k^i, \end{aligned} \quad (35)$$

where now $k_i k^i = -k_1^2 + k_2^2 + k_3^2$. It is easy to verify that they satisfy constraint (18) but only cover the $P_0 + P_1 > 0$ region, half AdS (see [27]). The *spurious* embedding coordinate has to be timelike in this case and indeed it is easily verified that this must be P_0 since it diverges for $\ell \rightarrow 0$, the flat momentum space limit.

Let us note that k_0 is now one of the components of the spatial momentum, and k_1 is the energy. It is physically more transparent to write the new coordinates as:

$$\begin{aligned} P_0 &= \frac{1}{\ell} \cosh(\ell p_1) + \frac{\ell}{2} e^{\ell p_1} (-E^2 + p_2^2 + p_3^2), \\ P_4 &= e^{\ell p_1} E, \\ P_2 &= e^{\ell p_1} p_2, \\ P_3 &= e^{\ell p_1} p_3, \\ P_1 &= \frac{1}{\ell} \sinh(\ell p_1) - \frac{\ell}{2} e^{\ell p_1} (-E^2 + p_2^2 + p_3^2). \end{aligned} \quad (36)$$

In such coordinates the line element is given by

$$ds^2 = e^{2\ell p_1} (-dE^2 + dp_2^2 + dp_3^2) + dp_1^2, \quad (37)$$

and the associated integration measure is

$$d\mu = e^{3\ell p_1} dE dp. \quad (38)$$

Analogously to what we have done in the previous subsections, we find the mass-shell condition by requiring that the spurious timelike coordinate is constant, which in this case amounts to asking $P_0 = \text{const}$:

$$P_0^2 - \frac{1}{\ell^2} = -\vec{P}^2 + P_4^2 = -m^2. \quad (39)$$

Also here this implies that the mass cannot be arbitrarily large, $m \leq \frac{1}{\ell}$. In terms of the embedding coordinates the mass-shell condition reads:

$$\begin{aligned} -m^2 &= -e^{2\ell p_1} E^2 + e^{2\ell p_1} (p_2^2 + p_3^2) \\ &+ \left(\frac{1}{\ell} \sinh(\ell p_1) - \frac{\ell}{2} e^{\ell p_1} (-E^2 + p_2^2 + p_3^2) \right)^2. \end{aligned} \quad (40)$$

In Fig. 4 we plot the submanifold of AdS covered by horospherical coordinates, as well as the mass-shells. Despite the obvious anisotropy introduced by these coordinates, a deformed description of rotations exists, and can be derived in analogy of what was done for boosts in cosmological coordinates. This means that the submanifold is invariant under infinitesimal transformations. However, one can see that the further restriction of the manifold to the $P_1 > 0$, $P_0 > 0$ region has to be enforced in order to have worldlines not exiting the manifold (i.e. to have invariance under finite boosts). This in turn leads to a breakdown of invariance under rotations, because of the condition $P_1 > 0$ due to the fact that embedding coordinates transform according to the standard Lorentz transformations.




FIG. 4 (color online). Portion of AdS momentum space covered by horospherical coordinates. The condition for the allowed region is $P_0 + P_1 > 0$, which is the portion of the AdS manifold on the lower-right side of the blue plane. The mass-shell is given by $P_0 = \text{const.}$ and is in red. Remember that now the P_4 coordinate is the one related to energy and P_1 is a spatial coordinate.

1. Anisotropic speed of light

It is interesting to look at the behavior of the speed of light in this model. For notational simplicity we restrict to the case of 2 + 1-dimensional AdS momentum space and we write the spatial momenta in polar coordinates $p_1 = p \cos \theta$, $p_2 = p \sin \theta$. The dispersion relation for a massless particle reads

$$E^2 = \frac{1}{2\ell^2} (2e^{-2\ell p \cos \theta} + (2 + \ell^2 p^2 \sin^2 \theta) - e^{-\ell p \cos \theta}) \times \sqrt{16 - p^2 \ell^2 \sin^2 \theta (4 + e^{2\ell p \cos \theta} (4 + 3\ell^2 p^2 \sin^2 \theta))}. \quad (41)$$

The deformation of rotations which guarantee local invariance of the manifold leads to a direction-dependent dispersion relation and as a consequence to a direction-dependent speed of light.

The general expression is quite complicated (a plot of its angular dependence can be seen in Fig. 5), so here we only write down the two special cases $\theta = 0$ (speed of light along the p_1 direction) and $\theta = \pi/2$ (speed of light along the p_2 direction)

$$c(p, \theta = 0) = e^{-\ell p} \quad (42)$$

$$c(p, \theta = \pi/2) = \frac{\ell p (4 + 3\ell^2 p^2 + A)}{A \sqrt{8 + 2\ell^2 p^2 - 2A}} \quad (43)$$

where $A = \sqrt{16 - 8\ell^2 p^2 - 3\ell^4 p^4}$. Note that the speed of light becomes imaginary whenever p and θ are such that




FIG. 5 (color online). Speed of light in horospherical coordinates as a function of θ , for $\ell = 1$, $p = 0.1$.

the condition $P_1 > 0$ is violated. This makes it impossible to reach infinite speed of light in any direction. Strange as this model might be it could be a good framework for encoding fundamental anisotropy, with constraints on it encapsulating observational facts, like those derived from modern day versions of the Michelson-Morley experiment. Occasional claims for cosmological anisotropy (e.g. [28]) could also be embedded in this model, but the matter is beyond the scope of this paper.

C. “Static” coordinates

Static coordinates cover the full AdS manifold, and so clearly they do not break Lorentz invariance, but rather deform it. They can be defined from the embedding coordinates via:

$$\begin{aligned} P_0 &= \frac{1}{\ell} \sin(\ell E) \cosh(\ell p'), \\ P_r &= \frac{1}{\ell} \sinh(\ell p'), \\ P_4 &= \frac{1}{\ell} \cos(\ell E) \cosh(\ell p'), \end{aligned} \quad (44)$$

where $P_r \equiv \sqrt{P_1^2 + P_2^2 + P_3^2}$. The line element in these coordinates is

$$ds^2 = -\cosh^2(\ell p') dE^2 + dp'^2 + \frac{1}{\ell^2} \sinh^2(\ell p') d\Omega. \quad (45)$$

We can also use an areal coordinate:

$$p = \frac{\sinh(\ell p')}{\ell} \quad (46)$$

resulting in:

$$\begin{aligned}
P_0 &= \frac{1}{\ell} \sin(\ell E) \sqrt{1 + (\ell p)^2}, \\
P_r &= p, \\
P_4 &= \frac{1}{\ell} \cos(\ell E) \sqrt{1 + (\ell p)^2},
\end{aligned} \tag{47}$$

for which the line element is

$$ds^2 = -(1 + (\ell p)^2) dE^2 + \frac{dp^2}{1 + (\ell p)^2} + p^2 d\Omega. \tag{48}$$

Notice that invariance under (deformed) Lorentz transformations is not spoiled if we ask the energy to be positive, i.e. this requirement is compatible with the deformed transformation rules. This can be seen from (47): enforcing the positivity of E is equivalent³ to enforcing the positivity of P_0 . But the embedding coordinate P_0 transforms with the standard Lorentz transformations, so asking it to be positive works in the same way as in the usual special relativistic case. Another interesting feature of such coordinates is that the integration measure is undeformed:

$$d\mu = p^2 dp dE. \tag{49}$$

We can find the mass-shell relation again by requiring that $P_4 = \text{const}$:

$$-P_0^2 + \vec{P}^2 = -\frac{1}{\ell^2} + P_4^2 = -m^2, \tag{50}$$

which in static coordinates becomes:

$$-\frac{\sin^2(\ell E)}{\ell^2} [1 + (\ell p)^2] + p^2 = -m^2. \tag{51}$$

In order to explore the physics and UV limit we consider massless particles. We see that their spatial momentum is unbounded, but their energy tends to a maximum:

$$p \rightarrow \infty \tag{52}$$

$$E \rightarrow E_{\text{max}} = \frac{\pi}{2\ell}. \tag{53}$$

We take these limiting values as the UV limit of the model. The speed of light is given by:

$$c = \frac{dE}{dp} = \frac{1}{1 + (\ell p)^2} \tag{54}$$

and in the UV limit this goes to zero, what is known in the literature as the *Carroll limit* [25].

³In this work we always assume $\ell > 0$.

IV. UV DIMENSIONAL REDUCTION IN ADS MOMENTUM SPACE

In a series of recent papers [18,20] we have shown that it is possible to characterize the phenomenon of dimensional reduction in the UV dispensing with the concept of spectral dimension altogether. This is beneficial, as the latter appeals to a fictitious time parameter, requires the Euclideanization of the space, and is not always based on a properly defined probability distribution. Instead we showed that we could transfer all the nontrivial effects of the MDRs into the measure, adopting linearizing variables and then study the Hausdorff dimension of the energy-momentum space in these variables. This is a physically clearer procedure, and asymptotically (i.e. in the deep UV limit) it produces results coinciding with those using the spectral dimension in all known cases. Given the difficulties in defining asymptotic spectral dimension for AdS momentum space, we favor our procedure here.

As with [18,20] we shall be concerned with MDRs which in the UV limit have the form:

$$\Omega = f(\mathcal{C}) \approx \mathcal{C}^{1+\gamma} \tag{55}$$

where \mathcal{C} is the Casimir invariant of the theory. We will examine the UV running of the Hausdorff dimension for cosmological and static coordinates which have a clear UV limit leaving aside the case of horospherical coordinates whose anisotropic nature renders the notion of UV limit ambiguous.

A. Cosmological coordinates

Following [18,20], we find linearizing variables in 2 steps: first by assuming $\gamma = 0$, then generalizing to $\gamma \neq 0$. When $\gamma = 0$ (i.e. when the MDRs are just the Casimir) the linearizing coordinates are just the embedding coordinates, as in (20): $\tilde{E} \equiv P_0$, $\tilde{p} \equiv P$. In $D + 1$ space-time dimensions the momentum space integration measure in such variables is

$$d\tilde{\mu} = \frac{\tilde{p}^{D-1}}{\sqrt{1 + \ell^2(\tilde{p}^2 - \tilde{E}^2)}} d\tilde{E} d\tilde{p}, \tag{56}$$

which, in the UV limit (as defined above), becomes

$$d\tilde{\mu} \approx \tilde{p}^{D-2} d\tilde{E} d\tilde{p}. \tag{57}$$

Therefore the Hausdorff dimension is reduced by 1. We note that this is just the general measure studied in [18]:

$$d\mu(\tilde{E}, \tilde{p}) \propto \tilde{p}^{D_x-1} \tilde{E}^{D_t-1} d\tilde{E} d\tilde{p} \tag{58}$$

with values $D_t = 1$, and $D_x = D - 1$.

If $\gamma \neq 0$ the linearizing coordinates can be found by following the ‘‘step 2’’ described in [18], but we stress that the procedure here does not rely on Euclideanization. In

[18] we were dealing with an Euclideanized version of momentum space (because we wanted to study the *asymptotic* coincidence of spectral and Hausdorff dimensions), but the procedure carries through with a minimal adaptation if we remain Lorentzian. All we need to do is introduce hyperbolic (instead of spherical) polar coordinates:

$$\tilde{E} = r \cosh \theta \quad (59)$$

$$\tilde{p} = r \sinh \theta \quad (60)$$

so that the MDRs become

$$\Omega = r^{2(1+\gamma)}. \quad (61)$$

We can then define a linearizing variable

$$\hat{r} = r^{1+\gamma} \quad (62)$$

such that:

$$d\mu \propto \hat{r}^{\frac{D_t+D_x}{1+\gamma}-1} (\cos \theta)^{D_t-1} (\sin \theta)^{D_x-1} d\hat{r} d\theta \quad (63)$$

leading to the conclusion that in the UV:

$$d_H = \frac{D_t + D_x}{1 + \gamma}. \quad (64)$$

For the AdS model we are considering, this therefore becomes:

$$d_H = \frac{D}{1 + \gamma}. \quad (65)$$

B. Static coordinates

Similarly to what happens for the cosmological coordinates, the linearizing coordinates for the model are the embedding coordinates found in (47). The integration measure is the same as (56). However the UV limit now entails $\tilde{E} \approx \tilde{p}$ leading to an undeformed measure. This model therefore has nontrivial physical effects (e.g. it has a Carroll limit) but it does *not* present running of the dimensionality, if $\gamma = 0$. If $\gamma \neq 0$ one can straightforwardly calculate

$$d_H = \frac{1 + D}{1 + \gamma}, \quad (66)$$

and we therefore have a nontrivial running of the dimensionality.

V. DE SITTER MOMENTUM SPACE IN STATIC COORDINATES

The first two models above arise from attempts to construct four-momenta defined on AdS space using a *duality* approach to dS space of momenta associated to κ -Poincaré. The third model based on “static coordinates,” however, was proposed without reference to a dS construction, so one might wonder what the equivalent dS model would be. As we shall see, whilst static coordinates lead to an AdS momentum space model which does not break any symmetry, its dS counterpart breaks Lorentz invariance.

Static coordinates for dS may be built from:

$$\begin{aligned} P_0 &= \frac{1}{\ell} \sinh(\ell E) \sqrt{1 - (\ell p)^2}, \\ P_r &= p, \\ P_4 &= \frac{1}{\ell} \cosh(\ell E) \sqrt{1 - (\ell p)^2}, \end{aligned} \quad (67)$$

leading to metric:

$$ds^2 = -(1 - (\ell p)^2) dE^2 + \frac{dp^2}{1 - (\ell p)^2} + p^2 d\Omega, \quad (68)$$

and an undeformed integration measure. The Casimir is

$$\mathcal{C} = -\frac{\sinh^2(\ell E)}{\ell^2} [1 - (\ell p)^2] + p^2 = m^2, \quad (69)$$

and we see that the theory has a maximum spatial momentum, $p_{\max} = 1/\ell$, but unbounded energy, just like the κ -Poincaré case. (This maximal momentum coincides with the location of de Sitter’s horizon, in the counterpart position space version of the space.) The UV limit may be accordingly defined by:

$$p \rightarrow p_{\max} = \frac{1}{\ell} \quad (70)$$

$$E \rightarrow \infty. \quad (71)$$

and in such limit the speed of light

$$c = \frac{dE}{dp} = \frac{1}{1 - (\ell p)^2} \quad (72)$$

goes to infinity in the UV.

All of these features are very similar to what is found in κ -Poincaré in the *bicrossproduct basis*. However this model breaks Lorentz symmetry in a way that mimics closely what happens for cosmological AdS coordinates. Indeed looking at the embedding coordinates (67), it is clear that the maximum value of momentum is reflected into a maximum value of the embedding coordinate P_r . Since the embedding coordinates transform according to standard

Lorentz transformations, this is inconsistent with the relativity of inertial frames.

We conclude by noticing that this model also exhibits running of dimensionality. Working out the integration measure in linearizing coordinates, $\tilde{E} \equiv P_0$, $\tilde{p} \equiv P_r$, leads to:

$$d\tilde{\mu} = \frac{\tilde{p}^{D-1}}{\sqrt{1 - \ell^2(\tilde{p}^2 - \tilde{E}^2)}} d\tilde{E}d\tilde{p}, \quad (73)$$

that in the UV limit becomes

$$d\tilde{\mu} \approx \frac{\tilde{p}^{D-1}}{\tilde{E}} d\tilde{E}d\tilde{p}. \quad (74)$$

Following our standard calculation, we find for the UV Hausdorff dimension of momentum space

$$d_H = \frac{D}{1 + \gamma}. \quad (75)$$

This is suggestively similar to what we found for AdS in cosmological coordinates. It also matches the result obtained for dS linearizing directly from the embedding coordinates, as discussed in Sec. II.

VI. CONCLUSIONS

In this paper we took a first stab at defining a curved momentum space based on AdS geometry, in analogy with previous work for dS space. A number of significant novelties were uncovered in the process.

The equivalent of the bicrossproduct basis was sought in two ways. First, we noted that we can regard the bicrossproduct basis as the momentum space counterpart of the ‘‘cosmological’’ covering of dS, and sought similar coordinates for AdS. We found that the equivalent construction for momentum space AdS, while simpler than dS and superficially more elegant, in fact breaks Lorentz invariance instead of deforming it. The model must introduce a preferred frame because the boundary of the corresponding submanifold is no longer invariant under the action of the Lorentz group. The ensuing model may thus be useful as a way of encoding subtle frame-dependence due to the boundary effects: the frame dependence is only obvious with sufficiently large Lorentz transformations.

One can also look at the bicrossproduct momenta associated to κ -Poincaré as *horospherical coordinates* on dS momentum space. Such coordinates can be introduced also for AdS momentum space and we defined the associated energy and momentum. We find that the corresponding construction introduces spatial anisotropy in momentum space and thus one must deform not only Lorentz symmetry but also spatial rotations. However, similarly to what happens in the ‘‘cosmological’’

coordinates setting, the boundary is not invariant under rotations, breaking isotropy. The result is awkward in several other ways, including the fact that the speed of light and the MDRs are anisotropic. So these first two models do not present very compelling phenomenology and, one might argue in particular, that the breaking of isotropy introduced by the second model is not particularly appealing.

A third construction, based on ‘‘static coordinates,’’ whilst not mimicking the usual set up for κ -Poincaré space, proves to be the best one conceptually and in terms of simplicity. It leads to an undeformed integration measure and a very simple Casimir invariant. It models a maximal energy and unbounded spatial momentum without introducing a preferred frame. The speed of light goes to zero in the UV limit, and this is achieved isotropically. We advocate this construction as the most conservative model for AdS momentum space. For completeness, in this paper we have also considered a dS model based on static coordinates, the counterpart to the last AdS model proposed in this paper. Curiously the dS static model breaks Lorentz invariance in a way similar to what happens to the AdS model in cosmological coordinates.

As a first explicit application of these models we investigated here the phenomenon of running of the dimensionality. We did this by considering ‘‘linearizing’’ coordinates (i.e. coordinates which render the dispersion relations trivial) and evaluating the integration measure in terms of them, to find the associated Hausdorff dimension. This procedure was considered in the past [18,20], and found to match the spectral dimension in the UV limit in all cases studied. In this paper we found that the (Lorentz breaking) AdS model based on cosmological coordinates runs to:

$$d_H = \frac{D}{1 + \gamma} \quad (76)$$

in the UV limit, whereas the (non-Lorentz breaking) model based on static coordinates runs to:

$$d_H = \frac{1 + D}{1 + \gamma} \quad (77)$$

showing further that the two models are physically distinct models. It is curious that the equivalent result for dS in static coordinates matches the result found for AdS in cosmological coordinates [cf. Eq. (75) and Eq. (65)]. This also matches the result for the κ -Poincaré space [24] if we linearize the Casimir coming directly from the embedding variables, as explained in the discussion leading to Eq. (17). Could this be pointing us to an interesting duality?

ACKNOWLEDGMENTS

We were all supported by the John Templeton Foundation. The work of M. A. was also supported by a Marie Curie Career Integration Grant within the 7th European Community Framework Programme. J. M. was also funded by a STFC consolidated grant and the Leverhulme Trust.

APPENDIX: DERIVATION OF COSMOLOGICAL COORDINATES FOR ADS

One starts by setting [29]:

$$\begin{aligned} P_0 &= \frac{1}{\ell} \sin \ell E, \\ P_i &= \hat{P}_i \cos \ell E, \\ P_4 &= \hat{P}_4 \cos \ell E, \end{aligned} \quad (\text{A1})$$

where $i = 1, 2, 3$ and P_μ are the embedding coordinates. In this way Eq. (18) becomes a condition requiring the spatial homogeneous leaves to be hyperboloids:

$$\hat{P}_i^2 - \hat{P}_4^2 = -\frac{1}{\ell^2}. \quad (\text{A2})$$

In terms of these coordinates the metric induced on the 4-surface is the cosmological rendition of (a portion of) AdS:

$$ds^2 = -dE^2 + \cos^2(\ell E) d\sigma^2 \quad (\text{A3})$$

where the spatial metric is

$$d\sigma^2 = d\hat{P}_1^2 + d\hat{P}_2^2 + d\hat{P}_3^2 - d\hat{P}_4^2 \quad (\text{A4})$$

subject to (A2). Introducing polar coordinates in the $\{\hat{P}_i\}$ space:

$$\begin{aligned} \hat{P}_1 &= p \cos \theta, \\ \hat{P}_2 &= p \sin \theta \cos \phi, \\ \hat{P}_3 &= p \sin \theta \sin \phi, \end{aligned} \quad (\text{A5})$$

ensures that p will be a comoving areal coordinate. Indeed, then $d\sigma^2 = dp^2 + p^2 d\Omega^2 - dP_4^2$, and dP_4^2 can at most

correct the dp^2 component of the metric. Specifically we can solve (A2) as:

$$\hat{P}_4 = \frac{\sqrt{1 + \ell^2 p^2}}{\ell} \quad (\text{A6})$$

and by differentiating and inserting in $d\sigma^2$ we get the cosmological form of the AdS metric:

$$ds^2 = -dE^2 + \cos^2(\ell E) \left(\frac{dp^2}{1 + \ell^2 p^2} + p^2 d\Omega^2 \right). \quad (\text{A7})$$

The explicit expression relating the two sets of coordinates is therefore:

$$\begin{aligned} P_0(E, \vec{p}) &= \frac{1}{\ell} \sin \ell E, P_1(E, \vec{p}) \\ &= p \cos(\ell E) \cos \theta, P_2(E, \vec{p}) \\ &= p \cos(\ell E) \sin \theta \cos \phi, P_3(E, \vec{p}) \\ &= p \cos(\ell E) \sin \theta \sin \phi, P_4(E, \vec{p}) \\ &= \frac{1}{\ell} \sqrt{1 + (\ell p)^2} \cos \ell E. \end{aligned} \quad (\text{A8})$$

This transformation can be abbreviated using notation:

$$\begin{aligned} P_0(E, \vec{p}) &= \frac{1}{\ell} \sin \ell E, \\ P_r(E, \vec{p}) &= p \cos \ell E, \\ P_4(E, \vec{p}) &= \frac{1}{\ell} \sqrt{1 + (\ell p)^2} \cos \ell E, \end{aligned} \quad (\text{A9})$$

where the $\{P_i\}$ are to be obtained from P_r via the usual polar coordinate formulas. Then Eq. (A9) is valid in any number D of spatial dimensions, as long as we employ the standard polar coordinate $D - 1$ angles. In these coordinates the AdS metric is

$$ds^2 = -dE^2 + \cos^2(\ell E) \left(\frac{dp^2}{1 + \ell^2 p^2} + p^2 d\Omega_{D-1}^2 \right). \quad (\text{A10})$$

-
- [1] G. Amelino-Camelia, Quantum-spacetime phenomenology, *Living Rev. Relativity* **16**, 5 (2013).
 [2] D. Mattingly, Modern tests of Lorentz invariance, *Living Rev. Relativity* **8**, 5 (2005).

- [3] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos, and S. Sarkar, Tests of quantum gravity from observations of gamma-ray bursts, *Nature (London)* **393**, 763 (1998).

- [4] U. Jacob and T. Piran, Neutrinos from gamma-ray bursts as a tool to explore quantum-gravity-induced Lorentz violation, *Nat. Phys.* **3**, 87 (2007).
- [5] S. Alexander, R. Brandenberger, and J. Magueijo, Noncommutative inflation, *Phys. Rev. D* **67**, 081301 (2003).
- [6] S. Alexander and J. Magueijo, in *Proceedings of the XIIIrd Rencontres de Blois 'Frontiers of the Universe'* (The Gioi Publishers, 2004), p. 281.
- [7] G. Gubitosi, L. Pagano, G. Amelino-Camelia, A. Melchiorri, and A. Cooray, A constraint on Planck-scale modifications to electrodynamics with CMB polarization data, *J. Cosmol. Astropart. Phys.* **08** (2009) 021.
- [8] G. Amelino-Camelia, Relativity in space-times with short distance structure governed by an observer independent (Planckian) length scale, *Int. J. Mod. Phys. D* **11**, 35 (2002).
- [9] J. Magueijo and L. Smolin, Lorentz Invariance with an Invariant Energy Scale, *Phys. Rev. Lett.* **88**, 190403 (2002).
- [10] J. Kowalski-Glikman and S. Nowak, Noncommutative space-time of doubly special relativity theories, *Int. J. Mod. Phys. D* **12**, 299 (2003).
- [11] G. Amelino-Camelia, M. Arzano, S. Bianco, and R. J. Buonocore, The DSR-deformed relativistic symmetries and the relative locality of 3D quantum gravity, *Classical Quantum Gravity* **30**, 065012 (2013).
- [12] M. Arzano, D. Latini, and M. Lotito, Group momentum space and Hopf algebra symmetries of point particles coupled to 2 + 1 gravity, *SIGMA* **10**, 079 (2014).
- [13] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman, and L. Smolin, The principle of relative locality, *Phys. Rev. D* **84**, 084010 (2011).
- [14] J. Lukierski, A. Nowicki, and H. Ruegg, New quantum Poincare algebra and κ deformed field theory, *Phys. Lett. B* **293**, 344 (1992).
- [15] J. Lukierski and H. Ruegg, Quantum kappa Poincare in any dimension, *Phys. Lett. B* **329**, 189 (1994).
- [16] A. Borowiec and A. Pachol, Unified description for κ -deformations of orthogonal groups, *Eur. Phys. J. C* **74**, 2812 (2014).
- [17] J. Lukierski, H. Ruegg, A. Nowicki, and V.N. Tolstoi, Q deformation of Poincare algebra, *Phys. Lett. B* **264**, 331 (1991).
- [18] G. Amelino-Camelia, M. Arzano, G. Gubitosi, and J. Magueijo, Planck-scale dimensional reduction without a preferred frame, *Phys. Lett. B* **736**, 317 (2014).
- [19] J. Magueijo and L. Smolin, Generalized Lorentz invariance with an invariant energy scale, *Phys. Rev. D* **67**, 044017 (2003).
- [20] G. Amelino-Camelia, M. Arzano, G. Gubitosi, and J. Magueijo, Dimensional reduction in momentum space and scale-invariant cosmological fluctuations, *Phys. Rev. D* **88**, 103524 (2013).
- [21] J. Kowalski-Glikman and S. Nowak, Quantum κ -Poincaré Algebra from de Sitter space of momenta, [arXiv:hep-th/0411154](https://arxiv.org/abs/hep-th/0411154).
- [22] S. Majid and H. Ruegg, Bicrossproduct structure of kappa Poincare group and noncommutative geometry, *Phys. Lett. B* **334**, 348 (1994).
- [23] M. Arzano, J. Kowalski-Glikman, and A. Walkus, Lorentz invariant field theory on kappa-Minkowski space, *Classical Quantum Gravity* **27**, 025012 (2010).
- [24] M. Arzano and T. Trzesniewski, Diffusion on κ -Minkowski space, *Phys. Rev. D* **89**, 124024 (2014).
- [25] J. M. Lévy-Leblond, Une nouvelle limite non-relativiste du group de Poincaré, *Ann. Inst. Henri Poincaré, A* **3**, 1 (1965).
- [26] N. J. Vilenkin and A. U. Klimyk, in *Representation of Lie Groups and Special Functions* (Kluwer, Dordrecht, 1991), Vol. 3.
- [27] H. Lu, C. N. Pope, and P. K. Townsend, Domain walls from anti-de Sitter space-time, *Phys. Lett. B* **391**, 39 (1997).
- [28] K. Land and J. Magueijo, The Axis of Evil, *Phys. Rev. Lett.* **95**, 071301 (2005).
- [29] I. Bengtsson, <http://www.fysik.su.se/ingemar/Kurs.pdf>.