

Moduli stabilization to a natural MSSM with gravitino dark matter and inflation

Hajime Otsuka*

Department of Physics, Waseda University, Tokyo 169-8555, Japan

(Received 10 April 2015; published 3 August 2015)

We propose the gravitino dark matter in the gravity mediated supersymmetry breaking scenario. The mass hierarchies between the gravitino and other superparticles can be achieved by the nontrivial Kähler metric of the supersymmetry breaking field. As a concrete model, we consider the five-dimensional supergravity model in which moduli are stabilized, and then one of the moduli induces the slow-roll inflation. It is found that the relic abundance of the gravitino and the Higgs boson mass reside in the allowed range without a severe fine-tuning.

DOI: [10.1103/PhysRevD.92.045001](https://doi.org/10.1103/PhysRevD.92.045001)

PACS numbers: 04.65.+e, 12.60.Jv, 95.35.+d, 98.80.-k

I. INTRODUCTION

The low-scale supersymmetry (SUSY) is an attractive scenario which not only protects the mass of the Higgs boson from the large radiative corrections but also gives the dark matter candidates. In addition to it, the existence of supersymmetry is also motivated in the string theory which is expected as the ultraviolet completions of the standard model (SM). This is because the SUSY guarantees the absence of tachyons in the string theory.

In the minimal supersymmetric standard model (MSSM), the large radiative corrections are indicated by the observed Higgs boson mass within ranges between 124.4 and 126.8 GeV [1]. One of the solutions to raise the Higgs boson mass in the MSSM is the high-scale SUSY-breaking scenario, and then the SUSY flavor and CP problems can be also solved at the same time. However, this scenario brings the tuning problem to the MSSM in order to realize the successful electroweak (EW) symmetry breaking. By contrast, there is another solution to raise the Higgs boson mass by the nature of left-right mixing of the top squarks. As pointed out in Ref. [2], the nonuniversal gaugino masses at the grand unification theory (GUT) scale $M_{\text{GUT}} \simeq 2 \times 10^{16}$ GeV lead to the maximal mixing of the top squarks and then, the realistic Higgs boson mass can be achieved without a severe fine-tuning by the structure of the renormalization group equations in the MSSM. Throughout this paper, we focus on this low-scale SUSY-breaking scenario.

The SUSY-breaking scenarios are mainly categorized into the gravity mediation [3], gauge mediation [4], and the anomaly mediation [5]. For any mediation mechanisms, the gravitino mass is sensitive to the cosmological problem, e.g., the cosmological gravitino problem [6]. If the gravitino is not stable, the mass of the gravitino should be larger than $\mathcal{O}(10 \text{ TeV})$ in order to be consistent with the successful big bang nucleosynthesis (BBN). The lower limit of the gravitino mass depends on the reheating temperature; for

more details see Refs. [6–10].¹ Therefore, before discussing our considered situation, we comment on several SUSY-breaking scenarios, focusing on the mass of the gravitino.

In the gauge mediated SUSY-breaking scenario, the dark matter candidate is the ultralight gravitino of mass $m_{3/2} \ll \mathcal{O}(1 \text{ GeV})$ under the low-scale SUSY breaking. Note that if the gravitino mass is larger than this scale, it is expected that the gravitational interactions give the sizable effects to the dynamics of the SUSY-breaking sector as well as the visible sector. In the pure anomaly mediated SUSY-breaking scenario, the winolike neutralino is likely to be the dark matter candidate due to the structure of the beta functions in the MSSM [12]. However, the recent results of the LHC experiments [13] indicate the TeV scale gluino mass; in other words, the large mass of the gravitino $m_{3/2} \simeq \mathcal{O}(100 \text{ TeV})$ is required in the framework of anomaly mediation. In the mirage mediation [14], the mixed neutralino would be the dark matter candidate and the large gravitino mass above $\mathcal{O}(10 \text{ TeV})$ is expected in the light of the cosmological gravitino problem. In the gravity mediation, the neutralino dark matter is often considered under the large gravitino mass above $\mathcal{O}(10 \text{ TeV})$ with high-scale SUSY breaking; otherwise, SUSY flavor violations arise due to the flavor dependent interactions.

In this paper, we consider the gravity mediated SUSY-breaking scenario which is compatible with the low-scale SUSY and observed Higgs boson mass without the cosmological gravitino and SUSY flavor problems. In general, the gravity mediation connects the scale of the gravitino mass with that of the supersymmetric particle, because the origin of soft SUSY-breaking terms is only the gravitational interactions. Therefore, it seems to be difficult to solve the cosmological gravitino problem with the low-scale SUSY-breaking scenario. In order to realize the low-scale SUSY without the cosmological gravitino problem, we propose the mechanism to generate the mass hierarchies

*hajime.13.gologo@akane.waseda.jp

¹It is also possible to consider the light gravitino in the extension of the MSSM. See, e.g., Ref. [11].

between the gravitino and the other sparticles based on the framework of a four-dimensional $\mathcal{N} = 1$ supergravity (4D $\mathcal{N} = 1$ SUGRA). Especially, we focus on the case that the gravitino is the lightest supersymmetric particle (LSP) whose mass is of $\mathcal{O}(100 \text{ GeV})$. Since such a stable gravitino is much heavier than that predicted by the gauge mediated SUSY-breaking scenario, this would be the typical feature of the gravity mediation. There are some studies for the gravitino dark matter with assumed sparticle spectra that focus on the cosmological implications and it is then found that the next-to-the-lightest supersymmetric particle (NLSP) is severely constrained. (See, e.g., Refs. [10,15–17].) In order to determine the relevant higher-dimensional operators in 4D $\mathcal{N} = 1$ SUGRA, we consider a five-dimensional supergravity (5D SUGRA) compactified on an orbifold S^1/Z_2 . In the framework of 5D SUGRA, the successful inflation mechanism as well as the moduli stabilization can be realized as suggested in Refs. [18,19]. The dynamics of inflaton and moduli are important to evaluate the abundance of the gravitino produced via the inflaton and moduli decay into the gravitino. Furthermore, the Yukawa hierarchies of elementary particles can be realized without a severe fine-tuning by employing the localized wave function of quarks, leptons, and Higgs in the fifth dimension [20].

The following sections are organized as follows. In Sec. II, we discuss how to realize the mass hierarchies between the gravitino and other supersymmetric particles in 4D $\mathcal{N} = 1$ SUGRA. As a concrete model, in Sec. III, we briefly review the structure of 5D SUGRA on S^1/Z_2 and then the gravitino can be the dark matter candidate. Thanks to the detailed moduli stabilization as well as the inflation mechanism, one can discuss the nonthermal productions of the gravitino via the moduli and inflaton decay after the inflation. After that, in Sec. IV, we evaluate the relic abundance of the gravitino and the Higgs boson mass with a severe fine-tuning. The obtained results are consistent with the cosmological observations as well as the collider experiments. Finally, Appendixes A and B denote the detailed derivations of the scalar potential around the vacuum and during the inflation, respectively.

II. THE MASS HIERARCHIES BETWEEN THE GRAVITINO AND OTHER SPARTICES

In this section, we show how to realize the mass hierarchies between the gravitino and other sparticles in the framework of 4D $\mathcal{N} = 1$ SUGRA. The scalar potential in 4D $\mathcal{N} = 1$ SUGRA is given by

$$\begin{aligned} V &= e^K (K^{I\bar{J}} D_I W D_{\bar{J}} \bar{W} - 3|W|^2) \\ &= K_{I\bar{J}} F^I F^{\bar{J}} - 3e^{K/2} |W|^2, \end{aligned} \quad (1)$$

where K and W are the Kähler and superpotential, respectively. $D_I W = W_I + K_I W$ with $W_I = \partial W / \partial Q^I$, $K_I = \partial K / \partial Q^I$ are the Kähler covariant derivatives of

the superpotential for the scalar components of the chiral superfields Q^I , $F^I = -e^{K/2} K^{I\bar{J}} D_{\bar{J}} \bar{W}$ are the F -terms of Q^I and $K^{I\bar{J}}$ are the inverse of Kähler metric $K_{I\bar{J}} = \partial_I \partial_{\bar{J}} K$. Here and hereafter, we set the Planck unit $M_{\text{Pl}} = 1$, unless we specify it. The vanishing cosmological constant $\langle V \rangle = 0$ is rewritten in the following form:

$$m_{3/2}^2 = \frac{1}{3} \langle K_{X\bar{X}} F^X F^{\bar{X}} \rangle, \quad (2)$$

where $m_{3/2} = e^{\langle K \rangle / 2} \langle W \rangle$ is the gravitino mass. It is then assumed that the SUSY is broken by the single chiral superfield X ,² whereas the soft SUSY-breaking masses of the gauginos and scalar components of the chiral superfields Q^I are given by

$$\begin{aligned} M_a &= \langle F^X \partial_X \ln(\text{Ref}_a) \rangle, \\ m_{\Phi^I}^2 &= -\langle F^X \bar{F}^{\bar{X}} \partial_X \partial_{\bar{X}} \ln Y_{\Phi^I} \rangle, \end{aligned} \quad (3)$$

where f_a , $a = U(1)_Y, SU(2)_L, SU(3)_C$ are the gauge kinetic functions of the standard model gauge groups whose vacuum expectation values (VEVs) determine the size of gauge couplings. Y_{Φ^I} are some nontrivial functions for the kinetic term of Q^I which can be severely constrained by the flavor structure of elementary particles as can be seen later. From the above equations, (2) and (3), the nontrivial Kähler metric of the SUSY-breaking field X gives rise to two nontrivial possibilities:

- (i) The gravitino dark matter:

In the case of $\langle K_{X\bar{X}} \rangle \ll 1$, the gravitino mass is smaller than the soft SUSY-breaking masses for any value of the F -term $\langle F^X \rangle$. Then it is possible to consider the gravitino dark matter in the gravity mediated SUSY-breaking scenario with TeV scale gauginos and sparticles. It is then assumed that the derivatives of the gauge kinetic function $\partial_X \text{Ref}_a$ and the kinetic term of Φ^I , $\partial_X \partial_{\bar{X}} \ln Y_{\Phi^I}$ satisfy the certain conditions in order to obtain the gravitino dark matter. This is because the renormalization group (RG) effects are significant to discuss the sparticle spectrum. Such conditions are discussed in the case of constrained MSSM (CMSSM) [21].

The stable gravitino would be consistent with the thermal history of the Universe, even if the decays of NLSP do not spoil the success of BBN [10,15–17] and at the same time, the relic abundance of the gravitino should not be larger than that reported by the Planck Collaboration [22]. In any case, the stable gravitino is favored in the light of naturalness, because the F -term of the SUSY-breaking field can be taken as a usual low-scale SUSY-breaking scenario which softens the divergences for the Higgs boson mass. Note that the small Kähler metric of the field X should be ensured in

²It is straightforward to extend our situation in multiple SUSY-breaking fields.

order not to be below that generated by the loop and/or higher derivative corrections.

(ii) Other dark matter candidates:

By contrast, in the case of $\langle K_{X\bar{X}} \rangle \gg 1$, it is expected that the gravitino is heavier than the other sparticles for any value of the F -term $\langle F^X \rangle$ with $\partial_X \text{Re} f_a$ and $\partial_X \partial_{\bar{X}} \ln Y_{\phi^I}$ of order unity. Thus, one can solve the cosmological gravitino problem with the low-scale SUSY-breaking scenario. Then, the gravitino mass can be chosen as above 10 TeV; otherwise, the BBN is threatened by the gravitino decay into the electronic and hadronic showers. Although we do not pursue this possibility, it is interesting to work in this direction.³

III. GRAVITINO DARK MATTER IN 5D SUGRA

A. 4D effective Lagrangian and matter contents

The soft SUSY-breaking terms are sensitive to the ultraviolet completion of the SM. As a concrete model, we consider the 5D SUGRA on S^1/Z_2 and the flat 5D background metric,

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu - dy^2,$$

where x^μ with $\mu = 0, 1, 2, 3$ and y denote the 4D spacetime and fifth coordinates, respectively. $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$ and the fundamental region of the orbifold is chosen as $0 \leq y \leq L$ in which $y = 0, L$ correspond to the fixed points. The S^1/Z_2 orbifold restricts all fields $f(x, y)$ to two classes of them such as Z_2 -even and -odd fields, satisfying the following Z_2 transformations: $f(x, -y) = f(x, y)$ and $f(x, -y) = -f(x, y)$, respectively. Only Z_2 -even fields have zero modes which can appear in the low-energy effective theory.

First of all, we list the relevant matter contents of 5D SUGRA. From the structure of the orbifold, 5D SUSY is broken into the 4D $\mathcal{N} = 1$ SUSY. Correspondingly, 5D vector multiplets \mathbf{V}^I and hypermultiplets Φ_α are decomposed into 4D vector multiplets V^I and three types of chiral multiplets Σ^I , Φ_α , and Φ_α^C , that is, $\mathbf{V}^I = \{V^I, \Sigma^I\}$ with $I = 1, 2, \dots, n_v$ and $\mathbf{H}_\alpha = \{\Phi_\alpha, \Phi_\alpha^C\}$ with $\alpha = 1, 2, \dots, n_H + n_C$ where n_C is the number of compensator hypermultiplets and in this paper, it is chosen as $n_C = 1$, for simplicity. In addition to the usual Z_2 -even vector multiplets V^I involving the vector multiplets in the standard model, we consider $U(1)_{I'}$ Z_2 -odd vector multiplets $\mathbf{V}^{I'}$ with $I' = 1, 2, \dots, n'_v$. The zero modes of Z_2 -even chiral multiplets Σ^I are called as the moduli chiral multiplets T^I whose linear combination⁴ plays a role of the inflaton field as pointed out in Ref. [18]. In what follows, we define the zero mode of chiral multiplets Φ_α as Q_α which involve the quark chiral multiplets $(Q_i, \mathcal{U}_i, \mathcal{D}_i)$, lepton chiral multiplets

$(\mathcal{L}_i, \mathcal{E}_i, N_i)$ with the number of generations $i = 1, 2, 3$, Higgs chiral multiplets $(\mathcal{H}_u, \mathcal{H}_d)$, SUSY-breaking chiral multiplet X and the stabilizer multiplets $H_{I'}$. These multiplets have representations of the standard model gauge groups and extra $U(1)_{I'}$ gauge groups whose gauge fields $A_M^I, A_M^{I'}$ living in vector multiplets \mathbf{V}^I and $\mathbf{V}^{I'}$, respectively. It then assigns $U(1)_{I'}$ charges $c_{I'}^{(\alpha)}$ to these hypermultiplets \mathbf{H}_α . Here it is assumed that the visible sector consists of the MSSM plus right-handed (s)neutrinos and the same number of stabilizer hypermultiplets as that of moduli multiplets in order to generate the moduli and inflaton potential as can be shown later.

Next, we show the effective action obtained from the 5D conformal supergravity action for vector and hypermultiplets which is an off-shell description of 5D SUGRA [23,24]. The structure of the Kähler potential in 5D SUGRA on S^1/Z_2 can be characterized by the cubic polynomial of vector multiplets, the so-called *norm function*, $\mathcal{N}(M) = \sum_{I,J,K=1}^{n_v} C_{I,J,K} M^I M^J M^K$ with real coefficients $C_{I,J,K}$ for $I, J, K = 1, 2, \dots, n_v$, and the $U(1)_{I'}$ charges of hypermultiplets. After the off-shell dimensional reduction discussed in Refs. [25–27] based on the 4D $\mathcal{N} = 1$ superspace [28,29],⁵ the 4D effective Lagrangian is given by

$$\begin{aligned} \mathcal{L}_{\text{eff}} = & -\frac{1}{4} \left[\int d^2\theta \sum_a f_a(X, T) \text{tr}(\mathcal{W}^a \mathcal{W}^a) + \text{H.c.} \right] \\ & + \int d^4\theta |\phi|^2 \Omega_{\text{eff}}(|Q|^2, \text{Re}T) \\ & + \left[\int d^2\theta \phi^3 W(Q, T) + \text{H.c.} \right], \end{aligned} \quad (4)$$

where ϕ is the compensator multiplet, \mathcal{W}^a is the field strength supermultiplet for massless 4D vector multiplets V^a with $a = U(1)_Y, SU(2)_L, SU(3)_C$ originating from the 5D Z_2 -even multiplets V^a , Q_α are the 4D chiral multiplets, X is the 4D chiral multiplet which induces the SUSY breaking, and T^I are the moduli chiral multiplets.

Then, the gauge kinetic functions $f_a(X, T)$ in Eq. (4) are supposed as

$$f_a(X, T) = \xi_X^a X + \sum_{I'=1}^{n'_v} \xi_{I'}^a T^{I'}, \quad (5)$$

where $\xi_{I'}^a$ and ξ_X^a are real constants determined by the real coefficients $C_{I',J,K}$ in the norm function and the gauge kinetic functions at the orbifold fixed point $y = 0$, respectively. Since the gauge kinetic functions at the orbifold fixed points depend on the dynamics of the SUSY-breaking sector, we comment on the reason why we take the above ansatz later.

³We will discuss it in the separate work.

⁴In the case $n'_v = 1$, the radion multiplet corresponds to the single modulus $T^{I'=1}$.

⁵The more general 5D action, including Z_2 -odd fields, is discussed in Refs. [30,31].

On the other hand, the effective Kähler potential in Eq. (4) is given by

$$\Omega_{\text{eff}}(|Q|^2, \text{Re}T) = \mathcal{N}^{1/3}(\text{Re}T) \left[-3 + 2 \sum_a Y(c_a \cdot T) |Q_a|^2 + \sum_{\alpha, \beta} \tilde{\Omega}_{\alpha, \beta}^{(4)}(\text{Re}T) |Q_\alpha|^2 |Q_\beta|^2 + \mathcal{O}(|Q|^6) \right], \quad (6)$$

without the Kähler potential at the orbifold fixed points $y = 0, L$, where $\mathcal{N}(\text{Re}T)$ is the norm function, $Y(z) \equiv (1 - e^{-2\text{Re}z})/2\text{Re}z$ stands for the kinetic terms of Q_α which have appeared after solving their equation of motion in the fifth direction, and $c_\alpha^{I'}$ denote the $U(1)_{I'}$ charges of Q_α . The four-point couplings $\tilde{\Omega}_{\alpha, \beta}^{(4)}$ are defined as

$$\tilde{\Omega}_{\alpha, \beta}^{(4)} \equiv -\frac{(c_\alpha \cdot \mathcal{P}a^{-1} \cdot c_\beta) \{Y((c_\alpha + c_\beta) \cdot T) - Y(c_\alpha \cdot T)Y(c_\beta \cdot T)\}}{(c_\alpha \cdot \text{Re}T)(c_\beta \cdot \text{Re}T)} + \frac{Y((c_\alpha + c_\beta) \cdot T)}{3},$$

$$\mathcal{P}^I_J(\mathcal{X}) \equiv \delta^I_J - \frac{\mathcal{X}^I \mathcal{N}_J}{3\mathcal{N}}(\mathcal{X}), \quad (7)$$

where $\mathcal{P}^I_J(\mathcal{X})$ is the operator to project the moduli multiplets out the radion multiplet. The notable feature there is that the flavor structure of matter fields is characterized by the $U(1)_{I'}$ charges of them in the Kähler potential (6). By contrast, the superpotential can be allowed only at the orbifold fixed points where the SUSY is reduced to the 4D $\mathcal{N} = 1$. Therefore, we consider the superpotential including the Yukawa couplings and μ -term in the MSSM, moduli potential at $y = 0$, and the moduli potential at $y = L$, respectively. The explicit form of the superpotential in Eq. (4) is shown later.

B. Gravitino dark matter in 5D SUGRA

In this section, we show the realization of mass hierarchies between the gravitino and other sparticles in the framework of 5D SUGRA. As shown in Eq. (6) in Sec. III A, the bulk Kähler potential is rewritten as

$$K_{\text{bulk}} = -\ln \mathcal{N}(\text{Re}T) + \sum_a Z_{Q_a}(\text{Re}T) |Q_a|^2 + Z_X(\text{Re}T) |X|^2 + \mathcal{O}(|Q|^4), \quad (8)$$

where the Kähler metric $K_{X\bar{X}}$ for the SUSY-breaking field X is given by

$$K_{X\bar{X}} = Z_X(\text{Re}T) = \frac{1 - e^{-2c_X \cdot \text{Re}T}}{c_X \cdot \text{Re}T} \simeq \begin{cases} \frac{1}{c_X \cdot \text{Re}T}, & c_X \cdot \text{Re}T > 0, \\ \frac{1}{|c_X \cdot \text{Re}T|} |\exp(2|c_X \cdot \text{Re}T|)|, & c_X \cdot \text{Re}T < 0, \end{cases} \quad (9)$$

where $K_{X\bar{X}}$ depends on the $U(1)_{I'}$ charges of the field X for the Z_2 -odd vector multiplets $\mathbf{V}^{I'}$ and the VEVs of the moduli $T^{I'}$, except for the case of the vanishing $U(1)_{I'}$ charges. For the mild large volume of the fifth dimension, $L \simeq \mathcal{N}^{1/2}(\langle \text{Re}T \rangle) \gg 1$ and positive $U(1)_{I'}$ charges, the VEV

of the Kähler metric is smaller than $\mathcal{O}(1)$, that is, $\langle K_{X\bar{X}} \rangle \ll 1$, which is important to obtain so that the light gravitino can be lower than the other sparticles.

The soft SUSY-breaking masses for the scalar components of Q_α are given by the four-point couplings $\tilde{\Omega}_{\alpha, X}^{(4)}$ in Eq. (7). For typical $U(1)_{I'}$ charges of Q_α to realize the realistic Yukawa couplings, the soft SUSY-breaking masses are larger than the gravitino mass as shown later. Furthermore, the gauge kinetic functions in Eq. (5) lead to the following gaugino masses at the compactification scale by employing the formula (3),

$$M_a = \frac{F^X}{g_a^2} \xi_X^a + \sum_{I'=1}^{n_{I'}} \frac{F^{T^{I'}}}{g_a^2} \xi_{I'}^a. \quad (10)$$

When the compactification scale is close to the GUT scale, we obtain the gaugino masses at the EW scale after solving the one-loop RG equations from the GUT scale to the EW scale,

$$\begin{aligned} M_1(M_{\text{EW}}) &\simeq 0.4M_1(M_{\text{GUT}}), \\ M_2(M_{\text{EW}}) &\simeq 0.8M_2(M_{\text{GUT}}), \\ M_3(M_{\text{EW}}) &\simeq 2.9M_3(M_{\text{GUT}}). \end{aligned} \quad (11)$$

Then, the gravitino LSP occurs if these gaugino masses at the EW scale are larger than the gravitino, as pointed out in Ref. [21]. In the case of 5D SUGRA, such situations can be realized by properly choosing the parameters ξ_X^a , $\xi_{I'}^a$ and at the same time, the Higgsino mass should be larger than the gravitino mass. Thus, one can consider the gravitino dark matter in the gravity mediated SUSY-breaking scenario without changing the VEVs of the F -terms as discussed in Sec. II. In order to estimate thermal and nonthermal abundances of gravitino via the moduli and/or inflaton decay, we focus on the specific model which realizes the successful moduli inflation as well as the moduli stabilization [18] in the next section, III C.

The mild large volume also reduces the contribution from the Kähler potential at the orbifold fixed points $y = 0, L$ to be small compared with the bulk Kähler potential (8). Since their boundary terms are described by

$$K_{\text{boundary}} = \mathcal{N}^{-1/3} (K^{(0)}(|X|^2) + K^{(L)}(e^{-c_x(\tilde{T}+\tilde{T})}|X|^2) + \dots), \quad (12)$$

the overall factor $\mathcal{N}^{-1/3}$ suppress these contributions. The one-loop corrections to the moduli Kähler potential [32] are also suppressed by the mild large volume of the fifth dimension.

By contrast, in the case of negative $U(1)_{I'}$ charges, the VEV of the Kähler metric is bigger than $\mathcal{O}(1)$, that is, $\langle K_{\tilde{X}\tilde{X}} \rangle \gg 1$. From the mass formula of the gravitino and sparticles given by Eqs. (2) and (3), one can expect that the sparticles are lighter than the gravitino without changing the F -term of the SUSY-breaking field. Thus, it is possible to solve the gravitino and fine-tuning problems at the same time.

C. Moduli stabilization

Following the discussion about the small-field inflation in Ref. [18], we choose the norm function as

$$\mathcal{N}(\text{Re}T) = (\text{Re}T^1)(\text{Re}T^2)(\text{Re}T^3), \quad (13)$$

which leads to the diagonal moduli Kähler metric. Because it seems to be difficult to obtain the realistic masses and mixings of quarks and leptons in the case of two moduli as shown in Sec. IV A, we restrict ourselves to the case of three moduli $T^{I'=1,2,3}$ in what follows. In order to generate the moduli potential, we introduce the same number of stabilizer chiral multiplets H_i as that of moduli chiral multiplets as stated in Sec. III A. The effective Kähler potential, except for the SUSY-breaking field X and other matters in the MSSM are

$$K = -\ln \mathcal{N}(\text{Re}T) + \sum_{i=1}^3 Z_{H_i} (\text{Re}T^{I'=i}) |H_i|^2, \quad (14)$$

where it is then assumed that the stabilizer fields H_i have only the $U(1)_{I'=i}$ charge with $i = 1, 2, 3$, for simplicity. In addition to it, the relevant superpotential for the moduli inflation and stabilization is

$$\begin{aligned} W_{\text{mod}} &= \sum_{i=1}^3 J_0^{(i)} H_i^{(0)} - \sum_{i=1}^3 J_L^{(i)} H_i^{(L)} \\ &= \sum_{i=1}^3 (J_0^{(i)} - J_L^{(i)} e^{-c_{I'}^{(i)} T^{I'}}) H_i^{(0)}, \end{aligned} \quad (15)$$

where $J_{0,L}^{(i)}$ are constants at the orbifold fixed points $y = 0, L$ and the exponential factor $e^{-c_{I'}^{(i)} T^{I'}}$ comes from the profile of the wave function of the stabilizer fields in the

fifth direction, $H_i^{(L)} = e^{-c_{I'}^{(i)} T^{I'}} H_i^{(0)}$. Here we assume that these tadpole terms are dominant in the superpotential and the other terms are negligible due to some symmetries or dynamics.⁶ In the following, we omit the subscripts of the stabilizer fields at the fixed point $y = 0$, that is $H_i = H_i^{(0)}$.

In fact, from the 4D scalar potential (1) given by the Kähler and superpotential (14), (15), the expectation values of the moduli $T^{I'}$ and the stabilizer fields H_i are found as [19]

$$c_{I'}^{(i)} \langle T^{I'} \rangle = \ln \frac{J_L^{(i)}}{J_0^{(i)}}, \quad \langle H_i \rangle = 0, \quad (16)$$

which are determined by the stabilization conditions, $\langle D_{I'} W \rangle = \langle D_i W \rangle = \langle W \rangle = 0$, at which the supersymmetric Minkowski minimum can be realized, $\langle V \rangle = 0$. Their supersymmetric masses of moduli and stabilizer fields are estimated as

$$m_{I'i}^2 \simeq \frac{e^{\langle K \rangle} \langle W_{I'i} \rangle^2}{\langle K_{I'I'} \rangle \langle K_{ii} \rangle}, \quad (17)$$

where $\langle W_{I'i} \rangle = -c_{I'}^{(i)} J_L^{(i)} e^{-c_{I'}^{(i)} T^{I'}}$ and $W_{ij} = \partial_i \partial_j W$. Now there are no mixing terms between the moduli and stabilizer fields in the mass matrices due to the diagonal Kähler metric of them. From the exponential behaviors of supersymmetric masses (17), the mass scales of moduli and stabilizer fields are controlled by the sizes of $U(1)_{I'=i}$ charge and constants $J_{0,L}^{(i)}$.

So far, the SUSY is not broken in the superpotential (15). For the SUSY-breaking sector, we consider the O'Raifeartaigh model [34] which is simplified as the following Kähler and superpotential of the SUSY-breaking field X after integrating out the heavy modes:

$$K = Z_X (\text{Re}T^1, \text{Re}T^2) |X|^2 - \frac{1}{\Lambda^2} |X|^4, \quad W = w + \nu X, \quad (18)$$

where w, ν are the real parameters and the SUSY-breaking field X has no $U(1)_3$ charge, for simplicity. The Kähler potential receives the loop corrections from the heavy modes, whose mass scale is Λ [35].

In general, the true vacuum of the moduli and stabilizer fields are deviated from the supersymmetric one due to the SUSY-breaking effects and then the moduli and stabilizer fields obtain their F -terms at the true vacuum. Since their F -terms would change the cosmological history of the Universe through the moduli decay into the gravitinos, it is important to evaluate their F -terms at the true vacuum. For that reason, we adopt the perturbation method, known as the reference point method [36] to search for the true vacua of all the fields.

⁶A similar moduli stabilization was proposed in Ref. [33] in the case of $n_c = 2$.

First, as the reference points for the moduli and stabilizer fields, we take them as given in Eq. (16) satisfying as

$$D_{H_i} W|_{\text{ref}} = W_{H_i} + K_{H_i} W = 0, \quad D_{T^{i'}} W|_{\text{ref}} = K_{T^{i'}} w, \quad (19)$$

and for the SUSY-breaking field X , its reference point is taken as that satisfying the following stabilization condition:

$$\begin{aligned} e^{-K} V_X|_{\text{ref}} &= \partial_X \left(\sum_{I'} K^{T^{i'} \bar{T}^{i'}} |D_{T^{i'}} W|^2 + K^{X\bar{X}} |D_X W|^2 - 3|W|^2 \right) \\ &\simeq 3W_X \bar{W} + \partial_X (K^{X\bar{X}}) |W_X|^2 + K^{X\bar{X}} W_X K_{X\bar{X}} \bar{W} - 3W_X \bar{W} \\ &\simeq \frac{4|W_X|^2}{\Lambda^2 (Z_X)^2} \bar{X} + W_X \bar{W} = 0, \end{aligned} \quad (20)$$

in the limit $w \ll 1$, where $V_X = \partial_X V$. Thus, we obtain

$$X|_{\text{ref}} = -\frac{\Lambda^2 (Z_X)^2}{4} \left(\frac{W}{W_X} \right) \simeq -\frac{\Lambda^2 (Z_X)^2 w}{4\nu}. \quad (21)$$

Next, we expand these fields as $\phi \rightarrow \phi|_{\text{ref}} + \delta\phi$, $\phi = T^{i'}$, H_i , X with $I', i = 1, 2, 3$ and evaluate their perturbations from the reference points given by Eqs. (16) and (21) under the following conditions:

$$\begin{aligned} V &= V|_{\text{ref}} + V_I|_{\text{ref}} \delta\phi^I + V_{\bar{I}}|_{\text{ref}} \delta\bar{\phi}^{\bar{I}} + V_{IJ}|_{\text{ref}} \delta\phi^I \delta\phi^J + V_{I\bar{J}}|_{\text{ref}} \delta\phi^I \delta\bar{\phi}^{\bar{J}} + V_{\bar{I}J}|_{\text{ref}} \delta\bar{\phi}^{\bar{I}} \delta\phi^J + \mathcal{O}(\delta\phi^3), \\ |V_I|_{\text{ref}} \delta\phi^I + V_{\bar{I}}|_{\text{ref}} \delta\bar{\phi}^{\bar{I}} &\gg |V_{IJ}|_{\text{ref}} \delta\phi^I \delta\phi^J + V_{I\bar{J}}|_{\text{ref}} \delta\phi^I \delta\bar{\phi}^{\bar{J}} + V_{\bar{I}J}|_{\text{ref}} \delta\bar{\phi}^{\bar{I}} \delta\phi^J, \end{aligned} \quad (22)$$

where $V_I = \partial_I V$ and $V_{IJ} = \partial_I \partial_J V$ are the derivatives for the relevant fields ϕ , and then $\phi|_{\text{ref}} + \delta\phi$ are considered as the vacua of relevant fields. Note that these perturbations are valid even if the SUSY-breaking scale is smaller than the scale of supersymmetric masses given by Eq. (17). As a result, the deviations of the fields from the reference points (16), (21) are

$$\delta H_i \simeq \frac{w}{2\text{Re}T^{i'} W_{T^{i'} H_i}}, \quad \delta T^{i'} \simeq \left(\frac{w}{W_{T^{i'} H_i}} \right)^2, \quad \delta X \simeq \left(\frac{\Lambda^2 Z_X^2}{4w^2} \right) 5w W_X, \quad (23)$$

and the F -terms and squared masses of moduli, stabilizer, and SUSY-breaking fields are roughly estimated as

$$\begin{aligned} \sqrt{K_{T^{i'} \bar{T}^{i'}}} F^{T^{i'}} &\simeq \mathcal{O}\left(\frac{w^3}{m_{T^{i'}}^2} \right), & \sqrt{K_{H_i \bar{H}_i}} F^{H_i} &\simeq \mathcal{O}\left(\frac{w^3}{m_{H_i}^2} \right), & \sqrt{K_{X\bar{X}}} F^X &\simeq \frac{-\nu}{\mathcal{N}^{1/2} Z_X^{1/2}} \\ m_{T^{i'}}^2 &\simeq m_{H_i}^2 \simeq \frac{e^K W_{T^{i'} H_i}^2}{K_{T^{i'} \bar{T}^{i'}} K_{H_i \bar{H}_i}} \quad (I' = i), & m_X^2 &\simeq \frac{e^K}{K_{X\bar{X}}} \frac{4w^2}{\Lambda^2 Z_X^2}, \end{aligned} \quad (24)$$

at the vacuum, $\phi = \phi|_{\text{ref}} + \delta\phi$. The mass squares of real and imaginary parts of moduli, stabilizer, and SUSY-breaking fields are the same as each other and here and in what follows, they are denoted as $m_{T^{i'}}^2$, $m_{H_i}^2$, and m_X^2 , respectively. The details of these derivations are summarized in Appendix A. The mass differences between $m_{T^{i'=i}}$ and m_{H_i} are of the order of the gravitino mass. It is remarkable that the fields, except for the SUSY-breaking field X , have almost vanishing F -terms due to their large supersymmetric masses.

D. Moduli inflation

In this section, we briefly review the inflation mechanism in which the inflaton is identified as one

of the real parts of the moduli. Although in Ref. [18], both the small- and large-field inflation are discussed in the light of recent Planck results, in this paper, we restrict ourselves to the small-field inflation, for simplicity.⁷ The inflaton potential is generated by the Kähler and superpotential of the pair (T^3, H_3) in Eqs. (14) and (15),

$$\begin{aligned} K &= -\ln \mathcal{N} + Z_{H_3} (\text{Re}T^3) |H_3|^2, \\ W_{\text{inf}} &= (J_0^{(3)} - J_L^{(3)} e^{-c_3^{(3)} T^3}) H_3, \end{aligned} \quad (25)$$

where

⁷The extension to the large-field inflation is straightforward.

$$Z_{H_3}(\text{Re}T^3) = \frac{1 - e^{-2c_3^{(3)}\text{Re}T^3}}{c_3^{(3)}\text{Re}T^3}, \quad (26)$$

and the effective scalar potential is obtained from Eq. (1) with the above Kähler and superpotential (25),

$$V_{\text{inf}} = e^K K^{H_3\bar{H}_3} |W_{H_3}|^2 \simeq \frac{|J_0^{(3)} - J_L^{(3)} e^{-c_3^{(3)}T^3}|^2}{\langle \text{Re}T^1 \rangle \langle \text{Re}T^2 \rangle (1 - e^{-c_3^{(3)}T^3})}, \quad (27)$$

where $\text{Re}T^3$ is identified as the inflaton. Here, it is supposed that the other moduli T^I , stabilizer fields H_i with $I, i = 1, 2$ are heavier than the pair (T^3, H_3) and fixed at their minima. This is because the minima of them are fixed by their own superpotential in Eq. (16), and they can be decoupled from the pair (T^3, H_3) by choosing the parameters in the superpotential (15),

$$J_0^{(1)} = J_0^{(2)} = \frac{1}{9}, \quad J_L^{(1)} = J_L^{(2)} = 1, \quad (28)$$

and the nonvanishing $U(1)_{1,2,3}$ charges of $H_{1,2}$,

$$c_1^{(1)} = c_2^{(2)} = \frac{1}{50}, \quad (29)$$

whereas the constants $J_{0,L}^{(3)}$ are chosen to be small compared with $J_{0,L}^{(1),(2)}$ as shown later. Furthermore, in the following analysis, we omit the fluctuation of H_3 and X , because their minima are fixed around the origin by the Hubble-induced mass during the inflation. $\text{Im}T^3$ is also fixed at the origin during and after the inflation. They can be checked that the fluctuations of these fields are negligible to the inflaton dynamics as explicitly shown in Appendix B.

When $\text{Re}T^3$ is identified as the inflaton, the effective scalar potential (27) is similar to the one in the Starobinski model [37] and is drawn in Fig. 1 with the parameters given by Eqs. (28), (29), and (36). From Fig. 1, the inflaton, $\text{Re}T^3$ can roll its potential slowly down to its minimum from the large value of $\text{Re}T^3$. In order to evaluate the cosmological observables for the cosmic microwave background (CMB) observed by Planck, we define the slow-roll parameters for the inflaton, $\sigma \equiv \text{Re}T^3$,

$$\epsilon \equiv \frac{M_{\text{Pl}}^2}{2} \frac{\nabla_\sigma V_{\text{inf}} K^{\sigma\sigma} \nabla_\sigma V_{\text{inf}}}{V_{\text{inf}}^2}, \quad \eta \equiv \frac{\nabla^\sigma \nabla_\sigma V_{\text{inf}}}{V_{\text{inf}}}, \quad (30)$$

where ∇_σ is the Kähler covariant derivative for the field σ . With these slow-roll approximations, the power spectrum




FIG. 1 (color online). Inflaton potential V_{inf} on the $\text{Im}T^1 = 0$ hypersurface.

of the scalar curvature perturbation, its spectral index, and tensor-to-scalar ratio can be expressed as

$$P_\xi(k) = \frac{1}{24\pi^2} \frac{V_{\text{inf}}}{\epsilon}, \quad n_s = 1 + \frac{d \ln P_\xi(k)}{d \ln k} \simeq 1 - 6\epsilon + 2\eta, \quad r = 16\epsilon. \quad (31)$$

The recent data reported by the Planck Collaboration shows the almost scale invariant spectrum and the upper limit of r [22],

$$P_\xi(k) \simeq 2.20 \pm 0.10 \times 10^{-9}, \quad n_s = 0.9655 \pm 0.0062, \quad r < 0.11, \quad (32)$$

at the scale $k_* = 0.05$ [Mpc^{-1}]. The inflaton dynamics is obeyed by its equation of motion,

$$\sigma'' = - \left(1 - \frac{g_{\sigma\sigma}(\sigma')^2}{6} \right) \left(3\sigma' + 6\sigma^2 \frac{V'}{V} \right) + \frac{(\sigma')^2}{\sigma'}, \quad (33)$$

where $'$ denotes the d/dN by employing the number of e -foldings N rather than time;

$$a(t) = e^N, \quad \frac{d}{dt} = \frac{dN}{dt} \frac{d}{dN} = H \frac{d}{dN}, \quad (34)$$

where $a(t)$ is the scale factor of 4D spacetime. The metric $g_{\sigma\sigma}$ is connected to the Kähler metric $K_{T^3\bar{T}^3}$ such that $\frac{1}{2}g_{\sigma\sigma}\partial\sigma\partial\sigma = K_{T^3\bar{T}^3}\partial T^3\partial\bar{T}^3$ and $\Gamma_{\sigma\sigma}^\sigma = -1/\sigma$ is the Christoffel symbol. As a result, it is found that the power spectrum of scalar curvature perturbation, its spectral index, and tensor-to-scalar ratio are consistent with the current cosmological data,

$$P_\xi(k) \simeq 2.2 \times 10^{-9}, \quad n_s \simeq 0.96, \quad r \simeq 10^{-5}, \quad (35)$$

with enough e -foldings $N \simeq 58$ and then the parameters are chosen as

$$J_0^{(3)} = \frac{1}{4000}, \quad J_L^{(3)} = \frac{3}{4000}, \quad c_3^{(3)} = \frac{1}{10}. \quad (36)$$

The running of the scalar spectral index is negligible, relative to the current observational sensitivity.

When the numerical values of the parameters are chosen as those in Eqs. (28), (29) and (36), the moduli VEVs become

$$(\text{Re}\langle T^1 \rangle, \text{Re}\langle T^2 \rangle, \text{Re}\langle T^3 \rangle) \simeq (110, 110, 11), \quad (37)$$

in the unit $M_{\text{Pl}} = 1$. According to these moduli VEVs, the typical Kaluza-Klein mass scale is found as

$$M_c = \frac{\pi}{L} \simeq \frac{\pi}{\langle \mathcal{N}^{1/2} \rangle} \simeq 2.1 \times 10^{16} \text{ GeV}, \quad (38)$$

which is close to the GUT scale due to the mild large volume of the fifth dimension, $\langle \mathcal{N}^{1/2} \rangle \simeq 364$. The mass of moduli T^I and stabilizer fields H_i are also given by

$$\begin{aligned} m_{T^1} &\simeq m_{T^2} \simeq m_{H_1} \simeq m_{H_2} \simeq 4.8 \times 10^{15} \text{ GeV}, \\ m_{T^3} &\simeq m_{H_3} \simeq 4 \times 10^{12} \text{ GeV}, \end{aligned} \quad (39)$$

and their F -terms become

$$\begin{aligned} F^{T^1} &\simeq F^{T^2} \simeq F^{H_1} \simeq F^{H_2} \simeq 1 \times 10^{-42}, \\ F^{T^3} &\simeq F^{H_3} \simeq 1.6 \times 10^{-36}, \end{aligned} \quad (40)$$

in the unit $M_{\text{Pl}} = 1$.

So far, we have specified the parameters relevant for the moduli and stabilizer fields. The parameters in the Kähler and superpotential (18) for the SUSY-breaking sector are considered as

$$\begin{aligned} \nu &\simeq -1.567 \dots \times 10^{-14}, \quad w = -6 \times 10^{-14}, \\ \Lambda &= 10^{-4}, \quad c_X^{(1)} = \frac{3}{10}, \quad c_X^{(2)} = \frac{1}{10}, \quad c_X^{(3)} = 0, \end{aligned} \quad (41)$$

where ν is properly chosen as realizing the Minkowski minimum. Then, the mass of the gravitino and the mass and F -term of X are obtained as

$$\begin{aligned} m_{3/2} &\simeq 395 \text{ GeV}, \quad m_X \simeq 6 \times 10^8 \text{ GeV}, \\ \frac{F^X}{M_{\text{Pl}}} &\simeq 4541 \text{ GeV}, \end{aligned} \quad (42)$$

which implies that the gravitino mass is suppressed by the Kähler metric of the SUSY-breaking field, $K_{X\bar{X}} \simeq 1/(c_X \cdot \text{Re}T) \simeq 0.023$ as discussed in Sec. III B and the concrete sparticle spectra are shown in Sec. IV C.

E. Moduli-induced gravitino problem and reheating temperature

As mentioned in Sec. III C, the moduli and stabilizer fields are so heavy that they decay into the particles in the MSSM before the BBN. However, even if they are much heavier than $\mathcal{O}(100 \text{ TeV})$, it has to be taken into account of the cosmological problem, e.g., the moduli-induced gravitino problem [38,39].

The moduli decay width into the gravitino pair can be evaluated by the couplings between moduli and gravitinos in the unitary gauge,

$$\begin{aligned} \mathcal{L}_{3/2} &= e^{\mu\nu\rho\sigma} \bar{\psi}_\mu \bar{\sigma}_\nu \tilde{D}_\rho \psi_\sigma - e^{K/2} W^* \psi_\mu \sigma^{\mu\nu} \psi_\nu \\ &\quad - e^{K/2} W \bar{\psi}_\mu \bar{\sigma}^{\mu\nu} \bar{\psi}_\nu, \end{aligned} \quad (43)$$

where ψ_μ denotes the gravitino in two-component formalism and the relevant covariant derivatives of the gravitino are $\tilde{D}_\rho \psi_\sigma = \partial_\rho \psi_\sigma + \frac{1}{4}(K_J \partial_\rho \phi^J - K_{\bar{J}} \partial_\rho \bar{\phi}^{\bar{J}})$. After carrying out a field-dependent chiral transformation,

$$\psi_\mu \rightarrow \left(\frac{W}{\bar{W}} \right)^{-1/4} \psi_\mu, \quad (44)$$

the Lagrangian (43) is simplified as

$$\mathcal{L}_{3/2} = e^{\mu\nu\rho\sigma} \bar{\psi}_\mu \bar{\sigma}_\nu \partial_\rho \psi_\sigma + \frac{e^{\mu\nu\rho\sigma}}{4} (G_J \partial_\rho \phi^J - G_{\bar{J}} \partial_\rho \bar{\phi}^{\bar{J}}) \bar{\psi}_\mu \bar{\sigma}_\nu \psi_\sigma - e^{G/2} (\psi_\mu \sigma^{\mu\nu} \psi_\nu + \bar{\psi}_\mu \bar{\sigma}^{\mu\nu} \bar{\psi}_\nu), \quad (45)$$

where $G = K + \ln |W|^2$ and $G_J = \partial_J G$. When we expand the moduli T^I around the vacuum given by employing the reference point method (16) and (23), the Lagrangian (43) reduces to

$$\begin{aligned} \mathcal{L}_{3/2} &= -\frac{e^{\mu\nu\rho\sigma}}{2} \bar{\Psi}_\mu \gamma_5 \gamma_\nu \partial_\rho \Psi_\sigma + \frac{e^{\mu\nu\rho\sigma}}{8} (\langle G_{T^{J'}} \rangle \partial_\rho \delta T^{J'} - \langle G_{\bar{T}^{\bar{J}'}} \rangle \partial_\rho \delta \bar{T}^{\bar{J}'}) \bar{\Psi}_\mu \gamma_\nu \Psi_\sigma \\ &\quad - \frac{1}{4} \langle e^{G/2} \rangle \bar{\Psi}_\mu [\gamma^\mu, \gamma^\nu] \Psi_\nu - \frac{1}{8} \langle e^{G/2} \rangle (\langle G_{T^{J'}} \rangle \delta T^{J'} + \langle G_{\bar{T}^{\bar{J}'}} \rangle \delta \bar{T}^{\bar{J}'}) \bar{\Psi}_\mu [\gamma^\mu, \gamma^\nu] \Psi_\nu, \end{aligned} \quad (46)$$

in the four-component formalism of the gravitino Ψ_μ . As shown in Eq. (39), the moduli and stabilizer fields, except for the pair (T^3, H_3) , are decoupled from the inflaton dynamics due to their heavy masses. Therefore, their decays can be neglected and do not give the sizable effects in the thermal history of the Universe. In this respect, we focus on the decay processes of T^3 , H_3 , and SUSY-breaking field X .

1. The inflaton decay

First, we concentrate on the inflaton decay into the gravitino pair. Since the gravitino wave function is described in terms of helicity $\pm 1/2$ components of the gravitino at a high-energy limit by the equivalence theorem, the inflaton decay width into the gravitino pair is estimated as

$$\begin{aligned} \Gamma(\sigma^3 \rightarrow \Psi_{3/2} \bar{\Psi}_{3/2}) &\simeq \frac{1}{288\pi \langle K_{T^3 \bar{T}^3} \rangle} \left| \left\langle \frac{D_{T^3} W}{W} \right\rangle \right|^2 \frac{m_{T^3}^5}{m_{3/2}^2 M_{\text{Pl}}^2} \\ &\simeq \frac{1}{288\pi \langle K_{T^3 \bar{T}^3} \mathcal{N} \rangle} \frac{m_{T^3} m_{3/2}^2}{M_{\text{Pl}}^2} \\ &\simeq 1.6 \times 10^{-18} \text{ GeV}, \end{aligned} \quad (47)$$

by employing the F -term of modulus (24) and numerical values of the mass, F -term of inflaton, and gravitino mass given by Eqs. (39) and (40). Here, the reduced Planck mass has been explicitly written. When the inflaton has the sizable F -term at the vacuum, the enhancement factor $m_{3/2}^{-2}$, as the longitudinal mode of the gravitino, induces the significant amount of gravitinos which would threaten to destroy the success of BBN. However, in our moduli inflation, this direct decay is so suppressed due to the almost vanishing F -term of the inflaton. Therefore, the dominant decay process of inflaton comes from the interactions with the gauge bosons,

$$\begin{aligned} \mathcal{L}_{Tgg} &= -\frac{1}{4(g_a)^2} F_{\mu\nu}^a F^{a\mu\nu} - \frac{1}{4} \xi_{J'}^a \delta T_R^{J'} F_{\mu\nu}^a F^{a\mu\nu} \\ &\quad - \frac{1}{8} \xi_{J'}^a \delta T_L^{J'} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^a F_{\rho\sigma}^a, \end{aligned} \quad (48)$$

where $T_R^{J'} = \text{Re}T^{J'}$, $T_L^{J'} = \text{Im}T^{J'}$. Now, the gauge kinetic functions $f_a(X, T)$ are considered as in Eq. (5). In general, $\xi_X^a X$ in the gauge kinetic function (5) could appear, because the R -symmetry is explicitly broken by the constant superpotential w in Eq. (18).

The inflaton decay width into the gauge bosons are

$$\begin{aligned} \sum_{a=1}^3 \Gamma(\sigma^3 \rightarrow g^{(a)} g^{(a)}) &\simeq \sum_{a=1}^3 \frac{N_G^a}{128\pi} \left\langle \frac{\xi_3^a}{\sqrt{2K_{T^3 \bar{T}^3}}} \right\rangle^2 \frac{m_{T^3}^3}{M_{\text{Pl}}^2} \\ &\simeq 3.95 \text{ GeV}, \end{aligned} \quad (49)$$

with the numerical values of mass and VEV of modulus (37), (39), where N_G^a are the number of the gauge bosons for the gauge groups $U(1)_Y, SU(2)_L, SU(3)_C$ and the nonvanishing coefficients in the gauge kinetic function are chosen as $\xi_3^1 = \xi_3^2 = \xi_3^3 = 0.22$ to realize the gauge coupling unification at the GUT scale $M_{\text{GUT}} \simeq 2 \times 10^{16}$ GeV. Although there are the other decay processes via the inflaton decay into the gauginos λ^a given by the interactions,

$$\begin{aligned} \mathcal{L}_{T\lambda\lambda} &= -\frac{i}{2} \sum_a \text{Re} f_a (\lambda^a \sigma^\mu D_\mu \bar{\lambda}^a + (\text{H.c.})) \\ &\quad + \frac{i}{2} \sum_a \text{Im} f_a D_\mu (\lambda^a \sigma^\mu \bar{\lambda}^a) \\ &\quad + \sum_a \left(\frac{1}{4} \frac{\partial f_a}{\partial T^{J'}} F^{T'J'} \lambda^a \lambda^a + (\text{H.c.}) \right), \end{aligned} \quad (50)$$

where D_μ is the covariant derivative for gaugino, such decay channels are suppressed by the small masses of gauginos and almost vanishing F -term of inflaton such as

$$\begin{aligned} \sum_{a=1}^3 \Gamma(\sigma^3 \rightarrow \tilde{\lambda}^a \tilde{\lambda}^a) &\simeq \sum_{a=1}^3 \frac{m_{T^3} (\xi_3^a)^2 m_{\lambda^a}^2}{16\pi M_{\text{Pl}}^2} \\ &\simeq 1.5 \times 10^{-21} \text{ GeV}, \end{aligned} \quad (51)$$

with $m_{\lambda^3} \simeq 1.5$ TeV and the derivative of F -term for the inflaton,

$$\begin{aligned} \left\langle \frac{\partial F^{T^3}}{\partial T^3} \right\rangle &= \left\langle \frac{\partial}{\partial T^3} e^{K/2} (K^{T^3 \bar{T}^3} |D_{T^3} W|^2 \right. \\ &\quad \left. + K^{T^3 \bar{H}^3} D_{T^3} W D_{\bar{H}^3} \bar{W}) \right\rangle \sim \mathcal{O} \left(\frac{m_{3/2}^4}{m_{T^3}^2} \right). \end{aligned} \quad (52)$$

The decays from the inflaton into sfermions are also suppressed because of the factor, $m_{\text{sfermion}}/m_{T^3}$, if the masses of sfermions are of $\mathcal{O}(1 \text{ TeV})$. Other decays from the inflaton into the fermion pairs and quark-quark-gluon are negligible due to their small masses and phase factors, respectively, as pointed out in Ref. [39]. The μ -term does not give the sizable effects for the inflaton decay process, because we consider the tiny μ -term (~ 500 GeV) in the light of naturalness as shown in Sec. IV A. Finally, we comment on a single gravitino production via the inflaton decay into the modulino and gravitino. Since the mixing terms between T^3 and H_3 in the mass squared matrices are controlled by the SUSY-breaking scale, i.e., the gravitino mass, the mass difference between the inflaton and modulino as its superpartner is of the order of the gravitino mass. Therefore, the inflaton decay width into the modulino $\tilde{\sigma}^3$ and gravitino is suppressed by the phase factor $m_{3/2}/m_{T^3}$,

$$\begin{aligned} \Gamma(\sigma^3 \rightarrow \tilde{\sigma}^3 \Psi_{3/2}) &\simeq \frac{1}{48\pi} \left(\frac{m_{T^3}}{M_{\text{Pl}}} \right)^2 \left(\frac{m_{3/2}}{m_{T^3}} \right) m_{3/2} \\ &\simeq 7.2 \times 10^{-22} \text{ GeV}, \end{aligned} \quad (53)$$

with $m_{3/2} = 395$ GeV, $m_{T^3} = 4 \times 10^{12}$ GeV given by Eqs. (39) and (42). The inflaton decay into the SUSY-breaking field X is also suppressed, because there is no tree-level interaction between X and T^3 due to the vanishing $U(1)_3$ charge of X . As a result, the branching ratios of the moduli decaying into the gravitino(s) are summarized as

$$\begin{aligned}\Gamma_{\text{all}}^{\sigma^3} &\equiv \Gamma(\sigma^3 \rightarrow \text{all}) \simeq \sum_{a=1}^3 \Gamma(\sigma^3 \rightarrow g^{(a)} g^{(a)}) \simeq 3.95 \text{ GeV}, \\ \text{Br}(\sigma^3 \rightarrow \Psi_{3/2} \Psi_{3/2}) &\simeq \frac{\Gamma(\sigma^3 \rightarrow \Psi_{3/2} \Psi_{3/2})}{\Gamma_{\text{all}}^{\sigma^3}} \simeq 1.4 \times 10^{-20}, \\ \text{Br}(\sigma^3 \rightarrow \tilde{\sigma}^3 \Psi_{3/2}) &\simeq \frac{\Gamma(\sigma^3 \rightarrow \tilde{\sigma}^3 \Psi_{3/2})}{\Gamma_{\text{all}}^{\sigma^3}} \simeq 1.8 \times 10^{-22},\end{aligned}\quad (54)$$

and then the reheating temperature is roughly estimated by equating the expansion rate of the Universe to the total decay width of inflaton,

$$\begin{aligned}\Gamma_{\text{all}}^{\sigma^3} = H_R \Leftrightarrow T_R &= \left(\frac{\pi^2 g_*}{90} \right)^{-1/4} \sqrt{\Gamma_{\text{all}} M_{\text{Pl}}} \\ &\simeq 1.38 \times 10^9 \text{ GeV},\end{aligned}\quad (55)$$

where $H_R = H(T_R)$ and $g_*(T_R) = 915/4$ is the effective degrees of freedom of the radiation in the MSSM at the reheating. The gravitino yield $Y_{3/2}$ via the inflaton decay is suppressed due to the tiny branching ratio of the inflaton decay into the gravitino(s),

$$Y_{3/2} = \frac{n_{3/2}}{s} \simeq \text{Br}(\sigma^3 \rightarrow \tilde{\sigma}^3 \Psi_{3/2}) \frac{3T_R}{4m_{T^3}} \simeq 3.8 \times 10^{-24}, \quad (56)$$

with $m_{3/2} = 395 \text{ GeV}$, $T_R = 1.38 \times 10^9 \text{ GeV}$, $s = 4\rho/3T$, where $n_{3/2}$, s , and ρ are the number density of the gravitino, entropy, and energy density of the Universe, respectively. Now it is supposed that the coherent oscillation of the inflaton field dominates the energy density of the Universe after the inflation and there is no entropy production after the inflation as shown later.

It is remarkable that the supersymmetric moduli stabilization is important to suppress the direct decays from the inflaton into the gravitino(s) which give the solution to the cosmological moduli problem, especially the moduli-induced gravitino problem. The other gravitino production from the stabilizer fields, the SUSY-breaking field, and the thermal bath can be estimated in the next section.

2. The decay of stabilizer and SUSY-breaking fields

The stabilizer field H_3 is stabilized at the origin during the inflation and after that, $\text{Re } H_3$ oscillates around its vacuum (23) deviated from the supersymmetric one (16). On the other hand, $\text{Im } H_3$ and $\text{Im } X$ evolve to the origin during inflation and do not oscillate after the inflation as shown in Appendixes A and B. Similarly, the SUSY-breaking field $\text{Re } X$ oscillates around its vacuum after the inflation. From the analyses in Appendixes A and B, the amplitudes of both fields are found as

$$\Delta h_3 \simeq \frac{m_{3/2}}{m_{H_3}}, \quad \Delta x \simeq \left(\frac{m_{3/2}}{m_X} \right)^2, \quad (57)$$

with $h_3 = \text{Re} H_3$ and $x = \text{Re} X$. By comparing their masses given in Eq. (39) with the reheating temperature (55), the coherent oscillations of both fields h_3 and x start before the reheating process. When H_3 does not couple to the fields in the MSSM, the dominant decay process is

$$\begin{aligned}\Gamma_{\text{all}}^{h_3} &\equiv \Gamma(h_3 \rightarrow \Psi_{3/2} \Psi_{3/2}) \\ &\simeq \frac{1}{288\pi K_{H_3 \tilde{H}_3}} \left| \left\langle \frac{D_{H_3} W}{W} \right\rangle \right|^2 \frac{m_{H_3}^5}{m_{3/2}^2 M_{\text{Pl}}^2} \\ &\simeq \frac{1}{288\pi \langle K_{H_3 \tilde{H}_3} \mathcal{N} \rangle} \frac{m_{H_3}^3}{M_{\text{Pl}}^2} \\ &\simeq 0.02 \text{ GeV},\end{aligned}\quad (58)$$

which implies the decay time of h_3 is smaller than the time of the coherent oscillation of h_3 and reheating, that is, $H_{\text{osc}}^{h_3} > H(T_R) > H_{\text{dec}}^{h_3}$, with $H_{\text{osc}}^{h_3} \simeq m_{h_3}$ and $H_{\text{dec}}^{h_3} \simeq \Gamma(h_3 \rightarrow \Psi_{3/2} \Psi_{3/2})$. Here and in what follows, H_R , H_{osc}^Φ , and H_{dec}^Φ refer to the Hubble parameters at the time of reheating, beginning of oscillation of relevant fields Φ , and decay of Φ . The scale factors of 4D spacetime a_R , a_{osc}^Φ , and a_{dec}^Φ are also defined in the same way as the Hubble parameters, H_R , H_{osc}^Φ , and H_{dec}^Φ . The energy density of coherent oscillation h_3 is

$$\rho_{h_3} \simeq \frac{1}{2} m_{H_3}^2 (\Delta h_3)^2 \simeq \frac{1}{2} m_{3/2}^2 \left(\frac{a}{a_{\text{osc}}^{h_3}} \right)^{-3}, \quad (59)$$

where $a_{\text{osc}}^{h_3}$ stands for the scale factor at the time when h_3 begins to oscillate and ρ_{h_3} is converted into the gravitino yield hereafter,

$$Y_{3/2}^{h_3} = \frac{2\rho_{h_3}}{m_{H_3} s} \simeq \frac{1}{4} \frac{m_{3/2}^2 T_R}{m_{H_3}^3} \simeq 8.2 \times 10^{-25}, \quad (60)$$

with $m_{3/2} = 395 \text{ GeV}$, $T_R = 1.38 \times 10^9 \text{ GeV}$, and $m_{H_3} = 4 \times 10^{12} \text{ GeV}$. Here we employed that the entropy production from h_3 can be neglected. In our model, the following inequality is satisfied due to the tiny mass of the gravitino and then h_3 does not dominate the Universe and release the significant entropy,

$$1 \gg \frac{\rho_{h_3}}{\rho_R} \Big|_{T=T_{\text{dec}}^{h_3}} = \frac{\rho_{h_3}}{\rho} \Big|_{\text{end}} \left(\frac{T_R}{T_{\text{dec}}^{h_3}} \right) \simeq \frac{m_{3/2}^2 M_{\text{Pl}}^2}{2V_{\text{inf}}} \left(\frac{T_R}{T_{\text{dec}}^{h_3}} \right), \quad (61)$$

where ρ_{h_3} , ρ_R are the energy densities of h_3 and radiation, respectively. $\rho|_{\text{end}} = V_{\text{inf}} \simeq \mathcal{O}(10^{-13})$ denotes the energy density at the end of inflation and $T_{\text{dec}}^{h_3}$ is the decay temperature of h_3 ,

$$T_{\text{dec}}^{h_3} = \left(\frac{\pi^2 g_*}{90}\right)^{-1/4} \sqrt{\Gamma_{\text{all}}^{h_3} M_{\text{Pl}}} \approx 8.6 \times 10^7 \text{ GeV}. \quad (62)$$

Furthermore, the SUSY-breaking field also produces the gravitinos through the following dominant decay channel:

$$\begin{aligned} \Gamma(x \rightarrow \Psi_{3/2} \Psi_{3/2}) &\simeq \frac{1}{288\pi \langle K_{X\bar{X}} \rangle} \left| \left\langle \frac{D_X W}{W} \right\rangle \right|^2 \frac{m_X^5}{m_{3/2}^2 M_{\text{Pl}}^2} \\ &\simeq \frac{1}{288\pi \langle K_{X\bar{X}} \rangle} \left| \frac{\nu}{w} \right|^2 \frac{m_X^5}{m_{3/2}^2 M_{\text{Pl}}^2}. \end{aligned} \quad (63)$$

With the parameters (41), the VEVs of moduli (37), and mass of $\Psi_{3/2}$ and X (42), the total decay width of X then becomes

$$\Gamma_{\text{all}}^x \equiv \Gamma(x \rightarrow \Psi_{3/2} \Psi_{3/2}) \approx 3.7 \times 10^8 \text{ GeV}. \quad (64)$$

Therefore, the decay time of x is smaller than that of reheating, that is, $H_{\text{osc}}^x > H_{\text{dec}}^x \gg H(T_R)$, with $H_{\text{osc}}^x \simeq m_X$ and $H_{\text{dec}}^x \simeq \Gamma(x \rightarrow \Psi_{3/2} \Psi_{3/2})$. The energy density of the coherent oscillation x is converted into that of the gravitino as

$$\rho_x \simeq \frac{1}{2} m_x^2 (\Delta x)^2 \simeq \frac{1}{2} \left(\frac{m_{3/2}^4}{m_x^2}\right) \left(\frac{a_{\text{dec}}^x}{a_{\text{osc}}^x}\right)^{-3} \left(\frac{a_R}{a_{\text{dec}}^x}\right)^{-4}, \quad (65)$$

at the time of reheating, where the gravitino is relativistic at the time of production. By employing the scale factors,

$$\begin{aligned} \frac{a_R}{a_{\text{osc}}^{\sigma^3}} &= \left(\frac{\sqrt{6}\Gamma_{\text{all}}^{\sigma^3}}{m_{T^3}}\right)^{-2/3}, \\ \frac{a_{\text{osc}}^x}{a_{\text{osc}}^{\sigma^3}} &= \left(\frac{\sqrt{6}m_X}{m_{T^3}}\right)^{-2/3}, \\ \frac{a_{\text{dec}}^x}{a_{\text{osc}}^{\sigma^3}} &= \left(\frac{6(\Gamma_{\text{all}}^x m_X)^2}{m_{3/2}^4}\right)^{-1/3}, \end{aligned} \quad (66)$$

the gravitino yield is

$$Y_{3/2}^x = \frac{2\rho_x}{m_{xS}} \simeq \frac{3}{2} \frac{T_R}{m_X} \left(\frac{m_{3/2}}{m_X}\right)^{16/3} \left(\frac{\Gamma_{\text{all}}^{\sigma^3}}{\Gamma_{\text{all}}^x}\right)^{2/3} \simeq 2 \times 10^{-32}, \quad (67)$$

with the numerical values given by Eqs. (42), (54), (55), and (64). It is found that the gravitino production via x decay is suppressed by the tiny mass of the gravitino and it is not the dominant source for the relic abundance of the gravitino. The entropy production from x can be also neglected in the same way as that of h_3 . As pointed out in Ref. [40], under $m_{3/2} \ll m_X \ll m_{T^3} \leq \Lambda$, the gravitino production is significantly relaxed and this condition is satisfied in our model. Note that when Λ is smaller than the inflaton mass, we have to take account of the inflaton decay into the fields in the hidden sector.

IV. GRAVITINO DARK MATTER AND THE HIGGS BOSON MASS

A. Yukawa couplings and naturalness

Before estimating the relic abundance of the gravitino, we specify the Yukawa couplings and μ -term in the superpotential which can be only introduced at the orbifold fixed points $y = 0, L$, where the SUSY is reduced to 4D $\mathcal{N} = 1$. As stated in Sec. III A, we consider the Yukawa interactions in the MSSM at the orbifold fixed point $y = 0$,

$$\begin{aligned} W_{\text{Yukawa}} &= \lambda_{ij}^u Q_i \mathcal{H}_u \mathcal{U}_j + \lambda_{ij}^d Q_i \mathcal{H}_d \mathcal{D}_j + \lambda_{ij}^e \mathcal{L}_i \mathcal{H}_d \mathcal{E}_j \\ &\quad + \lambda_{ij}^n \mathcal{L}_i \mathcal{H}_u \mathcal{N}_j, \end{aligned} \quad (68)$$

where $\lambda_{ij}^{u,d,e,n}$ are the holomorphic Yukawa coupling constants and are supposed to be of $\mathcal{O}(1)$. After the canonical normalization of fields in the MSSM, the physical Yukawa couplings are expressed as

$$\begin{aligned} y_{ij}^u &= \frac{\lambda_{ij}^u}{\sqrt{\langle Y_{Q_i} Y_{\mathcal{H}_u} Y_{\mathcal{U}_j} \rangle}}, & y_{ij}^d &= \frac{\lambda_{ij}^d}{\sqrt{\langle Y_{Q_i} Y_{\mathcal{H}_d} Y_{\mathcal{D}_j} \rangle}}, \\ y_{ij}^e &= \frac{\lambda_{ij}^e}{\sqrt{\langle Y_{\mathcal{L}_i} Y_{\mathcal{H}_d} Y_{\mathcal{E}_j} \rangle}}, & y_{ij}^n &= \frac{\lambda_{ij}^n}{\sqrt{\langle Y_{\mathcal{L}_i} Y_{\mathcal{H}_u} Y_{\mathcal{N}_j} \rangle}}, \end{aligned} \quad (69)$$

where

$$Y_a \equiv 2\mathcal{N}^{1/3} (\text{Re}T) \{Y(c_a \cdot T) + \tilde{\Omega}_{a,X}^{(4)} (\text{Re}T) |X|^2 + \mathcal{O}(|X|^4)\}. \quad (70)$$

The function $Y(z)$ is always positive, and approximated as

$$Y(z) \equiv \frac{1 - e^{-2\text{Re}z}}{2\text{Re}z} \simeq \begin{cases} \frac{1}{2\text{Re}z}, & \text{Re}z > 0 \\ \frac{1}{2|\text{Re}z|} \exp(2|\text{Re}z|), & \text{Re}z < 0. \end{cases} \quad (71)$$

In the 5D viewpoint, the wave functions of fields are localized toward $y = 0$ ($y = L$) in the case that $c_a \cdot (\text{Re}T)$ is positive (negative). As can be seen in Eq. (71), $y_{ij}^{u,d,e,n}$ are of $\mathcal{O}(1)$ or exponentially small when all the relevant fields are localized toward $y = 0$ or $y = L$, respectively.

Therefore, we expect that the mass hierarchies of elementary particles and the extreme smallness of the neutrino masses can be realized even in the case of Dirac neutrinos. In fact, when we choose the $U(1)_F$ charges and $\mathcal{O}(1)$ values of the holomorphic Yukawa couplings $\lambda_{i,j}^{u,d,e,n}$ in Tables I and II, the observed masses and mixing angles of quarks and leptons at the electroweak scale can be realized. Here, we employ the full one-loop RG equations of the MSSM from the GUT to the EW scale. It is remarkable that the flavor structure of soft SUSY-breaking terms is determined by the $U(1)_F$ charge assignment as can be seen in the Kähler potential (6). In fact, the soft

TABLE I. $U(1)_{F'}$ flavor charges of the quarks, leptons, and Higgs.

$c_{Q_i}^{I=1} = (0.1, 0.1, 1.1)$	$c_{L_i}^{I=1} = (0.1, 0.1, 1.6)$	$c_{H_u}^{I=1} = 0$
$c_{Q_i}^{I=2} = (-0.1, -0.1, 0.8)$	$c_{L_i}^{I=2} = (-0.1, -0.1, 0)$	$c_{H_u}^{I=2} = 0.1$
$c_{Q_i}^{I=3} = (0.1, 0.4, 1)$	$c_{L_i}^{I=3} = (0.1, 0.5, 0)$	$c_{H_u}^{I=3} = -0.9$
$c_{U_i}^{I=1} = (0.1, 0.1, 0.6)$	$c_{E_i}^{I=1} = (0.1, 0.2, 0.2)$	$c_{H_d}^{I=1} = 0$
$c_{U_i}^{I=2} = (-0.1, -0.1, 0.3)$	$c_{E_i}^{I=2} = (-0.1, -0.1, 0)$	$c_{H_d}^{I=2} = 0$
$c_{U_i}^{I=3} = (-0.2, 0.2, 1)$	$c_{E_i}^{I=3} = (-0.2, 0, -0.5)$	$c_{H_d}^{I=3} = -0.1$
$c_{D_i}^{I=1} = (0.1, 0.1, 0.2)$	$c_{N_i}^{I=1} = (0.1, 0.1, 0.1)$	
$c_{D_i}^{I=2} = (-0.1, -0.1, 0)$	$c_{N_i}^{I=2} = (-0.3, -0.3, -0.3)$	
$c_{D_i}^{I=3} = (0.3, 0.2, -0.5)$	$c_{N_i}^{I=3} = (-0.7, -0.7, -0.7)$	

TABLE II. $\mathcal{O}(1)$ values of the holomorphic Yukawa couplings $\lambda_{ij}^{u,d,e,n}$ in the superpotential (68).

$ \lambda_{ij}^u $	$ \lambda_{ij}^d $
$\begin{pmatrix} 0.32 & 0.35 & 0.95 \\ 0.22 & 0.42 & 0.33 \\ 0.51 & 0.48 & 1.5 \end{pmatrix}$	$\begin{pmatrix} 0.45 & 0.5 & 0.59 \\ 0.28 & 0.24 & 0.38 \\ 1.03 & 1.02 & 0.81 \end{pmatrix}$
$ \lambda_{ij}^e $	$ \lambda_{ij}^n $
$\begin{pmatrix} 0.28 & 0.22 & 0.52 \\ 0.4 & 1.15 & 0.31 \\ 0.8 & 1.02 & 1.05 \end{pmatrix}$	$\begin{pmatrix} 0.77 & 0.85 & 0.69 \\ 0.25 & 0.98 & 0.58 \\ 0.34 & 0.26 & 1.03 \end{pmatrix}$

SUSY-breaking terms at the GUT scale are determined by the following formula [41,42]:

$$\begin{aligned}
M_a &= \langle F^I \partial_I \ln(\text{Re} f^a) \rangle, \\
m_{Q_a}^2 &= -\langle F^I \bar{F}^{\bar{J}} \partial_I \partial_{\bar{J}} \ln Y_{Q_a} \rangle, \\
A_{ij}^u &= \langle F^I \partial_I \ln(Y_{\mathcal{H}_u} Y_{Q_i} Y_{U_j}) \rangle, \\
A_{ij}^d &= \langle F^I \partial_I \ln(Y_{\mathcal{H}_d} Y_{Q_i} Y_{D_j}) \rangle, \\
A_{ij}^e &= \langle F^I \partial_I \ln(Y_{\mathcal{H}_e} Y_{L_i} Y_{E_j}) \rangle, \\
A_{ij}^n &= \langle F^I \partial_I \ln(Y_{\mathcal{H}_n} Y_{L_i} Y_{N_j}) \rangle, \quad (72)
\end{aligned}$$

where indices I and J run over all the chiral multiplets. Then, the $U(1)_{F'}$ charge assignment in Table I and the F -term of the SUSY-breaking field X given by Eq. (42) give rise to the soft scalar masses and gluino mass in Table III. By contrast, the A -terms are almost vanishing due to the tiny F -terms of moduli. Here and hereafter, we parametrize the ratios of gaugino masses at the GUT scale as

$$r_1 = \frac{M_1(M_{\text{GUT}})}{M_3(M_{\text{GUT}})}, \quad r_2 = \frac{M_2(M_{\text{GUT}})}{M_3(M_{\text{GUT}})}, \quad (73)$$

where $M_1(M_{\text{GUT}})$, $M_2(M_{\text{GUT}})$, and $M_3(M_{\text{GUT}})$ are the bino, wino, and gluino masses at the GUT scale, $M_{\text{GUT}} \approx 2 \times 10^{16}$ GeV. The ratios of gaugino masses are controlled

TABLE III. The soft scalar masses $m_{\tilde{Q}_\alpha}$, the up- and down-type Higgs masses $M_{\mathcal{H}_{u,d}}$, and the gluino mass M_3 at the GUT scale. The subscripts \tilde{Q}_α denote the mass eigenvalues for the left-handed \tilde{Q}_i , up-type right-handed \tilde{U}_i , down-type right-handed \tilde{D}_i squarks, left-handed \tilde{L}_i , right-handed \tilde{E}_i charged sleptons, and right-handed sneutrinos \tilde{N}_i with the three-generation $i = 1, 2, 3$.

Sparticles	Mass [GeV]	(S)Particles	Mass [GeV]
$m_{\tilde{Q}_1}$	1682	$m_{\tilde{L}_3}$	2834
$m_{\tilde{Q}_2}$	1530	$m_{\tilde{E}_1}$	1157
$m_{\tilde{Q}_3}$	581	$m_{\tilde{E}_2}$	2390
$m_{\tilde{U}_1}$	1157	$m_{\tilde{E}_3}$	2298
$m_{\tilde{U}_2}$	1698	$m_{\tilde{N}_1}$	414.5
$m_{\tilde{U}_3}$	799	$m_{\tilde{N}_2}$	414.5
$m_{\tilde{D}_1}$	1636	$m_{\tilde{N}_3}$	414.5
$m_{\tilde{D}_2}$	1698	$M_{\mathcal{H}_u}$	1100
$m_{\tilde{D}_3}$	2298	$M_{\mathcal{H}_d}$	298.5
$m_{\tilde{L}_1}$	1682	M_3	550
$m_{\tilde{L}_2}$	1396		

by the parameters ξ_X^a in the gauge kinetic function (5) without spoiling the gauge coupling unification due to the tiny VEV of X .

On the other hand, the μ -term can be generated by the following superpotential:

$$W_{\mu\text{-term}} = \sum_{i=1}^3 \kappa_i H_i \mathcal{H}_u \mathcal{H}_d, \quad (74)$$

where κ_i are the $\mathcal{O}(1)$ dimensionless couplings, H_i are the stabilizer fields with R -charge 2, whereas Higgs chiral superfields do not have the R -charge. These cubic interactions do not affect the moduli stabilization as well as the moduli inflation due to the almost vanishing VEVs of the Higgs fields. Thus, it is possible to consider the VEVs of the stabilizer fields H_i as the origin of the μ -term. After the canonical normalization of the relevant fields, the μ -term at the GUT scale becomes

$$\mu = \sum_{i=1}^3 \frac{\kappa_i \langle H_i \rangle}{\langle Y_{H_i} Y_{\mathcal{H}_u} Y_{\mathcal{H}_d} \rangle}. \quad (75)$$

Especially, in the case of $\kappa_2 = \kappa_3 = 0$, the scale of the μ -term is chosen as TeV scale,

$$\mu \approx 3.8 \times 10^{-3} \frac{m_{3/2}}{m_{H_1}} M_{\text{Pl}} \approx \mathcal{O}(500 \text{ GeV}), \quad (76)$$

where $\kappa_1 = 2/3$, $m_{3/2} = 395 \text{ GeV}$, $m_{H_1} \approx 4.8 \times 10^{15} \text{ GeV}$, and $\langle H_1 \rangle \approx m_{3/2}/m_{H_1}$ are given by Eq. (23) and the factor 3.8×10^{-3} comes from the mild large volume of the fifth dimension and normalization factors for H_1 , \mathcal{H}_u , and \mathcal{H}_d . The EW symmetry breaking requires the following relation between the mass of the Z-boson, m_Z and soft SUSY-breaking masses of the up-type Higgs $m_{\mathcal{H}_u}$:

$$\frac{m_Z^2}{2} \approx -m_{\mathcal{H}_u}^2(M_{\text{EW}}) - |\mu(M_{\text{EW}})|^2 + \mathcal{O}\left(\frac{1}{\tan^2 \beta}\right), \quad (77)$$

in the limit of large value of $\tan \beta$, where $\mu(M_{\text{EW}})$ and $m_{\mathcal{H}_u}(M_{\text{EW}})$ are the μ -term and $m_{\mathcal{H}_u}$ at the EW scale, respectively. The VEVs of up- and down-type Higgs fields are denoted by $v_u = v \sin \beta$ and $v_d = v \cos \beta$ with $v = 174 \text{ GeV}$. Thus, the observed Z-boson mass indicates $|\mu(M_{\text{EW}})| \sim |m_{\mathcal{H}_u}(M_{\text{EW}})| \sim m_Z$; otherwise μ and $m_{\mathcal{H}_u}$ have to be properly tuned to obtain the EW vacuum. We adopt the measure of the degree of tuning the μ -term at the GUT scale as

$$\Delta_\mu = \frac{1}{2} \frac{\partial \ln m_Z^2}{\partial \ln |\mu|}, \quad (78)$$

and then $100 \times |\Delta_\mu^{-1}| \%$ represents the degree of tuning to obtain the Z-boson mass $m_Z = 91.2 \text{ GeV}$ [43]. Although the conventional CMSSM scenario requires more severe tuning than the degree of 0.1%, as pointed out in Ref. [2], certain ratios of the nonuniversal gaugino masses at the GUT scale relax the degree of tuning and observed 125 GeV Higgs boson mass at the same time.

B. Relic abundance of the gravitino

We are now ready to estimate the relic abundance of the gravitino. As stated in Sec. III E, there are no significant gravitino productions from the inflaton, moduli, stabilizer, and SUSY-breaking fields after the inflation. However, there are two processes to produce the gravitinos associated with the decay of other particles in the MSSM.

One of them is the decay from the thermal bath which is constituted of the relativistic particles after the reheating process. On the thermal bath, the dominant decay process comes from gauginos into gravitinos, because the couplings between the gravitino and other sparticles are more suppressed than those of gauginos as discussed in Refs. [44,45]. The abundance of the gravitino is estimated as




FIG. 2 (color online). Contours of the thermal abundance of the gravitino in the (r_1, r_2) -plane.

$$\Omega_{3/2}^{TP} h^2 = \sum_{a=1}^3 \left(1 + \frac{M_a(T_R)^2}{3m_{3/2}^2} \right) w_a g_a(T_R)^2 \times \ln \left(\frac{k_a}{g_a(T_R)} \right) \left(\frac{m_{3/2}}{100 \text{ GeV}} \right) \left(\frac{T_R}{10^{10} \text{ GeV}} \right), \quad (79)$$

where w_a and k_a are the parameters whose values are defined in Ref. [45] and h is a dimensionless Hubble parameter. The thermal production of the gravitino is drawn in Fig. 2 in terms of the ratios of gaugino masses at the GUT scale M_{GUT} , $r_1 = M_1(M_{\text{GUT}})/M_3(M_{\text{GUT}})$ and $r_2 = M_2(M_{\text{GUT}})/M_3(M_{\text{GUT}})$ with $M_3(M_{\text{GUT}}) = 550 \text{ GeV}$. The gaugino masses at the reheating temperature $M_a(T_R)$ can be expressed as $M_a(M_{\text{GUT}})$ by employing the one-loop RG equations in the MSSM. The Planck Collaboration reported that the abundance of dark matter resides in the range of $0.1175 \leq \Omega_{3/2}^{TP} h^2 \leq 0.1219$ [22], where the upper and lower limits correspond to the dotted curves in Fig. 2. Here we assume that the dark matter only consists of the abundance of the thermally produced gravitino.

The other process is the nonthermal gravitino productions from the NLSP and/or next-to-next-to-lightest supersymmetric particle (NNLSP). As shown in Table IV, when we take the ratios of gaugino masses $(r_1, r_2) = (6, 3.5)$ consistent with the observed relic abundance of dark matter in Fig. 2, the NLSPs and NNLSP correspond to the degenerated sneutrinos and Higgsino-like neutralino, respectively. The relevant sparticle spectra are obtained by employing the one-loop RG equations in the MSSM from the GUT to EW scale with $(r_1, r_2) = (6, 3.5)$ and the input parameters in Table III. The full sparticle spectra are shown in the next section. Note that the degenerated sneutrinos do not have sizable interactions with the other

TABLE IV. The masses of NNLSP, NLSPs, and the gravitino at the EW scale for the reference point $(r_1, r_2) = (6, 3.5)$. The subscripts denote the mass eigenvalues for the sneutrinos ($\tilde{\nu}$), the Higgsino-like neutralino ($\tilde{\chi}$).

NNLSP (Higgsino-like Neutralino)	Mass [GeV]
$\tilde{\chi}_3^0$	441
NLSPs (right-handed sneutrinos)	Mass [GeV]
$\tilde{\nu}_{e_2}$	415
$\tilde{\nu}_{\mu_2}$	415
$\tilde{\nu}_{\tau_2}$	415
LSP (gravitino)	Mass [GeV]
$\Psi_{3/2}$	395

(s)particles due to the tiny Yukawa couplings of Dirac-type neutrinos and then the soft SUSY-breaking masses of right-handed sneutrinos do not receive significant loop corrections.

Since the gravitino and right-handed sneutrinos are weakly coupled with the other (s)particles, they are not thermalized. Thus, the nonthermal gravitino productions from the Higgsino-like neutralino and sneutrinos are roughly estimated as

$$\Omega_{3/2}^{NTP} h^2 = \frac{m_{3/2}}{m_{\tilde{\chi}_3^0}} \Omega_{\tilde{\chi}_3^0} h^2, \quad (80)$$

where $m_{\tilde{\chi}_3^0}$ and $\Omega_{\tilde{\chi}_3^0}$ are the mass and the thermal abundance of the Higgsino-like neutralino $\tilde{\chi}_3^0$, respectively. The thermal abundance of the Higgsino-like neutralino is known to be small when the μ -term is smaller than wino and bino masses. Since the chargino and Higgsino-like neutralino are degenerated, both decay into the particles of the SM at almost the same decoupled time, which leads to the smallness of the thermal abundance of $\tilde{\chi}_3^0$. After all, the nonthermal abundance of the gravitino can be neglected,

$$\Omega_{3/2}^{NTP} h^2 \ll 0.11, \quad (81)$$

and the total relic abundance of the gravitino is approximated by the thermal abundance of it,⁸

$$\Omega_{3/2} h^2 \simeq \Omega_{3/2}^{TP} h^2. \quad (82)$$

However, the decays of neutralino and sneutrinos into the gravitino dark matter would threaten to spoil the successful BBN. The produced right-handed neutrinos via the sneutrino decay into the gravitino are suppressed due to the thermal abundance of $\tilde{\chi}_3^0$, and then they are harmless for the BBN. On the other hand, the Higgsino-like neutralino decay into the gravitino affects the BBN. The authors of Ref. [47] suggest a way to relax the constraints from the BBN by assuming that the NLSP is the Dirac-type

right-handed sneutrino. Although they consider the binolike neutralino as the NNLSP, the sparticle spectra are almost the same as our obtained one. Because of the small thermal abundance of the Higgsino-like neutralino, it is then expected that our spectra are consistent with the BBN.

Note that the nonthermal production of the gravitino is enhanced when the binolike neutralino is NNLSP which corresponds to the small value of $|r_1|$ in Fig. 2. In this case, it would break the successful BBN because of the large thermal abundance of the binolike neutralino [10,17,47].

C. The Higgs boson mass, gravitino dark matter, and sparticle spectra

The ratios of gaugino masses at the GUT scale, r_1 and r_2 , are severely constrained by the relic abundance of the gravitino as can be seen in Fig. 2. In this section, we show that the mass of the Higgs boson further constrains the ratios of gaugino masses, r_1 and r_2 . The lightest CP -even Higgs boson corresponds to the SM-like Higgs in the framework of MSSM. Without the loop corrections, the Higgs boson mass is much lower than the observed mass of the Higgs reported by the LHC experiment [48]. Although, the high-scale SUSY-breaking scenario is a simple solution as one of the possibilities to raise the Higgs mass, it requires the tuning to obtain the EW vacuum. Therefore, we consider the maximal mixing of left- and right-handed top squarks to raise the Higgs boson mass without a severe fine-tuning.

With an approximation that the mass eigenstates of top squarks are nearly degenerate, the mass of the lightest CP -even Higgs boson cannot be realized. Thus, as pointed out in [2], we add the contribution from the mass differences between left- and right-handed top squarks in order to realize the observed Higgs boson mass and relax the degree of tuning at the same time. By employing the full one-loop RG equations of the MSSM from the GUT to the EW scale, we numerically calculate the Higgs boson mass which resides in the range of $124.4 \leq m_h \leq 126.8$ [1], which is represented as the blue colored region in Fig. 3 and the degree of tuning a μ -term, $|\Delta_\mu| \times 100\%$, is also given by the green dashed (1%) and solid lines (10%), respectively. From Fig. 3, there are the parameter spaces which are consistent with the relic abundance of the gravitino and the Higgs boson mass reported by the current cosmological observations [22] as well as the collider experiments [1] without a severe fine-tuning.

In particular, at the reference point $(r_1, r_2) = (6, 3.5)$, the sparticle spectra, the Higgs boson mass m_h , and the degree of tuning a μ -term $|\Delta_\mu| \times 100(\%)$ are summarized in Tables IV, V, and VI. It is then satisfied by all the experimental lower bounds from the LHC experiments for the masses of all sparticles in Refs. [13] and [49]. In general, the SUSY flavor violations are dangerous in the gravity mediated SUSY-breaking scenario due to the flavor dependent interactions. In our setup, there are vanishing A -terms and no flavor dependent soft SUSY-breaking terms

⁸In this paper, we do not take the gravitino production by the primordial black hole into account [46].




FIG. 3 (color online). The Higgs boson mass, the degree of tuning a μ -term, $|\Delta_\mu| \times 100(\%)$, and the relic abundance of the gravitino $\Omega_{3/2} h^2$ on the (r_1, r_2) -plane. In the blue shaded regions, the Higgs boson mass resides in the allowed range, $124.4 \leq m_h \leq 126.8$ GeV [1]. The green dashed and solid lines show the 1% and 10% tuning, respectively. The red dashed curves show the relic abundance of the gravitino within ranges $0.1175 \leq \Omega_{3/2} h^2 \leq 0.1215$, reported by the Planck Collaboration [22].

at the GUT scale because the moduli do not have the F -terms. Even if the moduli have the F -terms at the vacuum, the SUSY flavor violations can be suppressed from the structure of the $U(1)_F$ charge assignments [50]. Thus, there

TABLE V. A typical sparticle spectra at the EW scale for the reference point, $(r_1, r_2) = (6, 3.5)$. The subscripts denote the mass eigenvalues for the following: up (\tilde{u}), charm (\tilde{c}), top (\tilde{t}), down (\tilde{d}), strange (\tilde{s}), bottom (\tilde{b}) squarks, the scalar electron (\tilde{e}), muon ($\tilde{\mu}$), tauon ($\tilde{\tau}$), neutrino ($\tilde{\nu}$), the neutralino ($\tilde{\chi}$), and the chargino ($\tilde{\chi}^\pm$).

Sparticles	Mass [GeV]	Sparticles	Mass [GeV]
\tilde{u}_1	2618	\tilde{e}_1	3241
\tilde{u}_2	2359	\tilde{e}_2	2525
\tilde{c}_1	2520	$\tilde{\mu}_1$	2421
\tilde{c}_2	2011	$\tilde{\mu}_2$	2331
\tilde{t}_1	1735	$\tilde{\tau}_1$	2133
\tilde{t}_2	974	$\tilde{\tau}_2$	1447
\tilde{d}_1	2625	$\tilde{\nu}_{e_1}$	3240
\tilde{d}_2	2620	$\tilde{\nu}_{e_2}$	415
\tilde{s}_1	2522	$\tilde{\nu}_{\mu_1}$	2330
\tilde{s}_2	2189	$\tilde{\nu}_{\mu_2}$	415
\tilde{b}_1	2117	$\tilde{\nu}_{\tau_1}$	2132
\tilde{b}_2	1724	$\tilde{\nu}_{\tau_2}$	415
$\tilde{\chi}_1^0$	1723	$\tilde{\chi}_1^\pm$	444
$\tilde{\chi}_2^0$	1135	$\tilde{\chi}_2^\pm$	1723
$\tilde{\chi}_3^0$	448		
$\tilde{\chi}_4^0$	441		

TABLE VI. The neutral and charged Higgs boson masses m_h , m_H , m_A , and m_{H^\pm} , the degree of tuning a μ -term, $|\Delta_\mu| \times 100(\%)$, and the gaugino masses at the EW scale for the reference point $(r_1, r_2) = (6, 3.5)$.

m_h [GeV]	m_H [GeV]	m_A [GeV]	m_{H^\pm} [GeV]
125.4	1423	1423	1425
$\Delta_\mu^{-1} \times 100(\%)$	$M_1(m_Z)$ [GeV]	$M_2(m_Z)$ [GeV]	$M_3(m_Z)$ [GeV]
2.1	1133	1719	1575

are no serious SUSY flavor violations; especially, the decay rates such as $\mu \rightarrow e\gamma$ and $b \rightarrow s\gamma$ evade the present limits [51,52].

V. CONCLUSION

In this paper, we proposed the gravitino dark matter in the gravity mediated SUSY-breaking scenario based on the 4D $\mathcal{N} = 1$ SUGRA. The nontrivial Kähler metric of the SUSY-breaking field induces the mass hierarchies between the gravitino and the other sparticles for any value of the F -term of the SUSY-breaking field. Especially, the small Kähler metric of the SUSY-breaking field leads to the stable gravitino of mass $\mathcal{O}(100)$ GeV with TeV scale gauginos and sparticles which would be the typical features in the natural MSSM with gravity mediation, if the gauge kinetic functions and the kinetic terms of the matter fields satisfy certain conditions. (See Ref. [21] for the case of CMSSM.) In the stable gravitino scenario, one can consider the low-scale SUSY without the cosmological gravitino problem, only if the NLSP decays do not spoil the success of BBN.

As a concrete model, we considered the 5D SUGRA model on S^1/Z_2 . Since the successful inflation mechanism as well as the moduli stabilization have been realized in 5D SUGRA [19,22], we have estimated the moduli and inflaton decays into the gravitino dark matter. Although the produced gravitinos via the moduli decays seem to be dangerous from the cosmological point of view, their decays can be suppressed only if the moduli do not have the F -terms. Such a situation can be applied in our model, because the moduli, inflaton, and stabilizer fields have supersymmetric masses at the vacuum. Even if the supersymmetry is broken in the SUSY-breaking sector, their F -terms are suppressed by the gravitino mass at the SUSY-breaking minimum. When the NLSP and NNLSP are taken as the sneutrino and Higgsino-like neutralino, the non-thermal productions of the gravitino are negligible due to the small thermal abundance of the Higgsino-like neutralino. The smallness of the thermal abundance of NNLSP also relaxes the constraints from the BBN [10,17,47], and at the same time, the amount of neutrinos via the sneutrino decay can be suppressed. Thus, the total relic abundance of the gravitino is approximated by the thermal abundance of it which depends on the gaugino masses. As pointed out in

[2], the certain ratios of gaugino masses are also important to raise the Higgs boson mass in the MSSM without a severe fine-tuning. From Fig. 3, it is found that certain ratios of gaugino masses are consistent with the relic abundance of the gravitino as well as the Higgs boson mass reported by the recent Planck and LHC data [1,22].

In this paper, we focus on the 5D SUGRA in order to show the realistic gravitino dark matter in the gravity mediation, and then the suppressed Kähler metric of the SUSY-breaking field is important to generate the mass hierarchies between the gravitino and other sparticles. When the 5D SUGRA is derived as the effective theory of superstring theories on a warped throat and/or M-theory on the Calabi-Yau manifold [53], the SUSY-breaking sector would be constructed from the gauge theory living on Dp-branes and/or NS5-branes. Especially, in the type II string theory, the visible and hidden sectors can be realized on the different D-branes which wrap the certain cycles in the internal manifold. In such cases, the different volumes of the internal cycles lead to the hierarchical Kähler metric between the SUSY-breaking field and matter fields in the visible sector.

ACKNOWLEDGMENTS

The author would like to thank H. Abe, T. Higaki, J. Kawamura, T. Kobayashi, and Y. Yamada for useful discussions and comments. H. O. is supported by Research Fellowships for Young Scientists (Grant No. 26-7296) of the Japan Society for the Promotion of Science.

APPENDIX A: THE F -TERMS OF FIELDS AT THE VACUUM

In this Appendix, we derive the F -terms of the moduli, stabilizer, and SUSY-breaking fields at the vacuum by employing the reference point method. As discussed in Sec. III C, when we expand the fields around the reference points given by Eqs. (16) and (21), $\phi \rightarrow \phi|_{\text{ref}} + \delta\phi$, $\phi = T^{I'}$, H_i , X with I' , $i = 1, 2, 3$, the Kähler metric with Kähler potentials (14) and (18) are expanded by

$$K_{I\bar{J}} = K_{I\bar{J}}^{(0)} + K_{I\bar{J}}^{(1)}, \quad (\text{A1})$$

where

$$K_{I\bar{J}}^{(0)} = \begin{pmatrix} 1/(2\text{Re}T^1)^2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1/(2\text{Re}T^2)^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1/(2\text{Re}T^3)^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & Z_{H_1} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & Z_{H_2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & Z_{H_3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & Z_X - 4|X|^2/\Lambda^2 \end{pmatrix}, \quad (\text{A2})$$

and

$$K_{I\bar{J}}^{(1)} = \begin{pmatrix} 0 & 0 & 0 & a_{H_1}H_1 & 0 & 0 & a_X^1 X \\ 0 & 0 & 0 & 0 & a_{H_2}H_2 & 0 & a_X^2 X \\ 0 & 0 & 0 & 0 & 0 & a_{H_3}H_3 & a_X^3 X \\ a_{H_1}\bar{H}_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & a_{H_2}\bar{H}_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{H_3}\bar{H}_3 & 0 & 0 & 0 & 0 \\ a_X^1\bar{X} & a_X^2\bar{X} & a_X^3\bar{X} & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (\text{A3})$$

in the field basis $(T^1, T^2, T^3, H_1, H_2, H_3, X)$, with

$$a_{H_i} \equiv \frac{\partial Z_{H_i}}{\partial T^{I'}} = \frac{1}{\text{Re}T^{I'}} \left(e^{-2c_{H_i}\text{Re}T^{I'}} - \frac{Z_{H_i}}{2} \right), \quad (I' = i),$$

$$a_X^i \equiv \frac{\partial Z_X}{\partial T^{I'}} = \frac{c_X^i}{c_X \cdot \text{Re}T} \left(e^{-2c_X \text{Re}T} - \frac{Z_X}{2} \right), \quad (\text{A4})$$

and the inverse of the Kähler metric is given by

$$\begin{aligned}
 K^{T^i \bar{T}^i} &\simeq (2\text{Re}T^i)^2 + 8\text{Re}T^i \text{Re}\delta T^i, & K^{T^i \bar{H}_i} &\simeq -A_{H_i} \delta H_i, \\
 K^{T^i \bar{X}} &\simeq -A_X^i X - A_X^i \delta X, & K^{H_i \bar{H}_i} &\simeq \frac{1}{Z_{H_i}} - \frac{2a_{H_i}}{(Z_{H_i})^2} \text{Re}\delta T^i, \\
 K^{X\bar{X}} &\simeq \frac{1}{Z_X} - \frac{2a_X^i}{(Z_X)^2} \text{Re}\delta T^i + \frac{4}{\Lambda^2 (Z_X)^2} |\delta X|^2,
 \end{aligned} \tag{A5}$$

where

$$A_{H_i} \equiv (2\text{Re}T^i)^2 \frac{a_{H_i}}{Z_{H_i}}, \quad A_X^i \equiv (2\text{Re}T^i)^2 \frac{a_X^i}{Z_X}. \tag{A6}$$

Here and hereafter, we omit the subscript of ϕ at the reference point, that is, $\phi = \phi|_{\text{ref}}$. From the relevant expansions in the scalar potential (1) with the Kähler and superpotential (14), (15), and (18),

$$\begin{aligned}
 D_{T^i} W &\simeq K_{T^i} w + W_{T^i H_i} \delta H^i + K_{T^i \bar{T}^i} w (\delta T^i + \delta \bar{T}^i) + K_{T^i} W_X \delta X \\
 &\quad + W_{T^i T^j H_i} \delta T^j \delta H_i + \sum_{J=k} K_{T^i} W_{T^j H_k} \delta T^k \delta H^k, \\
 D_{H_i} W &\simeq W_{T^i H_i} \delta T^i + K_{H_i \bar{H}_i} w \delta \bar{H}_i + \frac{W_{T^i T^j H_i}}{2} (\delta T^i)^2, \\
 D_X W &\simeq W_X + K_{X\bar{X}} w \delta \bar{X} + \frac{1}{2} \partial_X (K_{X\bar{X}}) w (2|\delta X|^2 + (\delta X)^2 + (\delta \bar{X})^2), \\
 W &\simeq w + W_X \delta X + \sum_{I=i}^3 W_{T^i H_i} \delta T^i \delta H_i, \\
 K &\simeq \sum_{I=1}^3 \left(-\ln(\text{Re}T^I) - \frac{\text{Re}\delta T^I}{\text{Re}T^I} + \frac{1}{2} \left(\frac{\text{Re}\delta T^k}{\text{Re}T^I} \right)^2 \right),
 \end{aligned} \tag{A7}$$

we obtain the scalar potential at the second order $\delta\phi$,

$$\begin{aligned}
 V &\simeq \frac{W_X^2}{Z_X} - 2w W_X (\delta X + \delta \bar{X}) - \sum_{I=i} (2\text{Re}T^I) w W_{T^i H_i} (\delta H_i + \delta \bar{H}_i) \\
 &\quad + \frac{4w^2}{\Lambda^2 Z_X^2} |\delta X|^2 + \sum_{I=i} \frac{W_{T^i H_i}^2}{Z_{H_i}} |\delta T^I|^2 + \sum_{I=i}^3 (2\text{Re}T^I)^2 W_{T^i H_i}^2 |\delta H_i|^2 \\
 &\quad + \sum_{I=i}^3 (-2\text{Re}T^I w W_{T^i T^j H_i} + w W_{T^i H_i}) (\delta T^I \delta H_i + \delta \bar{T}^I \delta \bar{H}_i) \\
 &\quad + \sum_{I=i}^3 \frac{A_{H_i}}{2\text{Re}T^I} w W_{T^i H_i} (\delta T^I \delta \bar{H}_i + \delta \bar{T}^I \delta H_i) - \sum_{I=1}^3 w W_{T^i H_i} (\delta T^I + \delta \bar{T}^I) (\delta H_i + \delta \bar{H}_i) \\
 &\quad + \sum_{I=i}^3 2\text{Re}T^I W_X W_{T^i H_i} (\delta H_i \delta \bar{X} + \delta \bar{H}_i \delta X) \\
 &\quad + \sum_{I=i}^3 \sum_{J'=1}^3 \frac{T^I + \bar{T}^I}{T^{J'} + \bar{T}^{J'}} w W_{T^i H_i} (\delta T^{J'} + \delta \bar{T}^{J'}) (\delta H_i + \delta \bar{H}_i).
 \end{aligned} \tag{A8}$$

Finally, the extremal conditions for the relevant fields lead to the following variations of them:

$$\begin{aligned}\delta H_i &\simeq \frac{w}{2\text{Re}T^{I'}W_{T^{I'}H_i}} \sim \mathcal{O}\left(\frac{m_{3/2}}{m_{H_i}}\right), & \delta X &\simeq \left(\frac{\Lambda^2 Z_{\bar{X}}^2}{4w^2}\right) 5wW_X, \\ \delta T^{I'} &\simeq \left(\frac{w}{W_{T^{I'}H_i}}\right)^2 Z_{H_i} \left(\frac{1+A_{H_i}K_{T^{I'}}}{2\text{Re}T^{I'}} + \frac{W_{T^{I'}T^{I'}H_i}}{W_{T^{I'}H_i}} - \frac{3}{\text{Re}T^{I'}}\right) \sim \mathcal{O}\left(\frac{m_{3/2}}{m_{T^{I'}}}\right),\end{aligned}\quad (\text{A9})$$

and their F -terms become

$$\begin{aligned}\sqrt{K_{T^{I'}\bar{T}^{I'}}}F^{T^{I'}} &= -e^{K/2}\sqrt{K_{T^{I'}\bar{T}^{I'}}}K^{T^{I'}\bar{J}}D_{\bar{J}}\bar{W} \sim \mathcal{O}\left(\frac{m_{3/2}^3}{m_{T^{I'}}^2}\right), \\ \sqrt{K_{H_i\bar{H}_i}}F^{H_i} &= -e^{K/2}\sqrt{K_{H_i\bar{H}_i}}K^{H_i\bar{J}}D_{\bar{J}}\bar{W} \sim \mathcal{O}\left(\frac{m_{3/2}^3}{m_{H_i}^2}\right), \\ \sqrt{K_{X\bar{X}}}F^X &\simeq -e^{K/2}\sqrt{K_{X\bar{X}}}K^{X\bar{X}}D_X W \simeq \frac{-W_X}{(\text{Re}T^1\text{Re}T^2\text{Re}T^3)^{1/2}Z_X^{1/2}},\end{aligned}\quad (\text{A10})$$

where

$$\begin{aligned}D_{T^{I'}}W &= \min\left(\mathcal{O}\left(\frac{m_{3/2}^3}{m_{T^{I'}}^2}\right), \mathcal{O}\left(\frac{m_{3/2}^3}{m_X^2}\right)\right), \quad (I' = 1, 2), \\ D_{T^3}W &= \mathcal{O}\left(\frac{m_{3/2}^3}{m_{T^3}^2}\right), \quad D_{H_i}W = \mathcal{O}\left(\frac{m_{3/2}^3}{m_{H_i}^2}\right), \quad (i = 1, 2, 3), \quad D_X W \simeq \nu.\end{aligned}\quad (\text{A11})$$

We also numerically checked these results, and then their F -terms can be suppressed by the tiny mass of the gravitino.

APPENDIX B: THE MINIMA OF FIELDS DURING THE INFLATION

By contrast, the minima of fields during the inflation are different from those at the true vacuum. In this section, we derive the minima of fields by employing the reference point method. The reference points of fields during the inflation are chosen in the same way as those at the vacuum.

Similarly, we expand the fields except for the inflaton $\text{Re}T^3$ around the reference points given by Eqs. (16) and (21), $\phi \rightarrow \phi|_{\text{ref}} + \delta\phi$, $\phi = T^{I'}, \text{Im}T^3, H_i, X$ with $I' = 1, 2, i = 1, 2, 3$. It is then supposed that H_3 is fixed at the origin due to the Hubble-induced mass. From the scalar potential (1) with the Kähler and superpotentials (14), (15), and (18) given by the following expansions:

$$\begin{aligned}D_{T^{I'}}W &\simeq K_{T^{I'}}w + W_{T^{I'}H_i}\delta H_i + K_{T^{I'}\bar{T}^{I'}}w(\delta T^{I'} + \delta\bar{T}^{I'}) + K_{T^{I'}}(W_{H_3}\delta H_3 + W_X\delta X) \\ &\quad + K_{T^{I'}\bar{T}^{I'}}W_{H_3}(\delta T^{I'} + \delta\bar{T}^{I'})\delta H^3 + W_{T^{I'}T^{I'}H_i}\delta T^{I'}\delta H_i + K_{T^{I'}}\sum_{J'=j}^3 W_{T^{I'}H_j}\delta T^{J'}\delta H^j, \\ D_{H_1}W &\simeq W_{T^1H_1}\delta T^1 + K_{H_1\bar{H}_1}w\delta\bar{H}_1 + K_{H_1\bar{H}_1}W_{H_3}\delta\bar{H}_1\delta H_3, \\ D_{H_2}W &\simeq W_{T^2H_2}\delta T^2 + K_{H_2\bar{H}_2}w\delta\bar{H}_2 + K_{H_2\bar{H}_2}W_{H_3}\delta\bar{H}_2\delta H_3, \\ D_{H_3}W &\simeq W_{H_3} + W_{T^3H_3}\delta T^3 + K_{H_3\bar{H}_3}W\delta\bar{H}_3 + \partial_{T^3}(K_{H_3\bar{H}_3})w(\delta T^3 + \delta\bar{T}^3)\delta\bar{H}_3, \\ &\quad + K_{H_3\bar{H}_3}(W_{H_3}|\delta H_3|^2 + W_X\delta\bar{H}_3\delta X) + \frac{W_{T^3T^3H_3}}{2}(\delta T^3)^2, \\ D_X W &\simeq W_X + K_{X\bar{X}}w\delta\bar{X} + \frac{1}{2}\partial_X(K_{X\bar{X}})w(2|\delta X|^2 + (\delta X)^2 + (\delta\bar{X})^2) + K_{X\bar{X}}W_{H_3}\delta H_3\delta\bar{X}, \\ W &\simeq w + W_X\delta X + W_{H_3}\delta H_3 + \sum_{I'=i}^3 W_{T^{I'}H_i}\delta T^{I'}\delta H_i, \\ K &\simeq \sum_{I'=i}^3 \left(-\ln \text{Re}T^{I'} - \frac{\text{Re}\delta T^{I'}}{\text{Re}T^{I'}} + \frac{1}{2}\left(\frac{\text{Re}\delta T^{I'}}{\text{Re}T^{I'}}\right)^2\right) + \sum_i Z_{H_i}|\delta H_i|^2 + Z_X|\delta X|^2,\end{aligned}\quad (\text{B1})$$

we obtain the extremal conditions for the relevant fields, and then their variations become

$$\begin{aligned}
\delta\tau^1 &= \delta\tau^2 = \delta\tau^3 = \delta k_1 = \delta k_2 = \delta k_3 = \delta y = 0, \\
\delta\sigma^{I'} &\sim \mathcal{O}\left(\frac{Z_{H_i}}{W_{T^{I'}H_i}^2 \text{Re}T^{I'}} \frac{|W_{H_3}|^2}{Z_{H_3}}\right) \simeq \mathcal{O}\left(\left(\frac{H_{\text{inf}}}{m_{T^{I'}}}\right)^2\right), \quad (I' = 1, 2), \\
\delta h_i &\sim \mathcal{O}\left(\frac{K_{T^{I'}W}}{W_{T^{I'}H_i} (2\text{Re}T^{I'})^2}\right) \simeq \mathcal{O}\left(\frac{m_{3/2}}{m_{H_i}}\right), \quad (i = 1, 2), \\
\delta h_3 &\sim \mathcal{O}\left(\frac{w}{W_{H_3}}\right) \simeq \mathcal{O}\left(\frac{m_{3/2}}{m_{H_3}}\right), \\
\delta x &\sim \mathcal{O}\left(\frac{\Lambda^2 Z_X^2}{4W_X^2} W_X w\right) \simeq \mathcal{O}\left(\left(\frac{m_{3/2}}{m_X}\right)^2\right), \tag{B2}
\end{aligned}$$

where

$$\delta T^{I'} \equiv \delta\sigma^{I'} + i\delta\tau^{I'}, \quad \delta H_i \equiv \delta h_i + i\delta k_i, \quad \delta X \equiv \delta x + i\delta y, \tag{B3}$$

with $I', i = 1, 2, 3$.

-
- [1] G. Aad *et al.* (ATLAS Collaboration), Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012); S. Chatrchyan *et al.* (CMS Collaboration), Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
- [2] H. Abe, T. Kobayashi, and Y. Omura, Relaxed fine-tuning in models with nonuniversal gaugino masses, *Phys. Rev. D* **76**, 015002 (2007); H. Abe, J. Kawamura, and H. Otsuka, The Higgs boson mass in a natural MSSM with nonuniversal gaugino masses at the GUT scale, *Prog. Theor. Exp. Phys.* **2013**, 013B02 (2013).
- [3] L. J. Hall, J. D. Lykken, and S. Weinberg, Supergravity as the messenger of supersymmetry breaking, *Phys. Rev. D* **27**, 2359 (1983); S. K. Soni and H. A. Weldon, Analysis of the supersymmetry breaking induced by $\mathcal{N} = 1$ supergravity theories, *Phys. Lett.* **126B**, 215 (1983).
- [4] G. F. Giudice and R. Rattazzi, Theories with gauge mediated supersymmetry breaking, *Phys. Rep.* **322**, 419 (1999).
- [5] L. Randall and R. Sundrum, Out of this world supersymmetry breaking, *Nucl. Phys. B* **557**, 79 (1999); G. F. Giudice, M. A. Luty, H. Murayama, and R. Rattazzi, Gaugino mass without singlets, *J. High Energy Phys.* **12** (1998) 027.
- [6] H. Pagels and J. R. Primack, Supersymmetry, Cosmology and New TeV Physics, *Phys. Rev. Lett.* **48**, 223 (1982); S. Weinberg, Cosmological Constraints on the Scale of Supersymmetry Breaking, *Phys. Rev. Lett.* **48**, 1303 (1982).
- [7] M. Y. Khlopov and A. D. Linde, Is it easy to save the gravitino?, *Phys. Lett.* **138B**, 265 (1984).
- [8] T. Moroi, H. Murayama, and M. Yamaguchi, Cosmological constraints on the light stable gravitino, *Phys. Lett. B* **303**, 289 (1993).
- [9] M. Kawasaki, K. Kohri, and T. Moroi, Big-Bang nucleosynthesis and hadronic decay of long-lived massive particles, *Phys. Rev. D* **71**, 083502 (2005); K. Kohri, T. Moroi, and A. Yotsuyanagi, Big-bang nucleosynthesis with unstable gravitino and upper bound on the reheating temperature, *Phys. Rev. D* **73**, 123511 (2006).
- [10] T. Kanzaki, M. Kawasaki, K. Kohri, and T. Moroi, Cosmological constraints on gravitino LSP scenario with sneutrino NLSP, *Phys. Rev. D* **75**, 025011 (2007); M. Kawasaki, K. Kohri, T. Moroi, and A. Yotsuyanagi, Big-bang nucleosynthesis and gravitino, *Phys. Rev. D* **78**, 065011 (2008).
- [11] C. Kelso, S. Profumo, and F. S. Queiroz, Nonthermal WIMPs as “dark radiation” in Light of ATACAMA, SPT, WMAP9, and Planck, *Phys. Rev. D* **88**, 023511 (2013); R. Allahverdi, B. Dutta, F. S. Queiroz, L. E. Strigari, and M. Y. Wang, Dark matter from late invisible decays to/of gravitinos, *Phys. Rev. D* **91**, 055033 (2015).
- [12] T. Moroi and L. Randall, Wino cold dark matter from anomaly mediated SUSY breaking, *Nucl. Phys. B* **570**, 455 (2000).
- [13] G. Aad *et al.* (ATLAS Collaboration), Search for strong production of supersymmetric particles in final states with missing transverse momentum and at least three b -jets at $\sqrt{s} = 8$ TeV proton-proton collisions with the ATLAS detector, *J. High Energy Phys.* **10** (2014) 24.
- [14] K. Choi, K. S. Jeong, and K. i. Okumura, Phenomenology of mixed modulus-anomaly mediation in fluxed string compactifications and brane models, *J. High Energy Phys.* **09** (2005) 039; M. Endo, M. Yamaguchi, and K. Yoshioka, Bottom-up approach to moduli dynamics in heavy gravitino

- scenario: Superpotential, soft terms, and sparticle mass spectrum, *Phys. Rev. D* **72**, 015004 (2005).
- [15] J. L. Feng, A. Rajaraman, and F. Takayama, Superweakly Interacting Massive Particles, *Phys. Rev. Lett.* **91**, 011302 (2003); J. L. Feng, A. Rajaraman, and F. Takayama, SuperWIMP dark matter signals from the early universe, *Phys. Rev. D* **68**, 063504 (2003).
- [16] J. R. Ellis, K. A. Olive, Y. Santoso, and V. C. Spanos, Gravitino dark matter in the CMSSM, *Phys. Lett. B* **588**, 7 (2004).
- [17] J. L. Feng, S. f. Su, and F. Takayama, SuperWIMP gravitino dark matter from slepton and sneutrino decays, *Phys. Rev. D* **70**, 063514 (2004).
- [18] H. Abe and H. Otsuka, Moduli inflation in five-dimensional supergravity models, *J. Cosmol. Astropart. Phys.* **11** (2014) 027.
- [19] N. Maru and N. Okada, Supersymmetric radius stabilization in warped extra dimensions, *Phys. Rev. D* **70**, 025002 (2004).
- [20] N. Arkani-Hamed and M. Schmaltz, Hierarchies without symmetries from extra dimensions, *Phys. Rev. D* **61**, 033005 (2000); D. E. Kaplan and T. M. P. Tait, Supersymmetry breaking, fermion masses and a small extra dimension, *J. High Energy Phys.* **06** (2000) 020.
- [21] J. Kersten and O. Lebedev, Gravitino dark matter in gravity mediation, *Phys. Lett. B* **678**, 481 (2009).
- [22] P. A. R. Ade *et al.* (Planck Collaboration), Planck 2015 results. XX. Constraints on inflation, [arXiv:1502.02114](https://arxiv.org/abs/1502.02114).
- [23] M. Zucker, Minimal off-shell supergravity in five-dimensions, *Nucl. Phys.* **B570**, 267 (2000); M. Zucker, Gauged $N = 2$ off-shell supergravity in five-dimensions, *J. High Energy Phys.* **08** (2000) 016; M. Zucker, Supersymmetric brane world scenarios from off-shell supergravity, *Phys. Rev. D* **64**, 024024 (2001); M. Zucker, Off-shell supergravity in five-dimensions and supersymmetric brane world scenarios, *Fortschr. Phys.* **51**, 899 (2003).
- [24] T. Kugo and K. Ohashi, Off-shell $D = 5$ supergravity coupled to matter Yang-Mills system, *Prog. Theor. Phys.* **105**, 323 (2001); T. Fujita and K. Ohashi, Superconformal tensor calculus in five-dimensions, *Prog. Theor. Phys.* **106**, 221 (2001); T. Fujita, T. Kugo, and K. Ohashi, Off-shell formulation of supergravity on orbifold, *Prog. Theor. Phys.* **106**, 671 (2001); T. Kugo and K. Ohashi, Superconformal tensor calculus on orbifold in 5D, *Prog. Theor. Phys.* **108**, 203 (2002).
- [25] H. Abe and Y. Sakamura, Dynamical radion superfield in 5D action, *Phys. Rev. D* **71**, 105010 (2005); H. Abe and Y. Sakamura, Consistent dimensional reduction of five-dimensional off-shell supergravity, *Phys. Rev. D* **73**, 125013 (2006).
- [26] F. Paccetti Correia, M. G. Schmidt, and Z. Tavartkiladze, 4D superfield reduction of 5-D orbifold SUGRA and heterotic M-theory, *Nucl. Phys.* **B751**, 222 (2006).
- [27] H. Abe and Y. Sakamura, Roles of $Z(2)$ -odd $N = 1$ multiplets in off-shell dimensional reduction of 5D supergravity, *Phys. Rev. D* **75**, 025018 (2007).
- [28] H. Abe and Y. Sakamura, Superfield description of 5-D supergravity on general warped geometry, *J. High Energy Phys.* **10** (2004) 013.
- [29] F. Paccetti Correia, M. G. Schmidt, and Z. Tavartkiladze, Superfield approach to 5D conformal SUGRA and the radion, *Nucl. Phys.* **B709**, 141 (2005).
- [30] Y. Sakamura, Direct relation of linearized supergravity to superconformal formulation, *J. High Energy Phys.* **12** (2011) 008.
- [31] Y. Sakamura, Superfield description of gravitational couplings in generic 5D supergravity, *J. High Energy Phys.* **07** (2012) 183.
- [32] Y. Sakamura, One-loop Kaehler potential in 5D gauged supergravity with generic prepotential, *Nucl. Phys.* **B873**, 165 (2013); **B873**, 728(E) (2013); Y. Sakamura and Y. Yamada, Impacts of non-geometric moduli on effective theory of 5D supergravity, *J. High Energy Phys.* **11** (2013) 090; **01** (2014) 181(E).
- [33] H. Abe and Y. Sakamura, Moduli stabilization in 5D gauged supergravity with universal hypermultiplet and boundary superpotentials, *Nucl. Phys.* **B796**, 224 (2008).
- [34] L. O’Raifeartaigh, Spontaneous symmetry breaking for chiral scalar superfields, *Nucl. Phys.* **B96**, 331 (1975).
- [35] R. Kallosh and A. D. Linde, O’KKLT, *J. High Energy Phys.* **02** (2007) 002; R. Kitano, Gravitational gauge mediation, *Phys. Lett. B* **641**, 203 (2006).
- [36] H. Abe, T. Higaki, T. Kobayashi, and Y. Omura, Moduli stabilization, F -term uplifting, and soft supersymmetry breaking terms, *Phys. Rev. D* **75**, 025019 (2007).
- [37] A. A. Starobinsky, A new type of isotropic cosmological models without singularity, *Phys. Lett.* **91B**, 99 (1980).
- [38] M. Endo, K. Hamaguchi, and F. Takahashi, Moduli-Induced Gravitino Problem, *Phys. Rev. Lett.* **96**, 211301 (2006).
- [39] S. Nakamura and M. Yamaguchi, Gravitino production from heavy moduli decay and cosmological moduli problem revived, *Phys. Lett. B* **638**, 389 (2006).
- [40] K. Nakayama, F. Takahashi, and T. T. Yanagida, Eluding the gravitino overproduction in inflaton decay, *Phys. Lett. B* **718**, 526 (2012).
- [41] K. Choi, A. Falkowski, H. P. Nilles, and M. Olechowski, Soft supersymmetry breaking in KKLT flux compactification, *Nucl. Phys.* **B718**, 113 (2005).
- [42] V. S. Kaplunovsky and J. Louis, Model independent analysis of soft terms in effective supergravity and in string theory, *Phys. Lett. B* **306**, 269 (1993).
- [43] R. Barbieri and G. F. Giudice, Upper bounds on supersymmetric particle masses, *Nucl. Phys.* **B306**, 63 (1988).
- [44] M. Bolz, A. Brandenburg, and W. Buchmuller, Thermal production of gravitinos, *Nucl. Phys.* **B606**, 518 (2001); **B790**, 336(E) (2008).
- [45] J. Pradler and F. D. Steffen, Thermal gravitino production and collider tests of leptogenesis, *Phys. Rev. D* **75**, 023509 (2007).
- [46] M. Y. Khlopov, A. Barrau, and J. Grain, Gravitino production by primordial black hole evaporation and constraints on the inhomogeneity of the early universe, *Classical Quantum Gravity* **23**, 1875 (2006).
- [47] K. Ishiwata, S. Matsumoto, and T. Moroi, Gravitino dark matter with weak-scale right-handed sneutrino, *Phys. Rev. D* **77**, 035004 (2008).
- [48] Y. Okada, M. Yamaguchi, and T. Yanagida, Upper bound of the lightest Higgs boson mass, *Prog. Theor. Phys.* **85**, 1 (1991).

- [49] J. Beringer *et al.* (Particle Data Group Collaboration), Review of Particle Physics, *Phys. Rev. D* **86**, 010001 (2012).
- [50] H. Abe, H. Otsuka, Y. Sakamura, and Y. Yamada, SUSY flavor structure of generic 5D supergravity models, *Eur. Phys. J. C* **72**, 2018 (2012).
- [51] J. Adam *et al.* (MEG Collaboration), New Constraint on the Existence of the $\mu^+ \rightarrow e^+\gamma$ Decay, *Phys. Rev. Lett.* **110**, 201801 (2013).
- [52] Y. Amhis *et al.* (Heavy Flavor Averaging Group Collaboration), Averages of b -hadron, c -hadron, and τ -lepton properties as of summer 2014, [arXiv:1412.7515](https://arxiv.org/abs/1412.7515).
- [53] A. Lukas, B. A. Ovrut, K. S. Stelle, and D. Waldram, Universe as a domain wall, *Phys. Rev. D* **59**, 086001 (1999); A. Lukas, B. A. Ovrut, K. S. Stelle, and D. Waldram, Heterotic M-theory in five-dimensions, *Nucl. Phys.* **B552**, 246 (1999).