

G_2 monopolesYa. Shnir^{1,2} and G. Zhilin²¹*BLTP, JINR, Dubna, Russia*²*Department of Theoretical Physics and Astrophysics, Belarusian State University, Minsk 220004, Belarus*
(Received 2 June 2015; published 20 August 2015)

We investigate some aspects of Bogomolny-Prasad-Sommerfield monopole solutions in the Yang-Mills-Higgs theory with the exceptional gauge group G_2 spontaneously broken to $U(1) \times U(1)$. The corresponding homotopy group is $\pi_2(G_2/U(1) \times U(1))$, and similarly to the $SU(3)$ theory, the G_2 monopoles are classified by two topological charges (n_1, n_2) . In the fundamental representation, these yield a subset of $SO(7)$ monopole configurations. Through inspection of the structure of $Alg(G_2)$, we propose an extension of the Nahm construction to the $(n, 1)_{G_2}$ monopoles. For the $(1, 1)_{G_2}$ monopole, the Nahm data are written explicitly.

DOI: [10.1103/PhysRevD.92.045025](https://doi.org/10.1103/PhysRevD.92.045025)

PACS numbers: 14.80.Hv, 11.27.+d

I. INTRODUCTION

Classical monopole solutions of spontaneously broken Yang-Mills-Higgs theories have long been the objects of detailed study.¹ These topologically nontrivial field configurations may exist in gauge theories for an arbitrary semisimple compact Lie group [4,5]. The simplest example is the 't Hooft-Polyakov monopole in the $SU(2)$ theory [6,7]. In the Bogomolny-Prasad-Sommerfield (BPS) limit [8,9], the potential of the scalar field is vanishing and the monopole solution is given by the first-order equation which is integrable. Furthermore, the Bogomolny equation can be treated as a dimensionally reduced self-duality equation, and there is a duality between the monopole solutions of the Bogomolny equation and the matrix valued Nahm data [10]. Nahm's construction is a very powerful tool for constructing various multimono poles in different models [11–16]; it also has a very interesting realization in the context of the construction of D-branes [17].

Nahm's construction can be generalized for all classical groups, as $SU(N)$ [14], symplectic and orthogonal groups [15,16,18]. Here we will concentrate on the case of the smallest simply connected compact exceptional group with a trivial center G_2 . Topologically nontrivial boundary conditions of the scalar field yield a nontrivial second homotopy group of the vacuum where the symmetry is broken to a residue group H , thus there are monopole solutions of the G_2 Yang-Mills-Higgs theory.

Gauge theories with symmetry group G_2 have attracted much attention recently [19–25]. One of the reasons is that such a theory is similar to usual $SU(3)$ gluodynamics; thus it is useful to investigate how the center symmetry is relevant for deconfinement phase transition in the lattice G_2 gluodynamics [19,20,22,23]. Recently, it was shown that in supersymmetric Yang-Mills theory, confinement-deconfinement transition does not break the symmetry of

the G_2 ground state although the expectation value of the Wilson line exhibits a discontinuity [25].

On the other hand, the gauge group G_2 is the automorphism group of the division algebra of octonions. This property allows one to construct an octonionic instanton solution to the seven-dimensional G_2 Yang-Mills theory [24]. Also, the massless monopole states in the $N = 2$ supersymmetric Yang-Mills theory with symmetry group G_2 were considered recently [21].

Note that the coupling of the gauge sector to the Higgs field in the seven-dimensional fundamental representation of G_2 may break this symmetry to $SU(3)$; however, in this case some fundamental monopoles, i.e. the monopoles associated with simple roots of the gauge group G_2 , become massless (see e.g. Ref. [2]). In this paper we will mainly consider another, more simple situation, when the gauge symmetry is broken maximally by an adjoint Higgs mechanism to $U(1) \times U(1)$. In this case the monopoles have two topological charges with respect to either of the unbroken Abelian groups $U(1)$; thus the monopoles can be labeled by two integers (n_1, n_2) .

The organization of the paper is as follows: Section II is a review of the basic properties of the first exceptional group G_2 ; there we also review the Nahm formalism. Section III contains our results of construction of the $(1, 1)_{G_2}$ monopoles. In Sec. IV we conclude with some additional remarks. In additional appendixes, we summarize the relevant information about the \mathfrak{g}_2 algebra and its representation.

II. EXCEPTIONAL GROUP G_2 AND THE NAHM CONSTRUCTION

We start with some introductory remarks about the Lie group G_2 . It is the smallest of the five exceptional simple Lie groups with a trivial central element. Mathematically, it can be thought as the group of automorphisms of the octonions or as a subgroup of the real orthogonal group $SO(7)$, which leaves one element of the eight-dimensional

¹For a review, see Refs. [1–3].

real spinor representation invariant. It is one of three simple Lie groups of rank two: $SU(3)$, $O(5)$ and G_2 . The fundamental representation of G_2 is seven-dimensional; the number of generators of the corresponding algebra is 14 (we refer to Table I, Appendix A for details). Thus, the Cartan subgroup contains two commuting generators H_1, H_2 . The roots and coroots of the G_2 are shown in Fig. 1.

Explicitly, we can take the elements of the Cartan subalgebra \mathbf{H}

$$\begin{aligned} H_1 &= \frac{1}{4} \text{diag}(-1, 1, -2, 0, 2, -1, 1), \\ H_2 &= \frac{1}{4\sqrt{3}} \text{diag}(0, -1, 1, 0, -1, 1, 0) \end{aligned} \quad (1)$$

so that the Killing form $\mathcal{K}(H_i, H_j) = \frac{1}{2} \delta_{ij}$.

Thereafter, we consider the Yang-Mills-Higgs theory in the BPS limit. Then the monopoles are solutions of the first-order Bogomolny equation

$$D_k \Phi = B_k. \quad (2)$$

The asymptotic value of the Higgs field along the positive direction of the third axis lies in the Cartan subalgebra: $\Phi_\infty = \mathbf{h} \cdot \mathbf{H}$. If the G_2 symmetry is maximally broken to $U(1) \times U(1)$, all roots have nonvanishing inner products with vector \mathbf{h} and, since $\pi_1(G_2) = 0$, the monopole solutions are classified according to the homotopy group $\pi_2(G_2/U(1) \times U(1)) = \pi_1(U(1) \times U(1)) = \mathbb{Z} \times \mathbb{Z}$. Recall that the magnetic field of the monopole configuration asymptotically also lies in the Cartan subalgebra

$$B_k = \mathbf{g} \cdot \mathbf{H} \frac{r_k}{4\pi r^3}. \quad (3)$$

Therefore, the quantized magnetic charge is

$$\mathbf{g} = \frac{4\pi}{e} (n_1 \alpha_1^* + n_2 \alpha_2^*), \quad (4)$$




FIG. 1. Root diagram of G_2 theory.

where two integers n_1, n_2 are topological charges of the monopoles given by embedding along the corresponding simple roots, and there are two distinct charge-1 fundamental monopoles which correspond to embeddings along the roots α_1 and α_2 ; they are (1,0) and (0,1), respectively. Thus, any (n_1, n_2) G_2 monopole can be viewed as a collection of n_1 individual α_1 fundamental monopoles and n_2 α_2 fundamental monopoles.

Then, making use of an explicit seven-dimensional representation of \mathfrak{g}_2 , the asymptotic of the Higgs field is of the form

$$\begin{aligned} \Phi &= \text{diag}(-s_1 - s_2, -s_2, -s_1, 0, s_1, s_2, s_1 + s_2) \\ &\quad - \frac{1}{2er} \text{diag}(-n_2, -n_1 + n_2, n_1 - 2n_2, 0, -n_1 \\ &\quad + 2n_2, n_1 - n_2, n_2) + O(r^{-1}), \end{aligned} \quad (5)$$

where $s_2 > s_1 > 0$ to follow the conventional ordering. The mass of the corresponding (n_1, n_2) configuration is given by

$$\begin{aligned} M &= \frac{4\pi}{e} [n_1 \mathbf{h} \cdot \alpha_1^* + n_2 \mathbf{h} \cdot \alpha_2^*] \\ &= \frac{4\pi}{e} [8n_1(s_2 - s_1) + 24n_2 s_1]. \end{aligned} \quad (6)$$

Let us briefly discuss the special case of nonmaximal symmetry breaking. Clearly, there are two situations when one of the G_2 monopoles becomes massless, $s_1 = s_2$ and $s_1 = 0$. The first case corresponds to the situation when the vector of the Higgs field is orthogonal to the long root α_1 and the symmetry is broken to $SU(2) \times U(1)$. In the second case, the Higgs field is orthogonal to the short root α_2 and the symmetry is broken to $U(1) \times SU(2)$. The total magnetic charge of these configurations is Abelian when the configuration remains invariant with respect to the transformations from the unbroken subgroup; such configurations are $([3n], 2n)$ and $(2n, [n])$, where the square brackets denote the holomorphic charge which counts the number of massless monopoles [26].

The Nahm construction can be considered as a duality between the Bogomolny equation (2) in \mathbb{R}^3 and solutions of the Nahm equation in one-dimensional space

$$\frac{dT_i}{ds} = \frac{1}{2} \epsilon_{ijk} [T_j, T_k], \quad (7)$$

where the Nahm data $T_k(s)$ are matrix-valued functions of a variable s over the finite interval given by the eigenvalues of the Higgs field on the spatial boundary. The first step of the Nahm construction is to find a solution of the linear differential equation (7) which must satisfy certain boundary conditions imposed on the end points of the interval of values of variable s . The second step is to solve the

construction equation² on the eigenfunctions $\omega(\mathbf{r}, s)$ of the linear operator which includes the Nahm data:

$$\left[-\mathbb{I}_{2k} \frac{d}{ds} + (r_i \mathbb{I}_k - T_i^{(k)}) \otimes \sigma_i \right] \omega(\mathbf{r}, s) + (v^{(k)})^\dagger S^{(k)}(\mathbf{r}) = 0. \quad (8)$$

Finally, the normalizable eigenfunctions allow us to recover the spacetime fields of the BPS monopole as

$$\begin{aligned} \Phi_{nm} &= \int_{s_1}^{s_2} ds s \omega_n^\dagger(s, \mathbf{r}) \omega_m(s, \mathbf{r}), \\ A_{nm}^k &= -i \int_{s_1}^{s_2} ds \omega_n^\dagger(s, \mathbf{r}) \partial^k \omega_m(s, \mathbf{r}), \end{aligned} \quad (9)$$

where s_1, s_2 are the endpoints of the interval of values of variable s .

This kind of duality was investigated in many papers; for a review, see Ref. [2], especially in the case of the gauge group $SU(2)$. In such a case it is possible to prove the isometry between the hyper-Kähler metrics of the moduli spaces of Nahm data and BPS monopoles. The conjecture about general equivalence of the metric on

the moduli space of the Hahm data and the metric on the monopole moduli space was used, for example, to calculate the metric on the moduli space of (2,1) $SU(3)$ monopoles [12].

The Nahm approach can be generalized to all classical groups [11,26]. The asymptotic Higgs field of the $SU(N)$ monopoles has N eigenvalues s_p , $p = 1, 2, \dots, N$, where the usual ordering is imposed: $s_1 \leq s_2 \leq \dots \leq s_N$. Thus, if the symmetry is broken to a maximal torus, there are $N - 1$ fundamental monopoles and the dimension of the corresponding moduli space is $4(N - 1)$. The Nahm data are defined over the interval $s \in [s_1, s_N]$; this range is subdivided into six subintervals $[s_p, s_{p+1}]$, on each of which the Nahm matrices $T_k(s)$ of dimension $n_p \times n_p$ satisfy Eq. (7) [14]. Thus, each of these subintervals corresponds to a different fundamental monopole, the length of the subinterval defines its mass, and the dimension of the matrices $T_k(s)$ yields the number of monopoles of that type.

The boundary conditions on the endpoint of the subintervals are

- (1) $n_p > n_{p+1}$: $T^{(p+1)}$ should have a well-defined limit at s_{p+1} , and

$$T^{(p)} = \begin{pmatrix} T^{(p+1)}(s_{p+1}) + O(s - s_{p+1}) & O[(s - s_{p+1})^{(n_p - n_{p+1} - 1)/2}] \\ O[(s - s_{p+1})^{(n_p - n_{p+1} - 1)/2}] & -\frac{L^{(p)}}{s - s_{p+1}} + O(1) \end{pmatrix} \quad (10)$$

near the boundary. Here the $n_p \times n_p$ matrix forms an irreducible n_p -dimensional representation of $SU(2)$.

- (2) $n_p < n_{p+1}$: The roles of the left and right end points of the subintervals are reversed, and the residue submatrix $L^{(p)}$ appears in the upper-left corner.
- (3) $n_p = n_{p+1}$: The Nahm data at the endpoint can be discontinuous; one has to introduce the jumping data, $n_p \times 2$ -sized matrix a , and require that at the junction

$$(T_j^{(p+1)} - T_j^{(p)})_{rs} = -\frac{1}{2} a_{s\alpha}^\dagger (\sigma_j)_{\alpha\beta} a_{\beta r}. \quad (11)$$

Here σ_j are the usual Pauli matrices.

III. CONSTRUCTION OF THE G₂ MONOPOLES

Apart from simple embedding of the properly rescaled $SU(2)$ monopole in the 2×2 block of the G_2 matrices, there is another, less trivial embedding into G_2 . Indeed, as \mathfrak{g}_2 algebra possesses $\mathfrak{su}(3)$ subalgebra, it can be decomposed as

$$\mathfrak{g}_2 = \mathfrak{su}(3) \oplus \mathfrak{G}, \quad (12)$$

with \mathfrak{G} forming a module under adjoint action of $\mathfrak{su}(3)$, $[\mathfrak{su}(3), \mathfrak{G}] = \mathfrak{G}$.

This observation leads to a curious consequence regarding zero modes of the $SU(3)$ embedded monopole configuration. Indeed, let us consider the corresponding linearized Bogomolny equation for monopole zero modes

$$\mathcal{D} \delta A = 0. \quad (13)$$

Since \mathcal{D} is $\mathfrak{su}(3)$ valued, these modes clearly separate into purely $\mathfrak{su}(3)$ -valued modes and purely \mathfrak{G} -valued ones. The former are just zero modes of the embedded $SU(3)$ monopole, while the latter appear since G_2 is larger than $SU(3)$. However, we can see that the norm of the Higgs field is not affected by excitation of the \mathfrak{G} -valued zero modes:

$$\delta \frac{1}{2} \text{Tr} \Phi^2 = \text{Tr} \Phi \delta \Phi. \quad (14)$$

By Ward's formula for energy density of the BPS monopoles [27], the excitation of these modes does not change the energy density distribution either. Note that physically these \mathfrak{G} -valued zero modes correspond to the decay of a certain kind of $SU(3)$ monopoles into a pair of different G_2 monopoles.

Let the simple roots of $\mathfrak{su}(3)$ subalgebra be β_1, β_2 . Their corresponding coroots can be decomposed in coroots of G_2 as

²Here we consider the $SU(N)$ model.

$$\beta_1^* = \alpha_1^*, \quad \beta_2^* = \alpha_1^* + \alpha_2^*. \quad (15)$$

Thus, we can set a correspondence between the monopoles as $(n_1, n_2)_{SU(3)} \rightarrow (n_1 + n_2, n_2)_{G_2}$. In other words, the first fundamental $SU(3)$ monopole can be viewed as the first fundamental G_2 monopole, and the second as a stack of both fundamental G_2 monopoles. Such identification is somewhat akin to the construction of the SO, Sp monopoles by restriction of the corresponding $SU(N)$ configurations [11]; however, the identification of some monopole species in this case happens without a reduction in the number of species.

This kind of embedding can be used to obtain some nontrivial configurations. For instance, consider the embedding $(1, 1)_{SU(3)} \rightarrow (2, [1])_{G_2}$ [for the second G_2 monopole to be massless, original $SU(3)$ monopoles should be of equal masses]. The result is the axially symmetric subset of the $(2, [1])_{G_2}$ configurations, i.e. two separated identical monopoles with a cloud of minimal size. We immediately arrive at the conclusion that $(2, [1])_{G_2}$ moduli space interpolates between Taub-NUT (which corresponds to the case of the non-Abelian cloud of minimal size) and Atiyah-Hitchin (the cloud of infinite size) geometries. The same result was obtained earlier by another method in Ref. [15] via the identification of certain species of the $SO(8)$ monopoles.

Axially symmetric $(2, [1])_{SU(3)}$ configurations were studied in detail in Ref. [28]. Such configurations can be of two types. The first one corresponds to the trigonometric axially symmetric Nahm data; it can be considered as the system of two coincident monopoles surrounded by a non-Abelian cloud of finite size. The configuration of the second type corresponds to the hyperbolic axially symmetric Nahm data, in which case the system is composed of two separated monopoles with a non-Abelian cloud of minimal size. By the embedding $(1, 1)_{SU(3)} \rightarrow (2, [1])_{G_2}$, we obtain precisely the latter configuration. Calculation of the energy density profile of the $(1, 1)_{SU(3)}$ embedded monopole then immediately yields the profile of the corresponding axially symmetric $(2, [1])_{G_2}$ configuration.

Apart from this simple embedding, there are different G_2 monopoles which can be constructed directly from the Nahm data. First, let us overview how this formalism can be extended to the classical groups other than $SU(N)$. Since both $SO(N)$ and $Sp(N)$ groups can be represented by unitary matrices with a unit determinant, the corresponding monopole configurations can be obtained by imposing constraints on a general $SU(N)$ solution. In effect, these constraints force some species of $SU(N)$ monopoles to merge, reducing the total number of fundamental monopoles.

Our approach to G_2 monopoles is essentially the same. Making use of the fundamental seven-dimensional representation, we have established the asymptotic behavior (5)

of G_2 monopoles. From a Nahm construction point of view, the leading term of (5) specifies the intervals on which Nahm matrices are defined. The subleading term tells us the number of fundamental $SU(7)$ [or $SO(7)$, since $G_2 \subset SO(7)$] monopoles involved. That is, $(n_1, n_2)_{G_2}$ monopoles lie in the $(n_2, n_1, 2n_2, 2n_2, n_1, n_2)_{SU(7)}$ sector [more precisely, its $(n_2, n_1, n_2)_{SO(7)}$ subsector]. Thus, similarly to the case of an orthogonal group, we need to merge further the $(1, 0, 0)_{SO(7)}$ and $(0, 0, 1)_{SO(7)}$ monopoles to form the $(0, 1)_{G_2}$ monopole.

Note that we can look at the G_2 monopoles from both the $SU(7)$ and $SO(7)$ points of view. The former approach seems to be more natural in the context of Nahm construction; however, the latter approach allows us to deal with a lower number of moduli parameters. Also, G_2 is a subgroup of the group $SO(7)$.

Finally, knowing the intervals on which Nahm matrices reside and their dimensions, we need to place a constraint on the Nahm data directly to merge some monopole species. The transition from $SU(7)$ to $SO(7)$ is well known; the Nahm matrices should possess a reflection symmetry

$$T_j(-s) = C(s)T_j'(s)C^{-1}(s), \quad (16)$$

where the matrix $C(s)$ satisfies $C(-s) = -C^t(s)$. The transition from $SO(7)$ to G_2 , similar to the construction of the $SO(N)$ and $Sp(N)$ monopoles via restriction of the $SU(N)$ Nahm data, should relate the Nahm matrices in the first and the third subintervals [since $(0, 1)_{G_2} \cong (1, 0, 1)_{SO(7)} \cong (1, 0, 2, 2, 0, 1)_{SU(7)}$]. However, the matrices in these intervals are of different sizes; thus, any constraint of the type (16) will not be sufficient.

Some progress can be made if we consider the $(n, 1)_{G_2} \cong (1, n, 1)_{SO(7)}$ sector. There is only one monopole each of the first and of the third kind, and their coordinates enter the Nahm data explicitly (due to reflection symmetry only, we restrict ourselves to $s \leq 0$):

$$T_j(s) = x_j, s \in [-s_1 - s_2, -s_2], \quad (17)$$

$$T_j(s) = I_2 y_j + \dots, s \in [-s_1, 0], \quad (18)$$

where the ellipsis denotes the traceless part, determined by the moduli of the n monopoles of the second kind. Coordinates of the monopoles to be nested are given by x_j and y_j ; it is natural to conjecture that the transition from $SO(7)$ to G_2 is accomplished by setting $x_j = y_j$. This automatically leaves us with a correct number of monopole moduli in the Nahm data.

Let us now see how the construction works for the simplest nontrivial case, $(1, 1)_{G_2}$. The skyline diagram and the corresponding Nahm matrices are given in Fig. 2. For the sake of simplicity, the second monopole is placed at the origin.




FIG. 2. Skyline diagram of $(1,1)_{G_2}$ monopole and its Nahm data.

Here $\sigma'_j = U\sigma_j U^\dagger$ ($U^\dagger = U^{-1}$) are rotated Pauli matrices. The parameters of the rotation and the value s_0 are fixed by the matching condition across the boundaries of the subintervals

$$t_i = -\frac{1}{2s_0}(\sigma'_i)_{22} = \frac{1}{2s_0}(\sigma'_i)_{11}. \quad (19)$$

The Nahm matrices are supplemented by the jumping data

$$\begin{aligned} s = 0: a_{r\alpha} &= \sqrt{\frac{2}{s_0}} U \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \\ s = -s_2: a_\alpha &= \sqrt{2|t_i|} \begin{pmatrix} \sin \theta/2 e^{-i\varphi/2} \\ -\cos \theta/2 e^{i\varphi/2} \end{pmatrix}, \\ s = +s_2: a_\alpha &= \sqrt{2|t_i|} \begin{pmatrix} \cos \theta/2 e^{i\varphi/2} \\ \sin \theta/2 e^{i\varphi/2} \end{pmatrix}, \end{aligned} \quad (20)$$

where θ and φ specify the direction of t_i .

It is a trivial matter to carry out the construction in the $t_i = 0$ case. The two fundamental G_2 monopoles now coincide; they are spherically symmetric. This case corresponds to the $SU(3)$ composite monopole embedded along the root $\beta_3 = \beta_1 + \beta_2$. Then the complete orthonormal set of construction equation solutions can be taken to be

$$\begin{aligned} \omega_1 &= \sqrt{\frac{r}{\sinh vr}} \exp\left(s\sigma_i \cdot r_i - \frac{rs_2}{2}\right) \eta_-^{\text{down}}; \\ \omega_2 &= 0, \quad S(-s_2) = 1; \\ \omega_3 &= \sqrt{\frac{r}{\sinh vr}} \exp\left(s\sigma_i \cdot r_i + \frac{rs_2}{2}\right) \eta_-^{\text{up}}; \\ \omega_4 &= 0, \quad S(0) = 1; \\ \omega_5 &= \sqrt{\frac{r}{\sinh vr}} \exp\left(s\sigma_i \cdot r_i + \frac{rs_2}{2}\right) \eta_+^{\text{down}}; \\ \omega_6 &= 0, \quad S(s_2) = 1; \\ \omega_7 &= \sqrt{\frac{r}{\sinh vr}} \exp\left(s\sigma_i \cdot r_i - \frac{rs_2}{2}\right) \eta_+^{\text{up}}; \end{aligned} \quad (21)$$

where $\eta_\pm^{\text{up/down}}$ are the usual eigenvectors of $\sigma \cdot \mathbf{r}$. These solutions give rise to the Higgs field of the G_2 monopole

$$\begin{aligned} \Phi &= s_2 \text{diag}\left(-\frac{1}{2}, -1, \frac{1}{2}, 0, -\frac{1}{2}, 1, \frac{1}{2}\right) \\ &+ \frac{1}{2} \left[(2s_1 + s_2) \coth(2s_1 + s_2) r - \frac{1}{r} \right] \\ &\times \text{diag}(-1, 0, -1, 0, 1, 0, 1). \end{aligned} \quad (22)$$

One can readily recognize the Higgs profile of a spherically symmetric monopole in the string gauge. We obtain the fields of an embedded $SU(2)$ monopole, just as expected.

For nonzero separation, the construction equation can be solved analytically; however, picking an orthonormal basis of its solutions is a technically difficult task.

In this simple case we can check the correctness of the construction indirectly. The $(1,1)_{G_2}$ solution is obtained by placing a constraint on a generic $(1,1,1)_{SO(7)}$ monopole. Both configurations contain no more than one monopole of each kind. Thus, the corresponding asymptotic metrics, which include monopole coordinates \mathbf{x}_i and phases ξ_i , turn out to be exact. This conclusion can be proven rigorously for two monopoles, since hyper-Kähler structure and asymptotic interaction completely determine the metric on the moduli space. On the other hand, the constraint we imposed selects a submanifold in $(1,1,1)_{SO(7)}$ moduli space (by setting $\mathbf{x}_1 = \mathbf{x}_3$), and hence gives us an expression for the metric of $(1,1)_{G_2}$. Direct computation confirms that the metric obtained by such identification is the correct one.

IV. CONCLUSIONS

The main purpose of this work was to present the application of the Nahm construction to the case of the BPS monopoles in the Yang-Mills-Higgs theory with the exceptional gauge group G_2 spontaneously broken to $U(1) \times U(1)$. As a particular example we considered the Abelian spherically symmetric $(1,1)_{G_2}$ monopole. We have shown that the G_2 monopoles can be constructed by identification of certain set of $SU(7)$ [or $SO(7)$] fundamental monopoles; in particular, the first G_2 fundamental monopole $(1,0)$ represents a set of two nested $SU(7)$ monopoles location and orientation of those coincide, while the second G_2 fundamental monopole $(0,1)$ represents another collection of six aligned and nested $SU(7)$ monopoles.

Perhaps the most interesting feature of the Nahm construction is its realization in the terms of Dirichlet branes. It was pointed out by Diakonessky [17] that there is one-to-one correspondence between the $SU(N)$ monopole embedded along the simple roots as $\mathbf{g} = \frac{4\pi}{e} \sum_i n_i \alpha_i^*$ and the one-branes stretching between the three-branes separated in a transverse direction. This sort of duality has been explicitly realized in $\mathcal{N} = 4$ $SU(N)$ super-Yang-Mills theory [29]. From that point of view, the construction of the Nahm data for G_2 monopoles corresponds to the

configuration of the D-branes, some of which must be identified according to the restrictions (15) [30].

There are various possible applications of the G_2 monopole solutions discussed in this work. An interesting task would be to study the contribution of these configurations in the confinement-deconfinement phase transitions. Note that this transition in the supersymmetric G_2 Yang-Mills theory recently was discussed in Ref. [25]. In particular, it was shown that deconfinement transition does not break the symmetry of the G_2 ground state, although the expectation value of the Wilson line exhibits a discontinuity.

Certainly, this is a first step towards comprehensive study of the monopoles in the gauge models with exceptional groups. As a direction for future work, it would be interesting to study in more detail the moduli space of G_2 monopoles, considering in particular various cases of nonmaximal symmetry breaking. It would allow us to better understand the role of the corresponding massless G_2 monopoles (non-Abelian clouds). Explicit construction of the $(n_1, n_2)_{G_2}$ moduli space metric, which determines the low energy of the monopoles, remains our first goal. We hope to report elsewhere on these problems.

ACKNOWLEDGMENTS

We thank Sasha Gorsky, Derek Harland, Evgeny Ivanov, Olaf Lechtenfeld, Nick Manton, Andrey Smilga and Paul Sutcliffe for many useful discussions and valuable comments. This work is supported in part by the A. von Humboldt Foundation in the framework of the Institutes linkage program and by the JINR Heisenberg-Landau program (Y.S.). We are grateful to the Institute of Physics at the Carl von Ossietzky University Oldenburg for hospitality.

APPENDIX A: \mathfrak{g}_2 ALGEBRA AND ITS REPRESENTATION

Our choice of simple roots is $\alpha_1 = g_{1,-2}$ (long root) and $\alpha_2 = g_{-2}$ (short root); $h_{\alpha_1} = h_1$, $h_{\alpha_2} = h_2 - h_1$. The representation is chosen in such a way that the elements of the Cartan subgroup h with $\alpha_{1,2}(h) \geq 0$ have properly ordered eigenvalues.

$$\begin{aligned}
h_1 &= -e_{22} + e_{33} - e_{55} + e_{66}, \\
h_2 &= -e_{11} - e_{33} + e_{55} + e_{77}, \\
g_{1,-2} &= -e_{32} + e_{65}, \\
g_{1,-3} &= e_{61} - e_{72}, \\
g_{2,-3} &= e_{51} - e_{73}, \\
g_1 &= e_{13} - \sqrt{2}e_{24} + \sqrt{2}e_{46} - e_{57}, \\
g_2 &= -e_{12} - \sqrt{2}e_{34} + \sqrt{2}e_{45} + e_{67}, \\
g_3 &= \sqrt{2}e_{41} + e_{52} - e_{63} - \sqrt{2}e_{74}, \\
g_{2,-1} &= (g_{1,-2})^T, \\
g_{3,-1} &= (g_{1,-3})^T, \\
g_{3,-2} &= (g_{2,-3})^T, \\
g_{-1} &= -(g_1)^T, \\
g_{-2} &= -(g_2)^T, \\
g_{-3} &= -(g_3)^T,
\end{aligned}$$

where e_{nm} is a 7×7 matrix with the only nonzero element $(e_{nm})_{nm} = 1$.

TABLE I. The generators of the \mathfrak{g}_2 algebra.

	h_1	h_2	$g_{1,-2}$	$g_{2,-3}$	$g_{3,-1}$	$g_{2,-1}$	$g_{3,-2}$	$g_{1,-3}$	g_1	g_2	g_3	g_{-1}	g_{-2}	g_{-3}
h_1	0	0	$2g_{1,-2}$	$-g_{2,-3}$	$-g_{3,-1}$	$-2g_{2,-1}$	$g_{3,-2}$	$g_{1,-3}$	$-g_1$	$+g_2$	0	g_{-1}	$-g_{-2}$	0
h_2		0	$-g_{1,-2}$	$2g_{2,-3}$	$-g_{3,-1}$	$g_{2,-1}$	$-2g_{3,-2}$	$g_{1,-3}$	0	$-g_2$	g_3	0	g_{-2}	$-g_{-3}$
$g_{1,-2}$			0	$g_{1,-3}$	$-g_{3,-2}$	h_1	0	0	$-g_2$	0	0	0	g_{-1}	0
$g_{2,-3}$				0	$g_{2,-1}$	0	h_2	0	0	$-g_3$	0	0	0	g_{-2}
$g_{3,-1}$					0	0	0	$-h_{-1} - h_2$	0	0	$-g_1$	g_{-3}	0	0
$g_{2,-1}$						0	$g_{3,-1}$	$-g_{2,-3}$	0	$-g_1$	0	g_{-2}	0	0
$g_{3,-2}$							0	$-g_{1,-2}$	0	0	$-g_2$	0	g_{-3}	0
$g_{1,-3}$								0	$-g_3$	0	0	0	0	$-g_{-1}$
g_1									0	$2g_{-3}$	$-2g_{-2}$	$2h_1 + h_2$	$3g_{2,-1}$	$3g_{3,-1}$
g_2										0	$2g_{-1}$	$3g_{1,-2}$	$-h_1 + h_2$	$3g_{3,-2}$
g_3											0	$3g_{1,-3}$	$3g_{2,-3}$	$-h_1 - 2h_2$
g_{-1}												0	$2g_3$	$-2g_2$
g_{-2}													0	$2g_1$
g_{-3}														0

APPENDIX B: REPRESENTATION OF $\mathfrak{su}(3)$ SUBGROUP

The representation is chosen so that the vacuum expectation value of the Higgs field has properly ordered eigenvalues.

$$\begin{aligned}
 h_1 &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & h_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & g_{2,-1} &= \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & g_{1,-2} &= \begin{pmatrix} 0 & 0 & 0 \\ \frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\
 g_{3,-2} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & 0 & 0 \end{pmatrix}, & g_{2,-3} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & 0 \end{pmatrix}, & g_{3,-1} &= \begin{pmatrix} 0 & 0 & \frac{1}{\sqrt{2}} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & g_{1,-3} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \frac{1}{\sqrt{2}} & 0 & 0 \end{pmatrix}.
 \end{aligned}$$

-
- [1] N. S. Manton and P. Sutcliffe, *Topological Solitons* (Cambridge University Press, Cambridge, England, 2004).
- [2] E. J. Weinberg and P. Yi, Magnetic monopole dynamics, supersymmetry, and duality, *Phys. Rep.* **438**, 65 (2007).
- [3] Y. M. Shnir, *Magnetic Monopoles* (Springer, Berlin, 2005).
- [4] A. S. Schwarz, Magnetic monopoles in gauge theories, *Nucl. Phys.* **B112**, 358 (1976).
- [5] A. N. Leznov and M. V. Saveliev, Exact monopole solutions in gauge theories for an arbitrary semisimple compact group, *Lett. Math. Phys.* **3**, 207 (1979).
- [6] G. 't Hooft, Magnetic monopoles in unified gauge theories, *Nucl. Phys.* **B79**, 276 (1974).
- [7] A. M. Polyakov, Particle spectrum in the quantum field theory, *Pis'ma Zh. Eksp. Teor. Fiz.* **20**, 430 (1974) [*JETP Lett.* **20**, 194 (1974)].
- [8] E. B. Bogomolny, Stability of classical solutions, *Yad. Fiz.* **24**, 861 (1976) [*Sov. J. Nucl. Phys.* **24**, 449 (1976)].
- [9] M. K. Prasad and C. M. Sommerfield, An Exact Classical Solution for The 't Hooft Monopole and The Julia-Zee Dyon, *Phys. Rev. Lett.* **35**, 760 (1975).
- [10] W. Nahm, A simple formalism for the BPS monopole, *Phys. Lett.* **90B**, 413 (1980); in *Monopoles in Quantum Field Theory* edited by (World Scientific, Singapore, 1982).
- [11] J. Hurtubise and M. K. Murray, On the construction of monopoles for the classical groups, *Commun. Math. Phys.* **122**, 35 (1989).
- [12] C. Houghton, P. W. Irwin, and A. J. Mountain, Two monopoles of one type and one of another, *J. High Energy Phys.* **04** (1999) 029.
- [13] P. Irwin, SU(3) monopoles and their fields, *Phys. Rev. D* **56**, 5200 (1997).
- [14] E. J. Weinberg and P. Yi, Explicit multimonopole solutions in SU(N) gauge theory, *Phys. Rev. D* **58**, 046001 (1998).
- [15] K. M. Lee and C. Lu, Two massive and one massless Sp(4) monopoles, *Phys. Rev. D* **57**, 5260 (1998).
- [16] C. J. Houghton and E. J. Weinberg, Multicloud solutions with massless and massive monopoles, *Phys. Rev. D* **66**, 125002 (2002).
- [17] D. E. Diaconescu, D-branes, monopoles, and Nahm equations, *Nucl. Phys.* **B503**, 220 (1997).
- [18] C. Lu, Two monopole systems and the formation of non-Abelian cloud, *Phys. Rev. D* **58**, 125010 (1998).
- [19] B. H. Wellegehausen, A. Wipf, and C. Wozar, Effective Polyakov loop dynamics for finite temperature G(2) gluodynamics, *Phys. Rev. D* **80**, 065028 (2009).
- [20] B. H. Wellegehausen, A. Wipf, and C. Wozar, Casimir scaling and string breaking in G(2) gluodynamics, *Phys. Rev. D* **83**, 016001 (2011).
- [21] K. Landsteiner, J. M. Pierre, and S. B. Giddings, Moduli space of $N = 2$ supersymmetric G(2) gauge theory, *Phys. Rev. D* **55**, 2367 (1997).
- [22] G. Cossu, M. D'Elia, A. Di Giacomo, B. Lucini, and C. Pica, G(2) gauge theory at finite temperature, *J. High Energy Phys.* **10** (2007) 100.
- [23] E. M. Ilgenfritz and A. Maas, Topological aspects of G(2) Yang-Mills theory, *Phys. Rev. D* **86**, 114508 (2012).
- [24] M. Gunaydin and H. Nicolai, Seven-dimensional octonionic Yang-Mills instanton and its extension to an heterotic string soliton, *Phys. Lett. B* **351**, 169 (1995); **376**, 329 (E) (1996).
- [25] E. Poppitz, T. Schäfer, and M. Ünsal, Universal mechanism of (semiclassical) deconfinement and theta-dependence for all simple groups, *J. High Energy Phys.* **03** (2013) 087.
- [26] K. M. Lee, E. J. Weinberg, and P. Yi, Massive and massless monopoles with non-Abelian magnetic charges, *Phys. Rev. D* **54**, 6351 (1996).
- [27] R. S. Ward, A Yang-Mills-Higgs monopole of charge 2, *Commun. Math. Phys.* **79**, 317 (1981).
- [28] A. S. Dancer, Nahm data and SU(3) monopoles, *Non-linearity* **5**, 1355 (1992).
- [29] A. Hanany and E. Witten, Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics, *Nucl. Phys.* **B492**, 152 (1997).
- [30] K. G. Selivanov and A. V. Smilga, Effective Lagrangian for 3D $N = 4$ SYM theories for any gauge group and monopole moduli spaces, *J. High Energy Phys.* **12** (2003) 027.