

Resummation effects in the forward production of $Z_0 + \text{jet}$ at the LHC

A. van Hameren,¹ P. Kotko,² and K. Kutak¹

¹*The H. Niewodniczański Institute of Nuclear Physics PAN, Radzikowskiego 152, 31-342 Kraków, Poland*

²*Department of Physics, The Pennsylvania State University, University Park, Pennsylvania 16802, USA*

(Received 21 May 2015; published 8 September 2015)

We calculate several differential cross sections for Z_0 and high- p_T jet production in the forward rapidity region at the LHC using the hybrid high energy factorization. We test various unintegrated gluon distributions involving subleading Balitski-Fadin-Kuraev-Lipatov effects (such as kinematic constraint, running strong coupling and Dokshitzer-Gribov-Lipatov-Altarelli-Parisi correction) and compare the results with experimental data obtained by the LHCb experiment. We find that the hard scale dependence of unintegrated gluon distributions, which effectively resums the Sudakov-type logarithms on the top of the resummation of the small x logarithms, is essential to describe the normalized azimuthal decorrelations between the Z_0 -boson and the jet.

DOI: 10.1103/PhysRevD.92.054007

PACS numbers: 13.85.Hd

I. MOTIVATION

The Large Hadron Collider opens an opportunity to explore kinematic regions where particles produced in high-energy collisions possess large transverse momenta and rapidities. The production of electroweak bosons and jets is a vital part of tests of the standard model as well as searches of physics beyond the standard model. Furthermore it has been recognized in [1] that studies of associated production of electroweak bosons and jets may provide a new insight into the transverse partonic structure of hadrons at small x , where x is the momentum fraction of the hadron taken by a parton participating in the hard collision. Furthermore, such a final state, being a combination of colorful and colorless particles, gives the opportunity for particularly interesting investigations complementary to results obtained in studies of pure jet final states [2–6] and Drell-Yan pairs [7]. In particular the final state rescatterings due to soft color exchanges should have less impact on the properties of the produced final state as compared to pure jet final states.

This work is motivated by a recent LHCb measurement [8] at $\sqrt{s} = 7$ TeV of the process

$$pp \rightarrow Z_0(\rightarrow \mu^+ \mu^-) + \text{jet} \quad (1)$$

in the forward direction within the pseudorapidity range $2.0 < \eta < 4.5$. The final-state muon and antimuon were required to have transverse momenta $p_{T\mu} > 20$ GeV while the leading jet was considered with two different cuts: $p_{Tj} > 10$ GeV and $p_{Tj} > 20$ GeV. The jets were reconstructed using the anti- k_T algorithm with radius $R = 0.5$ and they were required to be separated from muon tracks on the $\phi - \eta$ plane (azimuthal angle-pseudorapidity) by a distance $R = 0.4$. The muon-pair was required to have an invariant mass within the range $60 \text{ GeV} < M_{\mu\mu} < 120 \text{ GeV}$. The rapidity constraint assures that in the

partonic picture of the process, one of the initial state partons carries a rather small fraction x_A of the corresponding hadron momentum p_A , while the other must have a fraction $x_B \gg x_A$ (cf. Fig. 1).

In order to describe the process perturbatively, one definitely needs to go beyond the pure collinear factorization and support the calculation by a resummation. In the modern advanced approaches this is achieved by parton showers and hadronization as implemented for example in pythia [9]. In the present work we consider another approach, namely a resummation of logarithms of $\ln(1/x_A)$ and $\ln(\mu/k_T)$, where μ is a hard scale and k_T is a certain additional scale given by the imbalance of the final states on the transverse plane. This approach captures certain aspects of the process more accurately already at lowest order of the strong coupling constant in the hard process.

In the present paper, we will therefore attempt to study the process within so called high energy factorization, or




FIG. 1. Diagrammatic representation of the hybrid high energy factorization for forward $Z_0 + \text{jet}$ production. The upper blob corresponds to a collinear PDF, whereas the lower one corresponds to the unintegrated gluon distribution. The gluon entering the hard scattering is off-shell with virtuality k_T^2 .

more precisely, using so-called hybrid high energy factorization motivated by the works [10–13]. Within the asymmetric kinematic situation $x_B \gg x_A$ described above, the cross section for the process under consideration can be expressed by the following formula

$$\begin{aligned}
& d\sigma_{AB \rightarrow \mu^+ \mu^- + \text{jet} + X} \\
&= \int d^2 k_{TA} \int \frac{dx_A}{x_A} \int dx_B \sum_b \\
&\quad \times \mathcal{F}_{g^*/A}(x_A, k_{TA}, \mu) f_b(x_B, \mu) d\hat{\sigma}_{g^* q_b \rightarrow q_b \mu^+ \mu^-}(x_A, x_B, k_{TA}, \mu),
\end{aligned} \tag{2}$$

where $\mathcal{F}_{g^*/A}$ is the unintegrated gluon distribution for hadron A , f_b is a collinear PDF and $d\hat{\sigma}_{g^* q_b \rightarrow q_b \mu^+ \mu^-}$ is the hard cross section obtained from a gauge invariant tree-level off-shell amplitude for the process $g^* q_b \rightarrow q_b \mu^+ \mu^-$, and where q_b refers to quarks as well as antiquarks (Fig. 1).

Let us note, that the original high energy factorization prescription was designed to study inclusive small x processes and the corresponding unintegrated gluon distribution was assumed to undergo the Balitski-Fadin-Kuraev-Lipatov (BFKL) evolution equation (see e.g. [14]). For more exclusive processes it is however necessary to include the subleading BFKL effects, such




FIG. 2 (color online). Azimuthal decorrelations for $p_{Tj} > 10$ GeV normalized to the total cross section.

as kinematic constraint ensuring energy conservation, large- x correction, running strong coupling constant, and—notably—the hard scale dependence. The last, denoted in Eq. (2) as μ , turns out to be essential to describe the data under consideration. In the present work we shall not discuss the validity of the model (2) on the theoretical level. Instead, we shall test it phenomenologically against the existing data. For a more detailed review of various approaches to the small x factorization and related issues see, e.g., [15]. For a derivation of hybrid high energy factorization from the dilute limit of the color glass condensate approach see [16].

II. RESULTS

Using the formalism described in the preceding section we have computed the cross sections for $Z_0 + \text{jet}$ production. We have used two programs to calculate off-shell $qg^* \rightarrow q\mu^+\mu^-$ amplitude (with Z_0 and γ exchange) and to cross-check the results. The first program is A VERY HANDY LIBRARY (AVHLIB) [17] in FORTRAN, in which the approach of [18] is implemented. It computes amplitudes entirely numerically and includes a full Monte Carlo program. The second is the electroweak extension of the program OGIME [19]. It calculates amplitudes analytically in a form that can be interfaced with a Monte Carlo




FIG. 3 (color online). Azimuthal decorrelations for $p_{Tj} > 20$ GeV normalized to the total cross section.

program. More specifically, the analytic expressions were implemented in the C++ code `LXJet` [20]. Note, however, since there are no final state gluons, the ordinary Feynman diagram depicted in Fig. 1 (with appropriate high energy projector for the off-shell gluon as described in [10]) is enough to obtain the gauge invariant amplitude for this process, and all the complications discussed in [18,21–24] do not have any impact here.

For the numerical computation we use the kinematic cuts as described in the previous section, i.e. the ones used in [8]. We use standard values for the electroweak parameters: electroweak coupling $g_{ew} = 0.308$, Z_0 boson mass

$M_Z = 91.2$ GeV and width $\Gamma_Z = 2.495$ GeV. We work with the 5 flavor scheme and use the CTEQ10NLO set [25] for the collinear PDFs. For the unintegrated gluon distributions we consider the following models (in brackets we give our abbreviations for the models):

- (i) BFKL evolution with kinematic constraint, running strong coupling constant, DGLAP effects including the contribution from the quark sea [26] and supplemented with the nonlinear term [27]; the initial condition has been fitted to HERA data [28] (KS)
- (ii) hard scale dependent KS; the hard scale dependence is achieved by implementing the Sudakov form




FIG. 4 (color online). Transverse momentum spectrum for Z_0 -boson for $p_{Tj} > 10$ GeV normalized to the total cross section.

- factor on the generated events in such a way that the total cross section remains unchanged [5] (KS+Sudakov)
- (iii) similar to above, but the Sudakov form factor is implemented to the KS density in such a way that the integrated gluon density remains unchanged [29] (Kutak-nonlinear-PRD15)
- (iv) the simplified Kimber-Martin-Ryskin model [30] applied to MRSTW08 PDFs; this model gives unintegrated gluon distribution that is hard scale dependent (KMR)
- (v) BFKL evolution with kinematic constraint, running strong coupling constant and the DGLAP

contribution coming only from gluons and fitted to the LHC jet data [15] (LHC-fit)

- (vi) hard scale dependent LHC fit (LHC-fit+Sudakov).

The hard scale was chosen to be the average of the three large scales appearing in the problem: the mass of the Z_0 boson, its p_T and the p_T of the jet. We present the results for numerical simulations in Figs. 2–10 and compare them with LHCb data. Figures 2, 3 present normalized azimuthal decorrelations for $p_{Tj} > 10$ GeV and $p_{Tj} > 20$ GeV, respectively (the azimuthal decorrelation is defined as the differential cross section in the difference of azimuthal angles of the reconstructed Z_0




FIG. 5 (color online). Transverse momentum spectrum for Z_0 -boson for $p_{Tj} > 20$ GeV normalized to the total cross section.

boson and the leading jet). We present also results for normalized differential cross sections in the transverse momentum p_{TZ} of Z_0 boson (Figs. 4, 5), in rapidity y_Z of Z_0 (Figs. 6, 7), in the transverse momentum p_{Tj} of the jet (Fig. 10), and—finally—in the rapidity separation between the Z_0 -boson and the jet Δy (Figs. 8, 9). The shaded boxes in Figs. 2–10 represent the theoretical uncertainty obtained by varying the hard scale by a factor of two. The differential cross sections are normalized to the total cross section, as in [8]. We list the total cross sections for different models in Table I.

III. DISCUSSION AND CONCLUSIONS

Let us first discuss the normalized differential cross sections. We can conclude that the azimuthal decorrelations are described reasonably well for both jet p_T cuts for most of the models. It was however essential for this observable to include the hard scale dependence in unintegrated gluon distributions. As this observable is the most sensitive one for the small x effects this underlines the importance of the resummation of the logarithms $\ln(\mu/k_T)$, where μ is the hard scale provided by the hard process and k_T is the transverse momentum of the gluon in unintegrated gluon distributions, on the top of the small x logarithms. The




FIG. 6 (color online). Rapidity spectrum of Z_0 -boson for $p_{Tj} > 10$ GeV normalized to the total cross section.

effect of this Sudakov-type resummation was also important for the transverse momentum spectrum of Z_0 boson. For the other observables it has had much less impact.

Since our calculations are not interfaced with any sort of final state parton shower we expected a rather rough description of transverse momenta spectra. Our actual study shows however that the situation is relatively good for the spectrum of Z_0 boson (except very low transverse momentum), Figs. 4–5, and indeed fails for the jet spectrum, Fig. 10. This can be attributed to the fact that all unintegrated gluon distributions we have used contain contribution from pieces of splitting functions

subleading at low x (see [15] for an analysis of the impact of this correction on jet p_T spectra). This correction seems to be enough for the colorless final state while the color rescattering for the final state jet is evidently missing. Also, the next-to-leading correction in the hard process is necessary to improve the description of the transverse momentum spectra. The LHC-jet-motivated unintegrated gluon densities, which were actually fitted to the LHC jet transverse momentum spectra behave similar to the other gluon densities. This gives one more clue that the improvement in the hard process in terms of higher order corrections or/and a resummation is necessary.




FIG. 7 (color online). Rapidity spectrum of Z_0 -boson for $p_{Tj} > 20$ GeV normalized to the total cross section.

For $p_{Tj} > 20$ GeV the *Kutak-nonlinear-PRD15* and the *KMR* models overestimate the normalized transverse momentum spectrum of the Z_0 boson. These models implement a hard scale dependence but the spectra they give are almost the same as from the *KS* model, which is hard-scale-independent (the difference is however in the uncertainty which is much bigger for the hard-scale-dependent models). This means that the Sudakov-type resummation in the former models has no effect for this observable with such large p_T cut. On the contrary, the Sudakov-type resummation with unitarity constraint (in the sense of preserving the total cross section) applied to the *KS* or *LHC-fit* improves the results.

Interestingly, the unintegrated gluon distribution which comes from the fit to the LHC jet data does not perform better than the distributions obtained from the inclusive deep inelastic scattering data. This suggests that the effects of factorization breaking due to the lack of universality is rather weak at the phenomenological level. As already mentioned above, this suggests also that one needs higher order corrections in the hard process. Indeed, as observed in [15] the p_T spectra cannot be described by improving the evolution of unintegrated gluon distribution itself. One should mention, that the evolution equations that were fitted to the LHC data did not include a contribution from the sea quarks on the dense hadron side, whereas this




FIG. 8 (color online). Differential cross section as a function of rapidity separation between the Z_0 -boson and the jet for $p_{Tj} > 10$ GeV normalized to the total cross section.

contribution is present in the other approaches. It is however unlikely that this can improve the situation.

Present calculations did not take into account the situation where the off-shell initial-state parton is a quark or an antiquark. We expect this contribution to be small, as at small x gluons dominate significantly. On the other hand, the process $\bar{q}q \rightarrow Z_0(\rightarrow \mu^+\mu^-) + g$ gives an important contribution in the collinear approach. Since in reality the probed values of x are not extremely small, such process might be important in the high energy factorization. Practical applications require however a set of unintegrated quark distributions, consistent with the

unintegrated gluon distributions. Inclusion of those is left for future studies.

As seen from the Table I our calculations strongly underestimate the total cross section as compared to the data. This should be probably attributed to the hard multiparton interaction (MPI) effects which are not taken into account in the hybrid high energy factorization. This statement is supported by the observation that the description of the azimuthal decorrelations is very good up to the normalization, which, when corrected, will only shift the data eventually forming a “pedestal.” It is also known that MPIs indeed contribute only a pedestal to the azimuthal




FIG. 9 (color online). Differential cross section as a function of rapidity separation between the Z_0 -boson and the jet for $p_{Tj} > 20$ GeV normalized to the total cross section.




FIG. 10 (color online). Transverse momentum spectrum of the jet normalized to the total cross section.

TABLE I. Total cross sections obtained from different models for the unintegrated gluon density. The model uncertainties are defined through variations of the hard scale. For the data the total uncertainty is estimated as an average square error from statistical, systematic and luminosity errors as given in [8]. We do not include models *KS+Sudakov* and *LHC-fit+Sudakov* as the Sudakov-type resummation applied there does not change the total cross section.

Unintegrated gluon distribution model	$p_{Tj} > 10$ GeV	$p_{Tj} > 20$ GeV
KS	$4.1^{+1.0}_{-0.7}$ pb	$2.3^{+0.6}_{-0.4}$ pb
Kutak-nonlinear-PRD15	$4.1^{+1.0}_{-1.0}$ pb	$2.4^{+0.6}_{-0.5}$ pb
KMR	$5.8^{+1.5}_{-1.3}$ pb	$3.3^{+0.9}_{-0.6}$ pb
LHC-fit	$3.4^{+0.8}_{-0.6}$ pb	$2.0^{+0.5}_{-0.4}$ pb
LHCb data	16.0 ± 1.4 pb	6.3 ± 0.5 pb

decorrelations (see e.g. [31,32]), as the two partons coming from independent scatterings are completely decorrelated (to leading order) and thus the azimuthal decorrelation distribution is flat. The subject of MPIs in the high energy factorization is however rather complicated and needs a separate study. In principle, on the dense hadron side (i.e. for the one probed at small x), the soft MPIs can be partially taken into account by means of the nonlinear term in the evolution equation, as for example in the KS unintegrated gluon density or its extensions. Therefore, one has to be careful not to make a double counting when using some phenomenological models by an inclusion of double hard scattering mechanism. Actually the problem extends beyond the saturation regime as the high energy factorization contributes to higher twists as well as MPIs, at least

for certain observables [33]. These subjects are however beyond the scope of the present work.

ACKNOWLEDGMENTS

We acknowledge useful correspondence and discussions with Albert Bursche, and we thank Rafał Maciuła for

providing us the necessary grid for the KMR pdf. The work of K. K. has been supported by Narodowe Centrum Nauki with Sonata Bis Grant No. DEC-2013/10/E/ST2/00656. P. K. acknowledges the support of the US Department of Energy Grants No. DE-SC-0002145 and No. DE-FG02-93ER40771.

-
- [1] S. Dooling, F. Hautmann, and H. Jung, Hadroproduction of electroweak gauge boson plus jets and TMD parton density functions, *Phys. Lett. B* **736**, 293 (2014).
- [2] M. Deak, F. Hautmann, H. Jung, and K. Kutak, Forward-central jet correlations at the large hadron collider, [arXiv:1012.6037](#).
- [3] A. van Hameren, P. Kotko, and K. Kutak, Three jet production and gluon saturation effects in p-p and p-Pb collisions within high-energy factorization, *Phys. Rev. D* **88**, 094001 (2013).
- [4] M. Nefedov, V. Saleev, and A. V. Shipilova, Dijet azimuthal decorrelations at the LHC in the parton Reggeization approach, *Phys. Rev. D* **87**, 094030 (2013).
- [5] A. van Hameren, P. Kotko, K. Kutak, and S. Sapeta, Small-x dynamics in forward-central dijet decorrelations at the LHC, *Phys. Lett. B* **737**, 335 (2014).
- [6] A. van Hameren, P. Kotko, K. Kutak, C. Marquet, and S. Sapeta, Saturation effects in forward-forward dijet production in p + Pb collisions, *Phys. Rev. D* **89**, 094014 (2014).
- [7] A. Lipatov, M. Malyshev, and N. Zotov, Drell-Yan lepton pair production at high energies in the k_T -factorization approach, *J. High Energy Phys.* **12** (2011) 117.
- [8] R. Aaij *et al.* (LHCb Collaboration), Study of forward Z + jet production in pp collisions at $\sqrt{s} = 7$ TeV, *J. High Energy Phys.* **01** (2014) 033.
- [9] T. Sjostrand, S. Mrenna, and P.Z. Skands, PYTHIA 6.4 physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [10] S. Catani, M. Ciafaloni, and F. Hautmann, High-energy factorization and small x heavy flavor production, *Nucl. Phys.* **B366**, 135 (1991).
- [11] S. Catani, M. Ciafaloni, and F. Hautmann, Gluon contributions to small x heavy flavor production, *Phys. Lett. B* **242**, 97 (1990).
- [12] S. Catani and F. Hautmann, High-energy factorization and small x deep inelastic scattering beyond leading order, *Nucl. Phys.* **B427**, 475 (1994).
- [13] M. Deak, F. Hautmann, H. Jung, and K. Kutak, Forward jet production at the Large Hadron Collider, *J. High Energy Phys.* **09** (2009) 121.
- [14] L. Lipatov, Small x physics in perturbative QCD, *Phys. Rep.* **286**, 131 (1997).
- [15] P. Kotko, W. Slominski, and D. Toton, Unintegrated gluon distributions for forward jets at LHC, [arXiv:1504.00823](#).
- [16] P. Kotko, K. Kutak, C. Marquet, E. Petreska, S. Sapeta *et al.*, Improved TMD factorization for forward dijet production in dilute-dense hadronic collisions, [arXiv:1503.03421](#).
- [17] M. Bury and A. van Hameren, Numerical evaluation of multi-gluon amplitudes for High Energy Factorization, [arXiv:1503.08612](#).
- [18] A. van Hameren, P. Kotko, and K. Kutak, Helicity amplitudes for high-energy scattering, *J. High Energy Phys.* **01** (2013) 078.
- [19] P. Kotko, OGIME—Off-shell Gauge Invariant Matrix Elements, FORM program, 2013.
- [20] P. Kotko, LxJet, C++ Monte Carlo program, 2013.
- [21] A. van Hameren, P. Kotko, and K. Kutak, Multi-gluon helicity amplitudes with one off-shell leg within high energy factorization, *J. High Energy Phys.* **12** (2012) 029.
- [22] A. van Hameren, K. Kutak, and T. Salwa, Scattering amplitudes with off-shell quarks, *Phys. Lett. B* **727**, 226 (2013).
- [23] P. Kotko, Wilson lines and gauge invariant off-shell amplitudes, *J. High Energy Phys.* **07** (2014) 128.
- [24] A. van Hameren, BCFW recursion for off-shell gluons, *J. High Energy Phys.* **07** (2014) 138.
- [25] H.-L. Lai, M. Guzzi, J. Huston, Z. Li, P. M. Nadolsky, J. Pumplin, and C.-P. Yuan, New parton distributions for collider physics, *Phys. Rev. D* **82**, 074024 (2010).
- [26] J. Kwiecinski, A. D. Martin, and A. Stasto, A unified BFKL and GLAP description of F2 data, *Phys. Rev. D* **56**, 3991 (1997).
- [27] K. Kutak and A. Stasto, Unintegrated gluon distribution from modified BK equation, *Eur. Phys. J. C* **41**, 343 (2005).
- [28] K. Kutak and S. Sapeta, Gluon saturation in dijet production in p-Pb collisions at Large Hadron Collider, *Phys. Rev. D* **86**, 094043 (2012).
- [29] K. Kutak, Hard scale dependent gluon density, saturation and forward-forward dijet production at the LHC, *Phys. Rev. D* **91**, 034021 (2015).
- [30] M. Kimber, A. D. Martin, and M. Ryskin, Unintegrated parton distributions, *Phys. Rev. D* **63**, 114027 (2001).
- [31] M. Strikman and W. Vogelsang, Multiple parton interactions and forward double pion production in pp and dA scattering, *Phys. Rev. D* **83**, 034029 (2011).
- [32] A. Stasto, B.-W. Xiao, and F. Yuan, Back-to-back correlations of di-hadrons in dAu collisions at RHIC, *Phys. Lett. B* **716**, 430 (2012).
- [33] M. Diehl, D. Ostermeier, and A. Schafer, Elements of a theory for multiparton interactions in QCD, *J. High Energy Phys.* **03** (2012) 089.