

Addendum to “Electroweak Sudakov effects in W , Z and γ production at large transverse momentum”

Thomas Becher and Xavier Garcia i Tormo

*Albert Einstein Center for Fundamental Physics, Institut für Theoretische Physik, Universität Bern,
Sidlerstrasse 5, CH-3012 Bern, Switzerland*

(Received 10 September 2015; published 19 October 2015)

In this addendum to T. Becher and X. Garcia i Tormo, Phys. Rev. D **88**, 013009 (2013), we give results for the electroweak Sudakov corrections in gauge-boson production at large transverse momentum p_T at proton colliders. For the results to be easily usable, we provide a simple and accurate parametrization of the corrections as a function of p_T and the center-of-mass energy \sqrt{s} . Additionally, we also discuss the dependence of the electroweak corrections on the rapidity of the produced boson and comment on the complications that arise in the photon-production case due to isolation requirements.

DOI: [10.1103/PhysRevD.92.073011](https://doi.org/10.1103/PhysRevD.92.073011)

PACS numbers: 12.38.Bx, 14.70.Fm, 14.70.Hp

In Ref. [1] we presented a study of electroweak Sudakov corrections to the cross sections for single W , Z , and γ production at large transverse momentum p_T , within the framework of Soft Collinear Effective Theory. Our results are based on a factorization formula for the hadronic cross section near the partonic threshold and include both QCD and electroweak corrections. To quantify the effect of including electroweak corrections in the cross section, it is convenient to look at the relative difference $\Delta\sigma^{ew}$ of the cross section computed with and without electroweak effects, i.e.

$$\Delta\sigma^{ew} := \frac{\sigma_{ew}^i - \sigma^i}{\sigma^i}, \quad (1)$$

where σ^i represents the cross section computed including only QCD effects, at a certain order denoted by i , and σ_{ew}^i represents the same cross section including electroweak effects. As shown in Ref. [1], the relative importance of the electroweak corrections, given by $\Delta\sigma^{ew}$, is essentially independent of the order i which we use to compute the QCD effects. This implies that, to good accuracy, one can include electroweak effects in existing pure-QCD computations simply as an overall multiplicative prefactor $1 + \Delta\sigma^{ew}$.

The inclusion of electroweak Sudakov effects is essential in order to obtain precise predictions for the p_T spectrum in the region $p_T \gg M_{W,Z}$, since their effect can be as large as 20% for $p_T \sim 1$ TeV. Therefore, any LHC analysis of these processes, as well as studies of future colliders at higher center-of-mass energies, must include them. In Ref. [1] we presented the correction factor $\Delta\sigma^{ew}$ for a center-of-mass energy $\sqrt{s} = 7$ TeV. Results at $\sqrt{s} = 33$ TeV and $\sqrt{s} = 100$ TeV were presented in Ref. [2], in the context of the Snowmass study of future colliders. It was found that the relative correction $\Delta\sigma^{ew}$ depends only mildly on the center-of-mass energy. For further Run I and future Run II LHC analyses, it would be useful to have an expression that allows one to obtain $\Delta\sigma^{ew}$ for different values of \sqrt{s} .

Providing such an expression is the main motivation for this addendum. To obtain it, we compared the correction factors $\Delta\sigma^{ew}$ at different \sqrt{s} values and found that their ratios are largely independent of p_T . For this reason, we first provide a simple parametrization of the results at $\sqrt{s} = 7$ TeV, which serves as a reference, and then supply the necessary prefactor to convert them to other center-of-mass energies.

The dependence on p_T of $\Delta\sigma^{ew}$ at our reference scale $\sqrt{s_0} = 7$ TeV can be very accurately parametrized by a third-order polynomial, according to

$$\Delta\sigma_V^{ew}(p_T, s_0) = a_0^V + a_1^V x + a_2^V x^2 + a_3^V x^3 =: g_V(p_T), \quad (2)$$

where x represents the p_T of boson V expressed in TeV and the a_i^V coefficients for the different gauge bosons are given in Table I, both for the central value and the associated error band. The accuracy of the parametrization in Eq. (2) is better than 1% for most of the p_T values between 100 and 1500 GeV, and the difference between the parametrization and the exact result is always much smaller than the theoretical uncertainties of $\Delta\sigma^{ew}$. Equation (2) can therefore be safely used in the whole p_T range we study but should not be employed beyond $p_T = 1500$ GeV. To illustrate this, we show in Fig. 1 the exact result for $\Delta\sigma^{ew}$ (black points) together with the outcome of Eq. (2) (red line), as well as the theoretical uncertainty due to scale variations (gray band).

Next, we provide prefactors $f_V(\sqrt{s})$ that convert Eq. (2) to other center-of-mass energies, according to

$$\Delta\sigma_V^{ew}(p_T, s) = g_V(p_T)/f_V(\sqrt{s}), \quad (3)$$

where $\Delta\sigma_V^{ew}(p_T, s)$ represents the difference $\Delta\sigma_V^{ew}$ at a center-of-mass energy \sqrt{s} , expressed in TeV. The $f_V(\sqrt{s})$, for each of the different bosons, are given by

$$f_Z(\sqrt{s}) = 1 - 1.0376 \times 10^{-2} y + 5.4060 \times 10^{-4} y^2, \quad (4)$$

TABLE I. Values of the a_i^V coefficients in Eq. (2), parametrizing the p_T dependence of $\Delta\sigma_V^{ew}$ at $\sqrt{s} = 7$ TeV, for the different gauge bosons V .

	Central value			
	a_0^V	a_1^V	a_2^V	a_3^V
Z	2.763	-40.76	21.95	-5.356
γ	1.713	-21.68	12.16	-3.050
W^+	3.816	-45.25	23.74	-5.899
W^-	4.074	-47.60	25.97	-6.414
	Upper edge			
	a_0^V	a_1^V	a_2^V	a_3^V
Z	7.640	-42.41	22.88	-5.609
γ	4.556	-22.06	12.30	-3.072
W^+	9.592	-46.81	24.09	-5.976
W^-	9.906	-49.00	26.63	-6.590
	Lower edge			
	a_0^V	a_1^V	a_2^V	a_3^V
Z	2.302	-41.49	22.53	-5.504
γ	0.824	-23.23	13.44	-3.428
W^+	2.307	-45.99	24.97	-6.244
W^-	2.622	-48.51	26.84	-6.637

$$f_\gamma(\sqrt{s}) = 1 - 2.3355 \times 10^{-2}y + 1.2310 \times 10^{-3}y^2, \quad (5)$$

$$f_{W^-}(\sqrt{s}) = 1 - 9.5067 \times 10^{-3}y + 4.9809 \times 10^{-4}y^2, \quad (6)$$

$$f_{W^+}(\sqrt{s}) = 1 - 1.9924 \times 10^{-2}y + 1.0891 \times 10^{-3}y^2 + p_T(1.2802 \times 10^{-2}y - 7.6955 \times 10^{-4}y^2), \quad (7)$$




FIG. 1 (color online). $\Delta\sigma^{ew}$ for Z production at $\sqrt{s} = 7$ TeV computed using the exact expressions from Ref. [1], black points, and the parametrization in Eq. (2), red line. For reference, we also show the theoretical uncertainty of the result as the gray band (see Ref. [1] for further details).

where $y = (\sqrt{s} - \sqrt{s_0})/\sqrt{s_0}$. Only in the case of W^+ , one observes a slight dependence of the ratio on p_T , and for this reason we decided to include also a term linear in p_T in f_{W^+} ; it is understood that p_T is expressed in TeV. Using the f_V factors to obtain $\Delta\sigma^{ew}$ at different energies provides an accuracy better than 1% for most of the p_T values between 100 and 1500 GeV, and the difference with the exact result is always much smaller than the theoretical uncertainties. We illustrate this in Fig. 2, by comparing ratios of $\Delta\sigma^{ew}$ computed at different energies using the exact result (black points) with the f_V factors in Eqs. (4)–(7) (red line). One can also use the f_V factors above to convert the upper and lower limits of the uncertainty bands at 7 TeV to other \sqrt{s} values.

Our finding that $\Delta\sigma^{ew}$ given above has only a mild dependence on \sqrt{s} is a reflection of the fact that most of the electroweak corrections arise in the hard function in the factorization formula, as discussed in Ref. [1]. The hard function evolves from its characteristic scale, which is of order p_T , to the factorization scale, which is of order $M_{Z,W}$, and the electroweak corrections generated by this evolution are independent of \sqrt{s} . We note that the predictions of our effective-theory setup become less accurate as $p_T \rightarrow M_{Z,W}$, since the hierarchy of scales on which it is based is no longer present. In particular we cannot use it to compute electroweak corrections below $p_T = M_{Z,W}$, and, consequently, one should just set our $\Delta\sigma^{ew}$ factor to zero for $p_T \lesssim M_{Z,W}$. Note also that we estimated the uncertainties of our results by varying the different scales appearing in the factorization formula independently. When $p_T \rightarrow M_{Z,W}$ the different scales are not really independent, and this procedure may result in an overestimate of the errors, but since the method itself deteriorates in this limit, we recommend using the error bands given above all the way down to $p_T = M_{Z,W}$.

Our approximate results are for cross sections integrated over the rapidity of the bosons, but to good accuracy the corrections can also be used for cross sections with rapidity cuts. To understand this, consider Fig. 3, which shows the dependence of the corrections on rapidity for a few fixed p_T values. The left panel of the figure shows the cross section differential in rapidity, normalized to $N = d\sigma/dy|_{y=0}$, the value of the cross section at $y = 0$. The right panel shows the size of the electroweak corrections $\Delta\sigma^{ew}(y)$ as a function of rapidity, normalized to $\Delta\sigma^{ew}$ integrated over rapidity. In the horizontal axis, the rapidity is normalized to the maximum value y_{\max} , which is given by

$$y_{\max} = -\ln \frac{s + M_V^2 - \sqrt{(M_V^2 - s)^2 - 4p_T^2 s}}{2\sqrt{s}\sqrt{p_T^2 + M_V^2}} \approx \frac{1}{2} \ln \frac{s}{p_T^2}, \quad (8)$$

where M_V is the mass of the produced boson. Only positive rapidities are shown because the cross section is symmetric in rapidity for pp collisions. From the figure we learn that, first of all, the bulk of the cross section comes from the central rapidity region, $y/y_{\max} \lesssim 0.8$ and, second, that the




FIG. 2 (color online). Ratios of $\Delta\sigma^{ew}$ at different energies, for a pp collider. Left: $\sqrt{s} = 7$ TeV over $\sqrt{s} = 33$ TeV for W^+ . Right: $\sqrt{s} = 7$ TeV over $\sqrt{s} = 33$ TeV for Z . The black points are always the exact result using the expressions from Ref. [1], and the red lines are the f_V factors in Eqs. (4)–(7).




FIG. 3 (color online). Left panel: Cross section differential in rapidity, normalized to the value at $y = 0$. Right panel: Size of the electroweak corrections as a function of rapidity, normalized to the value of the correction integrated over rapidity.

electroweak corrections are essentially flat in this region. Therefore, $\Delta\sigma^{ew}$ is mostly independent of rapidity. Furthermore, the deviations from unity in Fig. 3 mostly arise for bosons at low p_T and therefore correspond to very high rapidities (e.g. at $p_T = 300$ GeV and $\sqrt{s} = 8$ TeV, $y_{\max} \approx 3.3$) which are typically not included in the experimental measurements. The fact that the corrections are largely independent of the vector-boson rapidity is demonstrated in Fig. 4, where we have computed them for different rapidity ranges relevant in the ATLAS experiment and find that they are almost identical in every case.

As a final, important point, we briefly want to discuss the effect of experimental cuts beyond the boson rapidity. Our computation is based on threshold resummation; i.e. it assumes that the final state only contains the hard vector boson, recoiling against a low-mass jet, plus soft gluons and soft photons. In experimental measurements a variety of phase-space cuts are imposed, and while we hope to

capture the bulk of the electroweak (EW) corrections, it is clear that the setup of our calculation is not fully realistic. In particular, the EW corrections can change significantly if the experimental setup is such that additional, unobserved massive EW bosons can be radiated into the final state since their contributions can cancel some of the EW Sudakov logarithms. A second issue related to experimental cuts is that near the partonic threshold the fragmentation contribution in photon production is power suppressed [3], but in practice one needs to define what is meant by a final-state photon, and depending on the photon isolation requirements, a fragmentation contribution will also arise and will involve different EW corrections. The experimental definition of the photon final state will also affect which value of electromagnetic coupling should be used when computing the cross section. [The value is obviously irrelevant for the ratio in Eq. (1) we present in this paper, but its choice is important when computing the cross section.] In our




FIG. 4. $\Delta\sigma^{\text{ew}}$ at 8 TeV for γ production at different rapidity ranges. In the highest rapidity range shown in the last plot, the cross section is essentially zero for $p_T > 1200 \text{ GeV}$.

computation, the appropriate value of the electromagnetic coupling is the value at the invariant mass of the vector-boson final state. If the experiment would be able to resolve photon jets with arbitrary small mass, the proper value would be $\mu = 0$, and one would use the on-shell, low-energy value of $\alpha_{\text{e.m.}} \approx 1/137$; see e.g. the discussion in Ref. [4]. However, with the photon isolation requirements at the LHC, the photon jet has a nonzero mass, and a value such as $\alpha_{\text{e.m.}}(M_Z) \approx 1/128$ might be more appropriate.

ACKNOWLEDGMENTS

We thank Marat Freytsis, Matt Schwartz, and Matt Strassler for discussions. This work is supported by the Swiss National Science Foundation under the Sinergia Grant No. CRSII2 141847 1 and the US Department of Energy under Grant No. DE-SC0013607. The authors thank the high-energy theory group at Harvard for hospitality and support.

-
- [1] T. Becher and X. Garcia i Tormo, Electroweak Sudakov effects in W , Z and γ production at large transverse momentum, *Phys. Rev. D* **88**, 013009 (2013).
 [2] K. Mishra *et al.*, Electroweak corrections at high energies, [arXiv:1308.1430](https://arxiv.org/abs/1308.1430).

- [3] T. Becher and M. D. Schwartz, Direct photon production with effective field theory, *J. High Energy Phys.* **02** (2010) 040.
 [4] A. Czarnecki and W. J. Marciano, Electroweak radiative corrections to $b \rightarrow s\gamma$, *Phys. Rev. Lett.* **81**, 277 (1998).