

ATLAS $Z +$ missing transverse energy excess in the MSSMM. Cahill-Rowley,^{1,*} J. L. Hewett,^{1,†} A. Ismail,^{2,3,‡} and T. G. Rizzo^{1,§}¹*SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA*²*Argonne National Laboratory, Argonne, Illinois 60439, USA*³*University of Illinois at Chicago, Chicago, Illinois 60607, USA*

(Received 26 June 2015; published 21 October 2015)

We demonstrate that the 3σ excess observed by ATLAS in the $Z +$ MET channel can be explained within the context of the MSSM. Using the freedom inherent in the pMSSM, we perform a detailed analysis of the parameter space and find a scenario that describes the excess while simultaneously complying with all other search constraints from the Run I data at 7 and 8 TeV, including the $Z +$ MET analysis by CMS. We generate a small sample of simplified models, using promising models from our existing pMSSM sample as seeds, and study their properties. The successful region is described by the production of 1st/2nd generation squark pairs, followed by their decay into a bino-like neutralino which in turn decays into a Higgsino-like LSP triplet by emitting a Z boson, i.e., $\tilde{q} \rightarrow \tilde{B} \rightarrow \tilde{h}$ with $\tilde{q} = \tilde{Q}_L, \tilde{u}_R, \text{ or } \tilde{d}_R$. The sweet spot for the sparticle spectrum is found to have squark masses in the 500–750 GeV range, with bino masses near 350 GeV with a mass splitting of 150–200 GeV with the Higgsino LSP. If this excess holds, then this scenario predicts that a signal will be observed in the $0l +$ jets and/or $1l +$ jets searches in the early operations of Run II.

DOI: [10.1103/PhysRevD.92.075029](https://doi.org/10.1103/PhysRevD.92.075029)

PACS numbers: 12.60.Jv, 14.80.Ly

I. INTRODUCTION

It is well-known that physics beyond the standard model (SM) must exist in order to address a number of outstanding questions such as the nature of dark matter, the generation of neutrino masses, the origin of the observed baryon asymmetry and the solution to the hierarchy problem—all of which remain unanswered. The nature of this new physics is presently mysterious: Not only is its form unknown, so is the energy scale at which it will first be revealed. Although constrained by data from Run I at the LHC, dark matter searches, and flavor physics observables, Supersymmetry (SUSY)[1] remains the leading theoretical framework to address at least some of these important puzzles. However, supersymmetry has so far been frustratingly elusive at the LHC, with numerous searches setting strong constraints on the simplest SUSY scenarios [2]. Nonetheless, the continual exploration of the SUSY parameter space remains mandatory, with missing transverse energy (MET) based searches at the LHC continuing to be the most promising avenue for discovery.

Along these lines, the ATLAS experiment recently announced [3] the observation of a 3σ excess in one of their Run I SUSY search channels, $Z +$ MET with $\geq 2j$, while a similar analysis by CMS [4] observed a result consistent with the expected SM background. Importantly, the detailed nature of the cuts employed by these two

experiments in this channel are sufficiently different, as we will discuss below, so that the apparent null result from CMS does not necessarily exclude the possibility of a signal being observed by ATLAS. However, an explanation of this potential signal within supersymmetry remains challenging, since any proposed scenario must also satisfy the constraints imposed by the plethora of ATLAS and CMS searches [2]. Nonetheless, a few new physics scenarios have been proposed [5] that could give rise to the observed ATLAS excess with varying degrees of success. In this work, we suggest a natural supersymmetric scenario, based on the pMSSM, which comfortably explains the ATLAS excess in the $Z +$ MET channel while evading all other searches.

Given the simple nature of the search channel, and the apparent rate of the excess, several features are clearly necessary for a supersymmetric model to provide a successful description of the data. Since the Z -boson is observed in the dilepton mode, the signal rate demands a strong production cross section, implying the production of relatively light gluinos or squarks which then decay to an intermediate state accompanied by jets. This intermediate, apparently neutral state, e.g., a neutralino, then decays via the emission of a Z plus the lightest supersymmetric particle (LSP), which produces the MET in a R-parity conserving scenario.¹ However, it is likely that such a spectrum would be easily excluded by, e.g., the $0l, 2-6$ jets + MET searches if the jets from the hadronic decay of the Z were sufficiently hard. Clearly the details of the

* mrowley@slac.stanford.edu† hewett@slac.stanford.edu‡ aismail@anl.gov§ rizzo@slac.stanford.edu¹In the analysis considered here, we will assume the LSP to be the lightest neutralino.

SUSY spectrum in such a scenario, in particular the relative masses and compositions of the sparticles, are highly constrained by multiple requirements and finding the right “balance” presents a significant model building challenge.

To perform this study, it is necessary to incorporate a detailed analysis of the available SUSY parameter space that remains viable after the Run I data at 7 and 8 TeV; there is no better way to accomplish this than to employ the 19-parameter p(henomenological)MSSM [6] which we have already studied in detail elsewhere [7]. In particular, this recent work contains a large sample of pMSSM models that are presently allowed, providing a viable playground for exploration. As will be discussed in detail below, an examination of these models reveals an intriguing scenario that describes the excess, while complying with all the constraints. Specifically, the 1st/2nd squarks, \tilde{Q}_L , \tilde{u}_R or \tilde{d}_R are identified as the leading candidates for the objects that initiate the ATLAS $Z + \text{MET}$ signal via a cascade decay. Once the other search constraints are taken into account, the primary production of gluinos in this role is found to yield too small of an event rate to explain the signal. The 1st/2nd generation squark scenario benefits from having both of the first two generations of squarks being produced simultaneously, as they are assumed to be degenerate in the pMSSM framework, yielding a large enough production rate. Whereas, within our pMSSM model sample, the 3rd generation squarks are also too highly constrained by specialized searches to play the role of the strong initiator of this signal. Within our successful scenario, the 1st/2nd generation squarks decay to a mostly binolike neutralino which then subsequently decays to a Higgsino-like LSP triplet by emitting a Z -boson. The masses and splittings dictated by this spectrum control the overall production rates for the different sparticles, the hardness of the jets and leptons, and the branching fractions for the intermediate neutralino decaying to the three light Higgsino states. We note that considering only a single set of squarks presents a somewhat simplified picture and that other states (such as \tilde{t} and \tilde{b}) may also contribute to the total signal, albeit in a secondary capacity. In the analysis below we use the successful models contained in our existing pMSSM sample as seeds to generate a small sample of simplified models that describe the excess while remaining consistent with the many other LHC searches. We then study the detailed properties of these simplified models and discuss the Run II analyses that can be used to elucidate this scenario more fully, or exclude it from further consideration.

II. ANALYSIS

The pMSSM [6] is the most general version of the R-parity conserving MSSM subjected to the guiding principles of CP -conservation, minimal flavor violation, and degenerate 1st and 2nd generation squark masses. Imposing these criteria reduces the number of free parameters in the MSSM to 19 (assuming a neutralino LSP):

$m_{\tilde{Q}_{L,1,2}}, m_{\tilde{Q}_3}, m_{\tilde{u}_{R,1,2}}, m_{\tilde{d}_{R,1,2}}, m_{\tilde{t}_1}, m_{\tilde{t}_2}, m_{\tilde{b}_1}, m_{\tilde{b}_2}, m_{\tilde{e}_{R,1,2}}, M_{1,2,3}, \mu, A_{t,b,\tau}, M_A$, and $\tan\beta$. In our previous work [7], we generated a large set of models (with a “model” describing a point in the 19-dimensional parameter space) by randomly scanning the parameter space, setting the upper limit on the scan at 4 TeV for the dimensionful parameters and taking $\tan\beta = 1\text{--}60$. The 4 TeV upper bound was chosen to facilitate collider studies at the 7, 8 and 13, 14 TeV LHC. We subjected these models to a global data set of collider, flavor, precision, dark matter and theoretical constraints. In particular, we have examined this model sample in light of the SUSY search results from Run I of the LHC [7], subjecting them to roughly 40 separate analyses performed by the experiments at 7 and 8 TeV. The result is a sample of approximately 125k models that remain viable at the end of Run I, including many models with light squarks and gluinos, providing an ample playground for further studies.

Here, we investigate our pMSSM model sample to determine whether a region of the parameter space could adequately describe the excess in the $Z + \text{MET}$ channel observed by ATLAS. For each model, we created SUSY production samples using Pythia 6.4 [8] for event generation and PGS [9] for detector simulation, normalizing to NLO cross sections from Prospino [10]. The details of this procedure are the same as for our previous pMSSM studies [7]. In particular, we have used the K-factors from Prospino assuming degenerate squarks at a mass equal to the average of all squark masses (including those that are decoupled). As our squarks in our scenario are not degenerate, this should be considered a source of systematic uncertainty. Other studies have considered the effect of nondegenerate squarks and gluinos [11], including squark decay [12]. For points that reproduce the $Z + \text{MET}$ excess, varying the masses of the decoupled squarks and gluino tends to change the resulting NLO cross sections by 20%–40%, which should be considered a systematic uncertainty on our predictions.

We then applied the cuts for the on- Z region of the ATLAS search for final states containing a pair of




FIG. 1 (color online). Sparticle spectrum for a representative pMSSM model that reproduces the ATLAS $Z + \text{MET}$ excess.

TABLE I. Scan ranges for the 3 variable parameters in each of the 3 grid scans described in the text.

Grid	μ (22 GeV steps)	M_1 (25 GeV steps)	$m_{\tilde{Q}_L}$ (35 GeV steps)
\tilde{Q}_L	100 GeV—254 GeV	$\mu + 100$ GeV— $m_{\tilde{Q}_L}$	MAX(350 GeV, $\mu + 150$ GeV)—900 GeV
\tilde{u}_R	100 GeV—254 GeV	$\mu + 100$ GeV— $m_{\tilde{Q}_L}$	MAX(300 GeV, $\mu + 150$ GeV)—800 GeV
\tilde{d}_R	100 GeV—254 GeV	$\mu + 100$ GeV— $m_{\tilde{Q}_L}$	MAX(250 GeV, $\mu + 150$ GeV)—700 GeV

opposite-sign dileptons, jets and MET [3] to these simulated SUSY samples. In particular, events were required to have at least two leptons with $p_T > 25$, 10 GeV respectively, two jets with $p_T > 35$ GeV, and MET > 225 GeV. The two hardest leptons were required to form a same-flavor opposite sign pair with invariant mass $m_{\ell\ell} = m_Z \pm 10$ GeV, and the scalar sum H_T of their transverse momenta and the transverse momenta of all jets with $p_T > 35$ GeV was required to be at least 600 GeV. We also imposed standard cuts on the rapidity and isolation of jets and leptons, and required an angular separation $\Delta\phi(j, \text{MET}) > 0.4$ between each of the two leading jets and the missing transverse momentum, as described by ATLAS.

Several of our pMSSM models predict significant numbers of events passing the ATLAS Z+MET cuts. The sparticle spectrum for such a representative pMSSM model is shown in Fig. 1. We observe a common pattern in these models, with light-flavor squarks decaying through gaugino cascades producing Z-bosons. The direct decay of the squarks to the lightest neutralino is usually suppressed by weak couplings to a Higgsino-like LSP (or alternatively a wino-like LSP if the squark is right-handed). Additionally, we find that these models tend to predict an observable excess in jets + MET searches, creating some tension with the null results in other Run I LHC SUSY searches that we have considered previously [7]. In particular, it is challenging to reproduce the ATLAS Z+MET excess in the leptonic channel while simultaneously satisfying bounds from the ATLAS jets + MET search [13]. However, the pMSSM models from our sample that reproduce the Z+MET excess often predict a jets + MET event rate that is near the boundary of the existing limits. In particular, the number of events in the jets + MET search signal regions is typically reduced because the squarks decay mainly to a heavier binolike neutralino rather than directly to the LSP.

Encouraged by these results, we are motivated to consider points with similar spectra to these promising pMSSM models, which may predict a significant number of Z+MET events while fully evading constraints from the other LHC SUSY searches. Given the results of our pMSSM analysis, we focus on simplified spectra with the dominant decay pattern $\tilde{q} \rightarrow \tilde{B} \rightarrow \tilde{h}$, where \tilde{q} is a light-flavor squark \tilde{Q}_L , \tilde{u}_R or \tilde{d}_R . Starting with a seed point taken from one of our successful pMSSM models, we vary the 3 most relevant Lagrangian parameters ($m_{\tilde{Q}_{L,2}}$, μ , and M_1) in

a grid around the region of interest. This corresponds to adjusting the most relevant physical sparticle masses, specifically $m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_3^0}$, and the squark masses $m_{\tilde{q}}$. Note that neither the sign of μ or the value of $\tan\beta$ are varied, as this would modify the details of the Higgsino spectrum yet leave the gaugino branching fractions mostly unaltered.² We expect the remaining pMSSM parameters to have a negligible impact on the simplified model phenomenology, as long as they are sufficiently heavy. We thus leave their values as given in the pMSSM seed model, with the exception of A_t which is adjusted to produce the observed value of the Higgs mass within theoretical and experimental uncertainties. In all cases, the squark masses not being scanned are set above 2 TeV. We use SOFTSUSY [14] to generate spectra, and SUSY-HIT [15] to calculate sparticle decays, which includes QCD corrections as well as the most important electroweak corrections.

To produce this grid, we scan μ between 100 GeV and 254 GeV (with the lower limit set by LEP constraints and upper limit set by kinematic considerations), with 22 GeV steps. Note that given this small value of μ we would expect these models to exhibit low values of fine-tuning from this source. We then scan the *physical* squark mass $m_{\tilde{Q}_L} \approx m(\tilde{u}_L) \approx m(\tilde{d}_L)$ between 350 GeV or $\mu + 150$ GeV (whichever is larger) and 900 GeV in increments of 35 GeV.³ Finally, we scan M_1 between $\mu + 100$ GeV and $m_{\tilde{Q}_L}$ with 25 GeV increments. We employ the same procedure (with slightly different scan ranges noted in Table I) to construct two additional grids, one each for \tilde{u}_R/\tilde{c}_R and \tilde{d}_R/\tilde{s}_R .

Figure 2 shows the relevant spectrum and branching fractions for one of our grid points (from the \tilde{Q}_L grid), which predicts 21 events in the ATLAS Z+MET search and is consistent with all other searches; this model is illustrative of the typical decay patterns for scenarios that reproduce the excess. The key features to note are the large branching fractions for squarks decaying to $\tilde{\chi}_3^0$, and the

²Except for cases where some of the Higgsinos are kinematically inaccessible to decays of the binolike $\tilde{\chi}_3^0$.

³Since we do not know the physical squark mass before spectrum generation, we estimate the soft mass required to give the desired physical mass. While approximate, this estimation is easily accurate enough to ensure that our scan grid is covering the region of interest.




FIG. 2. Spectrum and decay patterns for a model in the \tilde{Q}_L grid, which predicts 21 events in the ATLAS 20 fb^{-1} $Z + \text{MET}$ analysis and is consistent with all other implemented searches. Numbers indicate the branching fraction in percent for each decay mode (only branching fractions larger than 5% are shown for simplicity).

multiple possible decays of $\tilde{\chi}_3^0$, about a quarter of which result in Z boson production.

III. RESULTS

We now examine the results of our scan over the simplified pMSSM spectra. As noted above, the strongest restrictions on the parameter space arise from the null results of other LHC SUSY searches, which are generally in tension with our goal of producing a large signal rate in the ATLAS 20 fb^{-1} $Z + \text{MET}$ analysis. Clearly, we require that a successful model point produces $\sim 15\text{--}20$ signal events for the ATLAS 20 fb^{-1} $Z + \text{MET}$ analysis. In addition, we also require the point to simultaneously satisfy the limit from the corresponding CMS search with different selection criteria. Finally, a successful model point must satisfy all of the null ATLAS search results in other channels. In particular, it is clear that both the ATLAS $1l + \text{jets}$ search (arising in the spectra we consider from the heavy bino decay producing a W instead of a Z) and the $0l, 2\text{--}6 \text{ jets}$ search (when W, Z , or the Higgs are produced and decay hadronically) will also be important in determining the detailed nature of a successful parameter space point.

The impact of these other searches will be discussed in more detail below.

To get an initial handle on the preferred parameter regions, Fig. 3 shows the LSP mass versus the mass splitting between the intermediate binolike $\tilde{\chi}_3^0$ and the LSP for fixed values of the squark masses in the 450–700 GeV range before imposing any additional constraints.⁴ The various colored regions show the anticipated ATLAS $Z + \text{MET}$ analysis event yields and, as we would naively expect, we see that lighter squarks will generally lead to larger signal rates due to their significantly larger production cross sections. Also, we see that the Higgsino-like LSP prefers to be relatively light, below $\sim 180\text{--}190$ GeV. Perhaps, most interestingly, we observe that the most favored range for the electroweakino mass splitting lies above ~ 150 GeV. This might be counterintuitive since we would naively expect that a mass splitting, Δm_{31} , in the range of $\sim 90\text{--}125$ GeV would be most desirable, since decays to the Higgs in this region would be kinematically forbidden, thereby increasing the branching fraction for decays through the Z . Clearly, in all cases we see that the largest signal rates are obtained when the $\tilde{\chi}_3^0$ is kinematically allowed to decay through both the Z and the Higgs, due to the increased visibility of the decay products. Of course the preferred range of Δm_{31} is somewhat sensitive to the nature of the parent squark. In the \tilde{Q}_L case, a value of $\Delta m_{31} \sim 150\text{--}200$ GeV is preferred while for $\tilde{u}_R(\tilde{d}_R)$ this value is significantly larger $\Delta m_{31} \sim 200\text{--}300(250\text{--}350)$ GeV (as we will see more clearly below) in obvious correlation with the production cross sections, i.e., the parent squark with the largest (smallest) production cross section prefers the smallest (largest) corresponding value of Δm_{31} .

Figures 4, 5 and 6 present the results from our scans of the simplified pMSSM spectra for each parent squark type, where the vertical bars represent the scanned regions in our somewhat coarsely spaced grid. The location of the vertical bars is set by the approximate mass of the parent squark and the LSP mass. The lowest slice of each of the vertical bars corresponds to the smallest value of the bino-LSP mass splitting, Δm_{31} , scaled by a factor of 0.02, while each successive higher slice (going up the bar) corresponds to increasing this value. The color code indicates the number of events predicted for the ATLAS 20 fb^{-1} $Z + \text{MET}$ analysis, with yellow to red tones indicating a higher event rate in agreement with the observed rate. Black regions represent points which are excluded by the corresponding CMS $Z + \text{MET}$ search, or by any of the ~ 40 null SUSY searches described in [7]. In the \tilde{Q}_L grid, the $0l + \text{jets}$ channel results gives by far the most important constraints, while for \tilde{u}_R , and particularly \tilde{d}_R , other searches also play

⁴Note that we have interpolated between grid points, smoothing out modest fluctuations in event yields.




FIG. 3 (color online). Signal event rate contours for the ATLAS Z + MET analysis in the $\chi_3^0 - \chi_1^0$ mass difference and χ_1^0 mass plane. The top three panels correspond to the case of $\tilde{Q}_L = 500, 600, 700$ GeV from left to right, while the bottom panels are for $\tilde{u}_R = 450, 500$ GeV and $\tilde{d}_R = 450$ GeV, left to right.




FIG. 4 (color online). Results from the simplified spectra scan for a parent Q_L -squark in the \tilde{Q}_L and $\chi_1^0 + (1/50)\Delta m_{31}$ mass plane. The vertical bars represent the coarse grid in our scan, with the value of the mass splitting Δm_{31} increasing for successively higher slices of the bar. The color code indicates the predicted event rate for the ATLAS 20 fb^{-1} Z + MET channel. Black slices in a vertical bar correspond to points excluded by any of the simulated searches.




FIG. 5 (color online). Results from the simplified spectra scan for a parent u_R -squark in the \tilde{u}_R and $\chi_1^0 + (1/50)\Delta m_{31}$ mass plane. The vertical bars represent the coarse grid in our scan, with the value of the mass splitting Δm_{31} increasing for successively higher slices of the bar. The color code indicates the predicted event rate for the ATLAS 20 fb^{-1} Z + MET channel. Black slices in a vertical bar correspond to points excluded by any of the simulated searches.




FIG. 6 (color online). Results from the simplified spectra scan for a parent d_R -squark in the \tilde{d}_R and $\chi_1^0 + (1/50)\Delta m_{31}$ mass plane. The vertical bars represent the coarse grid in our scan, with the value of the mass splitting Δm_{31} increasing for successively higher slices of the bar. The color code indicates the predicted event rate for the ATLAS 20 fb^{-1} $Z + \text{MET}$ channel. Black slices in a vertical bar correspond to points excluded by any of the simulated searches.

an important role. The results are seen to be quite different for the three parent squark cases we consider. In particular, we see that the \tilde{Q}_L case provides the best fit to the excess. However, even in this case the most successful points are close to the black excluded regions, indicating that the other LHC SUSY searches are providing important constraints on this scenario. We also find the strong constraints at larger values of Δm_{31} , due to larger contributions to the $0l + \text{jets}$ channel when this splitting is too large. In general, we expect the $0l + \text{jets}$ rate to place strong restrictions on the \tilde{Q}_L scenario because of either a large production rate for relatively light squark masses or because the rate is still reasonably large for heavier masses where the jets from the decay of the W, Z and Higgs bosons appearing in the bino to LSP transitions are becoming sufficiently hard to pass the $0l + \text{jets}$ cuts. For surviving points explaining the $Z + \text{MET}$ excess, we find that low LSP masses and moderate values of Δm_{31} are preferred, and that the production rate falls off too quickly for squark masses above ~ 800 GeV to generate a sufficient number of events. In the case of a \tilde{u}_R parent, both the favored region and the region excluded by the other searches are smaller (a simple consequence of the lower production cross section) and, overall, lower signal rates are obtained. These same features are seen to be further emphasized for the case where the parent squark is a \tilde{d}_R . In all cases, we find that the sweet spot for describing the excess is in the region where the parent squark is 500–700 GeV with a LSP mass of 100–200 GeV and a bino-LSP mass splitting of 100–250 GeV.

For each parent squark type, we next examine the impact of the $0l, 1l + \text{jets}$ and CMS $Z + \text{MET}$ channels. Here, we

only study the set of models from our simplified grid scan that are consistent with the constraints in these channels. For each of these analyses, we compute the expected number of events in each signal region, and show the ratio of the expected number of events to the 95% C.L. event exclusion limit for the most important signal region for that channel (i.e., the signal region with the largest value of this ratio). For example, a ratio of 0.5 indicates that the model predicts 1/2 as many events as are allowed by the relevant null search result at 95% C.L. Figure 7 displays these results for the case of the $0l + \text{jets}$ ATLAS search [13], i.e., the event rate ratio R_{0l} , as a function of the number of predicted events for the ATLAS 20 fb^{-1} $Z + \text{MET}$ analysis for all three parent squark types. The color code indicates the value of Δm_{31} . The top-left, top-right, and bottom panels correspond to the parent squarks $\tilde{Q}_L, \tilde{u}_R,$ and \tilde{d}_R , respectively. In the \tilde{Q}_L parent case, we see that many of the models lie close to the $0l + \text{jets}$ exclusion boundary. In particular, we see that for model points with at least 15 ATLAS 20 fb^{-1} $Z + \text{MET}$ signal events, the values of R_{0l} lie in the range 0.6–1. Generally an increase in the number of ATLAS 20 fb^{-1} $Z + \text{MET}$ signal events corresponds to a larger value of R_{0l} , so that at some point consistency with the $0l + \text{jets}$ search prevents larger signal rates from being obtained. Also, we see that as Δm_{31} decreases for a fixed signal rate, the points are farther away from the $0l + \text{jets}$ exclusion boundary since the jets produced by W, Z and Higgs decays are becoming correspondingly softer. Considering the \tilde{u}_R parent case, we find that the model points are a bit further away from the $0l + \text{jets}$ boundary (due to the smaller production cross section), but we also find, correspondingly, fewer models that produce a significant signal in the $Z + \text{MET}$ analysis. This trend continues for the case of the \tilde{d}_R parent.

Figure 8 displays our results for the case of the $1l + \text{jets}$ ATLAS search [16] where the y-axis now shows the event rate ratio R_{1l} . For all three squark parent cases we see that the models tend to mostly lie reasonably far away from the exclusion boundary for this search, implying that it has little impact on shaping the parameter region for successful models. In fact, we find that few models are excluded by the $1l + \text{jets}$ analysis after the other null search results have been applied. Figure 9 shows the results for the case of the CMS $Z + \text{MET}$ analysis [4], expressed as the ratio R_{CMS} ; clearly for all squark parents there is a rough linear correlation between the value of R_{CMS} and the number of predicted ATLAS 20 fb^{-1} $Z + \text{MET}$ signal events. From this one might expect that requiring $R_{\text{CMS}} \leq 1$ cuts off the corresponding ATLAS signal. However, this region is already restricted by the $0l + \text{jets}$ ATLAS search, with the result that the CMS $Z + \text{MET}$ analysis has only a small additional impact on our model selection beyond the effect of the $0l + \text{jets}$ ATLAS search.

In addition to the $0l, 1l + \text{jets}$ and CMS $Z + \text{MET}$ searches, other ATLAS searches which are less clearly




FIG. 7 (color online). Ratio of the predicted number of events for models in our simplified grid scan to the ATLAS 95% C.L. event limit for the $0l + \text{jets}$ channel, R_{0l} , as a function of the number of signal events for the ATLAS 20 fb^{-1} $Z + \text{MET}$ search. The color code corresponds to the value of the $\chi_3^0 - \chi_1^0$ mass splitting. The top-left, top-right, and bottom panels correspond to the three grid scans, with \tilde{Q}_L , \tilde{u}_R , and \tilde{d}_R parent squarks, respectively.

targeted for these types of models can still have an important impact on the allowed parameter space, especially for the \tilde{u}_R and \tilde{d}_R grids, where the $0l$ search is less dominant. In particular, the ATLAS $3l$ gaugino search [17], $4l$ search [18], and same-sign dilepton search [19] all make unique contributions to the combined exclusion region. Since the $3l$ and $4l$ searches are targeted at electroweak production, it is unsurprising that they are particularly sensitive to the lower mass regions allowed for the right-handed squarks, particularly \tilde{d}_R .

It is worth a short discussion to compare the event selection between the ATLAS and CMS $Z + \text{MET}$ analyses. While both searches select events with a leptonic Z , at least two jets, and missing energy, the 600 GeV H_T cut of the ATLAS search is highly effective at reducing Drell-Yan background, leaving $t\bar{t}$ as the dominant background process. The CMS analysis considers multiple search regions with missing energy bins to gain increased sensitivity, but even in the highest bin, requiring $\text{MET} > 300$ GeV, Drell-Yan production is still the most significant background. For

comparison, the ATLAS analysis imposes the tight H_T cut stated above and simply cuts on missing energy, $\text{MET} > 225$ GeV. As a result of these cut choices, the overlap between the ATLAS and CMS $Z + \text{MET}$ search regions is small [20].

We now comment on the effect of 13 TeV searches for supersymmetry on the parameter space we have outlined. As mentioned above, Fig. 7 shows a correlation between the number of events in the ATLAS $Z + \text{MET}$ search region predicted by any given point and the capability of the $0l$ search for supersymmetry to probe the associated spectrum. Now, for the squark masses that provide the best fit to the $Z + \text{MET}$ excess, the pair production cross section from pp collisions is higher at 13 TeV than at 8 TeV by a factor of 5–10 [21]. On the other hand, the backgrounds for the $0l + \text{jets}$ search regions are dominated by $V + \text{jets}$ and $t\bar{t}$, which increase much more slowly in going from 8 to 13 TeV; the top pair production cross section rises by only a factor of roughly 3 [22], and that for $V + \text{jets}$ by even less. It is thus reasonable to expect that 13 TeV searches for




FIG. 8 (color online). Ratio of the predicted number of events for models in our simplified grid scan to the ATLAS 95% C.L. event limit for the $1l + \text{jets}$ channel, R_{1l} , as a function of the number of signal events for the ATLAS $20 \text{ fb}^{-1} Z + \text{MET}$ search. The color code corresponds to the value of the $\chi_3^0 - \chi_1^0$ mass splitting. The top-left, top-right, and bottom panels correspond to the parent squark cases of \tilde{Q}_L , \tilde{u}_R , and \tilde{d}_R , respectively.

supersymmetry in the $0l + \text{jets} + \text{MET}$ channel will serve as a test of the squark scenario that we have found as a potential explanation for the ATLAS $Z + \text{MET}$ excess. Similar arguments apply for the $1l$ channel; while there is less correlation in Fig. 8 between the results of the $1l$ search and the number of $Z + \text{MET}$ events, most points giving sufficient $Z + \text{MET}$ events are within a factor of 2–3 of being excluded by the $1l$ search at 8 TeV. For both searches, a full study would be worthwhile to evaluate the impact of 13 TeV supersymmetry searches on the parameter space that can explain the ATLAS $Z + \text{MET}$ excess.

As discussed above, our successful models kinematically allow the decay of the intermediate binolike state into the lighter Higgsinos by W , Z and Higgs boson emission. Apart from phase space considerations, these relative branching fractions are controlled by the bino and Higgsino content of the gauginos.⁵ Since $\tan\beta$ is being held fixed in our grid

⁵We essentially treat the winos as being decoupled with a correspondingly large value of M_2 .

scans, the bino and Higgsino content of these states are only being regulated (at tree-level) by the values of M_1 and μ . Clearly as the mass splitting between these states, Δm_{31} , increases, the purity of each state increases. If the intermediate state were to be *pure* bino, then its decay via either the W or Z -boson would be forbidden (as these channels require both the initial and final states to have a nonzero Higgsino content), while decays through the Higgs would remain allowed (as this is controlled instead by the product of the bino and Higgsino content of both the initial and final states). Figure 10 shows the distributions for the χ_3^0 branching fractions for the \tilde{Q}_L parent squark case for all the models in the initial grid, as well as after applying constraints from the CMS $Z + \text{MET}$ search and null results in other ATLAS channels and requiring the point to predict 12 or more events in the ATLAS $20 \text{ fb}^{-1} Z + \text{MET}$ search. Here we see several things: (i) the typical W -boson branching fraction is rather large, although models with the largest values for this branching fraction are unable to satisfy the constraints applied to the right panel. (ii) In both




FIG. 9 (color online). Ratio of the predicted number of events for models in our simplified grid scan to the CMS 95% C.L. event limit for the Z + MET channel, R_{CMS} , as a function of the number of signal events for the ATLAS 20 fb⁻¹ Z + MET search. The color code corresponds to the value of the $\tilde{\chi}_3^0 - \tilde{\chi}_1^0$ mass splitting. The top-left, top-right, and bottom panels correspond to the parent squark cases of \tilde{Q}_L , \tilde{u}_R , and \tilde{d}_R , respectively.




FIG. 10 (color online). Distributions of the branching fractions for the decay modes, as indicated, of the $\tilde{\chi}_3^0$ intermediate state for the full set of models from our \tilde{Q}_L grid scan (left panel) and for models predicting 12 or more events in the ATLAS 20 fb⁻¹ Z + MET search while remaining consistent with all simulated searches, including the ATLAS 0l, 1l + jets and CMS Z + MET searches (right panel).




FIG. 11 (color online). Mass distributions of the parent squark, χ_3^0 and χ_1^0 states for the models from our grid scan that agree with all null search results and yield at least 5 events in the ATLAS $20 \text{ fb}^{-1} Z + \text{MET}$ channel. The top-left, top-right, and bottom panels correspond to the parent squark cases of \tilde{Q}_L , \tilde{u}_R , and \tilde{d}_R , respectively.

panels, the Z -boson branching fraction is more than twice as large for the decay into $\tilde{\chi}_2^0$ than for decays to the LSP. The reverse is true for decays producing a Higgs boson. (iii) The χ_3^0 decays mediated via the Z -boson and the Higgs, to either the LSP or to $\tilde{\chi}_2^0$, are seen to have similar branching fractions. Clearly it is not advantageous to completely suppress the Higgs mode, which can only be accomplished by reducing Δm_{31} to values below the Higgs mass. Interestingly, this scenario would then also predict a signal in the $h + \text{MET}$ channel. We obtain similar results for the other squark parent scenarios.

Lastly, it is interesting to examine the mass distributions of the squarks, binolike $\tilde{\chi}_3^0$ and Higgsino-like LSP states in the models that successfully reproduce the ATLAS $20 \text{ fb}^{-1} Z + \text{MET}$ signal (here defined to be $N \geq 5 Z + \text{MET}$ events). This is presented for the three parent squark types in Fig. 11. Here we see that the overall spectra of these three particles gradually become lighter as we compare the \tilde{Q}_L parent to \tilde{u}_R and then to \tilde{d}_R , reflecting the corresponding

falling squark pair production cross sections, with these distributions peaking at 750, 650, and 600 GeV, respectively. In all three cases the peak of the χ_3^0 distribution is near ~ 350 GeV resulting in a softening of the jets on average, for the \tilde{u}_R/\tilde{d}_R cases compared to the \tilde{Q}_L parent squark, due to a compression of the spectrum. The peak of the LSP mass distribution lies roughly near ~ 200 GeV for the \tilde{Q}_L parent squark and near ~ 150 GeV for both the \tilde{u}_R and \tilde{d}_R cases, implying that Δm_{31} prefers to lie near $\sim 150\text{--}200$ GeV in all cases.

IV. CONCLUSIONS

We have examined the $3\sigma Z + \text{MET}$ excess observed by the ATLAS collaboration in Run I of the LHC in the context of a supersymmetric framework. We have employed the freedom inherent in the pMSSM parameter space to explore whether supersymmetric models can be constructed that generate the observed excess, while simultaneously being

consistent with the numerous other null SUSY searches at the LHC. Using a large pMSSM model sample that we had previously generated, we found a handful of points that satisfied our criteria, demonstrating the power of this approach. These points shared characteristics for the sparticle spectrum that are crucial for describing the data, namely relatively light 1st/2nd generation squarks that decay into a bino-like neutralino, which in turn decays into a light Higgsino multiplet, i.e., $\tilde{q} \rightarrow \tilde{B} \rightarrow \tilde{h}$. Using these points as seeds, we performed three grid scans, corresponding to the possible types of the parent squark, \tilde{Q}_L , \tilde{u}_R and \tilde{d}_R . We scanned over the set of relevant parameters, $m_{\tilde{q}}$, M_1 and μ , and generated three sets of simplified pMSSM models within a limited kinematic range. All other strongly-interacting sparticles were set to the same value as in the parent pMSSM model (~ 2 TeV) and A_t was varied to reproduce the observed Higgs boson mass. In principle it is possible that light stop and sbottom squarks could also contribute to the signal, but we limited our analysis here to the simplest scenario.

We then examined the properties of these simplified models in detail. They predict a range of event rates, up to 21 events, for the ATLAS 20 fb^{-1} Z + MET channel, in agreement with the ATLAS measurement. Several hundred of our grid points were found to be consistent with the 95% C.L. bounds from all simulated searches, including the ATLAS $0l$, $1l$ + jet and CMS Z + MET search channels. The case of a left-handed doublet parent squark, \tilde{Q}_L , is

found to yield the best fit to the data, with the other scenarios giving slightly smaller event rates. The sweet spot for the sparticle spectrum is found to have squark masses in the 500–750 GeV range, with bino masses near 350 GeV with a mass splitting of 150–200 GeV with the Higgsino LSP. The bino χ_3^0 state has important decays involving W and Higgs bosons, as well as the Z-boson. The predicted event rates for these models are close to the 95% C.L. limits from the $0l$ + jets search and the CMS Z + MET analysis, but lie somewhat further from the $1l$ +jet search results.

In conclusion, we have constructed a simplified supersymmetric model based on the pMSSM, with specific characteristic features that successfully yields an excess for the 20 fb^{-1} Z + MET ATLAS analysis, while evading all other SUSY searches at the Run I LHC. The operations at the 13 TeV LHC currently underway will be able to quickly discover, or exclude, this scenario. If the ATLAS 3σ excess is confirmed with the new data set, it could very well be a signal for supersymmetry.

ACKNOWLEDGMENTS

The authors would like to thank Brian Petersen for communications. This work was supported by the Department of Energy, Contracts No. DE-AC02-06CH11357, No. DE-AC02-76SF00515 and No. DE-FG02-12ER41811.

-
- [1] For an overview of MSSM physics and phenomenology as well as mSUGRA, see M. Drees, R. Godbole, and P. Roy, *Theory and Phenomenology of Sparticles* (World Scientific, Singapore, 2004), p. 555; H. Baer and X. Tata, *Weak Scale Supersymmetry: From Superfields to Scattering Events* (Cambridge University Press, Cambridge, England, 2006), p. 537; S. P. Martin, A supersymmetry primer, *Adv. Ser. Dir. High Energy Phys.* **21**, 1 (2010); Perspectives on supersymmetry, *Adv. Ser. Dir. High Energy Phys.* **18**, 1 (1998).
- [2] For recent overviews of SUSY searches at the LHC by ATLAS and CMS, see the talk given by Z. Marshall (ATLAS Collaboration) and those given by S. Folgueras and H. W. Wulsin (CMS Collaboration) in 50th Recontres de Moriond QCD, La Thuile, Italy, 2015 (to be published).
- [3] G. Aad *et al.* (ATLAS Collaboration), Search for supersymmetry in events containing a same-flavour opposite-sign dilepton pair, jets, and large missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector, *Eur. Phys. J. C* **75**, 318 (2015).
- [4] V. Khachatryan *et al.* (CMS Collaboration), Search for physics beyond the standard model in events with two leptons, jets, and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV, *J. High Energy Phys.* **04** (2015) 124.
- [5] See, for example, B. A. Dobrescu, Leptophobic boson signals with leptons, jets and missing energy, [arXiv:1506.04435](https://arxiv.org/abs/1506.04435); J. Cao, L. Shang, J. M. Yang, and Y. Zhang, Explanation of the ATLAS Z-peaked excess in the NMSSM, *J. High Energy Phys.* **06** (2015) 152; A. Kobakhidze, A. Saavedra, L. Wu, and J. M. Yang, ATLAS Z-peaked excess in MSSM with a light sbottom or stop, [arXiv:1504.04390](https://arxiv.org/abs/1504.04390) [*Phys. Rev. D* (to be published)]; B. Allanach, A. Raklev, and A. Kvellestad, Consistency of the recent ATLAS $Z + E_T^{\text{miss}}$ excess in a simplified GGM model, *Phys. Rev. D* **91**, 095016 (2015); U. Ellwanger, Possible explanation of excess events in the search for jets, missing transverse momentum and a Z boson in pp collisions, *Eur. Phys. J. C* **75**, 367 (2015); N. Vignaroli, Z-peaked excess from heavy gluon decays to vector-like quarks, *Phys. Rev. D* **91**, 115009 (2015); G. Barenboim, J. Bernabeu, V. A. Mitsou, E. Romero, E. Torro, and O. Vives, METing SUSY on the Z peak, [arXiv:1503.04184](https://arxiv.org/abs/1503.04184).
- [6] A. Djouadi *et al.* (MSSM Working Group Collaboration), The minimal supersymmetric standard model: group summary report [arXiv:hep-ph/9901246](https://arxiv.org/abs/hep-ph/9901246); C. F. Berger, J. S.

- Gainer, J. L. Hewett, and T. G. Rizzo, Supersymmetry without prejudice, *J. High Energy Phys.* **02** (2009) 023.
- [7] M. Cahill-Rowley, J. L. Hewett, A. Ismail, and T. G. Rizzo, Lessons and prospects from the pMSSM after LHC Run I, *Phys. Rev. D* **91**, 055002 (2015). For earlier work, see, J. A. Conley, J. S. Gainer, J. L. Hewett, M. P. Le, and T. G. Rizzo, Supersymmetry without prejudice at the LHC, *Eur. Phys. J. C* **71**, 1697 (2011); M. W. Cahill-Rowley, J. L. Hewett, S. Hoeche, A. Ismail, and T. G. Rizzo, The new look pMSSM with neutralino and gravitino LSPs, *Eur. Phys. J. C* **72**, 2156 (2012); M. W. Cahill-Rowley, J. L. Hewett, A. Ismail, and T. G. Rizzo, The Higgs sector and fine-tuning in the pMSSM, *Phys. Rev. D* **86**, 075015 (2012); M. W. Cahill-Rowley, J. L. Hewett, A. Ismail, and T. G. Rizzo, More energy, more searches, but the pMSSM lives on, *Phys. Rev. D* **88**, 035002 (2013).
- [8] T. Sjostrand, S. Mrenna, and P. Z. Skands, PYTHIA 6.4 physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [9] J. Conway, PGS4, Pretty Good detector Simulation, <http://www.physics.ucdavis.edu/~conway/research/software/pgs/pgs.html>.
- [10] W. Beenakker, R. Hopker, M. Spira, and P. M. Zerwas, Squark and gluino production at hadron colliders, *Nucl. Phys.* **B492**, 51 (1997); W. Beenakker, R. Hopker, and M. Spira, PROSPINO: A program for the production of supersymmetric particles in next-to-leading order QCD, [arXiv: hep-ph/9611232](https://arxiv.org/abs/hep-ph/9611232); W. Beenakker, M. Kramer, T. Plehn, M. Spira, and P. M. Zerwas, Stop production at hadron colliders, *Nucl. Phys.* **B515**, 3 (1998); W. Beenakker, M. Klasen, M. Kramer, T. Plehn, M. Spira, and P. M. Zerwas, The Production of Charginos/Neutralinos and Sleptons at Hadron Colliders, *Phys. Rev. Lett.* **83**, 3780 (1999); **100**, 029901(E) (2008).
- [11] D. Goncalves-Netto, D. Lopez-Val, K. Mawatari, T. Plehn, and I. Wigmore, Automated squark and gluino production to next-to-leading order, *Phys. Rev. D* **87**, 014002 (2013).
- [12] W. Hollik, J. M. Lindert, and D. Pagani, NLO corrections to squark-squark production and decay at the LHC, *J. High Energy Phys.* **03** (2013) 139; W. Hollik, J. M. Lindert, and D. Pagani, On cascade decays of squarks at the LHC in NLO QCD, *Eur. Phys. J. C* **73**, 2410 (2013); R. Gavin, C. Hangst, M. Krmer, M. Mhlleitner, M. Pellen, E. Popena, and M. Spira, Matching squark pair production at NLO with parton showers, *J. High Energy Phys.* **10** (2013) 187; R. Gavin, C. Hangst, M. Krmer, M. Mhlleitner, M. Pellen, E. Popena, and M. Spira, Squark production and decay matched with parton showers at NLO, *Eur. Phys. J. C* **75**, 29 (2015).
- [13] G. Aad *et al.* (The ATLAS Collaboration), Report No. ATLAS-CONF-2013-047.
- [14] B. C. Allanach, SOFTSUSY: A program for calculating supersymmetric spectra, *Comput. Phys. Commun.* **143**, 305 (2002).
- [15] A. Djouadi, M. M. Muhlleitner, and M. Spira, Decays of supersymmetric particles: The program SUSY-HIT (SUSpect-SdecaY-Hdecay-InTerface), *Acta Phys. Pol. B* **38**, 635 (2007).
- [16] G. Aad *et al.* (ATLAS Collaboration), Search for squarks and gluinos in events with isolated leptons, jets and missing transverse momentum at $\sqrt{s} = 8$ TeV with the ATLAS detector, *J. High Energy Phys.* **04** (2015) 116.
- [17] G. Aad *et al.* (ATLAS Collaboration), Reports No. ATLAS-CONF-2012-154, No. ATLAS-COM-CONF-2012-192.
- [18] G. Aad *et al.* (ATLAS Collaboration), Reports No. ATLAS-CONF-2012-153, No. ATLAS-COM-CONF-2012-189.
- [19] G. Aad *et al.* (ATLAS Collaboration), Reports No. ATLAS-CONF-2012-105, No. ATLAS-COM-CONF-2012-141.
- [20] A. Haas (ATLAS Collaboration), in *Les Rencontres de Physique de la Vallée d'Aoste, La Thuile, Aosta Valley, Italy, 2015* (to be published).
- [21] M. Kramer, A. Kulesza, R. van der Leeuw, M. Mangano, S. Padhi, T. Plehn, and X. Portell, Supersymmetry production cross sections in pp collisions at $\sqrt{s} = 7$ TeV, [arXiv:1206.2892](https://arxiv.org/abs/1206.2892); C. Borschensky, M. Krmer, A. Kulesza, M. Mangano, S. Padhi, T. Plehn, and X. Portell, Squark and gluino production cross sections in pp collisions at $\sqrt{s} = 13, 14, 33$ and 100 TeV, *Eur. Phys. J. C* **74**, 3174 (2014).
- [22] M. Czakon and A. Mitov, Top++: A program for the calculation of the top-pair cross-section at hadron colliders, *Comput. Phys. Commun.* **185**, 2930 (2014).