

Accretion in the presence of an axisymmetric solitary vortexV. S. Beskin^{*}*P.N. Lebedev Physical Institute, Leninsky prospekt, 53, Moscow 119991, Russia,
and Moscow Institute of Physics and Technology, 9 Institutskiy pereulok, Dolgoprudny,
Moscow Region 141700, Russian Federation*K. V. Lezhnin[†]*Moscow Institute of Physics and Technology, 9 Institutskiy pereulok, Dolgoprudny,
Moscow Region 141700, Russian Federation*

(Received 19 May 2015; published 21 October 2015)

We analyze the quasispherical accretion in the presence of axisymmetric curl. The condition allowing the curl to reach the central object is formulated. Possible implications of the solution to the theory of turbulent accretion are discussed.

DOI: [10.1103/PhysRevD.92.083012](https://doi.org/10.1103/PhysRevD.92.083012)

PACS numbers: 97.10.Gz

I. INTRODUCTION

An activity of many astrophysical sources (active galactic nuclei, young stellar objects, galactic x-ray sources, microquasars) is associated with the accretion. For this reason, the accretion onto compact objects (neutron stars or black holes) is the classical problem of modern astrophysics (see, e.g., Lipunov [1] Shapiro and Teukolsky [2], and references therein). At present the analytical approach, whose foundation was laid back in the mid-twentieth century [3,4], began to be supplanted by numerical simulations [5–9]. Analytical solutions were found only in exceptional cases [10–16].

It should be emphasized that last time the focus of the research has been shifted to numerical magnetohydrodynamic simulations, within which framework it has become possible to take into account the turbulent processes associated with magnetic reconnection, magnetorotational instability, etc. [17–19]. However, in our opinion, some important features of the turbulent accretion can still be understood on the ground of simple analytical model.

In our paper we discuss a problem of dynamics of a solitary vortex against the background of spherical Bondi flow. We consider the case of the vorticity similar to Sec. 5.1 from Kovalenko and Eremin [20] corresponding to the second order of expansion in terms of our small parameter. As to linear perturbations, their role was clarified by Foglizzo [21,22]. Due to the isentropicity of the flow and the specific case of the turbulence [$v \cdot (\nabla \times (\nabla \times v)) = 0$ in the linear approximation] described below linear perturbations are absent.

In the first part, we formulate the basic equations of ideal steady-state axisymmetric hydrodynamics, which are known to be reduced to one second-order equation for the

stream function. Then, in the second part, the structure of the solitary curl is discussed in detail. Finally, in the third part we consider two toy models describing axisymmetric turbulence. It is shown that the turbulence changes mainly the effective gravity potential but not the effective pressure.

II. BASIC EQUATIONS

First of all, let us formulate basic hydrodynamical equations describing axisymmetric stationary flows using the spherical coordinate system. Then, as is well known (see classical textbooks [23,24]), it is convenient to introduce the potential $\Phi(r, \theta)$ connected with the poloidal velocity \mathbf{v}_p (which, by its definition, equals to $\vec{v} - \vec{v}_\varphi$) and the number density n as [25–27] [here and below we assume that all base vectors ($\vec{e}_r, \vec{e}_\theta, \vec{e}_\varphi$) are normalized to unity]

$$n\mathbf{v}_p = \frac{\nabla\Phi \times \mathbf{e}_\varphi}{2\pi r \sin\theta}. \quad (1)$$

This definition results in the following properties:

- (i) The continuity equation $\nabla \cdot (n\mathbf{v}) = 0$ is satisfied automatically.
- (ii) Multiplying both sides of Eq. (1) by the area element $d\mathbf{S} = \mathbf{e}_r r^2 \sin\theta d\theta d\varphi$ and integrating over the spherical element of radius r , strained on a circle that intersects the point (r, θ) , it is easy to verify that the potential $\Phi(r, \theta)$ has the meaning of the particle flux through the circle $r, \theta, 0 < \varphi < 2\pi$. In particular, the total flux through the surface of the sphere of radius r is $\Phi_{\text{tot}} = \Phi(r, \pi)$.
- (iii) As $\mathbf{v} \cdot \nabla\Phi = 0$, the velocity vectors \mathbf{v} are located on the surfaces $\Phi(r, \theta) = \text{const}$.

In this case, three conserved quantities for energy E_n , angular momentum L_n , and the entropy s can be formulated as

$$E_n = E_n(\Phi) = \frac{v^2}{2} + w + \varphi_g, \quad (2)$$

^{*}beskin@lpi.ru[†]lezhnin@phystech.edu

$$L_n = L_n(\Phi) = v_\varphi r \sin \theta, \quad (3)$$

$$s = s(\Phi). \quad (4)$$

Here w is the specific enthalpy, and φ_g is the gravitational potential.

In what follows we for simplicity consider the entropy $s(\Phi)$ to be constant. Then the equation for the stream function $\Phi(r, \theta)$ (which is no more than the projection of the Euler equation onto the axis perpendicular to the velocity vector \mathbf{v}) looks like (cf. Beskin [25,26], Heyvaerts [27] and classical textbooks [23,24])

$$\varpi^2 \nabla_k \left(\frac{1}{\varpi^2 n} \nabla^k \Phi \right) + 4\pi^2 n L_n \frac{dL_n}{d\Phi} - 4\pi^2 \varpi^2 n \frac{dE_n}{d\Phi} = 0, \quad (5)$$

where $\varpi = r \sin \theta$. This equation represents the balance of forces in a normal direction to flow lines. In particular, for spherically symmetric flow, i.e., for $E_n(\Phi) = \text{const}$, $L_n(\Phi) = 0$, it has the solution

$$\Phi = \Phi_0(1 - \cos \theta), \quad (6)$$

where Φ_0 is the positive-valued constant in case of ejection and negative-valued in case of accretion.

In the following, we deal with the linear angular operator

$$\hat{L}_\theta = \sin \theta \frac{\partial}{\partial \theta} \left(\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \right), \quad (7)$$

originated from Eq. (5). It has eigenfunctions

$$Q_0 = 1 - \cos \theta, \quad (8)$$

$$Q_1 = \sin^2 \theta, \quad (9)$$

$$Q_2 = \sin^2 \theta \cos \theta, \quad (10)$$

...

$$Q_m = \frac{2^m m! (m-1)!}{(2m)!} \sin^2 \theta \mathcal{P}'_m(\cos \theta), \quad (11)$$

and the eigenvalues

$$q_m = -m(m+1). \quad (12)$$

Here $\mathcal{P}_m(x)$ are the Legendre polynomials and the dash indicates their derivatives.

Here and further we use the standard approach when the small linear disturbances are analyzed on the background of analytical solution, in our case—on the background of the solution from Eq. (6). Let us consider now the small disturbance of the spherically symmetric flow, so that one can write down the flux function as

$$\Phi = \Phi_0[1 - \cos \theta + \varepsilon^2 f(r, \theta)] \quad (13)$$

with the small parameter $\varepsilon \ll 1$. Then Eq. (5) can be linearized, while the equation for the perturbation function $f(r, \theta)$ is written as [25]

$$\begin{aligned} & -\varepsilon^2 D \frac{\partial^2 f}{\partial r^2} - \frac{\varepsilon^2}{r^2} (D+1) \sin \theta \frac{\partial}{\partial \theta} \left(\frac{1}{\sin \theta} \frac{\partial f}{\partial \theta} \right) + \varepsilon^2 N_r \frac{\partial f}{\partial r} \\ & = -\frac{4\pi^2 n^2 r^2}{\Phi_0^2} \sin \theta (D+1) \frac{dE_n}{d\theta} \\ & + \frac{4\pi^2 n^2}{\Phi_0^2} (D+1) \frac{L_n}{\sin \theta} \frac{dL_n}{d\theta} - \frac{4\pi^2 n^2 \cos \theta}{\Phi_0^2 \sin^2 \theta} L_n^2. \end{aligned} \quad (14)$$

Here $D = -1 + c_s^2/v^2$, and $N_r = 2/r - 4\pi^2 n^2 r^2 GM/\Phi_0^2$. This equation allows us to seek the solution in the form

$$f(r, \theta) = \sum_{m=0}^{\infty} g_m(r) Q_m(\theta). \quad (15)$$

Introducing now dimensionless variables

$$x = \frac{r}{r_*}, \quad u = \frac{n}{n_*}, \quad l = \frac{c_s^2}{c_*^2}, \quad (16)$$

where the $*$ -values correspond to the sonic surface (which can be taken from the zero approximation), we can write the ordinary differential equations describing the radial functions $g_m(r)$:

$$\begin{aligned} & (1 - x^4 l u^2) g_m'' + 2 \left(\frac{1}{x} - x^2 u^2 \right) g_m' + m(m+1) x^2 l u^2 g_m \\ & = k_m \frac{R^2}{r_*^2} x^4 l u^4 - \lambda_m \frac{R^2}{r_*^2} u^2 - \sigma_m x^6 l u^4. \end{aligned} \quad (17)$$

Here $g'_m = dg_m(x)/dx$, $g''_m = d^2g_m(x)/dx^2$, and the expansion coefficients σ_m , λ_m and k_m depend on the disturbances as

$$\sin \theta \frac{dE_n}{d\theta} = \varepsilon^2 c_*^2 \sum_{m=0}^{\infty} \sigma_m Q_m(\theta), \quad (18)$$

$$\frac{\cos \theta}{\sin^2 \theta} L_n^2 = \varepsilon^2 c_*^2 r_*^2 \sum_{m=0}^{\infty} \lambda_m Q_m(\theta), \quad (19)$$

$$\frac{L_n}{\sin \theta} \frac{dL_n}{d\theta} = \varepsilon^2 c_*^2 r_*^2 \sum_{m=0}^{\infty} k_m Q_m(\theta). \quad (20)$$

Finally, the functions $l(x)$ and $u(x)$ correspond to the spherically symmetric flow. For the polytropic equation of state $P(n, s) = A(s)n^{\Gamma-1}$ we use here they are connected by the relation $l = u^{\Gamma-1}$. As to the dimensionless number density $u(x)$, it can be found from the ordinary differential equation

$$\frac{du}{dx} = -2 \frac{u(1-x^3u^2)}{x(1-x^4lu^2)} \quad (21)$$

with the boundary conditions

$$u(x)|_{x=1} = 1, \quad (22)$$

$$\left. \frac{du}{dx} \right|_{x=1} = -\frac{4 + \sqrt{10 - 6\Gamma}}{\Gamma + 1}. \quad (23)$$

The second boundary condition from Eq. (23) could be easily derived by L'Hopital expansion of (21). The choice of sign corresponds to accretion, opposite to the case of [28], where the process of ejection has been considered.

As for the boundary conditions for the system of Eq. (20), they are taken analogous to the case of Bondi-Hoyle accretion [25,28]:

$$\varepsilon^2 g_m(1) = \frac{(2m)!}{2^m(m+1)m!} \left[\frac{(\delta E_n)_m}{c_*^2} - \frac{(L_n^2/\sin^2\theta)_m}{2c_*^2 r_*^2} \right], \quad (24)$$

$$g'_m(R/r_*) = 0, \quad (25)$$

where $(\dots)_m$ stands for the expansion in terms of the Legendre polynomials, which can be found from energy and angular momentum integral perturbations.

III. SOLITARY CURL

Now let us consider in detail the internal structure of the quasispherical accretion with a small axisymmetric perturbation localized near the axis $\theta = \pi$. In other words, we suppose that the angular size of the vortex is small enough ($\delta\theta_{\text{curl}} \ll 1$). The main goal of this numerical calculation is to find the disturbance function $f(r, \theta)$ which gives us the possibility to determine velocity components v_θ and v_φ , i.e., the main characteristics of the perturbed flow.

To model the internal structure of the vortex, we determine θ -dependent angular velocity $\Omega(\theta)$ in the form

$$\Omega(\theta) = \Omega_0 \exp[-\alpha^2(1 + \cos\theta)]. \quad (26)$$

Here Ω_0 gives the amplitude of considered curl and the coefficient $\alpha \approx 10$ (which are free parameters of our model) is an inversed curl width. Certainly, we assume that the perturbation is small in comparison with the main contribution of the radial accretion.

Then the flow structure can be described by the system (17), (21)–(25) formulated in the previous section. As to the expansion coefficients k_m , λ_m and σ_m , they are to be determined from Eqs. (18)–(20) and (24) on the outer boundary of a flow $r = R$. For our choice (26) the disturbances have the form

$$\delta E_n(\theta) = \frac{\Omega_0^2 \exp[-2\alpha^2(1 + \cos\theta)] R^2 \sin^2\theta}{2}, \quad (27)$$

$$\delta L_n(\theta) = \Omega_0 \exp[-\alpha^2(1 + \cos\theta)] R^2 \sin^2\theta. \quad (28)$$

As one can easily check, it gives $\varepsilon = \Omega_0 R / c_*$.

Expansions (18)–(20) in terms of $Q_m(\theta)$ contain some numerical difficulties because the set of this functions is not an orthogonal one, and, even though it converges, in our case of very small curl width we can neglect just summands with numbers larger than 50. Even in some trivial cases like $\Omega(\theta) \sim (1 - \theta^2)$ these polynomials call a number of numerical obstacles (e.g., bad-conditioned matrix of linear equation for coefficients k_m , λ_m and σ_m , etc.).

In order to expand functions of integrals, we used the auxiliary set of Chebyshev polynomials, which is orthogonal and possesses a feature of generally faster convergence. Using these polynomials, we could find all expansions with the accuracy no worse than 10^{-3} . As was shown in Sec. II, the normalized density function $u(x)$ can be derived from Eq. (21) and boundary conditions (22) and (23). The results of numerical calculations for different polytropic indices Γ are shown in Fig. 1. In particular, as one can see, the density is nearly constant in the subsonic regime ($r \gg r_*$).

An example of the numerical calculation of perturbation function $f(r, \theta)$ is shown in Fig. 2. We should stress that $f(r, \theta)$ turns actually zero outside the small region near the axial curl. This statement allows us to assume as a zero approximation that the turbulent accretion regime containing a number of curls can be considered as a set of noninteracting ones.




FIG. 1. Dimensionless number density $u(x) = n/n_*$ in region $1 \leq x \leq R/r_* = 10$, Γ from 1.1 to 5/3.




FIG. 2. Normalized v_θ in region $1 \leq x \leq R/r_* = 10$, $0 \leq \theta \leq \pi$, $\Gamma = 4/3$.

Apart from the numerical solution, we can also find the analytical asymptotic solutions in the supersonic region $x \ll 1$ where Eq. (17) can be rewritten as

$$g_m'' + \frac{3}{2x} g_m' + \frac{m(m+1)}{2^{\Gamma+1}} x^{-(3\Gamma+1)/2} g_m + \lambda_m \frac{R^2}{r_*^2} \frac{1}{4x^3} = O(x^{-3}). \quad (29)$$

Here we take into account that $\Gamma < 5/3$. Getting rid of all parameters from the right part of this equation, one can introduce a new function $y_m = g_m / (\lambda_m R^2 / r_*^2)$. Then Eq. (29) can be rewritten as

$$y_m'' + \frac{3}{2x} y_m' + \frac{m(m+1)}{2^{\Gamma+1}} x^{-(3\Gamma+1)/2} y_m + \frac{1}{4x^3} = O(x^{-3}). \quad (30)$$

Neglecting now all terms which are proportional to $x^{-\nu}$, where $\nu > -3$, we obtain that this equation has a universal solution independent of the boundary conditions on the outer boundary $r = R$,

$$y(x) = -\frac{8}{x}. \quad (31)$$

Remember that the same asymptotic behavior was obtained by [11] for homogeneously rotating flow.

As was already stressed, numerical results allow us to determine the v_θ/v_φ ratio around the curl. It is easy to show that

$$\frac{v_\theta}{v_\varphi} = 2\sqrt{2}\varepsilon\pi \left(\frac{r_*}{R}\right)^{7/2} p(r, \theta), \quad (32)$$

where

$$p(r, \theta) = \frac{\sum_{m=0}^{\infty} g_m'(r) Q_m(\theta)}{u(r) \sin^2 \theta \exp[-\alpha^2(1 + \cos \theta)]}. \quad (33)$$

In our calculation we put $r_*/R = 0.1$, so that the function (33) is limited in area near the curl ($|p(r, \theta)| < 200$). Taking now into account that ε is a small parameter of our expansion and $|f(r, \theta)| < 20$, one can show that for reasonable parameter ε the ratio $|v_\theta/v_\varphi|$ in the area of vortex has an order of $\sim 10^{-5}$ (see Fig. 3). Thus, we could claim that $v_\theta \ll v_\varphi$, and then one can neglect all terms in Navier-Stokes equations that consist of v_θ .

In the same time, one can determine the v_φ/v_r ratio, which will be useful in further consideration. Taking into account (28) and deriving v_r for transonic flow from the zero order approximation, we get

$$\frac{v_\varphi}{v_r} = \frac{n}{n_*} \frac{r}{r_*} \frac{R}{r_*} \varepsilon \sin \theta \exp[-\alpha^2(1 + \cos \theta)]. \quad (34)$$




FIG. 3. v_θ/v_φ ratio in region $1 \leq x \leq 10$, $19\pi/20 \leq \theta \leq \pi$.

In the subsonic region $r_* \leq r$ the absolute value of this ratio has an order of 10^{-4} . On the other hand, in the supersonic region near the star ($r \approx r_{\text{star}} \ll r_*$, where r_{star} is a star radius) the value of (34) is approximately $(r_{\text{star}}/r_*)^{-1/2} (R/r_*) \varepsilon \sin \theta \exp(-\alpha^2(1 + \cos \theta))$, which sufficiently depends on the accretor radius r_{star} .

Thus, according to (31), we cannot use our solution in the limit $r \rightarrow 0$. Indeed, analyzing the field line equation $r d\theta/dr = v_\theta/v_r$, we obtain that the asymptotic solution survives until $\theta_0 \approx \delta\theta$, where $\theta_0 \ll 1$ is an initial angular size of a curl and $\delta\theta$ is a broadening parameter of the streamline. Deriving the θ -component of the velocity from (1) and taking v_r of zero order, we get

$$\frac{d\theta}{dr} \sim \varepsilon^2 \frac{\partial f / \partial r}{\sin \theta}. \quad (35)$$

Assuming now that $\theta \ll 1$, one can expand Eq. (35) in θ and neglecting all summands of m with $m \geq 2$, we find

$$\frac{d\theta}{dr} \sim \varepsilon^2 \theta \frac{\partial R^2}{\partial r r_*}. \quad (36)$$

This equation can be simply integrated, and we obtain

$$\ln \frac{\theta_0 + \delta\theta}{\theta_0} \sim \varepsilon^2 \frac{R^2}{r_* r}. \quad (37)$$

Hence, under the radius $r \sim \varepsilon^2 R^2 / r_*$ we cannot use the solution (31) as the disturbance becomes larger than unity. In order to keep the solution up to star surface $r = r_{\text{star}}$, we should demand

$$\varepsilon^2 R^2 / r_* < r_{\text{star}}. \quad (38)$$

It gives us the general condition of applicability of the approach described above, unless we cannot use the method of linear expansion of the Grad-Shafranov equation, and the turbulent flow is to be described in another

way which lies outside the consideration of the current paper [20].

IV. TWO TOY MODELS

Let us suppose now that the turbulence in the accreting matter can be described by the large number of axisymmetric vortices with different parameters Ω_0 and α filling all the accreting volume. Within this approach one can construct two toy analytical models demonstrating how the turbulence can affect the structure of the spherically symmetric accretion.

A. Inviscid flow

The first model in which we neglect viscosity corresponds to classical ideal spherically symmetric Bondi accretion [3]. In this case one can consider the following system of equations:

$$\nabla(n\mathbf{v}) = 0, \quad (39)$$

$$(\mathbf{v} \cdot \nabla)\mathbf{v} = -\frac{\nabla P}{\rho} - \nabla\varphi_g, \quad (40)$$

$$(\mathbf{v} \cdot \nabla)s = 0. \quad (41)$$

Following Bondi [3] we consider the polytropic equation of state $P = P(n, s) = k(s)n^\Gamma$ resulting in polytropic index $\Gamma \neq 1$:

$$c_s^2 = \frac{\Gamma k}{m_p} n^{\Gamma-1}, \quad (42)$$

$$w = \frac{c_s^2}{\Gamma - 1}, \quad (43)$$

$$T = \frac{m_p}{\Gamma} c_s^2. \quad (44)$$

Here again n (1/cm³) is the number density, m_p (in g) is the mass of particles ($\rho = m_p n$ is the mass density), s is the entropy per one particle (dimensionless), w (in cm²/s²) is the specific enthalpy, T (in erg) is the temperature in energy units, and, finally, c_s (cm/s) is the sound velocity.

As was demonstrated above, for a weak enough turbulence level (38) for any individual curl one can neglect the θ -component of the velocity perturbation in comparison with the toroidal one v_φ up to the central body $r = r_{\text{star}}$. Thus, in zero approximation one can put $v_\theta = 0$, i.e., $\theta = \text{const}$. This implies that, according to the angular momentum conservation law $r \sin\theta v_\varphi = \text{const}$, we can write down

$$v_\varphi(r, \theta) = \Omega(\theta) \frac{R^2}{r} \sin\theta. \quad (45)$$

Here $\Omega(\theta)$ is a smooth function of θ that can be approximately described as

$$\Omega(\theta) \approx \begin{cases} \Omega_0, & \pi - \alpha^{-1} < \theta < \pi, \\ 0 & 0 < \theta < \pi - \alpha^{-1}. \end{cases} \quad (46)$$

In order to find the characteristic values of the accretion flow we have to use energy and momentum integrals conserving on streamlines. Taking into account an assertion $v_r \gg v_\varphi \gg v_\theta$, we can neglect the θ -component of velocity which gives

$$E_n(\theta) = \frac{v_r^2(r)}{2} + \omega(r) + \varphi_g(r) + \frac{L^2(\theta)}{2r^2 \sin^2\theta}, \quad (47)$$

$$L_n(\theta) = v_\varphi r \sin\theta = \Omega(\theta) r^2 \sin^2\theta. \quad (48)$$

Averaging now these integrals in θ and introducing a new value

$$L_{\text{av}}^2 \equiv \left\langle \frac{L^2(\theta)}{\sin^2\theta} \right\rangle \quad (49)$$

we obtain for the averaged energy integral

$$E_{\text{av}} \equiv \langle E_n(\theta) \rangle = \frac{v_r^2(r)}{2} + \omega(r) + \underbrace{\varphi_g(r)}_{\varphi_{\text{eff}}(r)} + \frac{L_{\text{av}}^2}{2r^2}. \quad (50)$$

As we see, the two last terms can be considered as an effective gravitational potential, as has also been proposed in a number of papers [29,30].

Further calculations are quite similar to the classical Bondi problem for the spherical flow. In other words, using another integrals of motion, i.e., the total particle flux

$$\Phi = 4\pi r^2 n(r) v_r(r) = \text{const}, \quad (51)$$

and the entropy s , one can rewrite the energy integral (50) as

$$E_{\text{av}} = \frac{\Phi^2}{32\pi^2 n^2 r^4} + \frac{\Gamma k(s) n^{\Gamma-1}}{\Gamma - 1} \frac{1}{m_p} - \frac{GM}{r} + \frac{L_{\text{av}}^2}{2r^2}. \quad (52)$$

It gives the following expression for the logarithmic r -derivative of the number density

$$\eta_1 = \frac{r}{n} \frac{dn}{dr} = \frac{2 - \frac{GM}{v_r^2 r} + \frac{L_{\text{av}}^2}{v_r^2 r^2}}{-1 + \frac{c_s^2}{v_r^2}}. \quad (53)$$

As for Bondi accretion, this derivative has a singularity on the sonic surface $v_r = c_s = c_*$. This implies that for a smooth transition through the sonic surface $r = r_*$, the additional condition is to be satisfied:

$$2 - \frac{GM}{c_*^2 r_*} + \frac{L_{\text{av}}^2}{c_*^2 r_*^2} = 0. \quad (54)$$

Solving now (54) in terms of r_* in this approximation, we find

$$r_* = \frac{GM}{2c_*^2} \left(1 - \frac{4L_{\text{av}}^2 c_*^2}{G^2 M^2} \right), \quad (55)$$

$$c_* = \sqrt{\frac{2}{5-3\Gamma} c_\infty} \left(1 + \frac{12(\Gamma-1)L_{\text{av}}^2 c_\infty^2}{(5-3\Gamma)^2 G^2 M^2} \right), \quad (56)$$

where c_∞ is evaluated from

$$E_{\text{av}} = \frac{c_\infty^2}{\Gamma-1}. \quad (57)$$

Accordingly, we obtain for r_*/r_{*B} and c_*/c_{*B} ratios

$$\frac{r_*}{r_{*B}} = 1 - \frac{16}{(5-3\Gamma)^2} \frac{L_{\text{av}}^2 c_\infty^2}{G^2 M^2}, \quad (58)$$

$$\frac{c_*}{c_{*B}} = 1 + \frac{12(\Gamma-1)}{(5-3\Gamma)^2} \frac{L_{\text{av}}^2 c_\infty^2}{G^2 M^2}, \quad (59)$$

where c_{*B} and r_{*B} correspond to the classical Bondi accretion.

The presence of vorticity not only changes sonic surface parameters, but also effectively diminishes accretion rate:

$$\frac{\dot{M}}{\dot{M}_B} = \frac{r_*^2 c_*}{r_{*B}^2 c_{*B}} = 1 - \frac{4(11-4\Gamma)}{(5-3\Gamma)^2} \frac{L_{\text{av}}^2 c_\infty^2}{G^2 M^2} \quad (60)$$

in our order of precision. The similar expression for Bondi-Hoyle accretion can be found in Krumholz *et al.* [31].

Finally, using the definition (49) for L_{av}^2 , we can rewrite our criteria of the applicability (38) as $L_{\text{av}}^2 \ll GM r_{\text{star}}$. As $r_{\text{star}} < r_*$, it can be finally rewritten as

$$L_{\text{av}} \ll \frac{GM}{c_*}. \quad (61)$$

To sum up, one can conclude that the nonzero angular momentum effectively decreases the gravitational force. In other words, the presence of the angular momentum does not allow matter to fall down as easy as in its absence. Roughly speaking, we substitute our gravitating center with one that possesses less mass. So, in the case of Bondi accretion with a small angular momentum perturbation we should modify the relations for sonic surface radius and velocity—they decrease and rise respectively [32]. It is important to note that we can consider the turbulent accretion regime as one with a modified gravity potential.

B. Viscous flow

In this subsection we consider stationary axisymmetric quasispherical flow of viscous fluid. Using the condition $|v_r| \gg |v_\phi| \gg |v_\theta|$, and neglecting all the terms containing v_θ , we obtain for the ϕ -component of the Euler equation [33]

$$v_r \frac{\partial v_\phi}{\partial r} + \frac{v_r v_\phi}{r} = \nu \left(\nabla^2 v_\phi - \frac{v_\phi}{r^2 \sin^2 \theta} \right). \quad (62)$$

For viscous flow it is convenient to determine the toroidal component of the velocity v_ϕ as

$$v_\phi = \Omega(r, \theta) r \sin \theta, \quad (63)$$

where we will use the following form for the angular velocity $\Omega(r, \theta)$:

$$\Omega(r, \theta) = \Omega_0(r) \exp \left[-\frac{\theta^2}{2\delta(r)} \right]. \quad (64)$$

Here $\Omega_0 = \Omega_0(r)$ is an amplitude, and $\delta = \delta(r)$ is a square of effective angular width of an individual curl. Substituting now v_ϕ into Eq. (62), we obtain

$$\dot{M} \frac{d}{dr} (\Omega r^2 \sin \theta) = \frac{4\pi r^2 \eta}{\sin^2 \theta} \frac{d}{d\theta} \left[\sin^3 \theta \frac{d\Omega}{d\theta} \right], \quad (65)$$

where

$$\dot{M} = 4\pi r^2 \rho v_r \quad (66)$$

is the accretion rate remaining constant in stationary flow, and $\eta = \rho\nu$ is a dynamic viscous coefficient which can be considered as a constant as well [33].

Using now Eq. (65), one can easily show that the total angular momentum of an individual vortex conserves ($dL/dr = 0$). Indeed, internal friction connecting with viscosity cannot change the total angular momentum of the accreting matter. For this reason, together with (66), the angular momentum

$$dL = \rho \Omega r^2 \sin^2 \theta d\phi \sin \theta d\theta r^2 dr \quad (67)$$

can be rewritten as a full θ -derivative. This implies that the right-hand side of Eq. (67) integrated over θ becomes zero.

Further, to determine the radial dependence of the curl amplitude $\Omega_0(r)$ and the squared width $\delta(r)$, we substitute the angular velocity $\Omega_0(r, \theta)$ (64) into ([33]) and expand it in terms of θ near the axis, neglecting all the terms with the power more than 3. As a result, we obtain two equations for $\Omega_0(r)$ and $\delta(r)$:

$$\frac{\dot{M}}{2\pi\eta} r \Omega_0(r) + \frac{4r^2 \Omega_0(r)}{\delta(r)} + \frac{\dot{M}}{4\pi\eta} r^2 \Omega_0'(r) = 0, \quad (68)$$

$$\begin{aligned}
& -18r^2\Omega_0(r) - \frac{3\dot{M}}{2\pi\eta} r\delta(r)\Omega_0(r) - 10r^2\delta(r)\Omega_0(r) \\
& - \frac{\dot{M}}{2\pi\eta} r\delta^2(r)\Omega_0(r) + \frac{3\dot{M}}{4\pi\eta} r^2\delta'(r)\Omega_0(r) \\
& - \frac{\dot{M}}{4\pi\eta} r^2\delta^2(r)\Omega_0'(r) = 0,
\end{aligned} \tag{69}$$

which have simple solutions

$$\delta(r) = \delta_0 + \frac{8\pi\eta(r-r_0)}{\dot{M}}, \tag{70}$$

$$\Omega_0(r) = \Omega_0 \frac{r_0^2}{r^2} \left[\frac{8\pi\eta(r-r_0)}{\dot{M}\delta_0} + 1 \right]^{-2}. \tag{71}$$

Introduction of a small vortex turbulence can be again treated by modifying a gravitational potential as

$$\varphi_{\text{eff}} = -\frac{GM}{r} + \frac{\Omega_0^2 r_0^4 \delta_0}{4\pi r^2} \left[1 + \frac{8\pi\eta}{\dot{M}\delta_0} (r-r_0) \right]^{-3}. \tag{72}$$

Thus, viscosity results in increasing of the vortex width ($\delta' < 0$ for $\dot{M} < 0$ corresponding to accretion) and diminishing of the angular rotation. On the other hand, the role of viscosity will be small if

$$\frac{8\pi\eta r_0}{|\dot{M}|\delta_0} \ll 1. \tag{73}$$

For $\eta \rightarrow 0$ we return to the previous result $\delta(r) = \text{const}$, $\Omega_0(r) \propto r^{-2}$. Introducing now the Reynolds number as

$$\text{Re} = \frac{\rho v l}{\eta}, \tag{74}$$

where ρ , v and $l = r_0\delta^{1/2}$ are characteristic values of a flow and using expression (66), we can rewrite (73) as

$$\text{Re} = \frac{|\dot{M}|}{4\pi r_0\eta} \gg \delta_0^{-1/2}. \tag{75}$$

This implies that the role of viscosity is small for turbulent flow.

Certainly, the analysis presented above allows us to take into consideration only an isolated set of curls. In reality, the dense cellular turbulent structure possesses a number of collective effects [34], which is to be described in another way. The easiest method to proceed with the minimal number of additional assumptions is to choose another angular velocity profile.

As the total angular momentum of the accreting matter is supposed to be zero, we will use the following expression for the angular velocity:




FIG. 4. Ratio $\Omega/\Omega_0 = F(\theta)$ for $0 \leq \theta \leq \pi/30$.

$$\Omega(r, \theta) = \Omega_0(r) \exp\left(-\frac{\theta^2}{2\delta}\right) \left(1 - \frac{\gamma^2}{2\delta}\theta^2\right). \tag{76}$$

Here the parameter γ is to be chosen from the condition of the zero total angular momentum

$$\int_{|\vec{r}| \leq R} dL = 0, \tag{77}$$

which is equivalent to

$$\int_0^\pi d\theta \sin^3\theta \Omega(r, \theta) = 0. \tag{78}$$

One of its realizations can be seen in Fig. 4 where the dashed line shows zero angular velocity level. A configuration like this one represents the unit of cellular turbulent structure, fulfilling the main requirements of its nature. To simplify our calculations, we hold γ and δ on constant values in order to get a simple equation on $\Omega_0(r)$. Again, we expand Eq. (65) in terms of θ to the first order. As a result, we obtain for $\Omega_0(r)$

$$\Omega_0(r) = \Omega_0 \left(\frac{r_0}{r}\right)^2 \exp\left[\frac{16\pi\eta(1+\gamma^2)}{\dot{M}\delta}(r_0-r)\right]. \tag{79}$$

In this case, the effective gravitational potential cannot be derived for arbitrary parameters without special functions. It can be written as

$$\varphi_{\text{eff}} = -\frac{GM}{r} + C \cdot \frac{\Omega_0^2 r_0^4}{r^2} \exp\left[-\frac{16\pi\eta(1+\gamma^2)}{|\dot{M}|\delta}(r_0-r)\right], \tag{80}$$

where

$$C = \frac{1}{2\pi} \int_0^\pi d\theta \exp(-\theta^2/\delta) \left(1 - \frac{\gamma^2}{2\delta}\theta^2\right)^2 \sin^2\theta. \tag{81}$$

Choosing, for instance, $\gamma = 1/\sqrt{2}$ and $\delta = 10^{-4}$, it gives $C \approx 3.4 \times 10^{-8}$.

The expression under the exponential function in (79) is always lower than zero, so the criterion of the importance of viscosity effects can be formulated as

$$\frac{16\pi\eta(1+\gamma^2)}{|\dot{M}|\delta}r_0 \gg 1, \quad (82)$$

which is more convenient to discuss in terms of Reynolds number

$$\delta^{-1/2} \gg \text{Re}. \quad (83)$$

Thus, for narrow vortex (i.e., for $\delta < 10^{-2}$) the viscosity effects must be taken into account.

V. DISCUSSION AND CONCLUSION

We have considered the dynamics of a solitary axisymmetric vortex against the background of Bondi accretion flow. It was shown that if condition (38) is not satisfied, the level of turbulence is high enough and the flow cannot be considered as radial [35–37]. It is necessary to stress that

we consider the special case of the turbulence which is nontrivial from the second order of the expansion only.

Scott and Lovelace [29] have already proposed an idea of the inclusion of vortex terms into the effective gravitational potential. In this paper the same approach is discussed along with the impact of the vorticity on the quasispherically symmetric accretion.

Further, we described two analytical toy models that show how the turbulence affects the structure of the spherically symmetric flow. In particular, it was shown that the sonic surface moves inwards because of effective diminishing of gravitational force. Finally, a criterion to analyze the importance of viscosity effects in the adiabatic flow filled either by isolated or dense set of curls was formulated.

ACKNOWLEDGMENTS

We thank K.P. Zybin and T. Foglizzo for fruitful discussions. This work was partially supported by Russian Foundation for Basic Research (Grant No. 15-02-03063). K. V. L. is grateful to ELI-Beamlines project for hospitality during the final stages of this work.

-
- [1] V. M. Lipunov, *Astrophysics of Neutron Stars* (Springer, Heidelberg, 1992).
 - [2] S. Shapiro and S. Teukolsky, *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact Objects* (Wiley, New York, 1983).
 - [3] H. Bondi, *Mon. Not. R. Astron. Soc.* **112**, 195 (1952).
 - [4] H. Bondi and F. Hoyle, *Mon. Not. R. Astron. Soc.* **104**, 273 (1944).
 - [5] R. Hunt, *Mon. Not. R. Astron. Soc.* **188**, 83 (1979).
 - [6] L. I. Petrich, S. Shapiro, R. F. Stark, and S. Teukolsky, *Astrophys. J.* **336**, 313 (1989).
 - [7] M. Ruffert and D. Arnett, *Astrophys. J.* **427**, 351 (1994).
 - [8] O. D. Toropina, M. M. Romanova Yu. M. Toropin, and R. V. E. Lovelace, *Astrophys. J.* **593**, 472 (2003).
 - [9] O. D. Toropina, M. M. Romanova, and R. V. D. Lovelace, *Mon. Not. R. Astron. Soc.* **420**, 810 (2012).
 - [10] M. Anderson, *Mon. Not. R. Astron. Soc.* **239**, 19 (1989).
 - [11] V. S. Beskin and L. M. Malyshekin, *Astron. Lett.* **22**, 475 (1996).
 - [12] V. S. Beskin and Yu. N. Pidoprygora, *J. Exp. Theor. Phys.* **80**, 575 (1995).
 - [13] G. S. Bisnovatyi-Kogan, Ya. M. Kazhdan, A. A. Klypin *et al.*, *Sov. Astron.* **23**, 201 (1979).
 - [14] V. I. Pariev, *Mon. Not. R. Astron. Soc.* **283**, 1264 (1996).
 - [15] L. I. Petrich, S. Shapiro, and S. Teukolsky, *Phys. Rev. Lett.* **60**, 1781 (1988).
 - [16] W. I. Axford and R. C. Newman, *Astrophys. J.* **147**, 230 (1967).
 - [17] S. A. Balbus and J. F. Hawley, *Astrophys. J.* **376**, 214 (1991).
 - [18] A. Brandenburg and D. D. Sokoloff, *Geophys. Astrophys. Fluid Dyn.* **96**, 319 (2002).
 - [19] J. Krolik and J. F. Hawley, *Astrophys. J.* **573**, 754 (2002).
 - [20] I. G. Kovalenko and M. A. Eremin, *Mon. Not. R. Astron. Soc.* **298**, 861 (1998).
 - [21] T. Foglizzo, *Astron. Astrophys.* **368**, 311 (2001).
 - [22] T. Foglizzo, *Astron. Astrophys.* **392**, 353 (2002).
 - [23] K. G. Guderley, *Theory of Transonic Flow* (Pergamon, Oxford, 1962).
 - [24] R. Mises, *Mathematical Theory of Compressible Fluid Flow* (Academic Press, New York, 1958).
 - [25] V. S. Beskin, *MHD Flows in Compact Astrophysical Objects* (Springer, Berlin, 2010).
 - [26] V. S. Beskin, *Phys. Usp.* **53**, 1199 (2010).
 - [27] J. Heyvaerts, in *Plasma Astrophysics*, edited by C. Chiuderi and G. Einaudi (Springer, Berlin, 1996), p. 31.
 - [28] V. S. Beskin and Yu. N. Pidoprygora, *Astron. Rep.* **42**, 71 (1998).
 - [29] H. A. Scott and R. V. E. Lovelace, *Astrophys. J.* **252**, 765 (1982).
 - [30] L. S. Sparke and F. H. Shu, *Astrophys. J.* **241**, L65 (1980).
 - [31] M. R. Krumholz, C. F. McKee, and R. I. Klein, *Astrophys. J.* **618**, 757 (2005).
 - [32] J. K. Bhattacharjee and A. K. Ray, *Astrophys. J.* **627**, 368 (2005).
 - [33] L. D. Landau and E. M. Lifshits, *Fluid Mechanics* (Pergamon Press, Oxford, 1987).
 - [34] M. E. Prokhorov, S. B. Popov, and A. V. Khoperskov, *Astron. Astrophys.* **381**, 1000 (2002).
 - [35] N. I. Shakura, K. A. Postnov, A. Yu. Kochetkova, and L. Yalmsdotter, *Phys. Usp.* **183**, 337 (2013).
 - [36] N. I. Shakura, K. A. Postnov, A. Yu. Kochetkova, and L. Yalmsdotter, *Mon. Not. R. Astron. Soc.* **420**, 216 (2012).
 - [37] R. V. Shcherbakov, *Astrophys. J. Suppl. Ser.* **177**, 493 (2008).