

de Sitter supergravity model buildingRenata Kallosh^{1,*} and Timm Wrase^{2,†}¹*SITP and Department of Physics, Stanford University, Stanford, California 94305, USA*²*Institute for Theoretical Physics, TU Wien, A-1040 Vienna, Austria*

(Received 17 September 2015; published 5 November 2015)

We present the explicit de Sitter supergravity action describing the interaction of supergravity with an arbitrary number of chiral and vector multiplets as well as one nilpotent chiral multiplet. The action has a non-Gaussian dependence on the auxiliary field of the nilpotent multiplet; however, it can be integrated out for an arbitrary matter-coupled supergravity. The general supergravity action with a given Kähler potential K , superpotential W and vector matrix f_{AB} interacting with a nilpotent chiral multiplet consists of the standard supergravity action defined by K , W and f_{AB} where the scalar in the nilpotent multiplet has to be replaced by a bilinear combination of the fermion in the nilpotent multiplet divided by the Gaussian value of the auxiliary field. All additional contributions to the action start with terms quartic and higher order in the fermion of the nilpotent multiplet. These are given by a simple universal closed form expression.

DOI: 10.1103/PhysRevD.92.105010

PACS numbers: 04.65.+e

I. INTRODUCTION

A supergravity action, including fermion interactions, with nonlinearly realized spontaneously broken local supersymmetry was derived in [1,2]. In the case of pure supergravity without scalars it was shown in [1] that de Sitter (dS) vacua form a simple set of classical solutions of this theory. For a long time only anti-de Sitter supergravity without scalars was known [3]. De Sitter vacua are also natural for the theory with a single chiral multiplet, constructed in [2]. The presence of a Volkov-Akulov (VA) type nonlinearly realized supersymmetry [4] in supergravity models [1,2] leads to a natural uplifting of the vacuum energy in all these models. This is in full agreement with the string theory realization of the KKLT de Sitter vacua that use an anti-D3 brane [5]. The construction in [5] was recently described as a supersymmetric realization of the KKLT uplifting with account of the fermions on the world-volume of the anti-D3 brane [6]. Furthermore, it was shown in [7] how this setup can be realized in explicit warped string compactifications.

The purpose of this paper is to derive general and explicit supergravity models with chiral and vector multiplets interacting with a nilpotent chiral multiplet based on the superconformal formulation of this theory given in [8]. This strategy was already used successfully in [1,2] where full actions with fermions in models with a nilpotent multiplet and without matter multiplets or with a single chiral multiplet were derived. Meanwhile in [9] it was proposed how to derive supergravities with a nilpotent multiplet for general classes of models with any number of chiral and vector multiplets and with generic K , W and f_{AB} . It was shown that one can use the same method as in [1], namely to perform a non-Gaussian

integration of the auxiliary field resulting in a closed form action. A complete action in the unitary gauge was presented in [9] for general matter coupling. Here we will derive the complete action with fermions and with local supersymmetry for general matter coupling.

The general action without a nilpotent multiplet is well known and we will use here the framework presented in [10]. Originally the corresponding general supergravity-Yang-Mills action was derived from the superconformal theory in [11–13].

The interest in a nilpotent chiral multiplet satisfying the nilpotency condition in supergravity in applications to cosmology was initiated in [14] for the VA-Starobinsky model. For a general supergravity, interacting with the VA model, the superconformal construction was presented in [8]. It has been shown in [15] that the approach of using the constrained curvature superfield is dual to the VA model coupled to supergravity. It was also shown recently that one can use a complex linear goldstino superfield and build de Sitter supergravity [16].

This interest in these new supergravity models was increased by the fact that they facilitate the construction of early universe inflationary models compatible with the data and the construction of de Sitter vacua for explaining dark energy and supersymmetry breaking, see for example [17] and references therein. Such supergravity models in application to cosmology require mostly the knowledge of the bosonic action of the theory, where the rules are very simple: for the complete models one has to construct the standard supergravity action defined by K , W and f_{AB} . Once the action is known, one has to take only the bosonic part of it and, moreover, set the scalar field in the nilpotent multiplet to zero since the scalar in the nilpotent multiplet, representing the VA theory, is a fermion bilinear. If the scalar in the nilpotent multiplet is z^1 , then the rule for obtaining the bosonic action is

*kallosh@stanford.edu

†timm.wrase@tuwien.ac.at

$$\begin{aligned}
 & e^{-1} \mathcal{L}_{\text{bosonic}} \\
 &= e^{-1} \mathcal{L}^{\text{book}} [K(z^\alpha, \bar{z}^{\bar{\alpha}}), W(z^\alpha), f_{AB}(z^\alpha)] \Big|_{z^1 = \chi^\alpha = \bar{\chi}^{\bar{\alpha}} = \lambda^A = \psi_\mu = 0}, \\
 & \quad \alpha = 1, \dots, n,
 \end{aligned} \tag{1.1}$$

for a given choice of Kähler potential, superpotential and vector metric. Here $\mathcal{L}^{\text{book}}$ is the standard supergravity action for a given choice of K , W and f_{AB} ; see also the next section for details.

However, the knowledge of the bosonic action may not be sufficient for studies of reheating of the universe in these models, as well as for studies of particle physics, where the role of the fermions is important. A complete action including fermions, with the bosonic part given in (1.1), will be derived here, following the proposal in [9].

II. SUPERGRAVITY ACTION WITH A NILPOTENT MULTIPLET

The general supergravity Lagrangian for an arbitrary number of chiral and vector multiplets coupled to supergravity is given for example in the book [10] in Eqs. (18.6)–(18.19) and takes the form

$$e^{-1} \mathcal{L}^{\text{book}} = \mathcal{L}_{\text{kin}} - V + \mathcal{L}_{\text{m}} + \mathcal{L}_{\text{mix}} + \mathcal{L}_{4f}, \tag{2.1}$$

where \mathcal{L}_{kin} contains all the terms with spacetime derivatives of the fields. All terms on the right-hand side of Eq. (2.1) are defined in [10]. The Lagrangian is a function of all the physical fields $\mathcal{L}^{\text{book}} = \mathcal{L}^{\text{book}}(e_\mu^\alpha, \psi_\mu, z^\alpha, \bar{z}^{\bar{\alpha}}, \chi^\alpha, \bar{\chi}^{\bar{\alpha}}, A_\mu^A, \lambda^A)$.

We are interested in the case in which one of the chiral superfields in the supergravity action is nilpotent. Without loss of generality we choose the first one which leads to the constraint [1]: $z^1 = \frac{(\chi^1)^2}{2F^1}$ (and likewise $\bar{z}^{\bar{1}} = \frac{(\bar{\chi}^{\bar{1}})^2}{2\bar{F}^{\bar{1}}}$).¹ What is the corresponding Lagrangian in this case? Clearly we cannot just plug $z^1 = \frac{(\chi^1)^2}{2F^1}$ into the on-shell Lagrangian (2.1) since we do not know what F^1 is in the general supergravity model. What we actually need to do is to start with an off-shell supergravity Lagrangian and then integrate out all the F^α to get the correct on-shell Lagrangian.

The corresponding off-shell supergravity action for an arbitrary number of chiral and vector multiplets coupled to gravity is given by

$$e^{-1} \mathcal{L}_{\text{off-shell}} = (F^\alpha - F_G^\alpha) g_{\alpha\bar{\alpha}} (\bar{F}^{\bar{\alpha}} - \bar{F}_G^{\bar{\alpha}}) + e^{-1} \mathcal{L}^{\text{book}}, \tag{2.2}$$

where

$$F_G^\alpha = -e^{\frac{\kappa}{2}} g^{\alpha\bar{\beta}} \bar{\nabla}_{\bar{\beta}} \bar{W} + (F_G^\alpha)^f, \tag{2.3}$$

with

$$(F_G^\alpha)^f = \frac{1}{2} \Gamma_{\beta\gamma}^\alpha \bar{\chi}^\beta \chi^\gamma + \frac{1}{4} \bar{f}_{AB\bar{\beta}} g^{\bar{\beta}\alpha} \bar{\lambda}^A P_R \lambda^B. \tag{2.4}$$

Alternatively, we could have started with the off-shell supergravity action in [12], eqs. (25)–(33), which is given in the form with all auxiliary fields not integrated out. This set up was used in [2]. When the auxiliary fields A_μ and F^0 are integrated out, the remaining action still has all auxiliary fields F^α as shown in our Eq. (2.2).

Now since the Lagrangian $\mathcal{L}^{\text{book}}$ does depend on $z^1 = \frac{(\chi^1)^2}{2F^1}$ it will also depend on F^1 and therefore it seems prohibitively difficult to explicitly integrate it out. However, using a proposal made in [9] which includes a simplifying assumption about the form of Kähler potential we will derive the explicit on-shell Lagrangian in the next section. In particular, we assume that the Kähler potential depends only on the product $z^1 \bar{z}^{\bar{1}}$ so that we have²

$$K(z^i, \bar{z}^{\bar{i}}, z^1 \bar{z}^{\bar{1}}) = K_0(z^i, \bar{z}^{\bar{i}}) + z^1 \bar{z}^{\bar{1}} g_{1\bar{1}}(z^i, \bar{z}^{\bar{i}}). \tag{2.5}$$

We also expand the holomorphic superpotential W and the holomorphic gauge kinetic function f_{AB} as follows,

$$\begin{aligned}
 W(z^1, z^i) &= W_0(z^i) + z^1 W_1(z^i) \equiv g(z^i) + z^1 f(z^i), \\
 f_{AB}(z^1, z^i) &= f_{AB0}(z^i) + z^1 f_{AB1}(z^i).
 \end{aligned} \tag{2.6}$$

With this assumption we will integrate out F^1 in Sec. III. This involves taking into account a non-Gaussian dependence of the action on the auxiliary field F^1 . This gives us the generic supergravity Lagrangian for a nilpotent chiral superfield coupled to an arbitrary number of chiral and vector multiplets:

$$\boxed{e^{-1} \mathcal{L}_{\text{final}} = [e^{-1} \mathcal{L}^{\text{book}}]_{z^1 = \frac{(\chi^1)^2}{2F^1}} - \frac{(\chi^1)^2 (\bar{\chi}^{\bar{1}})^2}{4g_{1\bar{1}} (f^1 \bar{f}^{\bar{1}})^2} \left| g_{1\bar{1}} \frac{\square (\chi^1)^2}{2f^1} + B^1 \right|^2} \tag{2.7}$$

¹We are using the conventions of [10] but we will set $\kappa = 1$ and we define the short-hand notations $(\chi^1)^2 = \bar{\chi}^{\bar{1}} P_L \chi^1$ and $(\bar{\chi}^{\bar{1}})^2 = \bar{\chi}^{\bar{1}} P_R \chi^{\bar{1}}$.

²Since $(z^1)^2 = (\bar{z}^{\bar{1}})^2 = 0$ the only other term that could arise in the fully general Kähler potential is $z^1 f(z^i, \bar{z}^{\bar{i}}) + \bar{z}^{\bar{1}} \bar{f}(z^i, \bar{z}^{\bar{i}})$. Note that linear terms in $z^1 (\bar{z}^{\bar{1}})$ that are multiplied by a holomorphic (anti-holomorphic) function can be removed by a Kähler transformation.

with

$$f^1 \equiv \frac{1}{g_{1\bar{1}}} \left(-e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}} + \frac{1}{4} \bar{f}_{AB\bar{1}} \bar{\lambda}^A P_R \lambda^B \right),$$

$$B^1 \equiv e^{-1} \frac{\delta \mathcal{L}^{\text{book}}}{\delta \bar{z}^{\bar{1}}} \Big|_{z^1 = \bar{z}^{\bar{1}} = 0}. \quad (2.8)$$

Note that the last term in Eq. (2.7) has already the maximal power of the (undifferentiated) spinor χ^1 so that we can drop all terms in B^1 that contain χ^1 or $\chi^{\bar{1}}$. We explicitly give the relevant part of B^1 for two examples in Sec. IV.

One can see that the complete Lagrangian (2.7) has the original form with z^1 replaced by its ‘Gaussian value’ and there are some additional terms that are at least quadratic in $\chi^1 \chi^{\bar{1}}$. On the basis of this expression we may conclude that the action with K , W and f_{AB} generic (but with K dependence on $z^1 \bar{z}^{\bar{1}}$) is given by the standard action with the following modifications:

After the standard action for K , W and f_{AB} is presented as a function of z^1 and $\bar{z}^{\bar{1}}$

- (1) Replace z^1 by $z^1 = \frac{(\chi^1)^2}{2F^1}$ and likewise for the conjugate $\bar{z}^{\bar{1}} = \frac{(\chi^{\bar{1}})^2}{2\bar{F}^{\bar{1}}}$.
- (2) Add the quartic and higher order in spinors terms shown in the second entry in Eq. (2.7), where B^1 is given in (2.8) and more explicitly for two examples in Sec. IV.

III. DERIVATION OF THE MATTER-COUPLED SUPERGRAVITY ACTION WITH A NILPOTENT MULTIPLET

We start with the off-shell action (2.2) and we would like to integrate out the auxiliary fields F^α . The dependence on F^1 is non-Gaussian since the Lagrangian depends on $z^1 = \frac{(\chi^1)^2}{2F^1}$. However, we can still trivially integrate out all the other F^i , $i = 2, \dots, n$ and get

$$g_{i\bar{1}}(\bar{F}^{\bar{1}} - \bar{F}_G^{\bar{1}}) + g_{i\bar{j}}(\bar{F}^{\bar{j}} - \bar{F}_G^{\bar{j}}) = 0. \quad (3.1)$$

The submatrix $g_{i\bar{j}}$ is invertible in order to have a non-degenerate kinetic term for the z^i . Thus, we find

$$(\bar{F}^{\bar{j}} - \bar{F}_G^{\bar{j}}) = -(g_{ji})^{-1} g_{i\bar{1}}(\bar{F}^{\bar{1}} - \bar{F}_G^{\bar{1}}), \quad (3.2)$$

and the Lagrangian after integrating out the F^i becomes

$$e^{-1} \mathcal{L}_{\text{off-shell}} = (F^1 - F_G^1)(g_{1\bar{1}} - g_{1\bar{j}}(g_{ji})^{-1} g_{i\bar{1}})(\bar{F}^{\bar{1}} - \bar{F}_G^{\bar{1}}) + e^{-1} \mathcal{L}^{\text{book}}. \quad (3.3)$$

Here $e^{-1} \mathcal{L}^{\text{book}}$ is defined as an expansion in z^1 , $\bar{z}^{\bar{1}}$:

$$e^{-1} \mathcal{L}^{\text{book}} = \bar{z}^{\bar{1}} A^1 z^1 + B^1 \bar{z}^{\bar{1}} + \bar{B}^1 z^1 + C^1. \quad (3.4)$$

In order to solve for F^1 , we follow the approach described in [9]. Using the Kähler potential in Eq. (2.5) which immediately implies that

$$g_{i\bar{1}} = z^1 \partial_i g_{1\bar{1}}, \quad g_{i\bar{i}} = \bar{z}^{\bar{1}} \partial_i g_{1\bar{1}}, \quad (3.5)$$

we rewrite the first term in the Lagrangian (3.3) as follows,

$$(F^1 - F_G^1) g_{1\bar{1}} (\bar{F}^{\bar{1}} - \bar{F}_G^{\bar{1}}) - (F^1 - F_G^1) z^1 \bar{z}^{\bar{1}} (\partial_j g_{1\bar{1}}) (K_{0,j\bar{i}})^{-1} (\partial_i g_{1\bar{1}}) (\bar{F}^{\bar{1}} - \bar{F}_G^{\bar{1}}), \quad (3.6)$$

where we used that $(g_{ji})^{-1} = (K_{0,j\bar{i}})^{-1} - z^1 \bar{z}^{\bar{1}} (K_{0,j\bar{l}})^{-1} \times (K_{0,i\bar{m}})^{-1} (\partial_l \partial_{\bar{m}} g_{1\bar{1}})$ and the fact that $(z^1)^2 = 0$.

When solving for F^1 , following [1], one finds that $F^1 = F_G^1$ up to powers of χ^1 (see Eq. (A.24) in [1]). Since the second term in Eq. (3.6) has already the maximal power of χ^1 because $z^1 \bar{z}^{\bar{1}} \propto (\chi^1)^2 (\chi^{\bar{1}})^2$, we find that the second term does not contribute to the action at all. We show this explicitly in Appendix A. Thus, we find that we are left with

$$e^{-1} \mathcal{L}_{\text{off-shell}} = (F^1 - F_G^1) g_{1\bar{1}} (\bar{F}^{\bar{1}} - \bar{F}_G^{\bar{1}}) + e^{-1} \mathcal{L}^{\text{book}}. \quad (3.7)$$

Now following [9] we want to separate all the explicit z^1 , $\bar{z}^{\bar{1}}$ dependence in the first term of (3.7) from the dependence on F^1 and $\bar{F}^{\bar{1}}$. This will then bring the Lagrangian into a form in which we can easily integrate out F^1 following the procedure developed in [1]. To do that we define the following expansion coefficients that are independent of z^1 and $\bar{z}^{\bar{1}}$,

$$F_G^1 = F_{G0}^1 + F_{G1}^1 z^1 + F_{G\bar{1}}^1 \bar{z}^{\bar{1}} + F_{G1\bar{1}}^1 z^1 \bar{z}^{\bar{1}}, \quad (3.8)$$

and likewise for $\bar{F}_G^{\bar{1}}$. We present some details on the moduli space geometry for our models in Appendix B. Using it, a straightforward calculation given in Appendix C, shows that $F_{G\bar{1}}^1 = \bar{F}_{G1}^{\bar{1}} = 0$ for the Kähler potential given in Eq. (2.5).

Now let us look at the term $(F^1 - F_G^1) g_{1\bar{1}} (\bar{F}^{\bar{1}} - \bar{F}_G^{\bar{1}})$. We can absorb the z^1 and $z^1 \bar{z}^{\bar{1}}$ dependent terms of F_G^1 into F^1 by defining

$$F'^1 = F^1 - F_{G1}^1 z^1 - z^1 \bar{z}^{\bar{1}} F_{G1\bar{1}}^1. \quad (3.9)$$

This does not affect the nilpotency condition that fixes $z^1 = \frac{(\chi^1)^2}{2F^1}$ since

$$z^1 = \frac{(\chi^1)^2}{2F^1} = \frac{(\chi^1)^2}{2F'^1} = \frac{(\chi^1)^2}{2F^1} \left(1 + \frac{z^1}{2F^1} (F^1_{G_1} + \bar{z}^1 F^1_{G_1\bar{1}}) \right). \quad (3.10)$$

Here we used that $(\chi^1)^2 z^1 \propto (\chi^1)^2 (\chi^1)^2 = 0$. Likewise, we can define

$$\bar{F}'^{\bar{1}} = \bar{F}^{\bar{1}} - \bar{F}^{\bar{1}}_{G_1} \bar{z}^1 - z^1 \bar{z}^1 \bar{F}^{\bar{1}}_{G_1\bar{1}}. \quad (3.11)$$

This leads to

$$(F^1 - F^1_{G_0}) g_{1\bar{1}} (\bar{F}^{\bar{1}} - \bar{F}^{\bar{1}}_{G_0}) = (F'^1 - F^1_{G_0}) g_{1\bar{1}} (\bar{F}'^{\bar{1}} - \bar{F}^{\bar{1}}_{G_0}). \quad (3.12)$$

Putting everything together, we can now rewrite the Lagrangian (3.7) as

$$e^{-1} \mathcal{L} = (F'^1 - F^1_{G_0}) g_{1\bar{1}} (\bar{F}'^{\bar{1}} - \bar{F}^{\bar{1}}_{G_0}) + \bar{z}^1 A z^1 + B \bar{z}^1 + \bar{B} z^1 + C^1. \quad (3.13)$$

We now introduce $F = F'^1 \sqrt{g_{1\bar{1}}}$, $F_{G_0} = F^1_{G_0} \sqrt{g_{1\bar{1}}}$, $z = z^1 / \sqrt{g_{1\bar{1}}} = \frac{(\chi^1)^2}{2F}$ and similarly for the conjugated quantities. We also define $A = g_{1\bar{1}} A^1$, $B = \sqrt{g_{1\bar{1}}} B^1$, $C = C^1$ to bring the Lagrangian into the form³

$$e^{-1} \mathcal{L}_{\text{off-shell}} = (F - F_{G_0}) (\bar{F} - \bar{F}_{G_0}) + \bar{z} A z + B \bar{z} + \bar{B} z + C. \quad (3.14)$$

Now we can solve the equation for F that is given by

$$\begin{aligned} \frac{\delta \mathcal{L}(z, \bar{z}, F, \bar{F})}{\delta F} + \frac{\partial z}{\partial F} \frac{\delta \mathcal{L}(z, \bar{z}, F, \bar{F})}{\delta z} \\ = \frac{\delta \mathcal{L}(z, \bar{z}, F, \bar{F})}{\delta F} - \frac{z}{F} \frac{\delta \mathcal{L}(z, \bar{z}, F, \bar{F})}{\delta z} = 0. \end{aligned} \quad (3.15)$$

This was done for the Lagrangian (3.14) in the paper [1] (see, in particular, Appendix A.5). The resulting on-shell Lagrangian is given by

$$e^{-1} \mathcal{L}_{\text{on-shell}} = \left[\bar{z} A z + z \bar{B} + B \bar{z} + C - \frac{1}{F_{G_0} \bar{F}_{G_0}} (\bar{z} A z + \bar{z} B) (z \bar{A} \bar{z} + z \bar{B}) \right]_{z = \frac{(\chi^1)^2}{2F_{G_0}}}. \quad (3.16)$$

Expressing this in our original variables, we have

³Note that our definition of C is different from the one used in [1].

$$\begin{aligned} e^{-1} \mathcal{L}_{\text{on-shell}} &= \left[\bar{z}^{\bar{1}} A^1 z^1 + z^1 \bar{B}^1 + B^1 \bar{z}^{\bar{1}} + C^1 \right. \\ &\quad \left. - \frac{1}{g_{1\bar{1}} F^1_{G_0} \bar{F}^{\bar{1}}_{G_0}} |\bar{z}^{\bar{1}} A^1 z^1 + \bar{z}^{\bar{1}} B^1|^2 \right]_{z^1 = \frac{(\chi^1)^2}{2F^1_{G_0}}} \\ &= \left[e^{-1} \mathcal{L}^{\text{book}} - \frac{1}{g_{1\bar{1}} F^1_{G_0} \bar{F}^{\bar{1}}_{G_0}} \right. \\ &\quad \left. \times |\bar{z}^{\bar{1}} A^1 z^1 + \bar{z}^{\bar{1}} B^1|^2 \right]_{z^1 = \frac{(\chi^1)^2}{2F^1_{G_0}}}. \end{aligned} \quad (3.17)$$

To make it fully explicit that the ‘‘new’’ terms in the Lagrangian contain the maximal power $(\chi^1)^2 (\chi^{\bar{1}})^2$ of the undifferentiated spinor χ^1 , we can rewrite the Lagrangian as

$$\begin{aligned} e^{-1} \mathcal{L}_{\text{final}} &= [e^{-1} \mathcal{L}^{\text{book}}]_{z^1 = \frac{(\chi^1)^2}{2F^1_{G_0}}} \\ &\quad - \frac{(\chi^1)^2 (\chi^{\bar{1}})^2}{4g_{1\bar{1}} (F^1_{G_0} \bar{F}^{\bar{1}}_{G_0})^2} \left| A^1 \frac{(\chi^1)^2}{2F^1_{G_0}} + B^1 \right|^2, \end{aligned} \quad (3.18)$$

where the explicit expression for $F^1_{G_0}$ is derived in Appendix C and given by

$$F^1_{G_0} = \frac{1}{g_{1\bar{1}}} \left(-e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}} + \frac{1}{2} (\partial_i g_{1\bar{1}}) \bar{\chi}^1 \chi^i + \frac{1}{4} \bar{f}_{AB\bar{1}} \bar{\lambda}^A P_R \lambda^B \right). \quad (3.19)$$

Note that A^1 in the above expression has to act with two derivatives on $(\chi^1)^2$ in order to give a nonzero expression. In the Lagrangian $\mathcal{L}^{\text{book}}$ in (3.4) the relevant part of the $\bar{z}^{\bar{1}} A^1 z^1$ term which appears in the second term in (3.18) is

$$\bar{z}^{\bar{1}} g_{1\bar{1}} g^{\mu\nu} \partial_\mu \partial_\nu z^1 \equiv \bar{z}^{\bar{1}} g_{1\bar{1}} \square z^1. \quad (3.20)$$

Likewise, from B^1 and $F^1_{G_0}$ the only parts that contribute are those independent of the undifferentiated χ^1 and $\chi^{\bar{1}}$ since they appear in the above action with the highest power of the spinor χ^1 . We denote these parts by b^1 and f^1 , respectively. We explicitly spell out b^1 for two examples in Sec. IV and f^1 in Appendix C. Taking all this into account, the Lagrangian (3.18) simplifies to our final result given in Eq. (2.7) above.

IV. EXAMPLES

In this section we discuss two examples of our general result. First we discuss the simplest case of pure supergravity without chiral and vector multiplets. The action in this case was first derived in [1] and [2]. Then we discuss a pretty general case in which we allow for an arbitrary number of chiral and vector multiplets and only make the assumption that the gauge kinetic function f_{AB} and the moment maps \mathcal{P}_A are independent of z^1 .

A. Pure dS supergravity

For the case of a single nilpotent chiral superfield coupled to supergravity, the most general Kähler and superpotential are given by

$$K = z^1 \bar{z}^1, \quad W = g + fz^1, \quad (4.1)$$

with g and f complex constants. The corresponding action was derived in [1] and [2] in two different superconformal gauges. The one we are using here is based on the framework in [10].

Here we show how the action follows from our general answer given in (2.7). For the simple Kähler and superpotential given in (4.1), we find that $F_{G0}^1 = -\bar{f}$. This leads to the action

$$e^{-1} \mathcal{L}_{\text{pure dS}} = [e^{-1} \mathcal{L}^{\text{book}}]_{z^1 = -\frac{(\chi^1)^2}{2\bar{f}}} - \frac{(\chi^1)^2 (\bar{\chi}^1)^2}{4|f|^4} \left| -\frac{\square(\chi^1)^2}{2\bar{f}} + b^1 \right|^2, \quad (4.2)$$

where the explicit expression for b^1 can be read off from the general Lagrangian in Eqs. (18.6)–(18.19) in [10]. In

particular, α, β only take the value 1 and we want to extract all terms that are linear in \bar{z}^1 and independent of z^1 as described in Eq. (2.8). Furthermore, we can drop all terms that contain the undifferentiated spinor χ^1 or $\bar{\chi}^1$ (after potentially integrating by parts). This leaves us with only three terms, one coming from the scalar potential V , one coming from the mass term of the gravitino and, finally, one term which comes from the term $+\frac{1}{\sqrt{2}}\bar{\psi}_\mu \partial \bar{z}^1 \gamma^\mu \chi^1$ in the action after partial integration. The resulting answer is

$$b^1 = 2g\bar{f} + \frac{1}{2}\bar{f}\bar{\psi}_\mu P_L \gamma^{\mu\nu} \psi_\nu - \frac{1}{\sqrt{2}}\bar{\psi}_\mu \gamma^\nu \gamma^\mu \partial_\nu \chi^1. \quad (4.3)$$

B. De Sitter supergravity coupled to chiral and vector superfields

In this subsection we spell out the action in the case of an arbitrary number of chiral and vector multiplets but we make the simplifying assumption that the gauge kinetic function f_{AB} and the moment maps \mathcal{P}_A are independent of z^1 (and \bar{z}^1). The action is given by

$$e^{-1} \mathcal{L}_{\text{final}} = [e^{-1} \mathcal{L}^{\text{book}}]_{z^1 = \frac{(\chi^1)^2}{2f^1}} - \frac{(\chi^1)^2 (\bar{\chi}^1)^2}{4g_{1\bar{1}} (f^1 \bar{f}^1)^2} \left| g_{1\bar{1}} \frac{\square(\chi^1)^2}{2f^1} + b^1 \right|^2, \quad (4.4)$$

with

$$f^1 = -\frac{e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}}}{g_{1\bar{1}}},$$

$$b^1 = e^{\kappa_0} \left[2W_0 \bar{W}_{\bar{1}} + (K_{0,i\bar{j}})^{-1} (D_{0,i} W_0) \left(\frac{\partial_j g_{1\bar{1}}}{g_{1\bar{1}}} \bar{W}_{\bar{1}} - \bar{D}_{0,j} \bar{W}_{\bar{1}} \right) \right] - \frac{1}{2} (\bar{\chi}^{\bar{j}} \partial \chi^1) [3\partial_j g_{1\bar{1}} + (\bar{z}^{\bar{k}} \partial_{\bar{k}} - z^k \partial_k) \partial_j g_{1\bar{1}}]$$

$$+ \frac{1}{2} e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}} \bar{\psi}_\mu P_L \gamma^{\mu\nu} \psi_\nu - \frac{1}{\sqrt{2}} \bar{\psi}_\mu \gamma^\nu \gamma^\mu \partial_\nu \chi^1 (g_{1\bar{1}} + (\partial_j g_{1\bar{1}}) \bar{z}^{\bar{j}}) + \frac{1}{4} e^{\frac{\kappa_0}{2}} f_{ABj\bar{\lambda}} \bar{\lambda}^A P_L \lambda^B (K_{0,j\bar{k}})^{-1} \left((\partial_{\bar{k}} + K_{0,\bar{k}}) - \frac{\partial_{\bar{k}} g_{1\bar{1}}}{g_{1\bar{1}}} \right) \bar{W}_{\bar{1}}$$

$$- \frac{1}{2} e^{\frac{\kappa_0}{2}} \bar{\chi}^{\bar{j}} \chi^{\bar{j}} (\bar{D}_{0,\bar{i}} \bar{D}_{0,\bar{j}} - \frac{\partial_{\bar{i}} \partial_{\bar{j}} g_{1\bar{1}}}{g_{1\bar{1}}}) + \left(K_{0,k\bar{l}} \right)^{-1} (K_{0,k\bar{i}\bar{j}}) \left(\frac{\partial_{\bar{i}} g_{1\bar{1}}}{g_{1\bar{1}}} - \bar{D}_{0,\bar{i}} \right) \bar{W}_{\bar{1}} + \frac{1}{\sqrt{2}} e^{\frac{\kappa_0}{2}} \bar{D}_{0,\bar{i}} \bar{W}_{\bar{1}} \bar{\chi}^{\bar{i}} \gamma \cdot \psi, \quad (4.5)$$

where we defined $D_{0,i} = \partial_i + K_{0,i}$ and likewise $\bar{D}_{0,\bar{i}} = \partial_{\bar{i}} + K_{0,\bar{i}}$. This explicit expression for b^1 can again be read off from the action given in Ref. [10]. In particular, the first line arises from the scalar potential V , the second line comes from the kinetic terms for the fermions and the third line contains the generalization of the terms that we already found above in Eq. (4.3). The fourth line comes from the gaugino mass term and the fifth from the mass term for the χ^i . The last line contains a term coming from \mathcal{L}_{mix} in Eq. (18.18) in [10].

Note that this fairly general case includes the dS supergravity action coupled to a single chiral multiplet that was derived in [2].

V. SUMMARY

We constructed the locally supersymmetric supergravity action for general models with chiral and vector multiplets, presented in Eq. (2.7). It depends on all chiral multiplets, z^α : the nilpotent one z^1 and all other z^i with $i = 2, \dots, n$. Our models are defined by $K(z^\alpha, \bar{z}^{\bar{\alpha}})$, $W(z^\alpha)$ and $f_{AB}(z^\alpha)$ given by the following expressions

$$K(z^\alpha, \bar{z}^{\bar{\alpha}}) = K_0(z^i, \bar{z}^{\bar{i}}) + z^1 \bar{z}^{\bar{1}} g_{1\bar{1}}(z^i, \bar{z}^{\bar{i}}),$$

$$W(z^\alpha) = W_0(z^i) + z^1 W_1(z^i),$$

$$f_{AB}(z^\alpha) = f_{AB0}(z^i) + z^1 f_{AB1}(z^i). \quad (5.1)$$

Here the choice of W and f_{AB} is most general, since $(z^1)^2 = 0$, whereas the Kähler potential is assumed to depend on $z^1 \bar{z}^{\bar{1}}$.

Interesting features of de Sitter supergravity coupled to generic chiral and vector multiplets and a nilpotent chiral multiplet became clear after the complete action in Eq. (2.7) was derived:

- (1) The bosonic action is the standard supergravity action defined by $K(z^\alpha, \bar{z}^{\bar{\alpha}})$, $W(z^\alpha)$ and $f_{AB}(z^\alpha)$ which depend on all chiral multiplets. In this bosonic action one has to take $z^1 = 0$.
- (2) The complete fermionic action up to terms quadratic in $\chi^1 \chi^{\bar{1}}$ is given by the standard supergravity action defined by $K(z^\alpha, \bar{z}^{\bar{\alpha}})$, $W(z^\alpha)$ and $f_{AB}(z^\alpha)$, in which z^1 has to be replaced by

$$z^1 = g_{1\bar{1}}(z^i, \bar{z}^{\bar{i}}) \frac{(\chi^1)^2}{2} \times \left(-e^{\frac{k_0}{2}} \bar{W}_{\bar{1}}(\bar{z}^{\bar{i}}) + \frac{1}{4} \bar{f}_{AB\bar{1}}(\bar{z}^{\bar{i}}) \bar{\lambda}^A P_R \lambda^B \right)^{-1}. \quad (5.2)$$

- (3) The complete fermionic action to all orders in fermions is given by the standard supergravity action defined by $K(z^\alpha, \bar{z}^{\bar{\alpha}})$, $W(z^\alpha)$ and $f_{AB}(z^\alpha)$, in which z^1 has to be replaced as shown in Eq. (5.2). In addition to this, a term quartic and higher order in fermions has to be added to the action. It is given in closed form by the second term in Eq. (2.7).
- (4) In the unitary gauge

$$\chi^1 = 0, \quad (5.3)$$

the action reduces to the standard supergravity action defined by $K(z^\alpha, \bar{z}^{\bar{\alpha}})$, $W(z^\alpha)$ and $f_{AB}(z^\alpha)$ taken at $\chi^1 = z^1 = 0$ but with $F^1 \neq 0$:

$$F^1|_{z^1=\chi^1=0} = \frac{1}{g_{1\bar{1}}(z^i, \bar{z}^{\bar{i}})} \times \left(-e^{\frac{k_0}{2}} \bar{W}_{\bar{1}}(\bar{z}^{\bar{i}}) + \frac{1}{4} \bar{f}_{AB\bar{1}}(\bar{z}^{\bar{i}}) \bar{\lambda}^A P_R \lambda^B \right). \quad (5.4)$$

The nonlinearly realized local supersymmetry of the action in the unitary gauge $\chi^1 = 0$ is broken. The extra terms due to F^1 from the nilpotent multiplet include the nonvanishing positive term in the potential

$$V = e^{K_0(z^i, \bar{z}^{\bar{i}})} |W_1(z^i)|^2 g^{1\bar{1}}(z^i, \bar{z}^{\bar{i}}) > 0, \quad (5.5)$$

as well as some fermionic terms in the case where f_{AB} depends⁴ on z^1 . This is the basic feature of all de Sitter supergravity models, the leftover of the positive energy term even in the unitary local supersymmetry gauge in which the VA fermion is absent.

In conclusion, in this paper we have constructed a locally supersymmetric supergravity action for the case with a nilpotent multiplet and generic chiral and vectors multiplets. This creates a consistent framework for the investigation of phenomenological consequences of theories with nilpotent fields for particle physics and cosmology.

ACKNOWLEDGMENTS

We are grateful to E. Bergshoeff, S. Ferrara, D. Freedman, A. Linde and F. Quevedo for stimulating discussions. The work of R. K. and T. W. is supported by the SITP, by the National Science Foundation Grant No. PHY-1316699 and by the Templeton Foundation Grant ‘‘Quantum Gravity Frontiers.’’ T. W. thanks the Department of Physics of Stanford University and the Templeton Foundation for the hospitality during a visit in which this work was initiated.

APPENDIX A: VANISHING OF THE EXTRA TERM

In this section we show explicitly that the second term in (3.6) does not contribute to the final on-shell action. We first recall that $F^1 z^1 = F^1 \frac{(\chi^1)^2}{2F^1} = \frac{(\chi^1)^2}{2}$ is F^1 independent (and likewise for $\bar{F}^{\bar{1}} \bar{z}^{\bar{1}}$). This allows us to rewrite the second term in the Lagrangian (3.6), which due to the factor $z^1 \bar{z}^{\bar{1}}$ can only depend on F^1_{G0} and $\bar{F}^{\bar{1}}_{G0}$ (see Eq. (3.8) for the definition)

$$\begin{aligned} & -(F^1 - F^1_{G0}) z^1 \bar{z}^{\bar{1}} (\partial_j g_{1\bar{1}}) (K_{0,j\bar{1}})^{-1} (\partial_i g_{1\bar{1}}) (\bar{F}^{\bar{1}} - \bar{F}^{\bar{1}}_{G0}) \\ & = (\partial_j g_{1\bar{1}}) (K_{0,j\bar{1}})^{-1} (\partial_i g_{1\bar{1}}) \left(-z^1 \bar{z}^{\bar{1}} F^1_{G0} \bar{F}^{\bar{1}}_{G0} + z^1 \frac{(\chi^1)^2}{2} F^1_{G0} \right. \\ & \quad \left. + \bar{z}^{\bar{1}} \frac{(\chi^1)^2}{2} \bar{F}^{\bar{1}}_{G0} - \frac{(\chi^1)^2 (\chi^{\bar{1}})^2}{4} \right). \end{aligned} \quad (A1)$$

So we see that the explicit F^1 , $\bar{F}^{\bar{1}}$ dependence completely disappears. The above terms then become a correction to (3.4) that of course also does not have an explicit dependence on F^1 and $\bar{F}^{\bar{1}}$. In particular, we find that the coefficient A^1 , B^1 , $\bar{B}^{\bar{1}}$ and C^1 get the following extra contributions,

⁴In string theory constructions of the nilpotent multiplet via the anti-D3 brane, one may argue that f_{AB} does not depend on z^1 [7]. In such case, the only effect of the nilpotent multiplet in the supergravity unitary gauge $\chi^1 = 0$ is the vacuum energy uplift (5.5).

$$\begin{aligned}
\Delta A^1 &= -\mathcal{M} F_{G0}^1 \bar{F}_{G0}^{\bar{1}}, \\
\Delta B^1 &= \mathcal{M} \frac{(\chi^1)^2}{2} \bar{F}_{G0}^{\bar{1}}, \\
\Delta \bar{B}^1 &= \mathcal{M} \frac{(\bar{\chi}^{\bar{1}})^2}{2} F_{G0}^1, \\
\Delta C^1 &= -\mathcal{M} \frac{(\chi^1)^2 (\bar{\chi}^{\bar{1}})^2}{4},
\end{aligned} \tag{A2}$$

with

$$\mathcal{M} = (\partial_j g_{1\bar{1}}) (K_{0,j\bar{i}})^{-1} (\partial_i g_{1\bar{1}}). \tag{A3}$$

Then we can proceed as in Sec. III and integrate out F^1 . The final action expressed in our original variables takes the form [cf. Eq. (3.17)]

$$\begin{aligned}
e^{-1} \mathcal{L}_{\text{on-shell}} &= \left[\bar{z}^{\bar{1}} (A^1 + \Delta A^1) z^1 + z^1 (\bar{B}^1 + \Delta \bar{B}^1) + (B^1 + \Delta B^1) \bar{z}^{\bar{1}} + (C^1 + \Delta C^1) \right. \\
&\quad \left. - \frac{1}{g_{1\bar{1}} F_{G0}^1 \bar{F}_{G0}^{\bar{1}}} |\bar{z}^{\bar{1}} (A^1 + \Delta A^1) z^1 + \bar{z}^{\bar{1}} (B^1 + \Delta B^1)|^2 \right]_{z^1 = \frac{(\chi^1)^2}{2F_{G0}^1}} \\
&= \left[e^{-1} \mathcal{L}^{\text{book}} + \bar{z}^{\bar{1}} \Delta A^1 z^1 + z^1 \Delta \bar{B}^1 + \Delta B^1 \bar{z}^{\bar{1}} + \Delta C^1 - \frac{1}{g_{1\bar{1}} F_{G0}^1 \bar{F}_{G0}^{\bar{1}}} |\bar{z}^{\bar{1}} A^1 z^1 + B^1 \bar{z}^{\bar{1}} + \bar{z}^{\bar{1}} \Delta A^1 z^1 + \Delta B^1 \bar{z}^{\bar{1}}|^2 \right]_{z^1 = \frac{(\chi^1)^2}{2F_{G0}^1}}.
\end{aligned} \tag{A4}$$

Now we note that using the explicit expressions in (A2) we find

$$\begin{aligned}
\bar{z}^{\bar{1}} \Delta A z^1 \Big|_{z^1 = \frac{(\chi^1)^2}{2F_{G0}^1}} &= \Delta C = -\mathcal{M} \frac{(\chi^1)^2 (\bar{\chi}^{\bar{1}})^2}{4}, \\
z^1 \Delta \bar{B} \Big|_{z^1 = \frac{(\chi^1)^2}{2F_{G0}^1}} &= \Delta B \bar{z}^{\bar{1}} \Big|_{z^1 = \frac{(\chi^1)^2}{2F_{G0}^1}} = +\mathcal{M} \frac{(\chi^1)^2 (\bar{\chi}^{\bar{1}})^2}{4}.
\end{aligned} \tag{A5}$$

This implies that

$$[\bar{z}^{\bar{1}} \Delta A^1 z^1 + z^1 \Delta \bar{B}^1 + \Delta B^1 \bar{z}^{\bar{1}} + \Delta C^1]_{z^1 = \frac{(\chi^1)^2}{2F_{G0}^1}} = 0. \tag{A6}$$

We also see from (A5) that

$$[\bar{z}^{\bar{1}} \Delta A z^1 + \Delta B \bar{z}^{\bar{1}}]_{z^1 = \frac{(\chi^1)^2}{2F_{G0}^1}} = 0. \tag{A7}$$

Thus, we have explicitly shown that the Lagrangian (A4) reduces to the one in Eq. (3.17).

APPENDIX B: INVERSE KÄHLER METRIC AND CHRISTOFFEL SYMBOLS

From the Kähler metric as given in Eq. (2.5) we trivially find the following expansion in z^1 and $\bar{z}^{\bar{1}}$

$$\begin{aligned}
g_{1\bar{1}} &= g_{1\bar{1}}(z^i, \bar{z}^{\bar{i}}), \\
g_{1\bar{i}} &= \bar{z}^{\bar{i}} \partial_i g_{1\bar{1}}, \\
g_{i\bar{1}} &= z^1 \partial_i g_{1\bar{1}}, \\
g_{i\bar{j}} &= K_{0,i\bar{j}} + z^1 \bar{z}^{\bar{i}} \partial_i \partial_j g_{1\bar{1}}.
\end{aligned} \tag{B1}$$

Contracting the Kähler metric $g_{\alpha\beta}$ with its inverse we find the following relations

$$\begin{aligned}
g_{1\bar{1}} g^{\bar{1}1} + g_{1\bar{j}} g^{\bar{j}1} &= 1, \\
g_{1\bar{1}} g^{\bar{1}k} + g_{1\bar{j}} g^{\bar{j}k} &= 0, \\
g_{i\bar{1}} g^{\bar{1}1} + g_{i\bar{j}} g^{\bar{j}1} &= 0, \\
g_{i\bar{1}} g^{\bar{1}k} + g_{i\bar{j}} g^{\bar{j}k} &= \delta_i^k.
\end{aligned} \tag{B2}$$

Now we recall that

$$(g_{j\bar{i}})^{-1} = (K_{0,j\bar{i}})^{-1} - z^1 \bar{z}^{\bar{i}} (K_{0,j\bar{l}})^{-1} (K_{0,i\bar{m}})^{-1} \partial_l \partial_{\bar{m}} g_{1\bar{1}}, \tag{B3}$$

and use the above (B2) to find the expansion of the inverse metric

$$\begin{aligned}
g^{1\bar{1}} &= \frac{1}{g_{1\bar{1}} - g_{1\bar{j}}(g_{ij})^{-1}g_{i\bar{1}}} \\
&= \frac{1}{g_{1\bar{1}}} + z^1 \bar{z}^{\bar{1}} \frac{(\partial_j g_{1\bar{1}})(K_{0,i\bar{j}})^{-1}(\partial_i g_{1\bar{1}})}{(g_{1\bar{1}})^2}, \\
g^{1\bar{j}} &= -(g_{i\bar{j}})^{-1}g_{i\bar{1}}g^{1\bar{1}} = -z^1 \frac{(K_{0,i\bar{j}})^{-1}(\partial_i g_{1\bar{1}})}{g_{1\bar{1}}}, \\
g^{k\bar{j}} &= (g_{i\bar{j}})^{-1}(\delta_i^k - g_{i\bar{1}}g^{1\bar{k}}) \\
&= (K_{0,k\bar{j}})^{-1} - z^1 \bar{z}^{\bar{1}} \left[(K_{0,j\bar{l}})^{-1}(K_{0,k\bar{m}})^{-1} \partial_l \partial_{\bar{m}} g_{1\bar{1}} \right. \\
&\quad \left. - \frac{(K_{0,i\bar{j}})^{-1}(\partial_i g_{1\bar{1}})(K_{0,k\bar{m}})^{-1}(\partial_{\bar{m}} g_{1\bar{1}})}{g_{1\bar{1}}} \right]. \tag{B4}
\end{aligned}$$

The Christoffel symbols for the Kähler manifold are given by $\Gamma_{\beta\gamma}^\alpha = g^{\alpha\bar{\delta}} \partial_\beta g_{\gamma\bar{\delta}}$. For the Christoffel symbols with upper index $\alpha = 1$ we explicitly find

$$\begin{aligned}
\Gamma_{1\bar{1}}^1 &= g^{1\bar{1}} \partial_1 g_{1\bar{1}} + g^{1\bar{i}} \partial_1 g_{i\bar{1}} = 0, \\
\Gamma_{1\bar{i}}^1 &= g^{1\bar{1}} \partial_1 g_{i\bar{1}} + g^{1\bar{j}} \partial_1 g_{j\bar{i}} \\
&= \frac{\partial_i g_{1\bar{1}}}{g_{1\bar{1}}} + z^1 \bar{z}^{\bar{1}} \left(\frac{(\partial_i g_{1\bar{1}})(\partial_j g_{1\bar{1}})(K_{0,k\bar{j}})^{-1}(\partial_k g_{1\bar{1}})}{(g_{1\bar{1}})^2} \right. \\
&\quad \left. - \frac{(\partial_i \partial_j g_{1\bar{1}})(K_{0,k\bar{j}})^{-1}(\partial_k g_{1\bar{1}})}{g_{1\bar{1}}} \right), \\
\Gamma_{i\bar{j}}^1 &= g^{1\bar{1}} \partial_i g_{j\bar{1}} + g^{1\bar{k}} \partial_i g_{j\bar{k}} \\
&= z^1 \left(\frac{\partial_i \partial_j g_{1\bar{1}}}{g_{1\bar{1}}} - \frac{(K_{0,i\bar{k}})^{-1}(\partial_l g_{1\bar{1}})}{g_{1\bar{1}}} K_{0,ij\bar{k}} \right). \tag{B5}
\end{aligned}$$

APPENDIX C: THE EXPANSION OF F_G^1

Using the equations from the previous appendix we can now expand F_G^1 to get the explicit expression for $F_{G0}^1 = F_G^1|_{z^1=\bar{z}^{\bar{1}}=0}$. We will also show that $F_{G\bar{1}}^1 = 0$, i.e. that F_G^1 has no terms that only depend on $\bar{z}^{\bar{1}}$ and not on z^1 . It then follows automatically that its complex conjugate $\bar{F}_{G1}^{\bar{1}}$ vanishes as well.

Let us first recall the definition of F_G^1 in Eqs. (2.3) and (2.4),

$$F_G^1 = -e^{\frac{\kappa}{2}} g^{1\bar{\beta}} \bar{\nabla}_{\bar{\beta}} \bar{W} + \frac{1}{2} \Gamma_{\beta\gamma}^1 \bar{\chi}^\beta \chi^\gamma + \frac{1}{4} \bar{f}_{AB\bar{\beta}} g^{\bar{\beta}1} \bar{\lambda}^A P_R \lambda^B. \tag{C1}$$

Expanding the bosonic part in powers of z^1 and $\bar{z}^{\bar{1}}$ we find using (B4) that

$$\begin{aligned}
&e^{\frac{\kappa}{2}} (g^{1\bar{1}} \bar{\nabla}_{\bar{1}} \bar{W} + g^{1\bar{i}} \bar{\nabla}_{\bar{i}} \bar{W}) \\
&= e^{\frac{\kappa_0}{2}} \left(1 + \frac{z^1 \bar{z}^{\bar{1}} g_{1\bar{1}}}{2} \right) \left(g^{1\bar{1}} (\bar{W}_{\bar{1}} + z^1 g_{1\bar{1}} \bar{W}) \right. \\
&\quad \left. - z^1 \frac{(K_{0,j\bar{1}})^{-1}(\partial_j g_{1\bar{1}})}{g_{1\bar{1}}} \bar{\nabla}_{\bar{j}} \bar{W} \right) \\
&= \frac{e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}}}{g_{1\bar{1}}} + \mathcal{O}(z^1, z^1 \bar{z}^{\bar{1}}). \tag{C2}
\end{aligned}$$

Since $\bar{W}_{\bar{1}}$ is independent of $\bar{z}^{\bar{1}}$ we see that the above expression contains no term linear in $\bar{z}^{\bar{1}}$ (and independent of z^1).

The term $\frac{1}{2} \Gamma_{\beta\gamma}^1 \bar{\chi}^\beta \chi^\gamma$ can only have a term linear in $\bar{z}^{\bar{1}}$ if $\Gamma_{\beta\gamma}^1$ has such a term. However, from the explicit expressions in Eq. (B5) we see that this is not the case. In particular, we find the expansion

$$\frac{1}{2} \Gamma_{\beta\gamma}^1 \bar{\chi}^\beta \chi^\gamma = \frac{(\partial_i g_{1\bar{1}}) \bar{\chi}^1 \chi^i}{2g_{1\bar{1}}} + \mathcal{O}(z^1, z^1 \bar{z}^{\bar{1}}). \tag{C3}$$

Finally, we see that the gaugino term can be expanded as

$$\begin{aligned}
\frac{1}{4} \bar{f}_{AB\bar{\beta}} g^{\bar{\beta}1} \bar{\lambda}^A P_R \lambda^B &= \frac{1}{4} (\bar{f}_{AB\bar{1}} g^{\bar{1}1} + \bar{f}_{AB\bar{i}} g^{\bar{i}1}) \bar{\lambda}^A P_R \lambda^B \\
&= \frac{\bar{f}_{AB\bar{1}} \bar{\lambda}^A P_R \lambda^B}{4g_{1\bar{1}}} + \mathcal{O}(z^1, z^1 \bar{z}^{\bar{1}}) \tag{C4}
\end{aligned}$$

and does not have a linear term in $\bar{z}^{\bar{1}}$ either. So we conclude that F_G^1 has no linear terms in $\bar{z}^{\bar{1}}$ and therefore by definition we have $F_{G\bar{1}}^1 = 0$.

Gathering the terms in (C2)–(C4) we find

$$F_{G0}^1 = \frac{1}{g_{1\bar{1}}} \left(-e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}} + \frac{1}{2} (\partial_i g_{1\bar{1}}) \bar{\chi}^1 \chi^i + \frac{1}{4} \bar{f}_{AB\bar{1}} \bar{\lambda}^A P_R \lambda^B \right). \tag{C5}$$

As discussed above, in the action F_{G0}^1 always appears multiplied by $(\chi^1)^2 = \bar{\chi}^1 \chi^1$ like for example in

$$\begin{aligned}
z_G^1 &= \frac{(\chi^1)^2}{2F_{G0}^1} \\
&= -\frac{g_{1\bar{1}} (\chi^1)^2}{2e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}}} \sum_{n \geq 0} \left(\frac{\frac{1}{2} (\partial_i g_{1\bar{1}}) \bar{\chi}^1 \chi^i + \frac{1}{4} \bar{f}_{AB\bar{1}} \bar{\lambda}^A P_R \lambda^B}{e^{\frac{\kappa_0}{2}} \bar{W}_{\bar{1}}} \right)^n. \tag{C6}
\end{aligned}$$

Since $(\chi^1)^2 \bar{\chi}^1 \chi^i = 0$ we can drop the term linear in χ^1 and find

$$z_G^1 = \frac{(\chi^1)^2}{2F_{G0}^1} = -\frac{g_{1\bar{1}}(\chi^1)^2}{2e^{\frac{\kappa_0}{2}}\bar{W}_{\bar{1}}} \sum_{n \geq 0} \left(\frac{\frac{1}{4}\bar{f}_{AB\bar{1}}\bar{\lambda}^A P_R \lambda^B}{e^{\frac{\kappa_0}{2}}\bar{W}_{\bar{1}}} \right)^n \equiv \frac{(\chi^1)^2}{2f^1}, \quad (C7)$$

where we defined $f^1 \equiv \frac{1}{g_{1\bar{1}}}(-e^{\frac{\kappa_0}{2}}\bar{W}_{\bar{1}} + \frac{1}{4}\bar{f}_{AB\bar{1}}\bar{\lambda}^A P_R \lambda^B)$ to contain all the terms in F_{G0}^1 that are independent of χ^1 .

-
- [1] E. A. Bergshoeff, D. Z. Freedman, R. Kallosh, and A. Van Proeyen, Pure de Sitter supergravity, [arXiv:1507.08264](#).
- [2] F. Hasegawa and Y. Yamada, Component action of nilpotent multiplet coupled to matter in 4 dimensional $\mathcal{N} = 1$ supergravity, *J. High Energy Phys.* **10** (2015) 106.
- [3] P. K. Townsend, Cosmological constant in supergravity, *Phys. Rev. D* **15**, 2802 (1977).
- [4] D. Volkov and V. Akulov, Is the neutrino a Goldstone particle?, *Phys. Lett.* **46B**, 109 (1973).
- [5] S. Kachru, R. Kallosh, A. D. Linde, and S. P. Trivedi, De Sitter vacua in string theory, *Phys. Rev. D* **68**, 046005 (2003). S. Kachru, R. Kallosh, A. D. Linde, J. M. Maldacena, L. P. McAllister, and S. P. Trivedi, Towards inflation in string theory, *J. Cosmol. Astropart. Phys.* **10** (2003) 013.
- [6] E. A. Bergshoeff, K. Dasgupta, R. Kallosh, A. Van Proeyen, and T. Wrase, $\overline{D3}$ and dS, *J. High Energy Phys.* **05** (2015) 058; R. Kallosh and T. Wrase, Emergence of spontaneously broken supersymmetry on an anti-D3-brane in KKLT dS vacua, *J. High Energy Phys.* **12** (2014) 117.
- [7] R. Kallosh, F. Quevedo, and A. M. Uranga, String theory realizations of the nilpotent Goldstino, [arXiv:1507.07556](#).
- [8] S. Ferrara, R. Kallosh, and A. Linde, Cosmology with nilpotent superfields, *J. High Energy Phys.* **10** (2014) 143.
- [9] R. Kallosh, Matter-coupled de Sitter supergravity, [arXiv:1509.02136](#).
- [10] D. Z. Freedman and A. Van Proeyen, *Supergravity* (Cambridge University Press, Cambridge, England, 2012).
- [11] E. Cremmer, S. Ferrara, L. Girardello, and A. Van Proeyen, Coupling supersymmetric Yang-Mills theories to supergravity, *Phys. Lett.* **116B**, 231 (1982); Yang-Mills theories with local supersymmetry: Lagrangian, transformation laws and super-Higgs effect, *Nucl. Phys.* **B212**, 413 (1983).
- [12] T. Kugo and S. Uehara, Improved superconformal gauge conditions in the $N = 1$ supergravity Yang-Mills matter system, *Nucl. Phys.* **B222**, 125 (1983).
- [13] R. Kallosh, L. Kofman, A. D. Linde, and A. Van Proeyen, Superconformal symmetry, supergravity and cosmology, *Classical Quantum Gravity* **17**, 4269 (2000); *Classical Quantum Gravity* **21**, 5017 (2004).
- [14] I. Antoniadis, E. Dudas, S. Ferrara, and A. Sagnotti, The Volkov-Akulov-Starobinsky supergravity, *Phys. Lett. B* **733**, 32 (2014).
- [15] E. Dudas, S. Ferrara, A. Kehagias, and A. Sagnotti, Properties of nilpotent supergravity, [arXiv:1507.07842](#); S. Ferrara, M. Porrati, and A. Sagnotti, Scale invariant Volkov-Akulov supergravity, [arXiv:1508.02939](#); I. Antoniadis and C. Markou, The coupling of non-linear supersymmetry to supergravity, [arXiv:1508.06767](#);
- [16] S. M. Kuzenko, Complex linear Goldstino superfield and de Sitter supergravity, *J. High Energy Phys.* **10** (2015) 006.
- [17] R. Kallosh and A. Linde, Inflation and uplifting with nilpotent superfields, *J. Cosmol. Astropart. Phys.* **01** (2015) 025; G. Dall'Agata and F. Zwirner, On sgoldstino-less supergravity models of inflation, *J. High Energy Phys.* **12** (2014) 172; J. J. M. Carrasco, R. Kallosh, A. Linde, and D. Roest, The hyperbolic geometry of cosmological attractors, *Phys. Rev. D* **92**, 041301 (2015); J. J. M. Carrasco, R. Kallosh, and A. Linde, α -Attractors: Planck, LHC and Dark Energy, [arXiv:1506.01708](#).