

Renormalization group optimized perturbation theory at finite temperatures

Jean-Loïc Kneur^{1,*} and Marcus B. Pinto^{2,†}¹*Laboratoire Charles Coulomb (L2C), UMR 5221 CNRS-Université Montpellier 2, 34095 Montpellier, France*²*Departamento de Física, Universidade Federal de Santa Catarina, 88040-900 Florianópolis, Santa Catarina, Brazil*

(Received 18 August 2015; published 23 December 2015)

A recently developed variant of the so-called optimized perturbation theory (OPT), making it perturbatively consistent with renormalization group (RG) properties, RGOPT, was shown to drastically improve its convergence for zero temperature theories. Here the RGOPT adapted to finite temperature is illustrated with a detailed evaluation of the two-loop pressure for the thermal scalar $\lambda\phi^4$ field theory. We show that already at the simple one-loop level this quantity is exactly scale-invariant by construction and turns out to qualitatively reproduce, with a rather simple procedure, results from more sophisticated resummation methods at two-loop order, such as the two-particle irreducible approach typically. This lowest order also reproduces the exact large- N results of the $O(N)$ model. Although very close in spirit, our RGOPT method and corresponding results differ drastically from similar variational approaches, such as the screened perturbation theory or its QCD-version, the (resummed) hard thermal loop perturbation theory. The latter approaches exhibit a sensibly degrading scale dependence at higher orders, which we identify as a consequence of missing RG invariance. In contrast RGOPT gives a considerably reduced scale dependence at two-loop level, even for relatively large coupling values $\sqrt{\lambda/24} \sim \mathcal{O}(1)$, making results much more stable as compared with standard perturbation theory, with expected similar properties for thermal QCD.

DOI: [10.1103/PhysRevD.92.116008](https://doi.org/10.1103/PhysRevD.92.116008)

PACS numbers: 11.10.Wx, 11.10.Gh, 12.38.Lg

I. INTRODUCTION

It is a well-established fact that evaluations devoted to describe quantum chromodynamics (QCD) phase transitions need to be done in a nonperturbative fashion, such as numerically solving it on the lattice (LQCD) [1], which nowadays is considered the most reliable way to tackle the problem at vanishing densities. Unfortunately, LQCD is still plagued by the so-called sign problem [2] which prevents the method from being used to describe the transitions expected to take place at lower temperatures and higher densities. On the other hand, multiloop perturbative results for many QCD physical quantities are within reach so that an appealing alternative would be to use them in conjunction with some resummation procedure in order to generate nonperturbative results. In this vein, different analytical techniques envisaged to combine the easiness of purely perturbative evaluations with nonperturbative optimization/resummation procedures have been proposed in the past decades [3]. Some of these methods are based on a reorganization of a given interacting Lagrangian, so that it becomes written in terms of an arbitrary mass parameter which, for massless theories, also works as an infrared regulator (as in hard thermal loop resummations [4,5]). One of these approaches is the so-called screened perturbation

theory (SPT) [6,7], in which the variational parameter is described by a thermal mass. The SPT was originally proposed to describe the thermodynamics of massless scalar theories, but it has been later generalized so that the equation of state of thermal gauge-invariant theories [4], such as QCD, could also be obtained. This gauge invariant generalization known as hard thermal loop (resummed) perturbation theory (HTLpt) [8], has been already used to calculate QCD thermodynamic functions up to three loop order at finite values of the temperature and chemical potential [9,10]. The SPT and HTLpt are actually conceptually similar to the so-called linear delta expansion (LDE) and optimized perturbation theory (OPT), developed earlier under various different names [11–13] mainly in the context of zero-temperature field theories. Within this technique, perturbative evaluations are performed using propagators written in terms of an arbitrary mass parameter, so that optimized nonperturbative results can be generated by requiring the mass parameter to satisfy a variational criterion. The two major problems the above mentioned methods try to solve are the poor convergence and the notoriously bad scale-dependence of the standard perturbative series both for the thermal mass and for the pressure at higher orders (see e.g. [3] for a review): not only the increasing perturbative orders show no clear sign of stability, but the scale-dependence worsens substantially at higher orders, at odds with what is intuitively expected for most known perturbative series at $T = 0$. Part of this

*jean-loic.kneur@univ-montp2.fr
†marcus.benghi@ufsc.br

bad behavior is commonly explained [3] by the unavoidable complicated interplay of soft and hard thermal contributions. Actually, the dynamical generation of a thermal screening mass $m_D \sim \sqrt{\lambda}T$ influences the relevant expansion of physical quantities, such as the pressure, which are then expressed in powers of $\sqrt{\lambda}$ rather than λ . In this case the predictions are, *a priori*, less convergent than for the $T = 0$ case, yet most of the interesting thermal physics happens at rather moderate coupling values, so that one could expect a better behavior. Despite the unavoidable $\sqrt{\lambda}$ “nonperturbative” dependence, both SPT [6,7] and OPT [14] applications to hot scalar theory show how these methods indeed improve the stability of the predictions when higher orders in the loop expansion are considered. Given the inherent technical difficulties associated with the (three loop) evaluation of the QCD pressure for the case of hot and dense quark matter, the recent results in [10] represent an impressive achievement. However, the same results exhibit a substantial *increasing* scale-dependence at increasing two- and three-loop orders, even for moderately large coupling values, which remains a surprising issue. This is more pronounced for HTLpt in QCD applications at the three loop level [9,10]. While the latter are sometimes remarkably close to lattice results for temperatures down to $T \gtrsim 2T_c$, for the central renormalization scale choice $\mu \sim 2\pi T$ in the $\overline{\text{MS}}$ -scheme, it appears puzzling that a moderate scale variation of a factor 2 dramatically affects the pressure and related thermodynamical quantities by relative variations of order 1 or more. It is argued [10] that resumming the logarithmic dependence of HTLpt results may improve this problem, but as we shall explain below the missing RG invariance properties is more basic within the SPT/HTLpt approach.

Recently, the standard OPT procedure at zero temperature has been modified to incorporate consistently perturbative renormalization group (RG) properties. It was shown to considerably improve the convergence of OPT, as tested for the Gross-Neveu (GN) model mass gap [15], and further used to determine with a good accuracy the basic QCD scale in the $\overline{\text{MS}}$ scheme, $\Lambda_{\overline{\text{MS}}}$, and corresponding value of the strong coupling $\bar{\alpha}_S$ [16,17]. Very recently the same method has been used to estimate [18] the QCD chiral quark condensate. Here, we extend the construction to the case of a scalar theory with quartic interaction to show how this RG improved OPT (RGOPT) can easily cope with the introduction of control parameters such as the temperature, developing in detail the construction presented very recently in [19]. Using this textbook example we aim to illustrate how the RGOPT takes care of the scale dependence problems of thermal theories, those being more clearly visible within the SPT [6,7] and HTLpt higher loop results [8–10], but also present in other resummation approaches.

In a nutshell, leaving aside technical details, our basic observation is that the arbitrary variational mass introduced in the SPT/OPT context can (and should) be treated as any

“proper” mass, from the RG viewpoint. In particular, this means that it involves its own standard anomalous dimension, which is unrelated to the coupling β function, and should be incorporated consistently within RG properties. This brings crucial consequences already in the form of an explicitly RG-invariant pressure, prior to any further improved/resummed perturbation approaches. It also more clearly separates the hard and soft modes from a RG/scale dependence perspective, at least in an intermediate stage of the calculations. In the more standard approach both contributions are mixed up since all relevant quantities are expressed as a function of the coupling and temperature only. That being accepted, one realizes that most of the observed worsening scale-dependence at higher orders is actually due to a manifest failure of RG invariance for a massive theory. This is at least particularly transparent in the $\overline{\text{MS}}$ scheme largely used within SPT/HTLpt methods, where we show that minimally subtracting the vacuum energy divergences, without finite vacuum energy contributions, explicitly misses RG invariance.

In addition, even when starting from an explicitly RG invariant perturbative expression, RG invariance is generally lost as a consequence of the standard (linear) modification of perturbative expansion implied in OPT/SPT, a fact that has been seldom appreciated in the relevant literature so far. However, RG properties can easily be restored by a consistent use of renormalization group properties for a massive theory, automatically incorporated in the RGOPT approach [17], whereby a drastically improved scale-dependence follows naturally. This opens up the possibility of rather simply exploiting many non-trivial SPT/HTLpt results performed up to three-loop order so far, by improving substantially their scale independence after applying appropriate RGOPT adaptations.

Incidentally, other resummation approaches, like typically the so-called “two-loop- Φ -derivable” approach, [related to the two-particle-irreducible (2PI) method] [20,21] are manifestly scale invariant when applied to the scalar model in a certain approximation. In fact one observes interesting analogies (but also important differences) between the first nontrivial (two-loop) 2PI order and our *one-loop* RGOPT results, as will be elaborated in more detail below. However, as it happens the scale-invariance of the above-mentioned 2PI results [20,21] is essentially due to a renormalization that appears peculiar to the scalar model, inspired by the properties of the $O(N)$ symmetric theory in the large- N limit [22] (for which indeed the two-loop- Φ -derivable 2PI results become exact). Thus, to the best of our understanding, this scale invariance appears somewhat accidental and difficult to translate to higher orders and to QCD. Incidentally, the 2PI approach has been pushed even to three-loop order for the ϕ^4 theory in Ref. [23], with remarkably stable results with respect to two-loop order, but the method becomes more involved, and a definite scale-dependence reappears, although much

more moderate than in the three-loop order SPT and HTLpt cases. Other approaches like the nonperturbative renormalization group (NPRG) [24], should be RG invariant by construction. But solving the relevant NPRG equations for thermal QCD beyond approximative truncation schemes becomes very involved in practice. We also remark that our approach shares some qualitative features with the ideas invoked and the framework developed very recently in Ref. [25], in which the authors explored some form of massive renormalization scheme and its consequences. However, the RGOPT differs substantially in its approach as it incorporates by construction [17,19] all relevant RG properties systematically order by order, relying basically on standard perturbation theory. Being largely based on (but not limited to) $\overline{\text{MS}}$ -scheme results, the RGOPT can easily be extended to any higher order calculations performed in the framework of different models, when already available, including thermal QCD. Therefore, it appears to us that the method presented here is conceptually simpler than other more sophisticated resummation approaches mentioned above.

The paper is organized as follows. In the next section we quickly review some basic results of thermal scalar theory for the free energy at the relevant two-loop level. Then, in Sec. III we address in some details the RG (non) invariance issue in the massive case, paying special attention to the $\overline{\text{MS}}$ scheme largely used in the SPT (or the similar HTLpt) method. In the same section we also explain how to restore the perturbative RG invariance at arbitrary orders in a simple fashion. Next, the resummation by optimized perturbation (OPT), with the crucial modification to maintain its perturbative RG invariance, RGOPT, is discussed in rather general terms in Secs. IV and V respectively. The method is then illustrated in details by evaluating the free energy of a hot scalar field theory at one- and two-loop level in Secs. VI and VII. We emphasize that all the construction developed in Secs. III to VII for the ϕ^4 model (for which we also briefly consider the large- N case) is actually more general, most of it being straightforwardly applicable to thermal QCD. We occasionally mention some properties anticipated to be similar (or eventually different) in the QCD case. Finally, our conclusions are presented in Sec. VIII, and one appendix deals with some technical details on the RG-invariant construction of counterterms.

II. FINITE TEMPERATURE SCALAR FIELD THEORY

We consider, as a starting point, the massive scalar field theory described by the Lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{m^2}{2} \phi^2 - \frac{\lambda}{4!} \phi^4; \quad (2.1)$$

where we introduce a (yet unspecified) generic mass term (m) which can be thought as a thermal mass generated by

higher perturbative orders in an originally massless theory. One may then evaluate the free energy using known results from ordinary perturbation theory for the massive case. Then, up to the two-loop level the basic expression of the (bare) free energy is formally [3,7,26]:

$$\mathcal{F}_0 = \frac{1}{2} \int_p \ln(p^2 + m^2) + \frac{\lambda}{8} \left(\int_p \frac{1}{p^2 + m^2} \right)^2 + \mathcal{F}_0^{\text{ct}}, \quad (2.2)$$

where the temperature is introduced via Matsubara's imaginary time formalism ($p^2 \equiv \omega_n^2 + \mathbf{p}^2$ with the bosonic Matsubara $\omega_n = 2\pi nT$) and we have also defined

$$\int_p \equiv \left(e^{\gamma_E} \frac{\mu^2}{4\pi} \right)^\epsilon T \sum_n \int \frac{d^{D-1}p}{(2\pi)^{D-1}}. \quad (2.3)$$

The divergent integrals are regulated using dimensional regularization techniques with $D = 4 - 2\epsilon$ while renormalization is carried out in the $\overline{\text{MS}}$ -scheme. The term $\mathcal{F}_0^{\text{ct}}$ represents all the relevant counterterms contributions to $O(\lambda)$ (see Ref. [7] for details). After all the mass, coupling, and vacuum energy counterterms have been consistently introduced to cancel the original divergences, one obtains the ($\overline{\text{MS}}$ -scheme) renormalized free energy [3,7]:

$$(4\pi)^2 \mathcal{F}_0 = \mathcal{E}_0 - \frac{1}{8} m^4 \left[3 + 2 \ln \left(\frac{\mu^2}{m^2} \right) \right] - \frac{1}{2} T^4 J_0 \left(\frac{m}{T} \right) + \frac{1}{8} \frac{\lambda}{16\pi^2} \left\{ \left[\ln \left(\frac{\mu^2}{m^2} \right) + 1 \right] m^2 - T^2 J_1 \left(\frac{m}{T} \right) \right\}^2, \quad (2.4)$$

where we have explicitly separated the thermal and non-thermal contributions for later convenience. Here and in all related renormalized expressions below, μ represents the arbitrary renormalization scale introduced by dimensional regularization in the $\overline{\text{MS}}$ -scheme, and $\lambda \equiv \lambda(\mu)$. Note carefully in Eq. (2.4) that $\mathcal{E}_0$ represents a possible *finite* vacuum energy term which is usually ignored, i.e. minimally set to zero in the (thermal) literature [7]. However, within our approach this quantity is necessarily nonzero and plays a crucial role as will become clear in the sequel.

The standard (dimensionless) thermal integrals appearing in Eq. (2.4) are given by

$$J_n(x) = 4 \frac{\Gamma[1/2]}{\Gamma[5/2 - n]} \int_0^\infty dt \frac{t^{4-2n}}{\sqrt{t^2 + x^2}} \frac{1}{e^{\sqrt{t^2 + x^2}} - 1}, \quad (2.5)$$

where $t = p/T$ and $x = m/T$. Different integrals can be easily related by employing derivatives such as

$$J_{n+1}(x) = -\frac{1}{2x} \frac{\partial}{\partial x} J_n(x). \quad (2.6)$$

Also, a high- T expansion such as

$$J_0(x) \simeq \frac{16}{45} \pi^4 - 4 \frac{\pi^2}{3} x^2 + 8 \frac{\pi}{3} x^3 + x^4 \left[\ln \left(\frac{x}{4\pi} \right) + \gamma_E - \frac{3}{4} \right] + \mathcal{O}(x^6), \quad (2.7)$$

is often useful since it represents a rather good approximation as long as $x \lesssim 1$, i.e., T larger than m .

Finally, for later comparison we recall that the thermal screening mass (m_D), defined [27] by the pole of the (static) propagator, is obtained for weak coupling for the massless theory as a perturbative series which to lowest orders reads [27]:

$$\frac{m_D^2}{T^2} = \frac{\lambda}{24} \left\{ 1 - \sqrt{6} \left(\frac{\lambda}{16\pi^2} \right)^{1/2} - \left(\frac{\lambda}{16\pi^2} \right) \times \left[3 \ln \frac{\mu}{2\pi T} - 2 \ln \frac{\lambda}{16\pi^2} - 6.4341 \right] + \mathcal{O}(\lambda^{3/2}) \right\}. \quad (2.8)$$

III. RG INVARIANT FREE ENERGY IN MASSIVE RENORMALIZATION SCHEMES

We now discuss the lack of RG invariance when $\mathcal{E}_0$ is minimally set to zero in Eq. (2.4). Remark first that to obtain (2.4), calculations have been performed with an arbitrary mass m in the dressed propagators (mainly to subsequently treat it variationally in the OPT/SPT approach), with no prejudice at this stage that it should be a thermal mass of order $m^2 \sim \lambda T^2$ in the actually massless theory. Thus from the RG viewpoint, everything in (2.4) should behave as if it was a genuine massive theory, in particular the mass should have its standard anomalous dimension. Recall that the (homogeneous) RG operator acting on a physical quantity with mass dependence, such as the free energy in the present case, is defined as

$$\mu \frac{d}{d\mu} = \mu \frac{\partial}{\partial \mu} + \beta(\lambda) \frac{\partial}{\partial \lambda} + \gamma_m(\lambda) m \frac{\partial}{\partial m}, \quad (3.1)$$

where our normalization for the β function is

$$\beta(\lambda) \equiv \frac{d\lambda}{d \ln \mu} = b_0 \lambda^2 + b_1 \lambda^3 + \dots \quad (3.2)$$

while for the anomalous mass dimension it is given by

$$\gamma_m(\lambda) \equiv \frac{d \ln m}{d \ln \mu} = \gamma_0 \lambda + \gamma_1 \lambda^2 + \dots \quad (3.3)$$

with [28]

$$(4\pi)^2 b_0 = 3; \quad (4\pi)^2 \gamma_0 = \frac{1}{2}; \\ (4\pi)^4 b_1 = -\frac{17}{3}, \quad (4\pi)^4 \gamma_1 = -\frac{5}{12}. \quad (3.4)$$

It is easy to see that the renormalized expression, Eq. (2.4), requires a finite $\mathcal{E}_0$ contribution to be RG-invariant, as one can readily see by considering the one-loop term which has an explicit $\ln \mu$ dependence. Thus, acting with the RG operator, Eq. (3.1), on the right-hand side (RHS) of Eq. (2.4) gives a nonzero contribution of order $\mathcal{O}(1)$: $-(1/2)m^4$, which is not compensated by terms in Eq. (3.1) coming from the lowest orders in $\beta(\lambda)\partial_\lambda$ or $m\gamma_m(\lambda)\partial_m \propto \lambda m^4$, those being at least of next order $\mathcal{O}(\lambda)$. This is a manifestation of the fact that (perturbative) RG invariance generally occurs from cancellations between terms coming from RG coefficients at order λ^k and the explicit μ dependence at the next order λ^{k+1} . This can also be understood alternatively by considering solely the original bare contribution to the free energy: although the latter only depends on the RG-invariant bare mass and coupling m_0, λ_0 (and on $2\epsilon = 4 - D$ in dimensional regularization), its *finite* part is not *a priori separately* RG invariant. In other words for a massive theory the T -independent vacuum energy divergences cannot be absorbed by an arbitrary redefinition of the vacuum energy without spoiling RG-invariance. Now, as we recall below the subtraction needed to recover RG invariance is perturbatively well defined and easy to construct order by order. The vacuum energy gets its own anomalous dimension which, within dimensional regularization, is essentially determined by the coefficients of the poles in $2\epsilon = 4 - D$, stemming from the remaining divergences once the mass and the coupling have been properly renormalized. This procedure had been exploited in an earlier application of the OPT to evaluate the vacuum energy of the Gross-Neveu (massive) model [29] and then extended to the QCD case [30,31]. Similarly, a well-known related result is that the Coleman-Weinberg effective potential for a general massive theory is not RG invariant without *finite* “vacuum energy” terms independent of the fields, as was originally carried out in Ref. [32] and in the $\overline{\text{MS}}$ -scheme in the context of RG-improvements of the effective potential [33]. Indeed, for the $O(N)$ ϕ^4 model, the vacuum energy anomalous dimension has even been computed up to four- and five-loop order in Ref. [34].

However, to the best of our knowledge, this point appears to have been overlooked in the context of thermal theories. In applications of improved/resummed massive perturbation schemes such as SPT [6], HTLpt [9], and the standard OPT [14] the calculations are mostly performed within the $\overline{\text{MS}}$ -scheme and the $T = 0$ vacuum energy divergence is minimally cancelled out by appropriate (zero

point) counterterms but missing out those extra finite subtractions required by RG properties. In fact, as far as the purely perturbative massless theory is concerned, the only mass is actually a thermal mass: $m_{\text{th}}^2 \sim \lambda T^2$, so that the lack of RG invariance pointed out above is rather postponed to higher (three-loop) perturbative order λ^2 , where it plainly resurfaces. Within the SPT, or the similar HTLpt, approaches the variational mass parameter is similarly perturbatively of order $m^2 \sim \lambda T^2$. It is thus not surprising that the scale dependence observed within SPT/HTLpt results appears to worsen at higher orders [7,9]. But more generally one wishes to use the nonperturbative mass gap resulting from such variational approaches possibly beyond standard perturbation for moderately large coupling values, as can be relevant near a critical temperature. Thus, the lack of RG invariance appears more serious since as we recall in next section [see Eq. (4.1)], the variational procedure formally treats the mass to be of the same perturbative order as the lowest order considered contributions, like e.g. the “hard” thermal one-loop contributions of order $\sim \lambda^0 T^4$. Moreover, in the standard procedure one makes the arbitrary renormalization scale μ effectively temperature dependent by choosing $\mu \sim 2\pi T$, such as to avoid large $\ln \mu / (2\pi T)$ contributions coming from the remnant scale-dependence. In this way the pressure can be studied as a function of T/T_c in QCD applications, where T_c is related to the basic QCD Λ_{QCD} , e.g. in the $\overline{\text{MS}}$ -scheme. But if the scale dependence appears not much reliable at higher loop orders, one may question as well the reliability of the corresponding T/T_c dependence of the pressure, even for the well-motivated central $\mu = 2\pi T$ prescription.

While those issues in $\overline{\text{MS}}$ or related schemes may perhaps not explain at once all the problems that thermal theories face with perturbative expansions at increasing coupling, a part of those problems are likely to be reduced if one adopts from the beginning a prescription fully consistent with RG properties. This problem appears partly circumvented (but are actually rather delayed to higher orders) in thermal perturbative calculations performed in some other renormalization schemes, where the zero-point energy, $\mathcal{F}_0(T=0)$, is subtracted for convenience prior to any subsequent calculations. Indeed, subtracting the $T=0$ contribution from Eq. (2.4) washes out all the first μ -dependent terms, making scale-independence (trivially) satisfied at one-loop order, and (less trivially) at the two loop level as well. But then the one-loop result becomes also trivial, with the only left contribution being the pure thermal, third term in the RHS of Eq. (2.4). So, there are no possible optimized solutions of the OPT/SPT/HTLpt form, which can only be obtained at the two-loop level. Moreover, applying subsequently the standard SPT/OPT procedure anyway spoils RG invariance. In any case it is most convenient to have a prescription generically valid both for zero and finite temperatures, so that subtracting $T=0$ contributions is not satisfactory for a more general

framework. Accordingly, the subtraction procedure we will consider next only depends on $T=0$ contributions but is generically valid also for $T \neq 0$. Moreover, a remarkable consequence is that the subsequent mass optimization, as implied by RG OPT, will give a nontrivial solution already at the one-loop order, and very similar to what is normally obtained at two-loop order with the other mentioned resummation schemes (SPT/OPT, HTLPT, 2PI,...), as we will examine in detail.

Following Refs. [16,17,29–31] the easiest way to construct an RG-invariant finite vacuum energy is to determine $\mathcal{E}_0$ order by order as a perturbative series from the reminder of acting with Eq. (3.1) on the non RG-invariant finite part of Eq. (2.4):

$$\begin{aligned} \mu \frac{d}{d\mu} \mathcal{E}_0(\lambda, m) &\equiv -\text{Remnant}(\lambda, m) \\ &= -\mu \frac{d}{d\mu} [\mathcal{F}_0(\mathcal{E}_0 \equiv 0)|_{\text{finite}}] \end{aligned} \quad (3.5)$$

where the RHS of (3.5) thus defines the anomalous dimension of the vacuum energy. As above mentioned, it is easy to see that, as a perturbative series, $\mathcal{E}_0$ has the convenient form in $\overline{\text{MS}}$ or similar schemes

$$\mathcal{E}_0(\lambda, m) = -\frac{m^4}{\lambda} \sum_{k \geq 0} s_k \lambda^k, \quad (3.6)$$

where the constant coefficients s_k are perturbatively determined order by order, being essentially determined by the coefficients of the (single) powers of $\ln \mu$ term at order $k+1$ (or equivalently by the single poles in $1/\epsilon$ of the unrenormalized expression) [17,30]. This procedure leaves non RG-invariant remnant terms of perturbative higher orders to be cared for similarly once higher order terms are considered. The apparently odd divergent behavior for $\lambda \rightarrow 0$ of this first order term is actually not a problem since, as we will see explicitly, it completely disappears from the final results.

We stress that all the previous considerations, being only dependent on the renormalization procedure, do not depend on the thermal contribution so that, at arbitrary perturbative orders, the subtraction function represented by Eq. (3.6) can be determined simply from the $T=0$ contributions only.¹ Note that Eq. (3.6) is not the only possible subtraction form in general, but a very convenient one in $\overline{\text{MS}}$ or related schemes to proceed systematically at higher orders.

¹However, when the $T=0$ and the $T \neq 0$ contributions are not explicitly separated, like in the case of two- and three-loop HTLpt [9], due to the systematic m/T expansion making such involved calculations tractable, caution will be needed to expand at a sufficient order in m/T so as to get all the relevant terms of the same perturbative order to construct the corresponding subtractions in Eq. (3.6).

Moreover, it is particularly convenient once introducing below the RGPT modification of perturbation, since the $1/\lambda$ term will be responsible for a nontrivial RGPT solution already at one-loop order.

Explicitly, one finds the RG-invariant form of Eq. (2.4) up to two-loop order with a nontrivial $\mathcal{E}_0$ given by

$$\mathcal{E}_0 = -m^4 \left[\frac{s_0}{\lambda} + s_1 + s_2 \lambda + \mathcal{O}(\lambda^2) \right]. \quad (3.7)$$

After some algebra one obtains

$$\begin{aligned} s_0 &= \frac{1}{2(b_0 - 4\gamma_0)} = 8\pi^2; & s_1 &= \frac{(b_1 - 4\gamma_1)}{8\gamma_0(b_0 - 4\gamma_0)} = -1; \\ s_2 &= \frac{96\pi^2(b_0 - 128\pi^2((1 + 4s_1)\gamma_1 - s_0(b_2 - 4\gamma_2)) - 41)}{12288\pi^4(b_0 + 4\gamma_0)} = \frac{23 + 36\zeta[3]}{480\pi^2} \simeq 0.01399, \end{aligned} \quad (3.8)$$

where the explicit RG dependence in the intermediate terms emphasizes the more general form of these results, while the last terms are specific to the $N = 1$ ϕ^4 theory. To derive s_2 according to the previous discussion we had to use the $(T = 0)$ $\ln \mu$ coefficient at three-loop order given e.g. in Ref. [7].

One may equivalently derive the finite subtraction in Eq. (3.7) in an alternative manner by RG invariance considerations solely on the bare expression of the free energy. For completeness, this is presented in the appendix. Instead of minimally subtracting the bare vacuum energy divergence, an RG-invariant counterterm can be added to cancel the remnant divergences, and necessarily incorporates also the same finite subtraction terms in Eq. (3.8). As a nontrivial crosscheck of our calculation, let us note that now acting with the RG operator, Eq. (3.5), on the results given by Eqs. (3.7) and (3.8) one recovers, for $N = 1$, the results up to λ^3 of the anomalous dimension $\beta_v(\lambda)$ [with $\mu d\mathcal{E}_0/d\mu \equiv 2m^4\beta_v(\lambda)$] which has been calculated up to four and five loops for arbitrary N in Ref. [34]. (Actually we could have used directly the results in [34] in the present scalar model case to derive the s_k in Eq. (3.8), but the above derivation using the basic available perturbative expressions shows precisely how to proceed for an arbitrary theory, where the vacuum energy anomalous dimension may not always be explicitly available.)

One sees that the subtraction with s_k , explicitly depending on RG coefficients, incorporates a nontrivial RG-dependence already at first (one-loop) order, only depending on already known one-loop standard RG coefficients. This result has important consequences for the subsequent OPT application. Now, there is a subtlety at this stage: while the s_k subtraction terms are strictly necessary to recover RG invariance at order λ^k , i.e. up to neglected λ^{k+1} terms, they enter the free energy expression at order λ^{k-1} as Eq. (3.6) indicates. For instance only s_0 is needed to recover RG-invariance at one-loop $\mathcal{O}(1)$, but the next term s_1 is of order $\mathcal{O}(1)$, so strictly speaking s_1 should also be included in the full “one-loop” free energy results. This appears as a complication *a priori*, meaning that at order k one needs

in principle the more demanding information from (the $\ln \mu$ coefficient of) perturbative order $k + 1$. On the other hand since RG invariance is constructed perturbatively, one may expect that the simplest minimal prescription of keeping only the s_k terms at order λ^k should already be a good enough approximation. Accordingly, we mainly consider below the simplest prescription but also examine both prescriptions, indicating the differences whenever relevant. We will see that, after the modification of perturbative expansion implied by RGPT, incorporating the higher order s_{k+1} at order λ^k makes no crucial differences, even at one-loop order, the resummation results being not very sensitive to such purely perturbative variations. The same stability with respect to such variations was also observed at vanishing temperature in Ref. [17] (where those different prescriptions were incorporated within the intrinsic theoretical errors of the method).

It should be clear from the previous derivation that by construction the subtraction terms make the free energy perturbatively RG-invariant. But just to crosscheck it in a more pedestrian way, let us now reexamine the result at one-loop order, with the $-s_0 m^4/\lambda$ subtraction included in Eq. (2.4), using the standard one-loop RG running coupling and mass. These are given from integrating respectively Eq. (3.2), (3.3) which yields the standard textbook result:

$$\begin{aligned} \lambda(\mu) &= \lambda(\mu_0) \left(1 - b_0 \lambda(\mu_0) \ln \frac{\mu}{\mu_0} \right)^{-1} \\ &\simeq \lambda(\mu_0) \left(1 + \frac{3}{16\pi^2} \lambda(\mu_0) \ln \frac{\mu}{\mu_0} + \mathcal{O}(\lambda^2) \right), \end{aligned} \quad (3.9)$$

$$\begin{aligned} m(\mu) &= m(\mu_0) \left(1 - b_0 \lambda(\mu_0) \ln \frac{\mu}{\mu_0} \right)^{-1/6} \\ &\simeq m(\mu_0) \left(1 + \frac{1}{32\pi^2} \lambda(\mu_0) \ln \frac{\mu}{\mu_0} + \mathcal{O}(\lambda^2) \right), \end{aligned} \quad (3.10)$$

where $\lambda(\mu_0)$ and $m(\mu_0)$ are the coupling and mass at some arbitrary reference scale μ_0 . Expanding Eq. (2.4) to first order in λ (i.e. to order λ^0) one obtains:

$$\begin{aligned}
(4\pi)^2 \mathcal{F}_0 &\simeq m^4(\mu_0) \left[-\frac{8\pi^2}{\lambda(\mu_0)} - \frac{1}{8} (4 \ln \mu + 3 + \dots) + 8\pi^2 \left(\frac{3}{16\pi^2} \ln \mu - \frac{2}{16\pi^2} \ln \mu \right) + \mathcal{O}(\lambda) \right] + \text{thermal part} \\
&\simeq m^4(\mu_0) \left[-\frac{8\pi^2}{\lambda(\mu_0)} - \frac{3}{8} + \mathcal{O}(\lambda) \right],
\end{aligned} \tag{3.11}$$

(where $\dots$ stands for μ -independent terms). This relation explicitly displays the cancellation, up to terms of higher order $\mathcal{O}(\lambda)$, of all $\ln \mu$ contributions coming respectively from the original one in Eq. (2.4), and from the running coupling and mass [third and fourth terms respectively in Eq. (3.11)]. We have neglected above the thermal contributions, but including these does not alter the results since the exact T -dependent contribution does not depend explicitly on μ . Alternatively, when the high- T expansion is considered, the explicit $\ln \mu/m$ in Eq. (2.4) is replaced by a $\ln \mu/(4\pi T)$ with the same coefficient consistently. It is instructive to push this exercise a step further now incorporating in the free-energy (2.4), restricted at one-loop order, all thermal contributions in the high- T limit, from Eq. (2.7), and plugging into the resulting expression $m = m_D$, the standard perturbative thermal mass, Eq. (2.8) also restricted at first order. Then, the expression only depends on the coupling, and using Eq. (3.9) gives the result:

$$\begin{aligned}
(4\pi)^2 \mathcal{F}_0 &\simeq T^4 \left[-\frac{8\pi^2}{45} + \frac{\pi^2}{72} \lambda(\mu_0) - \frac{\pi}{36\sqrt{6}} \lambda^{3/2}(\mu_0) + \mathcal{O}(\lambda^2, \ln \mu) \right],
\end{aligned} \tag{3.12}$$

where all μ dependence has been canceled out up to order λ^2 , since $m_D^4 \sim \lambda^2$. This was expected since the subtraction takes care of the $T = 0$ lowest orders μ -dependence as shown previously, while the thermal contribution $\propto J_0(m/T)$ in Eq. (2.4) does not depend on the scale. More interestingly, one recognizes from Eq. (3.12) the standard perturbative expansion for the pressure which, upon using the common normalization with the free gas pressure $P_0 = \pi^2 T^4/90$ and $\lambda_0 = 4!g^2$, can be expressed as:

$$\frac{P}{P_0} = 1 - \frac{15}{8} \frac{g^2}{\pi^2} + \frac{15}{2} \frac{g^3}{\pi^3}. \tag{3.13}$$

In particular, the term $-s_0/\lambda$ is incorporated in the final result Eq. (3.12) once using Eq. (2.8) since $-s_0 m_D^4/\lambda \sim \mathcal{O}(\lambda)$. The latter gives a contribution $(15/8)g^2$ to P/P_0 , making this complete one-loop expression consistent with the standard perturbative expression of the pressure up to order $\lambda^{3/2}$, while the original (unsubtracted) perturbative one-loop expression is not, giving an expansion similar to (3.13) but with a twice

too large g^2 term $-(15/4)g^2$. However, this agreement is merely an accident of one-loop order: at two-loop order, including the next order subtraction term from (3.7) does not give the correct *massless* perturbative pressure when replacing m by m_D . (In particular the subtraction $-s_0 m^4/\lambda$ happens to be exactly cancelled by the hard contribution of order λ , $\propto \lambda T^4$). But this is not surprising, since more generally the perturbative *massless* pressure [27] cannot be obtained consistently from simply replacing the perturbative thermal mass (2.8) in the expression of the *massive* pressure. However, when the mass is traded for a variational parameter, like in the OPT/SPT resummation approaches to be recalled next, one may recover the massless pressure results under specific conditions. This is made possible because the OPT construction and mass optimization drastically modifies the massive contributions as compared with the original perturbative expansion.

To summarize this section, starting with Eq. (2.4) including $\mathcal{E}_0$ from Eq. (3.7) to obtain a perturbatively RG-invariant free energy provides a sound basis for more elaborate resummation procedures like the OPT method to be addressed next. As above mentioned, an extra advantage of the subtraction terms (3.7) starting with $1/\lambda$, is that the optimization procedure will provide a nontrivial mass $\tilde{m}(\lambda)$ already at the lowest one-loop order, in contrast with the standard SPT and HTLpt approaches (where at one-loop order the mass optimization gives a trivial λ -independent solution [6–8]). This is a welcome feature specially for more involved theories like thermal QCD where higher order contributions are challenging to evaluate, and a comparison between successive orders is crucial to establish the stability of the resummation results.

IV. OPTIMIZED PERTURBATION THEORY (OPT)

The basic feature of the optimized perturbation theory (OPT) (appearing also under different names and variations [11–13]), is to introduce an extra parameter $0 < \delta < 1$, which interpolates between $\mathcal{L}_{\text{free}}$ and $\mathcal{L}_{\text{int}}$ in Eq. (2.1), so that the mass m is traded for an arbitrary trial parameter. This is perturbatively equivalent to taking any standard perturbative expansions in $\lambda(\mu)$, after renormalization, reexpanded in powers of δ after substituting:

$$m \rightarrow m(1 - \delta)^a, \quad \lambda \rightarrow \delta \lambda. \tag{4.1}$$

This procedure is consistent with renormalizability [29,35,36] and gauge invariance [31], whenever the latter is relevant, provided of course that the above redefinition of

the coupling is performed consistently for all interaction terms and counterterms appropriate for renormalizability and gauge invariance in a given theory.² Note that in Eq. (4.1) we have introduced an extra parameter, a , to reflect *a priori* a certain freedom in the interpolation form. As will be demonstrated below this parameter plays an essential role within our method for allowing compelling constraints to be imposed. Applying Eq. (4.1) to some given renormalized perturbative expansion for a physical quantity, $P(m, \lambda)$, reexpanding in δ to order k , and taking *afterwards* the $\delta \rightarrow 1$ limit (to recover the original *massless* theory) leaves a remnant m -dependence at any finite δ^k -order. The arbitrary mass parameter m is then most conveniently fixed by a variational optimization prescription known as the principle of minimal sensitivity [13]

$$\frac{\partial}{\partial m} P^{(k)}(m, \lambda, \delta = 1)|_{m \equiv \tilde{m}} \equiv 0, \quad (4.2)$$

thus determining a nontrivial optimized mass $\tilde{m}(\lambda)$, with nonperturbative λ -dependence, realizing dimensional transmutation (more precisely, e.g. for asymptotically free theories at vanishing temperatures, the optimized mass is automatically of the order of the basic scale $\Lambda \sim \mu e^{-1/(b_0 \lambda)}$, in contrast with the original vanishing mass).

In simpler ($D = 1$) models, at vanishing temperatures, the procedure may be seen as a particular case of “order-dependent mapping” [12], which has been proven [37] to converge exponentially fast for the $D = 1$ ϕ^4 oscillator energy levels. For higher dimensional $D > 1$ renormalizable models, no rigorous convergence proof exists, although the OPT was shown to partially damp the factorially divergent (infrared renormalons) perturbative behavior at large orders [38]. Nevertheless, this technique can give rather successful approximations to some nonperturbative quantities beyond the large- N (or mean field) approximations in a large variety of physical situations which include the study of phase transitions within condensed matter related renormalizable models [39–42] as well as within QCD nonrenormalizable effective models [43].

We emphasize that at finite temperatures the very same basic idea has been exploited by the SPT [5,6]/HTLpt [8] method, where in this thermal context the screening thermal

mass is treated as an arbitrary variational parameter, and in Eq. (4.2) P also depends on T like e.g., Eq. (2.4).

V. RENORMALIZATION GROUP COMPATIBILITY OF OPT

In most previous standard OPT (or similarly SPT and HTLpt) applications, the so-called linear δ -expansion is used, assuming $a = 1/2$ (i.e. $m^2 \rightarrow m^2(1 - \delta)$ for a scalar mass) in Eq. (4.1) mainly for simplicity and economy of parameters while the more recent approach, developed in Refs. [15–17], differs in two crucial aspects which turn out to drastically improve the convergence. First, it introduces a straightforward marriage between OPT and renormalization group (RG) properties, by requiring the (δ -modified) expansion to satisfy, in addition to the OPT Eq. (4.2), a standard RG equation:

$$\mu \frac{d}{d\mu} P^{(k)}(m, \lambda, \delta = 1) = 0, \quad (5.1)$$

where the RG operator was defined in Eq. (5.1). Moreover, once combined with Eq. (4.2), the RG equation takes the reduced massless form:

$$\left[\mu \frac{\partial}{\partial \mu} + \beta(\lambda) \frac{\partial}{\partial \lambda} \right] P^{(k)}(m, \lambda, \delta = 1) = 0. \quad (5.2)$$

Therefore, Eqs. (5.2) and (4.2) if used together, completely determine *optimized* $m \equiv \tilde{m}$ and $g \equiv \tilde{g}$ “variational” fixed point values.

Since interaction and free terms from the original perturbative series are rather drastically reshuffled by the modification implied by Eq. (4.1), the RG invariance is in general no longer perturbatively satisfied, even when the original perturbative series is RG-invariant prior to performing (4.1). This spoiled RG invariance has to be restored in some manner, and thus Eq. (5.1) gives a nontrivial additional constraint. This feature has been seldom appreciated and considered in many former applications of the δ -expansion/OPT method to renormalizable theories (perhaps in part because in many analyses with more elaborated theories the OPT is restricted to first order, where RG improvements are supposed to play a minor role). This important role of RG properties was recognized much earlier in Refs. [29,30] where to recover the RG consistency the standard linear δ -expansion was resummed to all orders. Indeed, this resummation can be done, at least for the pure RG dependence up to two-loop, but the result comes as a rather involved integral representation, not practically intuitive and making difficult to perform the mass optimization or to generalize to other physical quantities and other models. In contrast, the purely perturbative procedure together with Eq. (5.1) appears a considerable shortcut, straightforward to apply to any model. Intuitively, just as the stationary point solutions

²Contrary to what is sometimes claimed (and worked out) in the literature, the OPT/SPT/HTLpt does not need any extra counterterms besides the standard ones of the corresponding massive theory: in particular all seemingly new divergences generated at arbitrary orders from the first replacement in Eq. (4.1) are evidently related to the single standard mass counterterm. Moreover at arbitrary orders, temperature-dependent divergences and associated counterterms should not appear, as expected from general principles, provided that one keeps the mass as an arbitrary parameter and carefully subtract all nested subdivergences until renormalization has been completed, before using a gap-equation giving perturbatively $\tilde{m} \propto T\sqrt{\lambda}$.

from Eq. (4.2) are expected to give sensible approximations, at successive orders, to the actually massless theory, one similarly expects that combining the latter with the RG solutions should further give a sensible sequence of best approximations to the exactly scale invariant all order results.

Still, a well-known drawback of the standard OPT approach is that, beyond lowest order, solving Eq. (4.2) generally gives more and more solutions at increasing orders, some of which are likely to be complex-valued. More generally, without some insight on the nonperturbative behavior of the solutions, it may be difficult to select the right one, and the unphysical complex-valued optimized solutions at higher orders are embarrassing. This is incidentally a problem encountered first at three-loop order in SPT [7] and HTLpt applications to QCD[9]. The mass optimization is then replaced by alternative prescriptions, most often using simply the purely perturbative screening mass, but accordingly losing a more nonperturbative ingredient from the optimized mass. But RG considerations also provide a possible way out, which is the second main difference and new feature of the present RGOPT version. For QCD a compelling selection criterion was proposed, in which only the solution(s) *continuously matching* the standard perturbative (asymptotically free) RG behavior for vanishing coupling are retained [16,17]. This prescription can easily be generalized to any model, like the nonasymptotically free (AF) ϕ^4 theory, by similarly requiring to asymptotically match the solutions to the standard perturbative behavior for small coupling, namely for fixed $\tilde{m}$ and arbitrary scale μ :

$$\tilde{\lambda}(\mu \ll \tilde{m}) \sim \left(b_0 \ln \frac{\tilde{m}}{\mu}\right)^{-1} + \mathcal{O}\left(\ln^{-2}\left(\frac{\tilde{m}}{\mu}\right)\right). \quad (5.3)$$

At zero temperature this turns out to give a unique solution for both the RG and OPT equations, up to rather high orders. An additional welcome feature is that by requiring at least one RG solution to fulfill Eq. (5.3) leads to a strong necessary condition on the basic interpolation, Eq. (4.1), uniquely determining a from the universal (scheme-independent) first order RG coefficients: $a \equiv \gamma_0/b_0$, as we derive in more detail below. A connection of the OPT exponent a with RG anomalous dimensions/critical exponents had also been established in a very different context, in the $D = 3$ ϕ^4 model for the Bose-Einstein condensate (BEC) critical temperature shift by two independent OPT approaches [41,42], where it also led to real OPT solutions [42]. However, AF-compatibility and reality of solutions can appear to be mutually exclusive beyond lowest order, depending on the particular model. A simple way out is to further exploit the RG freedom, considering a perturbative renormalization scheme change to attempt to recover RGOPT solutions both AF-compatible and real [17]. We will see that this extra complication is not even necessary in

the present case, where (at least up to the two-loop order) the RG-compatible solutions remain real for a large range of relevant values of the coupling and temperature. All these features are easy to generalize at finite temperatures due to the fact that RG properties are essentially determined by the divergence structure of the $T = 0$ part. So, the only complication is technical since at finite temperature the previous Eqs. (4.2), (5.1), and (5.2) come with an extra T dependence. Let us now illustrate explicitly all those features by evaluating the RGOPT modification of the free energy of a thermal scalar field.

VI. 1-LOOP, $\mathcal{O}(\delta^0)$

A. $T = 0$

Let us first truncate Eq. (2.4) at strict one-loop order, and first restricting to $T = 0$ which is simpler and sufficient to determine the RG-exponent a in Eq. (4.1). We have

$$(4\pi)^2 \mathcal{F}_0^{\text{RGI}}(T = 0) = -\frac{s_0}{\lambda} m^4 - \frac{1}{8} m^4 \left(3 + 2 \ln \frac{\mu^2}{m^2}\right), \quad (6.1)$$

where the superscript RGI emphasizes the (perturbative) RG invariance of this quantity. At this order the calculation is elementary so it can best illustrate the main steps. Now, applying Eq. (4.1), performing the δ -expansion to order δ^0 consistently, and taking *afterwards* $\delta \rightarrow 1$, gives:

$$(4\pi)^2 \mathcal{F}_0^{\text{RGI}}(\delta^0, \delta = 1) = m^4 \left[-\frac{s_0}{\lambda} (1 - 4a) - \frac{1}{8} \left(3 + 2 \ln \frac{\mu^2}{m^2}\right) \right]. \quad (6.2)$$

Note that the $\mathcal{O}(1)$ term remained unmodified (this is a general property of OPT: expanding at order δ^k and taking $\delta \rightarrow 1$ leaves the order λ^k term unaffected due to the screening from $\lambda \rightarrow \delta\lambda$). Then, requiring Eq. (6.2) to be perturbatively RG-invariant after this modification of the perturbative series, i.e. applying the RG Eq. (5.2), on gets

$$m^4 \left[\left(1 - \frac{b_0}{\gamma_0} a\right) + \mathcal{O}(\lambda) \right] = 0, \quad (6.3)$$

which uniquely fixes

$$a = \frac{\gamma_0}{b_0} = \frac{1}{6}, \quad (6.4)$$

where, in Eqs. (6.3) and (6.4), we made the RG coefficient dependence explicit to emphasize the generality of these results. At this point several remarks are in order:

- (i) the very same result, Eq. (6.4), was obtained [16,17] (up to a trivial b_0 difference of normalization by a factor 2), while considering the RGOPT for QCD (with appropriate QCD values for those RG

coefficients). The exponent a is universal for a given model, in the sense that it only depends on the first-order RG coefficients, which are renormalization scheme independent. Furthermore, at vanishing temperature, Eq. (6.4) greatly improves the convergence of the procedure at higher orders: considering only the first RG coefficients b_0, γ_0 dependence (i.e., neglecting higher RG orders and non-RG terms), it gives the exact nonperturbatively resummed result at the first δ order and any successive order [17]. This is not the case for $a = 1/2$ (for a scalar mass), where the convergence appears very slow, if any.

- (ii) The standard linear δ -expansion interpolation, widely used for zero temperature models and for SPT/HTLpt, takes $a = 1/2$ [7,11,14,36]:

$$m^2 \rightarrow m^2(1 - \delta), \quad (6.5)$$

thus our RG-compatible exponent (6.4) is three times smaller.³ Indeed the standard OPT/SPT interpolation (6.5) leads to an unmatched RG equation, while the OPT equation (4.2) solved for $\lambda(m)$ (or equivalently for $m(\lambda)$ gives, in the perturbative regime $\mu \ll m$:

$$\lambda(\mu \ll m) \sim -\frac{16\pi^2}{\ln(\frac{m}{\mu})}, \quad (6.6)$$

in clear contradiction with the true running by a wrong overall sign plus a factor three too large.

- (iii) At the very first nontrivial δ^0 order, once having fixed $a = \gamma_0/b_0$ the RG equation is satisfied and thus does not give further constraint. We will see that at the next and higher orders in δ , Eq. (6.4) is always required for the RG equation to have at least one solution matching Eq. (5.3). In addition, it also fixes λ in terms of the other parameters (m , and the only remaining parameter μ/T when considering the thermal part).

Let us next consider the other OPT constraint given by Eq. (4.2). Still neglecting the thermal part, and pulling out an overall factor yields

$$m^3 \left[\frac{1}{b_0 \lambda} + \frac{1}{2} \left(1 + \ln \frac{\mu^2}{m^2} \right) \right] = 0. \quad (6.7)$$

One readily remarks the explicit exact scale-invariance of Eq. (6.7), thus of its solution, provided that one uses for $\lambda \equiv \lambda(\mu)$ the exact (one-loop resummed) running in Eq. (3.9), since the expression $1/\lambda(\mu) + b_0 \ln \mu$ is explicitly μ -independent. Letting apart the trivial $m = 0$ solution, Eq. (6.7) has the more interesting solution

³In QCD e.g for 3 light quark flavors $a = \gamma_0/(2b_0) = 4/9$ is also substantially smaller than the linear case value $a = 1$ for fermion masses.

$$\tilde{m}^2 = \mu^2 e^{1 + \frac{2}{b_0 \lambda}}, \quad (6.8)$$

which is seen to be compatible, for $\lambda \rightarrow 0$, with the perturbative first order standard RG solution obtained from solving Eq. (3.2) for $\lambda(\mu)$ at first order. Namely, for fixed m and arbitrarily small μ , it exhibits infrared freedom: $\lambda(\mu \ll m) \simeq (b_0 \ln(m/\mu))^{-1}$.

Moreover, plugging Eq. (6.8) within the modified vacuum energy expression (6.2) for the case of vanishing temperatures, gives a remarkably simple result:

$$\mathcal{F}_0(\tilde{m}, \lambda) = -\frac{\tilde{m}^4}{8(4\pi)^2}. \quad (6.9)$$

Despite its somewhat trivial look, Eq. (6.9) represents a nonperturbative result, in the sense that it only involves the expression of $\tilde{m}^4$ from Eq. (6.8). Apart from the one in Eq. (6.8) all λ -dependence disappeared [in particular the $-s_0/\lambda$ term has consistently disappeared upon using the OPT gap equation (6.7)]. Therefore, Eq. (6.9) gives a nontrivial $T = 0$ (negative) vacuum energy contribution and resembles much the large- N nonperturbative result, up to appropriate identification of $b_0 \lambda$, but here obtained from the OPT. When higher RG orders and non-RG contributions are included, they spoil this simple form result [17], as does also the thermal part, that we will consider next. In these cases there are remnant coupling dependences, once having used the solution of Eq. (4.2) within the physical expression of the vacuum energy.

B. $T \neq 0$

Let us now consider the thermal contributions in Eq. (2.4) still at one-loop order. After performing the δ -expansion to the lowest corresponding order δ^0 we obtain the $T \neq 0$ free energy, similarly as in (6.2):

$$(4\pi)^2 \mathcal{F}_0^{\text{RGI}}(T \neq 0, \delta^0, \delta = 1) = m^4 \left[-\frac{1}{2b_0 \lambda} - \frac{1}{8} \left(3 + 2 \ln \frac{\mu^2}{m^2} \right) \right] - \frac{1}{2} T^4 J_0 \left(\frac{m}{T} \right), \quad (6.10)$$

where the first term $-s_0 m^4 (1 - 4a)/\lambda = -m^4/(2b_0 \lambda)$ is the only one affected by (4.1) at this lowest order. Calculations are slightly more involved than for $T = 0$ but note that the δ -expansion and subsequent OPT minimization equation involve successive derivatives of the thermal function $J_n(m/T)$. For more generality and to make contact with various other resummation methods, it turns out to be particularly convenient to express all our RGOPT results in terms of the one-loop renormalized self-energy, including all thermal dependence:

$$\Sigma_R \equiv \frac{\lambda}{2} \int_p \frac{1}{p^2 + m^2} + \Sigma^{\text{ct}} = \gamma_0 \lambda \left[m^2 \left(\ln \frac{m^2}{\mu^2} - 1 \right) + T^2 J_1 \left(\frac{m}{T} \right) \right], \quad (6.11)$$

[where for completeness the mass counterterm reads Σ^{ct} reads $\gamma_0 \lambda m^2 / (2\epsilon)$]. This simple factorization is possible for the scalar ϕ^4 model up to the two-loop level, because the two-loop contribution [the last order λ term in Eq. (2.4)] factorizes as the square of one-loop expressions (i.e. graphs with a different “nested” topology only appear at the three-loop level for the $\lambda\phi^4$ interactions).

Then, noting that $\frac{\partial}{\partial m^2} \int_p \ln(p^2 + m^2) = 2\Sigma_R/\lambda$, the exact solution of the OPT Eq. (4.2) can easily be written in the form of a self-consistent “gap” equation for $\tilde{m}$:

$$\tilde{m}^2 = (4\pi)^2 b_0 \Sigma_R = b_0 \frac{\lambda}{2} \left[\tilde{m}^2 \left(\ln \frac{\tilde{m}^2}{\mu^2} - 1 \right) + T^2 J_1 \left(\frac{\tilde{m}}{T} \right) \right], \quad (6.12)$$

which like the $T = 0$ previous case, is exactly scale-invariant by construction, as we will illustrate more explicitly below.

1. Digression: Connection with large- N and 2PI results

As an important digression, we point out that Eq. (6.12) is recognized as the very same solution obtained for the large- N $O(N)$ ϕ^4 model in Ref. [22], upon appropriate b_0 definition for the large- N case. Indeed, in the leading $1/N$ approximation, the only contributing graphs have the one-loop structure, and the mass-gap equation can be solved exactly. More precisely it is easily found from the arbitrary N RG coefficients given e.g. in Ref. [28] that in our normalization,

$$(4\pi)^2 b_0(N) = \frac{N+8}{3}, \quad (4\pi)^2 \gamma_0(N) = \frac{N+2}{6}, \quad (6.13)$$

so that in particular we have for the crucial exponent in (4.1):

$$a \equiv \frac{\gamma_0}{b_0} = \frac{1}{2} \left(\frac{N+2}{N+8} \right) \rightarrow \frac{1}{2} \quad \text{for } N \rightarrow \infty \quad (6.14)$$

for which values all our previous construction, and the corresponding mass gap equation in (6.12), reproduce exactly the large N results in [22]. Note in particular that $a = 1/6$ for $N = 1$ while accidentally the large N value of $a = 1/2$ is the standard linear one, but here being fully consistent with RG properties. The fact that the one-loop RGOPT reproduces exactly the large N result can be seen as the finite temperature analog of similar RGOPT properties [15] obtained for the large N limit of the GN model.

Similarly, Eq. (6.12) is also recognized as the very same form of mass gap solution obtained in the 2PI formalism but at two-loop order [21] (or also in the tadpole approximation for the self-energy [20]), except that in [21] the correct $b_0 = 3/(16\pi^2)$ is effectively replaced by $b_0/3$ because, as explained by the authors, only one channel out of three is taken into account at this level of the 2PI approximation, similarly to the leading $1/N$ approximation. We will come back below on this apparent b_0 value mismatch when discussing the perturbative reexpansion of the pressure to make contact with standard perturbation results. In fact, the analogy with Ref. [21] goes further, in particular their expression of the mass gap solution is also exactly scale invariant at two-loop order, and the free energy involves, after renormalization, a term $-m^4/(2\lambda)$, once again identical to our subtraction terms $-m^4/(32\pi^2 b_0 \lambda)$ in Eq. (6.2), when taking $b_0 \rightarrow b_0/3$. But the origin of this $1/\lambda$ term in Ref. [21] is very different, emerging upon using the gap equation. Incidentally, the scale-invariance of the 2PI results is essentially due to a renormalization procedure that is peculiar to the scalar model, inspired by the large N limit [22], which thus appears to us as being accidental, and limited to the first nontrivial two-loop level. In contrast the mass gap (6.12) is obtained already at the one-loop level, and it should be clear from the previous construction that the RGOPT systematic procedure is applicable in any model at arbitrary orders.

2. One-loop RGOPT mass gap and pressure solutions

Although Eq. (6.12) may easily be solved numerically, it is instructive to consider next the approximation given by the high- T expansion, which is very precise as long as $T \gtrsim m$: in fact this condition can be easily checked *a posteriori* considering the optimized solution $\tilde{m}$. At one-loop level, it turns out that the optimized mass always satisfies this criterion for all the relevant range of coupling values. Therefore, we will not need to solve the exact Eq. (6.12) for all practical purposes. In the high- T approximation (2.7), the one-loop order OPT Eq. (4.2) produces (discarding the trivial solution $\tilde{m} = 0$) a simple quadratic equation for m :

$$\left(\frac{1}{b_0 \lambda} + L_T \right) x^2 + 2\pi x - 2 \frac{\pi^2}{3} = 0, \quad (6.15)$$

where $m \equiv xT$ and we defined for shorthand notations the μ/T dependent part $L_T \equiv \ln[\mu/(4\pi T)e^{\gamma_E}]$. As already explained above the RG Eq. (5.2) reduces to Eq. (6.3) which is already satisfied for Eq. (6.4), thus it gives no additional constraint.

Solving Eq. (6.15) gives two real solutions, but one is clearly unphysical, giving $\tilde{m} < 0$ for any λ . The other unique physical solution is thus

$$\frac{\tilde{m}^{(1)}}{T} = \pi \frac{\sqrt{1 + \frac{2}{3}(\frac{1}{b_0\lambda} + L_T)} - 1}{\frac{1}{b_0\lambda} + L_T}, \quad (6.16)$$

where for more generality we kept the dependence on b_0 explicit. Despite the apparently minor modification of the series represented by Eq. (6.2) at this first order, the solution given by Eq. (6.16) has clearly a nonperturbative dependence on λ . We stress that the variational mass (6.16) is unrelated to the physical screening mass [27] in Eq. (2.8), and thus has no reason to reproduce the latter. Moreover, as anticipated, Eq. (6.16) is strictly *exactly* scale-invariant, provided that one uses for $\lambda \equiv \lambda(\mu)$ the exact (one-loop resummed) running in Eq. (3.9) (now with μ being temperature-dependent as usual), since the expression $1/\lambda(\mu) + b_0 L_T \equiv 1/\lambda(\mu) + b_0 \ln \mu + \dots$ is explicitly μ -independent. In other words the mass gap in (6.16) actually only depends on the single parameter $b_0 \lambda(\mu_0)$, where μ_0 is some reference scale, typically $\mu_0 = 2\pi T$.

Before we proceed, it is worthwhile to comment a little more on this result: recall that prior to the δ -expansion, the basic one-loop expression [first line in Eq. (2.4)] with the first term s_0 in the subtraction $\mathcal{E}_0$, is by construction RG-invariant to one-loop order, i.e. up to neglected higher order terms $\mathcal{O}(\lambda)$. But what is more remarkable is that the optimal mass resulting from solving (4.2) is *exactly* scale-invariant to all orders (of course “all orders” but neglecting genuine higher orders in the running coupling, i.e. keeping only the b_0 dependence to all orders). This is a direct consequence of the value $a = \gamma_0/b_0$ in the interpolating relation, Eq. (4.1). This result is the finite temperature analog of what was similarly obtained generically for zero temperature QCD in Ref. [17]: namely, “all-order” (one-loop RG) resummed results are correctly obtained by the very first RGOPT δ order. Indeed, since this is a generic result, we anticipate that applying the same procedure to thermal QCD will give similar one-loop results, with an OPT equation and solution very similar to Eqs. (6.15) and (6.16) up to obvious changes in some factors, but also exhibiting exact scale-invariance). However, this exact scale-invariance is due to the form of the exact one-loop running of the coupling, perfectly matching Eqs. (6.15), (6.16), which does not generalize once including higher RG orders in the β function and non-RG dependence at higher orders. As we examine in next section, at the two-loop $\mathcal{O}(\lambda)$ order, the scale invariance resulting from RGOPT extends beyond the two-loop perturbative order at which it is imposed by construction, but a (moderate) scale dependence reappears unavoidably at a *finite* higher perturbative order, precisely at order λ^3 , thus one order higher than naively expected.

To proceed one may expand Eq. (6.16) perturbatively, which is easily seen to be an expansion in $\sqrt{\lambda}$, as expected. One then finds:

$$\begin{aligned} \frac{\tilde{m}^{(1)}}{T} \sim \pi \left[\sqrt{\frac{2}{3}} \sqrt{b_0 \lambda} - b_0 \lambda + \frac{1}{2\sqrt{6}} (3 - 2L_T) (b_0 \lambda)^{3/2} \right. \\ \left. + L_T (b_0 \lambda)^2 + \dots \right], \end{aligned} \quad (6.17)$$

where we kept the b_0 dependence explicit on purpose.

As an important side remark, up to now we have considered the simplest minimal prescription of incorporating only the $-m^4(s_0/\lambda)$ subtraction at one-loop order, strictly necessary for recovering perturbative RG invariance. It is thus opportune to mention what is changing if incorporating the next order subtraction $s_1 \neq 0$ term from (3.7), being formally also of one-loop order. In fact this simply amounts to the replacement $L_T \rightarrow L_T + 2s_1 = L_T - 2$ consistently in all previous expressions (6.15), (6.16), and (6.17), as could be easily traced by consistently introducing $-m^4 s_1$ into Eqs. (6.10), (6.12). Therefore, it means that at one-loop RGOPT order s_1 can be simply absorbed by a change of scale (or renormalization scheme) definition, $\mu \rightarrow \mu e^{2s_1} = \mu e^{-2}$. Thus apart from changing the reference scale with respect to the $\overline{\text{MS}}$ -scheme, it does not really change physical results: if redefining accordingly the coupling with a RG evolution $\mu \rightarrow \mu e^{-2}$, we obtain strictly identical results. With this in mind, for now on we proceed with the simplest choice $s_1 = 0$ at one-loop.

Coming back to Eq. (6.16) it is obviously an expansion solely in the single parameter $(b_0 \lambda)$. Using the b_0 value from Eq. (3.4) we have: $\pi \sqrt{2b_0/3} = 1/(2\sqrt{2})$, which tells that the first order coefficient differs from the standard Debye mass in (2.8), $m_D^2 \sim (\lambda/24)T^2$, being $\sqrt{3}$ larger. This originates directly from the correct value $b_0 = 3/(16\pi^2)$ used in $a = \gamma_0/b_0$ in (4.1), which is the only value compatible with RG invariance. The factor 3 in b_0 is the statistical factor originating from three similar graphs contributing to the β function, as is well known. Thus, the first perturbative coefficient of the Debye screening mass would be obtained from Eq. (6.16) if one would take $b_0/3 = 1/(16\pi^2)$ as given by a single loop contributing to the self-energy at one-loop, as argued in Ref. [21]. The standard perturbative term of order λ , comparing with Eq. (2.8), is also reproduced provided again that one takes $b_0 = 1/(16\pi^2)$ in Eq. (6.16). But, as mentioned above, Eqs. (6.16)–(6.17) reproduce exactly (at arbitrary orders) the large N -results (e.g. Eq. (5.7) of ref. [22]), as can be checked upon identifying the correct large- N value of $b_0 = 1/(16\pi^2)$ in the normalization of [22]. This factor 3 discrepancy in the optimized mass for the $N = 1$ ϕ^4 model from b_0 mismatch is not a problem, since the OPT nonperturbative variational mass, not being a physical parameter, has no physical connection with the perturbative physical screening mass in Eq. (2.8) and is therefore not required to reproduce the latter. Indeed, there are $\ln(\lambda)$ terms appearing at the three-loop order in the genuine

screening mass [3,7], Eq. (2.8), that are not present in the expansion of Eq. (6.16) which only involves λ and $\lambda^{1/2}$ powers. Incidentally the fact that the standard OPT or SPT/HTLpt correctly reproduces the first two orders of the thermal perturbative mass expansion [6,8] appears in retrospect merely accidental, due to the common canonical choice $a = 1/2$ in (4.1), i.e. as if one had taken $a = \gamma_0/b_0$ with $b_0 \rightarrow b_0/3$.⁴

The (exact) expression given by Eq. (6.16) is plotted as a function of the coupling in Fig. 1 in the very common normalization [3] $\lambda \equiv 24g^2$, where it is compared to the standard purely perturbative thermal screening mass, m_D/T , with scale dependence illustrations. In particular, we remark the saturation of the optimized mass for sufficiently large coupling, which agrees qualitatively well with what is obtained at the two-loop order in [21]. This saturation can be seen more explicitly by expanding Eq. (6.16) for strong coupling:

$$\frac{\tilde{m}^{(1)}}{T} \sim \pi \frac{\sqrt{1 + \frac{2}{3}L_T} - 1}{L_T} + \mathcal{O}(\lambda^{-1}). \quad (6.18)$$

The above relation reveals that even if we do not expect our approximation to be valid for arbitrarily large coupling the relation $\tilde{m}/T \lesssim 1$ is always valid while $\tilde{m}/T \ll 1$ in the more perturbative range (see Fig. 1). Therefore, the high- T approximation used to derive those analytic expressions is fully justified *a posteriori*. Concerning the scale dependence, in order to compare with standard results, we use the physical OPT solution, Eq. (6.16), replacing $\lambda \equiv \lambda(\mu)$ by its “exact” one-loop running coupling in Eq. (3.9). We take $\mu_0 \equiv 2\pi T$ as a reference scale, and vary as usual the scale μ in the range $[\pi T, 4\pi T]$. The plot in Fig. 1 is only made for illustration and comparison with the standard perturbative thermal mass, since as explained earlier Eq. (6.16) is *exactly* scale-invariant, i.e. the RGOPT mass gap in Fig. 1 has actually zero thickness, being valid for *any* scale.⁵ Even if one would use the approximate one-loop expanded running coupling [i.e. the last expression in the RHS of Eq. (3.9)], the scale dependence would be extremely moderate, barely visible on the same plot.

Next, coming to the pressure it has a rather simple expression at this one-loop order, in terms of the OPT mass

⁴We anticipate that for QCD, it happens accidentally that $\gamma_0^G(\text{QCD}) = b_0(\text{QCD})/2$, where $\gamma_0^G(\text{QCD})$ is the gluon anomalous mass dimension easily calculable from the relevant counterterm given e.g. in [8,9]. Thus $a = 1/2$ for the gluonic contributions, but the analogous RG-compatible OPT mass (6.17) will not necessarily coincide for the first few perturbative terms with the QCD gluon screening mass [3].

⁵Except for the fact that at some large scale μ , depending on $\lambda(2\pi T)$ value, one hits the naive (one-loop) Landau pole, more precisely at $\mu/(2\pi T) = e^{1/(b_0\lambda(2\pi T))}$: for instance for $g = \sqrt{\lambda(2\pi T)/24} = 1$ the Landau pole is reached at $\mu/(2\pi T) \approx 8.96$.




FIG. 1 (color online). RGOPT mass m/T at one-loop (δ^0) order (thick line), versus standard one-loop perturbative mass as functions of $g(\mu_0 = 2\pi T) = (\lambda(\mu_0)/24)^{1/2}$ with scale-dependence. Grey (light blue) bands: variation of the standard perturbative mass between $\mu = \pi T$ and $4\pi T$ using the exact one-loop running coupling in Eq. (3.9). NB: the RGOPT mass has actually zero thickness since it is exactly scale-invariant.

$\tilde{m}$ and normalized to the ideal gas pressure $P_0 = \pi^2 T^4/90$ (still keeping the general b_0 -dependence):

$$\begin{aligned} \frac{P^{(1)}}{P_0} = 1 - \frac{15}{4\pi^2} \frac{\tilde{m}^2}{T^2} + \frac{15}{2\pi^3} \frac{\tilde{m}^3}{T^3} \\ + \frac{45}{16\pi^4} \left(\frac{1}{b_0\lambda} + L_T \right) \frac{\tilde{m}^4}{T^4} + \mathcal{O}(\tilde{m}^6/T^6) \end{aligned} \quad (6.19)$$

where we recall that this is actually an approximation according to using (2.7), which we argue is however precise at the 10^{-3} level up to $x = m/T \lesssim 1$. One can thus plug the OPT mass expression, Eq. (6.16), into Eq. (6.19) to obtain the full λ -dependence. After some algebra it takes a compact form:

$$\begin{aligned} \frac{P^{(1)}}{P_0}(G) = 1 - \frac{5}{4}G - \frac{15}{2}G^2(1+G) \\ + \frac{5}{3}\sqrt{6} \left[G \left(1 + \frac{3}{2}G \right) \right]^{3/2} + \mathcal{O}(x^6), \end{aligned} \quad (6.20)$$

where we defined $1/G \equiv 1/(b_0\lambda(\mu)) + L_T = 1/(b_0\lambda(\mu_0)) + \gamma_E - \ln 2$, to emphasize that P is exactly scale-invariant and only depends on the single parameter $\lambda(\mu_0)$. Expression (6.20) also explicitly separates the “perturbative” first three terms from the clearly “non-perturbative” last term, and is valid implicitly in the high- T approximation as indicated, but very precise as long as $x \lesssim 1$, corresponding to $G \lesssim 3.3$ meaning very strong coupling for $\lambda(\mu_0) \sim G/b_0$. The first neglected term in (6.20) is actually $-15\zeta[3]/(128\pi^6)x^6 \approx -1.4710^{-4}x^6$ (which is indeed exactly the last term of Eq. (5.8) in [22] in the large- N normalization of b_0). One should be evidently cautious not to use (6.20) beyond its range of validity,

typically for $m \gg T$, in such a case one rather solves the exact mass gap, Eq. (6.12).

One may also easily derive the perturbative expansion of the pressure, which reads to few first orders

$$\begin{aligned} \frac{P^{(1)}}{P_0} \simeq & 1 - \frac{5}{4}\alpha + \frac{5}{3}\sqrt{6}\alpha^{3/2} + \frac{5}{4}(L_T - 6)\alpha^2 \\ & - \frac{5}{2}\sqrt{6}\left(L_T - \frac{3}{2}\right)\alpha^{5/2} - \frac{5}{4}[L_T(L_T - 12) + 6]\alpha^3 \\ & + \frac{5}{32}\sqrt{6}[20L_T(L_T - 3) + 9]\alpha^{7/2} \\ & + \frac{5}{4}L_T[L_T(L_T - 18) + 18]\alpha^4 + \mathcal{O}(\alpha^{9/2}), \end{aligned} \quad (6.21)$$

where $\alpha \equiv b_0\lambda$. We remark again that the one- and two-loop standard perturbative terms for the physical massless pressure [3,7,44] would be reproduced by Eq. (6.21) if effectively replacing, following Ref. [21], $b_0 \rightarrow b_0/3 = 1/(16\pi^2)$. But it is easily seen that the correct coefficients of the leading logarithm terms of the pressure at arbitrary order n : $\alpha^n \ln^{n-1}(\mu)$ and $\alpha^{n+1/2} \ln^{n-1}(\mu)$ (appearing first at three-loop order λ^2), are given by $(b_0\lambda/3)(b_0 \ln(\mu)\lambda)^{n-1}$ with the correct b_0 (comparing with Ref. [44] where the calculation was performed up to order $\alpha^4 \ln \alpha$ using RG techniques). Thus taking $b_0/3$ is not consistent beyond the first two perturbative terms. Equation (6.21) being RG invariant by construction correctly reproduces the leading logarithm structure to all orders beyond two-loop order. We stress also that Eq. (6.21) correctly reproduces all perturbative terms of the large- N result in [22], when taking the correct value of the large- N b_0 . Since our expressions (6.19), (6.20) are valid for arbitrary N we can in principle follow continuously the pressure from large N to $N = 1$, and while doing this there is no reason to abruptly modify the correct $b_0(N)$ to some other “effective” b_0 value. It is useful at this stage to compare this behavior with the SPT pressure up to two-loop or higher order [7], basically build on taking $a = 1/2$ in (4.1): it does reproduce the coefficients of the standard perturbative pressure up to second $\alpha^{3/2}$ order, but not the correct leading logarithm coefficients at order α^2 and beyond, as a consequence of missing RG invariance (it would need to rescale $\ln \mu \rightarrow 3 \ln \mu$ to reproduce those logarithms). So, the scale dependence of the SPT pressure is unmatched at order α^2 and beyond if using the standard running coupling with b_0 .

Thus, keeping the correct b_0 , which is compelling in our RG-based approach, the optimized results Eqs. (6.20) and (6.21) differ from the first two terms of the standard (massless) perturbative pressure for small coupling values by $\lambda(\mu_0) \rightarrow \lambda_{\text{pert}}(\mu_0)/3$, for the canonically normalized coupling of the $N = 1$ scalar model. But this is not a problem, simply a different calibration: the exactly scale-invariant RGOPT pressure (6.20) only depends on the single coupling G or equivalently $\lambda(\mu_0)$, still an arbitrary

parameter at this stage, since the model is not fully specified by any data fixing a physical input scale, μ_0 . So the pressure as a function of this coupling has a limited physical meaning. When going to higher loops, since the nonperturbative RGOPT approximations resum more higher orders, it is not surprising that they differ from standard perturbation when expressed in terms of the original perturbative coupling $\lambda(\mu)$, and one expects to obtain a better approximation for large coupling. The only mandatory feature of any such approximation is certainly the Stefan-Boltzmann limit $P \rightarrow P_0$ for $\lambda \rightarrow 0$, trivially fulfilled by (6.20). Indeed, those features do not contradict truly physical results, as this apparent discrepancy disappears if expressing the pressure in terms of the *physical* mass: to see it, we solve Eq. (4.2) now reciprocally, for $\tilde{\lambda}(m)$, and replacing it in (6.19). It gives simply:

$$P^{(1)}/P_0 = 1 - 15x^2/(8\pi^2) + 15x^3/(8\pi^3) + \mathcal{O}(10^{-4}x^6). \quad (6.22)$$

But here $x = m/T$ is arbitrary as we already used (6.15) to fix $\tilde{\lambda}(m)$. Now, taking for m the physical screening mass [27] m_D in Eq. (2.8), as easily checked it exactly reproduces the first two terms of the standard physical pressure [3]. Thus if we would plot our results in terms of the screening mass, $P(m_D^2/T^2)$, we would have very good agreement with standard results for sufficiently small screening mass m_D^2 , and deviations for larger m^2 , but the study of scale dependence which is our main concern, especially for large coupling values, would be much more difficult. In the sequel we keep the results (6.20) in terms of the $\overline{\text{MS}}$ coupling $\lambda(\mu_0)$, which scale dependence is well-defined at a given perturbative order, since our aim is mainly to compare the scale dependence with other results in the literature also mostly expressed in terms of the running coupling.

The exact expression for $P^{(1)}/P_0$, Eq. (6.19), is plotted in Fig. 2 where the RGOPT result is compared with the standard perturbative expansion at order $\lambda \sim g^2$. The pressure for the rescaled coupling $\lambda \rightarrow \lambda/3$ is also shown on the same figure just for the sake of illustration, which accordingly compares better with the standard perturbative pressure for small coupling values. The improvement of scale (in)dependence of RGOPT is once again drastic at this one-loop order: using the exact one-loop resummed coupling, Eq. (3.9), the RGOPT pressure is exactly scale-invariant, which is obvious in Eq. (6.19) since $\tilde{m}$ is itself exactly scale-invariant and the combination $1/\lambda(\mu) + b_0 L_T$ too, as discussed above. This feature is well illustrated by Fig. 2 by comparing the RGOPT with the standard perturbative pressure at one-loop which has a notoriously large scale dependence.

To conclude this section we stress that all the previous RGOPT one-loop results, reproducing among other things




FIG. 2 (color online). One-loop (δ^0) RGOPT pressure (thick), and rescaling $\lambda \rightarrow \lambda/3$ (dashed), versus standard one-loop (dotted, light blue bands) pressure as function of $g = \sqrt{\lambda(\mu)/24}$ with scale-dependence between $\mu = \pi T$ and $\mu = 4\pi T$. NB: the RGOPT pressure has actually zero thickness since it is exactly scale-invariant.

the exact large- N results, only rely so far on the very simple massive one-loop free energy graph and the knowledge of the first order RG coefficients b_0, γ_0 . But these results are not too surprising since the RG properties, if fully exploited, involve information on all orders “daisy” and “super-daisy” foam graphs like those explicitly resummed in the large- N limit in [22].

VII. 2-LOOP, $\mathcal{O}(\delta)$

We now switch to the two-loop order, thus incorporating all the terms in Eq. (2.4), adding the subtraction terms in Eqs. (3.7) and (3.8), and then performing the δ -expansion consistently to order δ before setting $\delta = 1$. For simplicity we first consider the minimal prescription taking only $s_1 \neq 0$, and will also later consider the relevant changes if the s_2 term, formally of order λ , is also included. The main novelty at two-loop order is that now the RG relation, Eq. (5.2), gives in general a nontrivial additional constraint, that can be used alternatively to the OPT equation, or combined with the latter to completely fix m and λ in terms of the only remaining free parameter, μ/T (apart from the overall dimensional dependence in T). This is another difference with standard OPT or SPT/HTLpt, in which the coupling remains undetermined and the generally adopted prescription is to take its perturbative value as a function of μ/T and a reference coupling $\lambda(\mu_0)$ value. In our case, as explained previously one of the RG solutions is matching this standard perturbative behavior for $\lambda \rightarrow 0$, but for moderate or larger coupling values it will give a non-perturbative dependence. One may thus follow at this stage different possible prescriptions: one can use any of the two OPT and RG equations, to be solved customarily for $\tilde{m}(\lambda)$, next using a two-loop order running coupling, in order to compare with other resummation methods. Alternatively,

one may consider the full RG and OPT combined solutions. Whatever way, neither the optimized mass $\tilde{m}$ solution of Eq. (4.2), nor the optimized coupling $\tilde{\lambda}$ when combining the former with the constraint given by Eq. (5.2), have intrinsic universal physical meaning. Both should better be viewed as intermediate stage values, to be used only within the physical quantities such as the pressure $P(\tilde{m}, \tilde{\lambda}, \mu/T)$. In particular there is no contradiction between the “fixed” optimized coupling $\tilde{\lambda}$ and the standard running coupling, obtained from a different (standard) perturbative RG equation.

Like for the one-loop approximation, we can express all two-loop RGOPT results in terms of the one-loop self-energy defined in Eq. (6.11). After some algebra, the $\mathcal{O}(\delta), \delta \rightarrow 1$ free energy takes a compact form:

$$\mathcal{F}_0 = -\frac{m^4}{(4\pi)^2} \left(\frac{1}{3b_0\lambda} + \frac{s_1}{3} + s_2\lambda \right) + \frac{1}{2} \int_{p,R} \ln(p^2 + m^2) - \frac{m^2}{\lambda} \left(\frac{2\gamma_0}{b_0} \right) \Sigma_R + \frac{1}{2\lambda} \Sigma_R^2, \quad (7.1)$$

where the index “ R ” in the integration means taking the finite part of this already renormalized expression. We also kept as much as possible a general dependence on RG coefficients. The first three terms originate from the subtraction terms s_i in Eqs. (3.7)–(3.8). Notice also the different coefficient $1/(3b_0)$ as a result of expanding to $\mathcal{O}(\delta^1)$, instead of previous $1/(2b_0)$ at one-loop δ^0 order. As already mentioned we first consider for simplicity $s_2 = 0$ in the sequel, while the effects from $s_2 \neq 0$ (that incorporates a RG part of the three-loop contributions) will be discussed later.

Next after straightforward manipulations the OPT Eq. (4.2), and the *reduced* RG operator, Eq. (5.2), acquires a compact neat form:

$$f_{\text{OPT}} \left(m, \lambda, \frac{\mu}{T} \right) = \frac{2}{3} h \left(-s_1 - \frac{1}{b_0\lambda} \right) + \frac{2}{3} S + \Sigma'_R \left(S - \frac{1}{3\lambda} \right) \equiv 0, \quad (7.2)$$

$$f_{\text{RG}} \left(m, \lambda, \frac{\mu}{T} \right) = h \left[\frac{1}{6} + \left(\frac{b_1}{3b_0} - S \right) \lambda \right] + \frac{1}{2} \beta^{(2)}(\lambda) S^2 \equiv 0, \quad (7.3)$$

with $h \equiv (4\pi)^{-2}$, $\beta^{(2)}(\lambda) \equiv b_0\lambda^2 + b_1\lambda^3$ is the standard β -function restricted to two-loops, and recalling also that $s_1 = -1$. We have defined for convenience the reduced (dimensionless) self-energy $S(m, \mu, T) \equiv \Sigma_R/(m^2\lambda)$ thus independent of λ , which makes the coupling dependence very transparent in Eqs. (7.2) and (7.3). For completeness and further use below we also have from Eq. (6.11):

$$\begin{aligned}\Sigma'_R &\equiv \frac{\partial \Sigma_R}{\partial m^2} = \lambda \left(S + m^2 \frac{\partial S}{\partial m^2} \right) \\ &= \gamma_0 \lambda \left(\ln \frac{m^2}{\mu^2} - J_2(m/T) \right).\end{aligned}\quad (7.4)$$

Note that in principle the reduced RG Eq. (7.3) is only valid when combined with Eq. (7.2) since the latter removes the ∂_m part of the complete RG operator in Eq. (5.1). Clearly, in the above normalization, the complete RG Eq. (5.1) reads

$$f_{\text{RG full}} \equiv f_{\text{RG}} + 2\gamma_m(\lambda)f_{\text{OPT}} \equiv 0, \quad (7.5)$$

where the anomalous mass dimension γ_m (truncated at the two-loop order) was defined in Eq. (3.3). Therefore, to obtain the most general solution $\tilde{m}(\lambda)$ consistent with arbitrary coupling values, Eq. (7.5) should be solved. As we shall see below, the solutions $\tilde{m}_{\text{RG}}(\lambda)$ and $\tilde{m}_{\text{RG full}}(\lambda)$ are very close for sufficiently small λ but can depart substantially from each other for arbitrarily large coupling. The advantage of using the reduced RG operator is that the solution can be more easily found when looking for the intersection between the two RG and OPT solutions $m_{\text{OPT}}(\lambda)$ and $m_{\text{RG}}(\lambda)$.

Before proceeding, another digressing remark is that the standard OPT/SPT would correspond to a much simpler OPT equation than Eq. (7.2), since in particular the first two (subtraction) terms s_0, s_1 would be absent, resulting in a simple OPT self-consistent solution: $\tilde{m}^2 \equiv \Sigma_R$. Moreover the modified RG Eqs. (7.3) or (7.5) are usually never considered within the standard OPT/SPT or HTLpt applications. The fact that the OPT and RG relations, Eq. (7.2) and Eq. (7.3), are more involved is expected since the one-loop RG OPT already gives nontrivial results qualitatively similar to two-loop standard OPT/SPT. Accordingly, the relative complexity of RG OPT equations at two-loop is due to the more information they carry on higher RG orders, and is a price for a more efficient and RG-consistent resummation procedure. At this stage the OPT and RG Eqs. (7.2), (7.3) could be solved exactly for $\lambda(m, \mu/T)$, being respectively quadratic and cubic algebraic equations in λ . But to compare with most results in the literature it is more customary to rather solve for a mass gap $m(\lambda, \mu/T)$, to obtain in a next stage the pressure or other thermodynamical quantities as a function of the coupling.

Considering first the (reduced) RG Eq. (7.3), it is best solved in a first stage as a simple quadratic equation for $S(m, \mu, T)$ whose mass gap solutions are

$$S_{\text{RG}}^{\pm}(\lambda) \equiv \frac{\Sigma_R}{\lambda m^2} = \frac{1 \mp \sqrt{1 - (1 + \frac{b_1}{b_0}\lambda)(1 + \frac{2b_1}{b_0}\lambda)}}{3\lambda(1 + \frac{b_1}{b_0}\lambda)}, \quad (7.6)$$

where again the explicit RG dependence has been kept for generality. Note that the coupling dependence is entirely contained in the RHS of Eq. (7.6) since by definition S does

not depend on λ , prior to using the RG Eq. (7.3). Just to see where we stand before considering the more involved exact mass gap solutions at two-loop order, let us consider Eq. (7.6) by crudely neglecting the two-loop β -function coefficient, $b_1 = 0$, or equivalently taking the leading term when expanding (7.6) for $\lambda \rightarrow 0$. It gives immediately $S_{\text{RG}}(\lambda) = 1/(3\lambda)$ which, recalling that $S(m, \mu, T) \equiv \Sigma_R/(\lambda m^2)$ and using (6.11), is nothing but the one-loop mass gap Eq. (6.8) consistently recovered. Once we use a nonzero $b_1 \neq 0$ in Eq. (7.6), at it should at two-loop RG order, one could consider a perturbative expansion of Eq. (7.6) to gradually include higher perturbative order corrections to the one-loop solution (6.8), but it is algebraically not more complicated to solve Eq. (7.6) exactly. Now in fact Eq. (7.6) also reflects a possible complication appearing at two-loop order for large coupling values, due to the two-loop ultraviolet fixed point (UVFP) at $\lambda = -b_0/b_1$ since $b_1 < 0$, see Eq. (3.4). This purely perturbative UVFP is totally spurious, not only since it disappears at three-loop level (where the next coefficient b_2 is positive [28] and large enough so that possible nontrivial fixed points are complex) but more generally since the existence of nonperturbative UVFP is excluded by the numerical evidence for the triviality [45] of the ϕ^4 theory. Nevertheless, since the RG OPT construction basically relies on perturbative RG properties, one may worry that some of our two-loop results could be affected, if driven by this spurious UVFP. Indeed the (+) solution in Eq. (7.6) is singular at the fixed point value of λ , which means that $\tilde{m}^2/\Sigma_R(\tilde{m}^2) \rightarrow 0$ i.e. $\tilde{m} \rightarrow 0$, while the (−) solution is regular: $S_{\text{RG}}(-b_0/b_1) \sim b_1/(6b_0)$, which means that it is *a priori* the solution not wrongly driven by the UVFP. Thus, in the sequel we should be careful to identify any behavior that could be an artifact of this perturbative fixed point. Now, in terms of the rescaled coupling $\lambda = 24g^2$, the UVFP is at $g \sim 1.866$, and the maximum of the β -function (beyond which the coupling is driven to a really wrong behavior), is at $\lambda = -2b_0/(3b_1)$ i.e. $g \simeq 1.524$. Both values are to be considered very large couplings, where the validity of a resummation procedure is anyhow questionable. Therefore, as long as one stays safely below these large coupling values, say not too much above $g \simeq 1$ in practice, our results should remain valid. Moreover, here we are basically focusing on the RG/scale invariance issues in very general terms, rather than on the peculiar nonperturbative dynamics of the ϕ^4 model, which is beyond the present scope. Another property to notice is that the exact RG solutions, Eq. (7.6), become complex at a coupling $\lambda_c = -3b_0/(2b_1) = (4\pi)^2 27/34$ corresponding to $g \simeq 2.28$, thus irrelevant since located beyond the fixed point anyway. Yet one should keep in mind that independently of the presence of nontrivial perturbative fixed points, complex optimized RG solutions are unavoidably expected to occur at some higher perturbative order from exactly solving the OPT and RG equations, as discussed above.




FIG. 3 (color online). $T = 0$ two-loop (δ^1) order OPT (dashed) and RG (thick) (real parts of) solutions $\ln \frac{m}{\mu}(g)$, $g \equiv \sqrt{\lambda/24}$. Also shown is the one-loop solution (dotted) from Eq. (6.7).

A. $T = 0$

To get some more feeling we first explore the $T = 0$ case, which is much simpler since both RG and OPT equations can be solved analytically, e.g. for $\ln m/\mu$ in terms of the (rescaled) coupling $g \equiv \sqrt{\lambda/24}$ (or its reciprocal function $\lambda(\ln m/\mu)$). The (real parts) of the OPT and RG exact solutions, expressed as $\ln[m(g)/\mu]$, are plotted in Fig. 3. We also show for comparison the one-loop solution (dotted) from Eq. (6.7). The physical branch solutions are clearly identified, i.e. those matching standard perturbation for $\mu \ll m$ with fixed m : $\ln m/\mu \rightarrow +\infty$ for $\lambda \rightarrow 0^+$ [see also the discussion after Eq. (6.8)]. One also sees the asymptotic value $\ln m/\mu \approx 1/2$ reached for large coupling g consistently with Eq. (6.7). The two-loop OPT (dashed) physical branch becomes complex for $g \gtrsim 2.094$ (where there is a corresponding bifurcation on the figure), in fact very close above the intersection between the two (real) physical branch solutions, occurring at $\tilde{g} \approx 2.08$, $\ln(\tilde{m}/\mu) \approx 0.083$, thus at $\tilde{m}/\mu$ close to 1 and a quite strong coupling value. The OPT branch is real again at about $g \gtrsim 6$ (where the dashed curve shows a little bump). The RG physical branch becomes also complex at a slightly higher $g \approx 2.28$ value, as already noted above after Eq. (7.6). There are also two other combined RG and OPT solutions (intersections) sitting on the complex branches, see Fig. 3, but these are to be considered unphysical solutions since not connected with the perturbative branches. One can also note that the RG branch has a pole behavior around $g \approx 2$, which is a consequence of the above discussed perturbative UVFP at $g = \sqrt{-b_0/(24b_1)} \approx 1.866$.⁶ The reciprocal function $g(\ln m/\mu)$ would show a frozen behavior at $g \approx 2$. In fact, we stress that the behavior around $g \approx 2$, including the solutions becoming complex, is all driven by

⁶The pole of $\ln m/\mu$ is not exactly at the fixed point of $\beta(\lambda)$ due to the mass-dependence entering the RG Eq. (5.2).

this naive perturbative two-loop UVFP, so that one should simply not trust what happens for g close to those values, say for $g \gtrsim 1$ to be on the safe side. (Notice however that on the figure the one-loop-like behavior is recovered for much larger g values). In particular, the above mentioned real OPT and RG intersection solution at $\tilde{g} \approx 2.08$ is beyond the UVFP, thus very untrustable. From examples in other theories [17], we expect that at higher orders the RGOPT intersection solution may decrease below the UVFP and stabilize to a more reasonably perturbative value.

Switching on the thermal contributions modifies coefficients of the relevant RG and OPT equations, which will result in real intersecting points for the RG and OPT solutions with somewhat lower coupling values $g \sim 1$ for generic μ/T values as we examine in next subsection below.⁷

B. $T \neq 0$

Considering now the thermal contributions, one may solve numerically Eq. (7.6) [or the full RG Eq. (7.5)] and the OPT Eq. (7.2), using the exact expression $S(m, T, \mu)$ from Eq. (6.11), to obtain $x \equiv m/T$ as function of $\lambda(\mu)$ at some chosen scale μ . Concretely, to solve the OPT gap-equation exactly for arbitrary temperature, it is convenient to first solve Eq. (7.2) as a linear equation for $S(m/T, \mu/T)$ in terms of λ and $S'(m/T, \mu/T)$, giving trivially

$$S \equiv \frac{\Sigma_R}{\lambda m^2} = \frac{1}{3\lambda} - \frac{1}{8\pi^2(2 + 3\lambda S')}, \quad (7.7)$$

to be then solved numerically as a mass gap $\tilde{m}(\lambda, \mu/T)$ as function of the coupling and scale, using the expressions of S, S' in Eqs (6.11), (7.4). In the right-hand side of Eq. (7.7), taking only the first term, dominant for $\lambda \rightarrow 0$, one immediately recovers again the one-loop RGOPT mass gap solution Eq. (6.8) (just like for the above discussed RG mass gap (7.6) when taking $b_1 = 0$), while the second term of Eq. (7.7) clearly gives higher order corrections if seen as a perturbative expansion. (However, for large coupling values of order $g = \sqrt{\lambda/24} \sim 1$ such a perturbative expansion of Eq. (7.7) would not give a very accurate mass gap solution so it is better to solve it exactly numerically).

The exact OPT Eq. (7.7) and RG Eq. (7.6) are illustrated with their roots for a rather strong coupling value $g = 1$ in Fig. 4. As one can see, in general both the two-loop order OPT and RG equations have three real solutions, until two solutions become complex (conjugates), which happens for

⁷When complex solutions occur on physical branches, one may recover real solutions by performing a perturbative scheme change, as done in [17]. But this more involved course of action can be avoided in the ϕ^4 case, at least at the two-loop level. We anticipate however that for thermal QCD, RGOPT will unavoidably give complex solutions, mainly due to the opposite signs of the b_0, b_1 coefficients due to asymptotic freedom.




FIG. 4 (color online). Roots of the two-loop (δ^1) exact OPT Eq. (7.2) and RG Eq. (7.3), as compared with one-loop OPT Eq. (6.12) for $g = \sqrt{\lambda(\mu_0)/24} = 1$, $\mu_0 = 2\pi T$. The y-axis values are those of f_{OPT} Eq. (7.2), f_{RG} Eq. (7.3), and Eq. (6.12) in convenient common units.

$g \gtrsim 2.09$ and $g \gtrsim 2.28$ respectively for the OPT and RG equations. Since this is well beyond the fake perturbative UVFP, we cannot trust the detailed consequences near such large coupling values. For more moderate coupling values as illustrated in Fig. 4, there is one RG and one OPT solution with very large $m/T \gg 1$, in fact behaving for small λ as $\tilde{m} \sim e^{1/(b_0\lambda)}$ as one can easily trace even from the one-loop mass gap Eq. (6.12): for $m \gg T$ the $T^2 J_1(m/T)$ term in (6.12) becomes negligible and one simply recovers the $T = 0$ solution for $m(\lambda)$. But for $T \neq 0$ this solution does not have the property of a thermal mass, $m \rightarrow 0$ for $T \rightarrow 0$, so that the other roots are the physically relevant ones. The behavior of the two intermediate and large values roots is qualitatively similar to the one-loop OPT also illustrated and large- N mass gap solution [22], indeed recovered as above explained at one-loop order. What is new as compared to one-loop order and seen on Fig. 4 are the two extra roots with the lowest $x = m/T$ values both for the OPT and RG equations. Concerning the RG root with the lowest $x \approx 0.7$ value, it is easily traced to be the one driven by the UV fixed point: for $\lambda \rightarrow -b_0/b_1$, it gives $x \rightarrow 0$, and we should reject it accordingly. The other OPT root with lowest $x \approx 0.6$ value is more special: it is more easily analyzed by solving the exact OPT Eq. (7.2) for $\lambda(m/T)$, which is a simple quadratic equation. It can also be seen to correspond to the perturbatively odd situation where the second term in Eq. (7.7) dominates over the first term. Then matching with the perturbative behavior, it gives an “ultrasoft” mass $m/T \approx (\pi b_0/2)\lambda + \mathcal{O}(\lambda^2)$ for $\lambda \rightarrow 0$, which contradicts the expected behavior of a thermal mass $m \sim \sqrt{\lambda}T$ on general grounds, that was indeed found at one-loop order. At higher orders this optimized mass has a standard power series in λ . So we consider this solution as a spurious unphysical one, an artifact of the more involved two-loop OPT equation. Therefore at two-loop order we identify unique physical OPT and RG solution, which are

real and positive for all relevant coupling values and compatible with the perturbative behavior of a thermal mass, corresponding to the two intermediate value roots near $x \approx 1$ illustrated on Fig. 4. Once correctly identified, those physical solutions have a qualitative behavior not drastically different from the one-loop order OPT solution of (6.12), apart from the more involved algebra. Note also that solving Eqs. (7.6) or (7.7) for $\tilde{\lambda}(m, \ln(\mu/T))$ one can check analytically that for fixed m/T values, the physical $\lambda(\mu/T)$ solution decreases logarithmically in the two extreme limits $\mu \ll 2\pi T$ and $\mu \gg 2\pi T$, having a maximum in between, a behavior qualitatively quite similar to the large- N or one-loop case.

Alternatively for any solutions, as long as $x \lesssim 1$, one can solve analytically both RG and OPT equations but using the high- T approximation (2.7) and derivatives for the relevant thermal integrals, giving

$$S(x \equiv m/T \lesssim 1) \simeq -\frac{1}{(4\pi)^2} \left(L_T + \frac{2\pi}{x} - \frac{2\pi^2}{x^2} \right). \quad (7.8)$$

In this way one obtains respectively quartic RG and cubic OPT equations in $x = m/T$, which are not particularly telling but gives algebraic solutions. We checked that all the approximate high- T RG solutions for $\tilde{m}$ and the pressure P/P_0 for relevant scale values $\pi T < \mu < 4\pi T$ are excellent, departing below 0.1% from the exact T -dependent solutions, at least up to a value of the (rescaled) coupling $g \equiv \sqrt{\lambda/24} \sim 1.5$, simply because $\tilde{x}$ remains always lower than about ~ 1 . Concerning the OPT solutions, they can give $\tilde{x} > 1$ at large coupling $g \gtrsim 1$, particularly for the higher scale choice $\mu = 4\pi T$, in which case the high- T approximation starts to fail and we better use the exact T -dependent numerical solutions⁸ of Eq. (7.7).

1. Comparison with standard perturbation theory

The OPT physical solution has the following perturbative expansion

$$\begin{aligned} \frac{\tilde{m}_{\text{OPT}}^{(2)}}{T} &\sim \pi \sqrt{\frac{2}{3}} \sqrt{b_0 \lambda} - \pi b_0 \lambda + \frac{3}{128\pi^2 \sqrt{2}} (5 - 2L_T) \lambda^{3/2} \\ &\quad - \frac{9}{1024\pi^3} (3 - 4L_T) \lambda^2 + \dots \end{aligned} \quad (7.9)$$

consistently with the first two terms of Eq. (6.17). Concerning the terms of order $\lambda^{3/2}$ and λ^2 , the (leading) logarithms coefficient of L_T are the same with respect to the one-loop expansion Eq. (6.17), as it should be to all orders. The differences appear only in the constant terms, due both to $s_1 \neq 0$ and other terms of order λ^0 in the original

⁸Such remarks are important to keep in mind in view of possible applications to thermal QCD, for which beyond the one-loop level only high- T expansion results are available [9].

expression of the free energy (7.1). Note that the relative weight of s_1 is changed at order δ^1 , with $s_1/3$ in (7.1). Quite similarly, incorporating $s_2 \neq 0$ gives differences only visible at order λ^3 not given in Eq. (7.9), so it has very little effect at least on the perturbative mass expansion. Plugging this in the pressure expression and expanding perturbatively (while keeping the coupling free for the moment) we obtain the ratio of the two-loop RGOPT pressure to the ideal gas one $P_0 = \pi^2 T^4/90$ as:

$$\begin{aligned} \frac{P_{\text{OPT}}^{(2)}}{P_0} \sim & 1 - \frac{5}{4}\alpha + \frac{5}{3}\sqrt{6}\alpha^{3/2} + \frac{5}{4}(L_T - 20/3)\alpha^2 \\ & - \frac{5}{2}\sqrt{6}(L_T - 13/6)\alpha^{5/2} - \frac{5}{4}[L_T(L_T - 40/3) + 44/3]\alpha^3 \\ & + \mathcal{O}(\lambda^{7/2}), \end{aligned} \quad (7.10)$$

where $\alpha \equiv b_0\lambda$, giving the same first two perturbative terms as the RGOPT one-loop result Eq. (6.21). We remark also, just like in the one-loop case, that if rather solving either the OPT or the RG Eqs. (7.2), (7.5) reciprocally for $\lambda(m)$, and then taking for m the physical screening mass Eq. (2.8), one consistently recovers the standard perturbative expression of the physical pressure up to two-loop order. Remark that the effect of incorporating $s_2 \neq 0$ instead of the simplest choice with only s_1 , appears first in the very last non-logarithmic term of order $\tilde{m}^4\lambda \sim \lambda^3$ in (7.10), replacing $44/3 \rightarrow 637/45 - 4\zeta[3]/5$, which has very little effects upon the final numerical results as long as λ remains reasonably perturbative. This is completely expected since perturbatively $m^4 s_2 \lambda$ in (3.7) is of lowest perturbative order $\sim \lambda^2 s_2 \lambda \sim s_2 \lambda^3$.

Alternatively, we also consider the solution $\tilde{m}(\lambda, \mu/T)$ obtained from the (full) RG Eqs. (7.3) and (7.5). The latter equations can be solved for the exact T -dependence first as quadratic equation for $S(m/T, \mu/T)$ and then numerically for $m(\lambda, \mu/T)$. But as mentioned above the high- T approximation is excellent in this case for all relevant coupling values. The high- T approximated RG equations gives two negative and two positive $\tilde{m}/T$ solutions, the latter having both the correct perturbative behavior, but with one saturating faster for large coupling, which is illustrated as “solution 2” in Fig. 5. In Fig. 5 we also show the scale dependence in the range $\pi T < \mu < 4\pi T$ for all solutions $\tilde{x} \equiv \tilde{m}(g, \mu)/T$. The “lower” values RG solution 2 happens to reach a maximum and then decreases toward zero for large coupling $g \sim 1.86$ (not shown in the figure), an odd behavior which is in fact completely driven by the perturbative two-loop UVFP. Incidentally this solution exhibits an extremely small, almost totally negligible scale dependence up to $g \sim 1$, as can be seen in Fig. 5, in agreement with what is intuitively expected near a RG fixed point. The corresponding pressure, if plugging this solution within (7.1), has even smaller scale dependence. Although we shall simply discard this solution, since the naive




FIG. 5 (color online). Two-loop (δ^1) optimized mass solutions obtained from OPT Eq. (4.2) (dashed, light blue on line) and RG Eq. (5.2) two solutions: thick lines (green on line) and dot-dashed (red on-line) respectively, as function of $g = \sqrt{\lambda(\mu)/24}$ with scale dependence $\pi T < \mu < 4\pi T$.

two-loop UVFP contradicts the genuine nonperturbative dynamics of the ϕ^4 model, it is worth remarking that the corresponding RGOPT solution faithfully reflects the quasi-scale-invariant behavior of the UVFP. The remaining (then unique physical) RG solution 1 exhibits a more pronounced scale dependence as seen in Fig. 5, which will be further explained below. It has a perturbative expansion for small coupling with the first two order terms identical with the other OPT solution above, Eq. (7.9):

$$\begin{aligned} \frac{\tilde{m}_{\text{RG}}^{(2)}}{T} \sim & \pi \sqrt{\frac{2}{3}} \sqrt{b_0\lambda} - \pi b_0\lambda + \frac{(13 - 12L_T)}{256\pi^2 \sqrt{2}} \lambda^{3/2} \\ & + \frac{9}{4096\pi^3} (5 + 16L_T) \lambda^2 + \dots \end{aligned} \quad (7.11)$$

where one can easily check that the coefficients of L_T , thus of $\ln \mu$, are identical, which should be the case to all orders consistently with RG invariance properties.

The perturbative expansion of P/P_0 is identical to (7.10) for the first few orders and for L_T coefficients at all orders if we use instead the $\tilde{m}(\lambda)/T$ solution, Eq. (7.11).

Remark in Fig. 5 the intersections, for different μ scale values, between the RG and OPT solutions, which only exist for the higher of the two RG solution branches (namely the physical solution not influenced by the perturbative UV fixed point). Those intersections can be considered as the full RGOPT solution, which is unique for a given μ input scale. We shall come back later on those full RGOPT solutions, while for the sake of comparison with standard perturbation results we consider at the moment the RG and OPT solutions separately, as given functions of the coupling.




FIG. 6 (color online). RGOPT one-loop (black line) and two-loop (dot-dashed, green on line) versus standard perturbative pressure $P/P_0(g \equiv \sqrt{\lambda/24})$ at one-loop and two-loop. The different bands give the scale dependence between $\mu = \pi T$ and $\mu = 4\pi T$. NB: the RGOPT one-loop curve has actually zero thickness since it is exactly scale-invariant.

In Fig. 6 we plot the exact two-loop RGOPT pressure $P/P_0(g \equiv \sqrt{\lambda/24})$ (dashed lines), as obtained from the mass solution of the full RG Eq. (7.5), compared with one-loop RGOPT and standard perturbative one- and two-loop results, with scale dependence in the range $\pi T < \mu < 4\pi T$. To study the scale dependence we use a standard running coupling exact at two-loop orders. The generic exact expression for the two-loop running $\lambda(\mu)$, generalizing the exact one-loop running in Eq. (3.9), can be expressed (see e.g. Ref. [31]) in terms of the (implicit) Lambert “function” [46] $W(x) \equiv \ln(x/W)$:

$$\lambda(\mu) = \frac{\lambda(\mu_0)}{f_W(\lambda(\mu_0), \ln \frac{\mu}{\mu_0})}, \quad (7.12)$$

with

$$\begin{aligned} f_W(\lambda, L) &= 1 - b_0 L \lambda + \frac{b_1}{b_0} \lambda \ln \left(f_W \frac{1 + \frac{b_1}{b_0} \lambda f_W^{-1}}{1 + \frac{b_1}{b_0} \lambda} \right) \\ &= -\frac{b_1}{b_0} \lambda \left\{ 1 + W \left[- \left(1 + \frac{b_0}{b_1 \lambda} \right) e^{-[1 + \frac{b_0}{b_1 \lambda} (1 - b_0 \lambda L)]} \right] \right\}, \end{aligned} \quad (7.13)$$

where $L \equiv \ln \mu / \mu_0$ and where we consider as usual the reference scale $\mu_0 = 2\pi T$. Actually Eq. (7.12) gives no visible difference (at least up to the relevant moderately large coupling values here studied and the moderate range

of scale variation) with a more handy perturbatively truncated expansion at order λ^3 :

$$\begin{aligned} \lambda^{-1}(\mu) &\approx \lambda^{-1}(\mu_0) - b_0 L - (b_1 L) \lambda - \left(\frac{1}{2} b_0 b_1 L^2 \right) \lambda^2 \\ &\quad - \left(\frac{1}{2} b_1^2 L^2 + \frac{1}{3} b_0^2 b_1 L^3 \right) \lambda^3 + \mathcal{O}(\lambda^4). \end{aligned} \quad (7.14)$$

The RGOPT improvement on scale dependence with respect to standard perturbative results is drastic for the pressure, as one can clearly see in Fig. 6, although scale invariance is not exact at two-loop like it is at one-loop order: there is a moderate residual scale dependence, clearly visible in Fig. 6 for moderately large (rescaled) coupling values $g \gtrsim 0.6$. What is also clearly seen in Fig. 6 is the much better stability of the RGOPT results, since up to $g \simeq .5$, both the scale dependence and the difference between one- and two-loop RGOPT pressure are hardly visible at the figure scale, in contrast with the already poorly convergent standard perturbative pressure for those values. Accordingly, there are important extra cancellations of the scale-dependence happening when $\tilde{m}(\lambda, \mu)$ optimized solutions from Fig. 5 are plugged into the pressure expression $P/P_0(\tilde{m}, \lambda, \mu)$ in Eq. (7.1). But we emphasize again that the optimized masses $\tilde{m}_{\text{OPT}}^{(k)}(\lambda, \mu)$ or $\tilde{m}_{\text{RG}}^{(k)}(\lambda, \mu)$ at order- δ^k are intermediate, *unphysical* quantities, therefore not expected to be themselves scale-invariant in general [although at one-loop order, $\tilde{m}_{\text{OPT}}^0(\lambda, \mu)$ in Eq. (6.16) is exactly (one-loop) scale-invariant as explained above]. Within the RGOPT procedure, RG-invariance is by construction required only for the pressure which represents the actual physical observable, resulting in Eqs. (7.3) and (7.2) optimizing the pressure. Accordingly, the further cancellation of scale-dependence of the two-loop OPT or RG masses $\tilde{m}_{\text{OPT}}(\lambda, \mu)$ or $\tilde{m}_{\text{RG}}(\lambda, \mu)$, once plugged into the pressure $P/P_0(m, \lambda, \mu)$, is expected from the resummation properties of the RGOPT procedure.

The residual scale dependence of the two-loop RGOPT pressure is unavoidable due to the RGOPT construction being not exact but resulting from the optimization of actually two-loop restricted basic free energy by construction, where terms of order λ^2 and higher are truncated. [In contrast the one-loop results above were exactly scale invariant because of the perfect matching of the exact one-loop running coupling with the mass gap (6.16)].

One can make those statements more precise by studying exactly at which perturbative order the scale dependence reappears: examining the perturbative expansion of the pressure above in Eq. (7.10) in which the coupling is replaced by its running expression at truncated two-loop order (so only with b_0, b_1 dependence) Eqs. (7.12) and (7.14), we have checked explicitly that the leading scale dependence reappears first at order λ^3 :

$$\frac{P_{\text{OPT}}^{(2)}(\mu)|_{\text{leading}}}{P_0} \simeq \frac{\lambda^3(\mu_0)}{16384\pi^6} \left(85 \ln \frac{\mu}{\mu_0} - 2190.5 \right) + \mathcal{O}(\lambda^{7/2}), \quad (7.15)$$

i.e. formally at four-loop order, that is one order higher than the naively expected three-loop λ^2 from standard RG invariance properties.⁹ This feature can be anticipated in fact with a little insight without involved explicit high order expansion calculations: recall that the RGOPT construction including the subtraction terms in (3.7), together with the RG invariance preserving interpolation (4.1), guarantees that the two-loop free energy (2.4) is RG invariant up to neglected three-loop terms, of order $\mathcal{O}(m^4\lambda^2)$, but for *arbitrary* mass m . This implies that the mass gap obtained either from the RG Eq. (7.6) or OPT Eq. (7.7) has a remnant scale dependence appearing perturbatively at order $\tilde{m}^2/T^2 \sim \lambda(1 + \dots + \lambda^2 \ln \mu)$, as could easily be checked by explicit expansion too. So it means that the lowest possible order at which a remnant scale dependence appears in the free energy (2.4) is given by the first subtraction terms $-s_0 m^4/\lambda$, giving remnant scale dependence at perturbative order $\lambda^3 \ln \mu$. In contrast the remnant scale-dependence of the standard perturbative two-loop pressure appears at the expected order λ^2 .

Using alternatively the RG solution $\tilde{m}_{\text{RG}}^{(2)}(\lambda)$, we find a residual perturbative scale-dependence reappearing also consistently at order λ^3 with the same coefficient of $\ln \mu$ than in Eq. (7.15), and a very similar constant term, with ~ -2190.5 in Eq. (7.15) replaced by ~ -2009.9 . However, we stress that Eq. (7.15) allows us to understand the perturbative behavior of the remnant scale dependence, but does not properly reflect the actual *nonperturbative* scale-dependence of the full RG or OPT solutions, which we used for the plots in Fig. 6. Indeed the actual values and scale dependence of the pressure for relatively large coupling $g > 0.6$ are very different than what would be obtained by a finite order perturbative truncation at order λ^3 . Accordingly, the moderate residual scale dependence seen on the plots for large g values appears much better than what (7.15) would give, as a result of further nonperturbative cancellations among successive higher orders (of course higher orders of the RGOPT resummation anyway based on a two-loop truncated free energy). Clearly when g becomes of order 1 $\tilde{m}/T$ is also of order 1 so that the previous perturbative reasoning with (7.15) no longer apply and we simply recover that the scale invariance is guaranteed by construction up to remnant terms of order $m^4\lambda^2 \ln \mu$.

As another illustration, in Fig. 7 the exact pressure obtained from the OPT mass solution of Eq. (7.2) is




FIG. 7 (color online). Two-loop pressure $P/P_0(g \equiv \sqrt{\lambda/24})$ obtained from the full RG solution of (7.5) (dot-dashed, green) and obtained from the OPT solution (4.2) (dotted, light red), with scale dependence between $\mu = \pi T$ and $\mu = 4\pi T$. The RGOPT one-loop pressure (black line) is also shown for comparison.

compared with the one obtained from the RG equation. Although both RG and OPT solutions have perturbative scale-dependence reappearing at the same λ^3 order, like in (7.15), the nonperturbative scale dependence of the OPT solution is almost negligible until $g \approx 0.8$, while for larger coupling values it becomes more important than the one of the RG solution (as one can see also for the corresponding OPT mass in Fig. 5). We also remark in Fig. 7 that the pressure obtained from the RG mass remains closer to the one-loop RGOPT pressure for large coupling values, so that the convergence appears better. This, together with the better scale independence for large coupling, is not very surprising since the RG mass solution originates from the RG Eq. (7.5) at the two-loop level. In contrast, the OPT solution results solely from the mass optimization, Eq. (4.2), which incorporates RG properties more indirectly: it also exhibits good scale invariance up to relatively large values $g \gtrsim 0.8$, beyond which it degrades quite rapidly. Also, the fact that the RG and OPT solutions are very close to each other until relatively large coupling values $g \approx 0.6$ shows an overall consistency, by quantifying the relatively small lack of exact RG invariance, since for an exact nonperturbative result the OPT and RG solutions would be identical.

We now consider incorporating the $s_2 \neq 0$ term from Eq. (3.7), which formally belongs to the two-loop order. We already mentioned above that perturbatively it evidently only affects the (re)-expanded pressure (7.10) at order $\tilde{m}^4\lambda \sim \lambda^3$. Accordingly, it does not affect the perturbative order at which the moderate residual scale-dependence first reappears, Eq. (7.15). However, the *nonperturbative* RG and OPT pressures are affected for larger coupling values, as intuitively expected since including $s_2 \neq 0$ incorporates a (RG) part of the three-loop contributions. Indeed, as

⁹When rescaling the coupling as $\lambda \rightarrow 24g^2$, the leading remnant scale-dependence in Eq. (7.15) gives: $\sim (0.075 \ln \mu/\mu_0 - 1.92)g^6$, where the lowest order coefficients are roughly of order $\mathcal{O}(1)$ in this normalization.




FIG. 8 (color online). Same as in Fig. 7, but with $s_2 \neq 0$ in Eq. (3.7).

shown in Fig. 8, it improves slightly the (nonperturbative) scale dependence of the RG pressure (long dashed), in addition the OPT and RG pressures are closer to each other for large coupling values (but the scale dependence is still better for the RG solution than for the OPT one). More remarkably, with $s_2 \neq 0$ the two-loop pressure obtained from the RG mass gap is seen to almost coincide with the one-loop pressure, up to relatively large coupling $g \sim 1$, thus improving the apparent convergence further more. Actually, this rather spectacular coincidence with the one-loop pressure for a large range of coupling values is partly accidental: after applying (4.1), there are some partial cancellations of the two-loop contributions happening for relatively large coupling values (due to opposite signs s_1, s_2), between $s_1/3 = -1/3$ and the dominant two-loop term $s_2\lambda = 24g^2s_2$ [see (7.1), (3.7), (3.8)], with a maximal cancellation for $g \approx 1$, with $24g^2s_2 \approx 1/3$. Since the s_k values depend on the particular RG coefficients, in a different theory s_1 and s_2 may have the same sign, or rather different magnitudes, possibly giving no such partial cancellations. Nevertheless the coincidence with the one-loop pressure is excellent even for relatively large intermediate coupling values e.g. $g \approx 0.6-0.7$, where $24g^2s_2 \approx .12-.16$ does not make the cancellation much effective. Accordingly, there is also clearly a more generic effective stabilization of the perturbative series resulting from the RGOPT construction, with an improved convergence and scale dependence when incorporating higher RG order dependence, as intuitively expected.

2. Comparison with other (SPT and HTLpt) variational resummation approaches

This stability and improved scale-dependence is also illustrated in the next Fig. 9, where the RG pressure is compared with a standard two-loop OPT/SPT [7,14]: more precisely, discarding $\mathcal{E}_0$ in (3.7), taking $a = 1/2$ in (4.1), and using solely Eq. (4.2). To compare with another mass prescription, instead of the mass optimization we use the screening mass, Eq. (2.8) (consistently truncated at two-loop order), plugged in the expression of the free energy, similarly




FIG. 9 (color online). One- and two-loop RGOPT pressure versus one and two-loop standard perturbative pressure and two-loop SPT pressure, with scale-dependence.

to the prescription mostly adopted for QCD HTLpt [9].¹⁰ One sees that using the optimized mass within the SPT/OPT gives a better scale dependence, although SPT with optimized or screening mass both have a definitely stronger scale dependence than the RGOPT for moderately large coupling values. To quantify what is illustrated in Fig. 9 more precisely, let us indicate the relative scale variation of the various methods for the relatively large (rescaled) coupling value $g(2\pi T) = 1$: the corresponding variation of P/P_0 between $\mu = \pi T$ and $4\pi T$ is $\approx 8\%$, 0.8% , 1.5% , 0.35% , and 0.3% respectively for the 2-loop standard perturbation, SPT (optimized mass), SPT (screening mass), RGOPT, and RGOPT with $s_2 \neq 0$. Beyond $g \gtrsim 1$ the two-loop RGOPT scale-dependence increases more rapidly, but is still only $\sim 1.6 \cdot 10^{-2}$ for $g \sim 1.5$ (while for such large coupling the relative variation of the standard perturbative two-loop pressure is as large as ~ 0.25 , and the scale variations of the HTLpt and standard OPT methods become very important too). It would also be interesting to compare quantitatively our scale variation results with the residual scale-dependence appearing in the 2PI approach at three-loop order [23]. However, a precise comparison appears difficult, since the renormalization scheme and scale used in [23] are completely different and not easy to translate into the present scale variation in the $\overline{\text{MS}}$ -scheme.

The RGOPT improvement on scale dependence at the two-loop order may not appear so spectacular as compared

¹⁰Using the perturbative screening mass instead of the optimized mass gap is essentially the procedure in HTLpt applications to QCD because the optimized mass has no real solutions [9].

with SPT, merely a factor $\sim 2-4$ for $g \simeq 1$, essentially because the two-loop order SPT scale dependence here illustrated is still moderate. But at the three-loop level the SPT scale dependence becomes much larger [7] (and similarly HTLpt QCD[9]), of order 1 for moderately large coupling. As previously explained this can be traced both to $a \neq \gamma_0/b_0$ in (4.1) together with the missing RG invariance from $m^4 \ln \mu$ terms for arbitrary m in (2.4): since perturbatively $m^4 \ln \mu \sim \lambda^2 \ln \mu$, the lack of scale invariance at formally one-loop order in the original massive free energy remains somewhat screened at one- and two-loop thermal perturbative expansion order, until it plainly resurfaces at three-loop λ^2 order. Remarkably the SPT pressure has even been calculated more recently to four-loop order [47], and these results show very good convergence with respect to two- and three-loop SPT for the central scale choice $\mu = 2\pi T$. But the remnant scale dependence is not illustrated in [47]. According to our general arguments we do not see how the missing RG-consistent one-loop terms could be compensated by going to higher orders if not present from the beginning. In contrast the RGOPT scale dependence remains very moderate, as illustrated in Fig. 9 when including a RG part of the three-loop contributions from $s_2 \neq 0$. Moreover, without explicitly calculating the full three-loop or higher order RGOPT pressure, we can be confident that the scale-dependence should further improve at higher orders: being built on perturbative RG invariance at order k for arbitrary m , the RGOPT mass gap will exhibit remnant scale dependence as $\tilde{m}^2 \sim \lambda T^2(1 + \dots + \mathcal{O}(\lambda^{k+1} \ln \mu))$, thus the dominant scale dependence in the free energy (namely the lowest perturbative order at which scale dependence will resurface), coming from the leading term $-s_0 m^4/\lambda$, should be of order $\mathcal{O}(\lambda^{k+2})$.

Coming back to the two-loop order pressure, we have also checked the variation of our results against various perturbative truncations: for instance given that all our calculation relies on the basic two-loop free energy (2.4), it may be unnecessary refinement to use the exact running coupling as in (7.12). Thus we looked for variation when truncating the running Eq. (7.14), at order λ^2 , or λ^3 only. Also, we studied the effect of truncating in the RG Eqs. (7.3) and (7.5) terms of order λ^3 , originating from the $b_1 \lambda^3 \partial_\lambda$ term in (3.2), as it is formally one order higher than perturbatively strictly required (and the b_1 -dependence enters anyway at lower orders in the RG Eq. due to the $-s_0/\lambda$ subtraction). We obtain very good stability, since the maximal resulting variations, for the relatively large coupling $g = 1$, and scale choice $\mu = 4\pi T$, is below 10^{-3} for P_{RG}/P_0 , and somewhat worst but reasonably below 10^{-2} for P_{OPT}/P_0 .

3. Full RGOPT two-loop solution

Finally, we can calculate $P_{\text{RGOPT}}^{(2)}/P_0$ for the complete two-loop RGOPT solution, given by the nontrivial intersection between the RG and OPT equations (as illustrated in Fig. 5), as function of the only free scale parameter, that

we choose as $t \equiv \mu/(2\pi T)$ (so the standard central scale choice corresponds to $t = 1$). Ideally for an all order calculation with exact scale invariance one would expect both RG and OPT equations to give identical $\tilde{m}(\lambda, \mu/T)$ solutions. This is indeed the case for simpler models, like typically [15] the large N -limit of the $O(N)$ GN-model. But for a more involved theory at a finite order one expects some differences between the RG and OPT solutions since RG properties are only imposed perturbatively, those remnant differences reflecting the lack of exact RG/scale invariance. Just like the stationary mass solution is expected to approximate the actually massless result, the intersection between the OPT and RG curves at a given order, defining (variational) “fixed-point” mass and coupling, is expected to give a sensible approximation to the exactly scale-invariant nonperturbative result.

For the standard central scale choice $t = 1$, the solution corresponding to the unique intersection of the physical RG and OPT branch solutions, readily seen for $\tilde{m}$ and $\tilde{g}$ in Fig. 5, gives:

$$\tilde{x} \equiv \frac{\tilde{m}}{T} \simeq 0.912; \quad \tilde{g} \equiv \sqrt{\frac{\tilde{\lambda}}{24}} \simeq 0.825; \quad \frac{P_{\text{RGOPT}}^{(2)}}{P_0} \simeq 0.907, \quad (7.16)$$

obtained using the simpler high- T expansion solutions. (NB for $t \gtrsim .5$ approximately, $x(t) \equiv \tilde{m}(\mu)/T$ remains smaller than 1, justifying *a posteriori* the high- T approximation. Typically, for $x = .9$ the high- T approximation is already correct at the level of 0.2% and differences are completely negligible for smaller x .)

The result in (7.16) needs further comments on its physical interpretation. Recall that the truly nonperturbative massless pressure expression $P(\lambda(\mu/T), \mu/T)/P_0$, if that was available, would actually be a function of the single coupling $\lambda(\mu_0 = 2\pi T_0)$ given at some input scale, thanks to exact scale invariance: for any renormalization scale choice μ the nonperturbative running $\lambda(\mu)$ would exactly compensate the explicit μ/T dependence. Incidentally, this is precisely the situation happening at the one-loop RGOPT order, where the exactly scale invariant P/P_0 in (6.20) only depends on the single parameter $b_0 \lambda(\mu_0)$. Now, at higher loop orders in the standard perturbative approaches (or similarly in the SPT/HTLpt approaches after expressing the mass gap $\tilde{m}(\lambda)$ in terms of the coupling), due to imperfect scale invariance giving remnant perturbative terms $\ln^p(\mu/(2\pi T))$, one avoids large logarithms by fixing μ of order $\sim 2\pi T$, which makes the scale effectively T -dependent and allows to study $P/P_0(\lambda(T/T_0))$ by varying the coupling as a function of the scale/temperature. In our case, using the second RG constraint Eq. (5.1) enforces RG invariance at a limited (here two-loop) perturbative order, mimicking exact RG invariance. At $T = 0$ two-loop order, as already mentioned in Sec. VII A there is also a nontrivial intersecting OPT and RG solution at $\tilde{g} \sim 2.08$,

$\ln(\tilde{m}/\mu) \approx 0.08$ (see also Fig. 3), that gives the free energy as $\mathcal{F}_0 \propto \mu^4$, which still requires a reference physical scale μ to be fully determined.¹¹ At $T \neq 0$, combining the OPT and RG equation fixes $\lambda(\mu/T)$ and $m(\mu/T)$ for a given μ/T input, which also fixes P/P_0 as in (7.16). But this combined solution is somewhat academic for the ϕ^4 model, especially in the symmetric phase studied here, where there is no particular physical input temperature to compare with. Moreover, since the scale invariance of the RGOPT pressure is still imperfect at two-loop order, the remnant scale dependence implies different P/P_0 values for different μ/T input choices. For instance, solving similarly the OPT and RG equations for $t = \mu/(2\pi T) = 1/2$ (respectively $t = 2$) gives $P/P_0 = 0.881$ (respectively 0.921). This is consistent with the previously examined remnant scale dependence of the RG and OPT pressures at such relatively large coupling values.

As above mentioned one may expect that the full RGOPT solution for arbitrary scale input could give a sensible approximation of the genuine nonperturbative scale dependence of the coupling. What Eq. (7.16) indicates is that, for the physically reasonable scale choice $\mu \sim 2\pi T$, the optimized coupling $\tilde{g}$ and mass $\tilde{m}/T$ are both of order 1, lying in the rather nonperturbative range where the soft modes $\sim gT$ become comparable with the hard modes $\sim T$. Had we rather found optimized values $\tilde{x}, \tilde{g} \ll 1$, we would not learn much beyond standard perturbation theory. However, we cannot easily follow the combined solution over an arbitrarily large range of scale: the physical branch solution remains real for relatively large variations of $t > 1$, but becomes complex for $t \lesssim 0.27$, which corresponds to the already mentioned complex RG solution for the large coupling $\lambda = -3b_0/(2b_1)$. Thus as already stressed we should not trust our results for g above $g \gtrsim 1$, which corresponds to $t \lesssim .5$. For $t > 1$ (respectively $t < 1$) within a reasonable range, the combined RGOPT solution gives slightly smaller (respectively larger) optimized $\tilde{g}$ values as compared with the one in (7.16), unlike the standard perturbative RG behavior of the ϕ^4 model at small coupling. Indeed for even larger $\mu \gg 2\pi T$, where one expects to recover the four-dimensional $T = 0$ ϕ^4 model properties, the RGOPT real solution gives a slowly (logarithmically) decreasing coupling and mass, which appears roughly consistent naively with the expected triviality property [45]. Conversely, for $\mu \ll 2\pi T$ one expects to recover the three-dimensional high- T limit, such that after eventually reaching a maximum, the optimized coupling is expected to decrease again for $\mu \ll 2\pi T$. But it is difficult to follow the full RGOPT solution becoming complex for $\mu \ll 2\pi T$. Nevertheless, even if they have no real

intersections, the OPT and RG solutions can be reliably determined for $\mu \ll 2\pi T$, as discussed in Sec. VII B, and for fixed $m \ll 2\pi T$, $\lambda_{\text{OPT}}(\mu/T)$ or $\lambda_{\text{RG}}(\mu/T)$ decrease logarithmically for $\mu \ll 2\pi T$.

Note that when incorporating the $s_2 \neq 0$ term from Eq. (3.7), the full RGOPT solutions similar to (7.16) are shifted to a different scale but with the same qualitative behavior, so the net effect of $s_2 \neq 0$ appears essentially as a renormalization scheme redefinition, without drastically changing the results. Incidentally, for $s_2 \neq 0$ the full RGOPT solution is already complex for $t = 1$, while real solutions appear for slightly larger $t \gtrsim 1.5$ values, with corresponding optimal P/P_0 values very close to the one in (7.16). This simply reflects that the occurrence of complex RGOPT solutions in a given theory depend much on the renormalization scheme, so that real solutions could be recovered in principle from appropriate perturbative scheme changes [17]. But this is a much more involved task in the present finite temperature case. Moreover, since the nontrivial RG and OPT intersecting solution happens first at two-loop order, it is probably safer not to take as a very firm prediction the result in (7.16), nor to vary in a wide range around the preferred value $\mu \sim 2\pi T$. From the example of $T = 0$ results in other models [17] the RGOPT variational fixed point solution is more likely to stabilize at the three-loop order, and probably with a more perturbative value of the optimized coupling.

VIII. CONCLUSIONS AND PROSPECTS

Let us summarize our main results. We have shown that the standard treatment of the free energy (pressure) in massive thermal theories, with minimally subtracted counterterms, as primarily done in resummation approaches like OPT/SPT,HTLpt typically, lacks RG invariance. We have then recalled a general simple recipe to restore RG invariance, leading unavoidably to additional finite, temperature-independent vacuum energy contributions, systematically derivable in a perturbative fashion. We have next explained that the OPT/SPT,HTLpt resummation methods based on the modification of the perturbative expansion from the linear δ -expansion in general do not preserve RG invariance, which can however be restored for a different interpolating prescription, Eq. (4.1), uniquely dictated by universal first order RG coefficients, $a = \gamma_0/b_0$. Moreover, the resulting RGOPT resummation can use the RG equation as an alternative or additional combined constraint to determine a self-consistent nonperturbative mass (and coupling), in addition to the sole standard OPT prescription using only the mass optimization. We have then illustrated the RGOPT in details by evaluating the free energy of a thermal scalar field theory at one- and two-loop level. The results show a substantial improvement regarding this type of nonperturbative approach. Namely, we obtain exact RG/scale invariance at the first nontrivial one-loop RGOPT order, which also reproduces all the exact large- N results [22] of the $O(N)$

¹¹Similarly for QCD at zero temperature [16,17] the pion decay constant has been used as a reference physical scale, and the analog of the combined RG and OPT solutions completely fixed $F_\pi/\Lambda_{\text{QCD}}$.

scalar model. At two-loop RGOPT order, the scale dependence and stability are drastically improved up to relatively large coupling values, with respect to most of the other available resummation approaches. Not surprisingly the pressure obtained from the RG mass gap equation happens to have better convergence and scale dependence properties for large coupling than the pressure obtained from the OPT mass gap. We have also illustrated the full RGOPT result obtained from combining the RG and OPT equations, therefore completely fixing the coupling and mass for a given input scale, with results that can be considered as a variational approximation to the truly scale invariant non-perturbative all order result. Beyond two-loop order, since all relevant perturbative results are available at three-loop order [7], the RGOPT procedure can be applied, but we leave this for future work as it becomes algebraically and numerically somewhat more involved. Besides, we are confident that it will further improve the scale dependence with respect to the two-loop results, without explicit three-loop calculations: as explained the RGOPT construction will guarantee the RGOPT free energy to be perturbatively RG invariant up to neglected four-loop $\mathcal{O}(m^4\lambda^3)$ terms, for *arbitrary* mass m . This implies, after using the (OPT or RG) gap equation, that perturbatively a remnant scale dependence appears in the free energy at order $\sim\lambda^4$. The same line of reasoning also explains why the lack of RG invariance of OPT/SPT at formally one-loop order m^4 for arbitrary m , remains partly hidden at one- and two-loop but resurfaces maximally at perturbative three-loop order λ^2 . We also see that the RGOPT anticipates the predictions by one perturbative order, the exact one-loop results being qualitatively similar to standard two-loop resummation results. Therefore, one may argue similarly that the two-loop RGOPT results should be qualitatively comparable to (standard) SPT three-loop results¹² (with a sensibly better scale invariance). Indeed, incorporating the s_2 subtraction term at two-loop order, which includes a (RG) part of the three-loop contributions, further improves the convergence and scale dependence in the nonperturbative coupling range. For these reasons considering the full three-loop contributions is certainly a welcome refinement but not necessary a crucial one in order to demonstrate the efficiency of the method which constitutes our main goal here.

It should be also clear that the whole construction illustrated in particular in Secs. III to V is actually more general, and that it is applicable to QCD. We have already mentioned some properties anticipated to be similar, or sometimes different, in the QCD case. One could expect that

¹²Of course with the limitations that our two-loop results, while resumming some RG dependence of higher order terms, does not reproduce the *full* three-loop contributions, in particular those given by the three-loop thermal “basketball” graph [7,48]. When the subtraction term $s_2\lambda$ from (3.7), (3.8) is included, it is related from RG properties to the single logarithm coefficient of this three-loop basketball graph.

the thermal QCD application may be *a priori* much more difficult than the traditionally simpler scalar model. But given that the difficult gauge dependence and related QCD issues have been solved by the HTL formalism [4,8], the elaborate perturbative HTLpt calculations performed for thermal QCD up to three-loop order in [9] should be readily adaptable to our RGOPT method, which in a first stage relies entirely on perturbative calculations. In HTLpt only vacuum energy, mass, and coupling counterterms are necessary to renormalize the thermodynamic potential. The quark mass anomalous dimension is just the standard one, while the gluon mass anomalous dimension is easily extracted from corresponding known counterterms, given e.g. in [8,9]. Thus our subtraction procedure to recover RG invariance will work just like in the scalar model, applying RG with Eq. (3.1), (3.3), and the modified interpolation Eq. (4.1). Moreover, the HTLpt formalism is inherently a high- T expansion, therefore it will give OPT and RG equations in m/T and g , simpler than the somewhat involved two-loop exact T -dependent mass gap Eqs. (7.2), (7.3) (except that for QCD, HTLpt at two-loop order involves $m \ln m$ terms, due to the two-loop QCD free energy graph structure). Finally, for QCD the known first four coefficients of the β -function, $b_0\dots b_3$, all have the same (negative) sign, such that no fake perturbative fixed points, such as the one present here in the scalar model, will obstruct the identification of correct RGOPT solutions. We are confident that a similarly improved scale-dependence and overall stability will be obtained also from RGOPT adaptation of HTLpt, which could potentially put confrontation with thermal QCD lattice results on even firmer grounds.

ACKNOWLEDGMENTS

M. B. P. is partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-Brazil).

APPENDIX: ALTERNATIVE DERIVATION OF SUBTRACTION TERMS FROM BARE RG-INVARIANCE

For completeness in this appendix we derive the subtraction terms in Eq. (3.8) alternatively by RG invariance considerations solely on the bare expression. The bare free energy at two-loop level consists of Eq. (2.4) plus the remnant divergent terms [7,34] (after mass and coupling renormalization at this order) with $D = 4 - 2\epsilon$:

$$(4\pi)^2 \mathcal{E}_0(\text{residual}) = -m^4 \left[\frac{1}{4\epsilon} + \frac{1}{8\epsilon^2} \left(\frac{\lambda}{16\pi^2} \right) \right]. \quad (\text{A1})$$

As amply explained, minimally subtracting Eq. (A1) would not produce a finite RG-invariant expression. Instead, an explicitly RG-invariant counterterm can be written [17,29,30] in the general (perturbative) form in terms of the RG invariant bare mass and couplings as:

$$(4\pi)^2 \Delta \mathcal{E}_0^{\text{RG}} \equiv -\frac{m_0^4}{\lambda_0} H_0(\epsilon), \quad (\text{A2})$$

with $H_0(\epsilon) \equiv \sum_{i \geq 0} h_i \epsilon^i$ an arbitrary series in ϵ , most conveniently determined perturbatively order by order. Now demanding Eq. (A2) to cancel the remnant divergent terms in (A1), upon using in Eq. (A2) the well-known expressions of the mass and coupling counterterms up to two-loop order, reading in our convention:

$$Z_\lambda \equiv \frac{\lambda_0}{\lambda} \simeq 1 + \frac{b_0}{2\epsilon} \lambda + \left[\left(\frac{b_0}{2\epsilon} \right)^2 + \frac{b_1}{4\epsilon} \right] \lambda^2 + \dots \quad (\text{A3})$$

$$Z_m \equiv \frac{m_0}{m} \simeq 1 + \frac{\gamma_0}{2\epsilon} \lambda + \left[\frac{\gamma_0(\gamma_0 + b_0)}{8\epsilon^2} + \frac{\gamma_1}{4\epsilon} \right] \lambda^2 + \dots \quad (\text{A4})$$

expanded in λ and ϵ series, after some algebra it uniquely determines $h_0 = s_0$ and $h_1 = s_1$, given in Eq. (3.8). But it leaves additional finite subtraction terms identical to Eq. (3.8). Note that it necessarily involves an $m^4 s_0 / \lambda$: using in Eq. (A2) simply m_0 cannot cancel the one-loop divergence in Eq. (A1) since the latter is $\mathcal{O}(\lambda^0)$. Of course for the present ϕ^4 model this construction is equivalent to the one performed in [34].

-
- [1] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz, and K. K. Szabo, *Nature (London)* **443**, 675 (2006); Y. Aoki, Z. Fodor, S. D. Katz, and K. K. Szabo, *Phys. Lett. B* **643**, 46 (2006); M. Cheng *et al.*, *Phys. Rev. D* **74**, 054507 (2006).
- [2] M. Cheng *et al.*, *Phys. Rev. D* **77**, 014511 (2008).
- [3] J. P. Blaizot, E. Iancu, and A. Rebhan, *arXiv:hep-ph/0303185*; U. Kraemmer and A. Rebhan, *Rep. Prog. Phys.* **67**, 351 (2004).
- [4] E. Braaten and R. D. Pisarski, *Nucl. Phys. B* **337**, 569 (1990).
- [5] R. R. Parwani, *Phys. Rev. D* **45**, 4695 (1992); **48**, 5965(E) (1993).
- [6] F. Karsch, A. Patkos, and P. Petreczky, *Phys. Lett. B* **401**, 69 (1997).
- [7] J. O. Andersen, E. Braaten, and M. Strickland, *Phys. Rev. D* **63**, 105008 (2001); J. O. Andersen and M. Strickland, *Phys. Rev. D* **64**, 105012 (2001); *Ann. Phys. (Amsterdam)* **317**, 281 (2005).
- [8] J. O. Andersen, E. Braaten, and M. Strickland, *Phys. Rev. Lett.* **83**, 2139 (1999); *Phys. Rev. D* **61**, 074016 (2000); J. O. Andersen, E. Braaten, E. Petitgirard, and M. Strickland, *Phys. Rev. D* **66**, 085016 (2002).
- [9] J. O. Andersen, M. Strickland, and N. Su, *Phys. Rev. Lett.* **104**, 122003 (2010); *J. High Energy Phys.* **08** (2010) 113; J. O. Andersen, L. E. Leganger, M. Strickland, and N. Su, *J. High Energy Phys.* **08** (2011) 053.
- [10] S. Mogliacci, J. O. Andersen, M. Strickland, N. Su, and A. Vuorinen, *J. High Energy Phys.* **12** (2013) 055; N. Haque, J. O. Andersen, M. G. Mustafa, M. Strickland, and N. Su, *Phys. Rev. D* **89**, 061701 (2014); N. Haque, A. Bandyopadhyay, J. O. Andersen, M. G. Mustafa, M. Strickland, and N. Su, *J. High Energy Phys.* **05** (2014) 027.
- [11] V. I. Yukalov, *Theor. Math. Phys.* **28**, 652 (1976); W. E. Caswell, *Ann. Phys. (N.Y.)* **123**, 153 (1979); I. G. Halliday and P. Suranyi, *Phys. Lett.* **85B**, 421 (1979); R. P. Feynman and H. Kleinert, *Phys. Rev. A* **34**, 5080 (1986); H. F. Jones and M. Moshe, *Phys. Lett. B* **234**, 492 (1990); A. Neveu, *Nucl. Phys. B, Proc. Suppl.* **18**, 242 (1991); V. I. Yukalov, *J. Math. Phys. (N.Y.)* **32**, 1235 (1991); S. Gandhi, H. F. Jones, and M. Pinto, *Nucl. Phys. B* **359**, 429 (1991); C. M. Bender, F. Cooper, K. A. Milton, M. Moshe, S. S. Pinsky, and L. M. Simmons, *Phys. Rev. D* **45**, 1248 (1992); H. Yamada, *Z. Phys. C* **59**, 67 (1993); A. N. Sisakian, I. L. Solovtsov, and O. P. Solovtsova, *Phys. Lett. B* **321**, 381 (1994); H. Kleinert, *Phys. Rev. D* **57**, 2264 (1998); *Phys. Lett. B* **434**, 74 (1998).
- [12] R. Seznec and J. Zinn-Justin, *J. Math. Phys. (N.Y.)* **20**, 1398 (1979); J. C. Le Guillou and J. Zinn-Justin, *Ann. Phys. (N.Y.)* **147**, 57 (1983).
- [13] P. M. Stevenson, *Phys. Rev. D* **23**, 2916 (1981); *Nucl. Phys. B* **203**, 472 (1982).
- [14] R. L. S. Farias, G. Krein, and R. O. Ramos, *Phys. Rev. D* **78**, 065046 (2008).
- [15] J.-L. Kneur and A. Neveu, *Phys. Rev. D* **81**, 125012 (2010).
- [16] J.-L. Kneur and A. Neveu, *Phys. Rev. D* **85**, 014005 (2012).
- [17] J.-L. Kneur and A. Neveu, *Phys. Rev. D* **88**, 074025 (2013).
- [18] J.-L. Kneur and A. Neveu, *Phys. Rev. D* **92**, 074027 (2015).
- [19] J.-L. Kneur and M. B. Pinto, *arXiv:1507.03508*.
- [20] A. Peshier, B. Kampfer, O. P. Pavlenko, and G. Soff, *Europhys. Lett.* **43**, 381 (1998).
- [21] J. P. Blaizot, E. Iancu, and A. Rebhan, *Phys. Rev. D* **63**, 065003 (2001).
- [22] I. T. Drummond, R. R. Horgan, P. V. Landshoff, and A. Rebhan, *Nucl. Phys. B* **524**, 579 (1998).
- [23] J. Berges, S. Borsanyi, U. Reinosa, and J. Serreau, *Phys. Rev. D* **71**, 105004 (2005); *Ann. Phys. (Amsterdam)* **320**, 344 (2005).
- [24] J. P. Blaizot, A. Ipp, and N. Wschebor, *Nucl. Phys. A* **849**, 165 (2011).
- [25] J. P. Blaizot and N. Wschebor, *Phys. Lett. B* **741**, 310 (2015).
- [26] J. I. Kapusta, *Finite-Temperature Field Theory* (Cambridge University Press, Cambridge, 1985); M. Le Bellac, *Thermal Field Theory* (Cambridge University Press, Cambridge, 1996).
- [27] E. Braaten and A. Nieto, *Phys. Rev. D* **51**, 6990 (1995).

- [28] E. Brezin, J. C. Le Guillou, and J. Zinn-Justin, *Phys. Rev. D* **9**, 1121 (1974); S. G. Gorishnii, S. A. Larin, F. V. Tkachov, and K. G. Chetyrkin, *Phys. Lett.* **132B**, 351 (1983); H. Kleinert, J. Neu, V. Schulte-Frohlinde, K. G. Chetyrkin, and S. A. Larin, *Phys. Lett. B* **272**, 39 (1991); **319**, 545 (1993).
- [29] C. Arvanitis, F. Geniet, M. Iacomini, J.-L. Kneur, and A. Neveu, *Int. J. Mod. Phys. A* **12**, 3307 (1997).
- [30] C. Arvanitis, F. Geniet, J. L. Kneur, and A. Neveu, *Phys. Lett. B* **390**, 385 (1997).
- [31] J.-L. Kneur, *Phys. Rev. D* **57**, 2785 (1998).
- [32] C. Ford, D. R. T. Jones, P. W. Stephenson, and M. B. Einhorn, *Nucl. Phys.* **B395**, 17 (1993).
- [33] B. M. Kastening, *Phys. Lett. B* **283**, 287 (1992); M. Bando, T. Kugo, N. Maekawa, and H. Nakano, *Phys. Lett. B* **301**, 83 (1993).
- [34] B. M. Kastening, *Phys. Rev. D* **54**, 3965 (1996); **57**, 3567 (1998).
- [35] S. Chiku and T. Hatsuda, *Phys. Rev. D* **58**, 076001 (1998).
- [36] M. B. Pinto and R. O. Ramos, *Phys. Rev. D* **60**, 105005 (1999); **61**, 125016 (2000).
- [37] R. Guida, K. Konishi, and H. Suzuki, *Ann. Phys. (N.Y.)* **241**, 152 (1995); **249**, 109 (1996).
- [38] J. L. Kneur and D. Reynaud, *Phys. Rev. D* **66**, 085020 (2002).
- [39] J. L. Kneur, M. B. Pinto, R. O. Ramos, and E. Staudt, *Phys. Rev. D* **76**, 045020 (2007); J. L. Kneur, M. B. Pinto, and R. O. Ramos, *Phys. Rev. D* **88**, 045005 (2013).
- [40] T. E. Restrepo, J. C. Macias, M. B. Pinto, and G. N. Ferrari, *Phys. Rev. D* **91**, 065017 (2015).
- [41] H. Kleinert, *Mod. Phys. Lett. B* **17**, 1011 (2003); B. Kastening, *Phys. Rev. A* **68**, 061601 (2003); **69**, 043613 (2004).
- [42] J.-L. Kneur, A. Neveu, and M. B. Pinto, *Phys. Rev. A* **69**, 053624 (2004).
- [43] J. L. Kneur, M. B. Pinto, and R. O. Ramos, *Phys. Rev. C* **81**, 065205 (2010); L. Ferroni, V. Koch, and M. B. Pinto, *Phys. Rev. C* **82**, 055205 (2010).
- [44] J. O. Andersen, L. Kyllingstad, and L. E. Leganger, *J. High Energy Phys.* **08** (2009) 066.
- [45] M. Luscher and P. Weisz, *Nucl. Phys.* **B290**, 25 (1987).
- [46] R. M. Corless, G. H. Gonnet, D. E. G. Hare, D. J. Jeffrey, and D. E. Knuth, *Adv. Comput. Math.* **5**, 329 (1996).
- [47] J. O. Andersen and L. Kyllingstad, *Phys. Rev. D* **78**, 076008 (2008).
- [48] J. O. Andersen, E. Braaten, and M. Strickland, *Phys. Rev. D* **62**, 045004 (2000).