

High-order tail in Schwarzschild spacetimeMarc Casals^{1,2,*} and Adrian Ottewill^{2,†}¹*Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro 22290-180, Brazil*
²*School of Mathematics and Statistics and Complex and Adaptive Systems Laboratory,
University College Dublin, Belfield, Dublin 4, Ireland*

(Received 22 October 2015; published 23 December 2015)

We present an analysis of the behavior at late times of linear field perturbations of a Schwarzschild black hole spacetime. In particular, we give explicit analytic expressions for the field perturbations (for a specific ℓ -multipole) of general spin up to the first four orders at late times. These expressions are valid at arbitrary radius and include, apart from the well-known power-law tail decay at leading order ($\sim t^{-2\ell-3}$), a new logarithmic behavior at third leading order ($\sim t^{-2\ell-5} \ln t$). We obtain these late-time results by developing an analytical formalism initially formulated by Mano, Suzuki and Takasugi (MST) [Prog. Theor. Phys. **95**, 1079 (1996); **96**, 549 (1996)] formalism and by expanding the various MST Fourier-mode quantities for small frequency. While we give explicit expansions up to the first four leading orders (for small frequency for the Fourier modes, for late time for the field perturbation), we give a prescription for obtaining expressions to arbitrary order within a “perturbative regime.”

DOI: 10.1103/PhysRevD.92.124055

PACS numbers: 04.70.Bw, 04.25.Nx, 04.30.-w, 97.60.Lf

I. INTRODUCTION

The study of linear field perturbations of black hole background spacetimes is important for several purposes. For example, it is important for the investigation of the (linear) stability of black holes, the effect that black holes have on fields propagating in their neighborhood or the binary inspiral of a black hole and another compact, astrophysical object. In 1955 and subsequent years [1–7], the equations describing linear field perturbations of a nonrotating (Schwarzschild) black hole spacetime were decoupled and separated. This rendered the equations treatable semianalytically as the full perturbation may be obtained as a sum of Fourier modes, with the radial part satisfying the so-called Regge-Wheeler (RW) ordinary differential equation. It was not until 1972 that a similar feat was achieved by Teukolsky [8,9] in the case of a rotating (Kerr) black hole spacetime. In the Schwarzschild limit, the radial part of the Teukolsky equation reduces to the so-called Bardeen-Press-Teukolsky (BPT) equation [10], which was also obtained in 1972.

The RW and the BPT radial equations are satisfied by different quantities (different combinations of field components and their derivatives) and their solutions have been studied thoroughly, both numerically as well as with asymptotic analyses. Of particular interest for this paper, is the result by Price [3,4] (obtained by studying the field perturbations without Fourier decomposing) for the behavior at late times of a RW field perturbation of any spin of a Schwarzschild black hole. Price found that its radiative ℓ -multipole decays to leading order in the form of a power law: $t^{-2\ell-3}$, where t the Schwarzschild time. The analysis in

[11] of the Teukolsky equation in Kerr shows that the same leading-order power-law tail behavior is satisfied by the radiative multipoles of BPT field perturbations in Schwarzschild. In this paper we present analytic expressions for the behavior of general-spin field perturbations in Schwarzschild up to the first *four* orders for late times, revealing a new logarithmic behavior in the third leading order ($\sim t^{-2\ell-5} \ln t$). Furthermore, although most analyses (though see, e.g., [12] for an exception) that give the radius-dependent coefficient of the leading-order power-law have been constrained to large radius, our results for all four orders are valid for *arbitrary* radius. We obtain this late-time behavior both for the field quantities satisfying the RW as well as the BPT equation. We obtain these results by developing a method which is valid not only at late times, but it may be used to obtain results valid in principle at any time regime. We now briefly introduce this method.

In 1986, Leaver [13] derived various analytical representations for the solutions of the RW and BPT equations in terms of infinite series of special functions. A series of Japanese researchers later “revamped” some of Leaver’s series representations and derived other new series representations for these radial solutions [14–16]. These latter series representations, to which we shall refer as MST expansions after the authors [14–16], are naturally adapted to carrying out small-frequency expansions. Small-frequency expansions yield the late-time behavior of the full linear perturbation after integration over frequency. We note, however, that the MST series in principle converge for any value of the frequency, although the speed of their convergence decreases as the magnitude of the frequency increases.

The MST method is a powerful method which only relatively recently researchers have been starting to use in order to obtain results in black hole perturbation theory. For

*mcasals@cbpf.br, marc.casals@ucd.ie
†adrian.ottewill@ucd.ie

example, in the case of a spherically symmetric background, the MST formalism has found applications in the calculation of the self-force [17] on a point particle [18–20], post-Newtonian coefficients and gauge-invariant quantities [21–25] and dynamical tidal interaction of compact objects [26]. The MST method has proven particularly useful for calculating the retarded Green function (GF) of the wave equation satisfied by the field perturbation, which is a fundamental quantity as it determines the evolution in time of any given initial data. The Fourier modes of the GF possess poles (the so-called quasinormal modes) and a branch cut (BC) in the complex-frequency plane [27]. It is known (e.g., [27,28]) that the small-frequency part of the BC is the contribution to the GF that gives rise to the late-time behavior; the non-small-frequency part of the BC contributes to the behavior at earlier times (see [20,29–32]).

In this paper we derive in detail some of the results on the GF Fourier modes and black hole perturbations that we briefly presented in the paper [33]. Namely, we derive a small-frequency expansion of the MST series in general, and of the BC in particular, which we then use to obtain the late-time behavior of the GF and field perturbations. We obtain the late-time behavior up to the first four leading orders at arbitrary radius in Schwarzschild spacetime and for general-integer spin¹ of the field. In [20] we used the calculation of the late-time GF derived in [33] in order to find its contribution to the self-force on a scalar charge in Schwarzschild. The MST method has also been used to calculate the quasinormal mode contribution to the GF in Schwarzschild spacetime [20] and in Kerr spacetime [34]; in the former case the GF calculation was applied to obtain the scalar self-force and, in the latter case, to obtain the radiation emitted given a specific perturbation source. The calculations of the quasinormal mode series in [20,34] involved evaluating the MST series at frequencies with “arbitrarily” large magnitude.

Apart from an explicit small-frequency/late-time analysis, we shall also generally develop the MST formalism in Schwarzschild spacetime. In particular, we shall present MST expressions (valid for general frequency) for the solutions of the RW equation for general integer spin. To the best of our knowledge, these expressions are new for spin-1 since the MST formalism has not yet been presented for the RW equation for spin-1 (it only has been for spin-0 and spin-2). Furthermore, we develop, also for the first time in the literature, the MST formalism for calculating the contribution to the GF of Fourier modes along the BC. We also give the relationships between the BPT quantities and the RW quantities via the so-called Chandrasekhar transformation. We then give explicit expressions for the main

¹We do not give the expansions explicitly for polar gravitational (Zerilli) perturbations nor for positive BPT spin but these can be derived directly from, respectively, axial (RW) gravitational perturbations or negative BPT spin, both of which we do derive explicitly.

MST quantities up to the first four leading orders for small frequency, for general spin and multipole number ℓ . We note that although small-frequency expansions have been given for some of these quantities already, that has typically been done within a post-Newtonian framework, and so for the radial solutions expanded about radial infinity. In here, instead, we give small-frequency expansions for the radial solutions which are valid for *arbitrary* radius. Our small-frequency expansions converge within a perturbative regime which we specify later in the text (end of Sec. IV).

For the reader who is not interested in the details and who just wants to use our results in order to obtain the late-time behavior of the Green function or a field perturbation to high order, the main result is in Sec. VII A. In particular, in Eq. (7.4) we give the late-time behavior of the ℓ -modes of the GF of the RW equation; above that equation we indicate how to find the coefficients appearing in the equation; below that equation we indicate how to obtain a similar late-time expansion in the BPT case. Of course, if one wants the late-time behavior of some given initial data for the field perturbation, one should convolute the obtained small-frequency expansions for the GF with the initial data as in Eq. (7.6).

The layout of this paper is as follows. In Sec. II we present the RW and BPT equations, expressions for their GFs and the BC contributions, as well as the relationships between the RW and BPT quantities. In Sec. III we develop the MST formalism both for the RW and BPT equations, including our new derivation in the specific case of the RW equation for spin-1. In Secs. IV and V we give explicit expansions of the various MST quantities, except for the radial functions, up to the first four leading orders for small frequency. In Sec. VI we extract the small-frequency expansions for the radial functions using a novel method. In Sec. VII we calculate the late-time behavior of the GF (Sec. VII A) and of a perturbation response (Sec. VII B) and compare it with highly accurate numerical results. In Appendix A we plot the small-frequency expansions of MST quantities and check that they match with an independent method (presented in [29]) which is valid in a “midfrequency” regime. In Appendix B we relate the radial coefficients of the solutions of the RW and BPT equations.

In this paper we use geometrized units: $c = G = 1$. We shall use a bar over a quantity to indicate that the quantity has been made dimensionless via an appropriate factor of “ $2M$ ” (except where otherwise indicated), where M is the black hole mass; e.g., $\bar{\omega} \equiv 2M\omega$ indicates a dimensionless frequency ω and $\bar{r} \equiv r/(2M)$ a dimensionless Schwarzschild radius r .

II. BRANCH CUT GREEN FUNCTION

A. Regge-Wheeler equation

Decoupled and separated equations for linear field perturbations of a Schwarzschild black hole spacetime were derived for axial—also called “odd”—gravitational perturbations (spin $s = 2$) in [1], for electromagnetic perturbations ($s = 1$) in [2,7], and for scalar perturbations ($s = 0$) in [3,4]. All these integer-spin-field perturbation

equations can be written compactly as one single partial differential equation, which in Schwarzschild coordinates reads

$$\left[\nabla^\alpha \nabla_\alpha + s^2 \frac{2M}{r^3} \right] \mathcal{X}_s = S_s, \quad (2.1)$$

where $\mathcal{X}_s = \mathcal{X}_s(t, r, \theta, \phi)$ is a scalar function that describes the field perturbation of spin s created by a source S_s and

$$\nabla^\alpha \nabla_\alpha = -\frac{r^2}{\Delta} \frac{\partial^2}{\partial t^2} + \frac{1}{r^2} \frac{\partial}{\partial r} \left(\Delta \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial}{\partial \theta} \left(\sin^2 \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial}{\partial \phi^2} \quad (2.2)$$

is the Klein-Gordon operator in Schwarzschild spacetime, where $\Delta \equiv r(r - 2M)$ and M is the mass of the black hole. Separating variables we may obtain a complete set of solutions of the form

$${}_s \mathcal{X}_{\ell m \omega}(t, r, \theta, \phi) = e^{-i\omega t} Y_{\ell m}(\theta, \phi) \frac{{}_s X_{\ell m}(r, \omega)}{r}, \quad (2.3)$$

where ℓ and m are, respectively, the multipolar and azimuthal numbers and $Y_{\ell m}(\theta, \phi)$ are the scalar (spin-weight 0) spherical harmonics. As in [35], we treat Eq. (2.1) as a *scalar* wave equation, although in the electromagnetic and gravitational cases we have to be aware that the nonradiative ($\ell < |s|$) modes would drop out when constructing the electromagnetic and gravitational potentials and so these modes would have to be included separately. The radial functions satisfy the ordinary differential equation

$$\left[\frac{1}{r^2} \frac{d}{dr} \left(\Delta \frac{d}{dr} \right) - 2 \frac{\Delta}{r^3} \frac{d}{dr} + \frac{\omega^2 r^2}{\Delta} - \left(\frac{\ell(\ell+1)}{r^2} + \frac{2M(1-s^2)}{r^3} \right) \right] {}_s X_{\ell m} = {}_s S_{\ell m}, \quad (2.4)$$

where ${}_s S_{\ell m} = {}_s S_{\ell m}(r, \omega)$ are the corresponding modes of the source S_s . We shall refer to Eq. (2.4) as the (radial) Regge-Wheeler (RW) equation and to Eq. (2.1) as the four-dimensional RW equation (as per [35]). Introducing the standard ‘‘tortoise’’ coordinate $r_* \equiv r + 2M \ln(r/(2M) - 1)$, Eq. (2.4) may also be written as

$$\left[\frac{d^2}{dr_*^2} + \omega^2 - \frac{\Delta}{r^4} \left(\ell(\ell+1) + \frac{2M(1-s^2)}{r} \right) \right] {}_s X_{\ell m} = \frac{\Delta}{r^2} {}_s S_{\ell m}. \quad (2.5)$$

The case of polar—or ‘‘even’’—gravitational perturbations was derived in [5,6]; the corresponding radial equation is the so-called Zerilli equation. Solutions of the Zerilli equation can be obtained as linear combinations of solutions and their radial derivatives of the RW equation for $s = 2$ [36].

We define the retarded Green function (GF) of the four-dimensional RW equation (2.1) as the solution of [35]

$$\left[\nabla^\alpha \nabla_\alpha + s^2 \frac{2M}{r^3} \right] {}_s G_{\text{ret}}(x, x') = -\delta(x, x') = -\frac{1}{rr'} \delta(t-t') \delta(r-r') \delta(\Omega - \Omega'), \quad (2.6)$$

which obeys appropriate causal boundary conditions, where x and x' are two spacetime points. Here, Ω is the solid angle of the 2-sphere. For notational simplicity, we use time translation invariance to henceforth take $t' = 0$. We may then use the symmetries of the spacetime to write

$${}_s G_{\text{ret}}(x, x') = \frac{1}{rr'} \int_{-\infty+ic}^{\infty+ic} \frac{d\omega}{2\pi} \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} Y_{\ell m}(\theta, \phi) Y_{\ell m}^*(\theta', \phi') \times e^{-i\omega t} {}_s G_{\ell}(r, r'; \omega), \quad (2.7)$$

where $c > 0$, and the Fourier modes ${}_s G_{\ell}(r, r'; \omega)$ must satisfy

$$\left\{ \frac{d^2}{dr_*^2} + \omega^2 - \frac{\Delta}{r^4} \left(\lambda + \frac{2M(1-s^2)}{r} \right) \right\} {}_s G_{\ell}(r, r'; \omega) = -\frac{\Delta}{r^2} \delta(r-r') = -\delta(r_* - r'_*), \quad (2.8)$$

where² $\lambda \equiv \ell(\ell+1)$. Correspondingly,

$${}_s G_{\ell}(r, r'; \omega) = -\frac{f_{\ell}(r_<, \omega) g_{\ell}(r_>, \omega)}{W(\omega)}, \quad (2.9)$$

where $r_> \equiv \max(r, r')$, $r_< \equiv \min(r, r')$, and the Wronskian is given by

$$W(\omega) \equiv W[f_{\ell}(r, \omega), g_{\ell}(r, \omega)] = f_{\ell} \frac{dg_{\ell}}{dr_*} - g_{\ell} \frac{df_{\ell}}{dr_*}. \quad (2.10)$$

Here, the ‘‘ingoing’’ $f_{\ell} = f_{\ell}(r, \omega)$ and ‘‘upgoing’’ $g_{\ell} = g_{\ell}(r, \omega)$ radial functions are solutions of the homogeneous version of the RW equation (2.5) which, for general spin, behave asymptotically as

$$f_{\ell} \sim \begin{cases} e^{-i\omega r_*}, & \bar{r}_* \rightarrow -\infty, \\ A_{\ell}^{\text{in}} e^{-i\omega r_*} + A_{\ell}^{\text{out}} e^{+i\omega r_*}, & \bar{r}_* \rightarrow +\infty, \end{cases} \quad (2.11)$$

where $\bar{r}_* \equiv r_*/(2M)$, A_{ℓ}^{in} and A_{ℓ}^{out} are complex-valued constant coefficients [we give higher-order terms of Eq. (2.11) in Appendix B], and

$$g_{\ell} \sim e^{+i\omega r_*}, \quad \bar{r}_* \rightarrow +\infty. \quad (2.12)$$

It is then easy to see that³

²We note that our definition of λ is slightly different from that in [16].

³Note that due to a typographical error the right-hand side of Eq. (2.3) of Ref. [29] should read $f_{\ell} g'_{\ell} - f'_{\ell} g_{\ell}$ instead of $g_{\ell} f'_{\ell} - f_{\ell} g'_{\ell}$ so that it is indeed $W = 2i\omega A_{\ell}^{\text{in}}$ as claimed below Eq. (2.7) [29].

$$W = 2i\omega A_\ell^{\text{in}}. \quad (2.13)$$

It will be convenient for the next section to define similar ingoing and upgoing solutions of Eq. (2.5) without choosing a specific overall normalization: ${}_s X_\ell^{\text{in}}$

$${}_s X_\ell^{\text{in}} \sim \begin{cases} {}_s X_\ell^{\text{in,tra}} e^{-i\omega r_*}, & \bar{r}_* \rightarrow -\infty, \\ {}_s X_\ell^{\text{in,inc}} e^{-i\omega r_*} + {}_s X_\ell^{\text{in,ref}} e^{+i\omega r_*}, & \bar{r}_* \rightarrow +\infty, \end{cases} \quad (2.14)$$

and

$${}_s X_\ell^{\text{up}} \sim \begin{cases} {}_s X_\ell^{\text{up,inc}} e^{+i\omega r_*} + {}_s X_\ell^{\text{up,ref}} e^{-i\omega r_*}, & \bar{r}_* \rightarrow -\infty, \\ {}_s X_\ell^{\text{up,tra}} e^{+i\omega r_*}, & \bar{r}_* \rightarrow +\infty, \end{cases} \quad (2.15)$$

where ${}_s X_\ell^{\text{inc, in/ref/tra}}$ and ${}_s X_\ell^{\text{up, inc/ref/tra}}$ are ingoing/reflection/transmission coefficients. It is clear that

$${}_s X_\ell^{\text{in}} = {}_s X_\ell^{\text{in,tra}} f_\ell, \quad {}_s X_\ell^{\text{up}} = {}_s X_\ell^{\text{up,tra}} g_\ell. \quad (2.16)$$

Using the standard spherical harmonic addition theorem, we may now rewrite Eq. (2.7) as

$${}_s G_{\text{ret}}(x, x') = \frac{1}{4\pi r r'} \sum_{\ell=0}^{\infty} (2\ell + 1) P_\ell(\cos \gamma) {}_s G_\ell^{\text{ret}}(r, r'; t), \quad (2.17)$$

where

$${}_s G_\ell^{\text{ret}}(r, r'; t) \equiv \frac{1}{2\pi} \int_{-\infty+ic}^{\infty+ic} d\omega {}_s G_\ell(r, r'; \omega) e^{-i\omega t}, \quad (2.18)$$

and γ is the angle between the spacetime points x and x' .

The upgoing radial solution g_ℓ , unlike f_ℓ , possesses a branch cut (BC) in the complex-frequency plane starting at the origin $\omega = 0$ and extending down the negative imaginary axis [13,27,33]. This BC is inherited by A_ℓ^{in} and by the Fourier modes ${}_s G_\ell(r, r'; \omega)$ of the GF. This BC in g_ℓ , however, only occurs as a change of sign in its imaginary part as the frequency crosses the negative imaginary axis, therefore $|g_\ell|$ and $|A_\ell^{\text{in}}|$ do not possess a BC. We will use a new “frequency variable” $\sigma > 0$, so that when $\omega = -i\sigma$ then it lies on the negative imaginary axis and when $\omega = +i\sigma$ then it lies on the positive imaginary axis.⁴ We define $\delta A(\bar{\sigma}) \equiv A_+(-i\bar{\sigma}) - A_-(-i\bar{\sigma})$ for any function $A = A(\bar{\omega})$ possessing a BC along the negative imaginary axis (NIA), where $A_\pm(-i\bar{\sigma}) \equiv \lim_{\epsilon \rightarrow 0^+} A(\pm\epsilon - i\bar{\sigma})$, with $\bar{\sigma} > 0$, where $\bar{\sigma} \equiv 2M\sigma$ and $\bar{\omega} \equiv 2M\omega$.

⁴We note that in [29,30,33] (and in the BC literature references therein) we used a different symbol for the frequency $\sigma = i\omega$. The symbol used there coincided with the symbol for the “renormalized angular momentum” parameter introduced later on that is used throughout the MST literature and which we also use in this paper; hence the reason for the change of symbol to σ .

The contribution from the BC to the ℓ -mode ${}_s G_\ell^{\text{ret}}$ is given by

$${}_s G_\ell^{\text{BC}}(r, r'; t) \equiv \frac{1}{2\pi i} \int_0^\infty d\sigma \delta_s G_\ell(r, r'; \sigma) e^{-\sigma t}. \quad (2.19)$$

Using the obvious symmetry $g_\ell(r, -\omega^*) = g_\ell^*(r, \omega)$ together with the boundary condition $g_\ell(r, +i\sigma) \sim e^{-\sigma r_*}$ as $\bar{r}_* \rightarrow \infty$ and the fact that all three functions $g_{\ell+}(r, -i\sigma)$, $g_{\ell-}(r, -i\sigma)$ and $g_\ell(r, +i\sigma)$ satisfy the same homogenous linear second-order differential equation (namely, the RW equation) with real-valued coefficients along both the negative and the positive imaginary axes (since $\omega = \pm i\sigma$ with $\sigma > 0$ there), it follows that

$$\delta g_\ell = iq(\sigma)g_\ell(r, +i\sigma), \quad (2.20)$$

for some real-valued function $q(\sigma)$, and that $W[g_{\ell-}(r, -i\sigma), g_{\ell+}(r, -i\sigma)] = -2i\sigma q(\sigma)$. We refer to $q(\sigma)$ as the “BC strength.” The symmetry $f_\ell(r, -\omega^*) = f_\ell^*(r, \omega)$ together with the fact that f_ℓ has no BC means that $f_\ell(r, -i\sigma)$, with $\sigma > 0$, is a real-valued function. Putting all these results together, we find that the discontinuity along the BC of the ℓ -modes of the GF is given by

$$\delta_s G_\ell(r, r'; \sigma) \equiv -2i\sigma \frac{q(\sigma)}{|W|^2} f_\ell(r, -i\sigma) f_\ell(r', -i\sigma), \quad (2.21)$$

which is a purely imaginary quantity. The reader may refer to [30,37] for details.

B. Bardeen-Press-Teukolsky equation

The Newman-Penrose formalism offers an alternative way of describing spin-field perturbations of a Schwarzschild black hole background spacetime to that provided by the RW formalism (one key advantage of the Newman-Penrose formalism, however, is that it generalizes to Kerr spacetime). The scaled Newman-Penrose scalars $\Psi_s = \Psi_s(t, r, \theta, \phi)$ obey the following equation [8–10]:

$$\left\{ \nabla^\alpha \nabla_\alpha + \frac{2s}{r^2} \left[\left(-r + \frac{Mr^2}{\Delta} \right) \frac{\partial}{\partial t} + (r - M) \frac{\partial}{\partial r} + i \frac{\cos \theta}{\sin^2 \theta} \frac{\partial}{\partial \phi} + \frac{1 - \text{sct}^2 \theta}{2} \right] \right\} \Psi_s = T_s, \quad (2.22)$$

where T_s is the matter source term. The scalings of the Newman-Penrose scalars here are given by

$$\Psi_s = \begin{cases} \psi_0 \text{ or } r^4 \psi_4, & s = +2 \text{ or } s = -2, \\ \phi_0 \text{ or } r^2 \phi_2, & s = +1 \text{ or } s = -1, \\ \varphi, & s = 0, \end{cases}$$

where $\psi_{0/4}$ are the radiative Weyl scalars, $\phi_{0/2}$ are the radiative Maxwell scalars and φ is a massless scalar field.

We note that in the scalar case ($s = 0$), Eq. (2.22) is the same as the four-dimensional RW equation (2.1).

As for the RW equation [Eq. (2.3)], we may obtain a complete set of solutions of Eq. (2.22) in the form

$$\Psi_s(t, r, \theta, \phi) = e^{-i\omega t} {}_s Y_{\ell m}(\theta, \phi) {}_s R_{\ell m}(r, \omega), \quad (2.23)$$

where ${}_s Y_{\ell m}(\theta, \phi)$ are the spin-weighted spherical harmonics [38,39]. These are given explicitly by ${}_0 Y_{\ell m}(\theta, \phi) = Y_{\ell m}(\theta, \phi)$,

$$\begin{aligned} \pm 1 Y_{\ell m}(\theta, \phi) &= \frac{1}{\sqrt{\ell(\ell+1)}} \left[\mp \partial_\theta - \frac{i}{\sin\theta} \partial_\phi \right] Y_{\ell m}(\theta, \phi), \\ &\ell \geq 1, \\ \pm 2 Y_{\ell m}(\theta, \phi) &= \frac{1}{\sqrt{(\ell-1)\ell(\ell+1)(\ell+2)}} \\ &\times \left[\partial_\theta^2 - \cot\theta \partial_\theta \pm \frac{2i}{\sin\theta} \partial_\theta \partial_\phi \mp \frac{2i \cos\theta}{\sin^2\theta} \partial_\theta \partial_\phi \right. \\ &\left. - \frac{1}{\sin^2\theta} \partial_\phi^2 \right] Y_{\ell m}(\theta, \phi), \quad \ell \geq 2. \end{aligned} \quad (2.24)$$

The spin-weighted spherical harmonics satisfy the following relations:

- (i) the conjugation relation ${}_s Y_{\ell m}^*(\theta, \phi) = (-1)^{m+s} {}_{-s} Y_{\ell(-m)}(\theta, \phi)$;
- (ii) the orthonormality relation

$$\int d^2\Omega {}_s Y_{\ell m}^*(\theta, \phi) {}_s Y_{\ell' m'}(\theta, \phi) = \delta_{\ell\ell'} \delta_{mm'};$$

- (iii) the completeness relation

$$\begin{aligned} \sum_{\ell=|s|}^{\infty} \sum_{m=-\ell}^{\ell} {}_s Y_{\ell m}^*(\theta, \phi) {}_s Y_{\ell m}(\theta', \phi') \\ = \delta(\cos\theta - \cos\theta') \delta(\phi - \phi'); \end{aligned}$$

- (iv) the parity relation ${}_s Y_{\ell m}(\pi - \theta, \pi + \phi) = (-1)^\ell {}_{-s} Y_{\ell(-m)}(\theta, \phi)$;
- (v) and the generalized addition relation

$$\begin{aligned} \sum_m {}_{s_1} Y_{\ell m}(\theta, \varphi) {}_{s_2} Y_{\ell m}^*(\theta', \varphi') \\ = \sqrt{\frac{2\ell+1}{4\pi}} {}_{s_1} Y_{\ell(-s_2)}(\gamma, \alpha) e^{-is_1\beta}, \end{aligned} \quad (2.25)$$

where

$$\begin{aligned} \alpha &= \tan^{-1}(\cos\theta \sin\theta' - \sin\theta \cos\theta' \cos(\phi - \phi')), \\ &\quad - \sin\theta \sin(\phi - \phi')), \\ \beta &= \tan^{-1}(\sin\theta \cos\theta' - \cos\theta \sin\theta' \cos(\phi - \phi')), \\ &\quad - \sin\theta' \sin(\phi - \phi')). \end{aligned}$$

In particular, when $s_1 = s_2 = s$,

$$\begin{aligned} \sum_m {}_s Y_{\ell m}(\theta, \varphi) {}_s Y_{\ell m}^*(\theta', \varphi') \\ = \frac{2\ell+1}{4\pi} (-1)^s \left(\frac{1+\cos\gamma}{2} \right)^{2s} P_{\ell-s}^{(0,2s)}(\cos\gamma) e^{-is(\alpha+\beta)}, \end{aligned} \quad (2.26)$$

where $P_k^{(a,b)}(x)$ are the Jacobi polynomials.

The radial part of the functions in Eq. (2.23) satisfies the ordinary differential equation

$$\begin{aligned} \left[\Delta \frac{d^2}{dr^2} + 2(r-M)(s+1) \frac{d}{dr} + \left(\frac{(r^2\omega - 2is(r-M))r^2\omega}{\Delta} \right. \right. \\ \left. \left. + 4is\omega r + s(s+1) - \lambda \right) \right] {}_s R_{\ell m} = r^2 {}_s T_{\ell m}, \end{aligned} \quad (2.27)$$

where ${}_s T_{\ell m} = {}_s T_{\ell m}(r, \omega)$ are the corresponding modes of the source T_s . We shall refer to Eq. (2.27) as the (radial) Bardeen-Press-Teukolsky (BPT) equation and to Eq. (2.22) as the four-dimensional BPT equation. We may write the former in self-adjoint form as

$$\begin{aligned} \left[\frac{d}{dr} \Delta^{s+1} \frac{d}{dr} + \Delta^s \left(\frac{(r^2\omega - 2is(r-M))r^2\omega}{\Delta} \right. \right. \\ \left. \left. + 4is\omega r + s(s+1) - \lambda \right) \right] {}_s R_{\ell m} = r^2 \Delta^s {}_s T_{\ell m}, \end{aligned} \quad (2.28)$$

or, writing ${}_s R_{\ell m}(r, \omega) = (\Delta^{-s/2}/r) {}_s \mathcal{X}_{\ell m}(r, \omega)$,

$$\begin{aligned} \left[\frac{d^2}{dr_*^2} + \omega^2 + \frac{2is(r-3M)\omega}{r^2} - \lambda \frac{\Delta}{r^4} - \frac{2M\Delta}{r^5} - \frac{M^2 s^2}{r^4} \right] {}_s \mathcal{X}_{\ell m} \\ = \frac{\Delta^{s/2+1}}{r} {}_s T_{\ell m}. \end{aligned} \quad (2.29)$$

Let us denote by ${}_s \mathcal{X}_{\ell}$ a general homogeneous solution of Eq. (2.29). Asymptotically, the homogeneous version of Eq. (2.29) takes the form

$$\begin{aligned} \left[\frac{d^2}{dr_*^2} + (\omega - is\kappa)^2 \right] {}_s \mathcal{X}_{\ell} \sim 0, \quad \bar{r}_* \rightarrow -\infty, \\ \left[\frac{d^2}{dr_*^2} + \omega^2 + \frac{2is\omega}{r} \right] {}_s \mathcal{X}_{\ell} \sim 0, \quad \bar{r}_* \rightarrow +\infty, \end{aligned}$$

where $\kappa \equiv 1/(4M)$ and correspondingly the solutions behave (omitting dimensionful constant factors) as linear combinations of

$$\begin{aligned} {}_s\mathcal{X}_\ell &\sim e^{\pm i(\omega - i\kappa)r_*} \sim e^{\pm s/2} \left(\frac{\Delta}{(2M)^2} \right)^{\pm s/2} e^{\pm i\omega r_*}, \quad \bar{r}_* \rightarrow -\infty, \\ {}_s\mathcal{X}_\ell &\sim r^{\mp s} e^{\pm i\omega r_*}, \quad \bar{r}_* \rightarrow +\infty. \end{aligned}$$

We may define ingoing solutions of the homogeneous version of Eq. (2.29) by their asymptotic behavior as

$${}_s\mathcal{X}_\ell^{\text{in}} \sim \begin{cases} {}_sR_\ell^{\text{in,tra}} (2M)^{1-s} \left(\frac{r}{2M} - 1 \right)^{-s/2} e^{-i\omega r_*}, & \bar{r}_* \rightarrow -\infty, \\ {}_sR_\ell^{\text{in,inc}} r^s e^{-i\omega r_*} + {}_sR_\ell^{\text{in,ref}} r^{-s} e^{+i\omega r_*}, & \bar{r}_* \rightarrow +\infty, \end{cases} \quad (2.30)$$

and the upgoing solutions as

$${}_s\mathcal{X}_\ell^{\text{up}} \sim \begin{cases} {}_sR_\ell^{\text{up,inc}} (2M)^{1+s} \left(\frac{r}{2M} - 1 \right)^{s/2} e^{+i\omega r_*} + {}_sR_\ell^{\text{up,ref}} (2M)^{1-s} \left(\frac{r}{2M} - 1 \right)^{-s/2} e^{-i\omega r_*}, & \bar{r}_* \rightarrow -\infty, \\ {}_sR_\ell^{\text{up,tra}} r^{-s} e^{+i\omega r_*}, & \bar{r}_* \rightarrow +\infty, \end{cases} \quad (2.31)$$

for general BPT spin. The corresponding ingoing and upgoing solutions⁵ of the homogeneous version of the BPT equation (2.27) respectively behave asymptotically as

$${}_s\mathcal{R}_\ell^{\text{in}} \sim \begin{cases} {}_sR_\ell^{\text{in,tra}} \Delta^{-s} e^{-i\omega r_*}, & \bar{r}_* \rightarrow -\infty, \\ {}_sR_\ell^{\text{in,inc}} r^{-1} e^{-i\omega r_*} + {}_sR_\ell^{\text{in,ref}} r^{-1-2s} e^{+i\omega r_*}, & \bar{r}_* \rightarrow +\infty, \end{cases} \quad (2.32)$$

and

$${}_s\mathcal{R}_\ell^{\text{up}} \sim \begin{cases} {}_sR_\ell^{\text{up,inc}} e^{+i\omega r_*} + {}_sR_\ell^{\text{up,ref}} \Delta^{-s} e^{-i\omega r_*}, & \bar{r}_* \rightarrow -\infty, \\ {}_sR_\ell^{\text{up,tra}} r^{-1-2s} e^{+i\omega r_*}, & \bar{r}_* \rightarrow +\infty, \end{cases} \quad (2.33)$$

where ${}_sR_\ell^{\text{in,inc/ref/tra}}$ are the incidence/reflection/transmission coefficients of the ingoing radial BPT solution; similarly for ${}_sR_\ell^{\text{up,inc/ref/tra}}$ for the upgoing solution. It is convenient to define the following ingoing and upgoing solutions and coefficients with a hat on, which are normalized with respect to the corresponding transmission coefficients: ${}_s\hat{R}_\ell^{\text{in}} \equiv {}_sR_\ell^{\text{in}} / {}_sR_\ell^{\text{in,tra}}$, ${}_s\hat{R}_\ell^{\text{up}} \equiv {}_sR_\ell^{\text{up}} / {}_sR_\ell^{\text{up,tra}}$, ${}_s\hat{R}_\ell^{\text{in,inc}} \equiv {}_sR_\ell^{\text{in,inc}} / {}_sR_\ell^{\text{in,tra}}$. Then it is easy to see that

$$\begin{aligned} W^T(\omega) &\equiv W^T[{}_s\hat{R}_\ell^{\text{in}}, {}_s\hat{R}_\ell^{\text{up}}] \\ &\equiv \Delta^{s+1} ({}_s\hat{R}_\ell^{\text{in}} {}_s\hat{R}_\ell^{\text{up}'} - {}_s\hat{R}_\ell^{\text{up}} {}_s\hat{R}_\ell^{\text{in}'}) = 2i\omega {}_s\hat{R}_\ell^{\text{in,inc}}, \end{aligned} \quad (2.34)$$

where primes denote differentiation with respect to r . The quantity W^T is a ‘‘generalized Wronskian’’ in the sense that it is independent of r . Similarly to the RW case, the solution

⁵We keep a spin s subindex in the solutions ${}_sR_\ell^{\text{in}}$ and ${}_sR_\ell^{\text{up}}$ of the BPT equation while we did not for the solutions f_ℓ and g_ℓ of the RW equation because of the explicit spin dependence in the asymptotic equations (2.32) and (2.33) for the former set of solutions [as opposed to Eqs. (2.11) and (2.12) for the latter set]; this leads to the explicit spin dependence in Eq. (2.36).

${}_s\hat{R}_\ell^{\text{up}}$ has a BC along the negative imaginary axis of the complex frequency plane, which is inherited by W^T , whereas ${}_s\hat{R}_\ell^{\text{in}}$ has no BCs [13].

The GF of the four-dimensional BPT equation is the solution of Eq. (2.22) with the source T_s replaced by the distribution $\delta(r - r')\delta(\cos\theta - \cos\theta')\delta(\phi - \phi')/\sqrt{-g}$. It may be expressed as

$$\begin{aligned} {}_sG_{\text{ret},T}(x, x') &= \sum_{\ell, m} \Delta^s(r') {}_sG_\ell^{\text{ret},T}(r, r'; t) {}_sY_{\ell m}(\theta, \varphi) {}_sY_{\ell m}^*(\theta', \varphi'), \end{aligned} \quad (2.35)$$

where

$$\begin{aligned} {}_sG_\ell^{\text{ret},T}(r, r'; t) &\equiv \int_{-\infty + ic}^{\infty + ic} d\omega {}_sG_\ell^T(r, r'; \omega) e^{-i\omega t}, \\ {}_sG_\ell^T(r, r'; \omega) &\equiv -\frac{{}_s\hat{R}_\ell^{\text{in}}(r_{<}, \omega) {}_s\hat{R}_\ell^{\text{up}}(r_{>}, \omega)}{W^T(\omega)}. \end{aligned}$$

In order to obtain an expression for the BC contribution ${}_sG_\ell^{\text{BC},T}$ to the ℓ -mode ${}_sG_\ell^{\text{ret},T}$ of the GF of the four-dimensional BPT equation we proceed similarly to the previous subsection for the RW equation. This contribution can be expressed as in Eq. (2.21) for the RW case, but with ${}_sG_\ell^{\text{BC}}$ and $\delta_s G_\ell$ replaced by ${}_sG_\ell^{\text{BC},T}$ and $\delta_s G_\ell^T$, respectively. We then note that ${}_s\hat{R}_\ell^{\text{up}}(r, -\omega^*) = {}_s\hat{R}_\ell^{\text{up}*}(r, \omega)$ and that $\Delta^{-s} {}_s\hat{R}_\ell^{\text{up}}(r, +i\sigma)$ goes like $e^{-\sigma r^*}/r$ as $\bar{r} \rightarrow \infty$ (neglecting a constant factor). It can be shown that all three functions ${}_s\hat{R}_\ell^{\text{up}}(r, -i\sigma)$, ${}_s\hat{R}_\ell^{\text{up}}(r, -i\sigma)$ and $\Delta^{-s} {}_s\hat{R}_\ell^{\text{up}}(r, +i\sigma)$ satisfy the same homogenous linear second-order differential equation (namely, the BPT equation) with real-valued coefficients along the imaginary- ω axis (since $\omega = \pm i\sigma$ with $\sigma > 0$ there). The two former solutions have the same large- \bar{r} behavior, which is linearly independent from that of the latter. Therefore, it follows that

$$\delta_s \hat{R}_\ell^{\text{up}} = iq^T(\sigma) \Delta^{-s} {}_s\hat{R}_\ell^{\text{up}}(r, +i\sigma), \quad (2.36)$$

for some real-valued function $q^T(\sigma)$, and that $W^T[{}_s\hat{R}_{\ell-}^{\text{up}}, {}_s\hat{R}_{\ell+}^{\text{up}}] = -2i\sigma q^T(\sigma)$. We emphasize that the ‘‘BC strength’’ function q^T is calculated using the upgoing radial function along the positive imaginary axis with the *opposite* spin sign, as opposed to the same spin sign for calculating q in the RW case [see Eq. (2.20)]. The property ${}_s\hat{R}_{\ell}^{\text{in}}(r, -\omega^*) = {}_s\hat{R}_{\ell}^{\text{in}*}(r, \omega)$ together with the fact that ${}_s\hat{R}_{\ell}^{\text{in}}$ has no BC means that ${}_s\hat{R}_{\ell}^{\text{in}}(r, -i\sigma)$, with $\sigma > 0$, is a real-valued function. Putting all these results and properties together, we find that the discontinuity along the BC of the ℓ -modes of the four-dimensional BPT GF is given by

$$\delta_s G_{\ell}^T(r, r'; \sigma) = -2i\sigma \frac{q^T(\sigma)}{|W^T|^2} {}_s\hat{R}_{\ell}^{\text{in}}(r, -i\sigma) {}_s\hat{R}_{\ell}^{\text{in}}(r', -i\sigma). \quad (2.37)$$

In the next subsection we relate the BPT quantities to the RW quantities and, in particular, we present an alternative way [namely via the RW ‘‘BC strength’’ $q(\sigma)$] of calculating the BPT discontinuity $\delta_s G_{\ell}^T$.

C. Relationship between RW and BPT quantities

The so-called Chandrasekhar transformation relates, in the homogeneous case, solutions of the RW equation to solutions of the BPT equation. We note that while the RW equations (2.1) and (2.4) are symmetric under $s \leftrightarrow -s$, the BPT equations (2.22) and (2.27) are not. We here write the Chandrasekhar transformation compactly for spin $s = 0, -1, -2$ (see [40] for spin-2 and, e.g. [41] for spin-1; there are similar transformations in the case of positive spin—which of course only changes the BPT equation, not the RW equation—but we do not deal with these in this paper).

Let us generically denote by ${}_s\mathcal{X}_{\ell}(t, r)$ a homogeneous solution of the 4D RW equation (2.1) after factorizing out the angle dependence via scalar spherical harmonics; similarly, we generically denote by ${}_s\Psi_{\ell}(t, r)$ a homogeneous solution of the 4D BPT equation (2.22) after factorizing out the angle dependence via spin-weighted

spherical harmonics. The corresponding Chandrasekhar transformation is then [35]

$${}_s\Psi_{\ell}(t, r) = \mathcal{O}(t, r) {}_s\mathcal{X}_{\ell}(t, r), \quad \mathcal{O}(t, r) \equiv \left(\frac{\Delta}{r}\right)^{|s|} \left(\partial_r - \frac{r^2}{\Delta} \partial_t\right)^{|s|} r^{|s|-1}, \quad (2.38)$$

up to a normalization constant. We already gave in Eq. (2.24) the angular counterpart of the above transformation, i.e., the transformation from the angular factor in the 4D RW ℓ -modes (namely, the scalar spherical harmonics) to the angular factor in the 4D BPT ℓ -modes (namely, the spin-weighted spherical harmonics). We note that if Eq. (2.24) were naively applied to the modes $\ell < |s|$ it would yield the zero function.

It is useful to write the Chandrasekhar transformation in the frequency domain explicitly for each spin $s = 0, -1$ and -2 separately. We now generically denote by ${}_sX_{\ell}$ and ${}_sR_{\ell}$ homogeneous solutions to the (radial) RW equation (2.4) and (radial) BPT equation (2.27), respectively. Introducing the operators

$$\mathcal{D} \equiv \frac{d}{dr} + \frac{i\omega}{f(r)}, \quad \mathcal{D}^{\dagger} = \frac{d}{dr} - \frac{i\omega}{f(r)},$$

where $f(r) \equiv \Delta/r^2$, we may express the BPT solutions in terms of the RW solutions as

$$\begin{aligned} {}_0R_{\ell} &= \frac{c_0}{r} {}_0X_{\ell}, \\ {}_{-1}R_{\ell} &= c_1(r - 2M)\mathcal{D}_{-1}X_{\ell} = c_1r[f(r) {}_{-1}X_{\ell}' + i\omega {}_{-1}X_{\ell}], \\ {}_{-2}R_{\ell} &= c_2(r - 2M)^2\mathcal{D}^2({}_{-2}X_{\ell}) \\ &= c_2[2(ir^2\omega + r - 3M)r(f(r) {}_{-2}X_{\ell}' + i\omega {}_{-2}X_{\ell}) \\ &\quad + (\ell(\ell + 1)r - 6M)f(r) {}_{-2}X_{\ell}], \end{aligned} \quad (2.39)$$

where c_0, c_1 and c_2 are constants of proportionality and primes denote differentiation with respect to r . Conversely, we have

$$\begin{aligned} {}_0X_{\ell} &= \frac{1}{c_0} r {}_0R_{\ell}, \\ {}_{-1}X_{\ell} &= \frac{1}{\lambda c_1} r^2 \mathcal{D}^{\dagger} \left(\frac{{}_{-1}R_{\ell}}{r} \right) = \frac{1}{\lambda c_1} \left[r {}_{-1}R_{\ell}' - \left(1 + \frac{i\omega r}{f(r)} \right) {}_{-1}R_{\ell} \right], \\ {}_{-2}X_{\ell} &= \frac{1}{((\ell - 1)\ell(\ell + 1)(\ell + 2) - 12iM\omega)c_2} r^3 \mathcal{D}^{\dagger 2} \left(\frac{{}_{-2}R_{\ell}}{r^2} \right) \\ &= \frac{1}{((\ell - 1)\ell(\ell + 1)(\ell + 2) - 12iM\omega)c_2} \left[\frac{2(3M - r - i\omega r^2)}{r - 2M} {}_{-2}R_{\ell}' \right. \\ &\quad \left. + \frac{r^2(\ell(\ell + 1) - 2r^2\omega^2 + 8i\omega r + 4) - 2Mr(\ell(\ell + 1) + 9i\omega r + 10) + 24M^2}{r(r - 2M)^2} {}_{-2}R_{\ell} \right]. \end{aligned} \quad (2.40)$$

We will now use the Chandrasekhar transformation in order to relate the BPT and RW Wronskians as well as the BPT and RW BC strengths. The specific normalizations (2.11) and (2.12) of the RW radial functions f_ℓ and g_ℓ yield specific normalizations for the corresponding BPT radial functions via the transformation equation (2.38). We will denote the BPT radial functions and coefficients with these specific normalizations with a tilde superscript, i.e., ${}_s\tilde{R}_\ell^{\text{in}} \equiv \mathcal{O}_\omega(r)f_\ell$, ${}_s\tilde{R}_\ell^{\text{up}} \equiv \mathcal{O}_\omega(r)g_\ell$ and ${}_s\tilde{R}_\ell^{\text{in,tra}}$, ${}_s\tilde{R}_\ell^{\text{up,tra}}$, ... are, respectively, the coefficients ${}_s\tilde{R}_\ell^{\text{in,tra}}$, ${}_s\tilde{R}_\ell^{\text{up,tra}}$, ... of ${}_s\tilde{R}_\ell^{\text{in}}$ and ${}_s\tilde{R}_\ell^{\text{up}}$. In Appendix B we use the Chandrasekhar transformation in order to find these BPT radial coefficients and Wronskian in terms of the RW radial coefficients and Wronskian. We are here using the obvious notation of $\mathcal{O}_\omega(r)$ for $\mathcal{O}(t, r)$ after replacing “ ∂_t ” by “ $-i\omega$ ” in it.

Let us now try to find an alternative expression to Eq. (2.37) for the BPT $\delta_s G_\ell^T$ in terms of the RW $q(\sigma)$. We first reexpress the BPT modes ${}_s G_\ell^T$ of Eq. (2.35) in terms of the RW solutions f_ℓ and g_ℓ by using the Chandrasekhar transformation equation (2.38):

$${}_s G_\ell^T(r, r'; \sigma) = -\mathcal{O}_\omega(r_<) \mathcal{O}_\omega(r_>) \frac{f_\ell(r_<, \omega) g_\ell(r_>, \omega)}{{}_s\tilde{R}_\ell^{\text{in,tra}} {}_s\tilde{R}_\ell^{\text{up,tra}} W^T}. \quad (2.41)$$

Using this expression we can find the discontinuity of the BPT ℓ -modes as

$$\begin{aligned} \delta_s G_\ell^T(r, r'; \sigma) &= \frac{W}{{}_s\tilde{R}_\ell^{\text{in,tra}} {}_s\tilde{R}_\ell^{\text{up,tra}} W^T} \mathcal{O}_\omega(r) \mathcal{O}_\omega(r') \delta_s G_\ell(r, r'; \sigma) \\ &= -\frac{2i\sigma q(\sigma) {}_s\tilde{R}_\ell^{\text{in,tra}}}{W^T \cdot W^* {}_s\tilde{R}_\ell^{\text{up,tra}}} \hat{R}_\ell^{\text{in}}(r, \omega) \hat{R}_\ell^{\text{in}}(r', \omega). \end{aligned} \quad (2.42)$$

In deriving Eq. (2.42) we have used the fact that $W/({}_s\tilde{R}_\ell^{\text{in,tra}} \cdot {}_s\tilde{R}_\ell^{\text{up,tra}} W^T)$ does not have a BC (as can be seen from the expressions in Appendix B) and we have made use of Eq. (2.21).

$$C = \frac{256M^6}{(\lambda-2)\lambda} \left(1 - \frac{6(\lambda^2 - 2\lambda - 1)\bar{\sigma}}{(\lambda-2)\lambda} + \frac{(13\lambda^4 - 52\lambda^3 + 16\lambda^2 + 72\lambda + 36)\bar{\sigma}^2}{(\lambda-2)^2\lambda^2} \right) + \mathcal{O}(\bar{\sigma}^3), \quad s = -2. \quad (2.48)$$

From Eqs. (2.37) and (2.46) and the fact that the radial functions f_ℓ and ${}_s\hat{R}_\ell^{\text{in}}$ are generically of the same leading order (order zero) in $\bar{\sigma}$ as $\bar{\sigma} \rightarrow 0$ (see Secs. IIIVI), it follows that the RW and the BPT BC modes $\delta_s G_\ell$ and $\delta_s G_\ell^T$ are of the same leading order in $\bar{\sigma}$ as $\bar{\sigma} \rightarrow 0$. As a

Comparing Eqs. (2.37) and (2.42) we immediately obtain a relationship between the RW and BPT BC strengths:

$$q^T(\sigma) = \left(\frac{W^T}{W} \right)^* \cdot \frac{{}_s\tilde{R}_\ell^{\text{in,tra}}}{{}_s\tilde{R}_\ell^{\text{up,tra}}} q(\sigma). \quad (2.43)$$

Using the results in Appendix B, we have that

$$q^T(\sigma) = q(\sigma) \begin{cases} 1 & s = 0, \\ -\frac{\alpha_\infty^* \alpha_+}{2\sigma \alpha_+^*}, & s = -1, \\ \frac{\beta_\infty^* (\alpha_+ + 2M^2 \beta_+)}{2\sigma^2 (\alpha_+^* + 2M^2 \beta_+^*)}, & s = -2, \end{cases} \quad (2.44)$$

and

$$\frac{q^T}{|W^T|^2} = \frac{{}_s\tilde{R}_\ell^{\text{in,tra}} W}{{}_s\tilde{R}_\ell^{\text{up,tra}} W^T} \frac{q}{|W|^2} = \frac{q}{|W|^2} \begin{cases} \frac{1}{4M^2} & s = 0, \\ -\frac{\alpha_+^2}{2\sigma \alpha_\infty}, & s = -1, \\ \frac{(\alpha_+ + 2M^2 \beta_+)^2}{2M^2 \sigma^2 \beta_\infty}, & s = -2. \end{cases} \quad (2.45)$$

Inverting the relationship and using the results in Appendix B, we explicitly obtain

$$\frac{q}{|W|^2} = C \frac{q^T}{|W^T|^2} \quad (2.46)$$

with

$$\begin{aligned} C &= 4M^2, \quad s = 0, \\ C &= -\frac{16\lambda M^4}{(\lambda - s^2 + 1)^2} (1 - 4\bar{\sigma} + 4\bar{\sigma}^2), \quad s = -1, \\ C &= \frac{256M^6 (1 - 2\bar{\sigma})^2 (1 - \bar{\sigma})^2}{\lambda^2 - 2\lambda - 6\bar{\sigma}}, \quad s = -2. \end{aligned} \quad (2.47)$$

The small-frequency behavior of C for $s = 0$ and -1 is already manifest in the exact result above; for $s = -2$ we expand it as

consequence, the RW and BPT GFs are of the same leading order in \bar{t} as $\bar{t} \rightarrow \infty$, as we explicitly see in Sec. VII. That is, the RW and the BPT quantities describing black hole perturbations decay at the same rate at late times.

III. MST FORMALISM FOR THE RW AND BPT EQUATIONS FOR GENERAL SPIN

The MST method for the Teukolsky equation in Kerr (and, therefore, for the BPT equation in Schwarzschild) was given in ST for general spin, and earlier in [14] (henceforth MSTa) just for spin-2. To the best of our knowledge, however, the MST method for the RW equation has only been given for spin-2, which was done in [15] (henceforth MSTb). Therefore, the MST method for the RW equation still has not been developed for spin-1 (obviously, the RW spin-0 case is essentially just the same as in ST with $s = 0$). In this section, we develop the MST method for the RW equation for general spin: for spin-2 we recover MSTb, for spin-0 we essentially recover ST with $s = 0$ and, for spin-1, to the best of our knowledge, the results are new. We also develop the MST method for the BPT equation which, although already existing in the literature, will allow us to emphasize the connections between the MST formalism for the RW and BPT equations for general spin $s = 0, -1$ and -2 . In particular, we write the expansions for the radial solutions of both the RW and BPT equations in terms of just one set of “universal” coefficients (namely, a_n). When referring to the literature, we shall use the generic term MST to refer to all MSTa, MSTb and ST.

We shall use the notation of $X_s^{\text{in/up}}$ and $R_s^{\text{in/up}}$ (i.e., with a slight change in the subindices with respect to the homogeneous RW solutions ${}_s X_\ell^{\text{in/up}}$ and the BPT solutions ${}_s R_\ell^{\text{in/up}}$, respectively) for the ingoing/upgoing solutions of the RW and BPT equations when using the specific normalization as in MST [i.e., the one in Eqs. (3.12) and (3.20) below]. A similar change in the subindices notation applies to their incidence, reflection and transmission coefficients.

A. Series of hypergeometric functions

In this section we shall assume that $s \leq 0$. For the RW functions we write

$$\begin{aligned} X_s^{\text{in}} &= e^{i\bar{\omega}\bar{r}^{s+1}} e^{-i\omega r_s} \check{p}_s^{\text{in}}(x) \\ &= \bar{r}^{s+1} e^{-i\bar{\omega}(\bar{r}-1)} (\bar{r}-1)^{-i\bar{\omega}} \check{p}_s^{\text{in}}(x) \\ &= (1-x)^{s+1} e^{i\bar{\omega}x} (-x)^{-i\bar{\omega}} \check{p}_s^{\text{in}}(x), \end{aligned} \quad (3.1)$$

where $x \equiv 1 - \bar{r}$. This leads to the ordinary differential equation

$$\check{p}_s^{\text{in}}(x) = N_s^{\text{in}} \sum_{n=-\infty}^{\infty} \check{a}_n \frac{\Gamma(-n-\nu+s-i\bar{\omega})\Gamma(n+\nu+s+1-i\bar{\omega})}{\Gamma(1-2i\bar{\omega})} {}_2F_1(-n-\nu+s-i\bar{\omega}, n+\nu+s+1-i\bar{\omega}; 1-2i\bar{\omega}; x), \quad (3.4a)$$

$$p_s^{\text{in}}(x) = N_s^{\text{in}} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(-n-\nu-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(1-s-2i\bar{\omega})} {}_2F_1(-n-\nu-i\bar{\omega}, n+\nu+1-i\bar{\omega}; 1-s-2i\bar{\omega}; x), \quad (3.4b)$$

$$\begin{aligned} x(1-x)\check{p}_s^{\text{in}''}(x) + (1-2i\bar{\omega}(1-x)^2 - 2(1+s)x)\check{p}_s^{\text{in}'}(x) \\ + ((\ell-s)(\ell+s+1) + 2i\bar{\omega}(1+s)(1-x))\check{p}_s^{\text{in}}(x) = 0. \end{aligned} \quad (3.2)$$

Correspondingly, for the BPT functions we write (as in ST)

$$\begin{aligned} R_s^{\text{in}} &= e^{i\bar{\omega}(\bar{r}-1)^{-s}} e^{-i\omega r_s} p_s^{\text{in}}(x) \\ &= e^{-i\bar{\omega}(\bar{r}-1)} (\bar{r}-1)^{-s-i\bar{\omega}} p_s^{\text{in}}(x) \\ &= e^{i\bar{\omega}x} (-x)^{-s-i\bar{\omega}} p_s^{\text{in}}(x), \end{aligned}$$

which leads to the equation

$$\begin{aligned} x(1-x)p_s^{\text{in}''}(x) + (4iM\omega x(1-x) + 2(2iM\omega - 1)x \\ + (1-s-4iM\omega))p_s^{\text{in}'}(x) + (\ell(\ell+1) \\ + 4iM\omega(1-s)(1-x))p_s^{\text{in}}(x) = 0. \end{aligned} \quad (3.3)$$

In terms of $\check{p}_s^{\text{in}}(x)$ and $p_s^{\text{in}}(x)$ the Chandrasekhar transformations take the form

$$\begin{aligned} p_0^{\text{in}}(x) &= c_0 \check{p}_0^{\text{in}}(x), \\ p_{-1}^{\text{in}}(x) &= c_1 \check{p}_{-1}^{\text{in}'}(x), \\ p_{-2}^{\text{in}}(x) &= c_2 \check{p}_{-2}^{\text{in}''}(x), \end{aligned}$$

and conversely

$$\begin{aligned} \check{p}_0^{\text{in}}(x) &= \frac{1}{c_0} p_0^{\text{in}}(x), \\ \check{p}_{-1}^{\text{in}}(x) &= \frac{1}{\ell(\ell+1)c_1} [x(1-x)p_{-1}^{\text{in}'}(x) \\ &\quad + (1-2i\bar{\omega}(1-x)^2)p_{-1}^{\text{in}}(x)], \\ \check{p}_{-2}^{\text{in}}(x) &= \frac{1}{((\ell-1)\ell(\ell+1)(\ell+2) - 6i\bar{\omega})c_2} \\ &\quad \times [x(1-x)(2x+1-2i\bar{\omega}(1-x)^2)p_{-2}^{\text{in}'}(x) \\ &\quad - (\ell(\ell+1)x(1-x) + 4\bar{\omega}^2(1-x)^4 \\ &\quad + 6i\bar{\omega}(x+1)(1-x)^2 - 2(2x+1))p_{-2}^{\text{in}}(x)]. \end{aligned}$$

We now follow MST and introduce the expansion in terms of hypergeometric functions,

where N_s^{in} is a normalization constant which we specify later on. Here, the parameter ν is referred to as the renormalized angular momentum and is determined by the requirement that these series converge both as $n \rightarrow \infty$ and $n \rightarrow -\infty$, it has the property that either $\nu = \ell + O(\bar{\omega})$ or $\nu = -\ell - 1 + O(\bar{\omega})$; see MSTa or ST for a full discussion. These series representations in terms of hypergeometric functions converge $\forall r \in [2M, \infty)$.

In these terms, the Chandrasekhar transformations follow from the standard identity [Eq. (15.5.1) [42]]

$$\frac{d}{dx} {}_2F_1(a, b; c; x) = \frac{ab}{c} {}_2F_1(a + 1, b + 1; c + 1; x),$$

for some parameters a , b and c . Inserting into their respective differential equations and using the standard hypergeometric function identities

$$\begin{aligned} x {}_2F_1(a, b; c; x) &= \frac{a(b-c)}{(a-b)(a-b+1)} {}_2F_1(a+1, b-1; c; x) + \frac{c(a+b-1)-2ab}{(a-b-1)(a-b+1)} {}_2F_1(a, b; c; x) \\ &\quad + \frac{b(a-c)}{(a-b-1)(a-b)} {}_2F_1(a-1, b+1; c; x), \\ x(1-x) \frac{d}{dx} {}_2F_1(a, b; c; x) &= \frac{ab(c-b)}{(a-b)(a-b+1)} {}_2F_1(a+1, b-1; c; x) + \frac{ab(2c-a-b-1)}{(a-b+1)(a-b-1)} {}_2F_1(a, b; c; x) \\ &\quad + \frac{ab(a-c)}{(a-b)(a-b-1)} {}_2F_1(a-1, b+1; c; x), \end{aligned}$$

we find that \check{a}_n and a_n must satisfy the same three-term recurrence relation:

$$\alpha_n a_{n+1} + \beta_n a_n + \gamma_n a_{n-1} = 0, \tag{3.5}$$

where

$$\begin{aligned} \alpha_n &= -\frac{i\bar{\omega}(n+\nu+1-i\bar{\omega})(n+\nu+1+s-i\bar{\omega})(n+\nu+1+s+i\bar{\omega})(n+\nu)}{(2n+2\nu+3)}, \\ \beta_n &= -\lambda(n+\nu)(n+\nu+1) + ((n+\nu)(n+\nu+1) + \bar{\omega}^2)^2 + s^2\bar{\omega}^2, \\ \gamma_n &= \frac{i\bar{\omega}(n+\nu+i\bar{\omega})(n+\nu-s+i\bar{\omega})(n+\nu-s-i\bar{\omega})(n+\nu+1)}{(2n+2\nu-1)}. \end{aligned}$$

The overall normalization of the coefficients in the homogeneous equation (3.5) is, of course, irrelevant for the value of the a_n but the above form has the advantage that all denominators are bounded away from 0 in the perturbative (small-frequency) regime. By perturbative regime we essentially mean the frequency regime where ν is real—see the end of Sec. IV for further details. We choose the normalization $\check{a}_0 = a_0 = 1$, so then we have $\check{a}_n = a_n$, $\forall n \in \mathbb{Z}$, and from now on we will write down all series using the universal set of coefficients a_n . These coefficients a_n are equal, for $s = -2$, to the a_n^ν in MSTb as long as the same normalization is chosen for the two sets.

We could alternatively choose to include the Γ -functions in the coefficients, that is, write

$$a_n^{\text{RW}} \propto a_n \frac{\Gamma(-n-\nu+s-i\bar{\omega})\Gamma(n+\nu+s-i\bar{\omega}+1)}{\Gamma(1-2i\bar{\omega})}, \tag{3.6}$$

$$a_n^{\text{T}} \propto a_n \frac{\Gamma(-n-\nu-i\bar{\omega})\Gamma(n+\nu-i\bar{\omega}+1)}{\Gamma(1-s-2i\bar{\omega})}, \tag{3.7}$$

where the constants of proportionality are independent of n and so just reflect the normalization of the series. A particularly convenient choice is

$$a_n^{\text{RW}} \equiv a_n (-\nu+s-i\bar{\omega})_{-n} (\nu+s-i\bar{\omega}+1)_n, \tag{3.8}$$

$$a_n^{\text{T}} \equiv a_n (-\nu-i\bar{\omega})_{-n} (\nu-i\bar{\omega}+1)_n = a_n \frac{(\nu-i\bar{\omega}+1)_n}{(\nu+i\bar{\omega}+1)_n} (-1)^n. \tag{3.9}$$

As we choose the normalization $a_0 = 1$, we also have $a_0^{\text{T}} = a_0^{\text{RW}} = 1$. Using this convention the corresponding three-term recurrence relations have coefficients

$$\begin{aligned} \alpha_n^{\text{RW}} &= -\frac{n+\nu+1-s+i\bar{\omega}}{n+\nu+1+s-i\bar{\omega}} \alpha_n, & \beta_n^{\text{RW}} &= \beta_n, \\ \gamma_n^{\text{RW}} &= -\frac{n+\nu+s-i\bar{\omega}}{n+\nu-s+i\bar{\omega}} \gamma_n, \end{aligned} \tag{3.10}$$

$$\begin{aligned}\alpha_n^T &= -\frac{n+\nu+1+i\bar{\omega}}{n+\nu+1-i\bar{\omega}}\alpha_n, & \beta_n^T &= \beta_n, \\ \gamma_n^T &= -\frac{n+\nu-i\bar{\omega}}{n+\nu+i\bar{\omega}}\gamma_n.\end{aligned}\quad (3.11)$$

Note that, in the perturbative regime, where ν is real, Eq. (3.11) may be reexpressed as

$$\alpha_n^T = \alpha_n^*, \quad \beta_n^T = \beta_n^*, \quad \gamma_n^T = \gamma_n^*.$$

Up to irrelevant overall normalization, the coefficients α_n^T , β_n^T and γ_n^T and corresponding a_n^T are the same as the corresponding quantities in Eq. (123) ST.

As the corresponding coefficients differ by a scaling that tends to 1 for large $|n|$, ν is the same for RW

and for BPT. In particular, ν depends only on $|s|$ which can be seen directly since under the transformation $s \rightarrow -s$, $n+\nu \rightarrow -n-\nu-1$, β_n is invariant while α_n and γ_n are simply interchanged.

As the event horizon of the Schwarzschild black hole is approached, we have

$$\begin{aligned}X_s^{\text{in}} &\sim e^{i\bar{\omega}} e^{-i\omega r_*} \check{p}_s^{\text{in}}(0) \quad \text{and} \\ R_s^{\text{in}} &\sim e^{i\bar{\omega}} (\bar{r}-1)^{-s} e^{-i\omega r_*} p_s^{\text{in}}(0), \quad r \rightarrow 2M.\end{aligned}$$

That is, our solutions (3.4a) and (3.4b) are normalized according to [see Eqs. (2.14) and (2.32)]

$$X_s^{\text{in,tra}} = e^{i\bar{\omega}} \check{p}_s^{\text{in}}(0) = N_s^{\text{in}} e^{i\bar{\omega}} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(-n-\nu+s-i\bar{\omega})\Gamma(n+\nu+s+1-i\bar{\omega})}{\Gamma(1-2i\bar{\omega})}, \quad (3.12a)$$

$$R_s^{\text{in,tra}} = e^{i\bar{\omega}} (2M)^{2s} p_s^{\text{in}}(0) = N_s^{\text{in}} e^{i\bar{\omega}} (2M)^{2s} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(-n-\nu-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(1-s-2i\bar{\omega})}. \quad (3.12b)$$

We note that the particular normalization choice,

$$N_s^{\text{in}} = \frac{\Gamma(1-s-2i\bar{\omega})}{\Gamma(-\nu-i\bar{\omega})\Gamma(1+\nu-i\bar{\omega})}, \quad (3.13)$$

yields the specific normalization used in ST for the ingoing BPT solutions, and so that is our choice henceforth.

B. Series of Coulomb wave functions

An alternative expansion is useful for the construction of the ‘‘up’’ solutions. In terms of the variable $z \equiv \omega r = \bar{\omega} \bar{r}$, the RW equation may be written as

$${}_s X_\ell''(z) + \left(\frac{1}{z-\bar{\omega}} - \frac{1}{z} \right) {}_s X_\ell'(z) + \left(1 + \frac{2\bar{\omega}}{z-\bar{\omega}} + \frac{\bar{\omega}^2}{(z-\bar{\omega})^2} - \frac{\ell(\ell+1)}{z(z-\bar{\omega})} - \frac{(1-s^2)\bar{\omega}}{z^2(z-\bar{\omega})} \right) {}_s X_\ell(z) = 0.$$

Writing ${}_s X_\ell(z) = (1-\frac{\bar{\omega}}{z})^{-i\bar{\omega}} \check{h}_s(z)$ this becomes

$$z^2 \check{h}_s'' + [z^2 + 2\bar{\omega}z - \ell(\ell+1)] \check{h}_s' = \bar{\omega} \left[z(\check{h}_s'' + \check{h}_s') - (1-2i\bar{\omega})\check{h}_s' - (s^2 - (1-i\bar{\omega})^2) \frac{1}{z} \check{h}_s - \bar{\omega} \check{h}_s \right]. \quad (3.14)$$

The left-hand side is the operator defining the Coulomb wave equation [Eq. (33.14.1) [42]] with the solution satisfying ‘‘up’’ boundary conditions at infinity given by

$$\hat{H}_\ell^+(-\bar{\omega}, z) = W_{i\bar{\omega}, \ell+\frac{1}{2}}(-2iz) = e^{iz} (-2iz)^{\ell+1} U(\ell+1-i\bar{\omega}, 2\ell+2, -2iz),$$

where $\hat{H}_\ell^+(-\bar{\omega}, z)$ denotes the (unnormalized) irregular Coulomb function [Sec. 33.2(iii) [42]] and W denotes the Whittaker function (Sec. 13.14 [42]), We use a ‘‘hat’’ to denote that in writing the above we have dropped the conventional normalization prefactor $e^{-i\pi\ell/2+i\sigma_\ell(\bar{\omega})} e^{-\pi\bar{\omega}/2}$, where $\sigma_\ell(-\bar{\omega})$ is the Coulomb phase shift, which is irrelevant to our current discussion. U denotes the irregular confluent hypergeometric function (Sec. 13.2 [42], Ψ in the notation of [44]). Again, following Leaver [13] and MST, this suggests that we introduce the expansion for the upgoing solution,

$$\begin{aligned}
 h_s^{\text{up}}(z) &= N_s^{\text{up}} \sum_{n=-\infty}^{\infty} \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})} a_n \hat{H}_{n+\nu}^+(-\bar{\omega}, z) \\
 &= N_s^{\text{up}} e^{iz} (-2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})} a_n \\
 &\quad \times (-2iz)^n U(n+\nu+1-i\bar{\omega}, 2n+2\nu+2, -2iz),
 \end{aligned} \tag{3.15}$$

where N_s^{up} is a normalization constant that we will specify later so that our normalization agrees with ST. Inserting into Eq. (3.14), eliminating second derivatives using the differential equation satisfied by $\hat{H}_L^+(-\eta, z)$, and noting the following identities which follow from standard properties of the functions U :

$$\begin{aligned}
 \frac{1}{z} \hat{H}_L^+(-\eta, z) &= -i \frac{(L+1-i\eta)}{(L+1)(2L+1)} i \hat{H}_{L+1}^+(-\eta, z) + \frac{\eta}{L(L+1)} \hat{H}_L^+(-\eta, z) + i \frac{(L+i\eta)}{L(2L+1)} i \hat{H}_{L-1}^+(-\eta, z), \\
 \frac{d}{dz} \hat{H}_L^+(-\eta, z) &= i \frac{L(L+1-i\eta)}{(L+1)(2L+1)} i \hat{H}_{L+1}^+(-\eta, z) + \frac{\eta}{L(L+1)} \hat{H}_L^+(-\eta, z) + \frac{(L+1)(L+i\eta)}{L(2L+1)} i \hat{H}_{L-1}^+(-\eta, z),
 \end{aligned}$$

we find that a_n must satisfy the same three-term recurrence relation Eq. (3.5).

Similarly, the BPT equation becomes

$${}_s R_\ell''(z) + (1+s) \left(\frac{1}{z-\bar{\omega}} + \frac{1}{z} \right) {}_s R_\ell'(z) + \left(1 + \frac{2(\bar{\omega}+is)}{z-\bar{\omega}} + \frac{\bar{\omega}(\bar{\omega}-is)}{(z-\bar{\omega})^2} - \frac{\ell(\ell+1)-s(s+1)}{z(z-\bar{\omega})} \right) {}_s R_\ell(z) = 0.$$

Writing ${}_s R_\ell(z) = z^{-1-s} (1-\frac{\bar{\omega}}{z})^{-s-i\bar{\omega}} h_s(z)$ this becomes

$$z^2 h_s'' + [z^2 + 2(\bar{\omega}+is)z - \ell(\ell+1)] h_s = \bar{\omega} [z(h_s'' + h_s) - (1-s-2i\bar{\omega})h_s' - (1-i\bar{\omega})(s-1+i\bar{\omega})\frac{1}{z}h_s + i(s+i\bar{\omega})h_s]. \tag{3.16}$$

The appropriate Coulomb function is now $\hat{H}_l^+(-\bar{\omega}-is, z)$ and the corresponding expansion for the upgoing solution is

$$\begin{aligned}
 h_s^{\text{up}} &= (-1)^s N_s^{\text{up}} \sum_{n=-\infty}^{\infty} \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})} a_n \hat{H}_{n+\nu}^+(-\bar{\omega}-is, z) \\
 &= (-1)^s N_s^{\text{up}} e^{iz} (-2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})} a_n \\
 &\quad \times (-2iz)^n U(n+\nu+1+s-i\bar{\omega}, 2n+2\nu+2, -2iz).
 \end{aligned}$$

The Chandrasekhar transformations in this case follow term by term from the Whittaker function identities:

$$\begin{aligned}
 z \left(1 - \frac{\bar{\omega}}{z} \right) \bar{D}_0 \left(\left(1 - \frac{\bar{\omega}}{z} \right)^{-i\bar{\omega}} W_{i\bar{\omega}, L+\frac{1}{2}}(-2iz) \right) &= - \left(1 - \frac{\bar{\omega}}{z} \right)^{1-i\bar{\omega}} W_{1+i\bar{\omega}, L+\frac{1}{2}}(-2iz), \\
 z^2 \left(1 - \frac{\bar{\omega}}{z} \right)^2 \bar{D}_0^2 \left(z \left(1 - \frac{\bar{\omega}}{z} \right)^{-i\bar{\omega}} W_{i\bar{\omega}, L+\frac{1}{2}}(-2iz) \right) &= z \left(1 - \frac{\bar{\omega}}{z} \right)^{2-i\bar{\omega}} W_{2+i\bar{\omega}, L+\frac{1}{2}}(-2iz),
 \end{aligned}$$

or equivalently

$$\begin{aligned}
 z \left(1 - \frac{\bar{\omega}}{z} \right) \bar{D}_0 \left(\left(1 - \frac{\bar{\omega}}{z} \right)^{-i\bar{\omega}} \hat{H}_{n+\nu}^+(-\bar{\omega}, z) \right) &= - \left(1 - \frac{\bar{\omega}}{z} \right)^{1-i\bar{\omega}} \hat{H}_{n+\nu}^+(-\bar{\omega}+i, z), \\
 z^2 \left(1 - \frac{\bar{\omega}}{z} \right)^2 \bar{D}_0^2 \left(z \left(1 - \frac{\bar{\omega}}{z} \right)^{-i\bar{\omega}} \hat{H}_{n+\nu}^+(-\bar{\omega}, z) \right) &= z \left(1 - \frac{\bar{\omega}}{z} \right)^{2-i\bar{\omega}} \hat{H}_{n+\nu}^+(-\bar{\omega}+2i, z),
 \end{aligned}$$

where

$$\bar{D}_0 = \bar{\omega} \left(\frac{d}{dz} + i \left(1 - \frac{\bar{\omega}}{z} \right)^{-1} \right).$$

Using the relationship between a_n and a_n^T we may write our solution in the alternate form,

$$h_s^{\text{up}} = (-1)^s \frac{1}{2} e^{-\pi\bar{\omega}} e^{iz} (-2z)^{\nu+1} \sum_{n=-\infty}^{\infty} \frac{(\nu+1+s-i\bar{\omega})_n}{(\nu+1-s+i\bar{\omega})_n} \times a_n^T (2iz)^n U(n+\nu+1+s-i\bar{\omega}, 2n+2\nu+2, -2iz), \quad (3.17)$$

where we have made the choice

$$N_s^{\text{up}} = \frac{1}{2} e^{-\pi\bar{\omega}} \frac{\Gamma(\nu+1-s+i\bar{\omega})\Gamma(\nu+1+i\bar{\omega})}{\Gamma(\nu+1+s-i\bar{\omega})\Gamma(\nu+1-i\bar{\omega})} e^{-i\frac{\pi}{2}(\nu+1)},$$

so as to agree with the normalization of ST. However our original form serves to highlight the boundary conditions and the link to the RW solution.

The corresponding upgoing BPT solution is given by

$$R_s^{\text{up}} = z^{-1-s} \left(1 - \frac{\bar{\omega}}{z} \right)^{-s-i\bar{\omega}} h_s^{\text{up}}(z). \quad (3.18)$$

This series representation in terms of irregular confluent hypergeometric functions converges $\forall r > 2M$. Since

$$k_s^{\text{up}}(z) = \sum_{n=-\infty}^{\infty} \frac{\Gamma(n+\nu-s+1-i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})}{\Gamma(n+\nu+s+1+i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})} a_n \hat{H}_{n+\nu}^+(-\bar{\omega}, z) \\ = e^{iz} (-2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} \frac{\Gamma(n+\nu-s+1-i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})}{\Gamma(n+\nu+s+1+i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})} a_n (-2iz)^n U(n+\nu+1-i\bar{\omega}, 2n+2\nu+2, -2iz).$$

While this expansion seems to more naturally capture the boundary conditions of the RW equation, the Chandrasekhar transformation is less natural in terms of it, so we will discuss it no further.

C. Relation between the two solutions

Using the standard relation Eq. (15.8.3) [42], we may reexpress X_s^{in} in terms that are better suited for discussing its behavior at radial infinity:

$$X_s^{\text{in}} = N_s^{\text{in}} (1-x)^{\nu+1+i\bar{\omega}} e^{i\bar{\omega}x} (-x)^{-i\bar{\omega}} \sum_{n=-\infty}^{\infty} \frac{\Gamma(-n-\nu+s-i\bar{\omega})\Gamma(2n+2\nu+1)}{\Gamma(n+\nu+1-s-i\bar{\omega})} a_n \\ \times (1-x)^n {}_2F_1 \left(-n-\nu+s-i\bar{\omega}, -n-\nu-s-i\bar{\omega}; -2n-2\nu; \frac{1}{1-x} \right) \\ + N_s^{\text{in}} (1-x)^{-\nu+i\bar{\omega}} e^{i\bar{\omega}x} (-x)^{-i\bar{\omega}} \sum_{n=-\infty}^{\infty} \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(-2n-2\nu-1)}{\Gamma(-n-\nu-s-i\bar{\omega})} a_n \\ \times (1-x)^{-n} {}_2F_1 \left(n+\nu+1+s-i\bar{\omega}, n+\nu+1-s-i\bar{\omega}; 2n+2\nu+2; \frac{1}{1-x} \right).$$

$U(a, c, x) \sim x^{-a}$ as $|x| \rightarrow \infty$, it is straightforward to write down the asymptotic forms

$$X_s^{\text{up}} \sim (2i)^s A_-^\nu e^{i(z+\bar{\omega} \ln z)}, \\ R_s^{\text{up}} \sim A_-^\nu z^{-1-2s} e^{i(z+\bar{\omega} \ln z)},$$

where

$$A_-^\nu \equiv (2i)^{-s} e^{-\frac{\pi}{2}\bar{\omega}} e^{-i\frac{\pi}{2}(\nu+1)} 2^{-1+i\bar{\omega}} \\ \times \sum_{n=-\infty}^{\infty} \frac{(\nu+1+s-i\bar{\omega})_n (\nu+1-i\bar{\omega})_n}{(\nu+1-s+i\bar{\omega})_n (\nu+1+i\bar{\omega})_n} a_n. \quad (3.19)$$

Clearly, then, from Eq. (2.33),

$$X_s^{\text{up,tra}} = (2i)^s A_-^\nu e^{i\bar{\omega} \ln \bar{\omega}}, \\ R_s^{\text{up,tra}} = A_-^\nu \bar{\omega}^{-1-2s} e^{i\bar{\omega} \ln \bar{\omega}}. \quad (3.20)$$

To conclude this subsection, we note that we could have used the ansatz ${}_s X_\ell(z) = (1 - \frac{\bar{\omega}}{z})^{i\bar{\omega}} k_s(z)$ for the RW equation, giving

$$z^2 k_s'' + [z^2 + 2\bar{\omega}z - \ell(\ell+1)] k_s \\ = \bar{\omega} [z(k_s' + k_s) - (1 + 2i\bar{\omega})k_s' \\ - (s^2 - (1+i\bar{\omega})^2) \frac{1}{z} k_s - \bar{\omega}k_s], \quad (3.21)$$

and the upgoing solution

The second term can be obtained from the first by the substitution $n \rightarrow -n$, $\nu \rightarrow -\nu - 1$ and correspondingly the terms are denoted by X_0^ν and $X_0^{-\nu-1}$ respectively, so

$$X_s^{\text{in}} = X_0^\nu + X_0^{-\nu-1}. \quad (3.22)$$

(Note that N_s^{in} is invariant under this transformation.) It is easily checked that each of these terms is independently a solution of the RW equation and moreover are linearly independent (MSTb). In terms of $z = \bar{\omega} \bar{r} = \bar{\omega}(1-x)$,

$$\begin{aligned} X_0^\nu(z) &= N_s^{\text{in}} e^{i\bar{\omega}} e^{-iz} \left(\frac{z}{\bar{\omega}} - 1\right)^{-i\bar{\omega}} \left(\frac{z}{\bar{\omega}}\right)^{\nu+i\bar{\omega}} \sum_{n=-\infty}^{\infty} \frac{\Gamma(-n-\nu+s-i\bar{\omega})\Gamma(2n+2\nu+1)}{\Gamma(n+\nu+1-s-i\bar{\omega})} a_n \\ &\times \left(\frac{z}{\bar{\omega}}\right)^n {}_2F_1\left(-n-\nu-i\bar{\omega}, -n-\nu-s-i\bar{\omega}; -2n-2\nu; \frac{\bar{\omega}}{z}\right). \end{aligned} \quad (3.23)$$

In an identical fashion we can write

$$R_s^{\text{in}} = R_0^\nu + R_0^{-\nu-1} \quad (3.24)$$

with

$$\begin{aligned} R_0^\nu &= N_s^{\text{in}} e^{i\bar{\omega}} e^{-iz} \left(\frac{z}{\bar{\omega}} - 1\right)^{-s-i\bar{\omega}} \left(\frac{z}{\bar{\omega}}\right)^{\nu+i\bar{\omega}} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(-n-\nu-i\bar{\omega})\Gamma(2n+2\nu+1)}{\Gamma(n+\nu+1-s-i\bar{\omega})} \\ &\times \left(\frac{z}{\bar{\omega}}\right)^n {}_2F_1\left(-n-\nu-i\bar{\omega}, -n-\nu-s-i\bar{\omega}; -2n-2\nu; \frac{\bar{\omega}}{z}\right). \end{aligned} \quad (3.25)$$

This series representation for R_s^{in} converges $\forall r > 2M$.

To obtain solutions in terms of confluent hypergeometric functions suitable for discussing the behavior at the horizon, we introduce the auxiliary solutions involving the regular Coulomb wave function,

$$\begin{aligned} \hat{F}_\ell(-\bar{\omega}, z) &= \frac{\Gamma(\ell+1+i\bar{\omega})}{\Gamma(2\ell+2)} M_{i\bar{\omega}, \ell+\frac{1}{2}}(-2iz) \\ &= \frac{\Gamma(\ell+1+i\bar{\omega})}{\Gamma(2\ell+2)} e^{iz} (-2iz)^{\ell+1} M(\ell+1-i\bar{\omega}, 2\ell+2, -2iz) \\ &= \frac{\Gamma(\ell+1+i\bar{\omega})}{\Gamma(2\ell+2)} e^{-iz} (-2iz)^{\ell+1} M(\ell+1+i\bar{\omega}, 2\ell+2, 2iz), \end{aligned}$$

where we include the prefactor for reasons that will become clear when we consider the Chandrasekhar transformation below. The function $\hat{F}_\ell(-\eta, z)$ then satisfies the following identities:

$$\begin{aligned} \frac{1}{z} \hat{F}_L(-\eta, z) &= i \frac{(L+1-i\eta)}{(L+1)(2L+1)} \hat{F}_{L+1}(-\eta, z) + \frac{\eta}{L(L+1)} \hat{F}_L(-\eta, z) - i \frac{(L+i\eta)}{L(2L+1)} \hat{F}_{L-1}(-\eta, z), \\ \frac{d}{dz} \hat{F}_L(-\eta, z) &= -i \frac{L(L+1-i\eta)}{(L+1)(2L+1)} \hat{F}_{L+1}(-\eta, z) + \frac{\eta}{L(L+1)} \hat{F}_L(-\eta, z) - i \frac{(L+1)(L+i\eta)}{L(2L+1)} \hat{F}_{L-1}(-\eta, z). \end{aligned}$$

Proceeding as before we construct the corresponding solutions:

$$\begin{aligned} X_C^\nu(z) &= N_C^\nu e^{-iz} \left(1 - \frac{\bar{\omega}}{z}\right)^{-i\bar{\omega}} (-2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})\Gamma(2n+2\nu+2)} (2iz)^n \\ &\times M(n+\nu+1+i\bar{\omega}, 2n+2\nu+2, 2iz) \end{aligned} \quad (3.26)$$

and

$$R_C^\nu(z) = N_C^\nu e^{-iz} \left(1 - \frac{\bar{\omega}}{z}\right)^{-s-i\bar{\omega}} z^{-1-s} (-2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1+i\bar{\omega})\Gamma(2n+2\nu+2)} (2iz)^n \times M(n+\nu+1-s+i\bar{\omega}, 2n+2\nu+2, 2iz), \quad (3.27)$$

where

$$N_C^\nu = \frac{\Gamma(\nu+1-s+i\bar{\omega})\Gamma(\nu+1+i\bar{\omega})}{2\Gamma(\nu+1+s-i\bar{\omega})\Gamma(\nu+1-i\bar{\omega})} e^{i\frac{\pi}{2}(\nu+1)}, \quad (3.28)$$

for agreement with Eq. (139) ST when $z > 0$. Our normalization for X_C^ν follows from that of R_C^ν via the Chandrasekhar transformation, and so for $s = -2$ it does not coincide with the normalization choice in MSTb. Specifically, for $s = -2$, our X_C^ν is equal to that in MSTb times

$$\frac{\Gamma(\nu+1-s+i\bar{\omega})\Gamma(\nu+1+i\bar{\omega})}{\Gamma(\nu-1-i\bar{\omega})\Gamma(\nu+1-i\bar{\omega})}.$$

The Chandrasekhar transformation when using these series representations follows term by term from the Whittaker function identities:

$$z \left(1 - \frac{\bar{\omega}}{z}\right) \bar{D}_0 \left(\left(1 - \frac{\bar{\omega}}{z}\right)^{-i\bar{\omega}} M_{i\bar{\omega}, L+\frac{1}{2}}(-2iz) \right) = (L+1+i\bar{\omega}) \left(1 - \frac{\bar{\omega}}{z}\right)^{1-i\bar{\omega}} M_{1+i\bar{\omega}, L+\frac{1}{2}}(-2iz),$$

$$z^2 \left(1 - \frac{\bar{\omega}}{z}\right)^2 \bar{D}_0^2 \left(z \left(1 - \frac{\bar{\omega}}{z}\right)^{-i\bar{\omega}} M_{i\bar{\omega}, L+\frac{1}{2}}(-2iz) \right) = (L+1+i\bar{\omega})(L+2+i\bar{\omega}) z \left(1 - \frac{\bar{\omega}}{z}\right)^{2-i\bar{\omega}} M_{2+i\bar{\omega}, L+\frac{1}{2}}(-2iz),$$

or equivalently

$$z \left(1 - \frac{\bar{\omega}}{z}\right) \bar{D}_0 \left(\left(1 - \frac{\bar{\omega}}{z}\right)^{-i\bar{\omega}} \hat{F}_{n+\nu}^+(-\bar{\omega}, z) \right) = \left(1 - \frac{\bar{\omega}}{z}\right)^{1-i\bar{\omega}} \hat{F}_{n+\nu}^+(-\bar{\omega} + i, z),$$

$$z^2 \left(1 - \frac{\bar{\omega}}{z}\right)^2 \bar{D}_0^2 \left(z \left(1 - \frac{\bar{\omega}}{z}\right)^{-i\bar{\omega}} \hat{F}_{n+\nu}^+(-\bar{\omega}, z) \right) = z \left(1 - \frac{\bar{\omega}}{z}\right)^{2-i\bar{\omega}} \hat{F}_{n+\nu}^+(-\bar{\omega} + 2i, z),$$

where

$$\bar{D}_0 = \bar{\omega} \left(\frac{d}{dz} + i \left(1 - \frac{\bar{\omega}}{z}\right)^{-1} \right).$$

These solutions can be related to X_s^{up} and R_s^{up} using the identity in Eq. (6.7(7)), Vol. 1 [45] (valid for $b \notin \mathbb{Z}$):

$$M(a, b, 2iz) = \begin{cases} \frac{\Gamma(b)}{\Gamma(b-a)} e^{a\pi i} U(a, b, 2iz) + \frac{\Gamma(b)}{\Gamma(a)} e^{(a-b)\pi i} e^{2iz} U(b-a, b, -2iz), & \text{Re}(z) > 0, \\ \frac{\Gamma(b)}{\Gamma(b-a)} e^{-a\pi i} U(a, b, 2iz) + \frac{\Gamma(b)}{\Gamma(a)} e^{-(a-b)\pi i} e^{2iz} U(b-a, b, -2iz), & \text{Re}(z) < 0, \end{cases}$$

the first and second terms on each line yielding the incoming and outgoing wave solutions at infinity, respectively. Thus, $X_C^\nu(z) = X_+^\nu(z) + X_-^\nu(z)$, where

$$X_+^\nu(z) = N_C^\nu e^{\mp\pi\bar{\omega}} e^{-iz} \left(1 - \frac{\bar{\omega}}{z}\right)^{-i\bar{\omega}} (2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(n+\nu+1+s-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})} (-2iz)^n \times U(n+\nu+1+i\bar{\omega}, 2n+2\nu+2, 2iz)$$

$$X_-^\nu(z) = N_C^\nu e^{\mp\pi\bar{\omega}} e^{\mp 2(\nu+1)\pi i} e^{iz} \left(1 - \frac{\bar{\omega}}{z}\right)^{-i\bar{\omega}} (2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})} (-2iz)^n \times U(n+\nu+1-i\bar{\omega}, 2n+2\nu+2, -2iz), \quad (3.29)$$

where the signs correspond to $\frac{\text{Re}(z)>0}{\text{Re}(z)<0}$. Here we take the BC on the complex z -plane to lie along $z < 0$.⁶ Similarly, $R_C^\nu(z) = R_+^\nu(z) + R_-^\nu(z)$, where

$$\begin{aligned}
 R_+^\nu(z) &= (-1)^s N_C^\nu e^{\mp \pi \bar{\omega}} e^{-iz} \left(1 - \frac{\bar{\omega}}{z}\right)^{-s-i\bar{\omega}} z^{-1-s} (2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1+i\bar{\omega})} (-2iz)^n \\
 &\quad \times U(n+\nu+1-s+i\bar{\omega}, 2n+2\nu+2, 2iz), \\
 R_-^\nu(z) &= (-1)^s N_C^\nu e^{\mp \pi \bar{\omega}} e^{\mp 2(\nu+1)\pi i} e^{iz} \left(1 - \frac{\bar{\omega}}{z}\right)^{-s-i\bar{\omega}} z^{-1-s} (2iz)^{\nu+1} \sum_{n=-\infty}^{\infty} a_n \frac{\Gamma(n+\nu+1+s-i\bar{\omega})\Gamma(n+\nu+1-i\bar{\omega})}{\Gamma(n+\nu+1-s+i\bar{\omega})\Gamma(n+\nu+1+i\bar{\omega})} (-2iz)^n \\
 &\quad \times U(n+\nu+1+s-i\bar{\omega}, 2n+2\nu+2, -2iz). \tag{3.30}
 \end{aligned}$$

Note that our naming convention here follows ST and is opposite in sign to MSTb. In particular, the minus solutions are just multiples of the corresponding ‘‘up’’ solutions.

Critically as noted by MST, The functions X_0^ν and X_C^ν solve the same differential equation and have the same analytical behavior as functions of z ; similarly for R_0^ν and R_C^ν . Therefore, they must be proportional:

$$X_0^\nu = K_\nu X_C^\nu, \quad \text{and} \quad R_0^\nu = K_\nu R_C^\nu, \tag{3.31}$$

where K_ν is the constant of proportionality. Equating the corresponding Laurent series we can obtain explicit expressions for K_ν in terms of ν and our a_n coefficients. The results are given for general spin by ST (based on their Teukolsky equation analysis), we repeat them here for completeness specialized to Schwarzschild spacetime:

$$\begin{aligned}
 K_\nu &= \frac{e^{i\bar{\omega}} (2\bar{\omega})^{s-\nu-r} 2^{-s} i^r \Gamma(1-s-2i\bar{\omega}) \Gamma(r+2\nu+2)}{\Gamma(r+\nu+1-s+i\bar{\omega}) \Gamma(r+\nu+1+i\bar{\omega}) \Gamma(r+\nu+1+s+i\bar{\omega})} \\
 &\quad \times \left(\sum_{n=r}^{\infty} \frac{\Gamma(n+r+2\nu+1)}{(n-r)!} \frac{\Gamma(n+\nu+1+s+i\bar{\omega})}{\Gamma(n+\nu+1-s-i\bar{\omega})} \frac{\Gamma(\nu+1+i\bar{\omega})}{\Gamma(\nu+1-i\bar{\omega})} a_n \right) \\
 &\quad \times \left(\sum_{n=-\infty}^r \frac{1}{(r-n)! (r+2\nu+2)_n} \frac{(\nu+1+s-i\bar{\omega})_n (\nu+1-i\bar{\omega})_n}{(\nu+1-s+i\bar{\omega})_n (\nu+1+i\bar{\omega})_n} a_n \right)^{-1}. \tag{3.32}
 \end{aligned}$$

The parameter r here is an arbitrary integer number.

The above expressions lay out the foundations for taking the $\bar{r} \rightarrow \infty$ limit of the ingoing solutions and thus finding their incidence and reflection coefficients. First we relate the quantities at $-\nu-1$ to those at ν . From Eq. (3.28) we have

$$N_C^{-\nu-1} = \frac{\sin(\pi(\nu+i\bar{\omega})) \sin(\pi(\nu+i\bar{\omega}-s))}{\sin(\pi(\nu-i\bar{\omega})) \sin(\pi(\nu-i\bar{\omega}+s))} e^{-i\pi\nu} e^{-i\frac{\pi}{2}} N_C^\nu. \tag{3.33}$$

Then, from Eq. (3.29) it follows that

$$X_+^{-\nu-1} = \frac{\sin(\pi(\nu+i\bar{\omega}))}{\sin(\pi(\nu-i\bar{\omega}))} e^{-i\pi\nu} e^{-i\frac{\pi}{2}} X_+^\nu \tag{3.34}$$

⁶This is different from the choice in the other sections of the paper of BC on the complex ω -plane along the NIA. However, that does not matter for $\text{Re}(\omega) > 0$, to which we restrict ourselves.

and

$$X_-^{-\nu-1} = e^{\pm 2i\pi\nu} e^{-i\pi\nu} e^{i\frac{\pi}{2}} X_-^\nu, \tag{3.35}$$

where the upper/lower sign corresponds to $\text{Re}(z)$ positive/negative. From now on we assume $\text{Re}(\omega) > 0$ [for $\text{Re}(\omega) < 0$ one may use the symmetries of the radial solutions].

From Eqs. (3.22) and (3.29) and (13.7.3) [42] we can take the $r \rightarrow \infty$ limit and obtain the incidence and reflection coefficients of the ingoing RW solution:

$$X_s^{\text{in,inc}} = \left(K_\nu - i e^{-i\pi\nu} \frac{\sin(\pi(\nu+i\bar{\omega}))}{\sin(\pi(\nu-i\bar{\omega}))} K_{-\nu-1} \right) \check{A}_+^\nu e^{-i\bar{\omega} \ln \bar{\omega}} \tag{3.36}$$

and

$$X_s^{\text{in,ref}} = (K_\nu + i e^{i\pi\nu} K_{-\nu-1}) (2i)^s A_-^\nu e^{i\bar{\omega} \ln \bar{\omega}}, \tag{3.37}$$

where

$$\begin{aligned} \check{A}_+^\nu &\equiv e^{-\pi\bar{\omega}/2} e^{i\frac{\pi}{2}(\nu+1)} 2^{-1-i\bar{\omega}} \frac{\Gamma(\nu+1-s+i\bar{\omega})\Gamma(\nu+i\bar{\omega}+1)}{\Gamma(\nu+1+s-i\bar{\omega})\Gamma(\nu-i\bar{\omega}+1)} \\ &\times \sum_{n=-\infty}^{\infty} (-1)^n \frac{\Gamma(n+\nu-i\bar{\omega}+1+s)}{\Gamma(n+\nu+i\bar{\omega}+1-s)} a_n. \end{aligned} \quad (3.38)$$

We can proceed similarly for the BPT solution. By taking $z \rightarrow \infty$ in Eq. (3.24) we obtain the incidence and reflection coefficients of the ingoing BPT solution:

$$\begin{aligned} R_s^{\text{in,inc}} &= \omega^{-1} \left(K_\nu - i e^{-i\pi\nu} \frac{\sin(\pi(\nu-s+i\bar{\omega}))}{\sin(\pi(\nu+s-i\bar{\omega}))} K_{-\nu-1} \right) \\ &\times A_+^\nu e^{-i\bar{\omega} \ln \bar{\omega}} \end{aligned} \quad (3.39)$$

and

$$R_s^{\text{in,ref}} = \omega^{-1-2s} (K_\nu + i e^{i\pi\nu} K_{-\nu-1}) A_+^\nu e^{i\bar{\omega} \ln \bar{\omega}}, \quad (3.40)$$

where

$$\begin{aligned} A_+^\nu &\equiv e^{-\pi\bar{\omega}/2} e^{i\frac{\pi}{2}(\nu+1-s)} 2^{-1+s-i\bar{\omega}} \\ &\times \frac{\Gamma(\nu+1-s+i\bar{\omega})\Gamma(\nu+i\bar{\omega}+1)}{\Gamma(\nu+1+s-i\bar{\omega})\Gamma(\nu-i\bar{\omega}+1)} \\ &\times \sum_{n=-\infty}^{\infty} (-1)^n \frac{\Gamma(n+\nu-i\bar{\omega}+1)}{\Gamma(n+\nu+i\bar{\omega}+1)} a_n. \end{aligned} \quad (3.41)$$

We note that one can obtain the radial incidence, reflection and transmission coefficients of the RW solution from those of the BPT solution (or vice versa) via the Chandrasekhar transformation Eq. (2.40) [or Eq. (2.39)]. In doing so, the leading order for large- r would be annihilated and one would require a higher order term.

IV. LOW-FREQUENCY EXPANSION OF THE COEFFICIENTS AND OF ν

In this section our goal is to provide the low-frequency behavior of the MST series renormalized angular momentum ν and series coefficients a_n . We will provide the expansions explicitly up to the first five leading orders. The behavior of the coefficients a_n^{RW} and a_n^{T} may be deduced immediately from Eqs. (3.8) and (3.9).

We start by noting that Eqs. (3.2) and (3.3) reduce to the hypergeometric equation when $\bar{\omega} = 0$, indeed it was

$$\begin{aligned} a_4 &= \frac{(\ell-s+1)^2(\ell-s+2)^2(\ell-s+3)^2(\ell-s+4)^2}{96(\ell+1)(\ell+2)(2\ell+1)(2\ell+3)^2(2\ell+5)^2(2\ell+7)} \bar{\omega}^4 + O(\bar{\omega}^5), \\ a_3 &= \frac{(\ell-s+1)^2(\ell-s+2)^2(\ell-s+3)^2}{24(\ell+1)(\ell+2)(2\ell+1)(2\ell+3)^2(2\ell+5)} i\bar{\omega}^3 - \frac{(3\ell^2+12\ell+11)(\ell-s+1)^2(\ell-s+2)^2(\ell-s+3)^2}{24(\ell+1)^2(\ell+2)^2(\ell+3)(2\ell+1)(2\ell+3)^2(2\ell+5)} \bar{\omega}^4 + O(\bar{\omega}^5), \\ a_2 &= -\frac{(\ell-s+1)^2(\ell-s+2)^2}{4(\ell+1)(2\ell+1)(2\ell+3)^2} \bar{\omega}^2 - \frac{(\ell-s+1)^2(\ell-s+2)^2}{4(\ell+1)^2(\ell+2)(2\ell+1)(2\ell+3)} i\bar{\omega}^3 \\ &+ \frac{P_{2,4}^{(15,8)}(\ell, s)}{48\ell(\ell+1)^3(\ell+2)(\ell+3)(2\ell-1)(2\ell+1)^3(2\ell+3)^4(2\ell+7)} \bar{\omega}^4 + O(\bar{\omega}^5), \end{aligned}$$

precisely for this reason that Leaver [13] and MST wrote them in this way. For the ‘‘in’’ RW and BPT solutions we want the regular solutions corresponding to, respectively,

$${}_2F_1(\ell+1+s, -\ell+s, 1; x) = {}_2F_1(-\ell+s, \ell+1+s, 1; x)$$

and

$${}_2F_1(\ell+1, -\ell, 1-s; x) = {}_2F_1(-\ell, \ell+1, 1-s; x).$$

The left-hand sides of these expressions correspond to $\nu = \ell$ and the right-hand sides to $\nu = -\ell - 1$ when $\bar{\omega} = 0$. In fact, under $\nu \rightarrow -\nu - 1$, α_n equals γ_{-n} and β_n equals β_{-n} and therefore, a_n satisfies the same recurrence relation as a_{-n} under $\nu \rightarrow -\nu - 1$; the equivalent symmetries hold for the RW counterparts (a_n^{RW} , α_n^{RW} , β_n^{RW} and γ_n^{RW}) and for the BPT counterparts (a_n^{T} , α_n^{T} , β_n^{T} and γ_n^{T}). This symmetry stems from the fact that the renormalized angular momentum ν was introduced into the ordinary differential equation [Eq. (119) ST] in the form $\nu(\nu+1)$, which is invariant under $\nu \rightarrow -\nu - 1$. In addition, we may determine the expansion of ν about ℓ from that about ‘‘ $-\ell - 1$.’’

With the natural ansatz that $a_n = O(\bar{\omega}^{|n|})$, the 3-term recurrence relation [Eq. (3.5)] can be solved directly yielding

$$\begin{aligned} \nu + \frac{1}{2} &= \left(\ell + \frac{1}{2} \right) \left[1 - \frac{\lambda(15\lambda-11)+3s^4+6(\lambda-1)s^2}{\lambda(4\lambda-3)(4\lambda+1)} \bar{\omega}^2 \right. \\ &\left. + \frac{P_{0,4}^{(8,4)}(\lambda, s^2)}{4(\lambda-2)\lambda^3(4\lambda-15)(4\lambda-3)^3(4\lambda+1)^2} \bar{\omega}^4 + O(\bar{\omega}^6) \right] \end{aligned} \quad (4.1)$$

for the renormalized angular momentum. Note that the expansion of ν must be even in s in Schwarzschild spacetime since it also arises through the expansion of the RW equation which manifestly has this property. As it will be needed later on, let us define ν_2 as minus the coefficient of $\bar{\omega}^2$ in the above expansion for ν , i.e.,

$$\nu = \ell - \nu_2 \bar{\omega}^2 + O(\bar{\omega}^3). \quad (4.2)$$

For the series coefficients themselves, the above procedure yields

$$\begin{aligned}
 a_1 &= -\frac{(\ell-s+1)^2}{2(\ell+1)(2\ell+1)}i\bar{\omega} + \frac{(\ell-s+1)^2}{2(\ell+1)^2(2\ell+1)}\bar{\omega}^2 + \frac{P_{1,3}^{(12,6)}(\ell,s)}{8\ell(\ell+1)^3(\ell+2)(2\ell-1)(2\ell+1)^3(2\ell+3)^2(2\ell+5)}i\bar{\omega}^3 \\
 &\quad + \frac{P_{1,4}^{(12,6)}(\ell,s)}{8\ell(\ell+1)^4(\ell+2)(2\ell-1)(2\ell+1)^3(2\ell+3)^2(2\ell+5)}\bar{\omega}^4 + O(\bar{\omega}^5), \\
 a_{-1} &= -\frac{(\ell+s)^2}{2\ell(2\ell+1)}i\bar{\omega} - \frac{(\ell+s)^2}{2\ell^2(2\ell+1)}\bar{\omega}^2 + \frac{P_{-1,3}^{(12,6)}(\ell,s)}{8(\ell-1)\ell^3(\ell+1)(2\ell-3)(2\ell-1)^2(2\ell+1)^3(2\ell+3)}i\bar{\omega}^3 \\
 &\quad + \frac{P_{-1,4}^{(12,6)}(\ell,s)}{8(\ell-1)\ell^4(\ell+1)(2\ell-3)(2\ell-1)^2(2\ell+1)^3(2\ell+3)}\bar{\omega}^4 + O(\bar{\omega}^5), \\
 a_{-2} &= -\frac{(\ell+s-1)^2(\ell+s)^2}{4\ell(2\ell-1)^2(2\ell+1)}\bar{\omega}^2 + \frac{(\ell+s-1)^2(\ell+s)^2}{4(\ell-1)\ell^2(2\ell-1)(2\ell+1)}i\bar{\omega}^3 \\
 &\quad + \frac{P_{-2,4}^{(15,8)}(\ell,s)}{48(\ell-2)(\ell-1)\ell^3(2\ell-5)(2\ell-1)^4(2\ell+1)^3(2\ell+3)}\bar{\omega}^4 + O(\bar{\omega}^5), \\
 a_{-3} &= \frac{(\ell+s-2)^2(\ell+s-1)^2(\ell+s)^2}{24(\ell-1)\ell(2\ell-3)(2\ell-1)^2(2\ell+1)}i\bar{\omega}^3 + \frac{P_{-3,4}(15,8)(\ell,s)}{24(\ell-2)(\ell-1)^2\ell^2(2\ell-3)(2\ell-1)^2(2\ell+1)}\bar{\omega}^4 + O(\bar{\omega}^5), \\
 a_{-4} &= \frac{(\ell+s-3)^2(\ell+s-2)^2(\ell+s-1)^2(\ell+s)^2}{96(\ell-1)\ell(2\ell-5)(2\ell-3)^2(2\ell-1)^2(2\ell+1)}\bar{\omega}^4 + O(\bar{\omega}^5), \tag{4.3}
 \end{aligned}$$

where $P^{(m,n)}(x, y)$ denotes a real polynomial with integer coefficients of degree m in x and n in y and recall $\lambda = l(l + 1)$. The precise form of the polynomials is easily determined to high order but is too long to be useful in printed form for general s . For completeness, we give, as the most important example, the term in the renormalized angular momentum for $s = 0, -1, -2$:

$$\begin{aligned}
 P_{0,4}^{(8,4)}(\lambda, 0) &= -\lambda^2(\lambda - 2)(3240 + 8733\lambda - 82625\lambda^2 + 155295\lambda^3 - 105000\lambda^4 + 18480\lambda^5), \\
 P_{0,4}^{(8,4)}(\lambda, 1) &= -(\lambda - 2)(405 + 1917\lambda - 3240\lambda^2 - 16305\lambda^3 + 19435\lambda^4 + 54495\lambda^5 - 84840\lambda^6 + 18480\lambda^7), \\
 P_{0,4}^{(8,4)}(\lambda, 4) &= 51840 + 102816\lambda - 953424\lambda^2 + 51522\lambda^3 + 123233\lambda^4 + 85775\lambda^5 - 2415\lambda^6 + 61320\lambda^7 - 18480\lambda^8.
 \end{aligned}$$

The naive pattern evident in the leading behavior under the ansatz $a_n = O(\bar{\omega}^{|n|})$ is

$$a_n^{\text{naive}} \equiv \begin{cases} \frac{((\ell+1-s)_n)^2(\ell+1)_n 2^n}{(2\ell+2)_n(2\ell+1)_{2n} n!} (-i\bar{\omega})^{|n|} + O(\bar{\omega}^{n+1}) & n > 0, \\ \frac{((-\ell-s)_n)^2(-\ell)_n 2^{|n|}}{(-2\ell)_{|n|}(-2\ell-1)_{2|n|} |n|!} (-i\bar{\omega})^{|n|} + O(\bar{\omega}^{n+1}) & n < 0, \end{cases} \tag{4.4}$$

where $(x)_n$ denotes the Pochhammer symbol $(x)_n = \Gamma(x+n)/\Gamma(x) = x(x+1)\dots(x+n-1)$. Note that reflecting the symmetry noted at the beginning of the section, we have $a_n \rightarrow a_{-n}$ under $\ell \rightarrow -\ell - 1$. We have also derived Eq. (4.4) via an alternative method: by imposing that Eq. (3.32) for K_ν is independent of the parameter r .

Critically, Eq. (4.4) reveals that the ansatz $a_n = O(\bar{\omega}^{|n|})$ is flawed since the denominator vanishes whenever $|n| \geq 2\ell + 1$ while, in addition, the numerator vanishes when $|n| \geq \ell + s + 1$. A detailed analysis reveals that the correct ansatz for $\nu = \ell + O(\bar{\omega}^2)$ and $s = 0$ is

$$a_n = \begin{cases} O(\bar{\omega}^{|n|}) & n \geq -\ell, \\ O(\bar{\omega}^{|n|+1}) & -(2\ell + 1) < n < -\ell, \\ O(\bar{\omega}^{|n|-1}) & n \leq -(2\ell + 1). \end{cases} \tag{4.5}$$

The same rules apply to $s \neq 0$ with the overrides:

$$s = -2: a_n = O(\bar{\omega}^{|n|+2}), \quad n = -\ell + 1 \text{ and } n = -\ell, \tag{4.6}$$

$$s = -1: a_n = O(\bar{\omega}^{|n|+2}), \quad n = -\ell, \tag{4.7}$$

$$s = +1: a_n = O(\bar{\omega}^{|n|-1}), \quad n = -\ell - 1, \tag{4.8}$$

$$s = +2: a_n = O(\bar{\omega}^{|n|-1}), \quad n = -\ell - 1 \text{ and } n = -\ell - 2. \tag{4.9}$$

Inserting this revised ansatz into Eq. (3.5), together with an expansion for ν , yields equations that can be solved

recursively along rising and falling diagonals (treating ν as if it were a_0) to very high order. Through the diagonal nature of this procedure, it becomes clear in all cases that the general terms given above work for a_n through the following orders while beyond this they must be supplemented by expansions based on the corrected ansatz:

$$\text{General expressions of Eq.(4.3) valid for } s \leq 0 \text{ to order } \begin{cases} O(\bar{\omega}^{n+2\ell+1}), & n \geq -\ell, \\ O(\bar{\omega}^{-n}), & -(2\ell + 1) < n < -\ell, \\ O(\bar{\omega}^{-n-2}), & n \leq -(2\ell + 1). \end{cases}$$

We may usefully turn this around, if we wish to work uniformly to order $\bar{\omega}^N$, the most anomalous behavior occurs at either $n = -\ell / -\ell - 1$ or at $n = -(2\ell + 1)$ and we may use the general expansions except when $\min(\ell + 1, 2\ell - 1) < N$. For example, the expansions to order $\bar{\omega}^4$ of Eq. (4.3) are valid for any $\ell > 2$ while we need to calculate the expansions for $\ell \leq 2$ using the revised, correct ansatz to order $\bar{\omega}^4$:

$$\begin{aligned} s=0: \ell=0 \quad & a_4 = \frac{1}{525}\bar{\omega}^4, \quad a_3 = \frac{1}{60}i\bar{\omega}^3 - \frac{11}{360}\bar{\omega}^4, \quad a_2 = -\frac{1}{9}\bar{\omega}^2 - \frac{1}{6}i\bar{\omega}^3 - \frac{943}{4536}\bar{\omega}^4, \\ & a_1 = -\frac{1}{2}i\bar{\omega} + \frac{1}{2}\bar{\omega}^2 - \frac{331}{360}i\bar{\omega}^3 + \frac{541}{360}\bar{\omega}^4, \quad \nu = -\frac{7}{6}\bar{\omega}^2 - \frac{9449}{7560}\bar{\omega}^4, \\ & a_{-1} = -\frac{2}{9} + \frac{7}{27}i\bar{\omega} - \frac{1591}{2835}\bar{\omega}^2 + \frac{31723}{34020}i\bar{\omega}^3 - \frac{73956143}{50009400}\bar{\omega}^4, \quad a_{-2} = \frac{1}{9}i\bar{\omega} + \frac{1}{54}\bar{\omega}^2 + \frac{853}{3780}i\bar{\omega}^3 + \frac{9679}{34020}\bar{\omega}^4, \\ & a_{-3} = \frac{2}{81}\bar{\omega}^2 + \frac{2}{243}i\bar{\omega}^3 + \frac{1243\bar{\omega}^4}{34020}, \quad a_{-4} = -\frac{1}{270}i\bar{\omega}^3 + \frac{1}{405}\bar{\omega}^4, \quad a_{-5} = -\frac{2}{4725}\bar{\omega}^4. \\ \ell=1 \quad & a_4 = \frac{1}{1323}\bar{\omega}^4, \quad a_3 = \frac{4}{525}i\bar{\omega}^3 - \frac{13}{1575}\bar{\omega}^4, \quad a_2 = -\frac{3}{50}\bar{\omega}^2 - \frac{1}{20}i\bar{\omega}^3 - \frac{1741}{67500}\bar{\omega}^4, \\ & a_1 = -\frac{1}{3}i\bar{\omega} + \frac{1}{6}\bar{\omega}^2 - \frac{2447}{18900}i\bar{\omega}^3 + \frac{2221}{18900}\bar{\omega}^4, \quad \nu = 1 - \frac{19}{30}\bar{\omega}^2 - \frac{1325203}{3591000}\bar{\omega}^4, \\ & a_{-1} = -\frac{1}{6}i\bar{\omega} - \frac{1}{6}\bar{\omega}^2 - \frac{169}{20520}i\bar{\omega}^3 - \frac{467}{4104}\bar{\omega}^4, \quad a_{-2} = -\frac{545}{4332}i\bar{\omega}^3 - \frac{5341}{25992}\bar{\omega}^4, \\ & a_{-3} = -\frac{25}{722}\bar{\omega}^2 + \frac{5}{228}i\bar{\omega}^3 - \frac{5070221}{32840892}\bar{\omega}^4, \quad a_{-4} = \frac{25}{2166}i\bar{\omega}^3 + \frac{5}{3249}\bar{\omega}^4, \quad a_{-5} = \frac{3}{1444}\bar{\omega}^4. \\ \ell=2 \quad & a_4 = \frac{5}{9702}\bar{\omega}^4, \quad a_3 = \frac{5}{882}i\bar{\omega}^3 - \frac{47}{10584}\bar{\omega}^4, \quad a_2 = -\frac{12}{245}\bar{\omega}^2 - \frac{1}{35}i\bar{\omega}^3 - \frac{87826}{9904125}\bar{\omega}^4, \\ & a_1 = -\frac{3}{10}i\bar{\omega} + \frac{1}{10}\bar{\omega}^2 - \frac{21793}{441000}i\bar{\omega}^3 + \frac{38383}{1323000}\bar{\omega}^4, \quad \nu = 2 - \frac{79}{210}\bar{\omega}^2 - \frac{708247}{9261000}\bar{\omega}^4, \\ & a_{-1} = -\frac{1}{5}i\bar{\omega} - \frac{1}{10}\bar{\omega}^2 - \frac{1613}{31500}i\bar{\omega}^3 - \frac{1399}{31500}\bar{\omega}^4, \quad a_{-2} = -\frac{1}{90}\bar{\omega}^2 + \frac{1}{60}i\bar{\omega}^3 - \frac{5104}{5599125}\bar{\omega}^4, \\ & a_{-3} = -\frac{7}{1422}\bar{\omega}^4, \quad a_{-4} = 0, \quad a_{-5} = -\frac{49}{112338}\bar{\omega}^4. \\ \ell=3 \quad & a_4 = \frac{35}{84942}\bar{\omega}^4, \quad a_3 = \frac{10}{2079}i\bar{\omega}^3 - \frac{37}{12474}\bar{\omega}^4, \quad a_2 = -\frac{25}{567}\bar{\omega}^2 - \frac{5}{252}i\bar{\omega}^3 - \frac{2323091}{468087984}\bar{\omega}^4, \\ & a_1 = -\frac{2}{7}i\bar{\omega} + \frac{1}{14}\bar{\omega}^2 - \frac{351521}{12224520}i\bar{\omega}^3 + \frac{117151}{9779616}\bar{\omega}^4, \quad \nu = 3 - \frac{169}{630}\bar{\omega}^2 - \frac{74380421}{2750517000}\bar{\omega}^4, \\ & a_{-1} = -\frac{3}{14}i\bar{\omega} - \frac{1}{14}\bar{\omega}^2 - \frac{6019}{205800}i\bar{\omega}^3 - \frac{29887}{1852200}\bar{\omega}^4, \quad a_{-2} = -\frac{3}{175}\bar{\omega}^2 + \frac{1}{70}i\bar{\omega}^3 - \frac{15979}{5145000}\bar{\omega}^4, \\ & a_{-3} = \frac{1}{2100}i\bar{\omega}^3 + \frac{11}{12600}\bar{\omega}^4, \quad a_{-4} = 0, \quad a_{-5} = 0. \end{aligned}$$

$$\begin{aligned}
 s = -1: \ell = 1 \quad & a_4 = \frac{1}{147}\bar{\omega}^4, \quad a_3 = \frac{1}{21}i\bar{\omega}^3 - \frac{13}{252}\bar{\omega}^4, \quad a_2 = -\frac{6}{25}\bar{\omega}^2 - \frac{1}{5}i\bar{\omega}^3 - \frac{463}{2500}\bar{\omega}^4, \\
 & a_1 = -\frac{3}{4}i\bar{\omega} + \frac{3}{8}\bar{\omega}^2 - \frac{14977}{33600}i\bar{\omega}^3 + \frac{24847}{67200}\bar{\omega}^4, \quad \nu = 1 - \frac{47}{60}\bar{\omega}^2 - \frac{43908007}{71064000}\bar{\omega}^4, \\
 & a_{-1} = -\frac{17}{282}i\bar{\omega}^3 - \frac{17}{282}\bar{\omega}^4, \quad a_{-2} = -\frac{5}{47}i\bar{\omega}^3 - \frac{107}{564}\bar{\omega}^4, \quad a_{-3} = -\frac{200}{2209}\bar{\omega}^2 + \frac{10}{141}i\bar{\omega}^3 - \frac{152936921}{307419903}\bar{\omega}^4, \\
 & a_{-4} = \frac{150}{2209}i\bar{\omega}^3 + \frac{85}{4418}\bar{\omega}^4, \quad a_{-5} = \frac{48}{2209}\bar{\omega}^4.
 \end{aligned}$$

$$\begin{aligned}
 \ell = 2 \quad & a_4 = \frac{5}{1782}\bar{\omega}^4, \quad a_3 = \frac{10}{411}i\bar{\omega}^3 - \frac{47}{2646}\bar{\omega}^4, \quad a_2 = -\frac{20}{147}\bar{\omega}^2 - \frac{5}{63}i\bar{\omega}^3 - \frac{440941}{14261940}\bar{\omega}^4, \\
 & a_1 = -\frac{8}{15}i\bar{\omega} + \frac{8}{45}\bar{\omega}^2 - \frac{32987}{330750}i\bar{\omega}^3 + \frac{56647}{992250}\bar{\omega}^4, \quad \nu = 2 - \frac{169}{420}\bar{\omega}^2 - \frac{6832249}{74088000}\bar{\omega}^4, \\
 & a_{-1} = -\frac{1}{20}i\bar{\omega} - \frac{1}{40}\bar{\omega}^2 - \frac{5707}{168000}i\bar{\omega}^3 - \frac{7397}{336000}\bar{\omega}^4, \quad a_{-2} = -\frac{11}{6760}\bar{\omega}^4, \quad a_{-3} = -\frac{7}{6084}\bar{\omega}^4, \quad a_{-4} = 0, \\
 & a_{-5} = -\frac{49}{57122}\bar{\omega}^4.
 \end{aligned}$$

$$\begin{aligned}
 s = -2: \ell = 2 \quad & a_4 = \frac{10}{891}\bar{\omega}^4, \quad a_3 = \frac{5}{72}i\bar{\omega}^3 - \frac{47}{864}\bar{\omega}^4, \quad a_2 = -\frac{15}{49}\bar{\omega}^2 - \frac{5}{28}i\bar{\omega}^3 - \frac{730781}{6338640}\bar{\omega}^4, \\
 & a_1 = -\frac{5}{6}i\bar{\omega} + \frac{5}{18}\bar{\omega}^2 - \frac{12029}{52920}i\bar{\omega}^3 + \frac{19519}{158760}\bar{\omega}^4, \quad \nu = 2 - \frac{107}{210}\bar{\omega}^2 - \frac{1695233}{9261000}\bar{\omega}^4, \quad a_{-1} = -\frac{1}{20}i\bar{\omega}^3 - \frac{1}{40}\bar{\omega}^4, \\
 & a_{-2} = \frac{11}{12840}\bar{\omega}^4, \quad a_{-3} = -\frac{7}{1926}\bar{\omega}^4, \quad a_{-4} = 0, \quad a_{-5} = -\frac{98}{11449}\bar{\omega}^4. \tag{4.10}
 \end{aligned}$$

In order to evaluate $q^T(\sigma)$ via Eqs. (5.10) and (5.11) we also need the sum of the a_n . Using Eq. (4.3) we obtain, to order $\bar{\omega}^4$,

$$\begin{aligned}
 \sum_{n=-\infty}^{\infty} a_n = & 1 + \frac{(\lambda + s^2)}{2\lambda}i\bar{\omega} - \frac{\lambda^2(4\lambda^2 - 7\lambda + 4) + \lambda(4\lambda + 9)s^4 + 6\lambda(4\lambda - 3)s^3 + 3(2\lambda - 1)(4\lambda^2 + \lambda - 6)s^2 + 2\lambda(4\lambda - 3)(7\lambda - 6)s}{4\lambda^2(4\lambda - 3)^2}\bar{\omega}^2 \\
 & + \frac{P_3}{24(3 - 4\lambda)^2(\lambda - 2)\lambda^3}i\bar{\omega}^3 + \frac{P_4}{48(\lambda - 2)^2\lambda^4(4\lambda - 15)^2(4\lambda - 3)^4}\bar{\omega}^4 + O(\bar{\omega}^5), \tag{4.11}
 \end{aligned}$$

where

$$\begin{aligned}
 P_3 = & (\lambda - 2)\lambda^2(4\lambda^3 + 213\lambda^2 - 309\lambda + 108) + i(4\lambda^3 + 69\lambda^2 - 81\lambda + 54)s^6 \\
 & - 18i(\lambda - 2)\lambda(4\lambda - 3)s^5 - i\lambda(36\lambda^3 - 175\lambda^2 + 432\lambda - 252)s^4 - 12i(\lambda - 2)\lambda(4\lambda - 3)(5\lambda - 3)s^3 \\
 & - i\lambda(36\lambda^4 - 279\lambda^3 + 686\lambda^2 - 738\lambda + 288)s^2 - 6i(\lambda - 2)\lambda^2(4\lambda - 3)(7\lambda - 6)s, \tag{4.12}
 \end{aligned}$$

$$\begin{aligned}
P_4 = & -(\lambda-2)^2\lambda^2(448\lambda^8 + 13584\lambda^7 - 68220\lambda^6 - 372693\lambda^5 + 2636097\lambda^4 - 5229081\lambda^3 + 4967622\lambda^2 - 2347380\lambda + 437400) \\
& - (\lambda-2)\lambda(448\lambda^6 + 6288\lambda^5 + 19620\lambda^4 - 334989\lambda^3 + 958311\lambda^2 - 841995\lambda + 109350)s^8 \\
& + 18(\lambda-2)^2\lambda(4\lambda-15)(4\lambda-3)^2(4\lambda^2-102\lambda+45)s^7 \\
& + (-256\lambda^9 - 24896\lambda^8 + 45680\lambda^7 + 1948740\lambda^6 - 12101346\lambda^5 + 28846314\lambda^4 \\
& - 34305849\lambda^3 + 21625542\lambda^2 - 7333740\lambda + 1312200)s^6 \\
& + 6(\lambda-2)\lambda(4\lambda-15)(4\lambda-3)^2(100\lambda^4 - 1194\lambda^3 + 4425\lambda^2 - 3744\lambda + 540)s^5 \\
& + (9600\lambda^{10} - 142432\lambda^9 + 532648\lambda^8 + 2117966\lambda^7 - 23736786\lambda^6 + 79670331\lambda^5 \\
& - 130327029\lambda^4 + 113471766\lambda^3 - 54566136\lambda^2 + 14346720\lambda - 1749600)s^4 \\
& + 2(\lambda-2)\lambda^2(4\lambda-15)(4\lambda-3)^2(540\lambda^4 - 6334\lambda^3 + 25641\lambda^2 - 40896\lambda + 15660)s^3 \\
& - \lambda(256\lambda^{10} + 57152\lambda^9 - 505072\lambda^8 + 7324\lambda^7 + 14370666\lambda^6 - 68251182\lambda^5 \\
& + 151062669\lambda^4 - 183278700\lambda^3 + 123738192\lambda^2 - 43759440\lambda + 6415200)s^2 \\
& + 2(\lambda-2)\lambda^2(4\lambda-15)(4\lambda-3)^2(84\lambda^5 - 3342\lambda^4 + 21295\lambda^3 - 47598\lambda^2 + 42048\lambda - 11880)s,
\end{aligned} \tag{4.13}$$

with special cases

$$\begin{aligned}
s = 0: \ell = 0 \quad & \sum_{n=-\infty}^{\infty} a_n = \frac{7}{9} - \frac{7}{54}i\bar{\omega} - \frac{122}{945}\bar{\omega}^2 + \frac{6343}{68040}i\bar{\omega}^3 + \frac{276257}{2500470}\bar{\omega}^4 + O(\bar{\omega}^5), \\
\ell = 1 \quad & \sum_{n=-\infty}^{\infty} a_n = 1 - \frac{1}{2}i\bar{\omega} - \frac{854}{9025}\bar{\omega}^2 - \frac{19669}{72200}i\bar{\omega}^3 - \frac{3079726441}{7982161250}\bar{\omega}^4 + O(\bar{\omega}^5), \\
\ell = 2 \quad & \sum_{n=-\infty}^{\infty} a_n = 1 - \frac{1}{2}i\bar{\omega} - \frac{53}{882}\bar{\omega}^2 - \frac{377}{3528}i\bar{\omega}^3 - \frac{167323193}{4855023684}\bar{\omega}^4 + O(\bar{\omega}^5), \\
s = -1: \ell = 1 \quad & \sum_{n=-\infty}^{\infty} a_n = 1 - \frac{3}{4}i\bar{\omega} + \frac{19643}{441800}\bar{\omega}^2 - \frac{2212413}{3534400}i\bar{\omega}^3 - \frac{10841072759029}{19128349520000}\bar{\omega}^4 + O(\bar{\omega}^5), \\
\ell = 2 \quad & \sum_{n=-\infty}^{\infty} a_n = 1 - \frac{7}{12}i\bar{\omega} + \frac{59}{3528}\bar{\omega}^2 - \frac{329}{1728}i\bar{\omega}^3 - \frac{3421065415}{236995065216}\bar{\omega}^4 + O(\bar{\omega}^5), \\
s = -2: \ell = 2 \quad & \sum_{n=-\infty}^{\infty} a_n = 1 - \frac{5}{6}i\bar{\omega} - \frac{25}{882}\bar{\omega}^2 - \frac{2045}{5292}i\bar{\omega}^3 - \frac{1706608417}{23750538336}\bar{\omega}^4 + O(\bar{\omega}^5).
\end{aligned}$$

Finally we should note that the power series for ν cannot be valid $\forall \bar{\omega} \in \mathbb{C}$. The reason is that, for example for $\bar{\omega} \in \mathbb{R}$, the exact value of ν is known to be real for small values of $\bar{\omega}$ but, as $\bar{\omega}$ increases, ν reaches some half-integer value and then it suddenly picks up an imaginary part. For real frequency, this behavior can be seen in, e.g., Table 1 in ST, and estimates for the frequency where it happens have been given in [46,47]; we have found a similar behavior for $\bar{\omega}$ on the negative imaginary axis. The power series for ν , however, is purely real for $\bar{\omega}$ real and so it cannot reproduce this behavior. In this paper, however, we are only interested in the small- $\bar{\omega}$ behavior, where the power series does converge. We have been referring to such a regime as the ‘‘perturbative (small-frequency) regime.’’

V. LOW-FREQUENCY EXPANSION OF THE BRANCH CUT INTEGRAND

It is clear from Eqs. (2.17) and (2.19) that the late-time asymptotics of the BC contribution to the GF is provided by the small- $\bar{\sigma}$ expansion of the modes $\delta_s G_\ell$. The small- $\bar{\sigma}$ expansion of $\delta_s G_\ell$, in its turn, is given by the small- $\bar{\sigma}$ asymptotics of the radius-independent quantity $q/|W|^2$ and the radial solution f_ℓ . In this section we will derive the small- $\bar{\sigma}$ asymptotics of $q/|W|^2$. We will use these asymptotics later in order to prove in Eq. (7.4) that ${}_s G_\ell^{\text{BC}} \sim t^{-2\ell-3}$, to leading order for large- t , for general integer spin.

Specifically, in the following subsections we provide explicit expansions for small- $\bar{\omega}$ up to the first three leading powers (which actually correspond to the first four leading orders, since the third order has the same power of $\bar{\omega}$ as the fourth order but, as we shall show, it contains a logarithm in

$\bar{\omega}$) for the various perturbation quantities (except for the a_n and ν , which we gave in the previous section, and for the radial functions, which we give in the next section). We provide these expansions for general integer-spin s and multipole number ℓ . These expansions will yield the first three leading powers (so four leading orders) for late times of the multipole- ℓ GF and field perturbations. The above comments apply to the RW GF and field perturbations but a similar argument applies to the BPT ones. In fact, we shall give the small- $\bar{\omega}$ expressions explicitly for BPT quantities; one can then readily find the corresponding expansions for the RW quantities via the transformations given in Sec. II C.

From now on and for the rest of the paper we shall restrict ourselves to the case that the BPT spin is a negative integer, $s = 0, -1, -2$ (RW spin, of course, is indistinctively positive or negative). BPT quantities for positive spin can be obtained from those for negative spin via the Teukolsky-Starobinskiĭ identities [36,48].

Also, as mentioned, from now on we will focus on small-frequency expansions up to the first three leading powers of the frequency. It is easy to see from the results in the previous section [e.g., compare the general- ℓ and $-s$ expressions in Eq. (4.3) with the specific-mode expressions in Eq. (4.10)] that the expressions that we shall obtain for general

multipole- ℓ and spin- s in principle are not necessarily valid, up to the first three leading powers of the frequency, for the three specific modes $\ell = 0$ (and $s = 0$) and $\ell = 1$ (and $s = 0, -1$). Indeed, some general- s and $-\ell$ expressions that we shall give appear to have singularities at the s and ℓ values for these modes. We have dealt with these three modes separately by carrying out small-frequency expansions after setting the corresponding values of s and ℓ right from the start. Remarkably, we have found that our general- s and $-\ell$ expansions up to the first three leading powers of the frequency, for q^T , ${}_s\hat{R}_\ell^{\text{in,inc}}$ and all subsequent quantities derived from these two quantities, actually give the correct result for two of these anomalous modes, namely for $\ell = 1$, with $s = 0$ and -1 .⁷ For the mode $s = \ell = 0$, the general- s and $-\ell$ expressions do not give the correct result and we present the results for this mode (as well as for $\ell = -s = 1$ and 2, for completeness) in Sec. V E.

A. K_ν

The quantity K_ν introduced in Eq. (3.31) is needed in order to obtain the Wronskian below [in Eq. (5.4)]. We obtain the following small- $\bar{\omega}$ expansion of K_ν from Eq. (3.32):

$$K_\nu = \frac{2^\ell \Gamma(\ell + \frac{1}{2}) \Gamma(2\ell + 2) \Gamma(1 - s)}{\sqrt{\pi} \Gamma^2(\ell - s + 1)} \bar{\omega}^{-\ell+s} + \frac{i 2^{\ell+1} \Gamma(2\ell) \Gamma(\ell + \frac{3}{2}) \Gamma(1 - s) (2(\ell + 1) \ell (H_\ell - 2H_{-\ell} + \gamma_E) + \ell^2 + \ell - s^2)}{\sqrt{\pi} (\ell + 1) \Gamma^2(\ell - s + 1)} \\ \times \bar{\omega}^{-\ell+s+1} + \frac{2^{-\ell-3} \Gamma(2\ell + 1) \Gamma(2\ell + 2) \Gamma(1 - s)}{\Gamma(\ell + 1) \Gamma(\ell - s + 1)^2} K_\nu^{(3)} \bar{\omega}^{-\ell+s+2} + O(\bar{\omega}^{-\ell+s+3}), \quad (5.1)$$

where

$$K_\nu^{(3)} \equiv \frac{4(6(\ell^2 + \ell - 1)s^2 + \ell(\ell + 1)(15\ell^2 + 15\ell - 11) + 3s^4)(2H_{\ell-s} + H_\ell - 2H_{2\ell} - 2H_{2\ell+1} + \ln(2\bar{\omega}) + \gamma_E)}{\ell(\ell + 1)(2\ell - 1)(2\ell + 1)(2\ell + 3)} \\ - 4 \left(-\frac{s^2}{\ell(\ell + 1)} + 2\gamma_E + 1 \right) (H_\ell - 2H_{-\ell} + \gamma_E) - 4(H_\ell - 2H_{-\ell})^2 - 4 \left(2H_{\ell-s}^{(2)} + H_\ell^{(2)} - 4H_{-\ell}^{(2)} + \frac{\pi^2}{6} - \gamma_E^2 \right) \\ - \frac{1}{\ell^2(\ell + 1)^2(2\ell - 1)^2(2\ell + 3)^2} \{ 2((\ell + 1)(8\ell^3 - 16\ell^2 - 24\ell + 9)s^4 + \ell^2(\ell + 1)^2(8\ell^4 + 16\ell^3 - 4\ell^2 - 13\ell + 4) \\ - (16\ell^6 + 104\ell^5 + 164\ell^4 - 16\ell^3 - 131\ell^2 - 3\ell + 18)s^2) \}. \quad (5.2)$$

Doing similarly for $K_{-\nu-1}$, we obtain

$$K_{-\nu-1} = \frac{(-1)^\ell i 2^{\ell-1} \Gamma(1 - s) \Gamma(\ell + 1) \Gamma^2(\ell + s + 1)}{\Gamma(2\ell + 1) \Gamma(2\ell + 2) \nu_2^2} \bar{\omega}^{s+\ell} + O(\bar{\omega}^{\ell+s-1}). \quad (5.3)$$

In obtaining Eq. (5.3), we have used Eq. (4.4) together with the fact that $\Gamma(-n + \bar{\omega}) \sim \frac{(-1)^n}{\Gamma(n+1)\bar{\omega}}$ as $\bar{\omega} \rightarrow 0$ when n is a nonnegative integer.

We note that the asymptotics of K_ν in Eq. (5.1) and those of $K_{-\nu-1}$ in Eq. (5.3) imply, via Eq. (168) ST (and since $\ell \geq |s|$), that $K_{-\nu-1}$ is not necessary for obtaining ${}_s\hat{R}_\ell^{\text{in,inc}}$ to the first four leading orders ($K_{-\nu-1}$ starts playing a part only in

⁷Specifically, for, e.g., $s = 0$ and $\ell = 1$, the general- s and $-\ell$ expression for K_ν gives the wrong coefficient in the third leading power of $\bar{\omega}$ and for $K_{-\nu-1}$ it gives the wrong leading order. However, for this mode, the two incorrections in K_ν and in $K_{-\nu-1}$ somehow miraculously cancel each other out to give the right result for ${}_s\hat{R}_\ell^{\text{in,inc}}$ up to the first three leading powers of $\bar{\omega}$.

the next order), except in the cases $\ell = 0$ and $\ell = 1$. As mentioned at the start of this section, though, Eqs. (5.1) and (5.3) are in principle not valid for $\ell = 0$ and 1, although the general- s and $-\ell$ expressions that we give for quantities from now on are, somewhat surprisingly, also valid for $\ell = 1$ (with $s = 0, 1$). The case $s = \ell = 0$ we treat separately in Sec. V E 1.

B. Radial coefficients and Wronskian

We now turn to the coefficients in Eq. (2.32) of the ingoing radial (BPT) solution. We have obtained the following expansion for the radial coefficient ${}_s\hat{R}_\ell^{\text{in,inc}}$ by carrying out small- $\bar{\sigma}$ asymptotics of $R_s^{\text{in,inc}}/R_s^{\text{in,tra}}$, where $R_s^{\text{in,inc/tra}}$ are the incidence/transmission coefficients of R_s^{in} (i.e., with the specific normalization used in Sec. III, which is the same normalization as in MST). The expression for the coefficient $R_s^{\text{in,tra}}$ is given in Eq. (3.12) whereas for $R_s^{\text{in,inc}}$ we used Eq. (168) ST (which requires K_ν and $K_{-\nu-1}$). We obtain

$$\begin{aligned} {}_s\hat{R}_\ell^{\text{in,inc}} = & M^{1-2s} \frac{2^{-\ell-s} e^{\frac{1}{2}i\pi(\ell-s+1)} \Gamma(2\ell+1) \Gamma(2\ell+2) \Gamma(1-s)}{\Gamma(\ell+1) \Gamma(\ell-s+1) \Gamma(\ell+s+1)} \frac{1}{\bar{\omega}^{\ell-s+1}} \\ & \times \left\{ 1 + \left[i \left(H_{\ell-s} + H_{\ell+s} + H_\ell - 2H_{-s} - \ln(2\bar{\omega}) + \frac{i\pi}{2} - \gamma_E \right) - \frac{i(\ell(\ell+1) + s^2)}{2\ell(\ell+1)} \right] \bar{\omega} \right. \\ & + \left[\frac{(6(\ell^2 + \ell - 1)s^2 + \ell(\ell+1)(15\ell^2 + 15\ell - 11) + 3s^4)(H_{\ell-s} + H_{\ell+s} + H_\ell - 2H_{2\ell} - 2H_{2\ell+1} + \ln(2\bar{\omega}) - \frac{i\pi}{2} + \gamma_E)}{2\ell(\ell+1)(2\ell-1)(2\ell+1)(2\ell+3)} \right. \\ & - \frac{1}{2} \left(\frac{s^2}{\ell(\ell+1)} + 1 \right) \left(-H_{\ell-s} - H_{\ell+s} - H_\ell + 2H_{-s} + \ln(2\bar{\omega}) - \frac{i\pi}{2} + \gamma_E \right) \\ & - \frac{1}{2} \left(-H_{\ell-s} - H_{\ell+s} - H_\ell + 2H_{-s} + \ln(2\bar{\omega}) - \frac{i\pi}{2} + \gamma_E \right)^2 + \frac{1}{2} \left(-H_{\ell-s}^{(2)} - H_{\ell+s}^{(2)} - H_\ell^{(2)} + 4H_{-s}^{(2)} - \frac{\pi^2}{6} \right) \\ & - \frac{(8\ell^3 - 16\ell^2 - 24\ell + 9)s^4}{4\ell^2(\ell+1)(2\ell-1)^2(2\ell+3)^2} - \frac{8\ell^4 + 16\ell^3 - 4\ell^2 - 13\ell + 4}{4(2\ell-1)^2(2\ell+3)^2} \\ & \left. \left. - \frac{(16\ell^6 - 8\ell^5 - 116\ell^4 - 48\ell^3 + 101\ell^2 + 21\ell - 18)s^2}{4\ell^2(\ell+1)^2(2\ell-1)^2(2\ell+3)^2} \right] \bar{\omega}^2 \right\} + o\left(\frac{1}{\bar{\omega}^{\ell-s-1}}\right). \end{aligned} \quad (5.4)$$

We note that the leading order of Eq. (5.4) agrees with Eq. (3.6.13) [49], which is obtained via an independent method based on Page's [50]. As a token example of the more simplified form that adopts the expansion for a particular value of s , we give Eq. (5.4) specifically for $s = 0$:

$$\begin{aligned} \frac{1}{M} {}_s\hat{R}_\ell^{\text{in,inc}} = & \frac{2^{-\ell} e^{\frac{1}{2}i\pi(\ell+1)} \Gamma(2\ell+1) \Gamma(2\ell+2)}{\Gamma^3(\ell+1)} \frac{1}{\bar{\omega}^{\ell+1}} \\ & + \frac{i2^{-\ell-1} e^{\frac{1}{2}i\pi(\ell+1)} \Gamma(2\ell+1) \Gamma(2\ell+2) (6\psi(\ell+1) - 2\ln(2\bar{\omega}) + i\pi + 4\gamma_E - 1)}{\Gamma^3(\ell+1)} \frac{1}{\bar{\omega}^\ell} \\ & \times \frac{2^{3\ell} e^{\frac{1}{2}i\pi(\ell+1)} \Gamma(\ell + \frac{1}{2}) \Gamma(\ell + \frac{3}{2})}{\pi \Gamma(\ell+1)} \left((\ln(2i\bar{\omega}) - i\pi - 2\gamma_E) \left(\frac{15\ell^2 + 15\ell - 11}{(2\ell-1)(2\ell+1)(2\ell+3)} + 6\psi(\ell+1) - 1 \right) \right. \\ & + \frac{(15\ell^2 + 15\ell - 11)(3\psi(\ell+1) - 2\psi(2\ell+1) - 2\psi(2\ell+2) + 2\gamma_E)}{(2\ell-1)(2\ell+1)(2\ell+3)} + \frac{2\ell+1}{4(2\ell-1)^2(2\ell+3)^2} \\ & \left. - 9\psi(\ell+1)^2 + 3\psi(\ell+1) + 3\psi^{(1)}(\ell+1) - \left(\ln(2i\bar{\omega}) - i\pi - 2\gamma_E \right)^2 - \frac{2\pi^2}{3} - \frac{1}{4} \right) \frac{1}{\bar{\omega}^{\ell-1}} + \left(\frac{1}{\bar{\omega}^{\ell-2}} \right), \\ & s = 0. \end{aligned} \quad (5.5)$$

Here, the function $\psi(z) \equiv \Gamma'(z)/\Gamma(z)$ is the digamma function [42] and $\psi^{(n)}(z)$ its n th derivative. We note that the digamma function may be expressed as $\psi(\ell) = H_{\ell-1} - \gamma_E$, in terms of the harmonic numbers used above.

C. BC strength q^T

In this section we will provide a small- $\bar{\sigma}$ for the BC strength q^T . The solution R_+^ν of the BPT equation has the following asymptotics, from Eqs. (3.39) and (3.30) (taking the upper sign),

$$\frac{R_+^\nu}{R_+^{\nu,\text{tra}}} \sim \frac{1}{r} e^{-i\omega r_s}, \quad \bar{r} \rightarrow \infty, \quad (5.6)$$

where

$$R_+^{\nu,\text{tra}}(\omega) \equiv A_+^\nu \omega^{-1} e^{-i\bar{\omega} \ln \bar{\omega}}, \quad (5.7)$$

and A_+^ν is given in Eq. (3.41). Comparing with Eq. (2.33), it follows that

$$\frac{R_+^\nu(r, \omega)}{R_+^{\nu,\text{tra}}(\omega)} = {}_{-s}\hat{R}_\ell^{\text{up}}(r, -\omega) \Delta^{-s}. \quad (5.8)$$

Now, from Eq. (3.18) together with the analytic continuation of the irregular hypergeometric U -functions on the complex- ω plane [Eq. (13.2.41) [42]], we find

$$R_s^{\text{up}}(r, \omega e^{2\pi i}) = e^{-2\pi\bar{\omega}} R_s^{\text{up}}(r, \omega) + [e^{-2\pi\bar{\omega}} - e^{-2\pi i\nu}] R_+^\nu(r, \omega). \quad (5.9)$$

Using the definition Eq. (2.36) of the BC strength, it then follows that

$$q^T(\sigma) = i[1 - e^{2\pi(\bar{\omega}-i\nu)}] \frac{R_+^{\nu,\text{tra}}(-i\sigma)}{R_s^{\text{up,tra}}(-i\sigma)}. \quad (5.10)$$

We can therefore calculate the ratio of coefficients on the right-hand side via

$$\frac{R_+^{\nu,\text{tra}}(\omega)}{R_s^{\text{up,tra}}(\omega)} = \frac{A_+^\nu}{A_-^\nu} \omega^{2s} \bar{\omega}^{-2i\bar{\omega}}, \quad (5.11)$$

where we have made use of Eqs. (5.6) and (3.20). From Eqs. (5.10) and (5.11) and expanding Eq. (3.41) for A_+^ν and Eq. (3.19) for A_-^ν , we find the first five leading orders for the BC strength:

$$\begin{aligned} q^T(\sigma) = & \frac{\bar{\sigma}^{2s+1} 2\pi(-1)^{\ell+s} \Gamma(\ell-s+1)}{M^{2s} \Gamma(\ell+s+1)} \left\{ 1 + \left[H_{\ell-s} + H_{\ell+s} + \frac{3(\ell^2 + \ell - 1)s^2}{\ell(\ell+1)(2\ell-1)(2\ell+1)(2\ell+3)} \right. \right. \\ & + \frac{16\ell^3 + 39\ell^2 + 11\ell - 17}{2(2\ell-1)(2\ell+1)(2\ell+3)} + \frac{3s^4}{2\ell(\ell+1)(2\ell-1)(2\ell+1)(2\ell+3)} - 2\ln(2\bar{\sigma}) - 2\gamma_E \Big] \bar{\sigma} \\ & + \left[\left(-2H_{\ell-s} - 2H_{\ell+s} - \frac{6(\ell^2 + \ell - 1)s^2 + \ell(\ell+1)(16\ell^3 + 39\ell^2 + 11\ell - 17) + 3s^4}{\ell(\ell+1)(2\ell-1)(2\ell+1)(2\ell+3)} + 4\gamma_E \right) \left(\gamma_E - H_{\ell-s} \right. \right. \\ & + \ln(2\bar{\sigma}) - \frac{1}{2} \Big) - 2 \left(\gamma_E - H_{\ell-s} - \frac{1}{2} \right)^2 + \frac{1}{2} (H_{\ell+s} - H_{\ell-s})^2 + \frac{1}{2} (H_{\ell+s}^{(2)} - H_{\ell-s}^{(2)}) \\ & \left. \left. + \frac{s(7\ell^2 + 7\ell + 3s^2 - 6)}{2\ell(\ell+1)(2\ell-1)(2\ell+3)} + 2\ln^2(2\bar{\sigma}) - \frac{\pi^2}{6} \right] \bar{\sigma}^2 \right\} + o(\bar{\sigma})^{2s+3}. \quad (5.12) \end{aligned}$$

Using Eq. (2.44) to relate BPT's q^T with RW's q , it is easy to check that the leading order in Eq. (5.12) agrees with Eq. (41) [27] (the “ iq ” here being “ K ” in [27]) for all spins $s = 0, -1, -2$ (the leading order of RW's q is independent of the spin).

Again, as a token example of the more simplified form for a particular value of s , we give Eq. (5.12) specifically for $s = 0$:

$$\begin{aligned} q^T(\sigma) = q(\sigma) = & i\pi(-1)^\ell \bar{\omega} \left\{ 2 + \left(\frac{i(16\ell^3 + 39\ell^2 + 11\ell - 17)}{8\ell^3 + 12\ell^2 - 2\ell - 3} + 4i\psi(\ell+1) - 4i\ln(2\bar{\sigma}) \right) \bar{\omega} \right. \\ & - \left(\frac{2(16\ell^3 + 39\ell^2 + 11\ell - 17)(\psi(\ell+1) - \ln(2\bar{\sigma}))}{(2\ell-1)(2\ell+1)(2\ell+3)} + \frac{8\ell^3 + 27\ell^2 + 13\ell - 14}{(2\ell-1)(2\ell+1)(2\ell+3)} \right. \\ & \left. \left. + 4 \left(\psi(\ell+1) - \ln(2\bar{\sigma}) \right)^2 - \frac{\pi^2}{3} \right) \bar{\omega}^2 \right\} + o(\bar{\omega}^3), \\ s = & 0. \quad (5.13) \end{aligned}$$

D. $q^T/|W^T|^2$

We are finally in a position to give an expansion for the main target of this section: the radius-independent quantity in the BC integrand Eq. (2.19). From Eqs. (5.4) and (5.12) it follows that

$$\begin{aligned} \frac{q^T(\sigma)}{|W^T|^2} = & M^{2s} \frac{(-1)^{\ell+s} \pi^2 2^{-2\ell+2s+1} \Gamma^3(\ell-s+1) \Gamma(\ell+s+1)}{\Gamma^2(\ell+\frac{1}{2}) \Gamma^2(2\ell+2) \Gamma^2(1-s)} \left[\bar{\sigma}^{2\ell+1} - \frac{\bar{\sigma}^{2\ell+2}}{2} \left(8H_{-s} - 2H_{\ell-s} - 2H_{\ell+s} - 4H_\ell \right. \right. \\ & + \frac{2(\ell+1)(\ell(8\ell+7)-6)s^2 + \ell(\ell+1)(\ell(32\ell+63)+7)-23}{\ell(\ell+1)(2\ell-1)(2\ell+1)(2\ell+3)} + 3s^4 \\ & \left. \left. + \frac{2(l+1)(8l^2+7l-6)s^2 + l(l+1)(32l^3+63l^2+7l-23)+3s^4}{l(l+1)(2l-1)(2l+1)(2l+3)} \right) - Q\bar{\sigma}^{2\ell+3} \right] + o(\bar{\sigma}^{2\ell+3}), \end{aligned} \quad (5.14)$$

where

$$\begin{aligned} Q \equiv & \frac{(-6(\ell^2+\ell-1)s^2 + \ell(\ell+1)(11-15\ell(\ell+1)) - 3s^4)(H_{\ell-s} - 4H_{2\ell} + 2H_{-s} + \ln(2\bar{\sigma}) + \gamma_E)}{\ell(\ell+1)(2\ell-1)(2\ell+1)(2\ell+3)} \\ & - 4H_{-s} \left(2H_{-s} + \frac{s^2}{\ell(\ell+1)} + 2 \right) + \left(-2H_\ell + 4H_{-s} + \frac{s^2}{\ell(\ell+1)} + 2 \right) (H_{\ell-s} + H_{\ell+s} + 2H_\ell) - \frac{1}{2} (H_{\ell-s} + H_{\ell+s})^2 \\ & + 2(H_\ell)^2 + \frac{3H_{\ell-s}^{(2)}}{2} + \frac{H_{\ell+s}^{(2)}}{2} + H_\ell^{(2)} - 4H_{-s}^{(2)} - \frac{3s^6}{2\ell^2(\ell+1)^2(2\ell-1)(2\ell+1)(2\ell+3)} - \frac{(7\ell^2+7\ell-6)s}{2\ell(\ell+1)(2\ell-1)(2\ell+3)} \\ & - \frac{(128\ell^6 + 768\ell^5 + 1088\ell^4 + 120\ell^3 - 364\ell^2 + 18\ell + 45)s^4}{4\ell^2(\ell+1)^2(2\ell-1)^2(2\ell+1)^2(2\ell+3)^2} \\ & + \frac{-512\ell^6 - 2016\ell^5 - 1616\ell^4 + 1472\ell^3 + 1128\ell^2 - 722\ell + 59}{4(2\ell-1)^2(2\ell+1)^2(2\ell+3)^2} \\ & - \frac{(256\ell^8 + 1464\ell^7 + 2660\ell^6 + 1082\ell^5 - 1361\ell^4 - 926\ell^3 + 122\ell^2 + 93\ell + 18)s^2}{2\ell^2(\ell+1)^2(2\ell-1)^2(2\ell+1)^2(2\ell+3)^2} \\ & - \frac{3s^3}{2\ell(\ell+1)(2\ell-1)(2\ell+3)} + \frac{\pi^2}{3}. \end{aligned} \quad (5.15)$$

Here we give the value of $q/|W|^2$ specifically for the case $s=0$. From Eqs. (5.5) and (5.13), or equivalently from Eq. (5.14) with $s=0$ (or, equivalently, from [33]⁸), we have

$$\begin{aligned} 4 \frac{q^T(\sigma)}{|W^T|^2} = & \frac{q(\sigma)}{|W|^2} = \frac{(-1)^\ell \pi}{2^{2\ell-3}} \left(\frac{(2\ell+1)\ell!}{((2\ell+1)!!)^2} \right)^2 \left[\bar{\sigma}^{2\ell+1} - \bar{\sigma}^{2\ell+2} \left(\frac{-32\ell^3 - 63\ell^2 - 7\ell + 23}{2(2\ell+3)(2\ell+1)(2\ell-1)} + 4H_\ell \right) \right] \\ & + \frac{(-1)^\ell \pi}{2^{2\ell-1}} \left(\frac{(2\ell+1)\ell!}{((2\ell+1)!!)^2} \right)^2 \bar{\sigma}^{2\ell+3} \left[\frac{4(15\ell^2 + 15\ell - 11)}{(2\ell-1)(2\ell+1)(2\ell+3)} (\ln(2\bar{\sigma}) + H_\ell - 4H_{2\ell} + \gamma_E) \right. \\ & \left. - 4(-8H_\ell^2 + 8H_\ell + 3H_\ell^{(2)} + 2H_\infty^{(2)}) + \frac{512\ell^6 + 2016\ell^5 + 1616\ell^4 - 1472\ell^3 - 1128\ell^2 + 722\ell - 59}{(2\ell-1)^2(2\ell+1)^2(2\ell+3)^2} \right] \\ & + o(\bar{\sigma}^{2\ell+3}), \end{aligned} \quad (5.16)$$

$s=0,$

where $H_\ell^{(r)}$ is the ℓ th harmonic number of order r .

We note the appearance in (5.14) of a logarithmic behavior in $\bar{\sigma}$ at order $\bar{\sigma}^{2\ell+3}$ for small frequency. It is worth pointing out the following ‘‘curious’’ fact. Even though the logarithmic behavior appears already at second leading order both in the BC strength q^T [see Eq. (5.12)] and in the radial coefficient ${}_s\hat{R}_\ell^{\text{in,inc}}$ [see Eq. (5.4)], there is a delicate cancellation between the terms which leads to the logarithmic behavior for $q^T/|W^T|^2$ appearing *not* at second order but at third leading order instead.

⁸We note a typo of an extra overall ‘‘-1’’ in Eq. (8) [33].

As shown later in Sec. VII, this implies that the logarithmic behavior of the Green function or a field perturbation will also appear at third—as opposed to second, as one might have expected—leading order for late times.

E. Cases $\ell = -s$

As pointed out at the start of this section, the expression in Eq. (5.16) might not be valid for the case $s = \ell = 0$. In this section we present the results obtained by carrying out a specific calculation for this case by setting $s = \ell = 0$ right at the start of the calculation. Although the BPT results for $\ell = -s = 1, 2$ can be obtained directly by putting in these values into the general s and $\ell (> 0)$ expressions found above, we also include these cases for completeness and because they require an extra step in order to obtain the RW results from the BPT results.

1. Case $s = \ell = 0$

We have carried out a specific calculation for the case $s = \ell = 0$ by setting these values right from the start and we have obtained

$$\begin{aligned} \frac{1}{M^s} \hat{R}_\ell^{\text{in,inc}} &= -\frac{1}{\bar{\sigma}} + \ln(2\bar{\sigma}) - i\pi + \gamma_E + \frac{1}{2} \\ &+ \frac{1}{36} \bar{\sigma} (15\pi^2 - 6\ln(2\bar{\sigma})(2\ln(2\bar{\sigma})) \\ &- 6i\pi + 6\gamma_E - 8) + 36i\gamma_E\pi - 48i\pi \\ &+ 6(8 - 3\gamma_E)\gamma_E - 82) + o(\bar{\sigma}). \end{aligned} \quad (5.17)$$

We note that the first two leading orders agree with Eq. (5.5) with $\ell = 0$ but the third leading orders differ slightly in the term which does not contain any “ $\ln(2\bar{\sigma})$.”

From Eq. (5.17) we readily obtain (assuming evaluation on the NIA, i.e., $\nu > 0$)

$$\begin{aligned} |W^T|^2 &= 4\sigma^2 |{}_s \hat{R}_\ell^{\text{in,inc}}|^2 = 1 - (2(\ln(2\bar{\sigma}) + \gamma_E) + 1)\bar{\sigma} \\ &+ \frac{1}{36} (72(\ln(2\bar{\sigma}) + \gamma_E)^2 - 60(\ln(2\bar{\sigma}) + \gamma_E) \\ &+ 6\pi^2 + 173)\bar{\sigma}^2 + o(\bar{\sigma}^2). \end{aligned} \quad (5.18)$$

For the BC strength we obtain

$$\begin{aligned} q^T(\sigma) = q(\sigma) &= 2\pi\bar{\sigma} + \frac{\bar{\sigma}^2}{3} (17\pi - 12\pi(\ln(2\bar{\sigma}) + \gamma_E)) \\ &+ \frac{\pi\bar{\sigma}^3}{3} (12\ln^2(2\bar{\sigma}) + 2(12\gamma_E - 17)\ln(2\bar{\sigma})) \\ &+ 2\gamma_E(6\gamma_E - 17) + 20 - \pi^2) + o(\bar{\sigma}^3). \end{aligned} \quad (5.19)$$

We note that if instead of calculating the BC strength directly for $s = \ell = 0$ we had used Eq. (5.12) [or Eq. (5.13)] with $s = \ell = 0$ we would have obtained the

result in Eq. (5.19) but with the number 14 in the place of the number 20 in the third leading order.

Finally, from Eqs. (5.18) and (5.19) it follows that

$$\begin{aligned} \frac{q^T(\sigma)}{|W^T|^2} &= 2\pi\bar{\sigma} + \frac{23\pi\bar{\sigma}^2}{3} - \frac{\pi\bar{\sigma}^3}{18} (12\pi^2 - 132(\ln(2\bar{\sigma}) \\ &+ \gamma_E) - 85) + o(\bar{\sigma}^4), \quad \ell = s = 0. \end{aligned} \quad (5.20)$$

The third leading order differs from Eq. (5.16) in that the “ -85 ” is a “ $+59$ ” in Eq. (5.16); this is a consequence of the corresponding discrepancies in both $q(\sigma)$ and A_ℓ^{in} .

2. Cases $\ell = -s = 1, 2$

By inserting the values $\ell = -s = 1$ and $\ell = -s = 2$ into Eq. (5.14) we respectively obtain

$$\begin{aligned} 4M^2 \frac{q^T(\sigma)}{|W^T|^2} &= \frac{4\pi}{9} \bar{\sigma}^3 + \frac{227\pi}{135} \bar{\sigma}^4 \\ &+ \frac{\pi(-1200\pi^2 + 5640(\ln(2\bar{\sigma}) + \gamma_E) + 10457)}{8100} \\ &\times \bar{\sigma}^5 + o(\bar{\sigma}^5), \quad \ell = -s = 1, \end{aligned} \quad (5.21)$$

and

$$\begin{aligned} 4M^4 \frac{q^T(\sigma)}{|W^T|^2} &= \frac{\pi}{75} \bar{\sigma}^5 + \frac{191\pi}{3500} \bar{\sigma}^6 \\ &- \frac{\pi(19600\pi^2 - 59920(\ln(2\bar{\sigma}) + \gamma_E) - 348727)}{4410000} \\ &\times \bar{\sigma}^7 + o(\bar{\sigma}^7), \quad \ell = -s = 2. \end{aligned} \quad (5.22)$$

VI. RADIAL SOLUTIONS

In this section we obtain the small-frequency behavior of the upgoing and ingoing radial solutions valid for *arbitrary* radius r . For this purpose, we use a special trick based on the Barnes integral representation of the hypergeometric function. As shown in Sec. II, the radial dependence of the BC contribution to the GF only comes in through the ingoing radial solution, not the upgoing one [see Eqs. (2.21) and (2.37)]. Therefore, the radial dependence at late times of the GF itself also only comes in through the ingoing radial solution. For this reason, we apply the mentioned Barnes trick to give explicitly the first three leading orders of the ingoing radial solution, whereas we only give the leading order (for which the Barnes trick is not necessary) of the upgoing solution. We show, however, how the Barnes trick can be used to obtain the behavior of the ingoing solution up to arbitrary order in the frequency within the perturbative regime and we note that it could be similarly applied to the upgoing solution.

The technique we use for obtaining the small-frequency expansion of the radial solutions can be applied just the same to the RW or to the BPT solutions. Besides, from the expansion for the RW solutions one can readily obtain the expansion for the BPT solutions via the Chandrasekhar transformation of Sec. II C. For this reason, we only give the expansions explicitly for one type of solutions: the BPT solutions for the upgoing modes and the RW solutions for the ingoing modes.

A. Upgoing radial solution

The results in this subsection correspond to the particular normalization of the BPT solutions that is chosen in ST. Taking the small- $\bar{\omega}$ asymptotics of Eq. (4.9) MSTa we obtain

$$R_s^{\text{up}} = \frac{\nu_2 \Gamma(2\ell + 1) \Gamma(2\ell + 2)}{2^\ell \Gamma(\ell + 1) \Gamma^2(\ell + s + 1) \bar{\omega}^{\ell+s-1}} R_0^\nu + o(\bar{\omega}^{-\ell-s+2}), \quad (6.1)$$

where we have used Eqs. (5.1) and (5.3). From Eq. (3.25) for R_0^ν we then obtain the leading-order asymptotics:

$$R_s^{\text{up}} = \frac{\nu_2 \Gamma^2(2\ell + 1) \Gamma(2\ell + 2)}{2^\ell \Gamma^2(\ell + 1) \Gamma^2(\ell + s + 1) \Gamma(\ell - s + 1)} \frac{(\bar{r} - 1)^{-s} {}_2F_1(-\ell, -\ell - s; -2\ell; \frac{1}{\bar{r}})}{\bar{\omega}^{\ell+s+1}} a_0^\top + o(\bar{\omega}^{-\ell-s}). \quad (6.2)$$

B. Ingoing radial solution

In this subsection we consider the ingoing radial solutions of the RW equation. We start with Eqs. (2.16), (3.12), (3.1) and (3.4a),⁹

$$f_\ell(r, \omega) = \frac{X_s^{\text{in}}}{N_s^{\text{in}} \sum_{k=-\infty}^{\infty} a_k \frac{\Gamma(a)\Gamma(b)}{\Gamma(c)}} = \bar{r}^{s+1} \frac{e^{-i\bar{\omega}\bar{r}} (\bar{r} - 1)^{-i\bar{\omega}}}{\sum_{k=-\infty}^{\infty} a_k \frac{\Gamma(a)\Gamma(b)}{\Gamma(c)}} \sum_{k=-\infty}^{\infty} a_k \frac{\Gamma(a)\Gamma(b)}{\Gamma(c)} {}_2F_1(a, b; c; 1 - \bar{r}), \quad (6.3)$$

where

$$a \equiv k + \nu + s + 1 - i\bar{\omega}, \quad b \equiv -k - \nu + s - i\bar{\omega}, \quad c \equiv 1 - 2i\bar{\omega}. \quad (6.4)$$

Let us here carry out a basic comparison of MST's leading order behavior with other results in the literature. We can use the small-frequency asymptotics for the ‘‘in’’ solution of the Teukolsky equation given by Eqs. (3.6.13) and (3.6.15) [49]¹⁰ (see also [50,51]). Setting $a = 0$, we obtain

$$\begin{aligned} {}_s R_\ell^{\text{in}}(r, \omega) &\sim r_h^{-2s} e^{-\bar{\omega}i} {}_s R_\ell^{\text{in,tra}}(\bar{r} - 1)^{-s - \bar{\omega}i} \bar{r}^{-s + \bar{\omega}i} {}_2F_1(-\ell - s, \ell - s + 1, 1 - s - 2\bar{\omega}i; 1 - \bar{r}) \\ &= r_h^{-2s} e^{-\bar{\omega}i} {}_s R_\ell^{\text{in,tra}}(\bar{r} - 1)^{-s - \bar{\omega}i} \bar{r}^{-\bar{\omega}i} {}_2F_1(-\ell - 2i\bar{\omega}, \ell + 1 - 2i\bar{\omega}, 1 - s - 2\bar{\omega}i; 1 - \bar{r}), \\ &(\bar{r} - 1)\bar{\omega} \ll \ell + 1. \end{aligned} \quad (6.5)$$

In order to obtain the RW ingoing solution from BPT's one, we need to apply the Chandrasekhar transformation as per Eq. (2.40). Specifically setting $s = 0$ in Eq. (6.5), and taking into account of the normalizations as per Eqs. (2.11) and (2.32), we obtain

$$f_\ell(r, \omega) \sim e^{-\bar{\omega}i} (\bar{r} - 1)^{-\bar{\omega}i} \bar{r}^{1 + \bar{\omega}i} {}_2F_1(-\ell, \ell + 1, 1 - 2\bar{\omega}i; 1 - \bar{r}), \quad (\bar{r} - 1)\bar{\omega} \ll \ell + 1, \quad s = 0. \quad (6.6)$$

The leading order ($\bar{\omega} = 0$) behavior of Eq. (6.6) clearly agrees with that from Eq. (6.3). Note that the asymptotics of Eq. (6.5) are not amenable to carrying out Fourier frequency integrations. We now proceed to give a prescription for expanding the ingoing RW solution to arbitrary orders in the frequency.

Now that we know from Sec. IV the behavior of the coefficients a_k as $\bar{\omega} \rightarrow 0$ our challenge is to obtain a suitable expansion for the hypergeometric functions in Eq. (6.3). To this end we employ the Barnes integral representation [e.g., Eq. (15.6.6) [42]] which gives

⁹We note that there is a typographical error on the right-hand side of Eq. (6) [33]: there is a factor \bar{r}^{s+1} missing and the j -sum in the denominator is missing $\frac{\Gamma(a)\Gamma(b)}{\Gamma(c)}$ as in Eq. (6.4) but with $k \rightarrow j$.

¹⁰Note that there is a typographical error in Eq. (3.6.15) [49]: a factor $\bar{r}^{-s+i\bar{\omega}}$ is missing on its right hand side.

$$\frac{\Gamma(a)\Gamma(b)}{\Gamma(c)} {}_2F_1(a, b, c, z) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{\Gamma(a+t)\Gamma(b+t)\Gamma(-t)}{\Gamma(c+t)} (-z)^t dt, \quad a, b \neq 0, -1, -2, \dots \quad (6.7)$$

where the path of integration must be chosen to separate the poles of $\Gamma(-t)$ at $t = 0, 1, 2, \dots$ from those of $\Gamma(a+t)$ and $\Gamma(b+t)$, respectively at $t = -a, -a-1, -a-2, \dots$ and $t = -b, -b-1, -b-2, \dots$

For small $\bar{\omega}$ we know from Eqs. (4.1) and (4.2) that

$$\nu \equiv \ell + \delta\nu = \ell - \nu_2 \bar{\omega}^2 + O(\bar{\omega}^3),$$

where ν_2 is positive so that the corresponding poles lie at unit intervals left from

$$\begin{aligned} -k - \ell - s - 1 + i\bar{\omega} + \nu_2 \bar{\omega}^2 + O(\bar{\omega}^3), \\ k + \ell - s + i\bar{\omega} - \nu_2 \bar{\omega}^2 + O(\bar{\omega}^3). \end{aligned}$$

Note that when $\omega = 0$ we have double poles and it is not possible to find a splitting contour and the representation breaks down.

In Fig. 1 we illustrate the poles and contour C_1 of the Barnes integral representation for the term of the MST series Eq. (6.3) for $k = 1, \ell = 2, s = 0, \bar{\omega} = -0.1i$. The proximity of the poles for small $\bar{\omega}$ leads to threading contour C_1 complicates evaluation of the integral however we may overcome this simply by deforming the contour to the contour C_2 lying to the right of all the poles of $\Gamma(a+t)$ and $\Gamma(b+t)$ collecting the residues from the poles of $\Gamma(-t)$ as we do. The contour C_2 can now be taken well away from any poles for example along the line $7/2 + it$ in Fig. 1 and is exponentially convergent.

Letting $N \equiv \max(k + \ell - s, -k - \ell - s - 1)$ denote the rightmost pole of $\Gamma(a+t)$ and $\Gamma(b+t)$, the sum over residues gives

$$F_1(k) \equiv \sum_{n=0}^N \frac{(-1)^n}{n!} \frac{\Gamma(n+k+\ell+s+1-i\bar{\omega}+\delta\nu)\Gamma(n-k-\ell+s-i\bar{\omega}-\delta\nu)}{\Gamma(n+1-2i\bar{\omega})} (\bar{r}-1)^n, \quad (6.8)$$

where the coefficients of the polynomial may readily be expanded about $\bar{\omega} = 0$. The corresponding contribution from C_2 is given by

$$\begin{aligned} F_2(k) \equiv \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\Gamma(N+k+\ell+s+3/2-i\bar{\omega}+\delta\nu+iy)\Gamma(N-k-\ell+s+1/2-i\bar{\omega}-\delta\nu+iy)\Gamma(-N-1/2-iy)}{\Gamma(N+3/2-2i\bar{\omega}+iy)} \\ \times (r-1)^{N+1/2+iy} dy. \end{aligned} \quad (6.9)$$

From the above, we obtain

$$f_\ell(r, \omega) = \bar{r}^{s+1} \frac{e^{-i\bar{\omega}\bar{r}} (\bar{r}-1)^{-i\bar{\omega}}}{\sum_{k=-\infty}^{\infty} a_k \frac{\Gamma(a)\Gamma(b)}{\Gamma(c)}} \sum_{k=-\infty}^{\infty} a_k \cdot (F_1(k) + F_2(k)). \quad (6.10)$$

We note that the integral in F_2 is exponentially convergent. All that one has to do is completely trivial small- $\bar{\omega}$ expansions of the various coefficients appearing in Eqs. (6.8) and (6.9). For that, the small- $\bar{\omega}$ expansions for ν and for a_k of the previous section are required.

C. Cases $\ell = -s = 0, 1, 2$

The results of the previous subsection are valid for *any* values of the spin s [except for, obviously, Eq. (6.6)] and multipole number $\ell \geq |s|$. In this subsection we use those results to give the small- $\bar{\omega}$ expansions for the ingoing RW solutions specifically for the modes $\ell = -s = 0, 1$ and 2 , as these are the modes that we will use in the next section in order to calculate the Schwarzschild black hole response to a specific perturbation.

Specifically, for $\ell = s = 0$ we have, for general r ,

$$f_\ell(r, -i\sigma) = \bar{r} + \bar{r} \left\{ -\log(\bar{r} - 1) - 1 + \int_{-\infty}^{\infty} dv \frac{(\bar{r} - 1)^{\frac{1}{2}+iv} \text{sech}(\pi v)}{1 + 2iv} \right\} \bar{\sigma} + \frac{\bar{r}}{6} \left\{ \bar{r}^2 + 8\bar{r} - 6 + 3 \log(\bar{r} - 1)(\log(\bar{r} - 1) + 2) \right. \\ \left. - \int_{-\infty}^{\infty} dv \frac{(\bar{r} - 1)^{\frac{1}{2}+iv} \text{sech}(\pi v)}{(4v^2 - 8iv - 3)} \left(21 - 3\bar{r} - 6i\bar{r}v + 10iv + \frac{24}{1 + 2iv} - 6(3 + 2iv) \log(\bar{r} - 1) \right) \right\} \bar{\sigma}^2 + o(\bar{\sigma}^2). \quad (6.11)$$

For $\ell = -s = 1$ we have, for general r ,

$$f_\ell(r, -i\sigma) = \bar{r}^2 - \left\{ \frac{\bar{r}}{2^4} (3 - 2\bar{r} + \bar{r} \log(\bar{r} - 1)) + \frac{1}{\pi} \int_{-\infty}^{\infty} dv (\bar{r} - 1)^{\frac{5}{2}+iv} \Gamma\left(-iv - \frac{5}{2}\right) \Gamma\left(iv + \frac{1}{2}\right) \right\} \bar{\sigma} \\ + \left\{ 20 - 168\bar{r} + 144\bar{r}^2 + 16\bar{r}^3 + 3\bar{r}^4 + 15\bar{r} \log(\bar{r} - 1)(6 - 4\bar{r} + \bar{r} \log(\bar{r} - 1)) \right. \\ \left. + \frac{1}{4} \int_{-\infty}^{\infty} dv \frac{(\bar{r} - 1)^{\frac{5}{2}+iv} \text{sech}(\pi v) \Gamma(iv + \frac{1}{2})}{\Gamma(iv + \frac{9}{2})} \left(809 - 15\bar{r}(1 + 2iv) - 60(7 + 2iv) \log(\bar{r} - 1) \right. \right. \\ \left. \left. + 154iv + \frac{720}{2iv + 1} + \frac{480}{2iv + 3} + \frac{240}{2iv + 5} \right) \right\} \frac{\bar{\sigma}^2}{30} + o(\bar{\sigma}^2). \quad (6.12)$$

For $\ell = -s = 2$ we have, for general r ,

$$f_\ell(r, -i\sigma) = \bar{r}^3 - \left\{ \frac{1}{6} (6\bar{r}^3 \log(\bar{r} - 1) - 19\bar{r}^3 + 30\bar{r}^2 - 15\bar{r} + 10) + \frac{12}{\pi\bar{r}} \int_{-\infty}^{\infty} dv (\bar{r} - 1)^{\frac{9}{2}+iv} \Gamma\left(-\frac{9}{2} - iv\right) \Gamma\left(\frac{1}{2} + iv\right) \right\} \bar{\sigma} \\ + \left\{ \frac{1}{5040\bar{r}} [840\bar{r} \log(\bar{r} - 1)(3\bar{r}^3 \log(\bar{r} - 1) - 19\bar{r}^3 + 30\bar{r}^2 - 15\bar{r} + 10) \right. \\ \left. + 360\bar{r}^6 + 1728\bar{r}^5 + 40385\bar{r}^4 - 67020\bar{r}^3 + 49470\bar{r}^2 - 397760\bar{r} + 24860] \right. \\ \left. + \frac{1}{35\pi\bar{r}} \int_{-\infty}^{\infty} dv \frac{(\bar{r} - 1)^{\frac{9}{2}+iv} \Gamma(iv + \frac{1}{2})}{\Gamma(iv + \frac{13}{2})} [175\pi(\bar{r} - 1)(13 + 2iv) \text{sech}(\pi v) \right. \\ \left. + \Gamma\left(-\frac{9}{2} - iv\right) \Gamma\left(\frac{13}{2} + iv\right) \left(420\bar{r} + 420 \log(\bar{r} - 1) + 420\psi\left(iv + \frac{1}{2}\right) - 420\psi\left(iv + \frac{11}{2}\right) - 1019 \right) \right] \right\} \bar{\sigma}^2 \\ + o(\bar{\sigma}^2), \quad (6.13)$$

where ψ is the digamma function.

VII. LATE-TIME TAIL

In this section we illustrate how one can apply the small-frequency expansions of the previous sections to obtain physically relevant results: the late-time behavior to high order of the black hole response, at an arbitrary point, to a field perturbation of arbitrary integer spin. Before we do that in Sec. VII B, we first derive the late-time tail of the Green function in the next subsection.

A. Late-time tail of the Green function

The late-time behavior of the RW Green function is, via Eq. (2.17), given by that of its ℓ -modes ${}_s G_\ell^{\text{ret}}$. In its turn, the late-time behavior of ${}_s G_\ell^{\text{ret}}$ is dominated by the BC ℓ -modes ${}_s G_\ell^{\text{BC}}$ in Eq. (2.19). Finally, it follows from the latter equation that ${}_s G_\ell^{\text{BC}}$ at late times is given by the small-frequency behavior of $\delta_s G_\ell$. We now give more detailed expressions for these expansions.

The radius-independent part of the BC integrand in the BPT case is given in general (except for $\ell = s = 0$) in Eq. (5.14), and specifically in Eqs. (5.20)–(5.22) for the lower modes. Let us generically write its small-frequency expansion as

$$\frac{q^T}{|W^T|^2} = v_0 \bar{\sigma}^{2\ell+1} + v_1 \bar{\sigma}^{2\ell+2} + (v_{2a} + v_{2b} \ln \bar{\sigma}) \bar{\sigma}^{2\ell+3} + o(\bar{\sigma}^{2\ell+3}), \quad (7.1)$$




FIG. 1 (color online). Plot of the contour deformation in the complex t -plane for the Barnes integral representation [Eq. (6.7)] for the hypergeometric function. The straight blue curve is the contour (C_1) in Eq. (6.7). The dashed red vertical line is the contour (C_2) in Eq. (6.9). The circled crosses correspond to the poles of the various Γ functions in the integrand of Eq. (6.7): the blue ones (at $t = 0, 1, 2, \dots$) are those of $\Gamma(-t)$; the green ones (at $t = -a, -a-1, -a-2, \dots$) are those of $\Gamma(a+t)$; the red ones (at $t = -b, -b-1, -b-2, \dots$) are those of $\Gamma(b+t)$. This plot is for the case $k = 1, \ell = 2, s = 0, \bar{\omega} = -0.1i$, which yields $a = 3.9 + O(\bar{\omega}^2)$, $b = -3.1 + O(\bar{\omega}^2)$ and $N = 3$.

where the $\bar{\sigma}$ -independent constant coefficients v_0, v_1 and $v_{2a/b}$ are readily readable from the mentioned equations. In order to obtain the corresponding RW quantity, we can

$$\begin{aligned}
 {}_s G_\ell^{\text{BC}} = & -\frac{(2\ell+2)!}{4\pi M^2} \{C_0 v_0 f_{00}(r, r') + (2\ell+3)[C_0 v_0 f_{01}(r, r') + (C_0 v_1 + C_1 v_0) f_{00}(r, r')] \bar{t}^{-1} \\
 & + 2(\ell+2)(2\ell+3)[C_0 v_0 f_{012}(r, r') + (v_0 C_1 + C_0 v_1) f_{01}(r, r') \\
 & + (C_0 v_{2b}(\psi(2\ell+5) - \ln \bar{t}) + C_0 v_{2a} + v_0 C_2 + C_1 v_1) f_{00}(r, r')] \bar{t}^{-2} \} \bar{t}^{-2\ell-3} + o(\bar{t}^{-2\ell-5}), \quad (7.4)
 \end{aligned}$$

where

$$\begin{aligned}
 f_{00}(r, r') & \equiv f_{\ell 0}(r) f_{\ell 0}(r'), & f_{01}(r, r') & \equiv f_{\ell 0}(r) f_{\ell 1}(r') + f_{\ell 0}(r') f_{\ell 1}(r), \\
 f_{012}(r, r') & \equiv f_{\ell 0}(r) f_{\ell 2}(r') + f_{\ell 0}(r') f_{\ell 2}(r) + f_{\ell 1}(r) f_{\ell 1}(r'). \quad (7.5)
 \end{aligned}$$

Here we are using the dimensionless time $\bar{t} \equiv t/(2M)$. The leading-order behavior at late times is, therefore, ${}_s G_\ell^{\text{BC}} \sim \bar{t}^{-2\ell-3}$ for all integer spins. We also note that the logarithmic behavior in Eq. (5.14) for $q^T/|W^T|^2$ for small frequencies led to the appearance of a logarithmic behavior in ${}_s G_\ell^{\text{BC}}$ as $\bar{t}^{-2\ell-5} \ln \bar{t}$. The late-time behavior of the GF of the four-dimensional RW equation is then obtained by replacing ${}_s G_\ell^{\text{ret}}$ in Eq. (2.17) by the expansion in Eq. (7.4).

A similar analysis can be done for the BPT GF, but using the corresponding equations instead: (2.37) and (2.35). The radius-independent part of the integrand, $q^T/|W^T|^2$, we have already given for the BPT case. The radius-dependent

trivially use Eq. (2.46). We write the small-frequency expansion of the proportionality constant in Eq. (2.46) as

$$C = C_0 + C_1 \bar{\sigma} + C_2 \bar{\sigma}^2 + O(\bar{\sigma}^3), \quad (7.2)$$

where the $\bar{\sigma}$ -independent constant coefficients $C_{0/1/2}$ can be read off from Eqs. (2.47) and (2.48).

The radius-dependent part of the BC integrand in the RW case is given in general in Eq. (6.10); all that one has to do is a trivial small- $\bar{\omega}$ expansion of the summand in Eq. (6.8) and the integrand in Eq. (6.9) [with the use of Eqs. (4.1) and (4.3) in general, and specifically Eq. (4.10) for the lower modes]. Again, let us write the small-frequency expansion of the ingoing RW solution as

$$f_\ell(r, -i\sigma) = f_{\ell 0}(r) + f_{\ell 1}(r) \bar{\sigma} + f_{\ell 2}(r) \bar{\sigma}^2 + O(\bar{\sigma}^3), \quad (7.3)$$

where the $\bar{\sigma}$ -independent (but radius-dependent) coefficients $f_{\ell,0/1/2}$ can be readily obtained in the manner just indicated.

The analytic small-frequency expansions of the radius-independent [Eq. (7.1) times Eq. (7.2)] and radius-dependent [Eq. (7.3) evaluated at r times the same expression but evaluated at r'] parts of the BC integrand in the RW case are then to be put together in Eq. (2.21) and integrated as per Eq. (2.19). The result, for late times is, straightforwardly,

part can be obtained from the expansions Eq. (7.3) of the ingoing RW solutions via the Chandrasekhar transformation of Eq. (2.39). The late-time behavior of the ℓ -modes of the BPT GF is of the same leading order as that in the RW case (i.e., $\bar{t}^{-2\ell-3}$) and the logarithmic term also generally appears at the same order as in the RW case (i.e., $\bar{t}^{-2\ell-5} \ln \bar{t}$).

B. Late-time tail of an initial perturbation

We shall now give a particular application of our late-time results for the GF. Let us here consider an initial field perturbation given by $u_\ell^{\text{ic}}(r'_*) \equiv u_\ell(r'_*, t' = 0)$ for the

ℓ -multipole of the field and $\dot{u}_\ell^{ic}(r'_*) \equiv \partial_t u_\ell(r'_*, t' = 0)$ for the ℓ -multipole of the time derivative of the field. Then the response of a Schwarzschild black hole is given by

$$u_\ell(r_*, t) = \int_{-\infty}^{\infty} dr'_* \left[{}_s G_\ell^{\text{ret}}(r, r'; t) \dot{u}_\ell^{ic}(r'_*) + u_\ell^{ic}(r'_*) \frac{\partial}{\partial t} {}_s G_\ell^{\text{ret}}(r, r'; t) \right], \quad (7.6)$$

where $G_\ell^{\text{ret}}(r, r'; t)$ is the ℓ -multipole of the retarded GF of the wave equation satisfied by the field. In this section we will take the RW equation as the wave equation. The late-time asymptotics of the perturbation u_ℓ are given by replacing ${}_s G_\ell^{\text{ret}}$ by ${}_s G_\ell^{\text{BC}}$ in Eq. (7.6) and approximating ${}_s G_\ell^{\text{BC}}$ by performing a small-frequency expansion of $\delta_s G_\ell$ in Eq. (2.19).

As in [33], let us consider the following initial perturbation for general spin¹¹

$$u_\ell^{ic}(r_*) = \frac{1}{M} \exp\left(-\frac{(r_* - x_0)^2}{2M^2}\right), \quad \dot{u}_\ell^{ic}(r_*) = -\frac{(r_* - x_0)}{M^2} u_\ell^{ic}(r_*), \quad (7.7)$$

with $x_0 \equiv r_*(10M)$.

In Figs. 2–6 we plot the time evolution via the RW equation of a spin- s field for various multipole numbers ℓ using the initial data Eq. (7.7), similarly to [33] where we only presented the case $s = 0$, $\ell = 1$. We calculate the time evolution numerically up to $T = 3000M$, where $T \equiv t - 2x_0$, and we compare it with late-time asymptotics. The numerical solution is obtained using the code in [53] for the $(1 + 1)$ -dimensional differential equation which results from the 4D RW equation after factorizing out the angle dependence of the solution via scalar spherical harmonics. We obtain the late-time asymptotics, up to four leading orders, from the results of the previous section. In order to obtain the small-frequency expansion of $\delta_s G_\ell$ via Eq. (2.21), we need the small-frequency expansion of $q/|W|^2$ and of the “in” radial solution f_ℓ . We now give the specific expansions of the perturbation response for different spins s and ℓ modes.

We found the late-time asymptotics for the case $s = 0$ and $\ell = 0$ using Eq. (6.11) for the radial solution and Eq. (5.20) for $q/|W|^2$; for $\ell = 1, 2$ we merely inserted the corresponding value of ℓ into Eq. (5.16). The results are the following (the approximation sign is due to the fact that we have rounded up the coefficients to seven significant figures):

$$u_\ell^{\text{BC}} \approx \frac{20.02467}{\bar{T}^3} - \frac{269.1100}{\bar{T}^4} + \frac{-881.0855 \ln(\bar{T}) + 7561.913}{\bar{T}^5} + o(\bar{T}^{-5}), \quad s = \ell = 0, \quad (7.8)$$

$$u_\ell^{\text{BC}} \approx -\frac{1144.318}{\bar{T}^5} + \frac{52385.24}{\bar{T}^6} + \frac{43484.10 \ln(\bar{T}) - 1746947}{\bar{T}^7} + o(\bar{T}^{-7}), \quad s = 0, \ell = 1, \quad (7.9)$$

$$u_\ell^{\text{BC}} \approx \frac{42634.22}{\bar{T}^7} - \frac{3114525}{\bar{T}^8} + \frac{-1796322 \ln(\bar{T}) + 146090980}{\bar{T}^9} + o(\bar{T}^{-9}), \quad s = 0, \ell = 2. \quad (7.10)$$

Figures 2–4 show excellent agreement at late times between the numerical solution of the RW equation for $s = 0$ and $\ell = 0, 1, 2$ and the late-time asymptotics of Eqs. (7.8)–(7.10).

The late-time asymptotics for the RW cases $|s| = \ell = 1$ and 2 are obtained similarly to those above for $s = 0$: we used Eqs. (7.1) and (7.2) for the small-frequency expansions of the radial solution and Eqs. (5.21) and (5.22) for that of $q^T/|W^T|^2$, which is then simply converted to RW version with the use of Eqs. (2.44) and (B4). The final results for the late-time perturbation are the following:

$$u_\ell^{\text{BC}} \approx -\frac{1338.214}{\bar{T}^5} + \frac{59157.01}{\bar{T}^6} - \frac{1970216 - 62896.05 \ln(\bar{T})}{\bar{T}^7} + o(\bar{T}^{-7}), \quad |s| = \ell = 1, \quad (7.11)$$

and

$$u_\ell^{\text{BC}} \approx \frac{60748.13}{\bar{T}^7} - \frac{4318418}{\bar{T}^8} + \frac{(198106997 - 3466693 \ln(\bar{T}))}{\bar{T}^9} + o(\bar{T}^{-9}), \quad |s| = \ell = 2. \quad (7.12)$$

¹¹As an alternative to the initial data of Eq. (7.7) we could consider the following initial data (as used in [52]): zero for the field and a Gaussian distribution of small width for the time derivative of the field. In this case, the perturbation response is an approximation to the Green function. However, we found that such an approximation is not as good as (7.7) for assessing the validity of the late-time asymptotics to the higher-than-leading orders as intended in this paper.




FIG. 2 (color online). Scalar field Mu_ℓ at $r_* = x_0$ as a function of time T/M for the initial data [Eq. (7.7)] with $x_0 = r_*(10M)$ and $s = \ell = 0$ (i.e., similar to Fig. 1 in [33] and with the same initial data but here it is for $s = \ell = 0$). Red curve: Numerical solution. Dashed black curve (overlapping with red curve): Late-time expression given by Eq. (7.8) [which comes from Eq. (5.20) together with the small- $\bar{\sigma}$ results in Sec. VI for the radial function]. Green curve: Numerical solution minus the leading order [i.e., $O(\bar{T}^{-3})$] in the late-time expression. Solid blue curve: Numerical solution minus the two leading orders [i.e., $O(\bar{T}^{-3})$ and $O(\bar{T}^{-4})$] in the late-time expression. Solid cyan curve: Numerical solution minus the four leading orders [i.e., $O(\bar{T}^{-3})$, $O(\bar{T}^{-4})$, $O(\bar{T}^{-5} \ln(\bar{T}))$ and $O(\bar{T}^{-5})$] in the late-time expression.

Figures 5 and 6 show excellent agreement at late times between the numerical solution of the RW equation for $|s| = \ell = 1, 2$ and the late-time asymptotics of Eqs. (7.11)–(7.12).

The above examples illustrate the appearance of a higher-order logarithmic behavior of a perturbation response, appearing after Price’s well-known power-law tail [3,4]. We note that, as already pointed out in Sec. VD,




FIG. 3 (color online). Similar to Fig. 2 but here it is for $s = 0$, $\ell = 1$ (the orders subtracted from the numerical solution are the corresponding leading order, next-to-leading order and four leading orders for this case). The late-time expression is given by Eq. (7.9).




FIG. 4 (color online). Similar to Fig. 2 but here it is for $s = 0$, $\ell = 2$ (the orders subtracted from the numerical solution are the corresponding leading order, next-to-leading order and four leading orders for this case).




FIG. 5 (color online). Similar to Fig. 2 but here it is the RW solution for $|s| = \ell = 1$ (the orders subtracted from the numerical solution are the corresponding leading order, next-to-leading order and four leading orders for this case).




FIG. 6 (color online). Similar to Fig. 2 but here it is the RW solution for $|s| = \ell = 2$ (the orders subtracted from the numerical solution are the corresponding leading order, next-to-leading order and four leading orders for this case).

this logarithmic behavior appears at *third* leading order at late times. It appears at third leading order, not at second leading order as one might have naively expected from the fact that the logarithmic behavior at small frequency appears at the second leading order both in the BC strength q^T and in the Wronskian. This is due to there being delicate cancellations between certain terms of q^T and the Wronskian, which we have been able to derive using the precise values of the coefficients in the expansions.

VIII. CONCLUSIONS

In this paper we have developed the MST formalism for the solutions of the radial Regge-Wheeler and Bardeen-Press-Teukolsky equations, which are obeyed by linear field perturbations of a Schwarzschild black hole spacetime. We have derived, for the first time, the MST formalism for the solutions of the RW equation for spin-1 as well as the MST expressions for the branch-cut-relevant quantities for general spin. We have given explicit expansions for small frequency up to the first four leading orders for the various MST quantities for general spin. In principle, the MST series could be expanded to arbitrarily large order in the frequency. The main difficulty in achieving that is the fact that the small-frequency expansion of the renormalized angular momentum parameter ν that we currently use does not reproduce the numerically observed behavior of a *sudden* appearance of an imaginary part in ν as the frequency is increased from 0 to larger real values.

We have used our small-frequency expansions in order to obtain the late-time behavior for arbitrary radius of spin-field perturbations of a Schwarzschild black hole up to the first four leading orders (for a specific multipole- ℓ). Our results explicitly reveal a new logarithmic behavior at third order for late times as $t^{-2\ell-5} \ln t$. We note that the appearance of a logarithmic behavior was already predicted by other works (see, e.g., [28,54–56]). However, to the best of our knowledge, the order at which the logarithmic behavior appears was not correctly predicted anywhere (nor was the calculation of the coefficients for general radius carried out). As we noted at the end of Sec. V D, there is a delicate cancellation between different logarithmic terms which averts the appearance of a logarithmic behavior at a lower order; this delicate cancellation is probably hard to predict unless an exact and detailed analysis is carried out such as the one in this paper (see also [57], where they find numerical evidence that the logarithmic behavior does not appear at first nor second leading orders).

We succinctly presented in the paper [33] the final results that we have derived in this paper. We already used some of these results in the calculation of the scalar self-force in Schwarzschild spacetime carried out in [20]. The natural extension of our results is the explicit calculation of the small-frequency expansions of the MST quantities for the

Teukolsky equation in Kerr spacetime, with the corresponding late-time analysis of perturbations of a Kerr black hole. We expect to present the Kerr analysis in the near future, together with its application to a self-force calculation.

Finally, while the calculation of the quasinormal modes has been applied to the modeling of radiation after the inspiral of two black holes via a matching to a numerical relativity solution, to the best of our knowledge, the branch cut has never been taken into account for such purposes. We would expect that the inclusion in the modeling of the branch cut, together with the quasinormal modes, would help match the analytical solution from perturbation theory with the numerical relativity one.

ACKNOWLEDGMENTS

We are thankful to Chris Kavanagh and Barry Wardell for useful discussions. A. C. O. acknowledges support from Science Foundation Ireland under Grant No. 10/RFP/PHY2847.

APPENDIX A: VALIDATION OF BC RESULTS

In this section we compare the small- $\bar{\sigma}$ results presented in this paper with an independent method which we presented in [29]. The latter method is naturally adapted to a midfrequency regime (which, although valid for all frequencies, is not practical to use in the asymptotic small- or large-frequency regimes). In this Appendix we illustrate with plots that there is a region of overlap between the results obtained with the small- $\bar{\sigma}$ method presented here and those obtained with the midfrequency method of [29].




FIG. 7 (color online). The radial function $e^{-i\omega r} \hat{f}_\ell(r, \omega)$ at $r = 10M$ as a function of $\bar{\sigma}$ for $s = 0$, $\ell = 1$. We calculated the straight blue curve using the so-called Jaffé series (see, e.g., [13,29]) and the dashed red curve using the small-frequency asymptotics presented in this paper up to $O(\bar{\sigma}^{15})$ [specifically, using Eq. (137) [16] as in this case we did not need to obtain the coefficients of the expansion analytically via the use of the Barnes integral method].




FIG. 8 (color online). BC strength $q(\sigma)$ of Eq. (2.20) [or, equivalently for $s = 0$, $q^T(\sigma)$ of Eq. (2.36)] as a function of $\bar{\sigma}$. In the straight blue curves we calculated $q(\sigma)$ using the midfrequency method of [29] (with the radial functions calculated at $r = 5M$). In the dashed red curves we calculated $q(\sigma)$ using the small-frequency asymptotics presented in this paper up to $O(\bar{\sigma}^{15})$. (a) For $s = 0$, $\ell = 0$. (b) For $s = 0$, $\ell = 1$.




FIG. 9 (color online). The Wronskian $W[\hat{f}_\ell, g_\ell]$ as a function of $\bar{\sigma}$ for $s = 0$, $\ell = 1$. The straight-blue and dotted-red curves respectively correspond to the real and imaginary parts of the Wronskian evaluated using the midfrequency method of [29] with the radial functions calculated at $r = 2.8M$; the convergence of the series used in the midfrequency method becomes slower as the frequency becomes smaller. The dashed blue and red curves respectively correspond to the real and imaginary parts of the Wronskian evaluated using the small-frequency asymptotics presented in this paper up to $O(\bar{\sigma}^{14})$. In (b) we plot the midfrequency to larger values of the frequency so as to show its general form; the green-dashed curve here corresponds to the absolute value of the Wronskian. Compare Fig. 13 [29] for $s = \ell = 2$.

In [29]¹² we already presented plots of quantities focused mainly on the $s = \ell = 2$ case. Therefore, for variety, here we will focus on the $s = 0$ and $\ell = 0, 1$ cases.

As explained in [29], the ingoing radial function $f_\ell(r, \omega)$ possesses simple poles along the NIA (which are ultimately irrelevant, as they are canceled out in the Fourier modes of the GF by the corresponding poles in the Wronskian). It is therefore useful to define the following radial function: $\hat{f}_\ell(r, \omega) \equiv \sin(-2\pi i \bar{\omega}) f_\ell(r, \omega)$.

¹²We note that there are two typos in Eq. (3.6) [29]: there should be an overall factor “−1” in the expression for $\Delta \hat{h}_\ell(r, \nu)$, using the notation of that paper, and, in the expression for $\Delta \hat{h}_{n,\ell}$, it should be \tilde{a}_{n+} instead of \tilde{a}_{n-} . The minus sign typo carried over to Fig. 10 [29], which is therefore a plot of “ $+ie^{\nu r_*} \Delta \tilde{g}_\ell$,” instead of “ $-ie^{\nu r_*} \Delta \tilde{g}_\ell$ ” as stated there.




FIG. 10 (color online). BC mode $\delta_s G_\ell(r, r', -i\sigma)$ of Eq. (2.21) as a function of $\bar{\sigma}$ at $r = r(r_* = 0.4M)$, $r' = r'(r_* = 0.2M)$ for $s = 0$ and $\ell = 1$. In the straight blue curve we calculated the BC mode using the midfrequency method of [29] and in the dashed red curve we calculated it using the small-frequency asymptotics presented in this paper up to $O(\bar{\sigma}^{15})$.

In Figs. 7–10 we respectively plot the radial function f_ℓ , the BC strength $q(\sigma)$, the Wronskian W and the BC mode $\delta_s G_\ell(r, r', -i\sigma)$, all as functions of the frequency $\bar{\sigma}$ along the NIA.

APPENDIX B: RADIAL COEFFICIENTS

In this Appendix we use the Chandrasekhar transformation [Eq. (2.38)] in order to relate the BPT and RW radial coefficients and Wronskians. Let us first add higher orders to the asymptotics in Eq. (2.11) of the RW solution:

$$f_\ell \sim \begin{cases} (1 + \alpha_+ \Delta + \beta_+ \Delta^2) e^{-i\omega r_*}, & r_* \rightarrow -\infty, \\ A_\ell^{\text{in}} \left(1 + \frac{\alpha_\infty}{r} + \frac{\beta_\infty}{r^2} \right) e^{-i\omega r_*} + A_\ell^{\text{out}} e^{+i\omega r_*}, & r_* \rightarrow +\infty, \end{cases} \quad (\text{B1})$$

where α_+ , β_+ , α_∞ and β_∞ are coefficients to be determined. By imposing that these asymptotics satisfy the RW equation, we find

$$\alpha_\infty = -\frac{i\lambda}{2\omega}, \quad \beta_\infty = \frac{\lambda(2-\lambda) + 4Mi(s^2-1)\omega}{8\omega^2},$$

$$\alpha_+ = \frac{i(\lambda+1-s^2)}{4M^2(4M\omega+i)}, \quad \beta_+ = \frac{4-s^4-\lambda^2 + \lambda(2-16iM\omega) + s^2(2\lambda+20iM\omega-3) - 20iM\omega}{64M^4(2M\omega+i)(4M\omega+i)}. \quad (\text{B2})$$

We note that α_∞ , α_+ , β_∞ , β_+ are all real valued when ω is purely imaginary.

Using the transformation Eq. (2.38) it is straightforward to find the following coefficients of BPT solutions:

$${}_s \tilde{R}_\ell^{\text{in,inc}} = \begin{cases} A_\ell^{\text{in}}, & s=0, \\ -\alpha_\infty A_\ell^{\text{in}}, & s=-1, \\ 2\beta_\infty A_\ell^{\text{in}}, & s=-2. \end{cases} \quad {}_s \tilde{R}_\ell^{\text{in,tra}} = \begin{cases} \frac{1}{2M}, & s=0, \\ \alpha_+, & s=-1, \\ \frac{2(\alpha_+ + 2M^2\beta_+)}{M}, & s=-2. \end{cases} \quad {}_s \tilde{R}_\ell^{\text{up,tra}} = \begin{cases} 1, & s=0, \\ 2i\omega, & s=-1, \\ -4\omega^2, & s=-2. \end{cases} \quad (\text{B3})$$

It then immediately follows, using Eqs. (2.13) and (2.34), that

$$W^T = W \cdot \begin{cases} 2M, & s=0, \\ \frac{-\alpha_\infty}{\alpha_+}, & s=-1, \\ \frac{M\beta_\infty}{\alpha_+ + 2M^2\beta_+}, & s=-2. \end{cases} \quad (\text{B4})$$

Note that W^T is, by definition, independent of the chosen normalization for ${}_s R_\ell^{\text{in}}$ and ${}_s R_\ell^{\text{up}}$.

-
- [1] T. Regge and J.A. Wheeler, *Phys. Rev.* **108**, 1063 (1957).
[2] J. A. Wheeler, *Phys. Rev.* **97**, 511 (1955).
[3] R. H. Price, *Phys. Rev. D* **5**, 2419 (1972).
[4] R. H. Price, *Phys. Rev. D* **5**, 2439 (1972).
[5] F. J. Zerilli, *Phys. Rev. Lett.* **24**, 737 (1970).
[6] F. J. Zerilli, *Phys. Rev. D* **2**, 2141 (1970).
[7] R. Ruffini, J. Tiomno, and C. Vishveshwara, *Lett. Nuovo Cimento* **3**, 211 (1972).
[8] S. A. Teukolsky, *Phys. Rev. Lett.* **29**, 1114 (1972).
[9] S. A. Teukolsky, *Astrophys. J.* **185**, 635 (1973).
[10] J. M. Bardeen and W. H. Press, *J. Math. Phys. (N.Y.)* **14**, 7 (1973).
[11] S. Hod, *Phys. Rev. D* **58**, 104022 (1998).
[12] L. Barack, *Phys. Rev. D* **61**, 024026 (1999).
[13] E. W. Leaver, *J. Math. Phys. (N.Y.)* **27**, 1238 (1986).
[14] S. Mano, H. Suzuki, and E. Takasugi, *Prog. Theor. Phys.* **95**, 1079 (1996). S. Mano, H. Suzuki, and E. Takasugi *Prog. Theor. Phys.* **96**, 549 (1996).
[15] S. Mano, H. Suzuki, and E. Takasugi, *Prog. Theor. Phys.* **96**, 549 (1996).
[16] M. Sasaki and H. Tagoshi, *Living Rev. Relativity* **6**, 6 (2003).
[17] E. Poisson, A. Pound, and I. Vega, *Living Rev. Relativity* **14**, 7 (2011).
[18] W. Hikida, S. Jhingan, H. Nakano, N. Sago, M. Sasaki, and T. Tanaka, *Prog. Theor. Phys.* **111**, 821 (2004).
[19] W. Hikida, S. Jhingan, H. Nakano, N. Sago, M. Sasaki, and T. Tanaka, *Prog. Theor. Phys.* **113**, 283 (2005).
[20] M. Casals, S. Dolan, A. C. Ottewill, and B. Wardell, *Phys. Rev. D* **88**, 044022 (2013).
[21] D. Bini and T. Damour, *Phys. Rev. D* **89**, 064063 (2014).

- [22] D. Bini and T. Damour, *Phys. Rev. D* **90**, 024039 (2014).
- [23] D. Bini and T. Damour, *Phys. Rev. D* **89**, 104047 (2014).
- [24] A. G. Shah, J. L. Friedman, and B. F. Whiting, *Phys. Rev. D* **89**, 064042 (2014).
- [25] A. G. Shah, *Phys. Rev. D* **90**, 044025 (2014).
- [26] S. Chakrabarti, T. Delsate, and J. Steinhoff, [arXiv:1304.2228](https://arxiv.org/abs/1304.2228)
- [27] E. W. Leaver, *Phys. Rev. D* **34**, 384 (1986).
- [28] E. S. C. Ching, P. T. Leung, W. M. Suen, and K. Young, *Phys. Rev. D* **52**, 2118 (1995).
- [29] M. Casals and A. C. Ottewill, *Phys. Rev. D* **87**, 064010 (2013).
- [30] M. Casals and A. Ottewill, *Phys. Rev. D* **86**, 024021 (2012).
- [31] A. Maassen van den Brink, *Phys. Rev. D* **62**, 064009 (2000).
- [32] A. Maassen van den Brink, *J. Math. Phys. (N.Y.)* **45**, 327 (2004).
- [33] M. Casals and A. Ottewill, *Phys. Rev. Lett.* **109**, 111101 (2012).
- [34] Z. Zhang, E. Berti, and V. Cardoso, *Phys. Rev. D* **88**, 044018 (2013).
- [35] H. Nakano and M. Sasaki, *Prog. Theor. Phys.* **105**, 197 (2001).
- [36] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Oxford University Press, New York, 1983).
- [37] P. T. Leung, A. Maassen van den Brink, K. W. Mak, and K. Young, [arXiv:gr-qc/0307024](https://arxiv.org/abs/gr-qc/0307024).
- [38] J. Goldberg, A. Macfarlane, E. T. Newman, F. Rohrlich, and E. Sudarshan, *J. Math. Phys. (N.Y.)* **8**, 2155 (1967).
- [39] E. T. Newman and R. Penrose, *J. Math. Phys. (N.Y.)* **7**, 863 (1966).
- [40] S. Chandrasekhar, *Proc. R. Soc. A* **343**, 289 (1975).
- [41] B. P. Jensen, J. G. McLaughlin, and A. C. Ottewill, *Phys. Rev. D* **43**, 4142 (1991).
- [42] DLMF, NIST Digital Library of Mathematical Functions, <http://dlmf.nist.gov/>, Release 1.0.5 of 2012-10-01, online companion to [43].
- [43] *NIST Handbook of Mathematical Functions*, edited by F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (Cambridge University Press, New York, 2010), print companion to [42].
- [44] I. Gradshteyn and I. Ryzhik, *Table of Integrals, Series, and Products* (Academic Press, New York, 2007).
- [45] A. Erdelyi, W. Magnus, F. Oberhettinger, and F. Tricomi, *Higher Transcendental Functions* (McGraw-Hill, New York, 1953).
- [46] R. Fujita and H. Tagoshi, *Prog. Theor. Phys.* **112**, 415 (2004).
- [47] R. Fujita and H. Tagoshi, *Prog. Theor. Phys.* **113**, 1165 (2005).
- [48] S. A. Teukolsky and W. Press, *Astrophys. J.* **193**, 443 (1974).
- [49] M. Casals, Electromagnetic quantum field theory on Kerr-Newman black holes, Ph.D. thesis, University College Dublin, 2004, [arXiv:0802.1885](https://arxiv.org/abs/0802.1885).
- [50] D. N. Page, *Phys. Rev. D* **13**, 198 (1976).
- [51] B. P. Jensen, J. G. McLaughlin, and A. C. Ottewill, *Phys. Rev. D* **51**, 5676 (1995).
- [52] B. Wardell, C. R. Galley, A. Zenginoğlu, M. Casals, S. R. Dolan, and A. C. Ottewill, *Phys. Rev. D* **89**, 084021 (2014).
- [53] B. Wardell, <https://github.com/barrywardell/scalarwave1d>.
- [54] E. S. C. Ching, P. T. Leung, W. M. Suen, and K. Young, *Phys. Rev. Lett.* **74**, 2414 (1995).
- [55] S. Hod, *Phys. Rev. D* **60**, 104053 (1999).
- [56] S. Hod, *Classical Quantum Gravity* **26**, 028001 (2009).
- [57] A. Z. Smith and L. M. Burko, *Phys. Rev. D* **74**, 028501 (2006).