

Spin asymmetries for vector boson production in polarized $p + p$ collisionsJin Huang,^{1,*} Zhong-Bo Kang,^{2,†} Ivan Vitev,^{2,‡} and Hongxi Xing^{2,§}¹*Physics Department, Brookhaven National Laboratory, Upton, New York 11973, USA*²*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

(Received 1 December 2015; published 28 January 2016)

We study the cross section for vector boson ($W^\pm/Z^0/\gamma^*$) production in polarized nucleon-nucleon collisions for low transverse momentum of the observed vector boson. For the case where one measures the transverse momentum and azimuthal angle of the vector bosons, we present the cross sections and the associated spin asymmetries in terms of transverse momentum dependent parton distribution functions (TMDs) at tree level within the TMD factorization formalism. To assess the feasibility of experimental measurements, we estimate the spin asymmetries for W^\pm/Z^0 boson production in polarized proton-proton collisions at the Relativistic Heavy Ion Collider by using current knowledge of the relevant TMDs. We find that some of these asymmetries can be sizable if the suppression effect from TMD evolution is not too strong. The W program at RHIC can, thus, test and constrain spin theory by providing unique information on the universality properties of TMDs, TMD evolution, and the nucleon structure. For example, the single transverse spin asymmetries could be used to probe the well-known Siverson function $f_{1T}^{\perp q}$, as well as the transversal helicity distribution g_{1T}^q via the parity-violating nature of W production.

DOI: 10.1103/PhysRevD.93.014036

I. INTRODUCTION

Spin-dependent observables for vector boson (W^\pm/Z^0) production in polarized nucleon-nucleon collisions offer excellent sensitivity to the spin-dependent parton distribution functions of the individual parton flavors in the nucleon [1,2]. For example, the violation of parity in the weak interactions gives rise to the single longitudinal spin asymmetries for W^\pm production in proton-proton collisions [3,4]. Such *longitudinal* spin asymmetries provide flavor separation for the quark helicity distributions for u, \bar{u}, d, \bar{d} , and, in particular, probe the antiquark polarization. For recent theoretical work and experimental measurements, see e.g. Refs. [5–8] and [9–13], respectively.

Single *transverse* spin asymmetries for vector bosons have also been proposed to probe the quark Siverson functions $f_{1T}^{\perp q}$ of individual flavors [14–17]. Quark Siverson functions represent the distributions of unpolarized quarks inside a transversely polarized nucleon through a correlation between the transverse momentum of the quark and the transverse spin of the nucleon, and they have a unique time-reversal modified universality property. It was shown from the parity and time-reversal invariance of QCD that the quark Siverson functions in semi-inclusive deep inelastic scattering (SIDIS) and those in Drell-Yan (DY) type process (e.g., $W^\pm/Z^0/\gamma^*$) should have the same functional form but an opposite sign—so-called “sign change” [16,18–21]. Quark Siverson functions have been

measured/extracted from SIDIS process, see, e.g. [22,23]. So far mainly the valence u and d quark Siverson functions were constrained, while sea quark Siverson functions remain largely unknown [24]. Single transverse spin asymmetries for vector bosons in proton-proton collisions can thus serve as these purposes, i.e., to test the sign change on one side, and to constrain the sea quark Siverson function at the same time. The sea quark distribution will be further measured with high precision in a future electron ion collider [25–27].

There are also theoretical studies of double spin asymmetries for vector boson production. For example, double transverse spin asymmetries for Drell-Yan production were originally proposed to extract the quark transversity distributions [28–32]. However, as pointed out already in [33], there could be additional terms involving the product of quark Siverson function $f_{1T}^{\perp q} f_{1T}^{\perp \bar{q}}$ (called double Siverson effect), as well as the product $g_{1T}^q g_{1T}^{\bar{q}}$ (called double worm-gear effect in [33] since g_{1T}^q is sometimes referred to as a worm-gear function). There are, of course, longitudinal-transverse double spin asymmetries for the vector boson production, which could provide complementary information on the quark helicity distribution functions [34,35].

The function g_{1T}^q is also called “transversal helicity” distribution function [36–38], as it gives the quark longitudinal polarization inside a transversely polarized proton. g_{1T}^q could also appear in single transverse spin asymmetry of W^\pm production due to the parity-violating nature of the weak interaction; see e.g. a study within the collinear twist-3 formalism [34]. Contrary to the quark Siverson function $f_{1T}^{\perp q}$ that changes sign from SIDIS to DY type processes, the transversal helicity distribution g_{1T}^q is universal between SIDIS and DY. As g_{1T}^q has been investigated in the SIDIS

*jhuang@bnl.gov

†zkang@lanl.gov

‡ivitev@lanl.gov

§hxing@lanl.gov

measurements [39–42] and is proposed to be measured with high precision in future SIDIS experiments [43], one in principle could test the universality of g_{1T}^q in W^\pm/Z^0 production in proton-proton collisions at RHIC.

RHIC has planned a dedicated transversely polarized proton-proton run to measure the spin asymmetries for W^\pm production [2]. Together with its longitudinal W^\pm physics, this will ensure a comprehensive W^\pm physics spin program. Besides the sensitivity to the spin-dependent parton distribution functions inside the nucleon, vector boson production also provides excellent constraints on the QCD evolution effects. This is because the mass of the vector boson, which sets the typical scale of the hard scattering, is usually much larger than the scales probed in the fixed-target SIDIS measurements. It is, thus, timely to present in a single, self-contained paper the results for the spin asymmetries for vector boson production in polarized nucleon-nucleon collisions. We will consider the production of vector bosons at small transverse momentum, and thus a transverse momentum dependent (TMD) factorization formalism is the appropriate framework [44–46]. Since it is now possible to perform a full reconstruction of the produced boson kinematics [2,47], we will integrate out the kinematics of the decayed leptons, and present the results at the level of the vector boson. This will simplify the physics results to facilitate the interpretation of the experimental measurements.

The rest of the paper is organized as follows. In Sec. II, we present the TMD formalism for all spin asymmetries for vector boson production ($W^\pm/Z^0, \gamma^*$) in polarized nucleon-nucleon collisions, where one measures the transverse momentum q_T and azimuthal angle ϕ_V of the vector bosons in the center-of-mass frame of the colliding nucleons. We point out that due to the parity-violation nature of the weak interaction for W^\pm/Z^0 production, besides the well-known Siverts term $\propto \sin(\phi_V - \phi_S)$, the single transverse spin asymmetry can have an additional term $\propto \cos(\phi_V - \phi_S)$ with ϕ_S the azimuthal angle of the transverse spin of the nucleon, which is related to the transversal helicity distribution g_{1T}^q [34]. In Sec. III we present the numerical estimate of these single and double spin asymmetries for polarized proton-proton collisions at RHIC and find that some asymmetries are sizable if TMD evolution does not lead to too strong a suppression. We conclude our paper in Sec. IV.

II. SPIN-DEPENDENT CROSS SECTION FOR VECTOR BOSON PRODUCTION

In this section, we derive the spin-dependent differential cross section for vector boson production ($V = W^\pm, Z^0$, or γ^*) in polarized nucleon-nucleon scattering

$$A(P_A, S_A) + B(P_B, S_B) \rightarrow V(y, q_T, \phi_V) + X, \quad (1)$$

where a polarized nucleon A of momentum P_A and spin S_A is colliding with another nucleon B of momentum P_B and spin S_B . We work in the center-of-mass (CM) frame of the colliding nucleons with $S = (P_A + P_B)^2$ the CM energy squared. In such a frame, we choose the nucleon A to be moving in the “+ z ” direction, while the nucleon B is moving in the “− z ” direction. In the final state, we have the full kinematics of the vector boson—its rapidity y and transverse momentum, with magnitude q_T and azimuthal angle ϕ_V . For a virtual photon we further observe its invariant mass Q . We will concentrate on the kinematic region where the transverse momentum q_T is much less than the mass M_V of the vector boson: $q_T \ll M_V$. This is the region where the usual TMD factorization formalism is expected to be valid [44–46].

We will take W boson production as an example to work out the leading order differential cross section in terms of the TMDs. The derivation for Z boson and virtual photon will be similar, and the corresponding results will be presented at the end of this section. The differential cross section for W boson production can be written as

$$\frac{d\sigma^W}{d^4q} = \frac{1}{2S} \left(\frac{g_W}{2\sqrt{2}} \right)^2 \sum_\lambda \epsilon_\mu^\lambda(q) \epsilon_\nu^{*\lambda}(q) W^{\mu\nu}(P_A, S_A, P_B, S_B) 2\pi \delta(q^2 - M_W^2), \quad (2)$$

where g_W is the weak charge and related to the electric charge as $e = g_W \sin \theta_W$ with θ_W the Weinberg angle. Using

$$\frac{g_W^2}{8M_W^2} = \frac{G_F}{\sqrt{2}}, \quad d^4q = \frac{1}{2} dq^2 dy d^2q_\perp, \quad (3)$$

$$\sum_\lambda \epsilon_\mu^\lambda(q) \epsilon_\nu^{*\lambda}(q) = -g_{\mu\nu} + \frac{q_\mu q_\nu}{M_W^2}, \quad (4)$$

with G_F the Fermi constant, we obtain

$$\frac{d\sigma^W}{dy d^2\vec{q}_T} = \frac{\pi G_F M_W^2}{2\sqrt{2}S} \left(-g_{\mu\nu} + \frac{q_\mu q_\nu}{M_W^2} \right) W^{\mu\nu}(P_A, S_A, P_B, S_B), \quad (5)$$

where $W^{\mu\nu}$ is the hadronic tensor and is given by

$$\begin{aligned} W^{\mu\nu}(P_A, S_A, P_B, S_B) &= \frac{1}{N_c} \sum_{q,q'} |V_{qq'}|^2 \int d^2\vec{k}_{aT} d^2\vec{k}_{bT} \delta^2(\vec{q}_T - \vec{k}_{aT} - \vec{k}_{bT}) \\ &\times \text{Tr}[\gamma^\mu (v_q - a_q \gamma^5) \Phi^q(x_a, \vec{k}_{aT}, S_A) \gamma^\nu \\ &\times (v_q - a_q \gamma^5) \bar{\Phi}^{q'}(x_b, \vec{k}_{bT}, S_B)]. \end{aligned} \quad (6)$$

Here $N_c = 3$ is the number of colors, $V_{qq'}$ are the CKM elements for weak interaction, v_q (a_q) is the vector (axial) coupling of the W boson to the quark with

$$v_q = 1, \quad a_q = 1. \quad (7)$$

The transverse momentum dependent quark-quark correlators are defined as [48]

$$\Phi^q(x_a, \vec{k}_{aT}, S_A) = \int \frac{dz^- d^2 \vec{z}_T}{(2\pi)^3} e^{ik_a^+ z^- - i\vec{k}_{aT} \cdot \vec{z}_T} \times \langle P_A, S_A | \bar{\psi}_j^q(0) \psi_i^q(z) | P_A, S_A \rangle, \quad (8)$$

$$\bar{\Phi}^q(x_b, \vec{k}_{bT}, S_B) = \int \frac{dz^+ d^2 \vec{z}_T}{(2\pi)^3} e^{ik_b^- z^+ - i\vec{k}_{bT} \cdot \vec{z}_T} \times \langle P_B, S_B | \psi_i^q(0) \bar{\psi}_j^q(z) | P_B, S_B \rangle, \quad (9)$$

where we have suppressed the gauge link dependence in the definitions. They can be expanded as the following parametrization in terms of the Dirac matrices at leading-twist

$$\Phi^q(x_a, \vec{k}_{aT}, S_A) = \Phi^{q[\gamma^+]} \frac{\gamma^-}{2} + \Phi^{q[\gamma^+ \gamma^5]} \frac{\gamma^5 \gamma^-}{2} + \Phi^{q[i\sigma^{\alpha+} \gamma^5]} \frac{-i\sigma^{\alpha+} \gamma^5}{2}, \quad (10)$$

where $\alpha = \{1, 2\}$ is a transverse index, $\Phi^{q[\Gamma]} \equiv \frac{1}{2} \text{Tr}[\Phi^q \Gamma]$ for the specific Dirac matrix Γ as given above, and likewise for $\bar{\Phi}^q$. With such an expansion at hand, we thus have the W boson cross section as

$$\frac{d\sigma^W}{dy d^2 \vec{q}_T} = \sigma_0^W \sum_{q,q'} |V_{qq'}|^2 \int d^2 \vec{k}_{aT} d^2 \vec{k}_{bT} \delta^2(\vec{q}_T - \vec{k}_{aT} - \vec{k}_{bT}) [(v_q^2 + a_q^2) (\Phi^{q[\gamma^+]} \bar{\Phi}^{q'[\gamma^-]} + \Phi^{q[\gamma^+ \gamma^5]} \bar{\Phi}^{q'[\gamma^- \gamma^5]}) - 2a_q v_q (\Phi^{q[\gamma^+]} \bar{\Phi}^{q'[\gamma^- \gamma^5]} + \Phi^{q[\gamma^+ \gamma^5]} \bar{\Phi}^{q'[\gamma^-]}) + (\Phi \leftrightarrow \bar{\Phi})], \quad (11)$$

where σ_0^W is given by

$$\sigma_0^W = \frac{\pi G_F M_W^2}{\sqrt{2} S N_c}. \quad (12)$$

It is instructive to note that there are no terms involving $\Phi^{q[\Gamma]}$ with $\Gamma = i\sigma^{\alpha\pm} \gamma^5$, i.e. the last term in Eq. (10) does not contribute to the final result. These terms are related to the transversely polarized quark distributions inside the proton. They contribute to the cross section only when one measures the angular dependence of the leptons decayed from the vector boson. When we integrate over the phase space of the decayed leptons, i.e. when one sums over the spin states of the vector boson, and contracts $W^{\mu\nu}$ with $\sum_\lambda \epsilon_\mu^\lambda(q) \epsilon_\nu^{*\lambda}(q)$ as specified in Eq. (4), they vanish.

We can further express the cross section in terms of leading-twist quark TMDs through the following standard expressions of $\Phi^{q[\Gamma]}$ for quark [48]:

$$\Phi^{q[\gamma^+]} = f_1^q(x_a, \vec{k}_{aT}^2) - \frac{\epsilon_T^{ij} k_{aT}^i S_{AT}^j}{M_A} f_{1T}^{\perp q}(x_a, \vec{k}_{aT}^2), \quad (13)$$

$$\Phi^{q[\gamma^+ \gamma^5]} = S_{AL} g_{1L}^q(x_a, \vec{k}_{aT}^2) + \frac{\vec{k}_{aT} \cdot \vec{S}_{AT}}{M_A} g_{1T}^q(x_a, \vec{k}_{aT}^2), \quad (14)$$

where f_1^q is the unpolarized quark TMD, and $f_{1T}^{\perp q}$ is the so-called quark Sivers function [49,50]. g_{1L}^q is the quark helicity distribution function, describing the quark longitudinal polarization in a longitudinally polarized proton. Finally, as we have mentioned already in the Introduction, g_{1T}^q is the “transversal helicity” distribution function,

describing the quark longitudinal polarization in a transversely polarized proton [36–38]. Similarly we have the expressions for $\bar{\Phi}^{q[\gamma^-]}$ for the antiquark:

$$\bar{\Phi}^{q[\gamma^-]} = f_1^{\bar{q}}(x_b, \vec{k}_{bT}^2) + \frac{\epsilon_T^{ij} k_{bT}^i S_{BT}^j}{M_B} f_{1T}^{\perp \bar{q}}(x_b, \vec{k}_{bT}^2), \quad (15)$$

$$\bar{\Phi}^{q[\gamma^- \gamma^5]} = -S_{BL} g_{1L}^{\bar{q}}(x_b, \vec{k}_{bT}^2) - \frac{\vec{k}_{bT} \cdot \vec{S}_{BT}}{M_B} g_{1T}^{\bar{q}}(x_b, \vec{k}_{bT}^2). \quad (16)$$

Here M_A (M_B) are the mass of the nucleon A (B), while S_{AL} , \vec{S}_{AT} and S_{BL} , \vec{S}_{BT} are the longitudinal and transverse components of the spin vector for nucleon A and B , respectively. It might be worthwhile to remind the reader that $\Phi^{q[\Gamma]}$ is associated with the incoming nucleon A , which moves in “+ z ” direction in the CM frame. On the other hand, the correlator $\bar{\Phi}^{q[\Gamma]}$ is associated with the incoming nucleon B , which moves in “− z ” direction. Their spin vectors in the CM frame have the following forms:

$$S_A^\mu = \left(S_{AL} \frac{|\vec{P}_A|}{M_A}, |\vec{S}_{AT}| \cos \phi_{S_A}, |\vec{S}_{AT}| \sin \phi_{S_A}, S_{AL} \frac{P_A^0}{M_A} \right), \quad (17)$$

$$S_B^\mu = \left(S_{BL} \frac{|\vec{P}_B|}{M_B}, |\vec{S}_{BT}| \cos \phi_{S_B}, |\vec{S}_{BT}| \sin \phi_{S_B}, -S_{BL} \frac{P_B^0}{M_B} \right). \quad (18)$$

Substituting the parametrization of $\Phi^{q[\Gamma]}$ and $\bar{\Phi}^{q[\Gamma]}$ into Eq. (11), we arrive at the final result:

$$\begin{aligned}
\frac{d\sigma^W}{dyd^2\vec{q}_T} = & \sigma_0^W \{ F_{UU} + S_{AL}F_{LU} + S_{BL}F_{UL} + S_{AL}S_{BL}F_{LL} + |\vec{S}_{AT}|[\sin(\phi_V - \phi_{S_A})F_{TU}^{\sin(\phi_V - \phi_{S_A})} + \cos(\phi_V - \phi_{S_A})F_{TU}^{\cos(\phi_V - \phi_{S_A})}] \\
& + |\vec{S}_{BT}|[\sin(\phi_V - \phi_{S_B})F_{UT}^{\sin(\phi_V - \phi_{S_B})} + \cos(\phi_V - \phi_{S_B})F_{UT}^{\cos(\phi_V - \phi_{S_B})}] \\
& + |\vec{S}_{AT}|S_{BL}[\sin(\phi_V - \phi_{S_A})F_{TL}^{\sin(\phi_V - \phi_{S_A})} + \cos(\phi_V - \phi_{S_A})F_{TL}^{\cos(\phi_V - \phi_{S_A})}] \\
& + S_{AL}|\vec{S}_{BT}|[\sin(\phi_V - \phi_{S_B})F_{LT}^{\sin(\phi_V - \phi_{S_B})} + \cos(\phi_V - \phi_{S_B})F_{LT}^{\cos(\phi_V - \phi_{S_B})}] \\
& + |\vec{S}_{AT}||\vec{S}_{BT}|[\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})F_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})} + \cos(\phi_{S_A} - \phi_{S_B})F_{TT}^1 \\
& + \sin(2\phi_V - \phi_{S_A} - \phi_{S_B})F_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})} + \sin(\phi_{S_A} - \phi_{S_B})F_{TT}^2] \}. \quad (19)
\end{aligned}$$

To write explicitly the expressions for all the structure functions F in the above equation, let us define the following shorthand notation:

$$\mathcal{C}^W[w(\vec{k}_{aT}, \vec{k}_{bT})f_1\bar{f}_2] = \sum_{q,q'} |V_{qq'}|^2 \int d^2\vec{k}_{aT} d^2\vec{k}_{bT} \delta^2(\vec{q}_T - \vec{k}_{aT} - \vec{k}_{bT}) w(\vec{k}_{aT}, \vec{k}_{bT}) [f_1^q(x_a, \vec{k}_{aT}^2) f_2^{\bar{q}'}(x_b, \vec{k}_{bT}^2) + (q \leftrightarrow \bar{q}')]. \quad (20)$$

One should be careful in the second term $q \leftrightarrow \bar{q}'$ when we interchange the roles of quarks and antiquarks. Due to the fact that [48,51]

$$\bar{\Phi}^{q[\Gamma]} = \pm \Phi^{\bar{q}[\Gamma]}, \quad \begin{cases} + & \text{for } \gamma^\mu, i\sigma^{\mu\nu}\gamma^5 \\ - & \text{for } \gamma^\mu\gamma^5, 1, i\gamma^5 \end{cases}, \quad (21)$$

when a term involves an odd number of g_{1L} and/or g_{1T} , there should be a minus sign when one interchanges $q \leftrightarrow \bar{q}'$. For example,

$$\begin{aligned}
\mathcal{C}^W[(v_q^2 + a_q^2)f_1\bar{f}_1] = & \sum_{q,q'} |V_{qq'}|^2 \int d^2\vec{k}_{aT} d^2\vec{k}_{bT} \delta^2(\vec{q}_T - \vec{k}_{aT} - \vec{k}_{bT}) (v_q^2 + a_q^2) \\
& \times [f_1^q(x_a, \vec{k}_{aT}^2) f_1^{\bar{q}'}(x_b, \vec{k}_{bT}^2) + f_1^{\bar{q}'}(x_a, \vec{k}_{aT}^2) f_1^q(x_b, \vec{k}_{bT}^2)], \quad (22)
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}^W\left[2v_q a_q \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} g_{1T} \bar{f}_1\right] = & \sum_{q,q'} |V_{qq'}|^2 \int d^2\vec{k}_{aT} d^2\vec{k}_{bT} \delta^2(\vec{q}_T - \vec{k}_{aT} - \vec{k}_{bT}) 2v_q a_q \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} \\
& \times [g_{1T}^q(x_a, \vec{k}_{aT}^2) f_1^{\bar{q}'}(x_b, \vec{k}_{bT}^2) - g_{1T}^{\bar{q}'}(x_a, \vec{k}_{aT}^2) f_1^q(x_b, \vec{k}_{bT}^2)], \quad (23)
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}^W\left[(v_q^2 + a_q^2) \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} g_{1T} \bar{g}_{1L}\right] = & \sum_{q,q'} |V_{qq'}|^2 \int d^2\vec{k}_{aT} d^2\vec{k}_{bT} \delta^2(\vec{q}_T - \vec{k}_{aT} - \vec{k}_{bT}) (v_q^2 + a_q^2) \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} \\
& \times [g_{1T}^q(x_a, \vec{k}_{aT}^2) g_{1L}^{\bar{q}'}(x_b, \vec{k}_{bT}^2) + g_{1T}^{\bar{q}'}(x_a, \vec{k}_{aT}^2) g_{1L}^q(x_b, \vec{k}_{bT}^2)]. \quad (24)
\end{aligned}$$

Here the first and the third equations, Eqs. (22) and (24), do not have a minus sign under $q \leftrightarrow \bar{q}'$. This is because they either do not involve any g_{1T} or g_{1L} at all as in Eq. (22), or they involve even total number of g_{1T} and g_{1L} as in Eq. (24). On the other hand, since Eq. (23) involves an odd total number of g_{1T} or g_{1L} , there is a minus sign under $q \leftrightarrow \bar{q}'$.

Using this shorthand notation and defining $\hat{q}_T = \vec{q}_T/q_T$, we have

$$F_{UU} = \mathcal{C}^W[(v_q^2 + a_q^2)f_1\bar{f}_1], \quad (25)$$

$$F_{LU} = -\mathcal{C}^W[2v_q a_q g_{1L} \bar{f}_1], \quad (26)$$

$$F_{UL} = \mathcal{C}^W[2v_q a_q f_1 \bar{g}_{1L}], \quad (27)$$

$$F_{LL} = -\mathcal{C}^W[(v_q^2 + a_q^2)g_{1L} \bar{g}_{1L}], \quad (28)$$

$$F_{TU}^{\sin(\phi_V - \phi_{S_A})} = C^W \left[(v_q^2 + a_q^2) \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} f_{1T}^\perp \bar{f}_1 \right], \quad (29)$$

$$F_{TU}^{\cos(\phi_V - \phi_{S_A})} = -C^W \left[2v_q a_q \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} g_{1T} \bar{f}_1 \right], \quad (30)$$

$$F_{UT}^{\sin(\phi_V - \phi_{S_B})} = -C^W \left[(v_q^2 + a_q^2) \frac{\hat{q}_T \cdot \vec{k}_{bT}}{M_B} f_{1T}^\perp \bar{f}_1 \right], \quad (31)$$

$$F_{UT}^{\cos(\phi_V - \phi_{S_B})} = C^W \left[2v_q a_q \frac{\hat{q}_T \cdot \vec{k}_{bT}}{M_B} f_{1T} \bar{g}_{1T} \right], \quad (32)$$

$$F_{TL}^{\sin(\phi_V - \phi_{S_A})} = C^W \left[2v_q a_q \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} f_{1T}^\perp \bar{g}_{1L} \right], \quad (33)$$

$$F_{TL}^{\cos(\phi_V - \phi_{S_A})} = -C^W \left[(v_q^2 + a_q^2) \frac{\hat{q}_T \cdot \vec{k}_{aT}}{M_A} g_{1T} \bar{g}_{1L} \right], \quad (34)$$

$$F_{LT}^{\sin(\phi_V - \phi_{S_B})} = C^W \left[2v_q a_q \frac{\hat{q}_T \cdot \vec{k}_{bT}}{M_B} g_{1L} \bar{f}_{1T}^\perp \right], \quad (35)$$

$$F_{LT}^{\cos(\phi_V - \phi_{S_B})} = -C^W \left[(v_q^2 + a_q^2) \frac{\hat{q}_T \cdot \vec{k}_{bT}}{M_B} g_{1L} \bar{g}_{1T} \right], \quad (36)$$

$$F_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})} = C^W \left[(v_q^2 + a_q^2) \frac{2\vec{k}_{aT} \cdot \hat{q}_T \vec{k}_{bT} \cdot \hat{q}_T - \vec{k}_{aT} \cdot \vec{k}_{bT}}{2M_A M_B} \right. \\ \left. \times (f_{1T}^\perp \bar{f}_{1T}^\perp - g_{1T} \bar{g}_{1T}) \right], \quad (37)$$

$$F_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})} = C^W \left[v_q a_q \frac{2\vec{k}_{aT} \cdot \hat{q}_T \vec{k}_{bT} \cdot \hat{q}_T - \vec{k}_{aT} \cdot \vec{k}_{bT}}{M_A M_B} \right. \\ \left. \times (f_{1T}^\perp \bar{g}_{1T} + g_{1T} \bar{f}_{1T}^\perp) \right], \quad (38)$$

$$F_{TT}^1 = -C^W \left[(v_q^2 + a_q^2) \frac{\vec{k}_{aT} \cdot \vec{k}_{bT}}{2M_A M_B} (f_{1T}^\perp \bar{f}_{1T}^\perp + g_{1T} \bar{g}_{1T}) \right], \quad (39)$$

$$F_{TT}^2 = -C^W \left[v_q a_q \frac{\vec{k}_{aT} \cdot \vec{k}_{bT}}{M_A M_B} (f_{1T}^\perp \bar{g}_{1T} - g_{1T} \bar{f}_{1T}^\perp) \right]. \quad (40)$$

To obtain the differential cross section for Z boson production in polarized proton-proton collisions, one simply replaces σ_0^W by σ_0^Z in Eq. (19) with

$$\sigma_0^Z = \frac{\sqrt{2}\pi G_F M_Z^2}{S N_c}. \quad (41)$$

All the structure functions will be given by the same expressions as above, with C^W in Eq. (20) replaced by C^Z :

$$C^Z[w(\vec{k}_{aT}, \vec{k}_{bT}) f_1 \bar{f}_2] \\ = \sum_q \int d^2 \vec{k}_{aT} d^2 \vec{k}_{bT} \delta^2(\vec{q}_T - \vec{k}_{aT} - \vec{k}_{bT}) w(\vec{k}_{aT}, \vec{k}_{bT}) \\ \times [f_1^q(x_a, \vec{k}_{aT}^2) f_2^{\bar{q}}(x_b, \vec{k}_{bT}^2) + (q \leftrightarrow \bar{q})]. \quad (42)$$

At the same time, for Z bosons we have

$$v_q = T_q^3 - 2e_q \sin^2 \theta_W, \quad a_q = T_q^3, \quad (43)$$

where e_q is the quark electric charge and T_q^3 is the third component of “weak isospin” of the quark. $T_q^3 = \frac{1}{2}$ for u, c, t quarks, while $T_q^3 = -\frac{1}{2}$ for d, s, b quarks.

Furthermore, we could also write down the differential cross section for the Drell-Yan lepton pair production through the virtual photon decay, which gives

$$\frac{d\sigma^{\text{DY}}}{dQ^2 dy d^2 \vec{q}_T} = \sigma_0^{\text{DY}} \{ F_{UU} + S_{AL} S_{BL} F_{LL} + |\vec{S}_{AT}| [\sin(\phi_V - \phi_{S_A}) F_{TU}^{\sin(\phi_V - \phi_{S_A})}] + |\vec{S}_{BT}| [\sin(\phi_V - \phi_{S_B}) F_{UT}^{\sin(\phi_V - \phi_{S_B})}] \\ + |\vec{S}_{AT}| S_{BL} [\cos(\phi_V - \phi_{S_A}) F_{TL}^{\cos(\phi_V - \phi_{S_A})}] + S_{AL} |\vec{S}_{BT}| [\cos(\phi_V - \phi_{S_B}) F_{LT}^{\cos(\phi_V - \phi_{S_B})}] \\ + |\vec{S}_{AT}| |\vec{S}_{BT}| [\cos(2\phi_V - \phi_{S_A} - \phi_{S_B}) F_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})} + \cos(\phi_{S_A} - \phi_{S_B}) F_{TT}^1] \}. \quad (44)$$

Here σ_0^{DY} is given by [24]

$$\sigma_0^{\text{DY}} = \frac{4\pi\alpha_{\text{em}}^2}{3SQ^2 N_c}, \quad (45)$$

where Q is the invariant mass of the lepton pair, and α_{em} is the electromagnetic coupling constant. At the same time, all the structure functions can be obtained from those for Z

boson production [including the convolution expression in Eq. (42)] but with the replacement

$$v_q = e_q, \quad a_q = 0. \quad (46)$$

Several comments are in order.

- (i) In our setup, for Drell-Yan production through the virtual photon channel (parity conserving channel)

only the Sivers effects survive in the single transverse spin asymmetry: as can be clearly seen from Eq. (44), they are related to $\sin(\phi_V - \phi_S)$ modulation. This is because we integrate out the full kinematics of the decayed lepton pair, and only measure the azimuthal angle of the virtual photon. If instead one further measures the kinematics of the decayed lepton pair, e.g., measure both polar and azimuthal angles in the so-called Collins-Soper frame [52], one could have additional terms, such as the product of quark transversity and Boer-Mulder functions [53]. For complete results in this case, see Ref. [48], as well as Ref. [54] where Z contribution is also included.

- (ii) For W^\pm/Z^0 production, with parity-violating interactions there are two azimuthal modulation terms which can contribute to the single transverse spin asymmetry, as can be seen from Eq. (19). Besides the term related to $\sin(\phi_V - \phi_S)$ modulation just like in the Drell-Yan dilepton production, there is another term related to $\cos(\phi_V - \phi_S)$ modulation. The $\sin(\phi_V - \phi_S)$ term is associated with the quark Sivers function $f_{1T}^{\perp q}$, and it is the usual Sivers effect. On the other hand, the $\cos(\phi_V - \phi_S)$ term is associated with the transversal helicity g_{1T}^q , which projects out the longitudinal quark distribution inside a transversely polarized proton. As in Eqs. (30) and (32), the relevant structure functions, $F_{TU}^{\cos(\phi_V - \phi_{S_A})}$ and $F_{UT}^{\cos(\phi_V - \phi_{S_B})}$, are directly proportional to a_q , and thus only exist when there is parity-violating interactions, i.e. they are unique to W^\pm/Z^0 production. The amplitudes of both $\sin(\phi_V - \phi_S)$ - and $\cos(\phi_V - \phi_S)$ modulations can be comparable, as they involve only one spin-dependent parton distribution. Therefore, in the experimental study of the single transverse spin asymmetries for W^\pm/Z^0 production, one should consider extracting both terms simultaneously to avoid cross contamination, e.g. by following a “maximum-likelihood fit” method widely used in the extraction of azimuthal spin asymmetries in the SIDIS process [41,55,56].
- (iii) There are terms contributing to the double transverse spin asymmetries. In particular, $F_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})}$ receives contributions from the product of the Sivers functions $f_{1T}^{\perp q} f_{1T}^{\perp \bar{q}}$, as well as the product of transversal helicity distribution $g_{1T}^q g_{1T}^{\bar{q}}$. They were referred to in Ref. [33] as double Sivers and double worm-gear effects, respectively. As pointed out already in [33], all these double transverse spin asymmetries do not involve quark transversity, contrary to the collinear factorization picture with q_T integrated, in which such asymmetries usually involve the contribution from the collinear quark transversity [28–31].

- (iv) For identical nucleon scattering, e.g., proton-proton collisions at RHIC, under the exchange of the rapidity of the vector boson $y \leftrightarrow -y$, we have the following relations:

$$\begin{aligned} F_{UU}(y) &= F_{UU}(-y), & F_{LL}(y) &= F_{LL}(-y), \\ F_{LU}(y) &= F_{UL}(-y), \end{aligned} \quad (47)$$

$$\begin{aligned} F_{TU}^{\sin(\phi_V - \phi_{S_A})}(y) &= -F_{UT}^{\sin(\phi_V - \phi_{S_B})}(-y), \\ F_{TU}^{\cos(\phi_V - \phi_{S_A})}(y) &= F_{UT}^{\cos(\phi_V - \phi_{S_B})}(-y), \end{aligned} \quad (48)$$

$$\begin{aligned} F_{TL}^{\sin(\phi_V - \phi_{S_A})}(y) &= -F_{LT}^{\sin(\phi_V - \phi_{S_B})}(-y), \\ F_{TL}^{\cos(\phi_V - \phi_{S_A})}(y) &= F_{LT}^{\cos(\phi_V - \phi_{S_B})}(-y), \end{aligned} \quad (49)$$

$$\begin{aligned} F_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})}(y) &= F_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})}(-y), \\ F_{TT}^1(y) &= F_{TT}^1(-y), \end{aligned} \quad (50)$$

$$\begin{aligned} F_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})}(y) &= -F_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})}(-y), \\ F_{TT}^2(y) &= -F_{TT}^2(-y). \end{aligned} \quad (51)$$

III. PHENOMENOLOGY AT THE RHIC ENERGY

In this section, we move on to discuss numerical estimates for the magnitude of the spin asymmetries in polarized proton-proton collisions at the top RHIC energy.

A. Definitions and parametrizations

To start, we first define various spin asymmetries for the vector boson production in polarized proton-proton collisions at RHIC. Longitudinal spin asymmetries do not involve any azimuthal angle dependence, and we define the single longitudinal spin asymmetry A_{LU} , A_{UL} and double longitudinal spin asymmetry A_{LL} as

$$A_{LU} = \frac{F_{LU}}{F_{UU}}, \quad A_{UL} = \frac{F_{UL}}{F_{UU}}, \quad A_{LL} = \frac{F_{LL}}{F_{UU}}. \quad (52)$$

All other spin asymmetries involve transverse spin of the incoming protons, and they can be defined similarly as follows:

$$A_{TU}^{\sin(\phi_V - \phi_{S_A})} = \frac{F_{TU}^{\sin(\phi_V - \phi_{S_A})}}{F_{UU}}, \quad A_{TU}^{\cos(\phi_V - \phi_{S_A})} = \frac{F_{TU}^{\cos(\phi_V - \phi_{S_A})}}{F_{UU}}, \quad (53)$$

$$A_{TL}^{\sin(\phi_V - \phi_{S_A})} = \frac{F_{TL}^{\sin(\phi_V - \phi_{S_A})}}{F_{UU}}, \quad A_{TL}^{\cos(\phi_V - \phi_{S_A})} = \frac{F_{TL}^{\cos(\phi_V - \phi_{S_A})}}{F_{UU}}, \quad (54)$$

$$A_{TT}^{\sin(2\phi_V-\phi_{S_A}-\phi_{S_B})} = \frac{F_{TT}^{\sin(2\phi_V-\phi_{S_A}-\phi_{S_B})}}{F_{UU}},$$

$$A_{TT}^{\cos(2\phi_V-\phi_{S_A}-\phi_{S_B})} = \frac{F_{TT}^{\cos(2\phi_V-\phi_{S_A}-\phi_{S_B})}}{F_{UU}}, \quad (55)$$

$$A_{TT}^1 = \frac{F_{TT}^1}{F_{UU}}, \quad A_{TT}^2 = \frac{F_{TT}^2}{F_{UU}}. \quad (56)$$

Likewise, we define $A_{UT}^{\sin(\phi_V-\phi_{S_B})}$, $A_{UT}^{\cos(\phi_V-\phi_{S_B})}$, $A_{LT}^{\sin(\phi_V-\phi_{S_B})}$, and $A_{LT}^{\cos(\phi_V-\phi_{S_B})}$ as the ratios of the corresponding structure functions to the unpolarized structure function F_{UU} , respectively. Following the relations in Eqs. (47)–(51), for the polarized proton-proton collisions, we have

$$A_{LU}(y) = A_{UL}(-y), \quad A_{LL}(y) = A_{LL}(-y), \quad (57)$$

$$A_{TU}^{\sin(\phi_V-\phi_{S_A})}(y) = -A_{UT}^{\sin(\phi_V-\phi_{S_B})}(-y),$$

$$A_{TU}^{\cos(\phi_V-\phi_{S_A})}(y) = A_{UT}^{\cos(\phi_V-\phi_{S_B})}(-y), \quad (58)$$

$$A_{TL}^{\sin(\phi_V-\phi_{S_A})}(y) = -A_{LT}^{\sin(\phi_V-\phi_{S_B})}(-y),$$

$$A_{TL}^{\cos(\phi_V-\phi_{S_A})}(y) = A_{LT}^{\cos(\phi_V-\phi_{S_B})}(-y). \quad (59)$$

$$A_{TT}^{\cos(2\phi_V-\phi_{S_A}-\phi_{S_B})}(y) = A_{TT}^{\cos(2\phi_V-\phi_{S_A}-\phi_{S_B})}(-y),$$

$$A_{TT}^1(y) = A_{TT}^1(-y), \quad (60)$$

$$A_{TT}^{\sin(2\phi_V-\phi_{S_A}-\phi_{S_B})}(y) = -A_{TT}^{\sin(2\phi_V-\phi_{S_A}-\phi_{S_B})}(-y),$$

$$A_{TT}^2(y) = -A_{TT}^2(-y). \quad (61)$$

Looking at the expressions for all the structure functions given in last section, we find that A_{LL} , $A_{TU}^{\sin(\phi_V-\phi_{S_A})}$, $A_{UT}^{\sin(\phi_V-\phi_{S_B})}$, $A_{TL}^{\cos(\phi_V-\phi_{S_A})}$, $A_{LT}^{\cos(\phi_V-\phi_{S_B})}$, $A_{TT}^{\cos(2\phi_V-\phi_{S_A}-\phi_{S_B})}$, and A_{TT}^1 exist for both W^\pm/Z^0 and γ^* , and they are parity-even spin asymmetries. On the other hand, A_{LU} , A_{UL} , $A_{TU}^{\cos(\phi_V-\phi_{S_A})}$, $A_{UT}^{\cos(\phi_V-\phi_{S_B})}$, $A_{TL}^{\sin(\phi_V-\phi_{S_A})}$, $A_{LT}^{\sin(\phi_V-\phi_{S_B})}$, $A_{TT}^{\sin(2\phi_V-\phi_{S_A}-\phi_{S_B})}$, and A_{TT}^2 are directly proportional to a_q (the axial coupling), and thus they exist because of the parity-violating nature of the weak interaction, i.e. they are all parity-odd spin asymmetries.

Single transverse spin asymmetries have also been denoted as A_N [16,17,23,57,58] in the literature, where usually one chooses a frame such that the transversely polarized proton A moves in the $+z$ -direction, and the spin vector \vec{S}_{AT} and the transverse momentum \vec{q}_T are along the y and x directions, respectively. It is important to realize that such definition of A_N is related to our definition of $A_{TU}^{\sin(\phi_V-\phi_{S_A})}$ by a minus sign [17]:

$$A_{TU}^{\sin(\phi_V-\phi_{S_A})} = -A_N. \quad (62)$$

It is worth mentioning that A_N (and thus $A_{TU}^{\sin(\phi_V-\phi_{S_A})}$) is usually referred to as “left-right” spin asymmetry [57,59–63]. In this sense, one could refer to the other single transverse spin asymmetry $A_{TU}^{\cos(\phi_V-\phi_{S_A})}$ as “up-down” spin asymmetry.

In this section we will present some numerical estimates for these spin asymmetries defined above. These spin asymmetries are all expressed in terms of various TMDs, as shown in the last section. In principle, to make precise quantitative predictions, one has to take into account the effect of the TMD evolution. The TMD evolution will likely lead to suppression of these spin asymmetries; see e.g. Refs. [24,64–70]. However, at this point the phenomenological implementation of TMD evolution formalism still has very large uncertainties. In fact, one of the motivations for the transverse spin asymmetry measurements of the vector boson production in polarized proton-proton collisions at RHIC is to constrain the TMD evolution formalism. Because of this, we will only present the numerical estimate of the spin asymmetries using the usual Gaussian model for all the quark TMDs, i.e., without TMD evolution. Since the TMD evolution is supposed to apply to all the TMDs roughly equally, the hope is that the relative magnitude of the various spin asymmetries could serve as a reasonable guidance for the experimental measurements.

Both the unpolarized quark TMD f_1^q and the helicity TMD distribution g_{1L}^q are k_T -even functions and we assume they have the following Gaussian forms:

$$f_1^q(x, k_T^2) = f_1^q(x) \frac{1}{\pi \langle k_T^2 \rangle_{f_1}} e^{-k_T^2 / \langle k_T^2 \rangle_{f_1}},$$

$$g_{1L}^q(x, k_T^2) = g_{1L}^q(x) \frac{1}{\pi \langle k_T^2 \rangle_{g_{1L}}} e^{-k_T^2 / \langle k_T^2 \rangle_{g_{1L}}}, \quad (63)$$

where $f_1^q(x)$ and $g_{1L}^q(x)$ are the collinear unpolarized parton distribution function and helicity distribution function, respectively. We will assume the Gaussian width for f_1^q and g_{1L}^q to be the same [71,72], and take the value of 0.25 GeV² [73]:

$$\langle k_T^2 \rangle_{f_1} = \langle k_T^2 \rangle_{g_{1L}} = 0.25 \text{ GeV}^2. \quad (64)$$

On the other hand, both the Sivers function $f_{1T}^{\perp q}$ and the transversal helicity distribution g_{1T}^q are the coefficients from the linear k_T -expansion term of the quark-quark correlator Φ^q ; see Eqs. (13) and (14). Quark Sivers functions have been extracted from semi-inclusive deep inelastic scattering. We take the parametrization from Ref. [22]

$$\frac{k_T}{M} f_{1T}^{\perp q}(x, k_T^2) = -\mathcal{N}_q(x) h(k_T) f_1^q(x, k_T^2), \quad (65)$$

where $\mathcal{N}_q(x)$ and $h(k_T)$ are given by

$$\mathcal{N}_q(x) = N_q x^{\alpha_q} (1-x)^{\beta_q} \frac{(\alpha_q + \beta_q)^{(\alpha_q + \beta_q)}}{\alpha_q^{\alpha_q} \beta_q^{\beta_q}},$$

$$h(k_T) = \sqrt{2e} \frac{k_T}{M_1} e^{-k_T^2/M_1^2}, \quad (66)$$

with the parameters N_q, α_q, β_q , and M_1 are given in [22]. At the same time, for g_{1T}^q , we have

$$\frac{1}{2M^2} g_{1T}^q(x, k_T^2) = g_{1T}^{q(1)}(x) \frac{1}{\pi \langle k_T^2 \rangle_{g_{1T}}^2} e^{-k_T^2 / \langle k_T^2 \rangle_{g_{1T}}}, \quad (67)$$

where we choose $\langle k_T^2 \rangle_{g_{1T}} = 0.15 \text{ GeV}^2$ [74], and $g_{1T}^{q(1)}(x)$ is the first k_T moment of $g_{1T}^q(x, k_T^2)$ defined as

$$g_{1T}^{q(1)}(x) = \int d^2 k_T \frac{\vec{k}_T^2}{2M^2} g_{1T}^q(x, k_T^2). \quad (68)$$

We note that there is a Wandzura-Wilczek-type approximation for $g_{1T}^{q(1)}(x)$, which relates $g_{1T}^{q(1)}(x)$ to the collinear helicity distribution function $g_{1L}^q(x)$ [75]:

$$g_{1T}^{q(1)}(x) \approx x \int_x^1 \frac{dz}{z} g_{1L}^q(z). \quad (69)$$

Now the parametrizations for all the TMDs involved in our calculations, $f_1^q, f_{1T}^{\perp q}, g_{1L}^q$, and g_{1T}^q , are given. One last thing one should keep in mind is that the quark Siverson function changes sign when probed in SIDIS and DY processes [16,18–21]:

$$f_{1T}^{\perp q}(x, k_T^2)|_{\text{DY/W/Z}} = -f_{1T}^{\perp q}(x, k_T^2)|_{\text{SIDIS}}, \quad (70)$$

whereas all other quark TMDs ($f_1^q, g_{1L}^q, g_{1T}^q$) are universal in SIDIS and Drell-Yan. As we have mentioned already in the Introduction, since $g_{1T}^q(x, k_T^2)$ was investigated in SIDIS measurements [39–42] and is proposed to be measured with high precision in future SIDIS experiments [43], one could in principle test the universality of $g_{1T}^q(x, k_T^2)$ in W^\pm/Z^0 production in proton-proton collisions at the RHIC. In our numerical studies below, we implement the sign change for the quark Siverson function and keep all other TMDs the same as in SIDIS when we present the asymmetries of W^\pm/Z^0 boson production at RHIC, to which we now turn.

B. Numerical estimate for the spin asymmetries

We now present the numerical estimates for the spin asymmetries. We use CTEQ6 [76] for the collinear

unpolarized parton distribution functions $f_1^q(x)$, DSSV [77,78] parametrization for the collinear helicity distribution functions $g_{1L}^q(x)$, and choose the factorization scale $\mu = M_V$. Within our approximation, where $\langle k_T^2 \rangle_{f_1} = \langle k_T^2 \rangle_{g_{1L}}$, there is no q_T dependence for the longitudinal spin asymmetries A_{LU}, A_{UL}, A_{LL} . This is because F_{LU}, F_{UL}, F_{LL} all have the same q_T dependence as the unpolarized structure function F_{UU} , which then cancels out in the longitudinal spin asymmetries. In this case, these longitudinal spin asymmetries will be the same as those for the inclusive (q_T -integrated) vector boson production, which are available in the literature; see e.g. Refs. [3–5,7,79,80]. We will, thus, not present numerical estimates for these longitudinal spin asymmetries here. Instead, we will focus on those spin asymmetries which involve transversely polarized proton in the collisions, where one has to measure the azimuthal angle of the vector boson in order to probe these asymmetries. In particular, we will present the numerical results for $A_{TU}^{\sin(\phi_V - \phi_{S_A})}, A_{TU}^{\cos(\phi_V - \phi_{S_A})}, A_{TL}^{\sin(\phi_V - \phi_{S_A})}, A_{TL}^{\cos(\phi_V - \phi_{S_A})}, A_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})}$, and $A_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})}$. The spin asymmetries involving the transverse spin vector \vec{S}_{BT} (i.e. ϕ_{S_B}) can be obtained from the results we present through the relations established in Eqs. (58) and (59).

Let us first discuss the single transverse spin asymmetry: $A_{TU}^{\sin(\phi_V - \phi_{S_A})}$ and $A_{TU}^{\cos(\phi_V - \phi_{S_A})}$. In Fig. 1, we plot single transverse spin asymmetry $A_{TU}^{\sin(\phi_V - \phi_{S_A})}$ as a function of the rapidity y of the vector boson (left), and as a function of the transverse momentum q_T of the vector boson at rapidity $y = 0.5$ (right) at the RHIC energy $\sqrt{s} = 510 \text{ GeV}$. In the left plot, we have integrated the vector boson transverse momentum in the region $0 < q_T < 3 \text{ GeV}$. The red solid curve is for W^+ , the blue dashed curve is for W^- , and the black dotted curve is for Z^0 production. These results are consistent with those in [16] and [17], respectively.¹ It is worthwhile to remind the reader that $A_{TU}^{\sin(\phi_V - \phi_{S_A})}$ is parity even, and can be used to probe the quark Siverson function $f_{1T}^{\perp q}(x, k_T^2)$. Measuring such a transverse spin asymmetry and testing the sign change of the quark Siverson function is one of the main goals of the transverse W program in the near future at RHIC [2]. One should also keep in mind that because the sea quark Siverson functions are not really constrained from the fixed-target SIDIS measurements, our theoretical curves could have very large uncertainties [47], especially in the backward rapidity region where the asymmetry is most sensitive to the sea quark distributions.

On the other hand, due to the parity-violating interaction, there is another single transverse spin asymmetry, $A_{TU}^{\cos(\phi_V - \phi_{S_A})}$ (parity odd). This term is related to $g_{1T}^q(x, k_T^2)$.

¹Of course one has to keep in mind that A_N were plotted in [16] and [17], while here $A_{TU}^{\sin(\phi_V - \phi_{S_A})}$ are plotted. So they should have an opposite sign because of Eq. (62).




FIG. 1. Single transverse spin asymmetry $A_{TU}^{\sin(\phi_V - \phi_{SA})}$ as a function of the rapidity y of the vector boson (left), and as a function of the transverse momentum q_T of the vector boson at rapidity $y = 0.5$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV. In the left plot, we have integrated vector boson transverse momentum in the region $0 < q_T < 3$ GeV. The red solid curve is for W^+ , the blue dashed curve is for W^- , and the black dotted curve is for Z^0 production. Note: $A_{TU}^{\sin(\phi_V - \phi_{SA})}$ is related to the parity-conserving interaction (parity-even), and can be used to probe the quark Sivers function $f_{1T}^{\perp q}(x, k_T^2)$.

Since g_{1T}^q is not fully constrained even for the valence quarks within the current SIDIS measurements [39–42], $A_{TU}^{\cos(\phi_V - \phi_{SA})}$ of W^\pm/Z^0 production at RHIC could on one hand serve as a complementary channel to constrain g_{1T}^q , at the same time, as we have emphasized, one in principle could test the universality of g_{1T}^q as to the future high precision SIDIS measurement [25,43]. In Fig. 2, we plot $A_{TU}^{\cos(\phi_V - \phi_{SA})}$ as a function of the rapidity y of the vector boson (left), and as a function of the transverse momentum q_T of the vector boson at rapidity $y = 1$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV. The asymmetry is sizable, in particular due to the fact that W^\pm production provides

maximum analyzing power for the quark longitudinal polarization. If TMD evolution only leads to a moderate suppression, this asymmetry should be measurable at the RHIC.

Let us now turn to the study of the transverse-longitudinal double spin asymmetry $A_{TL}^{\cos(\phi_V - \phi_{SA})}$ and $A_{TL}^{\sin(\phi_V - \phi_{SA})}$. While $A_{TL}^{\cos(\phi_V - \phi_{SA})}$ is related to the parity-conserving interaction and is sensitive to both g_{1T}^q and g_{1L}^q , $A_{TL}^{\sin(\phi_V - \phi_{SA})}$ is related to the parity-violating interaction and is sensitive to $f_{1T}^{\perp q}$ and g_{1L}^q . In Fig. 3, we plot the transverse-longitudinal double spin asymmetries $A_{TL}^{\cos(\phi_V - \phi_{SA})}$ as a function of rapidity y (left) and as a




FIG. 2. Single transverse spin asymmetry $A_{TU}^{\cos(\phi_V - \phi_{SA})}$ as a function of the rapidity y of the vector boson (left), and as a function of the transverse momentum q_T of the vector boson at rapidity $y = 1$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV. In the left plot, we have integrated vector boson transverse momentum in the range $0 < q_T < 3$ GeV. The red solid curve is for W^+ , the blue dashed curve is for W^- , and the black dotted curve is for Z^0 production. Note: $A_{TU}^{\cos(\phi_V - \phi_{SA})}$ is related to the parity-violating interaction (parity-odd), and can be used to probe the transversal helicity distribution $g_{1T}^q(x, k_T^2)$.




FIG. 3. Transverse-longitudinal double spin asymmetries $A_{TL}^{\cos(\phi_V - \phi_{SA})}$ as a function of rapidity y (left) and as a function of transverse momentum q_T of the vector boson at rapidity $y = 0.7$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV.




FIG. 4. Transverse-longitudinal spin asymmetries double spin asymmetries $A_{TL}^{\sin(\phi_V - \phi_{SA})}$ as a function of rapidity y (left) and as a function of transverse momentum q_T of the vector boson at rapidity $y = -0.7$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV.

function of the transverse momentum q_T of the vector boson at rapidity $y = 0.7$ (right). Since both g_{1T}^q and g_{1L}^q are suppressed when compared with the unpolarized parton distribution f_1^q , the double spin asymmetry is indeed much smaller than the single transverse spin asymmetry, where only one spin-dependent parton distribution (either $f_{1T}^{\perp q}$ or g_{1T}^q) is involved. In Fig. 4, we plot $A_{TL}^{\sin(\phi_V - \phi_{SA})}$ as a function of rapidity y (left) and as a function of transverse momentum q_T of the vector boson at rapidity $y = -0.7$ (right) at the top RHIC energy. Even though the asymmetry is in general not very large, $A_{TL}^{\sin(\phi_V - \phi_{SA})}$ for W^+ is quite sizable, $\sim 5\%$ – 10% . The reason lies in the fact that the Siverson function $f_{1T}^{\perp q}$ from Ref. [22] for \bar{d} quark is still sizable and g_{1L}^q for u quark is reasonably large. Thus, their product $f_{1T}^{\perp \bar{d}} g_{1L}^u$ leads to a large double spin asymmetry.

Finally, we study the double transverse spin asymmetries. As examples, we present the numerical results for

$A_{TT}^{\cos(2\phi_V - \phi_{SA} - \phi_{SB})}$ and $A_{TT}^{\sin(2\phi_V - \phi_{SA} - \phi_{SB})}$, which still involve the azimuthal angle ϕ_V of the vector boson (the other two asymmetries A_{TT}^1 and A_{TT}^2 do not). In Fig. 5 we plot $A_{TT}^{\cos(2\phi_V - \phi_{SA} - \phi_{SB})}$ as a function of rapidity y (left) and as a function of transverse momentum q_T of the vector boson at rapidity $y = 0.7$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV. Only the W^+ and the Z^0 bosons have reasonable large asymmetries $\lesssim 5\%$ but with opposite sign. This can be understood as follows. We have $f_{1T}^{\perp u} f_{1T}^{\perp \bar{d}}$ contributing to W^+ while $f_{1T}^{\perp d} f_{1T}^{\perp \bar{u}}$ to the Z boson. From the Siverson function parametrization we used [22], the u and d quark Siverson distributions have the opposite sign. Together with the fact that $f_{1T}^{\perp \bar{d}}$ is still sizable in this parametrization, this leads to opposite but reasonably large asymmetry for W^+ and Z^0 bosons. In Fig. 6 we plot $A_{TT}^{\sin(2\phi_V - \phi_{SA} - \phi_{SB})}$ as a function of rapidity y (left) and as




FIG. 5. Double transverse spin asymmetry $A_{TT}^{\cos(2\phi_V - \phi_{S_A} - \phi_{S_B})}$ as a function of rapidity y (left) and as a function of transverse momentum q_T of the vector boson at rapidity $y = 0$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV.




FIG. 6. Double transverse spin asymmetry $A_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})}$ as a function of rapidity y (left) and as a function of transverse momentum q_T of the vector boson at rapidity $y = 0.7$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV.

function of transverse momentum q_T of the vector boson at rapidity $y = 0.7$ (right) at the RHIC energy $\sqrt{s} = 510$ GeV. The slightly larger asymmetries $A_{TT}^{\sin(2\phi_V - \phi_{S_A} - \phi_{S_B})}$ for W^+/Z^0 can be understood similarly.

IV. SUMMARY

In this paper we studied the spin-dependent differential cross sections for vector boson ($W^\pm/Z^0/\gamma^*$) production in polarized nucleon-nucleon collisions for low transverse momentum of the observed vector boson. We considered the situation where the full kinematics of the vector boson could be reconstructed and both the magnitude q_T and azimuthal angle ϕ_V of the vector bosons are measured. We presented the cross sections and the associated single and double spin asymmetries in terms of the transverse momentum dependent parton distribution functions (TMDs) at tree

level within the TMD factorization formalism. We estimated these spin asymmetries for W^\pm/Z^0 boson production in polarized proton-proton collisions at the top RHIC energy, and found that if the TMD evolution effect does not lead to too strong a suppression, some of the asymmetries are rather sizable and should be measurable at RHIC. In particular, the single transverse spin asymmetries contain two large orthogonal azimuthal terms: a parity-conserving term that is sensitive to the quark Sivers function f_{1T}^\perp , as well as a parity-violating term that probes the quark transversal helicity distribution g_{1T}^q . While f_{1T}^\perp is predicted to change sign from semi-inclusive deep inelastic scattering to the Drell-Yan process, g_{1T}^q is supposed to be universal. Thus, the W spin physics program at RHIC could be viewed as truly multipurpose: one that tests the universality properties of TMDs, constrains the TMD evolution effects, and probes the sea quark TMDs.

ACKNOWLEDGMENTS

We thank E. C. Aschenauer, A. Metz, D. Pitonyak, and M. Schlegel for helpful comments. This work is supported by

the U.S. Department of Energy under Contracts No. DE-SC0012704 (J. H.) and No. DE-AC52-06NA25396 (Z. K., I. V., and H. X.), and in part by the LDRD program at LANL.

-
- [1] E. C. Aschenauer *et al.*, [arXiv:1304.0079](#).
 - [2] E.-C. Aschenauer *et al.*, [arXiv:1501.01220](#).
 - [3] C. Bourrely and J. Soffer, *Phys. Lett. B* **314**, 132 (1993).
 - [4] C. Bourrely and J. Soffer, *Nucl. Phys.* **B423**, 329 (1994).
 - [5] D. de Florian and W. Vogelsang, *Phys. Rev. D* **81**, 094020 (2010).
 - [6] D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, *Prog. Part. Nucl. Phys.* **67**, 251 (2012).
 - [7] Z.-B. Kang and J. Soffer, *Phys. Rev. D* **83**, 114020 (2011).
 - [8] F. Ringer and W. Vogelsang, *Phys. Rev. D* **91**, 094033 (2015).
 - [9] M. M. Aggarwal *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **106**, 062002 (2011).
 - [10] A. Adare *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **106**, 062001 (2011).
 - [11] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **113**, 072301 (2014).
 - [12] A. Adare *et al.* (PHENIX Collaboration), [arXiv:1504.07451](#).
 - [13] C. A. Aidala, S. D. Bass, D. Hasch, and G. K. Mallot, *Rev. Mod. Phys.* **85**, 655 (2013).
 - [14] S. J. Brodsky, D. S. Hwang, and I. Schmidt, *Phys. Lett. B* **553**, 223 (2003).
 - [15] I. Schmidt and J. Soffer, *Phys. Lett. B* **563**, 179 (2003).
 - [16] Z.-B. Kang and J.-W. Qiu, *Phys. Rev. Lett.* **103**, 172001 (2009).
 - [17] Z.-B. Kang and J.-W. Qiu, *Phys. Rev. D* **81**, 054020 (2010).
 - [18] J. C. Collins, *Phys. Lett. B* **536**, 43 (2002).
 - [19] D. Boer, P. J. Mulders, and F. Pijlman, *Nucl. Phys.* **B667**, 201 (2003).
 - [20] S. J. Brodsky, D. S. Hwang, and I. Schmidt, *Phys. Lett. B* **530**, 99 (2002).
 - [21] S. J. Brodsky, D. S. Hwang, and I. Schmidt, *Nucl. Phys.* **B642**, 344 (2002).
 - [22] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, S. Melis, F. Murgia, A. Prokudin, and C. Türk, *Eur. Phys. J. A* **39**, 89 (2009).
 - [23] L. Gamberg, Z.-B. Kang, and A. Prokudin, *Phys. Rev. Lett.* **110**, 232301 (2013).
 - [24] M. G. Echevarria, A. Idilbi, Z.-B. Kang, and I. Vitev, *Phys. Rev. D* **89**, 074013 (2014).
 - [25] A. Accardi *et al.*, [arXiv:1212.1701](#).
 - [26] E. C. Aschenauer *et al.*, [arXiv:1409.1633](#).
 - [27] S. Abeyratne *et al.*, [arXiv:1504.07961](#).
 - [28] J. P. Ralston and D. E. Soper, *Nucl. Phys.* **B152**, 109 (1979).
 - [29] X. Artru and M. Mekhfi, *Z. Phys. C* **45**, 669 (1990).
 - [30] R. L. Jaffe and X.-D. Ji, *Nucl. Phys.* **B375**, 527 (1992).
 - [31] J. L. Cortes, B. Pire, and J. P. Ralston, *Z. Phys. C* **55**, 409 (1992).
 - [32] D. Boer, *Phys. Rev. D* **62**, 094029 (2000).
 - [33] D. Boer, W. J. den Dunnen, and A. Kotzinian, *Phys. Rev. D* **83**, 114032 (2011).
 - [34] A. Metz and J. Zhou, *Phys. Lett. B* **700**, 11 (2011).
 - [35] A. Metz, D. Pitonyak, A. Schaefer, and J. Zhou, *Phys. Rev. D* **86**, 114020 (2012).
 - [36] P. J. Mulders and R. D. Tangerman, *Nucl. Phys.* **B461**, 197 (1996); **B484**, 538(E) (1997).
 - [37] A. Bacchetta, M. Diehl, K. Goeke, A. Metz, P. J. Mulders, and M. Schlegel, *J. High Energy Phys.* **02** (2007) 093.
 - [38] D. Boer and P. J. Mulders, *Phys. Rev. D* **57**, 5780 (1998).
 - [39] B. Parsamyan (COMPASS Collaboration), *Eur. Phys. J. Spec. Top.* **162**, 89 (2008).
 - [40] L. L. Pappalardo and M. Diefenthaler (HERMES Collaboration), [arXiv:1107.4227](#).
 - [41] J. Huang *et al.* (The Jefferson Lab Hall A Collaboration), *Phys. Rev. Lett.* **108**, 052001 (2012).
 - [42] B. Parsamyan, [arXiv:1504.01599](#).
 - [43] H. Gao *et al.*, *Eur. Phys. J. Plus* **126**, 2 (2011).
 - [44] J. C. Collins and D. E. Soper, *Nucl. Phys.* **B193**, 381 (1981); **B213**, 545 (1983).
 - [45] X.-d. Ji, J.-p. Ma, and F. Yuan, *Phys. Rev. D* **71**, 034005 (2005).
 - [46] X.-d. Ji, J.-P. Ma, and F. Yuan, *Phys. Lett. B* **597**, 299 (2004).
 - [47] L. Adamczyk *et al.* (STAR Collaboration), [arXiv:1511.06003](#).
 - [48] S. Arnold, A. Metz, and M. Schlegel, *Phys. Rev. D* **79**, 034005 (2009).
 - [49] D. W. Sivers, *Phys. Rev. D* **41**, 83 (1990).
 - [50] D. W. Sivers, *Phys. Rev. D* **43**, 261 (1991).
 - [51] R. D. Tangerman and P. J. Mulders, *Phys. Rev. D* **51**, 3357 (1995).
 - [52] J. C. Collins and D. E. Soper, *Phys. Rev. D* **16**, 2219 (1977).
 - [53] D. Boer, *Phys. Rev. D* **60**, 014012 (1999).
 - [54] D. Pitonyak, M. Schlegel, and A. Metz, *Phys. Rev. D* **89**, 054032 (2014).
 - [55] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Rev. Lett.* **103**, 152002 (2009).
 - [56] M. G. Alekseev *et al.* (COMPASS Collaboration), *Phys. Lett. B* **692**, 240 (2010).
 - [57] Z.-B. Kang, J.-W. Qiu, W. Vogelsang, and F. Yuan, *Phys. Rev. D* **83**, 094001 (2011).
 - [58] Z.-B. Kang and A. Prokudin, *Phys. Rev. D* **85**, 074008 (2012).
 - [59] D. L. Adams *et al.* (FNAL-E704), *Phys. Lett. B* **264**, 462 (1991).
 - [60] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **92**, 171801 (2004).
 - [61] B. I. Abelev *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **101**, 222001 (2008).

- [62] L. Gamberg, Z.-B. Kang, A. Metz, D. Pitonyak, and A. Prokudin, *Phys. Rev. D* **90**, 074012 (2014).
- [63] M. G. Perdekamp and F. Yuan, *Annu. Rev. Nucl. Part. Sci.* **65**, 429 (2015).
- [64] D. Boer, *Nucl. Phys.* **B874**, 217 (2013).
- [65] D. Boer, *Nucl. Phys.* **B603**, 195 (2001).
- [66] S. M. Aybat, J. C. Collins, J.-W. Qiu, and T. C. Rogers, *Phys. Rev. D* **85**, 034043 (2012).
- [67] S. M. Aybat, A. Prokudin, and T. C. Rogers, *Phys. Rev. Lett.* **108**, 242003 (2012).
- [68] C. A. Aidala, B. Field, L. P. Gamberg, and T. C. Rogers, *Phys. Rev. D* **89**, 094002 (2014).
- [69] P. Sun and F. Yuan, *Phys. Rev. D* **88**, 114012 (2013).
- [70] M. Anselmino, M. Boglione, and S. Melis, *Phys. Rev. D* **86**, 014028 (2012).
- [71] Z.-B. Kang, A. Prokudin, P. Sun, and F. Yuan, *Phys. Rev. D* **93**, 014009 (2016).
- [72] Z.-B. Kang, A. Prokudin, P. Sun, and F. Yuan, *Phys. Rev. D* **91**, 071501 (2015).
- [73] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, and A. Prokudin, *Phys. Rev. D* **71**, 074006 (2005).
- [74] A. Kotzinian, B. Parsamyan, and A. Prokudin, *Phys. Rev. D* **73**, 114017 (2006).
- [75] A. Metz, P. Schweitzer, and T. Teckentrup, *Phys. Lett. B* **680**, 141 (2009).
- [76] J. Pumplin, D. R. Stump, J. Huston, H.-L. Lai, P. Nadolsky, and W.-K. Tung, *J. High Energy Phys.* **07** (2002) 012.
- [77] D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, *Phys. Rev. Lett.* **101**, 072001 (2008).
- [78] D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, *Phys. Rev. D* **80**, 034030 (2009).
- [79] T. Gehrmann, *Nucl. Phys.* **B534**, 21 (1998).
- [80] C. von Arx and T. Gehrmann, *Phys. Lett. B* **700**, 49 (2011).