

**Tetraquark production in double parton scattering**F. Carvalho,<sup>1</sup> E. R. Cazaroto,<sup>2</sup> V. P. Gonçalves,<sup>3,4</sup> and F. S. Navarra<sup>2</sup><sup>1</sup>*Departamento de Ciências Exatas e da Terra, Universidade Federal de São Paulo, Campus Diadema, Rua Prof. Artur Riedel, 275, Jd. Eldorado, 09972-270 Diadema, São Paulo, Brazil*<sup>2</sup>*Instituto de Física, Universidade de São Paulo, CEP 05315-970 São Paulo, São Paulo, Brazil*<sup>3</sup>*Department of Astronomy and Theoretical Physics, Lund University, 223-62 Lund, Sweden*<sup>4</sup>*Instituto de Física e Matemática, Universidade Federal de Pelotas, CEP 96010-900 Pelotas, Rio Grande do Sul, Brazil*

(Received 25 November 2015; published 2 February 2016)

We develop a model to study tetraquark production in hadronic collisions. We focus on double parton scattering and formulate a version of the color evaporation model for the production of the  $X(3872)$  and of the  $T_{4c}$  tetraquark, a state composed by the  $c\bar{c}c\bar{c}$  quarks. We find that the production cross section grows rapidly with the collision energy  $\sqrt{s}$  and make predictions for the forthcoming higher energy data of the LHC.

DOI: [10.1103/PhysRevD.93.034004](https://doi.org/10.1103/PhysRevD.93.034004)**I. INTRODUCTION****A. Production mechanism**

Over the last years, the existence of exotic hadrons has been firmly established [1,2], and now the next step is to determine their structure. Among the proposed configurations, the meson molecule and the tetraquark are the most often discussed. So far almost all the experimental information about these states comes from their production in B decays. The production of exotic particles in proton proton collisions is one of the most promising testing grounds for our ideas about the structure of the new states. It has been shown [1,3,4] that it is extremely difficult to produce molecules in pp collisions. In the molecular approach, the estimated cross section for  $X(3872)$  production is 2 orders of magnitude smaller than the measured one. The present challenge for theorists is to show that these data can be explained by the tetraquark model. To the best of our knowledge, this has not been done so far. In this work, we give a step in this direction, considering the production of the  $X(3872)$  and of the  $T_{4c}$ , a state composed by two charm quark pairs:  $c\bar{c}c\bar{c}$ .

In recent high energy collisions at the LHC, it became relatively easy to produce [5,6] four charm quarks ( $c\bar{c}c\bar{c}$ ) in the same event. Events with four heavy quarks can be treated as a particular case of  $\alpha_s^2$  correction to the standard single gluon-gluon scattering, in which an extra  $c\bar{c}$  pair is produced, i.e., the process  $gg \rightarrow c\bar{c}c\bar{c}$ . This is usually called single parton scattering (SPS). Another possible way to produce  $c\bar{c}c\bar{c}$  is by two independent leading order gluon-gluon scatterings, i.e., two times the reaction  $gg \rightarrow c\bar{c}$ . This is usually called double parton scattering (DPS) [7,8]. In fact, apart from  $c\bar{c}c\bar{c}$ , DPS events may generate many other different final states, such as four jets, a  $c - \bar{c}$  pair plus two jets, etc. For our purposes, the other relevant DPS process is the production of a  $c - \bar{c}$  pair plus a light quark pair,  $q - \bar{q}$ , which will hadronize and form the

$X(3872)$ . Since DPS events are in the realm of perturbative physics, the light quark pair must be produced with large invariant mass, and the final state  $X(3872)$  will carry large transverse momentum. This seems to be appropriate to describe the CMS data [9], where the  $X(3872)$  was observed with a transverse momentum lying in the range  $10 \leq p_T \leq 25$  GeV. In Refs. [10,11], it has been shown that DPS charm production is already comparable to SPS production at LHC energies. DPS grows faster with the energy because it is proportional to  $g(x, \mu^2)^4$ , while SPS is proportional to  $g(x, \mu^2)^2$ . Here,  $g(x, \mu^2)$  is the gluon density in the proton as a function of the gluon fractional momentum  $x$  and of the scale  $\mu$ , and it grows quickly with increasing collision energies. In the present work, we shall consider the DPS events with the production of the two  $c\bar{c}$  pairs and also with a  $c\bar{c}$  and a light quark  $q\bar{q}$  pair. We will study tetraquark production only at LHC energies, using the lower energy to fix the parameters and make predictions for the higher one. If we would try to extrapolate back to the Tevatron energies, we would enter the domain where SPS production becomes dominant. At the Tevatron energies, adding the SPS contribution to the DPS contribution would certainly make the cross section larger and closer to the available data. We refrain from including the SPS contribution because, in this case, it is more difficult to make a reliable model. We are aware that at all the energies considered we are missing the SPS contribution, which is important especially at lower energies. This introduces some uncertainties in our calculations. However, when the DPS becomes dominant, these uncertainties are reduced.

Once we have generated all the quarks and antiquarks needed to form the  $X(3872)$  or the  $T_{4c}$  in DPS events, we need to bind them together. To this end, we shall use the main ideas of the CEM [12,13] of charmonium production, where the  $c - \bar{c}$  is “kinematically bound”; i.e., the charm pair sticks together because it does not have an invariant mass large




FIG. 1. The gluons with odd (even) label come from the upper (lower) hadron and carry momentum fraction  $x_i$ . The “gluon 1” scatters with “gluon 2,” making the state  $M_{12}$ . An analogous process occurs with gluons 3 and 4. Finally,  $M_{12}$  and  $M_{34}$  merge and form the  $T_{4c}$  with mass  $M$ .

enough to produce anything else. We shall use the CEM ideas to study  $T_{4c}$  and  $X(3872)$  production in DPS events. In the CEM formalism, one parton from the hadron target scatters with one parton from the hadron projectile forming a charmonium state, which can absorb (emit) additional gluons from (to) the hadronic color field to become color neutral. The energy involved in this soft gluon exchange is of the order of  $\Lambda_{\text{QCD}}$ . Hence, this process is not suppressed by powers of  $\alpha_s$  and occurs with probability close to 1. This gluon exchange occurs in the hadronization stage, long after the initial, short-time, production stage. It has no memory of the initial collision energy, being energy independent. These assumptions lead to the introduction of a constant factor in the cross section, which is fixed at only one energy. The model keeps its predictive power.

The hadron formation mechanism of the CEM may be improved once we know better the tetraquark wave function and once we know how to project the intermediate partonic state into the final tetraquark state, with the right color and right spatial size.

At high energies, the gluon density in the proton is much bigger than the sea quark density, and hence, in what follows, we shall consider particle production only from gluon-gluon collisions. Now, we are going to extend the CEM to the case where two gluons from the hadron target scatter independently with two gluons from the hadron projectile as depicted in Fig. 1, where we show DPS production of  $T_{4c}$ . In the figure, two gluons collide and form a  $c\bar{c}$  state with mass  $M_{12}$ , while other two gluons collide and form a second  $c\bar{c}$  state with mass  $M_{34}$ . The two objects bind to each other forming the  $T_{4c}$ . Additional gluon exchanges with the environment are not shown in the figure. Replacing one  $c\bar{c}$  pair by a light quark pair,  $q\bar{q}$ , the diagram would describe the production of  $X(3872)$ .

The main difference between a tetraquark and a meson molecule is that the former is compact and the interaction between the constituents occurs through color exchange forces, whereas the latter is an extended object and the interaction between its constituents happens through meson exchange forces. In what was said above, no explicit mention to size or color is made. However, when we speak about the initial gluon fusion and about the final color neutralization through gluon emission or absorption, it is understood that all this must happen within the confinement scale  $\approx 1$  fm. For this reason, we believe that our model is suitable to describe tetraquark production. Although not explicitly excluded, it seems very unlikely that the clusters with masses  $M_{12}$  and  $M_{34}$  will form color singlets interacting through meson exchange.

## B. Kinematics

Working with the usual CEM one-dimensional kinematics, the rapidities of the objects  $M_{12}$  and  $M_{34}$  are, respectively,

$$y_{12} = \frac{1}{2} \ln \frac{x_1}{x_2} \quad \text{and} \quad y_{34} = \frac{1}{2} \ln \frac{x_3}{x_4}, \quad (1)$$

and their invariant masses are

$$M_{12} = \sqrt{x_1 x_2 s} \quad \text{and} \quad M_{34} = \sqrt{x_3 x_4 s}. \quad (2)$$

In terms of these variables and in the low  $p_T$  regime, the invariant mass of the  $c\bar{c}c\bar{c}$  system is then given by

$$M^2 = M_{12}^2 + M_{34}^2 + 2M_{12}M_{34} \cosh(y_{12} - y_{34}). \quad (3)$$

The cosh function grows very rapidly with the argument, and hence even a modest rapidity difference between the two clusters with  $M_{12}$  and  $M_{34}$  will significantly increase the value of  $M$ . We will then assume that both clusters move with equal rapidity, i.e.  $y_{12} = y_{34}$ , and become bound to each other, forming a system with mass:

$$M = M_{12} + M_{34}. \quad (4)$$

Finally, in order to produce the final tetraquark state with right mass,  $M_T$ , the cluster with mass  $M$  emits or absorbs gluons carrying an energy  $\Delta$ , which will be discussed below. We have thus

$$M \pm \Delta = M_T. \quad (5)$$

A remarkable difference between the standard CEM for charmonium production and the model developed here is in the role played by the limits of the integral over the squared invariant mass  $M^2$ . In the case of the usual  $J/\psi$  production, it goes from  $(2m_c)^2$  to  $(2m_D)^2$ . This ensures that the  $c - \bar{c}$  can never decay into an open charm, not forming the

charmonium state, because it does not have enough invariant mass. The case of the tetraquark  $X(3872)$  is different. Suppose, for example, that we have the four-quark system with an invariant mass of 3740 MeV. While this system can only form the  $X$  resonance by absorbing some gluons (carrying energy  $\Delta$ ) from the target or from the projectile, it has sufficient mass to decay immediately into a  $D^+D^-$  pair and not form the resonance. Moreover, since the energy  $\Delta$  is carried by an undefined number of gluons, this decay is not hindered by parity (or charge conjugation) conservation. Therefore, in our case, the integration over  $M^2$  must be changed, becoming more restrictive,

$$\int_{(2m_c)^2}^{(2m_D)^2} dM^2 \rightarrow \int_{(M_T-\Delta)^2}^{(M_T+\Delta)^2} dM^2, \quad (6)$$

where the left side refers to the usual CEM and the right side refers to tetraquark states. We will use this restriction in Sec. III.

## II. TETRAQUARK PRODUCTION

### A. $T_{4c}$ : The all-charm tetraquark

The  $T_{4c}$  state was first discussed a long time ago by Iwasaki [14]. In the 1980s and early 1990s, many authors [15–18] addressed the subject, arriving at different conclusions concerning the existence of a  $c\bar{c}c\bar{c}$  bound state. More recently, with the revival of charmonium spectroscopy, Lloyd and Vary [19] investigated the four-body  $c\bar{c}c\bar{c}$  system obtaining several close-lying bound states. They found that deeply bound ( $\approx 100$  MeV) states may exist with masses around 6 GeV. In Ref. [20], the existence of  $c\bar{c}c\bar{c}$  states was discussed in the framework of the hyperspherical harmonic formalism. The results suggested the possible existence of three four-quark bound states with quantum numbers  $0^{+-}$ ,  $2^{+-}$ , and  $2^{++}$  and masses of the order of 6.50, 6.65, and 6.22 GeV. More recently, using the Bethe-Salpeter approach, the authors of Ref. [21] found an all-charm tetraquark with  $J^{PC} = 0^{++}$  and mass  $M_{T_{4c}} = 5.3 \pm 0.5$  GeV. This mass is considerably lower than the 6.0 GeV obtained in the previous model calculations [14,19]. It is also much lower than the  $2\eta_c$  threshold. Potential decay channels into D mesons and pairs of light mesons necessarily involve internal gluon lines. The resulting decay width may therefore be rather small. On the other hand, preliminary lattice QCD calculations [22,23] seem to disfavor the existence of a deeply bound  $c\bar{c}c\bar{c}$  state, being more compatible with a loosely bound  $2\eta_c$  molecular state. In the works [24,25],  $T_{4c}$  production was studied in SPS events.

### B. Production cross section

The cross section of the process shown in Fig. 1 can be calculated with the schematic DPS “pocket” formula,

$$\sigma_{\text{DPS}} \propto \frac{\sigma_{\text{SPS}}^{12} \sigma_{\text{SPS}}^{34}}{\sigma_{\text{eff}}}, \quad (7)$$

where  $\sigma_{\text{eff}} \approx 15$  mb is a constant extracted from data analysis and  $\sigma_{\text{SPS}}$  is the standard QCD parton model formula, i.e., the convolution of parton densities with partonic cross sections. To be more precise, we expand the above formula showing the kinematical constraints introduced to study tetraquark production. It reads

$$\begin{aligned} \sigma_{\text{DPS}} = & \frac{F_{T_{4c}}}{\sigma_{\text{eff}}} \left[ \int_0^1 dx_1 \int_0^1 dx_2 g(x_1, \mu^2) g(x_2, \mu^2) \sigma_{g_1 g_2 \rightarrow c\bar{c}} \right] \\ & \times \left[ \int_0^1 dx_3 \int_0^1 dx_4 g(x_3, \mu^2) g(x_4, \mu^2) \sigma_{g_3 g_4 \rightarrow c\bar{c}} \right] \\ & \times \Theta(1 - x_1 - x_3) \Theta(1 - x_2 - x_4) \\ & \times \Theta(M_{12}^2 - 4m_c^2) \Theta(M_{34}^2 - 4m_c^2) \\ & \times \delta(y_{34} - y_{12}), \end{aligned} \quad (8)$$

where  $g(x, \mu^2)$  is the gluon distribution in the proton with the gluon fractional momentum  $x$  and at the factorization scale  $\mu^2$  and  $\sigma_{gg \rightarrow c\bar{c}}$  is the  $gg \rightarrow c\bar{c}$  elementary cross section. The step functions  $\Theta(1 - x_1 - x_3)$  and  $\Theta(1 - x_2 - x_4)$  enforce momentum conservation in the projectile and in the target. The step functions  $\Theta(M_{12}^2 - 4m_c^2)$  and  $\Theta(M_{34}^2 - 4m_c^2)$  guarantee that the invariant masses of the gluon pairs 12 and 34 are large enough to produce two charm quark pairs. The delta function implements the “binding condition,” and  $F_{T_{4c}}$  is a constant, analogous to the one appearing in the CEM formula, which represents the probability of the four-quark system to evolve to a particular tetraquark state.

In the above formula, all the variables depend on the momentum fractions  $x_1 \dots x_4$ . Because of the delta function, we know that the two clusters shown in Fig. 1 are “flying together” and that they form a system with mass  $M = M_{12} + M_{34}$ , which can take any value. In order to improve our kinematical description of this bound state, we can impose constraints on the values of  $M$ , such as (6). This can be best done rewriting (8) and changing variables from  $x_1, x_2, x_3$ , and  $x_4$  to  $y_{12}, y_{34}, M_{12}$ , and  $M_{34}$ . We obtain

$$\begin{aligned} \sigma_{\text{DPS}} = & \frac{F_{T_{4c}}}{\sigma_{\text{eff}}} \left[ \frac{1}{s} \int dy_{12} \int dM_{12}^2 g(\bar{x}_1, \mu^2) g(\bar{x}_2, \mu^2) \sigma_{g_1 g_2 \rightarrow c\bar{c}} \right] \\ & \times \left[ \frac{1}{s} \int dy_{34} \int dM_{34}^2 g(\bar{x}_3, \mu^2) g(\bar{x}_4, \mu^2) \sigma_{g_3 g_4 \rightarrow c\bar{c}} \right] \\ & \times \Theta(1 - \bar{x}_1 - \bar{x}_3) \Theta(1 - \bar{x}_2 - \bar{x}_4) \Theta(M_{12}^2 - 4m_c^2) \\ & \times \Theta(M_{34}^2 - 4m_c^2) \delta(y_{34} - y_{12}), \end{aligned} \quad (9)$$

where

$$\begin{aligned}\bar{x}_1 &= \frac{M_{12}}{\sqrt{s}} e^{y_{12}}, & \bar{x}_2 &= \frac{M_{12}}{\sqrt{s}} e^{-y_{12}}, \\ \bar{x}_3 &= \frac{M_{34}}{\sqrt{s}} e^{y_{34}}, & \bar{x}_4 &= \frac{M_{34}}{\sqrt{s}} e^{-y_{34}}\end{aligned}\quad (10)$$

and consequently

$$\begin{aligned}\Theta(1 - \bar{x}_1 - \bar{x}_3) &= \Theta\left(1 - \frac{M_{12}}{\sqrt{s}} e^{y_{12}} - \frac{M_{34}}{\sqrt{s}} e^{y_{34}}\right), \\ \Theta(1 - \bar{x}_2 - \bar{x}_4) &= \Theta\left(1 - \frac{M_{12}}{\sqrt{s}} e^{-y_{12}} - \frac{M_{34}}{\sqrt{s}} e^{-y_{34}}\right).\end{aligned}\quad (11)$$

From the above expressions, it is easy to see that when  $y_{12} = y_{34} = y$  then (4) holds, and the theta functions give lower and upper limits for the integration in  $y$ :

$$-\ln \frac{\sqrt{s}}{M} \leq y \leq \ln \frac{\sqrt{s}}{M}.\quad (12)$$

The upper limit of  $M_{12}$  and  $M_{34}$  can be fixed imposing constraints on their sum,  $M$ . In the case of the  $X(3872)$ , we already know the mass of the state that we want to produce. In principle, we could just use (4) with a fixed value of  $M$ . However, following the spirit of the CEM, we will assume that when the system with mass  $M = M_{12} + M_{34}$  goes to the final state with mass  $M_T$  it can absorb or emit soft gluons to neutralize color. These gluons carry an energy going from almost zero to the QCD scale, given by  $\Delta = \mathcal{O}(\Lambda_{\text{QCD}})$ . Then, from (4) and (5), we have

$$M^{\min} = M_{12}^{\min} + M_{34}^{\min} = M_T - \Delta\quad (13)$$

and

$$M^{\max} = M_{12}^{\max} + M_{34}^{\max} = M_T + \Delta.\quad (14)$$

From these equations, we can see that, knowing the mass of the tetraquark state and fixing the amount of energy which can be exchanged in the formation of the final state, we constrain the limits in the integrations over  $M_{12}$  and  $M_{34}$ . In the symmetric case of  $T_{4c}$  production  $M_{12}^{\min} = M_{34}^{\min}$ ,  $M_{12}^{\max} = M_{34}^{\max}$ , Eqs. (13) and (14) completely fix these limits. In the case of the  $X(3872)$ , we may have different choices for  $M_{12}^{\min}(M_{12}^{\max})$  and  $M_{34}^{\min}(M_{34}^{\max})$ , but they will be correlated.

### III. NUMERICAL RESULTS AND DISCUSSION

#### A. $T_{4c}$

As mentioned in the Introduction, we take the production cross section of the  $T_{4c}$  as a baseline because it is heavy, and hence treatable in pQCD, and also to make some contact with the production of  $c\bar{c}c\bar{c}$  in DPS. In this subsection, we discuss the numerical results obtained for  $T_{4c}$ . Then, in the following subsection, after only a few

changes, we calculate the cross section for  $X(3872)$  production.

We now evaluate Eq. (9), replacing  $g(x, \mu^2)$  by the Martin-Roberts-Stirling-Thorne (MRST) gluon distribution [26] and  $\sigma_{gg \rightarrow c\bar{c}}$  by the standard leading order QCD result [13],

$$\begin{aligned}\sigma_{gg \rightarrow c\bar{c}} &= \frac{\pi\alpha_s^2(m^2)}{3m^2} \left\{ \left(1 + \frac{4m_c^2}{m^2} + \frac{m_c^4}{m^4}\right) \ln \left[\frac{1+\beta}{1-\beta}\right] \right. \\ &\quad \left. - \frac{1}{4} \left(7 + \frac{31m_c^2}{m^2}\right) \beta \right\}\end{aligned}\quad (15)$$

with

$$\beta = \left[1 - \frac{4m_c^2}{m^2}\right]^{1/2},$$

where  $m^2$  is equal to  $M_{12}^2$  or  $M_{34}^2$ . A difficulty in our calculation is the uncertainty in the normalization of the cross section. Whereas in the case of charmonium production in the CEM we have experimental information, which can be used to fix the nonperturbative constant  $F_H$ , in the case of the  $T_{4c}$ , nothing is known. For the time being, we can only try to make a simple estimate.

In the usual CEM, it is assumed that the nonperturbative probability for the  $Q\bar{Q}$  pair to evolve into a quarkonium state  $H$  is given by a constant  $F_H$  that is energy-momentum and process independent. Once  $F_H$  has been fixed by comparison with the measured total cross section for the production of the quarkonium  $H$  at one given energy, the CEM can predict, with no additional free parameters, the energy dependence of the production cross section and the momentum distribution of the produced quarkonium. Following the CEM strategy, we shall adjust  $\sigma_{T_{4c}}$ , connecting it to the experimentally measured cross section of  $X(3872)$  production at one single energy, and then make predictions for higher energies.

We know that the production cross section of  $T_{4c}$  must be smaller than the one for  $X(3872)$  production, and the latter has been measured by the CMS Collaboration [9] at  $\sqrt{s} = 7$  TeV. Moreover, assuming that the binding mechanism is the same, the only difference is that we must replace the light quark pair [which is in the  $X(3872)$ ] by the  $c\bar{c}$  pair, which is much more difficult to produce. Therefore, in order to estimate the cross section for producing the  $T_{4c}$ , we must multiply the  $X(3872)$  production cross section,  $\sigma_X$ , by a penalty factor,

$$\sigma_{T_{4c}} = \frac{\sigma_{c\bar{c}c\bar{c}}}{\sigma_{c\bar{c}q\bar{q}}} \sigma_X \approx \frac{\sigma_{c\bar{c}}\sigma_{c\bar{c}}}{\sigma_{c\bar{c}}\sigma_{q\bar{q}}} \sigma_X \approx \frac{\sigma_{c\bar{c}}}{\sigma_{\text{inel}}} \approx 0.12\sigma_X,\quad (16)$$

where  $\sigma_{c\bar{c}c\bar{c}}$  and  $\sigma_{c\bar{c}q\bar{q}}$  are the cross sections for the production of  $c\bar{c}c\bar{c}$  and  $c\bar{c}q\bar{q}$ , respectively. These cross sections can be measured in double parton scattering events. In the above expression, after using the factorization hypothesis,  $\sigma_{c\bar{c}}$  cancels out, and the ratio  $\sigma_{c\bar{c}}/\sigma_{q\bar{q}} \approx \sigma_{c\bar{c}}/\sigma_{\text{inel}}$  can be inferred from data [27,28], which at

TABLE I.  $X(3872)$  and  $T_{4c}$  production cross sections at  $\sqrt{s} = 7$  TeV and  $\sqrt{s} = 14$  TeV.

Energy (TeV)	$\sigma_{c\bar{c}}$ (mb)	$\sigma_{\text{inel}}$ (mb)	$\sigma_X$ (nb)	$\sigma_{T_{4c}}$ (nb)
7	8.5 [28]	73.2 [27]	30.0 [9]	$3.6 \pm 2.5$
14			$44.6 \pm 17.7$	$7.0 \pm 4.8$

7 TeV yield  $\approx 0.12$ . All the required numbers are collected in Table I. Finally, using the value of  $\sigma_X \approx 30$  nb [9], we have

$$\sigma_{T_{4c}}(\sqrt{s} = 7 \text{ TeV}) \approx (3.6 \pm 2.5) \text{ nb}. \quad (17)$$

Having fixed the numbers, we plot the cross section for  $T_{4c}$  production as a function of the energy in Fig. 2. In order to obtain an estimate of the theoretical error, we vary the parameters trying to scan the most relevant region in the parameter space. We choose  $\Delta \approx \Lambda_{\text{QCD}} \approx 200$  MeV, and we assume that the  $T_{4c}$  mass is given by  $M_{T_{4c}} = 5.4$  GeV, as obtained in Ref. [21]. With these two parameters fixed, we can choose different values for the charm mass  $m_c$ . However, there is an upper limit for  $m_c$ , which cannot be bigger than  $M_{12}^{\text{min}}/2$ . Substituting  $\Delta$  and  $M_{T_{4c}}$  in Eq. (13), we find that  $M_{12}^{\text{min}} = 2.6$  GeV, and consequently the maximum value for  $m_c$  is  $m_c = 1.3$  GeV. In Fig. 2, the upper line corresponds to  $m_c = 1.2$  GeV, and the lower line corresponds to  $m_c = 1.3$  GeV. The star in Fig. 2 corresponds to the central value at  $\sqrt{s} = 7$  TeV. Here, the constant  $F_{T_{4c}}$  was chosen so as to reproduce (17). Once all the parameters are fixed at  $\sqrt{s} = 7$  TeV, the energy dependence of the cross section is completely determined by the model. In Fig. 2, the cross represents the central value of our prediction for the energy  $\sqrt{s} = 14$  TeV:

$$\sigma_{T_{4c}}(\sqrt{s} = 14 \text{ TeV}) \approx (7.0 \pm 4.8) \text{ nb}. \quad (18)$$




FIG. 2. Cross section of  $T_{4c}$  production as a function of the energy.

The main feature of the curves is the rapid rise with  $\sqrt{s}$ , which might render the  $T_{4c}$  observable already at 14 TeV. This same fast growing trend was observed in other estimates with DPS [7,8].

## B. $X(3872)$

We now turn to the production cross section of  $X(3872)$ . We use the same parton densities as in the previous subsection and also the elementary cross section for heavy quark production (15). Note that we use this expression even for light quark production  $\sigma_{g_s g_s \rightarrow q \bar{q}}$ , which appears now in the second line of (8) or (9). Since this expression only holds for heavy enough quarks, its use here is questionable. In spite of this uncertainty, the existing experience in the literature is encouraging. In Ref. [29], the authors used (15) to compute the cross section of strange particle production and calculated the asymmetries in the production of  $K^+/K^-$ ,  $\Lambda/\bar{\Lambda}$ , ..., etc. They have used the convolution formula of the parton model and have taken the strange quark mass to be  $m_s \approx 500$  MeV. They could reproduce well the existing data on asymmetries. In our case, the invariant mass  $M_{34}$  defines the perturbative QCD scale, and hence we must have  $M_{34} > 1$  GeV. This can be achieved with the light quarks having masses close to zero and transverse momenta in the few GeV region. Since we are using the one-dimensional version of the CEM, instead of transverse momentum we will assign an effective mass to the light quarks,  $m_q = 0.5$  GeV, which guarantees that  $M_{34} > 2m_q > 1$  GeV. Moreover, choosing  $N_f = 2$  and  $\Lambda_{\text{QCD}} = 200$  MeV, we have typically

$$\alpha_s = \frac{12\pi}{(33 - 2N_f) \ln\left(\frac{(2m_q)^2}{\Lambda_{\text{QCD}}^2}\right)} \approx 0.4. \quad (19)$$

Although we may expect significant corrections, this number is still small enough for perturbation theory to make sense. As in the previous subsection, after fixing these parameters and knowing the tetraquark mass  $M_X = 3872$  MeV, the only remaining free parameters are the charm mass and the constant  $F_X$ . We show our results in Fig. 3, where the upper line corresponds to  $m_c = 1.2$  GeV and the lower line corresponds to  $m_c = 1.3$  GeV. The constant  $F_X$  was adjusted so that the central value of the cross section at  $\sqrt{s} = 7$  TeV (shown with a star) corresponds to  $\sigma_X = 30.0$  nb. With all the numbers fixed at the lower energy, the energy dependence is given by the model. At  $\sqrt{s} = 14$  TeV, the cross indicates the central value of our prediction:

$$\sigma_X(\sqrt{s} = 14 \text{ TeV}) \approx 44.6 \pm 17.7 \text{ nb}. \quad (20)$$

The error in the number given above is relatively large, but at least we can predict the order of magnitude of the cross section. As a first estimate with DPS, we think that the




FIG. 3.  $X(3872)$  production cross section as a function of the energy.

result is satisfactory. The model presented here can be improved in several aspects. Probably the most relevant one is the prescription to form the resonance, i.e., the hadronization of the multi-quark system. Progress in this direction would also benefit the SPS calculations of this process. Our prescription, based mostly on the kinematical aspects and using only the rapidities and invariant masses, is not accurate enough and is the largest source of uncertainties. Work along this line is in progress. The other sources of uncertainties are, as usual, the choice of parton densities, the choice of the energy scale at which they are computed, the choice of the scale at which  $\alpha_s$  is computed, the choice of  $\Lambda_{\text{QCD}}$ , and the charm and light quark masses.

#### IV. CONCLUSION

We have developed a model for tetraquark production which combines double parton scattering and the basic ideas of the color evaporation model. We have made predictions for the  $X(3872)$  production cross section, which may be confronted with the forthcoming LHC data taken at  $\sqrt{s} = 14$  TeV.

The results presented above contain some uncertainties:

- (i) They do not include tetraquark production in SPS events, which can be larger than the DPS cross section. The calculation of the SPS cross section requires some fragmentation function which is not known.
- (ii) The binding mechanism is probably too simple and insensitive to the quantum numbers of the involved particles.
- (iii) In the case of  $X(3872)$  production, the use of formula (15) for light quark production is questionable. This problem may be circumvented using the next-to-leading order version of the CEM, in which the transverse momentum is included. In this case, the light quarks can be really light, but they have large  $p_T$ , rendering plausible the use of the perturbative formula (15).

#### ACKNOWLEDGMENTS

We are grateful to J. M. Richard and to M. Nielsen for instructive discussions. This work was partially supported by the Brazilian funding agencies CAPES, CNPq, FAPESP, and FAPERGS.

- 
- [1] A. Esposito, A. L. Guerrieri, F. Piccinini, A. Pilloni, and A. D. Polosa, *Int. J. Mod. Phys. A* **30**, 1530002 (2015).
  - [2] M. Nielsen and F. S. Navarra, *Mod. Phys. Lett. A* **29**, 1430005 (2014); M. Nielsen, F. S. Navarra, and S. H. Lee, *Phys. Rep.* **497**, 41 (2010); F. S. Navarra, M. Nielsen, and S. H. Lee, *Phys. Lett. B* **649**, 166 (2007).
  - [3] C. Bignamini, B. Grinstein, F. Piccinini, A. D. Polosa, and C. Sabelli, *Phys. Rev. Lett.* **103**, 162001 (2009).
  - [4] A. Esposito, A. L. Guerrieri, L. Maiani, F. Piccinini, A. Pilloni, A. D. Polosa, and V. Riquer, *Phys. Rev. D* **92**, 034028 (2015).
  - [5] V. Khachatryan *et al.* (CMS Collaboration), *J. High Energy Phys.* **09** (2014) 094.
  - [6] R. Aaij *et al.* (LHCb Collaboration), *Phys. Lett. B* **707**, 52 (2012).
  - [7] R. Astalos *et al.*, [arXiv:1506.05829](https://arxiv.org/abs/1506.05829); M. Diehl, *Proc. Sci., DIS2013* (2013) 006.
  - [8] For recent reviews, see S. Bansal *et al.*, [arXiv:1410.6664](https://arxiv.org/abs/1410.6664); A. Szczurek, *Acta Phys. Pol. B Proc. Suppl.* **8**, 483 (2015); *Acta Phys. Pol. B* **46**, 1415 (2015).
  - [9] S. Chatrchyan *et al.* (CMS Collaboration), *J. High Energy Phys.* **04** (2013) 154.
  - [10] E. R. Cazaroto, V. P. Goncalves, and F. S. Navarra, *Phys. Rev. D* **88**, 034005 (2013).
  - [11] M. Luszczak, R. Maciula, and A. Szczurek, *Phys. Rev. D* **85**, 094034 (2012).
  - [12] See N. Brambilla *et al.*, *Eur. Phys. J. C* **71**, 1534 (2011) and references therein.
  - [13] R. Vogt, *Ultrarelativistic Heavy Ion Collisions* (Elsevier, Amsterdam, 2007), p. 396.
  - [14] Y. Iwasaki, *Prog. Theor. Phys.* **54**, 492 (1975).
  - [15] J. P. Ader, J.-M. Richard, and P. Taxil, *Phys. Rev. D* **25**, 2370 (1982); J. L. Ballot and J.-M. Richard, *Phys. Lett.* **123B**, 449 (1983); H. J. Lipkin, *Phys. Lett. B* **172**, 242 (1986).
  - [16] L. Heller and J. A. Tjon, *Phys. Rev. D* **32**, 755 (1985); **35**, 969 (1987).
  - [17] B. Silvestre-Brac, *Phys. Rev. D* **46**, 2179 (1992).
  - [18] B. Silvestre-Brac and C. Semay, *Z. Phys. C* **57**, 273 (1993); **59**, 457 (1993); **61**, 271 (1994).

- [19] R. J. Lloyd and J. P. Vary, *Phys. Rev. D* **70**, 014009 (2004).
- [20] N. Barnea, J. Vijande, and A. Valcarce, *Phys. Rev. D* **73**, 054004 (2006).
- [21] W. Heupel, G. Eichmann, and C. S. Fischer, *Phys. Lett. B* **718**, 545 (2012).
- [22] M. Wagner, A. Abdel-Rehim, C. Alexandrou, M. Dalla Brida, M. Gravina, G. Koutsou, L. Scorzato, and C. Urbach, *J. Phys. Conf. Ser.* **503**, 012031 (2014).
- [23] P. Bicudo, K. Cichy, A. Peters, B. Wagenbach, and M. Wagner, *Phys. Rev. D* **92**, 014507 (2015).
- [24] A. V. Berezhnoy, A. K. Likhoded, A. V. Luchinsky, and A. A. Novoselov, *Phys. Rev. D* **84**, 094023 (2011).
- [25] A. V. Berezhnoy, A. V. Luchinsky, and A. A. Novoselov, *Phys. Rev. D* **86**, 034004 (2012).
- [26] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, *Eur. Phys. J. C* **4**, 463 (1998).
- [27] B. Abelev *et al.* (ALICE Collaboration), *Eur. Phys. J. C* **73**, 2456 (2013).
- [28] B. Abelev *et al.* (ALICE Collaboration), *J. High Energy Phys.* **07** (2012) 191.
- [29] T. D. Gutierrez and R. Vogt, *Nucl. Phys.* **A705**, 396 (2002).