

**Diboson excesses in leptophobic  $U(1)_{LP}$  models from string theories**Tianjun Li,<sup>1,2</sup> James A. Maxin,<sup>3</sup> Van E. Mayes,<sup>4</sup> and Dimitri V. Nanopoulos<sup>5,6,7</sup><sup>1</sup>*State Key Laboratory of Theoretical Physics and Kavli Institute for Theoretical Physics China (KITPC), Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, People's Republic of China*<sup>2</sup>*School of Physical Electronics, University of Electronic Science and Technology of China, Chengdu 610054, People's Republic of China*<sup>3</sup>*Department of Physics and Engineering Physics, The University of Tulsa, Tulsa, Oklahoma 74104, USA*<sup>4</sup>*Department of Physics, University of Houston-Clear Lake, Houston, Texas 77058, USA*<sup>5</sup>*George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA*<sup>6</sup>*Astroparticle Physics Group, Houston Advanced Research Center (HARC), Mitchell Campus, Woodlands, Texas 77381, USA*<sup>7</sup>*Academy of Athens, Division of Natural Sciences, 28 Panepistimiou Avenue, Athens 10679, Greece*  
(Received 25 November 2015; published 5 February 2016)

The ATLAS Collaboration has reported excesses in the search for resonant diboson production with decay modes to hadronic final states at a diboson invariant mass around 2 TeV in boosted jets from  $WZ$ ,  $W^+W^-$ , and  $ZZ$  channels. Given potential contamination, we investigate the anomalies in leptophobic  $U(1)_{LP}$  models. We show that leptophobic models can be constructed in flipped  $SU(5) \times U(1)_X$  models from free fermionic string constructions and Pati-Salam models from D-brane constructions. Additionally, we perform a collider phenomenological analysis to study production cross sections for  $pp \rightarrow Z' \rightarrow jj/\bar{t}\bar{t}/WW/Zh$  and discover the excess can be interpreted in both the leptophobic flipped  $SU(5) \times U(1)_X$  models and intersecting D-branes.

DOI: [10.1103/PhysRevD.93.045007](https://doi.org/10.1103/PhysRevD.93.045007)**I. INTRODUCTION**

The ATLAS and CMS Collaborations have completed searches for massive resonances decaying into a pair of weak gauge bosons via jet substructure techniques, i.e., the  $pp \rightarrow V_1 V_2 \rightarrow 4j$  ( $V_{1,2} = W^\pm$  or  $Z$ ) channels [1–3]. The ATLAS analyses consisted of  $20.3 \text{ fb}^{-1}$  of data at 8 TeV LHC beam collision energies, indicating excesses for narrow widths around 2 TeV in the  $WZ$ ,  $WW$ , and  $ZZ$  channels with local significances of  $3.4\sigma$ ,  $2.6\sigma$ , and  $2.9\sigma$ , respectively [1]. Furthermore, CMS performed similar searches, though did not distinguish between  $W$ - and  $Z$ -tagged jets, uncovering a  $1.4\sigma$  excess near 1.9 TeV [2]. It is intriguing that CMS also reported about  $2\sigma$  and  $2.2\sigma$  excesses near 1.8 TeV and 1.8–1.9 TeV in the dijet resonance channel and the  $e\nu b\bar{b}$  channel, respectively, which could be accounted for by a  $W' \rightarrow Wh$  process [4,5]. Though these excesses are not yet statistically significant, consideration is warranted for potential interpretations of these anomalous events as new physics beyond the Standard Model (SM), as evidence mounts for a possible nontrivial explanation. In the intervening time since ATLAS and CMS first reported their findings, these diboson excesses have been extensively studied [6–45].

The ATLAS diboson excess is well fit by resonance peaks around 2 TeV and widths less than about 100 GeV. Narrow resonances such as this might imply new weakly interacting particles, therefore we shall consider the underlying theories to be perturbative in this work. Turning our

focus to the ATLAS excess in the  $WZ$ ,  $WW$ , and  $ZZ$  channels, the tagging selections for each mode used in the analysis are rather incomplete, as these channels share about 20% of the events. It may be difficult to pronounce that a single resonance is responsible for all excesses, although there does remain the possibility that one 2 TeV particle contributes to the excess in only one channel, whereas the additional excesses in the alternate channels are via contaminations. Approaching the analysis from this perspective provides motivation for not attempting to formulate a simultaneous explanation for all excesses, thereby studying only models with a new resonance in one channel. The reference ranges of the production cross-section times the decay branching ratio for the 2 TeV resonances in the  $WZ$ ,  $WW$ , and  $ZZ$  channels are approximately 4–8, 3–7, and 3–9 fb, respectively.

The goal in this work is to understand the diboson excesses in leptophobic  $U(1)_{LP}$  models from string theories. The leptophobic property aids the process of relaxing LHC search constraints on the leptonic decay channel  $Z' \rightarrow \ell^+ \ell^-$ , with string model building allowing for a deeper understanding of the particle physics. Consequently, we shall realize a leptophobic  $U(1)_{LP}$  in flipped  $SU(5) \times U(1)_X$  models from free fermionic string constructions [46–51] and in Pati-Salam models from D-brane constructions [52–63]. We conclude the study with exploration of the production cross sections for  $pp \rightarrow Z' \rightarrow jj/\bar{t}\bar{t}/WW/Zh$ , demonstrating a plausible interpretation of the ATLAS excess in both leptophobic flipped  $SU(5) \times U(1)_X$  models and intersecting D-branes.

## II. THE LEPTOPHOBIC $U(1)_{\text{LP}}$ MODEL FROM STRINGY FLIPPED $SU(5) \times U(1)_X$ MODELS

The convention we adopt here denotes the SM left-handed quark doublets, right-handed up-type quarks, right-handed down-type quarks, left-handed lepton doublets, right-handed charged leptons, and right-handed neutrinos as  $Q_i$ ,  $U_i^c$ ,  $D_i^c$ ,  $L_i$ ,  $E_i^c$ , and  $N_i^c$  respectively. Our analysis shall investigate the leptophobic  $U(1)_{\text{LP}}$  model from string theory as a viable explanation of the diboson excess. The leptophobic  $U(1)_{\text{LP}}$  cannot be realized in  $SU(5)$  models due to the fact the matter field representations  $\mathbf{10}_i$  contain  $\{Q_i, U_i^c, E_i^c\}$ , while the representations  $\bar{\mathbf{5}}_i$  contain  $\{D_i^c, L_i\}$ . Similar results are found for traditional  $SO(10)$  and  $E_6$  models as well. Of significant note, representations for three families of SM fermions in flipped  $SU(5) \times U(1)_X$  models [46] are

$$\begin{aligned} F_i &= (\mathbf{10}, \mathbf{1}) = \{Q_i, D_i^c, N_i^c\}, \\ \bar{f}_i &= (\bar{\mathbf{5}}, -\mathbf{3}) = \{U_i^c, L_i\}, \\ \bar{l}_i &= (\mathbf{1}, \mathbf{5}) = \{E_i^c\}, \end{aligned} \quad (1)$$

TABLE I. The massless matter fields and their transformation properties under  $G_{U(1)}$  in the observable sector. Under  $SU(5) \times U(1)_X$  gauge symmetry, these fields transform as  $F = (10, 1)$ ,  $\bar{f} = (\bar{5}, -3)$ ,  $\bar{l}^c = (1, 5)$ ,  $h = (5, -2)$ , and  $\bar{h} = (\bar{5}, 2)$ . Moreover, we present their charges under  $U_A$  and three orthogonal linear combinations of interest ( $U', U'', U'''$ ).

	$U_1$	$U_2$	$U_3$	$U_4$	$U_5$	$U_A$	$U'$	$U''$	$U'''$
$F_0$	$-\frac{1}{2}$	0	0	$-\frac{1}{2}$	0	$\frac{3}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0
$F_1$	$-\frac{1}{2}$	0	0	$\frac{1}{2}$	0	$\frac{3}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0
$F_2$	0	$-\frac{1}{2}$	0	0	0	$\frac{1}{2}$	0	$\frac{3}{2}$	$-\frac{1}{2}$
$F_3$	0	0	$\frac{1}{2}$	0	$-\frac{1}{2}$	$\frac{3}{2}$	1	$-\frac{1}{2}$	0
$F_4$	$-\frac{1}{2}$	0	0	0	0	$\frac{3}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0
$\bar{F}_4$	$\frac{1}{2}$	0	0	0	0	$-\frac{3}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0
$\bar{F}_5$	0	$\frac{1}{2}$	0	0	0	$-\frac{1}{2}$	0	$-\frac{3}{2}$	$\frac{1}{2}$
$\bar{f}_2, \bar{l}_2^c$	0	$-\frac{1}{2}$	0	0	0	$\frac{1}{2}$	0	$\frac{3}{2}$	$-\frac{1}{2}$
$\bar{f}_3, \bar{l}_3^c$	0	0	$\frac{1}{2}$	0	$\frac{1}{2}$	$\frac{1}{2}$	0	$\frac{3}{2}$	1
$\bar{f}_5, \bar{l}_5^c$	0	$-\frac{1}{2}$	0	0	0	$\frac{1}{2}$	0	$\frac{3}{2}$	$-\frac{1}{2}$
$h_1$	1	0	0	0	0	-3	1	1	0
$\bar{h}_1$	-1	0	0	0	0	3	-1	-1	0
$h_2$	0	1	0	0	0	-1	0	-3	1
$\bar{h}_2$	0	-1	0	0	0	1	0	3	-1
$h_3$	0	0	1	0	0	2	1	1	1
$\bar{h}_3$	0	0	-1	0	0	-2	-1	-1	-1
$h_{45}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	0	0	2	$-\frac{1}{2}$	1	$-\frac{1}{2}$
$\bar{h}_{45}$	$\frac{1}{2}$	$\frac{1}{2}$	0	0	0	-2	$\frac{1}{2}$	-1	$\frac{1}{2}$

where  $i = 1, 2, 3$ . Notice that  $F_i$  does not contain the charged leptons, thus the leptons can be charged under the leptophobic  $U(1)_{\text{LP}}$  gauge symmetry. It is also clear that  $\bar{f}_i$  and  $\bar{l}_i$  cannot be charged under the leptophobic  $U(1)_{\text{LP}}$  gauge symmetry.

In this work we consider flipped  $SU(5) \times U(1)_X$  models from four-dimensional free fermionic string constructions [47], which possess various favorable properties regarding vacuum energy, string unification, dynamical generation of all mass scales, top-quark mass, and the strong coupling [48]. The complete gauge group has three identifiable pieces  $G = G_{\text{obs}} \times G_{\text{hidden}} \times G_{U(1)}$ , where  $G_{\text{obs}} = SU(5) \times U(1)_X$ ,  $G_{\text{hidden}} = SU(4) \times SO(10)$ , and  $G_{U(1)} = U_1(1) \times U_2(1) \times U_3(1) \times U_4(1) \times U_5(1)$ . There are 63 massless matter fields present, annotated in detail in Tables I, II, III, and IV, including their charges under  $G_{U(1)}$ . In particular, there are five  $F$ , two  $\bar{F}$ , three  $\bar{f}$ , and three  $\bar{l}_i$ , which according to the original conventions, are denoted as  $F_0$ ,  $F_1$ ,  $F_2$ ,  $F_3$ ,  $F_4$ ,  $\bar{F}_4$ ,  $\bar{F}_5$ ,  $\bar{f}_2$ ,  $\bar{f}_3$ ,  $\bar{f}_5$ ,  $\bar{l}_2$ ,  $\bar{l}_3$ , and  $\bar{l}_5$ , respectively [47].

Special emphasis is warranted for the property  $\text{Tr}U_{1,2,3,5} \neq 0$ , whereas  $\text{Tr}U_4 = 0$ . The anomalous symmetries are artifacts of the truncation of the full string spectrum down to the massless sector. The low-energy effective theory is correctly specified by rotating all anomalies into a single anomalous  $U_A \propto \sum_{i=1,2,3,5} [\text{Tr}U_i] U_i$  [49],

TABLE II. The singlet fields and their transformation properties under  $G_{U(1)}$ ,  $U_A$ , and three orthogonal linear combinations of interest ( $U', U'', U'''$ ).

	$U_1$	$U_2$	$U_3$	$U_4$	$U_5$	$U_A$	$U'$	$U''$	$U'''$
$\Phi_{12}$	-1	1	0	0	0	2	-1	-4	1
$\bar{\Phi}_{12}$	1	-1	0	0	0	-2	1	4	-1
$\Phi_{23}$	0	-1	1	0	0	3	1	4	0
$\bar{\Phi}_{23}$	0	1	-1	0	0	-3	-1	-4	0
$\Phi_{31}$	1	0	-1	0	0	-5	0	0	-1
$\bar{\Phi}_{31}$	-1	0	1	0	0	5	0	0	1
$\phi_{45}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0	0	0	$\frac{3}{2}$	2	$\frac{3}{2}$
$\bar{\phi}_{45}$	$-\frac{1}{2}$	$-\frac{1}{2}$	-1	0	0	0	$-\frac{3}{2}$	-2	$-\frac{3}{2}$
$\phi^+$	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	1	-2	$-\frac{1}{2}$	4	$\frac{1}{2}$
$\bar{\phi}^+$	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	-1	2	$\frac{1}{2}$	-4	$-\frac{1}{2}$
$\phi^-$	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	-1	0	$\frac{3}{2}$	0	$-\frac{3}{2}$
$\bar{\phi}^-$	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	1	0	$-\frac{3}{2}$	0	$\frac{3}{2}$
$\phi_{3,4}$	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	0	-1	$\frac{1}{2}$	2	$-\frac{1}{2}$
$\bar{\phi}_{3,4}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	0	1	$-\frac{1}{2}$	-2	$\frac{1}{2}$
$\eta_{1,2}$	0	0	0	1	0	0	0	0	0
$\bar{\eta}_{1,2}$	0	0	0	-1	0	0	0	0	0
$\Phi_{0,1,3,5}$	0	0	0	0	0	0	0	0	0

TABLE III. The hidden  $SO(10)$  decaplets (**10**)  $T_i$  fields and their transformation properties under  $G_{U(1)}$ . In addition, we present their charges under  $U_A$  and three orthogonal linear combinations of interest ( $U', U'', U'''$ ).

	$U_1$	$U_2$	$U_3$	$U_4$	$U_5$	$U_A$	$U'$	$U''$	$U'''$
$T_1$	$-\frac{1}{2}$	0	$\frac{1}{2}$	0	0	$\frac{5}{2}$	0	0	$\frac{1}{2}$
$T_2$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	0	$-\frac{1}{2}$	$\frac{5}{2}$	0	0	-1
$T_3$	$-\frac{1}{2}$	0	$\frac{1}{2}$	0	0	$\frac{5}{2}$	0	0	$\frac{1}{2}$

then adding a one-loop correction to the D-term corresponding to  $U_A$ :  $D_A \rightarrow D_A + \epsilon M^2$ , where  $M$  is the reduced Planck scale and  $\epsilon = g^2 \text{Tr} U_A / 192 \pi^2$  [64].

The mass spectrum of all states in the accompanying tables can be obtained through a complex procedure by considering trilinear and nonrenormalizable contributions to the superpotential, and likewise to the masses and interactions [47]. However, this procedure does not provide a unique outcome since the vacuum expectation values (VEVs) of the singlet fields in Table II are unknown, though constrained by the anomalous  $U_A$  cancellation conditions. The objective here is to generate an electroweak-scale

TABLE IV. The hidden  $SU(4)$  fields and their transformation properties under  $G_{U(1)}$ .  $D_i$  represent sixplets (**6**), whereas ( $\tilde{F}_i, \tilde{\bar{F}}_i$ ) represent tetraplets (**4,  $\bar{4}$** ). Moreover, we present the charges under  $U_A$  and three orthogonal linear combinations of interest ( $U', U'', U'''$ ).

	$U_1$	$U_2$	$U_3$	$U_4$	$U_5$	$U_A$	$U'$	$U''$	$U'''$
$D_1$	0	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0	$\frac{3}{2}$	$\frac{1}{2}$	2	0
$D_2$	0	$-\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	0	$\frac{3}{2}$	$\frac{1}{2}$	2	0
$D_3$	$-\frac{1}{2}$	0	$\frac{1}{2}$	0	0	$\frac{5}{2}$	0	0	$\frac{1}{2}$
$D_4$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	0	$\frac{3}{2}$	$\frac{3}{2}$	-1	2	0
$D_5$	0	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	$\frac{3}{2}$	$\frac{1}{2}$	2	0
$D_6$	0	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	$-\frac{3}{2}$	$-\frac{1}{2}$	-2	0
$D_7$	$\frac{1}{2}$	0	$-\frac{1}{2}$	0	0	$-\frac{5}{2}$	0	0	$-\frac{1}{2}$
$\tilde{F}_1$	$-\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	0	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{9}{4}$	$-\frac{1}{2}$
$\tilde{F}_2$	$\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	0	$\frac{1}{2}$	-2	$-\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{2}$
$\tilde{F}_3$	$\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$	0	$\frac{1}{2}$	$-\frac{3}{2}$	$-\frac{1}{2}$	$\frac{7}{4}$	0
$\tilde{F}_4$	$-\frac{1}{4}$	$\frac{3}{4}$	$\frac{1}{4}$	0	0	$\frac{1}{2}$	0	$-\frac{9}{4}$	1
$\tilde{F}_5$	$-\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	0	$\frac{1}{2}$	$-\frac{1}{2}$	-1	$-\frac{1}{4}$	$\frac{1}{2}$
$\tilde{F}_6$	$-\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	0	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{9}{4}$	$-\frac{1}{2}$
$\tilde{\bar{F}}_1$	$-\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{3}{2}$	$\frac{1}{2}$	$-\frac{7}{4}$	0
$\tilde{\bar{F}}_2$	$-\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{3}{2}$	$\frac{1}{2}$	$-\frac{7}{4}$	0
$\tilde{\bar{F}}_3$	$\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{4}$	0	$-\frac{1}{2}$	$\frac{1}{2}$	1	$\frac{1}{4}$	$-\frac{1}{2}$
$\tilde{\bar{F}}_4$	$-\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	0	$-\frac{1}{2}$	$\frac{3}{2}$	$\frac{1}{2}$	$-\frac{7}{4}$	0
$\tilde{\bar{F}}_5$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{4}$	0	$-\frac{1}{2}$	2	$\frac{1}{2}$	$-\frac{1}{4}$	$-\frac{1}{2}$
$\tilde{\bar{F}}_6$	$-\frac{3}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	0	0	$\frac{3}{2}$	-1	$-\frac{7}{4}$	0

spectrum that is closely comparable to the Minimum Supersymmetric Standard Model (MSSM). Relevant studies can be found in Refs. [47,50]. Two scenarios are studied here that possess an anomaly free leptophobic  $U(1)_{LP}$  gauge symmetry [47,50].

### A. The first scenario

The  $U_4(1)$  gauge symmetry is traceless (anomaly-free), and hence does not participate in the  $U_A(1)$  cancellation mechanism (unbroken). Specifically,  $U_4(1)$  is leptophobic since the leptons  $\tilde{f}_{2,3,5}$  and  $\tilde{l}_{2,3,5}^c$  are not charged under it from Table I. It is however interesting to note that  $F_0$  and  $F_1$  are indeed charged under  $U_4(1)$ . The  $U_4(1)$  and  $U_Y(1)$  do not mix though: the Higgs doublets, which break the electroweak symmetry, are neutral under  $U_4$  (see  $h_i, \bar{h}_i$  in Table I). The mixing via gauge kinetic functions cannot be realized due to  $\text{Tr}(YU_4) = 0$ . This factor ‘‘protects’’ the leptophobia, as otherwise the leptons would experience their  $U_4$  charges shifted away from zero.

Under the assumption that  $F_0$  and  $F_1$  contain the first two generations of the SM quarks, the  $U_4$  can remain unbroken during the  $SU(5) \times U(1)_X$  symmetry breaking at the usual grand unified theory (GUT) scale. The  $U_4$  symmetry may be broken radiatively at low energy if the singlet fields  $\eta_{1,2}$  and  $\bar{\eta}_{1,2}$ , which solely carry the  $U_4$  charges (see Table II), acquire suitable dynamical VEVs. Although this  $Z'$  can explain the dijet excess, it cannot explain the  $WW$  excess since it clearly does not mix with the  $Z$  boson. However, the top-quark Yukawa coupling is forbidden by the  $U(1)_{LP}$  gauge symmetry.

### B. The second scenario

Three linear combinations of  $U_{1,2,3,5}$  are orthogonal to  $U_A = U_1 - 3U_2 + U_3 + 2U_5$  and traceless. Without loss of generality, we can choose the following basis:  $U'_1 = U_3 + 2U_5$ ,  $U'_2 = U_1 - 3U_2$ ,  $U'_3 = 3U_1 + U_2 + 4U_3 - 2U_5$ . Since the leptons transform as  $\tilde{f}_{2,5}, \ell_{2,5}^c: (0, \frac{3}{2}, -\frac{1}{2})$ ;  $\tilde{f}_3, \ell_3^c: (\frac{3}{2}, 0, 1)$  under  $U'_i$ , there is a unique leptophobic linear combination of  $U'_i$ :  $U' \propto 2U'_1 - U'_2 - 3U'_3 \propto U_1 + U_3 - U_5$ . The  $U'$  gauge symmetry is by construction anomaly-free and leptophobic, and some of the Higgs pentaplets are charged under it (i.e., mixed). The charges of all fields under  $U'$  are given in the tables, along with two extra traceless combinations which can be chosen as  $U'' = U_1 - 3U_2 + U_3 + 2U_5$  and  $U''' = U_2 + U_3 + U_5$ . From the tables we find that only a very limited set of fields is neutral under  $U'$

$$F_2, \bar{F}_5, \Phi_{31}, \bar{\Phi}_{31}, T_1, T_2, T_3, D_3, D_7, \quad (2)$$

and therefore their VEVs will not break the  $U'$  gauge symmetry. The challenge is whether the usual D- and F-flatness conditions can be satisfied with such a limited set of VEVs since it generally breaks the hidden sector gauge

groups. This problem may be solved if one introduces the nonrenormalizable superpotential [50].

It can be verified that if  $\text{Tr}(YU') = 0$ , then  $U'$  can indeed remain unbroken down to low energy, thus permitting the  $U'$  charges to remain unshifted and the leptophobia protected. Moreover, only  $F_2$  and  $\bar{F}_5$  can break the  $SU(5) \times U(1)_X$  gauge symmetry since they are not charged under  $U'$ . Unlike the previous studies [47] where  $F_4$  contains the third-generation quarks, we consider  $F_0$ ,  $F_1$ , and  $F_3$  as the first, second and third generations, respectively. Additionally,  $F_4$  and  $\bar{F}_4$  can form vectorlike particles at the intermediate scale such that string-scale gauge coupling unification can be achieved [48]. For the pentaplet Higgs fields  $h_i$  and  $\bar{h}_i$ , for simplicity, we assume that  $h_2$  and  $\bar{h}_2$  are vectorlike and have mass around the usual GUT or string scale. Moreover, the triplets in the  $(h_1, \bar{h}_1)$  and  $(h_3, \bar{h}_3)$  will be light at low energy since they are charged under  $U(1)_{\text{LP}}$ . Therefore, in the low-energy supersymmetric SM, there will be two pairs of Higgs doublets and two pairs of Higgs triplets. In particular,  $H_d$  is a linear combination of  $h_1$  and  $h_3$ , while  $H_u$  is a linear combination of  $\bar{h}_1$  and  $\bar{h}_3$  and its dominant component is  $\bar{h}_3$ . Notably,  $(h_1, \bar{h}_1)$  and  $(h_3, \bar{h}_3)$  are charged under  $U(1)_{\text{LP}}$ , and then  $Z$  and  $Z'$  are mixed after the Higgs fields acquire the VEVs. Given this scenario, we could explain the diboson and dijet excesses [7]. In particular, unlike the first scenario, the top-quark Yukawa coupling  $F_3 \bar{f}_3 \bar{h}_3$  is allowed by the  $U(1)_{\text{LP}}$  gauge symmetry. And the down-type quark Yukawa couplings such as  $F_0 F_0 h_1$ ,  $F_0 F_1 h_1$ , and  $F_1 F_1 h_1$  can be realized at renormalizable level. While all the other SM fermion Yukawa couplings should be generated via high-dimensional operators. Furthermore, for simplicity, we assume that all the Higgs fields except the SM-like Higgs field are heavy and then are still undetected at the LHC.

### III. LEPTOPHOBIC $Z'$ FROM $SU(4)_C \times SU(2)_L \times SU(2)_R \times U(1)_X$ ON D-BRANES

A phenomenologically interesting intersecting D-brane model has been studied in Refs. [57,58]. A variation of this model with a different hidden sector was also studied in Refs. [59,60]. The full gauge symmetry of the model is given by  $[U(4)_C \times U(2)_L \times U(2)_R]_{\text{observable}} \times [U(2) \times USp(2)^2]_{\text{hidden}}$ , with the matter content shown in Tables V and VI. Note that in Table V,  $a, b, c$ , etc. refer to different stacks of D-branes which wrap cycles of the compactified manifold and which generically intersect at angles. A stack of  $2N$  D-branes results in a  $U(N)$  gauge group in the world volume of each stack. Strings localized at the intersection between two stacks result in massless fermions in the bifundamental representation of the gauge group of each stack. Vectorlike matter may also be present between stacks which do not intersect, again in the bifundamental representation of each stack's gauge group.

Since  $U(N) = SU(N) \times U(1)$ , associated with each of the stacks  $a, b, c$ , and  $d$  are  $U(1)$  gauge groups, denoted as  $U(1)_a, U(1)_b, U(1)_c$ , and  $U(1)_d$ . In general, these  $U(1)$ s are anomalous. The anomalies associated with these  $U(1)$ s are canceled by a generalized Green-Schwarz (GS) mechanism that involves untwisted R-R forms. As a result, the gauge bosons of these Abelian groups generically become massive. The GS couplings determine the exact linear combinations of  $U(1)$  gauge bosons that become massive. Some linear combinations may remain massless if certain conditions are satisfied.

As shown in Ref. [59], precisely one linear combination of the present model remains massless and anomaly-free:

$$U(1)_X = U(1)_a + 2[U(1)_b + U(1)_c + 3U(1)_d]. \quad (3)$$

TABLE V. The chiral superfields, their multiplicities and quantum numbers under the gauge symmetry  $[U(4)_C \times U(2)_L \times U(2)_R]_{\text{observable}} \times [U(2) \times USp(2)^2]_{\text{hidden}}$ , where  $Q_X = Q_a + 2(Q_b + Q_c + 3Q_d)$ . Here  $a, b, c$ , etc. refer to different stacks of D-branes.

	Multiplicities	Quantum Number	$Q_a$	$Q_b$	$Q_c$	$Q_d$	$Q_X$	Field
$ab$	3	$(4, \bar{2}, 1, 1, 1, 1)$	1	-1	0	0	-1	$F_L(Q_L, L_L)$
$ac$	3	$(\bar{4}, 1, 2, 1, 1, 1)$	-1	0	1	0	1	$F_R(Q_R, L_R)$
$bd$	1	$(1, \bar{2}, 1, 2, 1, 1)$	0	-1	0	1	4	$X_{bd}$
$cd$	1	$(1, 1, 2, \bar{2}, 1, 1)$	0	0	1	-1	-4	$X_{cd}$
$b4$	3	$(1, \bar{2}, 1, 1, 1, 2)$	0	-1	0	0	-2	$X_{b3}^i$
$c3$	3	$(1, 1, 2, 1, \bar{2}, 1)$	0	0	1	0	2	$X_{c3}^i$
$d3$	1	$(1, 1, 1, \bar{2}, 2, 1)$	0	0	1	-1	-6	$X_{cd}$
$d4$	1	$(1, 1, 1, 2, 1, \bar{2})$	0	0	1	1	6	$X_{cd}$
$b_S$	2	$(1, 3, 1, 1, 1, 1)$	0	2	0	0	4	$T_L^i$
$b_A$	2	$(1, \bar{1}, 1, 1, 1, 1)$	0	-2	0	0	-4	$S_L^i$
$c_S$	2	$(1, 1, \bar{3}, 1, 1, 1)$	0	0	0	-2	-4	$T_R^i$
$c_A$	2	$(1, 1, 1, 1, 1, 1)$	0	0	2	0	4	$S_R^i$

TABLE VI. The vectorlike superfields, their multiplicities and quantum numbers under the gauge symmetry  $[U(4)_C \times U(2)_L \times U(2)_R]_{\text{observable}} \times [U(2) \times USp(2)^2]_{\text{hidden}}$ , where  $Q_X = Q_a + 2(Q_b + Q_c + 3Q_d)$ . Here  $a, b, c$ , etc. refer to different stacks of D-branes.

	Multiplicities	Quantum Number	$Q_a$	$Q_b$	$Q_c$	$Q_d$	$Q_X$	Field
$ab'$	3	(4,2,1,1,1,1)	1	1	0	0	3	$\Omega_L^i$
	3	( $\bar{4}, \bar{2}, 1, 1, 1, 1$ )	-1	-1	0	0	-3	$\bar{\Omega}_L^i$
$ac'$	3	(4,1,2,1,1,1)	1	0	1	0	3	$\Phi_i$
	3	( $\bar{4}, 1, \bar{2}, 1, 1, 1$ )	-1	0	-1	0	-3	$\bar{\Phi}_i$
$ad$	2	(4,1,1, $\bar{2}$ ,1,1)	1	0	0	-1	-5	$\varphi_i$
	2	( $\bar{4}, 1, 1, 2, 1, 1$ )	-1	0	0	1	5	$\bar{\varphi}_i$
$ad'$	1	(4,1,1,2,1,1)	1	0	0	1	7	$\zeta$
	1	( $\bar{4}, 1, 1, \bar{2}, 1, 1$ )	-1	0	0	-1	-7	$\bar{\zeta}$
$bc$	6	(1,2, $\bar{2}$ ,1,1,1)	0	1	-1	0	0	$H^i(H_u, H_d)$
	6	(1, $\bar{2}$ ,2,1,1,1)	0	-1	1	0	0	$\bar{H}^i(\bar{H}_u, \bar{H}_d)$
$bc'$	1	(1,2,2,1,1,1)	0	1	1	0	4	$\mathcal{H}(\mathcal{H}_u, \mathcal{H}_d)$
	1	(1, $\bar{2}$ , $\bar{2}$ ,1,1,1)	0	-1	-1	0	-4	$\bar{\mathcal{H}}(\bar{\mathcal{H}}_u, \bar{\mathcal{H}}_d)$
$bd'$	1	(1,2,1,2,1,1)	0	1	0	1	8	$\xi$
	1	(1, $\bar{2}$ ,1, $\bar{2}$ ,1,1)	0	0	-1	-1	-8	$\bar{\xi}$
$cd'$	1	(1,1,2,2,1,1)	0	1	0	1	8	$\psi$
	1	(1,1, $\bar{2}$ , $\bar{2}$ ,1,1)	0	0	-1	-1	-8	$\bar{\psi}$

Thus, the effective gauge symmetry of the model at the string scale is given by

$$SU(4)_C \times SU(2)_L \times SU(2)_R \times U(1)_X \times [SU(2) \times USp(2)^2]. \quad (4)$$

As can be seen from Table V, the superfields  $F_L^i(Q_L, L_L)$  carry charge  $Q_X = -1$ , the superfields  $F_R^i(Q_R, L_R)$  carry charge  $Q_X = +1$ . In addition, there are the Higgs superfields  $H_u^i, H_d^i$  in the  $bc$  sector which are uncharged under  $U(1)_X$  while the Higgs superfields  $\mathcal{H}_u, \mathcal{H}_d$  in the  $bc'$  sector carry charges  $Q_X = \pm 4$  respectively.

It should be noted that the Yukawa couplings with the Higgs superfields  $H_u^i, H_d^i$  are allowed by the global  $U(1)$  charges. The resulting Yukawa mass matrices for quarks and leptons are of rank 3, and it has been shown that it is possible to obtain the correct masses and mixings for all quarks and leptons [57,58]. On the other hand, the Yukawa couplings with the Higgs superfields from the  $bc'$  sector  $\mathcal{H}_u, \mathcal{H}_d$  are forbidden. In addition we may form a  $\mu$ -term in the superpotential of the form In addition we may form a  $\mu$ -term in the superpotential of the form

$$W_\mu = \frac{y^{ijkl}}{M_{St}} S_L^i S_R^j H_u^k H_d^l, \quad (5)$$

which is TeV-scale, where  $S_R^j$  receive string-scale VEVs,  $M_{St}$  is the string scale, and the VEVs of  $S_L^i$  are TeV-scale.

The  $\mu$ -term may be fine-tuned so that only a pair of Higgs eigenstates  $H_u$  and  $H_d$  remain light, as in the MSSM.

The gauge symmetry is first broken by splitting the D-branes as  $a \rightarrow a1 + a2$  with  $N_{a1} = 6$  and  $N_{a2} = 2$ , and  $c \rightarrow c1 + c2$  with  $N_{c1} = 2$  and  $N_{c2} = 2$ . After splitting the D6-branes, the gauge symmetry of the observable sector is

$$SU(3)_C \times SU(2)_L \times U(1)_{I3R} \times U(1)_{B-L} \times U(1)_{3B+L}, \quad (6)$$

where

$$U(1)_{I3R} = \frac{1}{2}(U(1)_{c1} - U(1)_{c2}),$$

$$U(1)_{B-L} = \frac{1}{3}(U(1)_{a1} - 3U(1)_{a2}), \quad (7)$$

and

$$U(1)_{3B+L} = -[U(1)_{a1} + U(1)_{a2} + 2(U(1)_b + U(1)_{c1} + U(1)_{c2} + 3U(1)_d)], \quad (8)$$

and  $U(1)_{3B+L} = -U(1)_X$ .

The gauge symmetry must be further broken to the SM, with the possibility of one or more additional  $U(1)$  gauge symmetries. In particular, the  $U(1)_{B-L} \times U(1)_{I3R} \times U(1)_{3B+L}$  gauge symmetry may be broken by assigning VEVs to the right-handed neutrino fields  $N_R^i$ . In this case, the gauge symmetry is broken to

TABLE VII. The chiral superfields, their multiplicities and quantum numbers under the gauge symmetry  $[SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_B]_{\text{observable}} \times [SU(2) \times USp(2)^2]_{\text{hidden}}$ .

	Multiplicities	Quantum Number	$Q_{I3R}$	$Q_{B-L}$	$Q_{3B+L}$	$Q_Y$	$Q_B$	Field
$a1b$	3	$(3, \bar{2}, 1, 1, 1, 1, 1)$	0	1/3	1	1/6	1/3	$Q_L$
$a1c2$	3	$(\bar{3}, 1, 1, 1, 1, 1, 1)$	-1/2	-1/3	-1	-2/3	-1/3	$U_R$
$a1c1$	3	$(\bar{3}, 1, 1, 1, 1, 1, 1)$	1/2	-1/3	-1	1/3	-1/3	$D_R$
$a2b$	3	$(1, \bar{2}, 1, 1, 1, 1, 1)$	0	-1	1	-1/2	0	$L$
$a2c1$	3	$(1, 2, 1, 1, 1, 1, 1)$	1/2	1	-1	1	0	$E_R$
$a2c2$	3	$(1, 2, 1, 1, 1, 1, 1)$	-1/2	1	-1	0	0	$N_R$
$bc1$	6	$(1, 2, 1, 1, 1, 1, 1)$	-1/2	0	0	-1/2	0	$H_d^i$
$bc2$	6	$(1, 2, 1, 1, 1, 1, 1)$	1/2	0	0	1/2	0	$H_u^i$
$bc1'$	1	$(1, \bar{2}, 1, 1, 1, 1, 1)$	-1/2	0	4	-1/2	1	$\mathcal{H}_d$
$bc2'$	1	$(1, \bar{2}, 1, 1, 1, 1, 1)$	1/2	0	4	1/2	1	$\mathcal{H}_u$

$$[SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_B]_{\text{observable}} [SU(2) \times USp(2)^2]_{\text{hidden}} \quad (9)$$

where

$$U(1)_Y = \frac{1}{6} [U(1)_{a1} - 3U(1)_{a2} + 3U(1)_{c1} - 3U(1)_{c2}] \\ = \frac{1}{2} U(1)_{B-L} + U(1)_{I3R}. \quad (10)$$

and

$$U(1)_B = \frac{1}{4} [U(1)_{B-L} + U(1)_{3B+L}] \\ = - \left[ \frac{1}{6} U(1)_{a1} + \frac{1}{2} (U(1)_{a2} + U(1)_b + U(1)_{c1} \right. \\ \left. + U(1)_{c2} + 3U(1)_d) \right]. \quad (11)$$

We will assume that all exotic matter, shown in Table VIII, may become massive, as shown in Ref. [60].

The resulting low-energy field content is shown in Tables VII and along with their charges under  $U(1)_{I3R}$ ,  $U(1)_{B-L}$ ,  $U(1)_{3B+L}$ ,  $U(1)_Y$ , and  $U(1)_B$ . Note that the quarks are charged under  $U(1)_B$ , but the leptons are not.

The extra gauge symmetry  $U(1)_B$  may then be spontaneously broken if the SM singlet fields  $S_L^i$ , which carry a charge of +1 under  $U(1)_B$  obtain VEVs at some scale  $\Lambda = \mathcal{O}(\text{TeV})$ . Thus, the model may possess a leptophobic  $Z'$  boson with an  $\mathcal{O}(\text{TeV})$  mass. In order to explain the diboson excess observed by ATLAS, we will take this scale to be 2–3 TeV.

The electroweak symmetry is broken when some of the Higgs fields obtain VEVs. In order to obtain masses and mixings for the quarks and leptons, we will assume that the dominate VEVs are acquired by the Higgs fields  $H_u^i, H_d^i$  in the  $bc$  sector which are uncharged under  $U(1)_B$ . As noted previously, the Yukawa couplings for quarks and leptons via these Higgs fields are present, and realistic masses and mixings may be obtained. The Yukawa couplings with the Higgs fields  $\mathcal{H}_u, \mathcal{H}_d$  from the  $bc'$  sector, which carry charges of  $\pm 1$  under  $U(1)_B \equiv U(1)_{LP}$  respectively, that

TABLE VIII. The chiral hidden sector superfields, their multiplicities and quantum numbers under the gauge symmetry  $[SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_B]_{\text{observable}} \times [SU(2) \times USp(2)^2]_{\text{hidden}}$ .

	Mult.	Quantum Number	$Q_{I3R}$	$Q_{B-L}$	$Q_{3B+L}$	$Q_Y$	$Q_B$	Field
$bd$	1	$(1, \bar{2}, 1, 2, 1, 1, 1)$	0	0	-4	0	-1	$X_{bd}$
$c1d$	1	$(1, 1, 1, \bar{2}, 1, 1, 1)$	1/2	0	4	1/2	1	$X_{c1d}$
$c2d$	1	$(1, 1, 1, \bar{2}, 1, 1, 1)$	-1/2	0	4	-1/2	1	$X_{c2d}$
$b4$	3	$(1, \bar{2}, 1, 1, 1, 2, 1)$	0	0	-2	0	-1/2	$X_{b3}^i$
$c13$	3	$(1, 1, 1, 1, 1, \bar{2}, 1)$	1/2	0	-2	1/2	-1/2	$X_{c13}^i$
$c23$	3	$(1, 1, 1, 1, 1, \bar{2}, 1)$	-1/2	0	-2	-1/2	-1/2	$X_{c23}^i$
$d3$	1	$(1, 1, 1, 1, \bar{2}, 2, 1)$	0	0	-6	0	-3/2	$X_{cd}$
$d4$	1	$(1, 1, 1, 1, 2, 1, \bar{2})$	0	0	-6	0	-3/2	$X_{cd}$
$b_S$	2	$(1, 3, 1, 1, 1, 1, 1)$	0	0	-4	0	-1	$T_L^i$
$b_A$	2	$(1, 1, 1, 1, 1, 1, 1)$	0	0	4	0	1	$S_L^i$
$c_S$	2	$(1, 1, 1, 1, 1, 1, 1)$	0	0	-4	0	-1	$T_R^i$

these fields may also obtain a subdominant VEV with respect to the Higgs fields in the  $bc$  sector. The  $Z$  and  $Z'$  bosons will then be mixed as a result. Thus, we might explain the diboson and dijet excesses [7].

Clearly, the requirement that a Higgs field be charged under the leptophobic  $U(1)$  in order to obtain mixing between the  $Z$  and  $Z'$  bosons results in the Yukawa couplings with this Higgs field being forbidden, which seems to be a generic problem for models of this type. In the present context, this leads to the requirement of an extended Higgs sector with some Higgs fields charged under  $U(1)_B$  and some which are not for which the Yukawa couplings are present. Specifically, the Yukawa couplings with Higgs fields in the  $bc$  sector are allowed by the global  $U(1)$  charges carried by these fields:

$$W_Y = y_U^{ijk} H_u^i Q_L^j U_R^k + y_D^{ijk} H_d^i Q_L^j D_R^k + y_\nu^{ijk} H_u^i L_L^j N_R^k + y_e^{ijk} H_d^i L^j E_R^k, \quad (12)$$

while the Yukawa couplings with the extra Higgs fields  $\mathcal{H}_u$  and  $\mathcal{H}_d$  from the  $bc'$  sector which are charged under  $U(1)_B$  are perturbatively forbidden. We assume that these extra Higgs fields have masses so that they have not been observed at the LHC. We shall defer a detailed study of such extra Higgs bosons to later work.

#### IV. DIBOSON SIGNALS AT THE 8 TEV LHC

The production of a leptophobic  $Z'$  boson at the 8 TeV LHC is strongly constrained by three independent search regions:  $Z' \rightarrow jj$  [14],  $Z' \rightarrow t\bar{t}$  [65], and  $Z' \rightarrow Zh$  [4]. Given the anomaly free nature of the  $Z'$  boson described in the models presented here, the  $Z' \rightarrow ZZ$  and  $Z' \rightarrow Z\gamma$  are forbidden. Therefore, we only apply the following three constraints:

$$\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow jj) \sim 91_{-45}^{+53} \text{ fb (1}\sigma \text{ fit)} \quad (13)$$

$$\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow t\bar{t}) \lesssim 11(18) \text{ fb (95\% C.L.)} \quad (14)$$

$$\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow Zh) \lesssim 7 \text{ fb (95\% C.L.)} \quad (15)$$

The  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow jj) \sim 91_{-45}^{+53}$  fb constraint consists of a  $1\sigma$  fitted cross section [14], thus we also consider a more relaxed alternative. Softening these  $1\sigma$  boundaries, we shall also observe the result of applying only a 90% C.L. upper bound of  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow jj) \lesssim 170$  fb [14]. The upper limit on  $t\bar{t}$  resonances of 11 fb established by the CMS experiment [65] corresponds to a decay width of 20 GeV for a 2 TeV  $Z'$  boson, whereas the 18 fb upper limit correlates to a 200 GeV decay width, though we shall generally only regard the less stringent 18 fb constraint in this analysis when phenomenologically constraining the gauge coupling  $g_{Z'}$ .

The calculation of the partial decay widths requires the quark and Higgs field charges on the leptophobic  $U(1)_{LP}$ . The quark decay width is given by [7]

$$\Gamma(Z' \rightarrow q\bar{q}) = \frac{g_{Z'}^2 N_C}{24\pi} M_{Z'} \left[ Q_{LP_{q_L}}^2 + Q_{LP_{q_R}}^2 - (Q_{LP_{q_L}} - Q_{LP_{q_R}})^2 \left( \frac{m_q}{M_{Z'}} \right)^2 \right] \sqrt{1 - 4 \left( \frac{m_q}{M_{Z'}} \right)^2} \quad (16)$$

where  $g_{Z'}$  is the  $U(1)_{LP}$  gauge coupling,  $N_C = 3$  represents the number of colors, and  $Q_{LP_{q_L}}$ ,  $Q_{LP_{q_R}}$  are the left- and right-handed charges of the quark content on  $U(1)_{LP}$ . Here we take  $M_{Z'} = 2$  TeV and  $m_t = 174.4$  GeV [66,67]. The  $\Gamma(Z' \rightarrow W^+W^-)$  decay width can be computed from [7]

$$\Gamma(Z' \rightarrow W^+W^-) = \frac{g_{Z'}^2}{48\pi} M_{Z'} Q_{LP_{H_u}}^2 \sin^4 \beta \quad (17)$$

with  $Q_{LP_{H_u}}$  as the Higgs field charge on  $U(1)_{LP}$ , and  $\beta$  the angle between the up and down Higgs VEVs. The equivalence theorem suggests for a heavy  $Z'$  boson in the decoupling limit that

$$\Gamma(Z' \rightarrow Zh) = \Gamma(Z' \rightarrow W^+W^-) \quad (18)$$

indicating that the branching ratio for these two decay modes are equivalent. As a result, application of the strong upper limit  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow Zh) \lesssim 7$  fb likewise tightly constrains our  $Z' \rightarrow W^+W^-$  production as well. Proper normalization is required for accurate  $U(1)_{LP}$  gauge coupling evolution, presumably achieving unification with the  $SU(3)_C$ ,  $SU(2)_L$ , and  $U(1)_Y$  gauge couplings at the GUT scale. For the flipped  $SU(5) \times U(1)_X$  models, the explicit normalization factor is  $\frac{1}{3}$ . All massless fields have conformal dimension 1, generating the transformation  $Q' \rightarrow Q'/\sqrt{3}$ , where  $Q'$  are the charges on  $U'$  given in Tables I, II, III, and IV. Therefore, the  $U'$  gauge coupling evolves according to the beta function  $b' = \frac{1}{3} \text{Tr}(Q')^2$ . For convenience, this factor of  $\frac{1}{3}$  in the decay widths is introduced directly into the calculation of the  $\sigma(pp \rightarrow Z')$  cross section in our numerical results via a normalization factor  $n$  in Eq. (19), explicitly implementing  $n = \frac{1}{3}$  for flipped models. For the intersecting D-brane model, an explicit normalization factor on  $U(1)_B$  is not yet known, though it is expected that it is of unity order, hence, we assume any normalization of  $\mathcal{O}(1)$  is already assimilated into the phenomenologically constrained value of the  $U(1)_B$  gauge coupling  $g_{Z'}$ . We thus apply  $n = 1$  in Eq. (19) for intersecting D-branes. An estimate of the 8 TeV LHC cross section for  $Z'$  boson production is given as [7,27,68]

TABLE IX. Computed branching ratios for  $Z'$  boson decay for the four channels realized in leptophobic flipped  $SU(5) \times U(1)_X$  and intersecting D-brane models. Given the absence of any electrically neutral particles at the electroweak scale in either model, there are no invisible decay modes.

	$\text{Br}(Z' \rightarrow jj)$	$\text{Br}(Z' \rightarrow t\bar{t})$	$\text{Br}(Z' \rightarrow W^+W^-)$	$\text{Br}(Z' \rightarrow Zh)$
Flipped $SU(5) \times U(1)_X$	0.870	0.078	0.026	0.026
Intersecting D-branes	0.670	0.130	0.100	0.100

$$\sigma(pp \rightarrow Z') \simeq n \times \left[ 5.2 \left( \frac{2\Gamma(Z' \rightarrow u\bar{u}) + \Gamma(Z' \rightarrow d\bar{d})}{\text{GeV}} \right) \right] \text{ fb} \quad (19)$$

using the decay widths given in Eq. (16).

The charges of the MSSM content on  $U'$  are given in Table I for leptophobic flipped  $SU(5) \times U(1)_X$  models. The matter fields  $F_3, F_1$ , and  $F_4$  represent the first, second, and third generations, respectively, and  $\bar{h}_1$  contains  $H_u$ . Thus, to compute the decay widths and branching ratios, the  $U' = U(1)_{LP}$  charges used are  $Q_{LP_{u_L}} = Q_{LP_{u_R}} = Q_{LP_{d_L}} = Q_{LP_{d_R}} = 1$ ,  $Q_{LP_{c_L}} = Q_{LP_{c_R}} = Q_{LP_{s_L}} = Q_{LP_{s_R}} = -\frac{1}{2}$ ,  $Q_{LP_{t_L}} = Q_{LP_{t_R}} = Q_{LP_{b_L}} = Q_{LP_{b_R}} = -\frac{1}{2}$ , and  $Q_{LP_{H_u}} = -1$ .

A review of Table I containing the MSSM content charges in the observable sector for the flipped  $SU(5) \times U(1)_X$  model shows there are no electrically neutral particles at the electroweak scale charged on  $U'$ , hence there are no invisible decays. The only  $Z'$  decay modes present then are  $Z' \rightarrow jj$ ,  $Z' \rightarrow t\bar{t}$ ,  $Z' \rightarrow WW$ , and  $Z' \rightarrow Zh$ . Though the No-scale supergravity boundary conditions at the unification scale are not applied here, we do implement a preferred no-scale supergravity angle  $\beta$  using  $\tan\beta = 25$  [69]. The decay widths were computed with both  $\tan\beta = 5$  and  $\tan\beta = 25$ , resulting in only a mere  $\sim 5\%$  increase in the cross section for the larger  $\tan\beta$ , a safely negligible delta for our purposes here. Given this lack of significant variation in the cross section as a function of  $\beta$ , there is essentially only




FIG. 1. Depiction of the cross section  $\sigma$  as a function of the  $Z'$  boson gauge coupling  $g_{Z'}$  on  $U(1)_{LP}$  for the flipped  $SU(5) \times U(1)_X$  models (left frame) and the intersecting D-brane model (right frame). Here we only consider the LHC constraints  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow jj) \sim 91_{-45}^{+53}$  fb,  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow Zh) \lesssim 7$  fb, and the 200 GeV decay width  $t\bar{t}$  constraint of  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow t\bar{t}) \lesssim 18$  fb. Those  $g_{Z'}$  satisfying these three constraints are shown as solid, thick sections of the curves, with the intersection marked by the cross-hatched region, where  $g_{Z'} \sim 0.25\text{--}0.45$  for the flipped models and  $g_{Z'} \sim 0.5$  for D-branes. If the LHC constraint on  $Z' \rightarrow jj$  is relaxed to only an upper limit of  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow jj) \lesssim 170$  fb, then loose model constraints of  $g_{Z'} \lesssim 0.49$  for flipped models and  $g_{Z'} \lesssim 0.50$  for D-branes are obtained.



one free-parameter remaining, the  $Z'$  gauge coupling  $g_{Z'}$ . Therefore, we phenomenologically constrain the value of  $g_{Z'}$  using the LHC constraints on  $Z' \rightarrow jj$ ,  $Z' \rightarrow t\bar{t}$ , and  $Z' \rightarrow Zh$ . The branching ratios are independent of variation in  $g_{Z'}$ , with the results of the computations for the flipped  $SU(5) \times U(1)_X$  models given in Table IX.

Observe in Table VII containing the MSSM content charges for intersecting D-branes that there are no electrically neutral particles at the electroweak scale charged on  $U(1)_B$ , hence there are also no invisible decays in the D-brane model. Likewise, the only  $Z'$  decay modes present then are  $Z' \rightarrow jj$ ,  $Z' \rightarrow t\bar{t}$ ,  $Z' \rightarrow WW$ , and  $Z' \rightarrow Zh$ . To calculate the decay widths and branching ratios, the  $U(1)_B = U(1)_{LP}$  charges employed are  $Q_{LP_{Q_L}} = 1/3$ ,  $Q_{LP_{U_R}} = Q_{LP_{D_R}} = -1/3$ , and  $Q_{LP_{\gamma_{tu}}} = 1$ . As with the flipped  $SU(5) \times U(1)_X$  models, the value of  $g_{Z'}$  is constrained via the LHC constraints. We use  $\tan\beta = 25$  for D-branes also. The intersecting D-brane branching ratios are included in Table IX.

The value of the  $Z'$  gauge coupling  $g_{Z'}$  is freely floated prior to application of the LHC constraints given in Eqs. (13)–(15). The individual cross sections as a function of the  $Z'$  gauge coupling are shown in Fig. 1 for both the flipped  $SU(5) \times U(1)_X$  and intersecting D-brane models. The dashed and dotted lines in Fig. 1 represent all values of the gauge couplings, while the thick, solid sections are only those values of  $g_{Z'}$  that satisfy the constraints of

Eqs. (13)–(15). It is clear that implementation of the  $Z' \rightarrow jj$  fitted  $1\sigma$  deviation very tightly constrains the value of the  $U(1)_{LP}$  gauge coupling to  $g_{Z'} \sim 0.25$ – $0.45$  for the flipped models and  $g_{Z'} \sim 0.50$  for D-branes. While these tight model constraints on  $g_{Z'}$  are rather predictive, particularly for intersecting D-branes, it does leave little room for deviation in the event future enhancements of the constraints are necessary. Given the modest accumulation of only about  $20 \text{ fb}^{-1}$  of luminosity thus far at 8 TeV, and awaiting the forthcoming deluge of data extracted from the 13/14 TeV beam collision energies, we alternatively relax the  $Z' \rightarrow jj$  constraint to only an upper limit on the cross section of  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow jj) \lesssim 170 \text{ fb}$ . This releases the lower values of  $g_{Z'}$  as viable candidates, providing loose model constraints of  $g_{Z'} \lesssim 0.49$  for flipped models and  $g_{Z'} \lesssim 0.50$  for D-branes. The 13/14 TeV data allocation arriving in the year 2015 and beyond shall reduce the experimental measurement uncertainties and should ultimately merge the tight and loose model constraints presented here.

## ACKNOWLEDGMENTS

This research was supported in part by the Natural Science Foundation of China under Grants No. 11135003, No. 11275246, and No. 11475238 (T.L.), and by the DOE Grant No. DE-FG02-13ER42020 (D.V.N.).

- 
- [1] G. Aad *et al.* (ATLAS Collaboration), Search for high-mass diboson resonances with boson-tagged jets in proton-proton collisions at  $\sqrt{s} = 8 \text{ TeV}$  with the ATLAS detector, *J. High Energy Phys.* **12** (2015) 055.
  - [2] V. Khachatryan *et al.* (CMS Collaboration), Search for massive resonances in dijet systems containing jets tagged as W or Z boson decays in pp collisions at  $\sqrt{s} = 8 \text{ TeV}$ , *J. High Energy Phys.* **08** (2014) 173.
  - [3] V. Khachatryan *et al.* (CMS Collaboration), Search for massive resonances decaying into pairs of boosted bosons in semi-leptonic final states at  $\sqrt{s} = 8 \text{ TeV}$ , *J. High Energy Phys.* **08** (2014) 174.
  - [4] V. Khachatryan *et al.* (CMS Collaboration), Search for a massive resonance decaying into a Higgs boson and a W or Z boson in hadronic final states in proton-proton collisions at  $\sqrt{s} = 8 \text{ TeV}$ , [arXiv:1506.01443](https://arxiv.org/abs/1506.01443).
  - [5] CMS Collaboration, Note No. PAS-EXO-14-010, 2015.
  - [6] H. S. Fukano, M. Kurachi, S. Matsuzaki, K. Terashi, and K. Yamawaki, 2 TeV walking technirho at LHC? *Phys. Lett. B* **750**, 259 (2015).
  - [7] J. Hisano, N. Nagata, and Y. Omura, Interpretations of the ATLAS diboson resonances, *Phys. Rev. D* **92**, 055001 (2015).
  - [8] D. Gerosa, M. Kesden, U. Sperhake, E. Berti, and R. O’Shaughnessy, A multi-timescale analysis of phase transitions in precessing black-hole binaries, *Phys. Rev. D* **92**, 064016 (2015).
  - [9] K. Cheung, W. Y. Keung, P. Y. Tseng, and T. C. Yuan, Interpretations of the ATLAS diboson anomaly, *Phys. Lett. B* **751**, 188 (2015).
  - [10] B. A. Dobrescu and Z. Liu,  $W'$  Boson near 2 TeV: Predictions for Run 2 of the LHC, *Phys. Rev. Lett.* **115**, 211802 (2015).
  - [11] A. Alves, A. Berlin, S. Profumo, and F. S. Queiroz, Dirac-fermionic dark matter in  $U(1)_X$  models, *J. High Energy Phys.* **10** (2015) 076.
  - [12] Y. Gao, T. Ghosh, K. Sinha, and J. H. Yu,  $SU(2) \times SU(2) \times U(1)$  interpretations of the diboson and  $Wh$  excesses, *Phys. Rev. D* **92**, 055030 (2015).
  - [13] A. Thamm, R. Torre, and A. Wulzer, Composite Heavy Vector Triplet in the ATLAS Diboson Excess, *Phys. Rev. Lett.* **115**, 221802 (2015).
  - [14] J. Brehmer, J. Hewett, J. Kopp, T. Rizzo, and J. Tattersall, Symmetry restored in dibosons at the LHC? *J. High Energy Phys.* **10** (2015) 182.

- [15] Q. H. Cao, B. Yan, and D. M. Zhang, Simple non-Abelian extensions and diboson excesses at the LHC, *Phys. Rev. D* **92**, 095025 (2015).
- [16] G. Cacciapaglia and M. T. Frandsen, Unitarity implications of diboson resonance in the TeV region for Higgs physics, *Phys. Rev. D* **92**, 055035 (2015).
- [17] T. Abe, R. Nagai, S. Okawa, and M. Tanabashi, Unitarity sum rules, three site moose model, and the ATLAS 2 TeV diboson anomalies, *Phys. Rev. D* **92**, 055016 (2015).
- [18] B. C. Allanach, B. Gripaios, and D. Sutherland, Anatomy of the ATLAS diboson anomaly, *Phys. Rev. D* **92**, 055003 (2015).
- [19] T. Abe, T. Kitahara, and M. M. Nojiri, Prospects for spin-1 resonance search at 13 TeV LHC and the ATLAS diboson excess, [arXiv:1507.01681](https://arxiv.org/abs/1507.01681).
- [20] A. Carmona, A. Delgado, M. Quiros and J. Santiago, Diboson resonant production in non-custodial composite Higgs models, *J. High Energy Phys.* **09** (2015) 186.
- [21] C. W. Chiang, H. Fukuda, K. Harigaya, M. Ibe, and T. T. Yanagida, Diboson resonance as a portal to hidden strong dynamics, *J. High Energy Phys.* **11** (2015) 015.
- [22] G. Cacciapaglia, A. Deandrea, and M. Hashimoto, A Scalar Hint from the Diboson Excess?, *Phys. Rev. Lett.* **115**, 171802 (2015).
- [23] H. S. Fukano, S. Matsuzaki, and K. Yamawaki, Conformal barrier for new vector bosons decay to the Higgs, [arXiv:1507.03428](https://arxiv.org/abs/1507.03428).
- [24] V. Sanz, On the compatibility of the diboson excess with a gg-initiated composite sector, [arXiv:1507.03553](https://arxiv.org/abs/1507.03553).
- [25] C. H. Chen and T. Nomura, 2 TeV Higgs boson and diboson excess at the LHC, *Phys. Lett. B* **749**, 464 (2015).
- [26] Y. Omura, K. Tobe, and K. Tsumura, Survey of Higgs interpretations of the diboson excesses, *Phys. Rev. D* **92**, 055015 (2015).
- [27] L. A. Anchordoqui, I. Antoniadis, H. Goldberg, X. Huang, D. Lust, and T. R. Taylor, Stringy origin of diboson and dijet excesses at the LHC, *Phys. Lett. B* **749**, 484 (2015).
- [28] W. Chao, ATLAS Diboson excesses from the stealth doublet model, *Phys. Lett. B* **753**, 117 (2016).
- [29] L. Bian, D. Liu and J. Shu, Low scale composite Higgs model and 1.8–2 TeV diboson excess, [arXiv:1507.06018](https://arxiv.org/abs/1507.06018).
- [30] D. Kim, K. Kong, H. M. Lee, and S. C. Park, Diboson excesses demystified in effective field theory approach, *J. High Energy Phys.* **11** (2015) 150.
- [31] K. Lane and L. Prichett, Heavy vector partners of the light composite Higgs, *Phys. Lett. B* **753**, 211 (2016).
- [32] A. E. Faraggi and M. Guzzi, Extra  $Z$ 's and  $W$ 's in heterotic-string derived models, *Eur. Phys. J. C* **75**, 537 (2015).
- [33] M. Low, A. Tesi, and L. T. Wang, Composite spin-1 resonances at the LHC, *Phys. Rev. D* **92**, 085019 (2015).
- [34] S. P. Liew and S. Shirai, Testing ATLAS diboson excess with dark matter searches at LHC, *J. High Energy Phys.* **11** (2015) 191.
- [35] H. Terazawa and M. Yasue, Excited gauge and Higgs bosons in the unified composite model, [arXiv:1508.00172](https://arxiv.org/abs/1508.00172).
- [36] P. Arnan, D. Espriu, and F. Mescia, Interpreting a 2 TeV resonance in WW scattering, *Phys. Rev. D* **93** 015020 (2016).
- [37] C. Niehoff, P. Stangl, and D. M. Straub, Direct and indirect signals of natural composite Higgs models, [arXiv:1508.00569](https://arxiv.org/abs/1508.00569).
- [38] D. Goncalves, F. Krauss, and M. Spannowsky, Augmenting the diboson excess for Run 2, *Phys. Rev. D* **92**, 053010 (2015).
- [39] S. Fichtel and G. von Gersdorff, Effective theory for neutral resonances and a statistical dissection of the ATLAS diboson excess, *J. High Energy Phys.* **12** (2015) 089.
- [40] C. Petersson and R. Torre, ATLAS diboson excess from low scale supersymmetry breaking, [arXiv:1508.05632](https://arxiv.org/abs/1508.05632).
- [41] F. F. Deppisch, L. Graf, S. Kulkarni, S. Patra, W. Rodejohann, N. Sahu, and U. Sarkar, Reconciling the 2 TeV excesses at the LHC in a linear seesaw left-right model, *Phys. Rev. D* **93**, 013011 (2016).
- [42] J. A. Aguilar-Saavedra, Triboson interpretations of the ATLAS diboson excess, *J. High Energy Phys.* **10** (2015) 099.
- [43] L. Bian, D. Liu, J. Shu, and Y. Zhang, Interference effect on resonance studies and the diboson excess, [arXiv:1509.02787](https://arxiv.org/abs/1509.02787).
- [44] P. S. Bhupal Dev and R. N. Mohapatra, Unified Explanation of the  $eejj$ , Diboson, and Dijet Resonances at the LHC, *Phys. Rev. Lett.* **115**, 181803 (2015).
- [45] D. B. Franzosi, M. T. Frandsen, and F. Sannino, Diboson signals via Fermi scale spin-one States, *Phys. Rev. D* **92**, 115005 (2015).
- [46] S. M. Barr, A new symmetry breaking pattern for SO(10) and proton decay, *Phys. Lett. B* **112**, 219 (1982); J. P. Derendinger, J. E. Kim, and D. V. Nanopoulos, Anti-Su(5), *Phys. Lett.* **139B**, 170 (1984); I. Antoniadis, J. R. Ellis, J. S. Hagelin, and D. V. Nanopoulos, Supersymmetric flipped SU(5) revitalized, *Phys. Lett. B* **194**, 231 (1987).
- [47] J. L. Lopez, D. V. Nanopoulos, and K. Yuan, The search for a realistic flipped SU(5) string model, *Nucl. Phys.* **B399**, 654 (1993).
- [48] J. L. Lopez and D. V. Nanopoulos, String no-scale supergravity, [arXiv:hep-ph/9412332](https://arxiv.org/abs/hep-ph/9412332); Vanishing Str  $M^2$  in the presence of anomalous  $U_A(1)$ , *Phys. Lett. B* **369**, 243 (1996); New Scenario for String Unification, *Phys. Rev. Lett.* **76**, 1566 (1996); J. L. Lopez, D. V. Nanopoulos, and A. Zichichi, String no-scale supergravity model and its experimental consequences, *Phys. Rev. D* **52** (1995) 4178; Experimental constraints on a stringy SU(5) $\times$ U(1) model, *Phys. Rev. D* **53**, 5253 (1996); J. Ellis, J. L. Lopez, and D. V. Nanopoulos, Lowering  $\alpha_s$  by flipping SU(5), *Phys. Lett. B* **371**, 65 (1996). For a review, see J. L. Lopez and D. V. Nanopoulos, Flipped SU(5): A Grand Unified Superstring Theory (GUST) Prototype, [arXiv:hep-ph/9511266](https://arxiv.org/abs/hep-ph/9511266).
- [49] J. L. Lopez and D. V. Nanopoulos, Hierarchical fermion masses and mixing angles from the flipped string, *Nucl. Phys.* **B338**, 73 (1990).
- [50] J. L. Lopez and D. V. Nanopoulos, Leptophobic  $Z'$  in stringy flipped SU(5), *Phys. Rev. D* **55**, 397 (1997).
- [51] J. Jiang, T. Li, and D. V. Nanopoulos, Testable flipped SU(5)  $\times$  U(1) $_X$  models, *Nucl. Phys.* **B772**, 49 (2007).
- [52] M. Cvetič, G. Shiu, and A. M. Uranga, Three-Family Supersymmetric Standardlike Models from Intersecting Brane Worlds, *Phys. Rev. Lett.* **87**, 201801 (2001); Chiral

- four-dimensional  $N = 1$  supersymmetric type IIA orientifolds from intersecting D6-branes, *Nucl. Phys.* **B615**, 3 (2001).
- [53] M. Cvetič, T. Li, and T. Liu, Supersymmetric Pati-Salam models from intersecting D6-branes: A road to the standard model, *Nucl. Phys.* **B698**, 163 (2004).
- [54] C. M. Chen, T. Li and D. V. Nanopoulos, Standard-like model building on Type II orientifolds, *Nucl. Phys.* **B732**, 224 (2006).
- [55] C. M. Chen, T. Li, and D. V. Nanopoulos, Type IIA Pati-Salam flux vacua, *Nucl. Phys.* **B740**, 79 (2006).
- [56] C. M. Chen, T. Li, and D. V. Nanopoulos, Flipped and unflipped SU(5) as type IIA flux vacua, *Nucl. Phys.* **B751**, 260 (2006).
- [57] C. M. Chen, T. Li, V. E. Mayes, and D. V. Nanopoulos, A realistic world from intersecting D6-branes, *Phys. Lett. B* **665**, 267 (2008).
- [58] C. M. Chen, T. Li, V. E. Mayes, and D. V. Nanopoulos, Towards realistic supersymmetric spectra and Yukawa textures from intersecting branes, *Phys. Rev. D* **77**, 125023 (2008).
- [59] J. A. Maxin, V. E. Mayes, and D. V. Nanopoulos, Proton stability and dark matter in a realistic string MSSM, *Phys. Rev. D* **84**, 106009 (2011).
- [60] C. M. Chen, T. Li, V. E. Mayes, and D. V. Nanopoulos, Variations of the hidden sector in a realistic intersecting brane model, *J. Phys. G* **35**, 095008 (2008).
- [61] C.-M. Chen, G. V. Kraniotis, V. E. Mayes, D. V. Nanopoulos, and J. W. Walker, A supersymmetric flipped SU(5) intersecting brane world, *Phys. Lett. B* **611**, 156 (2005).
- [62] C.-M. Chen, G. V. Kraniotis, V. E. Mayes, D. V. Nanopoulos, and J. W. Walker, A K-theory anomaly free supersymmetric flipped SU(5) model from intersecting branes, *Phys. Lett. B* **625**, 96 (2005).
- [63] C.-M. Chen, V. E. Mayes, and D. V. Nanopoulos, Flipped SU(5) from D-branes with type IIB fluxes, *Phys. Lett. B* **633**, 618 (2006).
- [64] M. Dine, N. Seiberg, and E. Witten, Fayet-Iliopoulos terms in string theory, *Nucl. Phys.* **B289**, 589 (1987); J. Atick, L. Dixon, and A. Sen, String calculation of fayet-iliopoulos D-terms in arbitrary supersymmetric compactifications, *Nucl. Phys.* **B292**, 109 (1987); M. Dine, I. Ichinose, and N. Seiberg, F terms and D terms in string theory, *Nucl. Phys.* **B293**, 253 (1987).
- [65] V. Khachatryan *et al.* (CMS Collaboration), Search for resonant  $t\bar{t}$  production in proton-proton collisions at  $\sqrt{s} = 8$  TeV, *Phys. Rev. D* **93**, 012001 (2016).
- [66] Tevatron Electroweak Working Group (CDF and D0 Collaborations), Combination of CDF and D0 results on the mass of the top quark using up to  $9.7 \text{ fb}^{-1}$  at the Tevatron, [arXiv:1407.2682](https://arxiv.org/abs/1407.2682).
- [67] T. Leggett, T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, Confronting electroweak fine-tuning with no-scale supergravity, *Phys. Lett. B* **740**, 66 (2015).
- [68] A. Leike, The phenomenology of extra neutral gauge bosons, *Phys. Rep.* **317**, 143 (1999).
- [69] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, No-Scale F-SU(5) in the Light of LHC, Planck and XENON, *J. Phys. G* **40**, 115002 (2013).