Approximation of the potential in scalar field dark energy models

Richard A. Battye[†] and Francesco Pace^{*}

Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester,

Manchester M13 9PL, United Kingdom

(Received 9 July 2016; published 13 September 2016)

We study the nature of potentials in scalar field based models for dark energy—with both canonical and noncanonical kinetic terms. We calculate numerically, and using an analytic approximation around $a \approx 1$, potentials for models with constant equation-of-state parameter w_{ϕ} . We find that for a wide range of models with canonical and noncanonical kinetic terms there is a simple approximation for the potential that holds when the scale factor is in the range $0.6 \leq a \leq 1.4$. We discuss how this form of the potential can also be used to represent models with nonconstant w_{ϕ} and, hence, how it could be used in reconstruction from cosmological data.

DOI: 10.1103/PhysRevD.94.063513

I. INTRODUCTION

The origin of the cosmic acceleration is one of the most significant open questions in cosmology and fundamental physics. A cosmological constant is still very much consistent with the data [1,2], but in order to either refute or confirm this simple hypothesis one needs to consider alternative models to explain the observations. One very simple idea is to postulate a dark energy component dominated by a scalar field either with a canonical or noncanonical kinetic term. Such models are known as quintessence models [3–10] and *k*-essence models [11–13], respectively.

The standard approach when constraining cosmological models with a dark energy component that is not the cosmological constant is to define an equation-of-state parameter $w_{\phi} = P_{\phi}/\rho_{\phi} \neq -1$, where P_{ϕ} is the pressure of dark energy and ρ_{ϕ} is its density, making no assumption as to the origin of the dark energy. In principle this is a general function of time, but it is often considered to be either constant or to be represented by a specific functional form, for example [14,15]. At the moment the data barely constrain anything beyond a constant w_{ϕ} , but this is likely to change in the near future as more observations probing the equation of state become available, such as Euclid [16–18], LSST [19,20] and SKA [21–25]. Various ideas have been put forward to extend to time-varying situations. These include various limited functional forms [14,15,26–28], the *Om* diagnostic [29,30], the state-finder approach [31,32] and even using principal component analysis on general piecewise linear parametrizations of w_{ϕ} [33]. For a review of the parametric and nonparametric methods to reconstruct the dark energy equation-of-state parameter, we refer to [34]. Since many of the observations are sensitive to perturbations in the dark energy it is also necessary to make some assumptions about the perturbations, but we will not consider this here.

An alternative is to presume that the origin of the dark energy is a model based on a scalar field. However, such models usually involve one or more arbitrary functions which would need to be specified before any model prediction could be made. One of these is the potential $V(\phi)$ of the scalar field which one might try to reconstruct from observations. One obvious suggestion [35], which extends the approach of [36] for inflation, is to represent the potential as a Taylor series expanded around the presentday value of the field ϕ_0 :

$$V(\phi) = V_0 + V_1(\phi - \phi_0) + V_2(\phi - \phi_0)^2 + \cdots$$
(1)

and attempt to fit for the coefficients V_i . However, it is not clear where to truncate this series in a controlled way. Similar and complementary methods have been proposed by [37–40]. Other reconstruction methods are valid in the slow-roll regime, that is, when $1 + w_{\phi} \approx 0$. For quintessence models, a one-parameter [41] or two-parameter [42–44] formula has been used and for *k*-essence models we refer to works by [11,45].

In this paper we first calculate potentials for a range of minimally coupled scalar field models with canonical (Sec. II) and noncanonical (Sec. III) kinetic terms assuming initially that w_{ϕ} is constant. It is possible to derive an analytic solution for the potential in quintessence models, but this is not possible in general for the case of *k*-essence models and therefore we resort to numerical calculations and an analytic approximation around the present day which is valid for $0.6 \leq a \leq 1.4$. Based on this analytic approximation we suggest a form of a potential with just four parameters which we demonstrate can lead to a wide range of behavior for w_{ϕ} as a function of time (Sec. IV) and, by design, includes models with constant w_{ϕ} . Of course,

^{*}Corresponding author.

francesco.pace@manchester.ac.uk

[†]richard.battye@manchester.ac.uk

this functional form will not include every possible behavior in a general model, but it does provide more physical insights and it is useful for models which are not significantly different from a linearly evolving equation-ofstate parameter. We conclude and discuss our results in Sec. V.

In the following, we will use natural units with $c = \hbar = 1$, the Planck mass is $M_{\rm pl} = G^{-1/2}$ and we assume a metric with signature (-, +, +, +).

II. MINIMALLY COUPLED SCALAR FIELDS WITH CONSTANT w_{ϕ}

The Lagrangian for minimally coupled scalar fields is

$$\mathcal{L} = -\frac{1}{2}\eta g^{\mu\nu} \nabla_{\mu} \phi \nabla_{\nu} \phi - V(\phi), \qquad (2)$$

and its corresponding stress-energy tensor

$$T_{\mu\nu} = g_{\mu\nu}\mathcal{L} + \eta \nabla_{\mu}\phi \nabla_{\nu}\phi$$
$$= \eta \nabla_{\mu}\phi \nabla_{\nu}\phi - g_{\mu\nu} \left[\frac{1}{2}\eta g^{\alpha\beta} \nabla_{\alpha}\phi \nabla_{\beta}\phi + V(\phi)\right]. \quad (3)$$

The constant η distinguishes between the quintessence case $(\eta = +1, -1 < w_{\phi} < 1)$ and the phantom case $(\eta = -1, w_{\phi} < -1)$ [46].

Density and pressure are given by

$$\rho_{\phi} = T^{0}{}_{0} = \frac{1}{2}\eta\dot{\phi}^{2} + V(\phi),
P_{\phi} = \frac{1}{3}T^{i}{}_{i} = \frac{1}{2}\eta\dot{\phi}^{2} - V(\phi),$$
(4)

and the conservation equation $\dot{\rho}_{\phi} + 3H(\rho_{\phi} + P_{\phi}) = 0$ gives rise to the Klein-Gordon equation, which describes the time evolution of the scalar field

$$\ddot{\phi} + 3H\dot{\phi} + \eta \frac{dV}{d\phi} = 0.$$
 (5)

To achieve an accelerated expansion, we require $w_{\phi} < -1/3$. In fact, observations require $w_{\phi} \simeq -1$ (due to the cosmological constant case) [1], and hence we can evaluate deviations of w_{ϕ} from -1 with the help of (5):

$$1 + w_{\phi} = \frac{V_{\phi}^2}{9H^2(\xi_s + 1)^2 \rho_{\phi}},\tag{6}$$

with $\xi_s = \ddot{\phi}/(3H\dot{\phi})$ [18]. Note that in a pure slow-roll approximation, $\xi_s = 0$.

By using Friedmann equations, we can determine the time evolution of the scalar field and its potential for a given $w_{\phi}(a)$ [47]:

$$\frac{\phi(a) - \phi_0}{M_{\rm pl}} = \pm \sqrt{\frac{3\Omega_{\rm de}}{8\pi}} \int_1^a \frac{\sqrt{\eta [1 + w_\phi(x)]g(x)}}{xE(x)} dx, \qquad (7)$$

$$V(a) = \frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}[1 - w_{\phi}(a)]g(a)}{16\pi},$$
(8)

where Ω_{de} is the dark energy density parameter today, H_0 the Hubble constant and ϕ_0 the value of the scalar field at a = 1. Finally, g(a) represents the time evolution of the dark energy component

$$g(a) = \exp\left(-3\int_{1}^{a} \frac{1+w_{\phi}(x)}{x} dx\right).$$
(9)

Assuming a flat geometry, the Hubble parameter is given by

$$H = H_0 E(a) = H_0 \left[\frac{\Omega_{\rm m}}{a^3} + \Omega_{\rm de} g(a) \right]^{\frac{1}{2}}, \qquad (10)$$

with Ω_m the matter density parameter today.

For a constant equation of state w_{ϕ} , integral (7) can be evaluated as

$$\frac{\phi - \phi_0}{M_{\rm pl}} = \mp \frac{2}{3w_{\phi}} \sqrt{\frac{3\eta(1 + w_{\phi})}{8\pi}} \times \left[\sinh^{-1}\left(\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}a^{-\frac{3w_{\phi}}{2}}\right) - \sinh^{-1}\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}\right], \quad (11)$$

and its inverse gives an expression for the scale factor in terms of the scalar field

$$a(\phi) = \left(\frac{\Omega_{\rm m}}{\Omega_{\rm de}}\right)^{-\frac{1}{3w_{\phi}}} \left[\sinh\left(\mp\frac{3}{2}w_{\phi}\sqrt{\frac{8\eta\pi}{3(1+w_{\phi})}}\left(\frac{\phi-\phi_{0}}{M_{\rm pl}}\right) + \sinh^{-1}\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}\right)\right]^{-\frac{2}{3w_{\phi}}}.$$
(12)

With these relations in hand, we can deduce the full expression for the potential $V(\phi)$:

$$V(\phi) = \frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}(1 - w_{\phi})}{16\pi} \left(\frac{\Omega_{\rm m}}{\Omega_{\rm de}}\right)^{\frac{1+w_{\phi}}{w_{\phi}}} \\ \times \left[\sinh\left(\mp \frac{3}{2}w_{\phi}\sqrt{\frac{8\eta\pi}{3(1+w_{\phi})}}\left(\frac{\phi - \phi_0}{M_{\rm pl}}\right) + \sinh^{-1}\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}\right)\right]^{\frac{2(1+w_{\phi})}{w_{\phi}}}.$$
 (13)

Since ϕ_0 just shifts the potential in the ϕ direction, we can make the choice

APPROXIMATION OF THE POTENTIAL IN SCALAR ...

$$\phi_0 = \pm \frac{2}{3w_{\phi}} M_{\rm pl} \sqrt{\frac{3\eta(1+w_{\phi})}{8\pi}} \sinh^{-1} \sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}},\qquad(14)$$

that simplifies the form of the potential

$$V(\phi) = \frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}(1 - w_{\phi})}{16\pi} \left(\frac{\Omega_{\rm m}}{\Omega_{\rm de}}\right)^{\frac{1+w_{\phi}}{w_{\phi}}} \sinh^{\frac{2(1+w_{\phi})}{w_{\phi}}} \times \left[\mp \frac{3w_{\phi}}{2} \sqrt{\frac{8\eta\pi}{3(1+w_{\phi})}} \frac{\phi}{M_{\rm pl}}\right],\tag{15}$$

in agreement with [34,48]. An analytic solution of (7) can be found also when dark matter has a constant equation-of-state parameter $w_m \neq 0$, as shown in [49].

While it is possible to find exact solutions for the potential of quintessence models with a constant equation of state, this is not the case when w_{ϕ} is a function of time or for more general scalar field models, such *k*-essence models. Moreover, the expressions for the potential at early and late times are not very useful from an observational point of view, since they assume one of the component to be dominant and are not relevant for modeling late-time observations. It is, therefore, worth-while to find approximate solutions valid for $a \approx 1$ that can be probed with data. To do this, we expand in series $\frac{d\phi}{da}$ for $a \approx 1$, but *a priori* it is not clear where to truncate the series. We have checked that a first-order expansion is a very good approximation, leading to a scalar field evolving quadratically with respect to the scale factor.

The differential equation describing the approximate evolution of the scalar field for $a \approx 1$ is

$$\frac{d\phi}{da} = \pm \sqrt{\frac{3\eta (1+w_{\phi})M_{\rm pl}^2 \Omega_{\rm de}}{8\pi}} \bigg\{ 1 - \frac{1}{2} (2 + 3\Omega_{\rm m} w_{\phi}) (a-1) \bigg\},$$
(16)

which implies the following approximate evolution for the scalar field:

$$\frac{\phi - \phi_0}{M_{\rm pl}} = \pm \sqrt{\frac{3\eta (1 + w_\phi)\Omega_{\rm de}}{8\pi}} \times \left[a - 1 - \frac{1}{4} (2 + 3\Omega_{\rm m} w_\phi) (a - 1)^2 \right].$$
(17)

By inverting this, we can find a relation between the scale factor and the scalar field

$$a = 1 + \frac{1}{2 + 3\Omega_{\rm m}w_{\phi}} \times \left\{ 1 - \sqrt{1 \mp [2 + 3\Omega_{\rm m}w_{\phi}]} \sqrt{\frac{8\eta\pi}{3\Omega_{\rm de}[1 + w_{\phi}]}} \frac{\phi - \phi_0}{M_{\rm pl}} \right\},$$
(18)

2

which leads to the following functional form for the approximate potential:

$$V(\phi) = AH_0^2 M_{\rm pl}^2 \left(1 - \sqrt{B + C\frac{\phi - \phi_0}{M_{\rm pl}}}\right)^D, \quad (19)$$

where the coefficients A, B, C and D are dimensionless constants depending on the cosmological parameters characterizing the model. For minimally coupled models with constant w_{ϕ} , the four coefficients assume the following values:

$$A = \frac{3\Omega_{de}(1 - w_{\phi})}{16\pi} \left(\frac{4 + 3\Omega_{m}w_{\phi}}{2 + 3\Omega_{m}w_{\phi}}\right)^{-3(1+w_{\phi})},$$

$$B = \frac{4}{(4 + 3\Omega_{m}w_{\phi})^{2}},$$

$$C = \mp 4 \frac{2 + 3\Omega_{m}w_{\phi}}{(4 + 3\Omega_{m}w_{\phi})^{2}} \sqrt{\frac{8\eta\pi}{3\Omega_{de}(1 + w_{\phi})}},$$

$$D = -3(1 + w_{\phi}).$$
(20)

(19) is an interesting result, showing that for constant equations of state, the potential can be represented by a very simple form.

The four parameters in (20) depend on two quantities, Ω_{de} and w_{ϕ} ; therefore, we can express two of them (*A* and *C*) in terms of *B* and *D*:

$$A = \frac{(D+6)[\sqrt{B}(D+1)-2]}{16\pi\sqrt{B}(D-3)(1-\sqrt{B})^{D}},$$

$$C = \mp 2\sqrt{\frac{8\eta\pi\sqrt{B}(3-D)}{D[\sqrt{B}(D+1)-2]}}\sqrt{B}(1-\sqrt{B}).$$
 (21)

To see how good our approximation is for $a \approx 1$, we compare the approximate expression for the potential to the exact solution for different values of the equation-of-state parameter w_{ϕ} in the top left panel of Fig. 1. We assumed the following cosmological parameters: $\Omega_{\rm m} = 0.3$, $\Omega_{\rm de} = 0.7$ and $H_0 = 70$ km s⁻¹ Mpc⁻¹. The value of ϕ at a = 1 ranges from $\phi/M_{\rm pl} \approx 0.02$ for $w_{\phi} = -0.99$ to $\phi/M_{\rm pl} \approx 0.2$ for $w_{\phi} = -0.7$. The approximate solution agrees very well with the analytic one over a range of values centred on a = 1 (by construction) and it deviates from it at both low



FIG. 1. Top left panel: Comparison between the exact solution for a constant equation of state for the potential (solid line) and its approximate expression (dashed line), for $a \approx 1$. Different colors refer to different values of w_{ϕ} . From top to bottom: the black, red, blue, yellow and violet lines correspond to $w_{\phi} = -0.7$, -0.8, -0.9, -0.95 and -0.99, respectively. Top right panel: Equation of state for the approximated potential of (19) for $w_{\phi} = -0.9$. The subscripts *a* and *e* represent the approximated (blue dashed line) and the exact (black solid line) solutions, respectively. Black horizontal dashed lines show differences of 1% with respect to the exact value. Bottom panel: Comparison between the exact solution for the scalar field with a constant equation of state $w_{\phi} = -0.9$ (solid line) and the approximate expression (dashed line), for $a \approx 1$.

and high values of the scale factor (corresponding to low and high values of the scalar field, respectively). In particular, by inspecting the top right panel of Fig. 1 we find an excellent agreement for $0.7 \leq a \leq 1.2$. We also note that a better agreement occurs when the equation-of-state parameter is not substantially different from $w_{\phi} = -1$: this is due to the fact that for the cosmological constant the scalar field and the potential are constant in time. If we require a tolerance of 1% in the equation of state derived from the approximate potential, then the confidence interval is $0.5 \leq a \leq 1.5$.

In the bottom panel of Fig. 1 we show the evolution of the scalar field with respect to the scale factor for $w_{\phi} = -0.9$. We show the time evolution of the scalar field rather than that of the potential because by construction, the latter evolves as $a^{-3(1+w_{\phi})}$. Note how the two expressions for the scalar field agree remarkably well over a range $0.5 \leq a \leq 1.7$. For values outside this range the approximate solution underestimates the exact one and it becomes negative for $a \leq 0.2$. This range is largely in agreement with what we found for the reconstructed equation-of-state parameter.

III. K-ESSENCE WITH CONSTANT w_{ϕ}

A straightforward extension of minimally coupled scalar fields is given by models with a noncanonical kinetic term. These models are described by a Lagrangian of the form $\mathcal{L} = \mathcal{L}(\phi, \chi)$ [12], where $\chi = -\frac{1}{2}g_{\mu\nu}\nabla^{\mu}\phi\nabla^{\nu}\phi$ is the canonical kinetic energy term. These models have been extensively used to describe dark energy scenario [12,13,50–53] and several works studied their dynamics and stability [54–58]. These models are dubbed "*k*-essence" models because the kinetic term χ can be responsible for the cosmic acceleration. A wide variety of models have been proposed and studied in different contests, such as low-energy effective string theory [59], tachyon models [52,60], ghost condensates [55,61–63], and Dirac-Born-Infeld theories [64–66].

The density and pressure are given by $\rho = 2\chi \mathcal{L}_{\chi} - \mathcal{L}$ and $P = \mathcal{L}$, respectively, where $\mathcal{L}_{\chi} = \frac{\partial \mathcal{L}}{\partial \chi}$ and we will also use $\mathcal{L}_{\chi\chi} = \frac{\partial^2 \mathcal{L}}{\partial \chi^2}$ and $\mathcal{L}_{\chi\phi} = \frac{\partial^2 \mathcal{L}}{\partial \chi \partial \phi}$. The sound speed for subhorizon modes is

APPROXIMATION OF THE POTENTIAL IN SCALAR ...

$$\alpha = \frac{P_{\chi}}{\rho_{\chi}} = \left(1 + \frac{2\chi \mathcal{L}_{\chi\chi}}{\mathcal{L}_{\chi}}\right)^{-1}.$$
 (22)

Using $P_{\phi} = w_{\phi}(a)\rho_{\phi}$, we can deduce that

$$2\chi \mathcal{L}_{\chi} = \frac{1 + w_{\phi}(a)}{w_{\phi}(a)} \mathcal{L}.$$
 (23)

From (23) we see that *k*-essence models can achieve $w_{\phi} \approx -1$ without $\chi \approx 0$. This means that such models need not be in the slow-roll regime to act as a dark energy component.

The energy-momentum tensor of *k*-essence is that of a perfect fluid

$$T_{\mu\nu} = \mathcal{L}_{\chi} \nabla_{\mu} \phi \nabla_{\nu} \phi + \mathcal{L} g_{\mu\nu} = (\rho + P) u_{\mu} u_{\nu} + P g_{\mu\nu} \qquad (24)$$

and velocity $u_{\mu} = \nabla_{\mu} \phi / \sqrt{2\chi}$. The equation of motion for the scalar field is

$$\dot{\chi}(\mathcal{L}_{\chi} + 2\chi\mathcal{L}_{\chi\chi}) + \sqrt{2\chi}(2\chi\mathcal{L}_{\chi\phi} - \mathcal{L}_{\phi}) + 6H\chi\mathcal{L}_{\chi} = 0,$$
(25)

and by rearranging the terms in (25), the equation of motion reads [18]

$$H^{\mu\nu}\nabla_{\mu}\nabla_{\nu}\phi + 2\chi\mathcal{L}_{\chi\phi} - \mathcal{L}_{\phi}$$

= $H^{\mu\nu}\nabla_{\mu}\nabla_{\nu}\phi - \mathcal{L}_{\chi\phi}g^{\mu\nu}\nabla_{\mu}\phi\nabla_{\nu}\phi - \mathcal{L}_{\phi} = 0,$ (26)

where

$$H^{\mu\nu} = \mathcal{L}_{\chi\chi} \nabla^{\mu} \phi \nabla^{\nu} \phi - \mathcal{L}_{\chi} g^{\mu\nu}.$$
 (27)

By inspecting (26), we notice that the equation of motion can be written in a very compact form as $\nabla_{\mu}J^{\mu} = -\mathcal{L}_{\phi}$, with $J^{\mu} = \mathcal{L}_{\chi}\nabla^{\mu}\phi$.

Many of the *k*-essence models proposed in the literature fall into one of the following types:

(A) Models of type A are given by [13,67–69]

$$\mathcal{L} = M^4 F(\chi) - V(\phi), \qquad (28)$$

where *M* has dimensions of mass and *F* is a dimensionless function. In the following, it is helpful to consider $F(\chi)$ to be a power law $F(\chi) = (\frac{\chi}{M^4})^n$, for *n* constant. Setting n = 1 implies $\mathcal{L} = \chi - V(\phi)$ which corresponds to the quintessence case discussed in Sec. II.

(B) Models of type B are given by [11,12,62,70]

$$\mathcal{L} = G(\chi)V(\phi), \tag{29}$$

where G is a dimensionless function.

Common Lagrangians proposed in the literature, which are mainly for purely kinetic *k*-essence model [i.e. $V(\phi) = M^4$ =const], are [53,71–76]

(1)
$$G(\chi) = -\sqrt{1 + 2\eta \frac{\chi}{M^4}},$$

(2) $G(\chi) = [2(\frac{\chi}{M^4})^n - 1]^{\frac{1}{2n}},$
(3) $G(\chi) = -[1 + 2\eta(\frac{\chi}{M^4})^n]^{\frac{1}{2n}},$
(4) $G(\chi) = A_1 \sqrt{\frac{\chi}{M^4}} - A_2(\frac{\chi}{M^4})^{\alpha},$
(5) $G(\chi) = -(1 - 2\frac{\chi}{M^4})^{\beta},$
(6) $G(\chi) = \frac{\chi}{M^4} - \sqrt{\frac{\chi}{M^4}},$
where n, α, β, A_1 and A_2 are constant

where n, α , β , A_1 and A_2 are constant and $\eta = \pm 1$. Typically it is not possible to transform between $\eta = +1$ and $\eta = -1$ via a simple redefinition of the scalar field.

An interesting Lagrangian to consider is the ghost condensate model [62]

$$\mathcal{L} = K(\phi)\chi + L(\phi)\frac{\chi^2}{M^4},$$
(30)

where $K(\phi) < 0$ and $L(\phi)$ are dimensionless potentials and *M*, again, has dimensions of mass. If one defines the scalar field ψ by

$$\left(\frac{d\psi}{d\phi}\right)^2 = \frac{L}{|K|} \tag{31}$$

and writes $X = -\frac{1}{2}g_{\mu\nu}\nabla^{\mu}\psi\nabla^{\nu}\psi$, then

$$\mathcal{L} = V(\phi) \left(-\frac{X}{M^4} + \frac{X^2}{M^8} \right), \tag{32}$$

where $V(\phi) = [K(\phi)]^2/L(\phi)$ if K < 0. Hence, this can be considered as a model of type B with $G(\chi) = -\chi/M^4 + \chi^2/M^8$.

(C) Models of type C are given by [63]

$$\mathcal{L} = -\chi - N(\chi)V(\phi), \qquad (33)$$

where $N = (\frac{\chi}{M^4})^n$ is a dimensionless function. The model represents a generalization of the dilatonic ghost condensate model and it is a special case of (30), where $K(\phi) = -1$, $L(\phi) = \frac{V(\phi)}{M^4}$ and $N(\chi) = \chi^2$.

A. Type A models with constant w_{ϕ}

For models of type A, with $F(\chi)$ being a power law, we have $\frac{\chi F_{\chi}}{F} = n$ and $\frac{\chi F_{\chi\chi}}{F_{\chi}} = n - 1$ and therefore $\alpha = (2n - 1)^{-1}$, constant. For a general $w_{\phi}(a)$ we have

$$V(\phi) = \frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}[1 - (2n - 1)w_{\phi}(a)]g(a)}{16\pi n},\qquad(34)$$

$$\frac{d\phi}{da} = \sqrt{2} \frac{M^2}{H_0} \left(\frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}}{16\pi n M^4} \right)^{\frac{1}{2n}} \frac{\{[1+w_\phi(a)]g(a)\}^{\frac{1}{2n}}}{aE(a)},\tag{35}$$

$$\frac{\phi - \phi_0}{M_{\rm pl}} = \sqrt{2} \frac{M^2}{H_0 M_{\rm pl}} \left(\frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}}{16\pi n M^4}\right)^{\frac{1}{2n}} \\ \times \int_1^a \frac{\{[1 + w_\phi(x)]g(x)\}^{\frac{1}{2n}}}{xE(x)} dx. \quad (36)$$

It is, therefore, possible in principle to find $\phi(a)$, at least numerically. Note that $n \neq 0$; otherwise the potential and the scalar field diverge. Our general results are consistent with [77] if we set $M^4 = 1$ and $F(\chi) = \chi^2$ and with [78].

If w_{ϕ} is constant, we can recover analogous results to the quintessence case. In this case, (36) becomes

$$\frac{\phi - \phi_0}{M_{\rm pl}} = -\frac{2\sqrt{2}M^2}{3w_{\phi}H_0M_{\rm pl}\Omega_{\rm m}^{1/2}} \left(\frac{3H_0^2M_{\rm pl}\Omega_{\rm de}(1+w_{\phi})}{16\pi nM^4}\right)^{\frac{1}{2n}} \times \left(\frac{\Omega_{\rm m}}{\Omega_{\rm de}}\right)^{\frac{1+w_{\phi}-n}{2nw_{\phi}}} \int_{\sinh^{-1}\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}}^{\sinh^{-1}\left(\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}a^{\frac{3w_{\phi}}{2}}\right)} dx \sinh^{\frac{(1+w_{\phi})(1-n)}{nw_{\phi}}}x.$$
(37)

It is not possible to compute this integral analytically for general w_{ϕ} and *n*, but it at least illustrates that a solution exists and the solution can be computed numerically. It is also important to notice that for a given equation-of-state parameter, the scalar field and its potential are not uniquely determined since for a given w_{ϕ} , these two quantities depend also on *n*. Note also that Eqs. (34), (36) and (37) reduce to the quintessence case for n = 1. At early and late times, the potential is given by $V_{\rm E}(\phi) \propto \phi^{\frac{2n(1+w_{\phi})}{|n-(1+w_{\phi})|}}$ and $V_{\rm L}(\phi) \propto \phi^{-\frac{2n}{(n-1)}}$, respectively, which are in agreement with [78].

Since analytical solutions are not possible, we find it useful to derive approximated expressions also for $a \approx 1$. In this case, (35) is approximated by

$$\frac{d\phi}{da} = \sqrt{2} \frac{M^2}{H_0} \left[\frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}(1+w_\phi)}{16\pi n M^4} \right]^{\frac{1}{2n}} \\ \times \left[1 + \frac{n-3+3(\Omega_{\rm de}n-1)w_\phi}{2n} (a-1) \right], \quad (38)$$

which leads to

$$\frac{\phi - \phi_0}{M_{\rm pl}} = \sqrt{2} \frac{M^2}{H_0 M_{\rm pl}} \left[\frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}(1 + w_\phi)}{16\pi n M^4} \right]^{\frac{1}{2n}} \\ \times \left[a - 1 + \frac{n - 3 + 3(\Omega_{\rm de}n - 1)w_\phi}{4n} (a - 1)^2 \right].$$
(39)

By inverting this expression to find $a(\phi)$, the potential can be written with the same functional form as (19), with the following coefficients:

$$A = \frac{3\Omega_{de}[1 - (2n - 1)w_{\phi}]}{16\pi n} \\ \times \left(\frac{3 + n - 3(\Omega_{de}n - 1)w_{\phi}}{3 - n - 3(\Omega_{de}n - 1)w_{\phi}}\right)^{-3(1+w_{\phi})}, \\ B = \frac{4n^2}{[3 + n - 3(\Omega_{de}n - 1)w_{\phi}]^2}, \\ C = -2\sqrt{2}n \frac{H_0 M_{pl}}{M^2} \frac{3 - n - 3(\Omega_{de}n - 1)w_{\phi}}{[3 + n - 3(\Omega_{de}n - 1)w_{\phi}]^2} \\ \times \left[\frac{16\pi n M^4}{3H_0^2 M_{pl}^2 \Omega_{de}(1 + w_{\phi})}\right]^{\frac{1}{2n}}, \\ D = -3(1 + w_{\phi}).$$
(40)

When n = 1, this reverts to the coefficients presented in (20).

In the top left panel of Fig. 2 we make a comparison between exact numerically generated solutions and the approximation around $a \approx 1$. As in the case of quintessence there is a good agreement between the two. We also show in the right panel that the potential for n > 2 quickly asymptotes to the $n \rightarrow \infty$ solution. This should be expected from the form of (37). In the lower panels we show the range of validity of the approximated scalar field (left) and potential (right) for the approximated expression found and described by the four coefficients listed above. As for quintessence models, the approximate potential recovers the exact one only for a limited range in the scale factor, hence also the reconstructed equation of state will be limited to the range of validity of the approximate potential.

We found that the behavior of the reconstructed equation of state is very similar to the quintessence case. More quantitatively, we match the true w_{ϕ} with $\Delta w_{\phi} = 0.01$ for $0.6 \leq a \leq 1.5$, a range that is largely in agreement with quintessence.

B. Type B models with constant w_{ϕ}

Type B models behave quite differently from type A models and it is not possible to make a direct comparison with quintessence or phantom models. These models are



FIG. 2. Top left panel: Comparison between the full solution, generated numerically, and the approximation for $a \approx 1$ for the scalar field potential for type A model with n = 2. Solid lines represent the full solution, dashed lines the approximate solution. Different colors show different equations of state, labeled as in Fig. 1. Top right panel: Scalar field potential for $w_{\phi} = -0.9$ for different values of n. From top to bottom we show n = 1, 2, 3, 4. We see that as n increases, the shape of the potential quickly asymptotes to that of $n \to \infty$. Bottom left (right) panel: Scalar field (potential) for the approximated solution compared with the exact expression [Eq. (34) together with Eq. (36)] for $w_{\phi} = -0.9$.

commonly studied in the literature because the kinetic term is completely factorized from the potential term, making the calculations relatively easy.

The evolution of the potential, the kinetic term and the sound speed are given, respectively, by

$$V(a) = \frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de} w_{\phi}(a) g(a)}{8\pi G(\chi)},$$
 (41)

$$\frac{\chi G_{\chi}}{G} = \frac{1 + w_{\phi}(a)}{2w_{\phi}(a)},\tag{42}$$

$$\alpha = \left(1 + 2\frac{\chi G_{\chi\chi}}{G_{\chi}}\right)^{-1}.$$
 (43)

One approach would be to solve for $G(\chi)$ from (42). When w_{ϕ} is constant it is given by

$$G(\chi) = \left(\frac{\chi}{M^4}\right)^{\frac{1+w_{\phi}}{2w_{\phi}}}.$$
(44)

However, if we do this, then we find that $\alpha = w_{\phi}$ which would mean that perturbations would be

unstable if $w_{\phi} < 0$. This is, therefore, not the correct approach for deducing a potential from constant w_{ϕ} .

The alternative is to specify $G(\chi)$ and consider (42) as a constraint on χ which will be constant. Let $\hat{\chi}$ be the constant value which solves (42) for a specific choice of $G(\chi)$, and then

$$\frac{d\phi}{da} = \frac{1}{H_0} \left(\frac{2\hat{\chi}a}{\Omega_{\rm m} + \Omega_{\rm de}a^{-3w_{\phi}}} \right)^{1/2},\tag{45}$$

whose solution is

$$\frac{\phi - \phi_0}{M_{\rm pl}} = -\frac{2}{3w\Omega_{\rm m}^{1/2}} \frac{\sqrt{2\hat{\chi}}}{H_0 M_{\rm pl}} \left(\frac{\Omega_{\rm de}}{\Omega_{\rm m}}\right)^{\frac{1}{2w_{\phi}}} \\ \times \int_{\sinh^{-1}\left(\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}a^{-\frac{3w_{\phi}}{2}}\right)} dx \sinh^{-\frac{1+w_{\phi}}{w_{\phi}}} x.$$
(46)

Again this at least proves the existence of an $\phi(a)$, and hence a $V(\phi)$, which gives rise to constant w_{ϕ} , although there is no analytic solution for general

 w_{ϕ} . Note that this expression is equivalent to (37) in the limit $n \to \infty$ and with $\sqrt{\hat{\chi}} = M^2$. We are able to find useful approximations at early and late times. In particular we find $V_{\rm E}(\phi) \propto \phi^{-2(1+w_{\phi})}$ at early times and $V_{\rm L}(\phi) \propto \phi^{-2}$ at late times, respectively.

Let us now consider the specific case of

$$G(\chi) = -\frac{\chi}{M^4} + \frac{\chi^2}{M^8}, \qquad (47)$$

from which we can deduce that $\frac{\hat{\chi}}{M^4} = \frac{1-w_{\phi}}{1-3w_{\phi}}$. Hence, we find that $\alpha = \frac{1-2\frac{\hat{\chi}}{M^4}}{1-6\frac{\hat{\chi}}{M^4}} = \frac{1+w_{\phi}}{5-3w_{\phi}}$. Note that $0 \le \alpha \le 1$ implies that $-1 \le w_{\phi} \le 1$. From (30), one finds that

$$\frac{\chi L(\phi)}{M^4 K(\phi)} = \frac{w_{\phi} - 1}{1 - 3w_{\phi}},$$
(48)

and the corresponding sound speed is

$$\alpha = \frac{1 + 2\frac{L}{K}\frac{\chi}{M^4}}{1 + 6\frac{L}{K}\frac{\chi}{M^4}} = \frac{1 + w_{\phi}}{5 - 3w_{\phi}},\tag{49}$$

which is what should be expected from the general discussion about the generalized ghost condensate model. Note that in type B models with constant w_{ϕ} , the sound speed becomes a function the of equation of state $\alpha = \alpha(w_{\phi})$; such models have been also studied in [79] where the authors used shear and cosmic microwave background lensing data to constrain dark energy perturbations.

For $a \approx 1$, the differential equation governing the evolution of the scalar field ϕ as a function of the scale factor a is

$$\frac{d\phi}{da} = \frac{\sqrt{2\hat{\chi}}}{H_0} \left[1 + \frac{1}{2} (1 + 3\Omega_{\rm de} w_{\phi})(a-1) \right], \quad (50)$$

which leads to

$$\frac{\phi - \phi_0}{M_{\rm pl}} = \frac{\sqrt{2\hat{\chi}}}{H_0 M_{\rm pl}} \left[a - 1 + \frac{1}{4} (1 + 3\Omega_{\rm de} w_\phi) (a - 1)^2 \right].$$
(51)

The relation between the scale factor and the scalar field is then



FIG. 3. Left panel: Approximation of the scalar field potential for $a \approx 1$ for different constant equation-of-state parameters w_{ϕ} . Solid lines represent the exact numerical solution, while the dashed line show the approximated solution. Colors are as in Fig. 1. Middle (right) panel: Scalar field (potential) for the approximated solution compared with the exact expression (41) together with (46) for $w_{\phi} = -0.9$. In all the panels we assume $G(\chi) = -\frac{\chi}{M^4} + \frac{\chi^2}{M^8}$.

APPROXIMATION OF THE POTENTIAL IN SCALAR ...

2

$$a = 1 - \frac{2}{1 + 3\Omega_{de}w_{\phi}} \times \left\{ 1 - \sqrt{1 + (1 + 3\Omega_{de}w_{\phi})\frac{H_0}{\sqrt{2\hat{\chi}}}(\phi - \phi_0)} \right\}.$$
(52)

Hence, also for type B models we will have an approximate potential of the form of (19) with dimensionless coefficients

$$A = \frac{3\Omega_{de}w_{\phi}}{8\pi G(\hat{\chi})} \left(\frac{3\Omega_{de}w_{\phi}-1}{3\Omega_{de}w_{\phi}+1}\right)^{-3(1+w_{\phi})},$$

$$B = \frac{4}{(1-3\Omega_{de}w_{\phi})^2},$$

$$C = 4\frac{1+3\Omega_{de}w_{\phi}}{(1-3\Omega_{de}w_{\phi})^2} \frac{H_0M_{pl}}{\sqrt{2\hat{\chi}}},$$

$$D = -3(1+w_{\phi}).$$
(53)

In the left panel of Fig. 3 we show the validity of the approximation for the scalar field potential for $a \approx 1$ for different constant equation-of-state parameters, as described in the caption, for a model with $G(\chi) = -\frac{\chi}{M^4} + \frac{\chi^2}{M^8}$. We obtain a similar level of agreement as for quintessence and type A models. The accuracy increases with the decrease of w_{ϕ} and is limited to an epoch centered on a = 1. In the right and middle panels of Fig. 3 we compare the approximated expression for the potential and the corresponding scalar field evolution with the exact

solution. Note that since (46) is a limiting case of (37), the range of agreement of the equation of state for type B models is similar to that found for type A models.

In the general discussion of type B models, we showed that $\alpha = \alpha(w_{\phi})$ and $\hat{\chi} = \hat{\chi}(w_{\phi})$; each model will have, therefore, its own particular functional form and a range of values for the parameters ensuring their stability. In Table I we show the specific functional form of $\hat{\chi}$ and α for several forms of $G(\chi)$ proposed in the literature and determine when their perturbations are stable ($\alpha \ge 0$) and subluminal ($\alpha \le 1$).

C. Type C models with constant w_{ϕ}

Type C models resemble phantom models discussed in Sec. II. They reduce to phantom models when the function $N(\chi)$ is constant, that is, n = 0 for the power-law choice of $N(\chi)$. Despite the apparent complexity, type C models, in contrast to type A and B models, have a general analytical solution when $N(\chi)$ is a power law. Key equations for general n > 0 and $w_{\phi}(a)$ are

$$V(a) = -\frac{[1 - w(a)]M^{4n}}{[1 - (2n - 1)w_{\phi}(a)]^n} (n - 1)^{n - 1} \\ \times \left(\frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de} g(a)}{16\pi}\right)^{1 - n},$$
(54)

$$\frac{d\phi}{da} = \left\{ \frac{3M_{\rm pl}^2 \Omega_{\rm de} [1 - (2n-1)w_{\phi}(a)]g(a)}{8\pi (n-1)a_{+-w_{\ell}}^2 E^2(a)} \right\}^{\frac{1}{2}}, \quad (55)$$

$$\alpha = \frac{1 + w_{\phi}}{2n + 1 - (2n - 1)w_{\phi}},$$
(56)

TABLE I. Dependence of $\hat{\chi}$ and α on the constant equation-of-state parameter w_{ϕ} and stability conditions for the model.

$G(\chi)$	$rac{\hat{\chi}(w_{\phi})}{M^4}$	$lpha(w_{oldsymbol{\phi}})$	Stability
$\left(\frac{\chi}{M^4}\right)^{\frac{1+w_\phi}{2w_\phi}}$		Wø	$w_{\phi} \ge 0$
$-\frac{\chi}{M^4}+\frac{\chi^2}{M^8}$	$\frac{1-w_{\phi}}{1-3w_{\phi}}$	$\frac{1+w_{\phi}}{5-3w_{\phi}}$	$-1 \le w_{\phi} \le 1$
$-rac{\chi}{M^4}+(rac{\chi}{M^4})^n$	$(\frac{1 - w_{\phi}}{1 - (2n - 1)w_{\phi}})^{\frac{1}{n - 1}}$	$\frac{1+w_{\phi}^{\varphi}}{(2n+1)-(2n-1)w_{\phi}}$	$-1 \le w_{\phi} \le 1$ for $n > 0$
$-\sqrt{1+2\etarac{\chi}{M^4}}$	$-\eta rac{1+w_\phi}{2}$	$-w_{\phi}$	$-1 \le w_{\phi} \le 0$
$[2(\frac{\chi}{M^4})^n - 1]^{\frac{1}{2n}}$	$\left(\frac{1+w_{\phi}}{2} ight)^{\frac{1}{n}}$	$-\frac{1}{2n-1}w_{\phi}$	$1 - 2n \le w_{\phi} \le 0$ for $n > \frac{1}{2}$
			$0 \le w_{\phi} \le 1 - 2n$ for $n < \frac{1}{2}$
$-[1+2\eta(\frac{\chi}{M^4})^n]^{\frac{1}{2n}}$	$(-\eta \frac{1+w_{\phi}}{2})^{\frac{1}{n}}$	$-\frac{1}{2n-1}W_{\phi}$	$1 - 2n \le w_{\phi} \le 0$ for $n > \frac{1}{2}$
			$0 \le w_{\phi} \le 1 - 2n$ for $n < \frac{1}{2}$
$rac{\chi}{M^4} - \sqrt{rac{\chi}{M^4}}$	$\frac{1}{(1-w_{\phi})^2}$	$\frac{1+w_{\phi}}{2}$	$-1 \le w_{\phi} \le 1$
$-(1-2\frac{\chi}{M^4})^{\beta}$	$\frac{1+w_{\phi}}{2[1+(1-2\beta)w_{\phi}]}$	$\frac{\beta w_{\phi}}{\beta - 1 + (2\beta - 1)w_{\phi}}$	$w_{\phi} \leq -1$ or $w_{\phi} \geq 0$ for $\beta < 0$ or $\beta > 1$
			$-1 \le w_{\phi} \le 0$ for $0 < \beta < 1$
$A_1 \sqrt{\frac{\chi}{M^2}} - A_2 (\frac{\chi}{M^2})^{\alpha}$	$\left\{\frac{A_2}{A_1}\left[1-(2\alpha-1)w_{\phi}\right]\right\}^{\frac{2}{1-2\alpha}}$	$\frac{1+w_{\phi}}{2\alpha}$	$-1 \le w_{\phi} \le 2\alpha - 1$ for $\alpha > 0$
• 			$2\alpha - 1 \le w_{\phi} \le -1$ for $\alpha < 0$

for the evolution of the potential, of the scalar field and of the sound speed, respectively. Note in particular that for n = 0 we recover results for the phantom models. The evolution of the scalar field as a function of the scale factor is

$$\frac{\phi - \phi_0}{M_{\rm pl}} = -\frac{2}{3w_\phi} \sqrt{\frac{3[1 + (1 - 2n)w_\phi]}{8\pi(n - 1)}} \times \left[\sinh^{-1}\left(\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}a^{-\frac{3w_\phi}{2}}\right) - \sinh^{-1}\left(\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}\right)\right],\tag{57}$$

and the corresponding potential is

$$V(\phi) = k \sinh^{\frac{2(n-1)(1+w_{\phi})}{w_{\phi}}} \times \left[-\frac{3w_{\phi}}{2} \sqrt{\frac{8\pi(n-1)}{3[1+(1-2n)w_{\phi}]}} \frac{\phi}{M_{\rm pl}} \right], \quad (58)$$

with the constant k:

$$k = \frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}(1 - w_{\phi})}{16\pi(1 - n)} \times \left[\frac{3H_0^2 M_{\rm pl}^2 \Omega_{\rm de}[1 - (2n - 1)w_{\phi}]}{16\pi(n - 1)M^4}\right]^{-n} \left(\frac{\Omega_{\rm m}}{\Omega_{\rm de}}\right)^{-\frac{(n - 1)(1 + w_{\phi})}{w_{\phi}}},$$
(59)

when we choose as we did in the quintessence case

$$\frac{\phi_0}{M_{\rm pl}} = -\frac{2}{3w_{\phi}} \sqrt{\frac{3[1+(1-2n)w_{\phi}]}{8\pi(n-1)}} \sinh^{-1}\left(\sqrt{\frac{\Omega_{\rm de}}{\Omega_{\rm m}}}\right).$$
(60)

These expressions are as expected similar to those for quintessence models.

Given the general form of the potential for type C models, we can expect a similar behavior to quintessence models for $a \ll 1$, $a \approx 1$ and $a \gg 1$. In particular, at early times $V_{\rm E}(\phi) \propto \phi^{\frac{2(1-n)(1+w_{\phi})}{w_{\phi}}}$ and at late times we recover the



FIG. 4. Top left panel: Comparison between the exact solution for constant equation of state for the absolute value of the potential (solid line) and its approximate expression, for $a \approx 1$ (dashed line) for type C models. Line styles and colors are as in Fig. 1. Top right (middle) panel: Scalar field (absolute value of the potential) for the approximated solution compared with the exact expression in (57) [(58)] for a model with $w_{\phi} = -0.9$. The subscripts *a* and *e* represent the approximated (blue dashed line) and the exact (black solid line) solutions, respectively.

usual exponential behavior $V_{\rm L}(\phi) \propto \exp(\phi/M_{\rm pl})$ typical of the minimally coupled models.

For $a \approx 1$, the scalar field is

$$\frac{\phi - \phi_0}{M_{\rm pl}} = \sqrt{\frac{3\Omega_{\rm de}[1 - (2n-1)w_\phi]}{8\pi(n-1)}} \times \left[a - 1 - \frac{1}{4}(2 + 3\Omega_{\rm m}w_\phi)(a-1)^2\right], \quad (61)$$

and the corresponding potential can be once again written in the general approximated form of (19), with coefficients:

$$A = \frac{3\Omega_{de}(1 - w_{\phi})}{16\pi(1 - n)} \left[\frac{3H_0^2 M_{pl}^2 \Omega_{de}[1 + (1 - 2n)w_{\phi}]}{16\pi(n - 1)M^4} \right]^{-n} \\ \times \left(\frac{4 + 3\Omega_m w_{\phi}}{2 + 3\Omega_m w_{\phi}} \right)^{-3(1 - n)(1 + w_{\phi})}, \\ B = \frac{4}{(4 + 3\Omega_m w_{\phi})^2}, \\ C = -4 \frac{2 + 3\Omega_m w_{\phi}}{(4 + 3\Omega_m w_{\phi})^2} \sqrt{\frac{8\pi(n - 1)}{3\Omega_{de}[1 + (1 - 2n)w_{\phi}]}}, \\ D = -3(1 - n)(1 + w_{\phi}).$$
(62)

As with quintessence, type A and B models, we show a comparison between the approximation and the exact solutions in Fig. 4. The picture is similar to the previous ones but the range of scale factor where the approximation is good is more restricted, $0.8 \leq a \leq 1.2$. This is due to the fact that for type C models, the potential has a stronger dependence on the scale factor with respect to the other models, given by the 1 - n power of g(a) in (54).

IV. POTENTIAL FOR NONCONSTANT w_{ϕ}

In the previous sections we have shown that a potential of the form (19) is a good approximation to that for scalar field models with constant w_{ϕ} —for both minimal and nonminimal kinetic terms—for some choice of the parameters *A*, *B*, *C* and *D* over a range of the scale factors around $a \approx 1$.

In Fig. 5 we have varied the parameters around their values for a specific constant $w_{\phi} = -0.9$ model with a minimal kinetic term. We do this by keeping three of the parameters fixed and vary the fourth one by $\pm 30\%$ while requiring that ϕ_0 coincides with the exact solution. We see that a wide range of behavior of the actual w(a) can be achieved in these models suggesting that this parametrization of the potential could be used as a proxy for a significant range of models, albeit with some restrictions.

We can attempt to generalize the set of coefficients of (19) to models with a nonconstant equation of state. It is not possible to adapt the exact method used for constant w_{ϕ} because usually there is not a general expression for g(a). However, we have been able to make some progress by realizing that $(1 + \alpha x)^{\beta} \approx 1 + \alpha \beta x$ for $x \ll 1$ and performing an expansion around $\phi \approx \phi_0$ in our set up. Expanding (19) to first order and matching the coefficients with a similar expansion derived from V(a) and $\phi(a)$, we can determine a new set of parameters A-D. As before, they will depend on the background cosmological parameters, $\Omega_{\rm m}$, w_{ϕ} and in this case, also on its derivative with respect to the scale factor, w'_{ϕ} , evaluated at a = 1. In the Appendix we report the explicit expression for the four coefficients for minimally coupled, type A and type C models, respectively. To understand them note that the relation between the scale factor and the scalar field is now, for quintessence models,



FIG. 5. Effect of the variations of the parameters A, B, C and D on the equation of state w_{ϕ} . In each panel, three of the coefficients are held constant at their exact value for $w_{\phi} = -0.9$ while the fourth one is varied by $\pm 30\%$ —left negative and right positive. The black curve shows the equation of state using the exact values of the coefficients as represented in Fig. 1. Effects of variations of A, B, C and D are shown with the red dashed, blue short dashed, brown dotted and green dot-dashed curve, respectively.



FIG. 6. Left (right) panel: Equation of state for the approximated potential of (19) for a quintessence (type A) model described by a CPL equation of state. The red (blue) curve represents a model with $w_{\phi}(a) = -0.9 + 0.02(1 - a) [w_{\phi}(a) = -0.9 + 0.15(1 - a)]$. Dotted lines represent a 1% difference from the true equation of state. Solid (dashed) lines show the true (approximated) equation of state.

$$a = 1 + \frac{2}{2 + 3\Omega_{\rm m}w_{\phi}(1) - \frac{w_{\phi}'(1)}{1 + w_{\phi}(1)}} \left\{ 1 - \sqrt{1 - \left[2 + 3\Omega_{\rm m}w_{\phi}(1) - \frac{w_{\phi}'(1)}{1 + w_{\phi}(1)}\right]} \sqrt{\frac{8\eta\pi}{3\Omega_{\rm de}[1 + w_{\phi}(1)]}} \frac{\phi - \phi_0}{M_{\rm pl}} \right\}.$$
 (63)

Also note that for $w'_{\phi}(1) = 0$, we recover the result in (18) for quintessence models. Similar expressions hold for type A and type C models.

To see how well this new parametrization performs we use the Chevallier-Polarski-Linder (CPL) parametrization [14,15]:

$$w_{\phi}(a) = w_0 + w_a(1-a), \tag{64}$$

where w_0 and w_a are constants. In Fig. 6 we show the comparison between the true and the approximated equation of state evaluated from the potential in (19) with the set of coefficients given in the Appendix for minimally coupled models and type A k-essence models. We use two different sets of coefficients (w_0, w_a) : one with a very gentle slope, $w_{\phi}(a) =$ -0.9 + 0.02(1 - a), and one with a more pronounced variation, $w_{\phi}(a) = -0.9 + 0.15(1 - a)$. At early times we find a better agreement for models not differing too much from a constant equation of state, while at late times, the agreement is better for models with $w_a = 0.15$. As it can be seen in Fig. 6, this is due to the fact that ϕ' shows fluctuations around the true value. Note that if we limit ourselves to a subpercent agreement between the true and the reconstructed equation of state, then the agreement is much more limited with respect to the case of constant w_{ϕ} . This is because we poorly approximate the function q(a); for a CPL model, it consists of two elements: a power law and an exponential and we only include the power law. When the exponential behavior dominates, our proposed potential is a less good fit to the true behavior. Note also that the range of agreement is similar for both quintessence and type A models. Deviations in type A models are suppressed with respect to quintessence models thanks to a higher value of n (2 in the example). From a quantitative point of view, for $w_a =$ $0.02 (w_a = 0.15)$, for quintessence models we reach a 1% agreement for $0.5 \leq a \leq 1.4$ ($0.5 \leq a \leq 1.7$), while for type A models we have $0.5 \leq a \leq 1.4$ and $0.7 \lesssim a \lesssim 1.7$, respectively. This is similar to what found before for a model with constant $w_{\phi} = -0.9$. One caveat to our approach is that the only knowledge of the evolution of the equation of state is given by its value and its time derivative, both evaluated at a = 1. Therefore nothing is known about its general time evolution and as a consequence, nothing is known about the functional form of q(a). This implies that our approach would work well for models with a monotonic equation of state [and hence a monotonic q(a)], but we expect it to fail and not be a good representation for the true potential for oscillating dark energy models (see e.g. [80] for a recent study of their properties and comparison with observations).

V. CONCLUSIONS

Scalar fields are an important field of research in cosmology and are one of the most studied candidates used to explain and describe the accelerated expansion of the Universe. In this work, we consider two main classes of models: minimally coupled models (both quintessence and phantom) and *k*-essence models. For this second class, we

specialize the Lagrangian to assume three particular functional forms, dubbed type A, type B and type C models. In each case, we have shown that specifying the scalar field potential $V(\phi)$, one can determine the evolution of the scalar field and the corresponding equation of state $w_{\phi}(a)$.

This is true generally but in order to make it clear, we have assumed the equation of state to be known and we calculated explicitly the time evolution of the scalar field and of the potential in some cases. We showed that it is possible to obtain an exact analytic solution for minimally coupled and the type C models with a constant equation of state. This is not possible for more general *k*-essence models or for models with a time-varying $w_{\phi}(a)$, but we have solutions for $\phi(a)$ as definite integrals and these can be used to establish the potentials, $V(\phi)$ numerically.

We have also derived useful approximate forms of the potential which are valid in different epochs, corresponding to the domination of one cosmic fluid. In particular we deduce the form of the potential at early times ($a \ll 1$, corresponding to the matter-dominated epoch) and at late times ($a \gg 1$, corresponding to the scalar-field-dominated regime), showing that in general the potential is often very well approximated by a power law.

From an observational point of view, the most important regime to understand the potential is around $a \approx 1$. Assuming initially a constant equation of state w_{ϕ} , we showed that the scalar field potential can be approximated by the expression given in (19). This expression depends only on four parameters and with the appropriate choice of coefficients can cover all the classes of models studied in this work. In Sec. IV we discussed how this expression might be applied to dynamical dark energy models, by appropriately choosing a new set of parameters which reduces to the correct expression in the limit of constant w_{ϕ} . Note that this cannot be done for type B models, since our formalism only works for constant equations of state.

In some respect our approach is similar to the work of [35]. To derive our expression in (19), we performed a Taylor expansion of the scalar field evolution, so the same critique could be applied: where to stop the series? Our

approximate potential, for $a \approx 1$ ($\phi \approx \phi_0$) can be expanded in powers of $\phi - \phi_0$ leading to the same form of the potential proposed by [35]. In contrast to that work, our proposed potential has well-motivated coefficients and in the regime of interest it would be possible to map the V_i coefficients of (1) in terms of our four parameters. For example, at zeroth order, we can write V_0 in (1) as $V_0 = A(1 - \sqrt{B})H_0^2M_{pl}^2$.

ACKNOWLEDGMENTS

We would like to thank Stefano Camera and Robert Reischke for useful comments. The work for this article was funded by an STFC postdoctoral fellowship.

APPENDIX: COEFFICIENTS FOR DYNAMICAL DARK ENERGY MODELS

In this section we write explicitly the generalization of the set of coefficients A-D for nonconstant equations of state using the approach discussed in the text. For minimally coupled models we find

$$\begin{split} A &= \frac{3\Omega_{\rm de}[1-w_{\phi}(1)]}{16\pi} \\ &\times \left\{ \frac{4+3\Omega_{\rm m}w_{\phi}(1)-\frac{w_{\phi}'(1)}{1+w_{\phi}(1)}}{2+3\Omega_{\rm m}w_{\phi}(1)-\frac{w_{\phi}'(1)}{1+w_{\phi}(1)}} \right\}^{-\left\{ \frac{w_{\phi}'(1)}{1-w_{\phi}(1)}+3[1+w_{\phi}(1)] \right\}}, \\ B &= \frac{4}{\left[4+3\Omega_{\rm m}w_{\phi}(1)-\frac{w_{\phi}'(1)}{1+w_{\phi}(1)} \right]^2}, \\ C &= -4\frac{2+3\Omega_{\rm m}w_{\phi}(1)-\frac{w_{\phi}'(1)}{1+w_{\phi}(1)}}{\left[4+3\Omega_{\rm m}w_{\phi}(1)-\frac{w_{\phi}'(1)}{1+w_{\phi}(1)} \right]^2} \sqrt{\frac{8\eta\pi}{3\Omega_{\rm de}[1+w_{\phi}(1)]}}, \\ D &= -\left\{ \frac{w_{\phi}'(1)}{1-w_{\phi}(1)}+3[1+w_{\phi}(1)] \right\}. \end{split}$$
(A1)

For models of type A we find

$$A = \frac{3\Omega_{de}[1 - (2n - 1)w_{\phi}(1)]}{16\pi n} \left\{ \frac{3 + n - 3(\Omega_{de}n - 1)w_{\phi}(1) - \frac{w'_{\phi}(1)}{1 + w_{\phi}(1)}}{3 - n - 3(\Omega_{de}n - 1)w_{\phi}(1) - \frac{w'_{\phi}(1)}{1 + w_{\phi}(1)}} \right\}^{-\left\{\frac{(2n - 1)w'_{\phi}(1)}{1 - (2n - 1)w_{\phi}(1)} + 3[1 + w_{\phi}(1)]\right\}},$$

$$B = \frac{4n^{2}}{[3 + n - 3(\Omega_{de}n - 1)w_{\phi}(1) - \frac{w'_{\phi}(1)}{1 + w_{\phi}(1)}]^{2}},$$

$$C = -2\sqrt{2}n \frac{3 - n - 3(\Omega_{de}n - 1)w_{\phi}(1) - \frac{w'_{\phi}(1)}{1 + w_{\phi}(1)}}{[3 + n - 3(\Omega_{de}n - 1)w_{\phi}(1) - \frac{w'_{\phi}(1)}{1 + w_{\phi}(1)}]^{2}} \frac{H_{0}M_{pl}}{M^{2}} \left[\frac{16\pi nM^{4}}{3H_{0}^{2}M_{pl}^{2}\Omega_{de}(1 + w_{\phi})}\right]^{\frac{1}{2n}},$$

$$D = -\left\{\frac{(2n - 1)w'_{\phi}(1)}{1 - (2n - 1)w_{\phi}(1)} + 3[1 + w_{\phi}(1)]\right\}.$$
(A2)

For models of type C we find

$$A = \frac{3\Omega_{de}(1-w_{\phi})}{16\pi(1-n)} \left[\frac{3H_0^2 M_{pl}^2 \Omega_{de}[1+(1-2n)w_{\phi}]}{16\pi(n-1)M^4} \right]^{-n} \left[\frac{4+3\Omega_{m}w_{\phi} - \frac{(1-2n)w_{\phi}(1)}{1+(1-2n)w_{\phi}(1)}}{2+3\Omega_{m}w_{\phi} - \frac{(1-2n)w_{\phi}(1)}{1+(1-2n)w_{\phi}(1)}} \right]^{-(1-n)\{\frac{[1+2n+(1-2n)w_{\phi}(1)]}{[1+(1-2n)w_{\phi}(1)][1-w_{\phi}(1)]} + 3[1+w_{\phi}(1)]\}},$$

$$B = \frac{4}{\left[4+3\Omega_{m}w_{\phi} - \frac{(1-2n)w_{\phi}(1)}{1+(1-2n)w_{\phi}(1)} \right]^2},$$

$$C = -4\frac{2+3\Omega_{m}w_{\phi} - \frac{(1-2n)w_{\phi}(1)}{1+(1-2n)w_{\phi}(1)}}{\left[4+3\Omega_{m}w_{\phi} - \frac{(1-2n)w_{\phi}(1)}{1+(1-2n)w_{\phi}(1)} \right]^2},$$

$$D = -(1-n)\left\{ \frac{[1+2n+(1-2n)w_{\phi}(1)]w_{\phi}(1)}{[1+(1-2n)w_{\phi}(1)][1-w_{\phi}(1)]} + 3[1+w_{\phi}(1)] \right\}.$$
(A3)

- [1] Planck Collaboration XIII, arXiv:1502.01589 [Astron. Astrophys. (to be published)].
- [2] Planck Collaboration XIV, arXiv:1502.01590.
- [3] L. H. Ford, Phys. Rev. D 35, 2339 (1987).
- [4] P. J. E. Peebles and B. Ratra, Astrophys. J. Lett. 325, L17 (1988).
- [5] B. Ratra and P. J. E. Peebles, Phys. Rev. D 37, 3406 (1988).
- [6] C. Wetterich, Nucl. Phys. B302, 668 (1988).
- [7] R. R. Caldwell, R. Dave, and P. J. Steinhardt, Phys. Rev. Lett. 80, 1582 (1998).
- [8] E. J. Copeland, A. R. Liddle, and D. Wands, Phys. Rev. D 57, 4686 (1998).
- [9] P.J. Steinhardt, L. Wang, and I. Zlatev, Phys. Rev. D 59, 123504 (1999).
- [10] T. Barreiro, E. J. Copeland, and N. J. Nunes, Phys. Rev. D 61, 127301 (2000).
- [11] C. Armendáriz-Picón, T. Damour, and V. Mukhanov, Phys. Lett. B 458, 209 (1999).
- [12] T. Chiba, T. Okabe, and M. Yamaguchi, Phys. Rev. D 62, 023511 (2000).
- [13] V. Mukhanov and A. Vikman, J. Cosmol. Astropart. Phys. 02 (2006) 004.
- [14] M. Chevallier and D. Polarski, Int. J. Mod. Phys. D 10, 213 (2001).
- [15] E. V. Linder, Phys. Rev. Lett. 90, 091301 (2003).
- [16] http://www.euclid-ec.org/.
- [17] R. Laureijs et al., arXiv:1110.3193.
- [18] L. Amendola et al., Living Rev. Relativ. 16, 6 (2013).
- [19] http://www.lsst.org.
- [20] LSST Dark Energy Science Collaboration, arXiv: 1211.0310.
- [21] https://www.skatelescope.org/.
- [22] P. Bull, S. Camera, A. Raccanelli, C. Blake, P. Ferreira, M. Santos, and D. J. Schwarz, arXiv:1501.04088.

- [23] S. Camera, A. Raccanelli, P. Bull, D. Bertacca, X. Chen, P. Ferreira, M. Kunz, R. Maartens, Y. Mao, M. Santos, P. R. Shapiro, M. Viel, and Y. Xu, arXiv:1501.03851.
- [24] A. Raccanelli, P. Bull, S. Camera, C. Blake, P. Ferreira, R. Maartens, M. Santos, P. Bull, D. Bacon, O. Doré, P. Ferreira, M. G. Santos, M. Viel, and G. B. Zhao, arXiv:1501.03821.
- [25] M. G. Santos, P. Bull, D. Alonso, S. Camera, P. G. Ferreira, G. Bernardi, R. Maartens, M. Viel, F. Villaescusa-Navarro, F. B. Abdalla, M. Jarvis, R. B. Metcalf, A. Pourtsidou, and L. Wolz, arXiv:1501.03989.
- [26] S. Hannestad and E. Mörtsell, J. Cosmol. Astropart. Phys. 09 (2004) 001.
- [27] H. K. Jassal, J. S. Bagla, and T. Padmanabhan, Phys. Rev. D 72, 103503 (2005).
- [28] E. M. Barboza and J. S. Alcaniz, Phys. Lett. B 666, 415 (2008).
- [29] V. Sahni, A. Shafieloo, and A. A. Starobinsky, Phys. Rev. D 78, 103502 (2008).
- [30] C. Zunckel and C. Clarkson, Phys. Rev. Lett. 101, 181301 (2008).
- [31] U. Alam, V. Sahni, T. D. Saini, and A. A. Starobinsky, Mon. Not. R. Astron. Soc. 344, 1057 (2003).
- [32] V. Sahni, T. D. Saini, A. A. Starobinsky, and U. Alam, JETP Lett. 77, 201 (2003).
- [33] R. G. Crittenden, L. Pogosian, and G.-B. Zhao, J. Cosmol. Astropart. Phys. 12 (2009) 025.
- [34] V. Sahni and A. Starobinsky, Int. J. Mod. Phys. D 15, 2105 (2006).
- [35] M. Sahlén, A. R. Liddle, and D. Parkinson, Phys. Rev. D 72, 083511 (2005).
- [36] I. J. Grivell and A. R. Liddle, Phys. Rev. D 61, 081301 (2000).
- [37] E. J. Copeland, E. W. Kolb, A. R. Liddle, and J. E. Lidsey, Phys. Rev. D 48, 2529 (1993).

- [38] R. A. Daly and S. G. Djorgovski, Astrophys. J. 597, 9 (2003).
- [39] R. A. Daly and S. G. Djorgovski, Astrophys. J. 612, 652 (2004).
- [40] J. Simon, L. Verde, and R. Jimenez, Phys. Rev. D 71, 123001 (2005).
- [41] Z. Slepian, J. R. Gott, and J. Zinn, Mon. Not. R. Astron. Soc. 438, 1948 (2014).
- [42] R. Crittenden, E. Majerotto, and F. Piazza, Phys. Rev. Lett. 98, 251301 (2007).
- [43] T. Chiba, Phys. Rev. D 79, 083517 (2009).
- [44] T. Chiba, Phys. Rev. D 80, 109902 (2009).
- [45] T. Chiba, S. Dutta, and R. J. Scherrer, Phys. Rev. D 80, 043517 (2009).
- [46] R. R. Caldwell, Phys. Lett. B 545, 23 (2002).
- [47] G. F. R. Ellis and M. S. Madsen, Classical Quantum Gravity 8, 667 (1991).
- [48] V. Sahni and A. Starobinsky, Int. J. Mod. Phys. D 9, 373 (2000).
- [49] A. Paliathanasis, M. Tsamparlis, S. Basilakos, and J. D. Barrow, Phys. Rev. D 91, 123535 (2015).
- [50] C. Armendariz-Picon, V. Mukhanov, and P.J. Steinhardt, Phys. Rev. Lett. 85, 4438 (2000).
- [51] C. Armendariz-Picon, V. Mukhanov, and P.J. Steinhardt, Phys. Rev. D 63, 103510 (2001).
- [52] T. Padmanabhan, Phys. Rev. D 66, 021301 (2002).
- [53] L. P. Chimento, Phys. Rev. D 69, 123517 (2004).
- [54] J. Garriga and V.F. Mukhanov, Phys. Lett. B 458, 219 (1999).
- [55] E. J. Copeland, M. R. Garousi, M. Sami, and S. Tsujikawa, Phys. Rev. D 71, 043003 (2005).
- [56] A. D. Rendall, Classical Quantum Gravity 23, 1557 (2006).
- [57] J. De-Santiago, J. L. Cervantes-Cota, and D. Wands, Phys. Rev. D 87, 023502 (2013).
- [58] J. De-Santiago and J. L. Cervantes-Cota, J. Phys. Conf. Ser. 485, 012017 (2014).

- [59] M. Gasperini and G. Veneziano, Phys. Rep. 373, 1 (2003).
- [60] L. R. Abramo and F. Finelli, Phys. Lett. B 575, 165 (2003).
- [61] J. M. Aguirregabiria and R. Lazkoz, Phys. Rev. D 69, 123502 (2004).
- [62] N. A. Hamed, H. S. Cheng, M. A. Luty, and S. Mukohyama, J. High Energy Phys. 05 (2004) 074.
- [63] F. Piazza and S. Tsujikawa, J. Cosmol. Astropart. Phys. 07 (2004) 004.
- [64] M. Alishahiha, E. Silverstein, and D. Tong, Phys. Rev. D 70, 123505 (2004).
- [65] E. Silverstein and D. Tong, Phys. Rev. D 70, 103505 (2004).
- [66] Z.-K. Guo and N. Ohta, J. Cosmol. Astropart. Phys. 04 (2008) 035.
- [67] N. Bose and A. S. Majumdar, Phys. Rev. D 80, 103508 (2009).
- [68] J. De-Santiago and J. L. Cervantes-Cota, Phys. Rev. D 83, 063502 (2011).
- [69] M. Sharif, K. Yesmakhanova, S. Rani, and R. Myrzakulov, Eur. Phys. J. C 72, 2067 (2012).
- [70] R. J. Scherrer, Phys. Rev. Lett. 93, 011301 (2004).
- [71] A. Sen, Mod. Phys. Lett. A 17, 1797 (2002).
- [72] A. Sen, J. High Energy Phys. 04 (2002) 048.
- [73] A. Sen, Int. J. Mod. Phys. A 20, 5513 (2005).
- [74] R.-J. Yang and S.-N. Zhang, Chin. Phys. Lett. 25, 344 (2008).
- [75] R. Yang and J. Qi, Eur. Phys. J. C 72, 2095 (2012).
- [76] R. Yang, B. Chen, J. Li, and J. Qi, Astrophys. Space Sci. 356, 399 (2015).
- [77] A. A. Mamon and S. Das, Eur. Phys. J. C 75, 244 (2015).
- [78] Z. Ossoulian, T. Golanbari, H. Sheikhahmadi, and K. Saaidi, Adv. High Energy Phys. 2016, 3047461 (2016).
- [79] R. A. Battye, A. Moss, and J. A. Pearson, J. Cosmol. Astropart. Phys. 04 (2015) 048.
- [80] F. Pace, C. Fedeli, L. Moscardini, and M. Bartelmann, Mon. Not. R. Astron. Soc. 422, 1186 (2012).