

Local effects of the quantum vacuum in Lorentz-violating electrodynamicsA. Martín-Ruiz^{*}¹*Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México,
04510 México, Distrito Federal, México*C. A. Escobar[†]²*CENTRA, Departamento de Física, Universidade do Algarve, 8005-139 Faro, Portugal
(Received 5 November 2016; published 15 February 2017)*

The Casimir effect is one of the most remarkable consequences of the nonzero vacuum energy predicted by quantum field theory. In this paper we use a local approach to study the Lorentz violation effects of the minimal standard model extension on the Casimir force between two parallel conducting plates in the vacuum. Using a perturbative method similar to that used for obtaining the Born series for the scattering amplitudes in quantum mechanics, we compute, at leading order in the Lorentz-violating coefficients, the relevant Green's function which satisfies given boundary conditions. The standard point-splitting technique allow us to express the vacuum expectation value of the stress-energy tensor in terms of the Green's function. We discuss its structure in the region between the plates. We compute the renormalized vacuum stress, which is obtained as the difference between the vacuum stress in the presence of the plates and that of the vacuum. The Casimir force is evaluated in an analytical fashion by two methods: by differentiating the renormalized global energy density and by computing the normal-normal component of the renormalized vacuum stress. We compute the local Casimir energy, which is found to diverge as approaching the plates, and we demonstrate that it does not contribute to the observable force.

DOI: [10.1103/PhysRevD.95.036011](https://doi.org/10.1103/PhysRevD.95.036011)**I. INTRODUCTION**

Symmetry principles play a fundamental role in theoretical physics. As an outstanding example, Lorentz invariance is one of the cornerstones of general relativity (GR) and the standard model (SM) of particle physics. Although both theories have been successful in explaining and predicting the observed physical phenomena with a high degree of accuracy, they come with their own set of deficiencies: ultraviolet divergences in quantum field theories and singularities in general relativity. It is generally believed that a unified quantum theory of gravity will solve these notable problems; and thus its search has become one of the most important challenges of theoretical physicists.

Interest in Lorentz violation [1–6] has grown rapidly in the last decades since many candidate theories of quantum gravity [7,8], such as string theory [9] and loop quantum gravity [10], possess scenarios involving deviations from Lorentz symmetry. Nowadays, investigations concerning Lorentz violation are mostly conducted under the framework of the standard model extension (SME), initiated by Kostelecký and Colladay [11,12]. The SME is an effective field theory that contains the standard model, general relativity, and all possible operators that break Lorentz invariance. The Lorentz-violating (LV) coefficients arise

as vacuum expectation values of some basic fields belonging to a more fundamental theory, such as string theory [13]. Some important features of the minimal SME comprise invariance under observer Lorentz transformations, energy-momentum conservation, gauge invariance, power-counting renormalizability [14–16], causality, stability, and Hermiticity [17].

A Lorentz-violating vacuum acts in many respects like a nontrivial optical medium. Therefore, one expects the photon sector of the SME to possess features similar to those of ordinary electrodynamics in macroscopic media. The electrodynamics limit of the SME has been widely studied in the literature [18]. Indeed, changes in the propagation of light, such as polarization, birefringence [18–20], and Čerenkov effect [21–23], have been predicted and used to place tight bounds on Lorentz violation. The main goal of this work is to provide additional contributions regarding the local effects of the quantum vacuum in a particular sector of the electrodynamics limit of the SME, namely, the *CPT*-odd Maxwell-Chern-Simons term [24]. Concretely, we study the Casimir effect (CE) between two parallel conducting plates using a local approach based on the calculation of the vacuum expectation value of the stress-energy tensor via Green's functions satisfying the suitable boundary conditions.

In its simple manifestation, the CE is a quantum force of attraction between two parallel uncharged conducting plates [25]. More generally, it refers to the stress on bounding surfaces when a quantum field is confined to a

^{*}alberto.martin@nucleares.unam.mx
[†]cruiz@ualg.pt

finite volume of space. The boundaries can be material media, interfaces between two phases of the vacuum, or topologies of space. In any case, the modes of the quantum fields are restricted, giving rise to a macroscopically measurable force [26]. The CE has been studied in different scenarios, including the standard model, the gravitational sector [27], the AdS-CFT correspondence [28], theories with minimal length [29] and extra dimensions [30], condensed matter systems [31], and chiral metamaterials [32], to name a few.

The CE has also been considered within the SME framework [33–35]. The authors in Ref. [35] used the zeta function regularization technique to compute the Casimir force between two parallel conducting plates within the $(3 + 1)$ D Maxwell-Chern-Simons theory. As a consistency check, they also evaluated the renormalized vacuum energy by a series summation (via the residue theorem) of one-particle energy eigenstates. The first attempt to tackle this problem was due to Frank and Turan [34]; however, as pointed out by Kharlanov and Zhukovsky [35], they used misinterpreted equations which led to an oversimplified treatment of the problem. More precisely, they considered that the photon dispersion relation corresponds to that for a massive photon; however, unlike the $(2 + 1)$ D case, in $(3 + 1)$ D the effect of the Maxwell-Chern-Simons term is a more complicated dispersion relation for the photon. Due to this wrong equation, Frank and Turan constructed also incorrectly the relevant Green's function (GF). One of the specific aims of this work is the construction of the correct Green's function within the $(3 + 1)$ D Maxwell-Chern-Simons theory and the calculation of the Casimir energy density and stress between two parallel conducting plates. Furthermore, the method can be further generalized to diverse geometries. The basics for the construction of the GF is that the Lorentz-violating field equations can be treated perturbatively due to the smallness of the LV coefficients [36]. Our Green's functions also provide information about the divergence of the local energy density near the plates.

The outline of this paper is as follows. Section II reviews some basics of the particular sector of the minimal SME to be considered in this work, namely, the $(3 + 1)$ D Maxwell-Chern-Simons model. Using a perturbative method similar to that used for obtaining the Born series for the scattering amplitudes in quantum mechanics, in Sec. III we compute the leading-order Green's function which satisfies given Dirichlet, Neumann, or Robin boundary conditions, provided the smallness of the LV coefficients. In Sec. IV we use the standard point-splitting technique to express the vacuum expectation value of the stress-energy tensor in terms of the Green's function. The concrete calculation of the renormalized vacuum stress (and the Casimir force) between two parallel conducting plates is performed in Sec. V. We also discuss the local energy density, which is found to diverge as approaching the plates. We demonstrate

that the divergent term does not contribute to the observable force. The conclusions are contained in Sec. VI. Details of technical computations are left to the Appendix. Here, Lorentz-Heaviside units are assumed ($\hbar = c = 1$), the metric signature will be taken as $(+, -, -, -)$, and the convention $\epsilon^{0123} = +1$ is adopted.

II. LORENTZ-VIOLATING ELECTRODYNAMICS

The renormalizable gauge-invariant photon sector of the SME consists of the usual Maxwell Lagrangian plus the additional terms $\frac{1}{2}(k_{AF})^\kappa \epsilon_{\kappa\lambda\mu\nu} A^\lambda F^{\mu\nu}$ and $-\frac{1}{4}(k_F)_{\alpha\beta\mu\nu} F^{\alpha\beta} F^{\mu\nu}$. The LV tensor coefficients $(k_{AF})^\kappa$ and $(k_F)_{\alpha\beta\mu\nu}$ are *CPT* odd and *CPT* even, respectively. Many components of these coefficients are strongly constrained by astrophysical spectropolarimetry [24]. Despite this, further investigations remain to be of great interest both for a better understanding of massless Lorentz-violating fields and for the potential complementary tighter bounds.

In this paper we are concerned with the *CPT*-odd sector. The relevant Lagrangian is

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (k_{AF})_\mu A_\nu \tilde{F}^{\mu\nu} - j_\mu A^\mu. \quad (1)$$

Here, $j^\mu = (\rho, \mathbf{J})$ is the 4-current source that couples to the electromagnetic 4-potential A^μ , $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$ is the electromagnetic field strength and $\tilde{F}^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\alpha\beta}F_{\alpha\beta}$ its dual. Since we take $(k_F)_{\mu\nu\alpha\beta} = 0$, we can omit the subscript *AF* of the Lorentz- and *CPT*-violating $(k_{AF})^\mu$ coefficients and set $(k_{AF})^\mu \equiv k^\mu = (k^0, \mathbf{k})$. A nondynamical fixed k^μ determines a special direction in spacetime. For example, certain features of plane wave propagating along \mathbf{k} might differ from those of waves perpendicular to \mathbf{k} . Thus, particle Lorentz transformations are violated.

Varying the action $\mathcal{S} = \int \mathcal{L} d^4x$ with respect to A^μ yields the equations of motion:

$$(\square \eta^\mu{}_\nu - \partial^\mu \partial_\nu - 2k_\beta \epsilon^{\mu\beta\alpha} \partial_\alpha) A^\nu = j^\mu, \quad (2)$$

which extend the usual covariant Maxwell equations to incorporate Lorentz violation. Of course, the homogeneous Maxwell equations that express the field-potential relationship

$$\partial_\mu \tilde{F}^{\mu\nu} = 0 \quad (3)$$

are not modified due to the $U(1)$ gauge invariance of the action. As in conventional electrodynamics, current conservation $\partial_\mu j^\mu = 0$ can be verified directly by taking the divergence at both sides of Eq. (2).

In noncovariant notation, the inhomogeneous equations of motion (2) read

$$\begin{aligned}\nabla \cdot \mathbf{E} - 2\mathbf{k} \cdot \mathbf{B} &= \rho, \\ \nabla \times \mathbf{B} - \dot{\mathbf{E}} - 2k^0 \mathbf{B} + 2\mathbf{k} \times \mathbf{E} &= \mathbf{J},\end{aligned}\quad (4)$$

while the homogeneous equations (3) are $\nabla \cdot \mathbf{B} = 0$ and $\nabla \times \mathbf{E} + \dot{\mathbf{B}} = 0$. Gauge invariance of physics is evident from Eqs. (4), and any of the usual conditions on A^μ , like Lorentz or Coulomb gauge, can be imposed. Interestingly, the microscopic equations (4) can be cast in the form of Maxwell equations for macroscopic media:

$$\nabla \cdot \mathbf{D} = \rho, \quad \nabla \times \mathbf{H} - \dot{\mathbf{D}} = \mathbf{J}, \quad (5)$$

with the modified constitutive relations

$$\mathbf{D} = \mathbf{E} + \theta(\mathbf{x}, t)\mathbf{B}, \quad \mathbf{H} = \mathbf{B} - \theta(\mathbf{x}, t)\mathbf{E}, \quad (6)$$

where $\theta(\mathbf{x}, t) = 2k_\mu x^\mu = 2k^0 t - 2\mathbf{k} \cdot \mathbf{x}$ is a spacetime dependent axion field. These equations reveal a remarkable feature of this theory, the magnetoelectric effect, which also occurs in the *CPT*-even photon sector of the SME [33]. Furthermore, Eqs. (5) together with the constitutive relations of Eq. (6) resemble those describing the electromagnetic response of condensed matter systems, in which case, the spatial and temporal dependence of the axion field defines a specific realization of a topologically nontrivial phase of matter. For example, $\theta = \pi$ in the case of 3D topological insulators [37], and $\theta(\mathbf{x}, t) = 2b^0 t - 2\mathbf{b} \cdot \mathbf{x}$ for Weyl semimetals [38], where the parameter $2b^0$ is interpreted as the separation between the nodes in energy, and $2\mathbf{b}$ denotes the separation between the Weyl nodes in momentum space. However, note that for condensed matter systems, θ is defined inside the material, while in the SME framework k_μ permeates the whole spacetime.

The stress-energy tensor for this theory is given by

$$\Theta^{\mu\nu} = -F^{\mu\alpha} F^\nu{}_\alpha + \frac{1}{4} \eta^{\mu\nu} F^{\alpha\beta} F_{\alpha\beta} - k^\nu \tilde{F}^{\mu\alpha} A_\alpha. \quad (7)$$

Here $\eta^{\mu\nu} = \text{diag}(1, -1, -1, -1)$ denotes the usual Minkowski flat space-time metric. Unlike the conventional case, $\Theta^{\mu\nu}$ cannot be symmetrized because its antisymmetric part is no longer a total derivative. By virtue of the equations of motion (2) and (3), the energy-momentum tensor obeys

$$\partial_\mu \Theta^{\mu\nu} = j_\mu F^{\mu\nu}, \quad (8)$$

which implies that it is conserved in the absence of sources. Although the energy-momentum tensor is gauge dependent, it only changes by a total derivative under the gauge transformation $A^\mu \rightarrow A^\mu + \partial^\mu \Lambda$, i.e.

$$\tilde{F}^{\mu\alpha} A_\alpha \rightarrow \tilde{F}^{\mu\alpha} A_\alpha - \partial_\alpha (\tilde{F}^{\mu\alpha} \Lambda). \quad (9)$$

Consequently, the integrals over all space are gauge invariant. Note that the energy

$$\mathcal{E} = \int \Theta^{00} d^3 \mathbf{x} = \int \frac{1}{2} (\mathbf{E}^2 + \mathbf{B}^2 - k^0 \mathbf{B} \cdot \mathbf{A}) d^3 \mathbf{x} \quad (10)$$

is not positive definite due to the term $k^0 \mathbf{B} \cdot \mathbf{A}$, which may be negative. The appearance of this term in the energy density can introduce instability in the theory [39], and it can be resolved by requiring that only spacelike components of k^μ are nonzero. However, this condition depends on the observer frame, so even an infinitesimal boost to another observer frame would reintroduce instability. Despite arising from a hitherto unobserved spontaneous breaking of the electromagnetic $U(1)$ gauge symmetry, the photon mass can be introduced in this theory to eliminate the linear instability. Although this idea might be physically acceptable, in this work we restrict ourselves to the minimal extension with a purely spacelike background $k^\mu = (0, \mathbf{k})$, which is fundamentally different from the theory with the purely timelike case $k^\mu = (k^0, \mathbf{0})$, as reviewed extensively in Refs. [40].

III. GREEN'S FUNCTION METHOD

Knowledge of Green's function (GF) allows one to compute the electromagnetic fields for an arbitrary distribution of sources, as well as to solve problems with given Dirichlet, Neumann, or Robin boundary conditions on arbitrary surfaces. To derive the GF for the previously discussed LV electrodynamics one can employ standard Fourier methods. As in conventional electrodynamics, the modified Maxwell operator appearing in parentheses in Eq. (2) is singular, as one can verify in Fourier space. To circumvent the noninvertibility of the corresponding Minkowski matrix one can further work in Lorentz gauge. The free-space GF (satisfying the standard boundary conditions at infinity) in momentum [23] and coordinate [40] representations can be obtained in a simple fashion. In this paper we are concerned with the effects of this Lorentz-violating electrodynamics on the Casimir force between two parallel conducting plates in the vacuum. To this end we employ a local approach consisting in the evaluation of the vacuum expectation value of the stress-energy tensor of the system, which can be expressed in terms of the appropriate Green's function. The presence of boundaries (e.g. the plates) makes the GF derived in Refs. [23,40] not suitable for our purposes. Thus the aim of this section is the construction of the Green's functions which incorporates the presence of boundaries.

In the Lorentz gauge $\partial_\mu A^\mu = 0$, the field equations (2) take the form

$$(\square \eta^\mu{}_\nu - 2k_\rho \epsilon^{\mu\beta\alpha}{}_\nu \partial_\alpha) A^\nu = j^\mu, \quad (11)$$

where $\square = \partial_\mu \partial^\mu = \partial_t^2 - \nabla^2$ is the D'Alembert operator. To obtain the general solution of Eq. (11) for arbitrary external sources, we introduce the GF matrix $G^\mu{}_\nu(x, x')$ solving Eq. (11) for a pointlike source,

$$(\square\eta^\mu{}_\nu - 2k_\beta\epsilon^{\mu\beta\alpha}{}_\nu\partial_\alpha)G^\nu{}_\gamma(x, x') = \eta^\mu{}_\gamma\delta^4(x - x'), \quad (12)$$

in such a way that the general solution for the 4-potential in the Lorentz gauge is

$$A^\mu(x) = \int G^\mu{}_\nu(x, x')j^\nu(x')d^4x. \quad (13)$$

Since the timelike theory appears to be inconsistent (that is, the theory violates unitarity and causality, or both), in this work we specialize in the purely spacelike case $k^\mu \equiv (0, \mathbf{k}) \equiv (0, 0, 0, \kappa)$. However, our subsequent analysis can be further generalized to a more complicated Chern-Simons background k^μ . Note that this condition makes the propagation of light anisotropic and defines a class of preferred inertial frames. Without loss of generality, we consider surfaces Σ_i which are orthogonal to \mathbf{k} in which Dirichlet, Neumann, or Robin boundary conditions have been imposed. In this way, the GF we consider has translational invariance in the directions x and y , while this invariance is broken in the z direction. Exploiting this symmetry we further introduce the Fourier transform in the direction parallel to the surfaces Σ_i , taking the coordinate dependence to be $\mathbf{R} = (x - x', y - y')$, and define

$$G^\mu{}_\nu(x, x') = \int \frac{d^2\mathbf{p}}{(2\pi)^2} e^{i\mathbf{p}\cdot(\mathbf{x}-\mathbf{x}')} \int \frac{d\omega}{2\pi} e^{-i\omega(t-t')} g^\mu{}_\nu(z, z'), \quad (14)$$

where $\mathbf{p} = (p_x, p_y)$ is the momentum parallel to Σ_i . In Eq. (14) we have suppressed the dependence of the reduced GF $g^\mu{}_\nu$ on ω and \mathbf{p} .

The substitution of Eq. (14) into Eq. (12) yields the reduced GF equation

$$(\tilde{\square}\eta^\mu{}_\nu - 2i\kappa\epsilon^{3\mu\beta}{}_\nu p_\beta)g^\nu{}_\alpha(z, z') = \eta^\mu{}_\alpha\delta(z - z'), \quad (15)$$

where $p^\alpha = (\omega, \mathbf{p}, 0)$ and $\tilde{\square} = \mathbf{p}^2 - \omega^2 - \partial_z^2$. We now must solve the reduced GF equation for the various components. At this point it is worth mentioning that the authors of Ref. [34] say that they derive the GF for a Chern-Simons-like theory; however, they used the wrong equation of motion $[\square + (k_{AF})^2]\epsilon^{\mu\nu\alpha\beta}\partial_\alpha A_\beta = 0$, which leads to a simple analysis in terms of a massive scalar field. Thus its solution is appropriate for the Proca theory, rather than for the Chern-Simons theory. Indeed, in Ref. [41], Milton presents a detailed derivation of the Casimir force for massive photons using different methods. In this section we derive the correct Green's function which solves the equations of motion for the (3 + 1)D Chern-Simons theory.

The solution to Eq. (15) is simple but not straightforward. Since the coefficient κ is assumed to be small, to solve it we employ a method similar to that used for obtaining the scattering amplitudes in quantum mechanics,

in which the Schrödinger equation can be written as an integral equation, the Lippmann-Schwinger equation, which can be iterated to obtain the Born series. Indeed, the Lippmann-Schwinger equation for Green's operator is called the resolvent identity. In the problem at hand let us consider that the free (with $\kappa = 0$) reduced GF is known, being the solution of $\tilde{\square}\mathbf{g}(z, z') = \delta(z - z')$ in the region $\mathfrak{D} \subseteq \mathbb{R}$ and satisfying appropriate boundary conditions on the surfaces $\Sigma_i \subseteq \mathbb{R}^2 = \{(x, y) : x, y \in \mathbb{R}\}$.

Now Eq. (15) can be directly integrated using the free reduced GF. We thus establish the integral equation

$$g^\mu{}_\nu(z, z') = \eta^\mu{}_\nu\mathbf{g}(z, z') + 2i\kappa\epsilon^{3\mu\alpha}{}_\beta p_\alpha \int_{\mathfrak{D}} \mathbf{g}(z, z'')g^\beta{}_\nu(z'', z')dz''. \quad (16)$$

Suppose we take this expression for $g^\beta{}_\nu$, and plug it under the integral sign. Iterating this procedure, we obtain a formal series for $g^\mu{}_\nu$. At leading order in the LV coefficient κ , the reduced GF can be written as the sum of two terms,

$$g^\mu{}_\nu(z, z') = \eta^\mu{}_\nu\mathbf{g}(z, z') + \mathbf{g}^\mu{}_\nu(z, z'). \quad (17)$$

The first term provides the propagation in the absence of Lorentz violation, while the second term, which can be shown to be

$$\mathbf{g}^\mu{}_\nu(z, z') = 2i\kappa\epsilon^\mu{}_\nu{}^{\alpha\beta} p_\alpha N(z, z') - 4\kappa^2[p^\mu p_\nu - (\eta^\mu{}_\nu + n^\mu n_\nu)p^2]M(z, z'), \quad (18)$$

encodes the Lorentz symmetry breakdown. In deriving Eq. (18) we have used the identity $\epsilon^{3\mu\beta}{}_\alpha\epsilon^{3\alpha\gamma}{}_\nu p_\beta p_\gamma = p^\mu p_\nu - (\eta^\mu{}_\nu + n^\mu n_\nu)p^2$, and

$$N(z, z') = \int_{\mathfrak{D}} \mathbf{g}(z, z'')\mathbf{g}(z'', z')dz'', \quad (19)$$

$$M(z, z') = \int_{\mathfrak{D}} \mathbf{g}(z, z'')N(z'', z')dz''. \quad (20)$$

Here $n^\mu = (0, 0, 0, 1)$ is the normal to the surfaces Σ_i . In the Appendix we present the evaluation of these functions for three simple cases which are relevant for the purposes of this paper: (a) in free-space, (b) for parallel conducting plates, and (c) for an infinite conducting plate.

Clearly, the full GF matrix $G^\mu{}_\nu$ can also be written as the sum of two terms,

$$G^\mu{}_\nu(x, x') = \eta^\mu{}_\nu\mathcal{G}(x, x') + \mathbf{G}^\mu{}_\nu(x, x'), \quad (21)$$

where \mathcal{G} and $\mathbf{G}^\mu{}_\nu$ are the Fourier transformations of $\mathbf{g}(z, z')$ and $\mathbf{g}^\mu{}_\nu$, respectively, as defined in Eq. (14). It is worth mentioning that the second term satisfies the Lorentz gauge condition, i.e. $\partial_\mu\mathbf{G}^\mu{}_\nu = 0$. The proof follows from the

reduced GF: $\partial_\mu \mathbf{G}^\mu{}_\nu \propto \int p_\mu \mathbf{g}^\mu{}_\nu$, which vanishes given that $\epsilon^\mu{}_\nu{}^{\alpha\beta} p_\mu p_\alpha = 0$ and $p_\mu n^\mu = 0$.

The reciprocity between the position of the unit charge and the position at which the GF is evaluated, $G_{\mu\nu}(x, x') = G_{\nu\mu}(x', x)$, is one of its most remarkable properties. From Eq. (14), this condition requires

$$g_{\mu\nu}(z, z', p^\alpha) = g_{\nu\mu}(z', z, -p^\alpha), \quad (22)$$

which we verify directly from Eq. (18).

IV. VACUUM STRESS-ENERGY TENSOR

In Sec. II we derived the stress-energy tensor (SET) for this theory and we showed that it can be written as the sum of two terms:

$$\Theta^{\mu\nu} = T^{\mu\nu} + \Xi^{\mu\nu}. \quad (23)$$

The first term,

$$T^{\mu\nu} = -F^{\mu\alpha} F^\nu{}_\alpha + \frac{1}{4} \eta^{\mu\nu} F^{\alpha\beta} F_{\alpha\beta}, \quad (24)$$

is the standard Maxwell stress-energy tensor, while the second,

$$\Xi^{\mu\nu} = -k^\nu \tilde{F}^{\mu\alpha} A_\alpha, \quad (25)$$

explicitly depends on the LV coefficients k^μ . Now we address the problem of the vacuum expectation value of the SET, to which we will refer simply as the vacuum stress (VS).

The local approach to compute the VS was initiated by Brown and Maclay who calculated the renormalized stress tensor by means of GF techniques [42]. Therein, the VS can be obtained from appropriate derivatives of the GF, in virtue of the formula

$$G^{\mu\nu}(x, x') = -i \langle 0 | \hat{T} A^\mu(x) A^\nu(x') | 0 \rangle. \quad (26)$$

Using the standard point splitting technique and taking the vacuum expectation value of the SET we find

$$\langle \Theta^{\mu\nu} \rangle = \langle T^{\mu\nu} \rangle + \langle \Xi^{\mu\nu} \rangle, \quad (27)$$

where the first term,

$$\begin{aligned} \langle T^{\mu\nu} \rangle = i \lim_{x' \rightarrow x} \left[-\partial^\mu \partial^\nu G^\lambda{}_\lambda + \partial^\mu \partial'_\lambda G^{\lambda\nu} + \partial^\lambda \partial^\nu G^\mu{}_\lambda \right. \\ \left. - \partial^{\lambda'} \partial_\lambda G^{\mu\nu} + \frac{1}{2} \eta^{\mu\nu} (\partial^\alpha \partial'_\alpha G^\lambda{}_\lambda - \partial^\alpha \partial'_\beta G^\beta{}_\alpha) \right] \end{aligned} \quad (28)$$

is the VS of the standard Maxwell SET, and

$$\langle \Xi^{\mu\nu} \rangle = -2ik^\nu \epsilon^{\mu\alpha\beta\gamma} \lim_{x' \rightarrow x} \partial'_\beta G_{\gamma\alpha}. \quad (29)$$

Here we have omitted the dependence of $G^{\mu\nu}$ on x and x' . This result can be further simplified as follows. Since the GF can be written as the sum of two terms, then Eq. (28) can also be written in the same way, i.e.

$$\langle T^{\mu\nu} \rangle = \langle t^{\mu\nu} \rangle + \langle \mathbf{T}^{\mu\nu} \rangle. \quad (30)$$

The first term,

$$\langle t^{\mu\nu} \rangle = -i \lim_{x' \rightarrow x} \left(2\partial^\mu \partial^\nu - \frac{1}{2} \eta^{\mu\nu} \partial^\lambda \partial'_\lambda \right) \mathcal{G}(x, x'), \quad (31)$$

is the vacuum stress in the absence of Lorentz violation. In obtaining Eq. (31) we used the fact that the zeroth-order GF (with $\kappa = 0$) is diagonal, i.e. it is equal to $\eta^\mu{}_\nu \mathcal{G}(x, x')$. The second term can be simplified since the GF $\mathbf{G}^\mu{}_\nu$ satisfies the Lorentz gauge condition $\partial_\mu \mathbf{G}^\mu{}_\nu = 0$. In this way,

$$\langle \mathbf{T}^{\mu\nu} \rangle = -i \lim_{x' \rightarrow x} \left[\partial^\mu \partial^\nu \mathbf{G} + \partial^{\lambda'} \partial_\lambda \left(\mathbf{G}^{\mu\nu} - \frac{1}{2} \eta^{\mu\nu} \mathbf{G} \right) \right], \quad (32)$$

where $\mathbf{G} = \mathbf{G}^\mu{}_\mu$ is the trace of $\mathbf{G}^\mu{}_\nu$. Finally the last term in Eq. (27) can be written as

$$\langle \Xi^{\mu\nu} \rangle = -2ik^\nu \epsilon^{\mu\alpha\beta\gamma} \lim_{x' \rightarrow x} \partial'_\beta \mathbf{G}_{\gamma\alpha}, \quad (33)$$

where we used again that the zeroth-order GF is diagonal.

V. CASIMIR EFFECT

Now let us consider the problem of calculating the renormalized VS $\langle \Theta^{\mu\nu} \rangle_{\text{ren}}$, which is obtained as the difference between the VS in the presence of boundaries and that of the vacuum. For two parallel conducting plates separated by a distance D in the z direction, one can construct the renormalized expectation value of the stress-energy tensor, using conservation, tracelessness, and symmetry arguments. The result is that it is uniform between the plates:

$$\langle t^{\mu\nu} \rangle_{\text{ren}} = -\frac{\pi^2}{720D^4} (\eta^{\mu\nu} + 4n^\mu n^\nu), \quad (34)$$

where $n^\mu = (0, 0, 0, 1)$ is the unit normal to the plates [42]. The Casimir stress is obtained by differentiating the Casimir energy $E_C = \langle t^{00} \rangle_{\text{ren}} D$ with respect to D , i.e. $F_C = -dE_C/dD = -\pi^2/240D^4$.

We turn now with the vacuum stress between two perfectly, conducting, infinite plates, separated by a distance D , embedded in the infinite Lorentz-violating vacuum. We simplify our configuration by orienting the coordinate frame so that the plates are perpendicular to the background LV vector $k^\mu = (0, \mathbf{k}) = \kappa n^\mu$; however, the general case proceeds along the same lines.

Using the Fourier representation of the GF in Eq. (14), together with the symmetry of the problem and the reduced Green's function given by Eq. (18), Eq. (32) can be written as

$$\langle \mathbf{T}^{\mu\nu} \rangle = -4i\kappa^2 \int \frac{d^2\mathbf{p}}{(2\pi)^2} \int \frac{d\omega}{2\pi} (p^\mu p^\nu + n^\mu n^\nu p^2) \lim_{z' \rightarrow z} (p^2 + \partial_z \partial'_z) M(z, z'). \quad (35)$$

From the rotational invariance around the z axis, the components of the stress perpendicular to n^μ , $\langle \Theta^{11} \rangle$ and $\langle \Theta^{22} \rangle$, are equal. In addition, from the mathematical structure of Eq. (33) we find the relation $\langle \mathbf{T}^{00} \rangle = -\langle \mathbf{T}^{11} \rangle$. These results, together with the tracelessness of $\langle \mathbf{T}^{\mu\nu} \rangle$, allow us to write the VS of Eq. (35) in the form

$$\langle \mathbf{T}^{\mu\nu} \rangle = (\eta^{\mu\nu} + 4n^\mu n^\nu) f(\kappa, z), \quad (36)$$

where

$$f(\kappa, z) = \frac{4\kappa^2}{i} \int \frac{d^2\mathbf{p}}{(2\pi)^2} \int \frac{d\omega}{2\pi} \omega^2 \lim_{z' \rightarrow z} (p^2 + \partial_z \partial'_z) M(z, z'). \quad (37)$$

Note that this term exhibits the same tensor structure as the result obtained by Brown and Maclay, but we obtain a z -dependent VS due to Lorentz violation. Using similar arguments, the last contribution to the VS given by Eq. (33) can be written as

$$\langle \Xi^{\mu\nu} \rangle = n^\mu n^\nu g(\kappa, z), \quad (38)$$

where

$$g(\kappa, z) = -8i\kappa^2 \int \frac{d^2\mathbf{p}}{(2\pi)^2} \int \frac{d\omega}{2\pi} p^2 \lim_{z' \rightarrow z} N(z, z'). \quad (39)$$

Therefore the renormalized vacuum expectation value of the stress-energy tensor $\Theta^{\mu\nu}$ between the conducting plates can be written as

$$\langle \Theta^{\mu\nu} \rangle_{\text{ren}} = \langle t^{\mu\nu} \rangle_{\text{ren}} + (\eta^{\mu\nu} + 4n^\mu n^\nu) f_{\text{ren}}(\kappa, z) + n^\mu n^\nu g_{\text{ren}}(\kappa, z), \quad (40)$$

where $\langle t^{\mu\nu} \rangle_{\text{ren}}$, which is given by Eq. (34), is the result obtained by Brown and Maclay in Lorentz symmetric electrodynamics [42]. Here, the renormalized functions $f_{\text{ren}}(\kappa, z)$ and $g_{\text{ren}}(\kappa, z)$ are defined as the difference between the functions in the presence of the plates and that of the vacuum. Note that the vacuum stress is symmetric, even when the SET is nonsymmetric. Next we derive the Casimir force between the plates.

A. Global Casimir energy density

The Casimir energy is defined as the energy per unit area stored in the electromagnetic field between the plates, i.e.

$$\mathcal{E}_C = \int_0^D \langle \Theta^{00} \rangle_{\text{ren}} dz, \quad (41)$$

where $\langle \Theta^{00} \rangle_{\text{ren}}$ is the renormalized 00-component of the stress-energy tensor, which according to the procedure described above, is obtained as the difference between the energy density in the presence of the plates and that of the free vacuum, i.e.

$$\langle \Theta^{00} \rangle_{\text{ren}} = \langle \Theta^{00} \rangle_{\parallel} - \langle \Theta^{00} \rangle_0. \quad (42)$$

Here, the labels \parallel and 0 mean that the expectation value is evaluated with the Green's function in the presence of the parallel plates and that of the vacuum, respectively. Using the formula (30), the Casimir energy (41) can be written as

$$\mathcal{E}_C = -\frac{\pi^2}{720D^3} + E_{\text{ren}}(\kappa), \quad (43)$$

where the first term is the usual Casimir energy density between two parallel conducting plates in Lorentz symmetric electrodynamics, i.e. $\int_0^D \langle t^{00} \rangle_{\text{ren}} dz$, and we define the function

$$E_{\text{ren}}(\kappa) = \int_0^D \langle \mathbf{T}^{00} \rangle_{\text{ren}} dz. \quad (44)$$

Here, $\langle \mathbf{T}^{00} \rangle_{\text{ren}} = \langle \mathbf{T}^{00} \rangle_{\parallel} - \langle \mathbf{T}^{00} \rangle_0$ is the renormalized Lorentz-violating contribution to the energy density. Here we interpret the labels in the same way as in Eq. (42). Now we must evaluate the function $E_{\text{ren}}(\kappa)$. First we consider the VS in the presence of the plates and define the function $E_{\parallel}(\kappa) \equiv \int_0^D \langle \mathbf{T}^{00} \rangle_{\parallel} dz$. Using the Green's function in the presence of the plates, i.e. Eq. (18) with the functions N_{\parallel} and M_{\parallel} given by Eqs. (A9) and (A10) respectively, we obtain

$$E_{\parallel}(\kappa) = -4i\kappa^2 \int \frac{d^2\mathbf{p}}{(2\pi)^2} \int \frac{d\omega}{2\pi} \frac{\omega^2}{8p^4} [4 - pD \cot(pD) - p^2 D^2 [1 + 2pD \cot(pD)] \csc^2(pD)]. \quad (45)$$

Notice that the first term in the integrand corresponds to a constant energy density, independent of D , so it may be discarded as irrelevant. The resulting integral can be evaluated as follows. We first write the momentum element as $d^2\mathbf{p} = |\mathbf{p}| d|\mathbf{p}| d\theta$ and integrate θ from 0 to 2π . Next we perform a Wick rotation such that $\omega \rightarrow i\zeta$, then replace ζ and $|\mathbf{p}|$ by the plane polar coordinates $\zeta = \frac{\xi}{D} \cos \varphi$, $|\mathbf{p}| = \frac{\xi}{D} \sin \varphi$ and finally integrate φ from 0 to $\pi/2$. In this way, Eq. (45) becomes

$$E_{\parallel}(\kappa) = \frac{\kappa^2}{24\pi^2 D} \int_0^{\infty} \xi [\coth \xi + \xi(1 + 2\xi \coth \xi) \operatorname{csch}^2 \xi] d\xi. \quad (46)$$

We can treat this result as containing two pieces. The first integral, which involves $\xi \coth \xi$, is divergent, so that we will retain such. The remaining part is convergent and it can be evaluated. The result is

$$E_{\parallel}(\kappa) = \frac{\kappa^2}{24\pi^2 D} \left[\frac{2\pi^2}{3} + \int_0^{\infty} \xi \coth \xi d\xi \right]. \quad (47)$$

Now we must evaluate the analogue function in vacuum. Using the Green's function in the free vacuum, i.e. Eq. (18) with the functions N_0 and M_0 given by Eqs. (A5) and (A6) respectively, we obtain

$$E_0(\kappa) = 4\kappa^2 \int \frac{d^2 \mathbf{p}}{(2\pi)^2} \int \frac{d\omega \omega^2 D}{2\pi 8p^3}. \quad (48)$$

The integral can be treated similarly. Integrating the angle in momentum space, performing a Wick rotation and introducing the plane polar coordinates defined before we find:

$$E_0(\kappa) = \frac{\kappa^2}{24\pi^2 D} \int_0^{\infty} \xi d\xi, \quad (49)$$

which clearly is a divergent integral. Subtracting both contributions to obtain the renormalized function (44), i.e. $E_{\text{ren}}(\kappa) = E_{\parallel}(\kappa) - E_0(\kappa)$, we obtain

$$E_{\text{ren}}(\kappa) = \frac{\kappa^2}{24\pi^2 D} \left[\frac{2\pi^2}{3} + \int_0^{\infty} \xi (\coth \xi - 1) d\xi \right], \quad (50)$$

and the resulting integral is perfectly convergent, with the result $\pi^2/12$. Therefore the final expression for the renormalized function is $E_{\text{ren}}(\kappa) = \kappa^2/32D$, and thus the Casimir energy becomes

$$\mathcal{E}_C = -\frac{\pi^2}{720D^3} + \frac{\kappa^2}{32D}. \quad (51)$$

The Casimir stress is obtained by differentiating the Casimir energy with respect to D , i.e.

$$\mathcal{F}_C = -\frac{d\mathcal{E}_C}{dD} = -\frac{\pi^2}{240D^4} + \frac{\kappa^2}{32D^2}. \quad (52)$$

The first term is recognized as the usual Casimir force in Lorentz-symmetric electrodynamics, while the second represents the Lorentz-violating contribution. We observe that unlike the $1/D^4$ dependence of the usual Casimir force, the LV part depends on $1/D^2$. This is so because of the Chern-Simons background field k^μ is dimensionfull, indeed $[k^\mu] = 1/[\text{length}]$. As discussed in Sec. I, the authors in

Ref. [35] analyzed the same Casimir configuration within the (3 + 1)D Chern-Simons extended electrodynamics.

They obtained an attractive correction of the form $-\frac{25\kappa_{AF}^2}{720D^2}$, while our result (52) is evidently repulsive. The origin of such disagreement is the Chern-Simons background tensor we work with; while the authors in Ref. [35] used a timelike background $k^\mu = (k_{AF}, \mathbf{0})$, we have chosen the spacelike vector $k^\mu = (0, \mathbf{k})$. Here we point out that the difference between our work and that of Ref. [35] is beyond the background tensor used. For example, they used standard techniques in Casimir physics (zeta function regularization and series summation via the residue theorem), while here we tackle the problem by means of Green's function techniques. The advantage of the local approach over the standard techniques is quite subtle: the former provides information about the local behavior of the quantum fields near a boundary, while the later only allows the calculation of the Casimir force. In Sec. V C we discuss the local effects in (3 + 1)D Chern-Simons electrodynamics, which is the main aim of the present work.

B. Stress on the plates

Now let us derive the same result by using the normal-normal component of the vacuum stress-energy tensor, i.e. $\langle \Theta^{zz} \rangle$. In the problem at hand, the pressure exerted by the electromagnetic field will be the difference between the $\langle \Theta^{zz} \rangle$ components outside the plates and that between the plates, calculated on the surface (at $z = D$, for instance)

$$\mathcal{F}_C = \langle \Theta^{zz} \rangle_{\parallel} - \langle \Theta^{zz} \rangle_{\perp}. \quad (53)$$

The first term, $\langle \Theta^{zz} \rangle_{\parallel}$, is the vacuum stress due to the electromagnetic field confined between the plates. The second term, $\langle \Theta^{zz} \rangle_{\perp}$, is the vacuum stress due to the electromagnetic field outside the plates. As before, the label \parallel means that the vacuum expectation value is calculated with the Green's function in the presence of the plates, which is given by Eqs. (17) and (18) with the functions N_{\parallel} and M_{\parallel} given by Eqs. (A9) and (A10), respectively. In order to calculate the vacuum expectation value due to the electromagnetic field outside the plates, $\langle \Theta^{zz} \rangle_{\perp}$, we should use the appropriate Green's function which vanishes at $z = D$ and has outgoing boundary conditions as $z \rightarrow \infty$, i.e. $\mathbf{g} \sim e^{ipz}$ [41]. Basically, we should use the Green's function due to a single conducting plate at $z = D$, and this is why we use the label \perp for this case. In the Appendix, we compute the corresponding N_{\perp} and M_{\perp} functions.

Using the formula (30), the Casimir stress (41) can be written as

$$\mathcal{F}_C = -\frac{\pi^2}{240D^4} + F_{\text{ren}}(\kappa), \quad (54)$$

where the first term in the Casimir stress in the absence of Lorentz violation, i.e. $\langle t^{zz} \rangle_{\parallel} - \langle t^{zz} \rangle_{\perp} = -\pi^2/240D^4$. The Lorentz-violating contribution is given by $F_{\text{ren}}(\kappa) = \langle \mathbf{T}^{zz} + \Xi^{zz} \rangle_{\parallel} - \langle \mathbf{T}^{zz} + \Xi^{zz} \rangle_{\perp}$. By virtue of the boundary conditions upon the plates, Ξ^{zz} does not contribute to the vacuum expectation values. Then we compute the normal-normal component of the vacuum stress tensor on the boundaries as

$$\langle \mathbf{T}^{zz} \rangle = -12i\kappa^2 \int \frac{d^2\mathbf{p}}{(2\pi)^2} \int \frac{d\omega}{2\pi} \omega^2 \lim_{z' \rightarrow z=0,D} \partial_z \partial'_z M(z, z'), \quad (55)$$

since $M(z, z') = 0$ at $z = 0$ and $z = D$. First we consider the contribution due to the electromagnetic field confined between the plates, i.e. we want to evaluate Eq. (55) using the function $M_{\parallel}(z, z')$ given by Eq. (A10). The resulting integral can be treated in the same way as that of Eq. (44). Integrating the angle in momentum space, performing a Wick rotation and introducing the plane polar coordinates, we find

$$\begin{aligned} \langle \mathbf{T}^{zz} \rangle_{\parallel} = & -\frac{\kappa^2}{32\pi^2 D^2} \int_0^{\infty} \xi \text{csch}^3 \xi [\cosh(3\xi) \\ & - (1 + 8\xi^2) \cosh \xi + 4\xi \sinh \xi] d\xi. \end{aligned} \quad (56)$$

This integral does not exist. However, it is appropriately regularized by subtracting the stress due to the electromagnetic field outside the plates, as discussed. To do this, let us evaluate Eq. (55) using the function $M_{\perp}(z, z')$ given by Eq. (A14). In this case, the normal-normal component of the vacuum stress at $z = D$ becomes

$$\langle \mathbf{T}^{zz} \rangle_{\perp} = -\frac{\kappa^2}{8\pi^2 D^2} \int_0^{\infty} \xi d\xi, \quad (57)$$

which also diverges. So, from the discontinuity in $\langle \mathbf{T}^{zz} \rangle$, that is, the difference $F_{\text{ren}} = \langle \mathbf{T}^{zz} \rangle_{\parallel} - \langle \mathbf{T}^{zz} \rangle_{\perp}$, we find the Lorentz-violating contribution to the force per unit area on the plate:

$$\begin{aligned} F_{\text{ren}}(\kappa) = & -\frac{\kappa^2}{32\pi^2 D^2} \int_0^{\infty} \xi \{ \text{csch}^3 \xi [\cosh(3\xi) \\ & - (1 + 8\xi^2) \cosh \xi + 4\xi \sinh \xi] - 4 \} d\xi. \end{aligned} \quad (58)$$

The resulting integral is perfectly convergent, with the result $-\pi^2$. In this way, the stress on the plate is $\mathcal{F}_C = -\frac{\pi^2}{240D^4} + \frac{\kappa^2}{32D^2}$, which is the same as obtained in the previous subsection.

C. Local effects

Heretofore, we have considered the global Casimir effect: the total energy of a field configuration or the force

per unit area on a bounding surface [41]. Local properties of the quantum vacuum induced by the presence of boundaries are of broad interest in quantum field theory [43] and they must be understood if one is to correctly interpret the inherent divergences in the theory.

The local energy density in Lorentz-symmetric electrodynamics has been discussed extensively in the literature [41,44]; however, the local effects in Lorentz-violating theories have not been considered. Here we aim to fill in this gap. We begin the analysis by considering an electromagnetic field confined between two parallel conducting plates at $z = 0$ and $z = D$, for which the energy density per unit volume between the plates is

$$\langle \Theta^{00} \rangle(z) = \langle t^{00} \rangle(z) + \langle \mathbf{T}^{00} \rangle(z). \quad (59)$$

A detailed analysis of the local effects due to the first term (in the absence of Lorentz violation) is presented in Ref. [41]. Here we concentrate on the Lorentz-violating contribution,

$$\langle \mathbf{T}^{00} \rangle(z) = \frac{4\kappa^2}{i} \int \frac{d^2\mathbf{p}}{(2\pi)^2} \int \frac{d\omega}{2\pi} \omega^2 \lim_{z' \rightarrow z} (p^2 + \partial_z \partial'_z) M_{\parallel}(z, z'), \quad (60)$$

where $M_{\parallel}(z, z')$ is given by Eq. (A10). This integral can be performed in an analytical fashion. Integrating the angle in momentum space, performing a Wick rotation and introducing the plane polar coordinates, we find

$$\langle \mathbf{T}^{00} \rangle(z) = -\frac{\kappa^2}{3\pi^2 D^2} \int_0^{\infty} \lambda^4 [F(\lambda) + R(\lambda, Z)] d\lambda, \quad (61)$$

where

$$F(\lambda) = -\frac{\text{csch}^3 \lambda}{32\lambda^3} \{ (8\lambda^2 - 1) \cosh \lambda + \cosh(3\lambda) + 4\lambda \sinh \lambda \}, \quad (62)$$

$$\begin{aligned} R(\lambda, Z) = & \frac{\text{csch}^3 \lambda}{16\lambda^3} \{ 2\lambda(1-Z) \sinh[\lambda(1+2Z)] \\ & + 2\lambda(1-2Z) \sinh[\lambda(1-2Z)] + 2\lambda Z \sinh[\lambda(3-2Z)] \\ & + \cosh[\lambda(1+2Z)] - 2\cosh[\lambda(1-2Z)] \\ & + \cosh[\lambda(3-2Z)] \}, \end{aligned} \quad (63)$$

with $Z = z/D$. Integrating we obtain

$$\langle \mathbf{T}^{00} \rangle(z) = \frac{\kappa^2}{32D^2} [1 - 2\text{csc}^2(\pi Z)]. \quad (64)$$

We observe that the z -independent term, $\frac{\kappa^2}{32D^2}$, corresponds to the global renormalized energy density $E_{\text{ren}}(\kappa)$ obtained




FIG. 1. The singular part of the Lorentz-violating correction to the local energy density between two parallel plates.

in Sec. V A. The second term encodes the local effects, and it can be expressed in terms of the generalized or Hurwitz zeta function as follows:

$$\begin{aligned} S(Z) &\equiv -\frac{\kappa^2}{16D^2} \csc^2(\pi Z) \\ &= -\frac{\kappa^2}{16\pi^2 D^2} [\zeta(2, Z) + \zeta(2, 1 - Z)]. \end{aligned} \quad (65)$$

This function is plotted in Fig. 1, where we observe that it diverges quadratically as $z \rightarrow 0, D$. Its z integral over the region between the plates diverges linearly. This result reveals a close analogy with the one obtained from the Lorentz-symmetric part. In that case, the singular part depends on $[\zeta(4, Z) + \zeta(4, 1 - Z)]/D^4$, thus implying that it diverges quartically as $z \rightarrow 0, D$. The less divergent Lorentz-violating contribution (65) can be understood as due to the dimensionfull Chern-Simons coupling κ .

To close this section, we show that the badly behaved function (65) does not contribute to the force on the plates. Using the integral representation of the Hurwitz zeta function in terms of the Mellin transform,

$$\zeta(s, q) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{t^{s-1} e^{-qt}}{1 - e^{-t}} dt, \quad \Re[s] > 1, \quad \Re[q] > 0, \quad (66)$$

Eq. (65) can be expressed as

$$S(Z) = -\frac{\kappa^2}{16\pi^2} \frac{1}{\Gamma(2)} \int_0^\infty \frac{e^{-z\rho} + e^{-(D-z)\rho}}{1 - e^{-\rho D}} \rho d\rho. \quad (67)$$

If we integrate this term over z ,

$$\int_0^D S(Z) dz = -\frac{\kappa^2}{16\pi^2} \frac{2}{\Gamma(2)} \int_0^\infty d\rho, \quad (68)$$

we obtain a divergent constant term (D independent), so it does not contribute to the observable force.

VI. SUMMARY AND CONCLUSIONS

The study of Lorentz violation is actively motivated by the search of quantum gravity effects. At presently attainable energies, such signatures are described by an effective field theory framework called the standard model extension (SME), which contains the standard model, general relativity and all possible operators that break Lorentz symmetry. The Lorentz-violating (LV) coefficients in the SME can arise in various underlying contexts, such as strings, spacetime-foam approaches, and noncommutative geometries, to name a few. The value of these coefficients can in principle be measured (or bounded) in experiments. This has allowed a systematic search for a large range of possible Lorentz-violating effects. Indeed, since the predictions of the standard model have been verified experimentally to an extremely high degree of accuracy, a possible route to test quantum gravitational effects is through high-sensitivity measurements of well-known particle physics phenomena, as any deviation from the standard theory is, at least in principle, experimentally testable.

The main aim of this paper is to analyze the local effects of the quantum vacuum in a particular sector of the minimal SME, namely, the $(3 + 1)$ D Maxwell-Chern-Simons term. Concretely, we use Green's function techniques to calculate the Casimir force between two parallel conducting plates, focusing on the local properties of the quantum vacuum induced by the presence of the boundaries. The study of divergences is a fundamental problem in quantum field theories, and the present work intends to fill in this gap in a particular gauge sector of the SME. The first derivation of the (global) Casimir energy stored between two parallel conducting plates within the SME framework is due to Frank and Turan [34]. However, as pointed out by Kharlanov and Zhukovsky [35], in the aforementioned work the authors used a wrong equation of motion for the electromagnetic potentials, namely, $(\square + k_{AF}^2) \epsilon^{\mu\nu\alpha\beta} \partial_\alpha A_\beta = 0$, instead of the correct equation $(\square \eta_\nu^\mu - 2k_\beta \epsilon^{\mu\beta\alpha} \partial_\alpha) A^\nu = 0$. This misinterpreted equation led to an oversimplification of the problem from both the conceptual and theoretical points of view. Conceptually we observe that the wrong equation implies that the photon dispersion relation corresponds to that for a massive photon, similar to what occurs in the $(2 + 1)$ D Maxwell-Chern-Simons theory. However, in the $(3 + 1)$ D theory the photon remains massless and the associated dispersion relation is more complicated. Mathematically, the wrong field equation leads to a simple analysis of the problem in terms of a single Green's function; however, the correct field equation suggests that a more subtle indexed (matrix of) Green's function(s) is required due to the magnetoelectric effect.

The authors in Ref. [35] evaluated the Casimir force within this theory using two methods, namely, the zeta function regularization technique and the summation and renormalization of the discrete sum involving the residue

theorem. In the present work, we used Green's function techniques to analyze the same Casimir configuration. Mathematically, the difference between our work and that of Ref. [35] is the Chern-Simons background tensor we work with; while the authors in Ref. [35] used a timelike background $k^\mu = (k_{AF}, \mathbf{0})$, we have chosen the spacelike vector $k^\mu = (0, \mathbf{k})$. Our choice is motivated by the fact that the timelike theory appears to be inconsistent, that is, the theory violates unitarity and causality, or both [39]. We remark that the difference between our work and that of Ref. [35] is beyond the Lorentz-violating background tensor used. From the physical point of view, the techniques used in Ref. [35] provides information about the global Casimir energy density, but they fail to describe the behavior of the quantum fields near the boundary, and this is precisely the gap we intend to fill with the present work.

Because the Green's function represents the vacuum expectation value of the time-ordered product of fields, it is possible to compute the vacuum expectation value of the stress-energy tensor, for example, in terms of the Green's function at coincident arguments. One of the contributions of the present work is the construction of the correct (indexed) Green's function for the theory we deal with: the $(3+1)$ D Maxwell-Chern-Simons theory with a spacelike Lorentz-violating background field. Using these results, we evaluate the renormalized vacuum expectation value of the stress-energy tensor and we compute the Casimir force between two parallel conducting plates separated by a distance D , with the result $\mathcal{F}_C = -\frac{\pi^2}{240D^4} + \frac{\kappa^2}{32D^2}$. We recognize the first term as the Casimir force in the absence of Lorentz violation, while the κ -dependent term is due to such LV coefficient, which has a different functional dependence on the distance between the plates ($1/D^2$). This result is expected from dimensional analysis for the second order LV contributions. A remarkable difference between our result and those of Ref. [35] emerges: while they found an attractive correction of the form $-25k_{AF}^2/720D^2$, our result (52) is evidently repulsive. The origin of such disagreement is that the Lorentz-violating background tensor we deal with is different. Even when we work within the same gauge sector of the SME, the choice of the LV background makes these theories fundamentally different. Due to the limited precision of the present experimental measurements of the Casimir force between parallel plates (15% precision in the 0.5–3 μm range), no useful bounds on the LV coefficients can be obtained from the results in this work. We also analyze the behavior of the local energy density when approaching the plates, which is found to be quadratically divergent according to $\zeta(2, z/D) + \zeta(2, 1 - z/D)$, where $\zeta(s, q)$ is the Hurwitz zeta function. In the standard case a quartically divergent term is obtained. The less divergent Lorentz-violating contribution can be understood as due to the dimensionfull Chern-Simons coupling κ . Using the integral representation of the Hurwitz zeta function in terms of the Mellin transform,

we demonstrate that such a term does not contribute to the measurable macroscopic force.

The present work can be further generalized in a variety of ways. For example, the Green's function for different geometries can also be constructed using the same perturbative procedure. On the other hand, our analysis can also be applied to ponderable media. More precisely, we can consider a semi-infinite planar material medium with dielectric constant ε for which the reduced Green's function $\mathbf{g}_\varepsilon(z, z')$ is known. Now we can use it to evaluate the associated $E_\varepsilon(z, z')$ and $M_\varepsilon(z, z')$ functions, which are what we require to study the Lorentz-violating effects. This scenario could be more useful to establish bounds on the SME coefficients, since in this case first-order LV contributions could appear.

ACKNOWLEDGMENTS

We thank A. Kostelecký and L. F. Urrutia for many valuable discussions, comments, and suggestions. This work has been partially supported by the project DGAPA-UNAM, Project No. IN-104815 and the CONACyT Project No. 237503. C. A. E. is supported by CONACyT Postdoctoral Grant No. 234745.

APPENDIX: EVALUATION OF THE FUNCTIONS $N(z, z')$ AND $M(z, z')$

Here we shall evaluate the functions $N(z, z')$ and $M(z, z')$, defined by Eqs. (19) and (20), respectively. To do this, we use the reciprocity symmetry of the reduced GF, $\mathbf{g}(z, z') = \mathbf{g}(z', z)$, which implies the same property for the functions $N(z, z')$ and $M(z, z')$, i.e.

$$N(z, z') = N(z', z), \quad M(z, z') = M(z', z). \quad (\text{A1})$$

In this way, we can compute the required integrals for a particular case, for example $z < z'$, and then generalize the result by replacing $z \rightarrow z_<$ and $z' \rightarrow z_<$, where $z_>$ ($z_<$) will be the greater (lesser) between z and z' .

Let $\mathcal{D} = [a, b]$ be the domain in which the reduced GF is defined. Assuming that $z < z'$, the domain of integration can be expressed as $\mathcal{D} = [a, z] + [z, z'] + [z', b]$. The function $N(z, z')$ can thus be written as

$$N(z, z') = \left(\int_a^z + \int_z^{z'} + \int_{z'}^b \right) \mathbf{g}(z, z'') \mathbf{g}(z'', z') dz'', \quad (\text{A2})$$

and the function $M(z, z')$ takes the form

$$M(z, z') = \left(\int_a^z + \int_z^{z'} + \int_{z'}^b \right) \mathbf{g}(z, z'') N(z'', z') dz''. \quad (\text{A3})$$

Now we compute these integrals for some cases of particular interest in this paper.

1. Free space

First we consider the free space reduced Green's function

$$g_0(z, z') = \frac{i}{2p} e^{ip(z_>-z_<)}, \quad (\text{A4})$$

which is defined on the real line, i.e. $\mathcal{D} = \mathbb{R}$. The associated N_0 function can thus be written as

$$N_0(z, z') = -\frac{1}{4p^2} \left[\int_{-\infty}^z e^{ip(z-z'')} e^{ip(z'-z'')} dz'' + \int_z^{z'} e^{ip(z''-z)} e^{ip(z'-z'')} dz'' + \int_{z'}^{\infty} e^{ip(z''-z)} e^{ip(z'-z'')} dz'' \right].$$

The result for this particular case ($z < z'$) can be obtained in a simple fashion. The general result takes the form

$$N_0(z, z') = -\frac{i + p(z_> - z_<)}{4p^3} e^{ip(z_>-z_<)}. \quad (\text{A5})$$

Now, using the reduced GF of Eq. (A4) and the previous result (A5), we can compute the associated M_0 -function according to Eq. (A3). The result is

$$M_0(z, z') = \frac{p^2(z_> - z_<)^2 + 3ip(z_> - z_<) - 3}{16ip^5} e^{ip(z_>-z_<)}. \quad (\text{A6})$$

Performing a Wick rotation one can further see that these functions satisfy the required boundary conditions at infinity, i.e. $N_0(z \rightarrow \infty, z'), M_0(z \rightarrow \infty, z') \rightarrow 0$.

2. Parallel conducting plates

Now let us consider the case of two parallel conducting plates separated by a distance D . The relevant reduced GF for this configuration is

$$g_{\parallel}(z, z') = \frac{\sin[pz_<] \sin[p(D - z_>)]}{p \sin[pD]}, \quad (\text{A7})$$

which is defined on the domain $\mathcal{D}_{\parallel} = [0, D]$ and satisfies the boundary conditions $g_{\parallel}(0, z') = g_{\parallel}(D, z') = 0$. The associated N_{\parallel} function can be written as

$$N_{\parallel}(z, z') = \frac{1}{p^2 \sin^2[pD]} \left[\int_0^z \sin^2[pz''] \sin[p(D - z)] \sin[p(D - z')] dz'' + \int_z^{z'} \sin[pz] \sin[p(D - z'')] \sin[pz''] \sin[p(D - z')] dz'' + \int_{z'}^D \sin[pz] \sin[pz'] \sin^2[p(D - z'')] dz'' \right], \quad (\text{A8})$$

with the final result

$$N_{\parallel}(z, z') = \frac{1}{4p^3 \sin^2[pD]} \{ \sin[pz_>] \sin[pz_<] [2p(D - z_>) - \sin[2p(D - z_>)]] + \sin[p(D - z_>)] \sin[pz_<] [\sin[p(D - 2z_<)] - \sin[p(D - 2z_>)] - 2p(z_> - z_<) \cos[pD]] + \sin[p(D - z_>)] \sin[p(D - z_<)] [2pz_< - \sin[2pz_<]] \}. \quad (\text{A9})$$

With the help of this result and the reduced GF (A7) we can compute the M_{\parallel} function. We obtain

$$M_{\parallel}(z, z') = \frac{\csc^2[pD]}{32p^5} \{ 2pz_< \sin[pz_>] \cos[pz_<] [4pD - 2pz_> + 2pz_> \cos[2pD] + 3 \sin[2pD]] - \csc[pD] \sin[pz_>] \sin[pz_<] [(3 + 8D^2p^2 - p^2(z_>^2 + z_<^2)) \cos[pD] + (-3 + p^2(z_>^2 + z_<^2)) \cos[3pD]] - \csc[pD] \sin[pz_>] \sin[pz_<] (6p(2D - z_> + z_> \cos[2pD]) \sin[pD]) - 4pz_< \cos[pz_>] \cos[pz_<] \sin[pD] (2pz_> \cos[pD] + 3 \sin[pD]) - 2 \cos[pz_>] \sin[pz_<] [3 + 4Dp^2z_> - p^2z_>^2 - p^2z_<^2 + (-3 + p^2(z_>^2 + z_<^2)) \cos[2pD] + 3pz_> \sin[2pD]] \}. \quad (\text{A10})$$

On can further check that this function satisfies the boundary conditions $M_{\parallel}(0, z') = M_{\parallel}(D, z') = 0$, which are inherited from the standard boundary conditions on the conducting plates.

3. Infinite conducting plate

Finally we consider the case of a single conducting planar plate located at $z = D > 0$. The reduced GF describing the propagation at the right-hand side ($z, z' > D$) is

$$\mathfrak{g}_|(z, z') = \frac{1}{p} \sin[p(z_{<} - D)] e^{ip(z_{>} - D)}, \quad (\text{A11})$$

which is defined in the domain $\mathcal{D}_| = [D, \infty)$ and satisfies the boundary conditions $\mathfrak{g}_|(D, z') = 0$ and $\mathfrak{g}_|(z \rightarrow \infty, z') \sim e^{ikz}$. Using Eq. (A2) we can further compute the associated $N_||$ function, i.e.

$$\begin{aligned} N_|| (z, z') = \frac{1}{p^2} & \left[\int_D^z \sin[p(z'' - D)] e^{ip(z - D)} \sin[p(z'' - D)] e^{ip(z' - D)} dz'' \right. \\ & + \int_z^{z'} \sin[p(z - D)] e^{ip(z'' - D)} \sin[p(z'' - D)] e^{ip(z' - D)} dz'' \\ & \left. + \int_{z'}^\infty \sin[p(z - D)] e^{ip(z'' - D)} \sin[p(z' - D)] e^{ip(z'' - D)} dz'' \right], \end{aligned} \quad (\text{A12})$$

and the final result is

$$\begin{aligned} N_|| (z, z') = \frac{1}{4p^3} & \{ e^{-ip(3D - z_{>})} \sin[p(D - z_{<})] [e^{2ipD} [1 - 2ip(z_{>} - z_{<})] - e^{2ipz_{<}}] \\ & + e^{-ip(2D - z - z_{<})} [2p(z_{<} - D) + \sin[2p(D - z_{<})]] \}. \end{aligned} \quad (\text{A13})$$

We close this section with the $M_||$ function, which can be computed with the help of Eqs. (A3), (A11), and (A13). The result is

$$\begin{aligned} M_|| (z, z') = \frac{1}{32p^5} & \{ e^{-ip(4D - z_{>} + z_{<})} [e^{2ip(D + z_{>})} [2p(z_{>} - z_{<}) + 3i] + 2pe^{4ipD} (z_{>} - z_{<}) [2 - ip(z_{>} - z_{<})] \\ & + e^{-ip(4D - z_{>} + z_{<})} [e^{4ipz_{<}} (-4pD + 4pz_{<} + 3i) - e^{2ip(z_{>} + z_{<})} [2p(-2D + z_{>} + z_{<}) + 3i]] \\ & + e^{-ip(4D - z_{>} + z_{<})} [ie^{2ip(D + z_{<})} [-3 + 2p(z_{>} - z_{<})] [p(-4D + z_{>} + 3z_{<}) + 2i]] \\ & - 2e^{-ip(3D - 2z_{>} - z_{<})} \sin[p(z_{>} - D)] [e^{2ip(D - z_{<})} [-3 + 2ip(z_{>} - z_{<})] + 4ipD - 2ip(z_{>} + z_{<}) + 3] \\ & + 4pe^{-ip(2D - z_{>} - z_{<})} (D - z_{<}) [2ip(D - z_{<}) + \cos[2p(D - z_{<})] + 2] \\ & \left. + 2e^{-ip(2D - z_{>} - z_{<})} \sin[2p(D - z_{<})] [-3 - 2ip(D - z_{<})] \}. \end{aligned} \quad (\text{A14})$$

The boundary conditions $N_|| (D, z') = M_|| (D, z') = 0$ can be directly verified.

-
- [1] I. Mocioiu, M. Pospelov, and R. Roiban, *Phys. Lett. B* **489**, 390 (2000); S. M. Carroll, J. A. Harvey, V. A. Kostelecký, C. D. Lane, and T. Okamoto, *Phys. Rev. Lett.* **87**, 141601 (2001); C. E. Carlson, C. D. Carone, and R. F. Lebed, *Phys. Lett. B* **518**, 201 (2001); A. Anisimov, T. Banks, M. Dine, and M. Graesser, *Phys. Rev. D* **65**, 085032 (2002).
- [2] C. P. Burgess, J. M. Cline, E. Filotas, J. Matias, and G. D. Moore, *J. High Energy Phys.* **03** (2002) 043; A. R. Frey, *J. High Energy Phys.* **04** (2003) 012; J. M. Cline and L. Valcárcel, *J. High Energy Phys.* **03** (2004) 032.
- [3] C. Barcelo, S. Liberati, and M. Visser, *Living Rev. Relativ.* **8**, 12 (2005).
- [4] J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos, A. S. Sakharov, and E. K. Sarkisyan, *Astropart. Phys.* **25**, 402 (2006); G. Amelino-Camelia, *Phys. Lett. B* **510**, 255 (2001); A. Connes and D. Kreimer, *Commun. Math. Phys.* **199**, 203 (1998).
- [5] M. R. Douglas and N. A. Nekrasov, *Rev. Mod. Phys.* **73**, 977 (2001); L. F. Urrutia, *Lect. Notes Phys.* **702**, 299 (2006); P. Horava, *Phys. Rev. D* **79**, 084008 (2009); G. Amelino-Camelia, *Phys. Lett. B* **510**, 255 (2001); G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman, and L. Smolin, *Phys. Rev. D* **84**, 084010 (2011).
- [6] C. A. Escobar and L. F. Urrutia, *Phys. Rev. D* **92**, 025042 (2015); **92**, 025013 (2015); *Eur. Phys. Lett.* **106**, 31002 (2014); C. A. Escobar and M. A. G. Garcia, *Phys. Rev. D* **92**, 025034 (2015); Y. Bonder and C. A. Escobar, *Phys. Rev. D* **93**, 025020 (2016); M. Gomes, J. R. Nascimento, A. Yu. Petrov, and A. J. da Silva, *Phys. Rev. D* **81**, 045018 (2010); L. H. C. Borges, A. G. Dias, A. F. Ferrari, J. R. Nascimento, and A. Yu. Petrov, *Phys. Rev. D* **89**, 045005 (2014).

- [7] R. Gambini and J. Pullin, *Phys. Rev. D* **59**, 124021 (1999); J. Alfaro, H. A. Morales-Técolt, and L. F. Urrutia, *Phys. Rev. D* **65**, 103509 (2002); *Phys. Rev. Lett.* **84**, 2318 (2000); S. M. Carroll, J. A. Harvey, V. A. Kostelecký, C. D. Lane, and T. Okamoto, *Phys. Rev. Lett.* **87**, 141601 (2001).
- [8] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos, and S. Sarkar, *Nature (London)* **393**, 763 (1998); J. Magueijo and L. Smolin, *Phys. Rev. D* **67**, 044017 (2003); S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **81**, 043001 (2010); G. Cognola, R. Myrzakulov, L. Sebastiani, S. Vagnozzi, and S. Zerbini, *Classical Quantum Gravity* **33**, 225014 (2016).
- [9] V. A. Kostelecký and S. Samuel, *Phys. Rev. D* **39**, 683 (1989); V. A. Kostelecký and R. Potting, *Nucl. Phys.* **B359**, 545 (1991); K. Hashimoto and M. Murata, *Prog. Theor. Exp. Phys.* **2013**, 043B01 (2013).
- [10] F. R. Klinkhamer and C. Rupp, *Phys. Rev. D* **70**, 045020 (2004); A. Martín-Ruiz, *Phys. Rev. D* **90**, 125027 (2014); A. Martín-Ruiz *et al.*, *Rev. Mex. Fís.* **61**, 182 (2015); A. Martín-Ruiz, A. Frank, and L. F. Urrutia, *Phys. Rev. D* **92**, 045018 (2015).
- [11] D. Colladay and V. A. Kostelecký, *Phys. Rev. D* **55**, 6760 (1997).
- [12] V. A. Kostelecký, *Phys. Rev. D* **69**, 105009 (2004).
- [13] V. A. Kostelecký and S. Samuel, *Phys. Rev. D* **40**, 1886 (1989); **39**, 683 (1989); *Phys. Rev. Lett.* **63**, 224 (1989).
- [14] V. A. Kostelecký, C. Lane, and A. Pickering, *Phys. Rev. D* **65**, 056006 (2002).
- [15] V. A. Kostelecký and A. Pickering, *Phys. Rev. Lett.* **91**, 031801 (2003).
- [16] D. Colladay and P. McDonald, *Phys. Rev. D* **75**, 105002 (2007); **77**, 085006 (2008); **79**, 125019 (2009).
- [17] V. A. Kostelecký and R. Lehnert, *Phys. Rev. D* **63**, 065008 (2001).
- [18] V. A. Kostelecký and M. Mewes, *Phys. Rev. D* **66**, 056005 (2002).
- [19] D. Colladay and V. A. Kostelecký, *Phys. Lett. B* **511**, 209 (2001); M. Perez, *J. High Energy Phys.* **04** (2001) 32; V. A. Kostelecký, C. D. Lane, and A. G. Pickering, *Phys. Rev. D* **65**, 056006 (2002); M. Cambiaso, R. Lehnert, and R. Potting, *Phys. Rev. D* **90**, 065003 (2014); **85**, 085023 (2012); B. Altschul, *Phys. Rev. D* **75**, 105003 (2007); M. Schreck, *Phys. Rev. D* **86**, 065038 (2012); **89**, 085013 (2014); F. R. Klinkhamer and M. Schreck, *Nucl. Phys.* **B848**, 90 (2011); R. Casana, M. M. Ferreira Jr., A. R. Gomes, and P. R. Pinheiro, *Eur. Phys. J. C* **62**, 573 (2009); F. R. Klinkhamer and M. Risse, *Phys. Rev. D* **77**, 016002 (2008); **77**, 117901(A) (2008); A. Kobakhidze and B. H. J. McKellar, *Phys. Rev. D* **76**, 093004 (2007).
- [20] P. Wolf, S. Bize, A. Clairon, G. Santarelli, M. E. Tobar, and A. N. Luiten, *Phys. Rev. D* **70**, 051902(R) (2004); P. L. Stanwix, M. E. Tobar, P. Wolf, C. R. Locke, and E. N. Ivanov, *Phys. Rev. D* **74**, 081101(R) (2006); V. A. Kostelecký and M. Mewes, *Phys. Rev. Lett.* **87**, 251304 (2001).
- [21] D. Colladay, P. McDonald, and R. Potting, *Phys. Rev. D* **93**, 125007 (2016).
- [22] V. A. Kostelecký and J. D. Tasson, *Phys. Lett. B* **749**, 551 (2015).
- [23] R. Lehnert and R. Potting, *Phys. Rev. Lett.* **93**, 110402 (2004); *Phys. Rev. D* **70**, 125010 (2004); **70**, 129906 (2004); C. Kaufhold and F. R. Klinkhamer, *Phys. Rev. D* **76**, 025024 (2007).
- [24] S. M. Carroll, G. B. Field, and R. Jackiw, *Phys. Rev. D* **41**, 1231 (1990).
- [25] H. B. G. Casimir, *Proc. K. Ned. Akad. Wet.* **51**, 793 (1948).
- [26] G. Bressi, G. Carugno, R. Onofrio, and G. Ruoso, *Phys. Rev. Lett.* **88**, 041804 (2002).
- [27] J. Q. Quach, *Phys. Rev. Lett.* **114**, 081104 (2015).
- [28] R. C. Myers, *Phys. Rev. D* **60**, 046002 (1999).
- [29] O. Panella, *Phys. Rev. D* **76**, 045012 (2007); A. M. Frassinio and O. Panella, *Phys. Rev. D* **85**, 045030 (2012); K. Nouicer, *J. Phys. A* **38**, 10027 (2005); U. Harbacha and S. Hossenfelder, *Phys. Lett. B* **632**, 379 (2006).
- [30] K. Poppenhaeager, S. Hossenfelder, S. Hofmann, and M. Bleicher, *Phys. Lett. B* **582**, 1 (2004); E. Elizalde, S. D. Odintsov, and A. A. Saharian, *Phys. Rev. D* **79**, 065023 (2009); L. P. Teo, *Phys. Lett. B* **682**, 259 (2009).
- [31] A. Martín-Ruiz, M. Cambiaso, and L. F. Urrutia, *Eur. Phys. Lett. B* **113**, 60005 (2016); A. G. Grushin, P. Rodríguez-Lopez, and A. Cortijo, *Phys. Rev. B* **84**, 045119 (2011); A. G. Grushin and A. Cortijo, *Phys. Rev. Lett.* **106**, 020403 (2011); P. Rodríguez-Lopez and A. G. Grushin, *Phys. Rev. Lett.* **112**, 056804 (2014).
- [32] R. Zhao, J. Zhou, Th. Koschny, E. N. Economou, and C. M. Soukoulis, *Phys. Rev. Lett.* **103**, 103602 (2009); F. S. S. Rosa, D. A. R. Dalvit, and P. W. Milonni, *Phys. Rev. A* **78**, 032117 (2008).
- [33] A. Martín-Ruiz and C. A. Escobar, *Phys. Rev. D* **94**, 076010 (2016).
- [34] M. Frank and I. Turan, *Phys. Rev. D* **74**, 033016 (2006).
- [35] O. G. Kharlanov and V. Ch. Zhukovsky, *Phys. Rev. D* **81**, 025015 (2010).
- [36] V. A. Kostelecký and N. Russell, *Rev. Mod. Phys.* **83**, 11–31 (2011).
- [37] A. Martín-Ruiz, M. Cambiaso, and L. F. Urrutia, *Phys. Rev. D* **92**, 125015 (2015); **93**, 045022 (2016); **94**, 085019 (2016).
- [38] X. L. Qi and S. C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011); M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [39] C. Adam and F. R. Klinkhamer, *Nucl. Phys. B* **607**, 247 (2001); J. Alfaro, M. Cambiaso, P. Giacconi, and R. Soldati, *Int. J. Mod. Phys. A* **25**, 3271 (2010).
- [40] R. Casana, M. M. Ferreira Jr., and C. E. H. Santos, *Phys. Rev. D* **78**, 025030 (2008); H. Belich Jr., M. M. Ferreira Jr., J. A. Helayël-Neto, and M. T. D. Orlando, *Phys. Rev. D* **68**, 025005 (2003); H. Belich Jr., T. Costa-Soares, M. M. Ferreira Jr., and J. A. Helayël-Neto, *Eur. Phys. J. C* **42**, 127 (2005).
- [41] K. A. Milton, *The Casimir Effect: Physical Manifestation of Zero-Point Energy* (World Scientific, Singapore, 2001); M. Bordag, G. L. Klimchitskaya, U. Mohideen, and V. M. Mostepanenko, *Advances in Casimir Effect* (Oxford University Press, England, 2009).
- [42] L. S. Brown and G. J. Maclay, *Phys. Rev.* **184**, 1272 (1969).
- [43] D. Deutsch and P. Candelas, *Phys. Rev. D* **20**, 3063 (1979).
- [44] V. Sopova and L. H. Ford, *Phys. Rev. D* **66**, 045026 (2002); **72**, 033001 (2005).