Continuous body dynamics and the Mathisson-Papapetrou-Dixon equations

S. P. Loomis^{*} and J. David Brown[†]

Department of Physics, North Carolina State University, Raleigh, North Carolina 27695, USA (Received 6 January 2017; published 17 February 2017)

We show that an effective particle Lagrangian yields the Mathisson-Papapetrou-Dixon (MPD) equations. The spin of the effective particle is defined without any reference to a fixed body frame or angular velocity variable. We then demonstrate that a continuous body, defined by a congruence of world lines and described by a general action, can be rewritten as an effective particle. We analyze the gauge freedom of the body and show that a natural center of mass condition is related to a spin supplementary condition.

DOI: 10.1103/PhysRevD.95.044025

I. INTRODUCTION

The motion of extended bodies in general relativity was first addressed by Mathisson in 1937 [1]. Mathisson defined multipole moments for the stress-energy tensor expanded about a central world line and formulated the conservation of stress-energy as a variational principle. He derived what we now know as the Mathisson-Papapetrou-Dixon (MPD) equations to "pole–dipole" order, and identified the quadrupole terms that had a nonrelativistic analogue. Papapetrou, using a different definition for multipole moments, derived the same equations in 1951 [2].

The analyses of Mathisson and Papapetrou yield ten equations for thirteen unknowns. The ten equations give the time evolution of the four components of momentum, P^{α} , and the six components of spin, $S^{\alpha\beta}$. The thirteen unknowns are the momentum, spin, and the three degrees of freedom contained in the particle's proper velocity U^{α} .

Mathisson and later Pirani [3] addressed the mismatch between the number of equations and number of unknowns by introducing the spin supplementary condition $U^{\alpha}S_{\alpha\beta} = 0$. Pirani justified this choice by analogy to a similar identity for the center of mass in special relativity. Papapetrou instead employed the spin supplementary condition $V^{\alpha}S_{\alpha\beta} = 0$ where V^{α} is an arbitrary time flow vector field. In 1959 Tulczyjew simplified Mathisson's multipole formulation, again deriving the same equations of motion at pole-dipole order but choosing the spin supplementary condition $P^{\alpha}S_{\alpha\beta} = 0$, arguing that Mathisson and Pirani's condition did not uniquely determine the world line. A survey of the various spin supplementary conditions and how they relate to one another can be found in [4]. A concrete analysis of the relationship between spin supplementary conditions and center of mass can be found in [5].

In a series of papers from 1970 to 1974, Dixon presented yet another reformulation of the multipole moments in

terms of a Fourier transformation of the stress-energy tensor [6–8]. The complete argument is also given in [9]. Dixon found that Mathisson's variational principle yields dynamical equations for P_{α} and $S_{\alpha\beta}$, but leaves the dynamical evolution of the quadrupole and higher order multipole moments undefined. His analysis places restrictions on the symmetries of these multipoles. Based on these symmetries, Dixon defines the reduced multipole moments $J^{\mu\nu\rho\sigma\alpha_1\cdots\alpha_n}$ for $n \ge 0$.

The final form of the MPD equations through quadrupole order, as given by Dixon, is [9]:

$$\frac{\mathrm{D}P_{\alpha}}{\mathrm{D}s} = -\frac{1}{2}R_{\alpha\beta\mu\nu}\dot{X}^{\beta}S^{\mu\nu} - \frac{1}{6}\nabla_{\alpha}R_{\mu\nu\rho\sigma}J^{\mu\nu\rho\sigma},\qquad(1.1\mathrm{a})$$

$$\frac{\mathrm{D}S_{\alpha\beta}}{\mathrm{D}s} = 2P_{[\alpha}\dot{X}_{\beta]} + \frac{4}{3}R_{\mu\nu\lambda[\alpha}J_{\beta]}{}^{\lambda\nu\mu}.$$
(1.1b)

Here, the world line is expressed as $x^{\alpha} = X^{\alpha}(s)$, where x^{α} are the spacetime coordinates and X^{α} are functions of a world line parameter *s*. The dot above a symbol denotes the time derivative d/ds and D/Ds is the covariant derivative along the world line; for example, $DP_{\alpha}/Ds = \dot{P}_{\alpha} - \Gamma^{\gamma}_{\alpha\beta}\dot{X}^{\beta}P_{\gamma}$. This will also be denoted with a circle above the symbol, so that $\mathring{P}_{\alpha} = DP_{\alpha}/Ds$.

Dixon defines the momentum and spin in terms of integrals over the leaves of a foliation $\Sigma(s)$ of the world tube. The integrals, which involve the stress-energy tensor $T^{\mu'\nu'}$, Synge's world function σ and the Jacobi propagators $H^{\mu'}{}_{\alpha}$ and $K^{\mu'}{}_{\alpha}$ (described in Appendix B) are [6]

$$P_{\alpha}(s) \equiv \int_{\Sigma(s)} \mathrm{d}\Sigma_{\nu'} K^{\mu'}{}_{\alpha} T^{\nu'}_{\mu'}, \qquad (1.2a)$$

$$S_{\alpha\beta}(s) \equiv 2 \int_{\Sigma(s)} \mathrm{d}\Sigma_{\nu'} H^{\mu'}{}_{[\alpha}\sigma_{\beta]} T^{\nu'}_{\mu'}. \qquad (1.2b)$$

Both $H^{\mu'}{}_{\alpha}$ and $K^{\mu'}{}_{\alpha}$ are bitensors with one primed index "located" at the point of integration $x^{\mu'} \in \Sigma(s)$ and one unprimed index located at the world line $X^{\alpha}(s)$. These

sloomis@ucdavis.edu

Present Address: Department of Physics, University of California, Davis, California 95616.

[†]david_brown@ncsu.edu

bitensors act to transfer vectors from one tangent space into the other. The derivative of Synge's world function $\sigma_{\beta} = \nabla_{\beta}\sigma$ points opposite to the vector tangent to the unique geodesic connecting $X^{\alpha}(s)$ and $x^{\mu'}$ and acts as a position vector. The corresponding integrals for the multipole moments are complicated and beyond the scope of this introduction.

The multipole method of Mathisson, Papapetrou, Tulzcyjew and Dixon is not the only approach that yields the MPD equations. Bailey and Israel [10], extending the work of Hanson and Regge [11], showed that a form of the MPD equations could be derived from any reparametrization–invariant Lagrangian involving a world line $X^{\alpha}(s)$, a set of Lorentz–orthonormal basis vectors $e_a^{\alpha}(s)$ transported along the world line, and a set of external tensor fields that interact with the body. The basis vectors $e_a^{\alpha}(s)$ define the orientation of the body, although the exact relationship for a physical (i.e. nonrigid) body is not addressed. By analogy with rigid body motion in classical mechanics, one says that the index *a* labels the legs of a "body-fixed frame."

These results were independently replicated without the external fields by Porto [12]. More recently, Steinhoff [13] has reformulated Bailey and Israel's full result in newer notation. We give a brief overview of Steinhoff's presentation below.

Steinhoff begins by considering an action of the form

$$S[X, e, \Phi_I] = \int ds L(g_{\alpha\beta}(X), \Phi_A(X), \dot{X}^{\alpha}(s), e_a^{\ \alpha}(s),$$
$$\Omega^{\alpha\beta}(s), \Phi_I(s)), \qquad (1.3)$$

where the integration is along the world line $X^{\alpha}(s)$. Here $g_{\alpha\beta}(x)$ is the metric of the spacetime manifold \mathcal{M} , $\Phi_A(x)$ are fields on this manifold and $\Phi_I(s)$ are scalar dynamical variables defined along the world line. Note that the Φ_I are functions of *s* only, and the index *I* can include a body frame index *a*. The angular velocity is defined in terms of the basis vectors $e_a{}^{\alpha}(s)$ by $\Omega^{\alpha\beta}(s) \equiv \eta^{ab} e_a{}^{\alpha} D(e_b{}^{\beta})/Ds$, where η^{ab} is the Minkowski metric.

Steinhoff defines the momentum and spin as

$$P_{\alpha} \equiv \frac{\partial L}{\partial \dot{X}^{\alpha}}, \qquad (1.4a)$$

$$S_{\alpha\beta} \equiv 2 \frac{\partial L}{\partial \Omega^{\alpha\beta}}.$$
 (1.4b)

With these definitions, variation of the action (1.3) with respect to $X^{\alpha}(s)$ and $e_a{}^{\alpha}(s)$ yields the equations of motion

$$\frac{\mathrm{D}P_{\alpha}}{\mathrm{D}s} = -\frac{1}{2}R_{\alpha\beta\mu\nu}\dot{X}^{\beta}S^{\mu\nu} + (\nabla_{\alpha}\Phi_{A})\frac{\partial L}{\partial\Phi_{A}},\qquad(1.5a)$$

$$\frac{\mathrm{D}S_{\alpha\beta}}{\mathrm{D}s} = 2P_{[\alpha}\dot{X}_{\beta]} - 2(\mathrm{G}_{[\alpha\beta]}\Phi_A)\frac{\partial L}{\partial\Phi_A}.$$
(1.5b)

Here, $G_{\alpha\beta}$ is a linear operator defined such that $\nabla_{\alpha}\Phi_{A} = \partial_{\alpha}\Phi_{A} + \Gamma^{\gamma}_{\alpha\beta}G^{\beta}_{\ \gamma}\Phi_{A}$. The action of $G^{\alpha}_{\ \beta}$ depends on the tensor type of Φ_{A} .

In Sec. II we consider a general action that depends only on a world line $x^{\alpha} = X^{\alpha}(s)$, a set of tensors $\psi_{I}(s)$ defined along the world line, and a set of external fields $\phi_A(x)$. We call this the "effective particle" model, and show that it yields the MPD equations.¹ Our action does not depend on an orthonormal body frame $e_a{}^{\alpha}$ or angular velocity $\Omega^{\alpha\beta}$. We show in Sec. III that continuous bodies (defined as a congruence of world lines minimizing a particular action) can be expressed in terms of the effective particle action. The definitions for momentum and spin that emerge from this analysis coincide with the definitions given by Dixon [6]. In Sec. IV we discuss the gauge constraints that can be placed on the effective particle, and explore the relation between spin supplementary condition and center of mass. Finally, in Sec. V we apply our results to analyze a continuous body of noninteracting particles-a "dust cloud." As an effective particle, the dust cloud satisfies the geodesic deviation equations.

II. THE MPD EQUATIONS FOR AN EFFECTIVE PARTICLE

In this section we generalize the method for deriving the MPD equations used by Bailey and Israel [10].

The system consists of an effective particle in a manifold \mathcal{M} with position $x^{\alpha} = X^{\alpha}(s)$ and a collection of tensors $\psi_I(s)$. The tensors ψ_I take the place of the orthonormal basis $e_a{}^{\alpha}$ and body-frame variables ϕ_I used in Eq. (1.3). The index *I* can denote tensor indices as well as functional dependence. The evolution of the effective particle system is described by the action

$$S[X, \psi_I] = \int_{s_0}^{s_1} \mathrm{d}s L(\phi_A(X), \dot{X}^{\alpha}(s), \psi_I(s), \ddot{\psi}_I(s)).$$
(2.1)

The $\phi_A(x)$ are any collection of spacetime fields. For example, ϕ_A can include the electromagnetic field and its derivatives, the metric tensor $g_{\alpha\beta}$, the curvature tensor $R_{\alpha\beta\gamma\delta}$ and its symmetrized derivatives $\nabla_{(\alpha} \cdots \nabla_{\gamma)} R_{\mu\nu\rho\sigma}$. In this paper we treat these fields as external sources—they are not varied in the variational principle.

Recall that the notation ψ_I is an abbreviation for the covariant derivative along the world line,

$$\mathring{\psi}_I \equiv \frac{\mathbf{D}\psi_I}{\mathbf{D}s} = \frac{\mathbf{d}\psi_I}{\mathbf{d}s} + \Gamma^{\nu}_{\alpha\mu} \dot{X}^{\alpha} \mathbf{G}^{\mu}_{\ \nu} \psi_I. \tag{2.2}$$

The operator G^{μ}_{ν} , discussed more fully in Appendix A, acts on tensor indices [15]. For example, we have

¹After obtaining this result, we became aware of a similar result by H. Fuchs [14].

CONTINUOUS BODY DYNAMICS AND THE MATHISSON- ...

$$G^{\mu}{}_{\nu}\psi^{\rho} = \delta^{\rho}_{\nu}\psi^{\mu}, \qquad (2.3a)$$

$$G^{\mu}{}_{\nu}\psi_{\rho} = -\delta^{\mu}_{\rho}\psi_{\nu}, \qquad (2.3b)$$

for contravariant and covariant vectors. The extension of $G^{\mu}{}_{\nu}$ to higher rank tensors is straightforward. Also note that the covariant derivative of X^{α} is defined by the vector $DX^{\alpha}/Ds \equiv \dot{X}^{\alpha}$. Thus, the world line coordinates behave as spacetime scalars under covariant differentiation.

The Lagrangian (2.1) is a function over a tensor bundle. That is, *L* depends on the position X(s) as well as tensors in the tangent space of x = X(s). The variation $\delta \psi_I$ is not covariant whenever the base point *x* is also varied. As a result, the functional derivatives $\delta S/\delta X^{\alpha}$ and $\delta S/\delta \psi_I$ yield the equations of motion in noncovariant combinations. Here we use the results of Appendix A to vary the action in a covariant manner.

To begin, let us define the momentum variables

$$P_{\alpha} \equiv \frac{\partial L}{\partial \dot{X}^{\alpha}}, \qquad (2.4a)$$

$$\pi^{I} \equiv \frac{\partial L}{\partial \dot{\psi}_{I}}.$$
 (2.4b)

Using the result (A15a) from Appendix A, the covariant variation of the action is

$$\delta S = \int_{s_0}^{s_1} \mathrm{d}s \left(P_{\alpha} \Delta \dot{X}^{\alpha} + \frac{\partial L}{\partial \phi_A} \nabla_{\alpha} \phi_A \delta X^{\alpha} + \pi^I \Delta \ddot{\psi}_I + \frac{\partial L}{\partial \psi_I} \Delta \psi_I \right).$$
(2.5)

Here, Δ is the covariant variation defined by [13]

$$\Delta \equiv \delta + \Gamma^{\nu}_{\alpha\mu} \delta X^{\alpha} \mathcal{G}^{\mu}{}_{\nu}. \tag{2.6}$$

Because ϕ_A are external sources, their variations are given by $\delta\phi_A = (\partial\phi_A/\partial x^{\alpha})\delta X^{\alpha}$. Then the covariant variations of these fields are $\Delta\phi_A = \nabla_{\alpha}\phi_A\delta X^{\alpha}$. Also observe that the covariant variation of X^{α} is defined by the vector $\Delta X^{\alpha} \equiv \delta x^{\alpha}$. That is, the world line coordinates behave as spacetime scalars under covariant variation.

To bring δS into a form that will provide the equations of motion, we must swap the order of the variations and time derivatives in $\Delta \dot{X}^{\alpha}$ and $\Delta \dot{\psi}^{\alpha}$. The covariant variation and covariant derivative do not commute; in general, we have the following relation from Eq. (A21):

$$\left[\Delta, \frac{\mathrm{D}}{\mathrm{D}s}\right] = R^{\sigma}{}_{\rho\mu\nu} \delta X^{\mu} \dot{X}^{\nu} \mathrm{G}^{\rho}{}_{\sigma}.$$
(2.7)

This yields the results

PHYSICAL REVIEW D 95, 044025 (2017)

$$\Delta \dot{X}^{\alpha} = \frac{\mathrm{D}}{\mathrm{D}s} (\delta X^{\alpha}), \qquad (2.8a)$$

$$\Delta \mathring{\psi}_{I} = \frac{\mathrm{D}}{\mathrm{D}s} (\Delta \psi_{I}) + R^{\sigma}{}_{\rho\mu\nu} \delta X^{\mu} \dot{X}^{\nu} \mathrm{G}^{\rho}{}_{\sigma} \psi_{I}, \qquad (2.8\mathrm{b})$$

since the world line coordinates X^{α} behave as scalar fields under covariant differentiation and covariant variation.

Now integrate by parts. The endpoint terms vanish if we assume that X^{α} and ψ_I are fixed at s_0 and s_1 . The variation of the action becomes

$$\delta S = \int_{s_0}^{s_1} \mathrm{d}s \left[\left(-\overset{\circ}{P}_{\alpha} + R^{\sigma}{}_{\rho\alpha\beta} \dot{X}^{\beta} \pi^I \mathrm{G}^{\rho}{}_{\sigma} \psi_I + \nabla_{\alpha} \phi_A \frac{\partial L}{\partial \phi_A} \right) \delta X^{\alpha} + \left(\frac{\partial L}{\partial \psi_I} - \overset{\circ}{\pi}^I \right) \Delta \psi_I \right]. \quad (2.9)$$

The coefficients of δX^{α} and $\Delta \psi_I$ give the equations of motion

$$\overset{\circ}{P}_{\alpha} = -R_{\alpha\beta}{}^{\rho\sigma}\dot{X}^{\beta}\pi^{I}G_{\rho\sigma}\psi_{I} + \nabla_{\alpha}\phi_{A}\frac{\partial L}{\partial\phi_{A}},\qquad(2.10a)$$

$$\overset{\circ}{\pi}^{I} = \frac{\partial L}{\partial \psi_{I}}.$$
(2.10b)

Equation (2.6) shows that the variation $\Delta \psi_I$ is a sum of noncovariant terms proportional to $\delta \psi_I$ and δX^{α} . From this we see that the equation of motion $\delta S / \delta \psi_I = 0$ (with X^{α} fixed) coincides with (2.10b), while the equation of motion $\delta S / \delta X^{\alpha} = 0$ (with ψ_I fixed) is a noncovariant combination of the covariant Eqs. (2.10)

Spin is defined as

$$S_{\rho\sigma} \equiv 2\pi^I \mathcal{G}_{[\rho\sigma]} \psi_I. \tag{2.11}$$

This puts the equation of motion (2.10a) into the form of the MPD equation (1.5a). The equation of motion for the spin variable itself is obtained from the covariant derivative of the definition (2.11), which yields $\mathring{S}_{\rho\sigma} = 2\mathring{\pi}^I G_{[\rho\sigma]} \psi_I + 2\pi^I G_{[\rho\sigma]} \mathring{\psi}_I$. With the equation of motion (2.10b) this becomes

$$\overset{\circ}{S}_{\rho\sigma} = 2 \frac{\partial L}{\partial \psi_I} \mathbf{G}_{[\rho\sigma]} \psi_I + 2\pi^I \mathbf{G}_{[\rho\sigma]} \overset{\circ}{\psi}_I.$$
(2.12)

We now make use of (A15b), which follows from the requirement of general covariance. This equation tells us that the operator $G^{\alpha}{}_{\beta}$ acting on the Lagrangian *L* of (2.1) follows a "chain rule". Since *L* is a scalar, we have

S. P. LOOMIS and J. DAVID BROWN

$$0 = \frac{\partial L}{\partial \phi_A} \mathbf{G}^{\alpha}{}_{\beta} \phi_A + \frac{\partial L}{\partial \dot{X}^{\mu}} \mathbf{G}^{\alpha}{}_{\beta} \dot{X}^{\mu} + \frac{\partial L}{\partial \psi_I} \mathbf{G}^{\alpha}{}_{\beta} \psi_I + \frac{\partial L}{\partial \dot{\psi}_I} \mathbf{G}^{\alpha}{}_{\beta} \dot{\psi}_I.$$
(2.13)

We can lower the index α on $G^{\alpha}{}_{\beta}$ and antisymmetrize. Using the notation (2.4) for the momenta, we have

$$0 = \frac{\partial L}{\partial \phi_A} \mathbf{G}_{[\alpha\beta]} \phi_A + P_{[\beta} \dot{X}_{\alpha]} + \frac{\partial L}{\partial \psi_I} \mathbf{G}_{[\alpha\beta]} \psi_I + \pi^I \mathbf{G}_{[\alpha\beta]} \ddot{\psi}_I.$$
(2.14)

This result can be used to rewrite the time derivative of the spin from (2.12) as the MPD equation (1.5b).

To summarize, we have shown that the equations of motion (2.10) that follow from the action (2.1), along with the definition (2.11) for spin, yield the MPD equations

$$\overset{\circ}{P}_{\alpha} = -\frac{1}{2} R_{\alpha\beta\mu\nu} \dot{X}^{\beta} S^{\mu\nu} + (\nabla_{\alpha}\phi_{A}) \frac{\partial L}{\partial \phi_{A}}, \qquad (2.15a)$$

$$\mathring{S}_{\alpha\beta} = 2P_{[\alpha}\dot{X}_{\beta]} - 2(\mathbf{G}_{[\alpha\beta]}\phi_A)\frac{\partial L}{\partial\phi_A}$$
(2.15b)

in the form (1.5).

The key difference between this result and previous analyses is that we do not require a body frame or basis vectors e_a^{α} to define the orientation of the body, and we have no need for an angular velocity variable $\Omega_{\alpha\beta}$. The spin, as defined in (2.11), does not rely on these constructions.

III. CONTINUOUS BODIES WRITTEN AS EFFECTIVE PARTICLES

Let \mathscr{M} be a Riemannian manifold with coordinates x^{μ} . We consider a continuous body in the sense defined by Carter and Quintana [16], as a congruence of world lines represented by the smooth mapping $X: \mathbb{R} \times \mathscr{S} \to \mathscr{M}$, where \mathscr{S} is a differentiable manifold whose points represent the world lines, and the real numbers \mathbb{R} label points along each world line. \mathscr{S} is called the "matter space" and is given coordinates ζ^i [17]. The coordinate on \mathbb{R} is *s*. Thus, functions over the congruence may be written in terms of coordinates (s, ζ^i) on $\mathbb{R} \times \mathscr{S}$, or in terms of the manifold coordinates x^{μ} . We introduce the inverse mapping $(Z^0(x), Z^i(x))$ that satisfies $s = Z^0(X(s, \zeta))$ and $\zeta^i = Z^i(X(s, \zeta))$.

The derivatives of X^{μ} and (Z^0, Z^i) satisfy [18]

$$\dot{X}^{\mu}Z^{0}_{,\nu} + X^{\mu}_{,i}Z^{i}_{,\nu} = \delta^{\mu}_{\nu}, \qquad (3.1a)$$

$$X^{\mu}_{,i}Z^{j}_{,\mu} = \delta^{j}_{i}, \qquad (3.1b)$$

$$\dot{X}^{\mu}Z^{0}_{,\mu} = 1,$$
 (3.1c)

$$X^{\mu}_{,i}Z^{0}_{,\mu} = 0, \qquad (3.1d)$$

$$\dot{X}^{\mu}Z^{i}_{,\mu} = 0,$$
 (3.1e)

where $\dot{X}^{\mu} = \partial X^{\mu} / \partial s$.

We introduce the notations

$$\frac{\mathcal{D}}{\mathcal{D}s} \equiv \dot{X}^{\mu} \nabla_{\mu} = \frac{\partial}{\partial s} + \Gamma^{\sigma}_{\mu\rho} \dot{X}^{\mu} G^{\rho}{}_{\sigma}, \qquad (3.2a)$$

$$\frac{\mathcal{D}}{\mathcal{D}\zeta^{i}} \equiv X^{\mu}_{,i} \nabla_{\mu} = \frac{\partial}{\partial \zeta^{i}} + \Gamma^{\sigma}_{\mu\rho} X^{\mu}_{,i} \mathbf{G}^{\rho}{}_{\sigma}, \qquad (3.2b)$$

using the generators G^{ρ}_{σ} from Appendix A, and following Vines' use of \mathcal{D} as a covariant partial derivative [19]. As before, we may also abbreviate $\mathcal{D}/\mathcal{D}s$ with the circle

(e.g. \tilde{A}) and $\mathcal{D}/\mathcal{D}\zeta^i$ with semicolons (e.g. $A_{;i}$).

We consider a fairly general action assuming that the body is subject to no external forces, though it may be subject to internal forces mediated by the first spatial derivative X_{j}^{μ} :

$$S[X] = \int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta \mathcal{L}(\zeta, g_{\mu\nu}(X), \dot{X}^{\mu}, X^{\mu}_{,i}).$$
(3.3)

We will generally restrict this action to be reparametrization invariant, although the results of this section do not depend on that assumption. When reparametrization invariance is enforced, the above action coincides with DeWitt's elastic body [20,21]

$$S[X] = -\int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta \alpha \rho(\zeta, f_{ij}), \qquad (3.4)$$

where $\alpha = \sqrt{-\dot{X}^{\mu}\dot{X}_{\mu}}$ and f_{ij} is the fleet metric $f_{ij} = (g_{\mu\nu} + U_{\mu}U_{\nu})X^{\mu}_{,i}X^{\nu}_{,j}$, with $U^{\mu} = \dot{X}^{\mu}/\alpha$ being the four-velocity field.

In this section we study some basic properties of the action (3.3) and show that it can be expressed in the form of the effective particle action (2.1). Doing so gives definitions of P_{α} and $S_{\alpha\beta}$ identical to Dixon's definitions (1.2) [6].

We start by varying the action to determine the equations of motion. With the exception of its dependence on ζ (which does not affect the variation), the Lagrangian density \mathcal{L} is a function of X through the metric field $g_{\mu\nu}$ and is also a function of vectors \dot{X}^{μ} and $X_{,i}^{\mu}$. This means we can use the methods of Appendix A, specifically Eq. (A15a), to vary the action. We define the canonical momentum density and stress density as

$$\mathcal{P}_{\mu} \equiv \frac{\partial \mathcal{L}}{\partial \dot{X}^{\mu}}, \qquad (3.5a)$$

$$S^{i}_{\mu} \equiv \frac{\partial \mathcal{L}}{\partial X^{\mu}_{\ i}}.$$
 (3.5b)

Then the variation with respect to X^{μ} is

$$\delta S = \int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta [\mathcal{P}_{\mu} \Delta \dot{X}^{\mu} + \mathcal{S}^i_{\mu} \Delta X^{\mu}_{,i}]. \quad (3.6)$$

Note that, since the metric is treated as an external field, its covariant variation $\Delta g_{\mu\nu}(X) = \nabla_{\alpha} g_{\mu\nu} \delta X^{\alpha}$ vanishes.

For the $\Delta \dot{X}^{\mu}$ term in Eq. (3.6), integration by parts gives

$$\int_{s_0}^{s_1} \mathrm{d}s \mathcal{P}_{\mu} \Delta \dot{X}^{\mu} = -\int_{s_0}^{s_1} \mathrm{d}s \frac{\mathcal{D} \mathcal{P}_{\mu}}{\mathcal{D}s} \delta X^{\mu} + [\mathcal{P}_{\mu} \delta X^{\mu}]_{s_0}^{s_1}.$$
 (3.7)

The last term in the above equation vanishes when we fix the initial and final configurations $X^{\mu}(s_0)$ and $X^{\mu}(s_1)$; that is, we set $\delta X^{\mu}(s_0) = \delta X^{\mu}(s_1) = 0$. We can similarly integrate by parts for the $X^{\mu}_{,i}$ term, with the result

$$\int_{\mathscr{S}} \mathrm{d}^{3} \zeta \mathcal{S}^{i}_{\mu} \Delta X^{\mu}_{,i} = -\int_{\mathcal{S}} \mathrm{d}^{3} \zeta \frac{\mathcal{D} \mathcal{S}^{i}_{\mu}}{\mathcal{D} \zeta^{i}} \delta X^{\mu} + \int_{\partial \mathscr{S}} \mathrm{d}^{2} \zeta \eta_{i} \mathcal{S}^{i}_{\mu} \delta X^{\mu}.$$
(3.8)

where η_i is the outward-pointing vector field normal to the boundary surface $\partial \mathscr{S}$. The variation of the action with X^{μ} fixed at s_0 and s_1 is then

$$\delta S = \int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta [-\overset{\circ}{\mathcal{P}}_{\mu} - \mathcal{S}^i_{\mu;i}] \delta X^{\mu} + \int_{s_0}^{s_1} \mathrm{d}s \int_{\partial \mathscr{S}} \mathrm{d}^2 \zeta [\eta_i \mathcal{S}^i_{\mu}] \delta X^{\mu}.$$
(3.9)

Since δX^{μ} is arbitrary, this gives the equations of motion and boundary conditions

$$-\overset{\circ}{\mathcal{P}}_{\mu}-\mathcal{S}^{i}_{\mu;i}=0\quad(\zeta\in\mathscr{S}), \tag{3.10a}$$

$$\eta_i \mathcal{S}^i_\mu = 0 \quad (\zeta \in \partial \mathscr{S}). \tag{3.10b}$$

By integrating (3.10a) over the matter space \mathscr{S} , and using the boundary (3.10b), we obtain a continuity equation for the total momentum within the body.

The stress-energy tensor is defined by

$$T^{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\partial \mathcal{L}}{\partial g_{\mu\nu}}.$$
 (3.11)

This can be rewritten in terms of the canonical momentum and stress densities as

$$T^{\mu\nu} = \frac{1}{\sqrt{-g}} \left(\mathcal{P}^{\mu} \dot{X}^{\nu} + \mathcal{S}^{\mu i} X^{\nu}_{,i} \right).$$
(3.12)

by using the chain rule (A15b) for $G^{\alpha}{}_{\beta}$. From Eq. (3.1) we see that $\mathcal{P}^{\mu} = -\sqrt{h}T^{\mu\nu}n_{\nu}$ where *h* is the determinant of the induced metric $h_{ij} = g_{\mu\nu}X^{\mu}_{,i}X^{\nu}_{,j}$ on surfaces $\Sigma(s)$ of constant *s*. Also, n_{μ} is the future-pointing unit vector field normal to $\Sigma(s)$.

We now turn to the primary goal of this section: to show that the continuous body action (3.3) can be written in the form of the effective particle action (2.1). This requires mapping the information about the congruence of geodesics to the tangent space of a point on a fiducial world line. To prepare for this analysis, we first make a change of notation by placing primes on coordinates and tensors associated with the continuous body. In particular, we will use $X' : \mathbb{R} \times \mathscr{S} \to \mathscr{M}$ rather than X to denote the congruence of world lines. For tensors associated with the body, we place the prime on the indices: for example, the (unnormalized) velocity, momentum density and stress density are $\dot{X}^{\mu'}$, $\mathcal{P}_{\mu'}$ and $\mathcal{S}^{i}_{\mu'}$, respectively.

The fiducial world line will be denoted $X^{\mu}(s)$. The fiducual worldline does not need to lie within the material body. If it does, the spacetime point $X^{\mu}(s)$ (for a given value of *s*) does not need to lie on the *s* = const surface.

The introduction of the fiducual worldline adds a gauge freedom to the system, in additional to the reparametrization invariance that is already present. In later sections we will discuss how the gauge freedom can be fixed.

We make use of the bitensor and exponential map formalism of Appendix B. We begin by defining the exponential map from a point $X^{\mu}(s)$ on the fiducial world line and a vector ξ^{μ} in the tangent space at $X^{\mu}(s)$, to the point $X^{\mu'}(s, \zeta)$ in the continuous body:

$$X'(s,\zeta) = \exp(X(s),\xi(s,\zeta)). \tag{3.13}$$

Here, $\xi^{\mu}(s,\zeta) = -\sigma^{\mu}(X(s), X'(s,\zeta))$ is the vector at X(s) tangent to the geodesic $\gamma(u)$ connecting X(s) and $X'(s,\zeta)$, affinely parametrized such that $\gamma(0) = X$ and $\gamma(1) = X'$. Note that the point $X'(s,\zeta)$ defined by the exponential map depends on the geometry in a neighborhood of X(s). To ensure that there is a unique geodesic connecting X(s) and $X'(s,\zeta)$, the size of the body must be restricted by the curvature and its derivatives at X(s). The region to which it is restricted is termed the "normal convex neighborhood" (NCN).

With the placement of primes on the coordinates and tensors associated with points in the continuous body, the action (3.3) becomes

$$S[X'] = \int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta \mathcal{L}(\zeta, g_{\mu'\nu'}, \dot{X}^{\mu'}, X_{,i}^{\mu'}).$$
(3.14)

Equation (3.13) defines a change of variables in the action from $X^{\mu'}(s,\zeta)$ to $X^{\mu}(s)$ and $\xi^{\mu}(s,\zeta)$. We can use Eq. (A7) to

write $\delta X^{\mu'}$ in terms of δX^{μ} and the covariant variation $\Delta \xi^{\mu}$. The result,

$$\delta X^{\mu'} = K^{\mu'}{}_{\alpha} \delta X^{\alpha} + H^{\mu'}{}_{\alpha} \Delta \xi^{\alpha}, \qquad (3.15)$$

makes use of the Jacobi propagators, $K^{\mu'}{}_{\alpha}$ and $H^{\mu'}{}_{\alpha}$. The Jacobi propagators are defined in terms of derivatives of the exponential map. They depend on $X^{\alpha}(s)$, $\xi^{\alpha}(s,\zeta)$ and the geometry in a neighborhood of x = X(s). They are introduced in Appendix A and further developed in Appendix B. Note that $H^{\mu'}{}_{\alpha}$ is invertible in the NCN, since there must be a one-to-one mapping between ξ^{α} and $X^{\mu'}$.

As a functional of $X^{\mu}(s)$ and $\xi^{\mu}(s, \zeta)$, the variation of the action is

$$\delta S = \int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta [-\mathcal{P}_{\mu'} - \mathcal{S}^i_{\mu';i}] (K^{\mu'}{}_{\alpha} \delta X^{\alpha} + H^{\mu'}{}_{\alpha} \Delta \xi^{\alpha}) + \int_{s_0}^{s_1} \mathrm{d}s \int_{\partial \mathscr{S}} \mathrm{d}^2 \zeta [\eta_i \mathcal{S}^i_{\mu'}] (K^{\mu'}{}_{\alpha} \delta X^{\alpha} + H^{\mu'}{}_{\alpha} \Delta \xi^{\alpha}),$$
(3.16)

where we have used the results from Eq. (3.9). Extremization of *S* with respect to ξ^{α} yields the equations of motion

$$[-\overset{\circ}{\mathcal{P}}_{\mu'} - \mathcal{S}^{i}_{\mu';i}]H^{\mu'}{}_{\alpha} = 0 \quad (\zeta \in \mathscr{S}),$$
(3.17a)

$$[\eta_i \mathcal{S}^i_{\mu'}] H^{\mu'}{}_{\alpha} = 0 \quad (\zeta \in \partial \mathscr{S}). \tag{3.17b}$$

The propagator $H^{\mu'}{}_{\alpha}$ is invertible in the NCN, so these equations are clearly equivalent to the equations of motion obtained by extremizing the action with respect to $X^{\mu'}$ [Eqs. (3.10) with primes on the spacetime indices].

The variation of the action with respect to the fiducial world line X^{α} must be handled with care, since δX^{α} on the boundary $\partial \mathscr{S}$ is not independent of δX^{α} in the bulk \mathscr{S} . To isolate δX^{α} in Eq. (3.16), we must convert the surface term to a volume integral. This yields

$$\int_{\mathscr{S}} \mathrm{d}^{3} \zeta[-\overset{\circ}{\mathcal{P}}_{\mu'} K^{\mu'}{}_{\alpha} + \mathcal{S}^{i}_{\mu'} K^{\mu'}{}_{\alpha;i}] = 0 \quad (\zeta \in \mathscr{S}) \quad (3.18)$$

for the equation of motion that comes from extremization of *S* with respect to X^{μ} . This equation is a combination of the equations (3.17). To see this, we first multiply Eq. (3.17a) by $\overline{H}^{a}_{\ \nu'}K^{\nu'}{}_{\beta}$ and integrate over the matter space \mathscr{S} . We then integrate by parts on the term $S^{i}_{\ \mu';i}K^{\mu'}{}_{\beta}$. The boundary term vanishes by virtue of Eq. (3.17b) (multiplied by $\overline{H}^{a}_{\ \nu'}K^{\nu'}{}_{\beta}$). The remaining volume integral is precisely the equation of motion (3.18).

The analysis above shows that the equations of motion obtained by varying X^{μ} and ξ^{μ} are not independent; this is a

consequence of the fact that X^{μ} and ξ^{μ} constitute a larger set of variables than $X^{\mu'}$. The variables X^{μ} and ξ^{μ} contain a new gauge freedom, not present in the original variables $X^{\mu'}$, which is the freedom to choose the fiducial world line.

The action (3.14) can be written explicitly in terms of the new variables X^{μ} and ξ^{μ} . To show this, we first use the general variation (3.15) to expand the derivatives of $X^{\mu'}$ as

$$\dot{X}^{\mu'} = K^{\mu'}{}_{\alpha}\dot{X}^{\alpha} + H^{\mu'}{}_{\alpha}\dot{\xi}^{\alpha}, \qquad (3.19a)$$

$$X_{,i}^{\mu'} = H^{\mu'}{}_{\alpha}\xi_{,i}^{\alpha}.$$
 (3.19b)

As a functional of $X^{\alpha}(s)$ and $\xi^{\alpha}(s,\zeta)$, the Lagrangian density from Eq. (3.14) becomes

$$\mathcal{L} = \mathcal{L}(\zeta, g_{\mu'\nu'}, K^{\mu'}{}_{\alpha}\dot{X}^{\alpha} + H^{\mu'}{}_{\alpha}\dot{\xi}^{\alpha}, H^{\mu'}{}_{\alpha}\xi^{\alpha}, i)$$
(3.20)

where the argument of $g_{\mu'\nu'}(X')$ is written in terms of X(s)and $\xi(s,\zeta)$ using the exponential map (3.13). Next, we make use of the parallel propagator $g^{\alpha}{}_{\mu'}$ defined in Appendix B. Since the Lagrangian density \mathcal{L} is a scalar, it depends only on scalar combinations of its arguments. Furthermore, it is a property of the propagator that contractions between contravariant and covariant indices are preserved; i.e. $A^{\mu'}g^{\alpha}{}_{\mu'}g^{\nu'}{}_{\alpha}B_{\nu'} = A^{\mu'}B_{\mu'}$. Therefore, we see that the Lagrangian density can be written as

$$\mathcal{L} = \mathcal{L}(\zeta, g_{\alpha\beta}, g^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta}\dot{X}^{\beta} + g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}\dot{\xi}^{\beta}, g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}\xi^{\beta}{}_{,i}).$$
(3.21)

The action for the continuous body is

$$S[X,\xi] = \int_{s_0}^{s_1} \mathrm{d}sL$$

= $\int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta \mathcal{L}(\zeta, g_{\alpha\beta}, g^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta}\dot{X}^{\beta}$
+ $g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}\dot{\xi}^{\beta}, g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}\xi^{\beta}{}_{,i}),$ (3.22)

with the Lagrangian defined by $L = \int_{\mathscr{S}} d^3 \zeta \mathcal{L}$.

The Lagrangian density (3.21) is constructed entirely from tensor fields defined in the tangent spaces of points $X^{\mu}(s)$ along the fiducial world line. Hence, the action (3.22) describes an effective particle with world line $X^{\mu}(s)$, carrying internal degrees of freedom described by the vectors $\xi^{\mu}(s)$. The effective particle interacts nonlocally with the geometry in a neighborhood of the world line; this dependence is contained in the tensors $g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}$ and $g^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta}$. In Eqs. (B7) we show the series expansions for $g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}$ and $g^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta}$. Expressed in this way, $g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}$ and $g^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta}$ depend on the Riemann tensor and its derivatives evaluated along the fiducial world line.

CONTINUOUS BODY DYNAMICS AND THE MATHISSON- ...

The effective particle action (3.22) takes the form of Eq. (2.1) with the correspondences:

$$\psi_I \leftrightarrow \xi^{\alpha}(\zeta),$$
(3.23a)

$$\phi_A \leftrightarrow \{g_{\alpha\beta}, R_{\mu\nu\rho\sigma}, \nabla_{\alpha}R_{\mu\nu\rho\sigma}, \dots\}, \qquad (3.23b)$$

where the dots denote higher order derivatives of the Riemann tensor. Note that the index *I* now represents the continuous labels ζ^i , as well as the discrete spacetime index α . This requires us to replace certain partial derivatives with functional derivatives; for example, $\partial L/\partial \psi_I$ becomes $\delta L/\delta \xi^{\alpha}(\zeta)$. Furthermore, a repeated index *I* must include an integral over the matter space \mathscr{S} .

The momenta as defined in Eqs. (2.4) are

$$P_{\alpha} \equiv \frac{\partial L}{\partial \dot{X}^{\alpha}} = \int_{\mathscr{S}} \mathrm{d}^{3} \zeta K^{\mu'}{}_{\alpha} \mathcal{P}_{\mu'}, \qquad (3.24a)$$

$$\pi_{\alpha}(\zeta) \equiv \frac{\delta L}{\delta \xi^{\alpha}(\zeta)} = H^{\mu'}{}_{\alpha} \mathcal{P}_{\mu'}.$$
 (3.24b)

These results are most easily derived with the Lagrangian density written in the form of Eq. (3.20). We also note the result

$$\frac{\delta L}{\delta \xi^{\alpha}_{,i}} = H^{\mu'}{}_{\alpha} \mathcal{S}^{i}_{\mu'} \tag{3.25}$$

satisfied by the momentum density. From the definition (2.11), the spin is

$$S_{\alpha\beta} = 2 \int_{\mathscr{S}} \mathrm{d}^{3} \zeta \pi_{\mu} \mathrm{G}_{[\alpha\beta]} \xi^{\mu} = 2 \int_{\mathscr{S}} \mathrm{d}^{3} \zeta \xi_{[\alpha} \pi_{\beta]}. \qquad (3.26)$$

Using $d\Sigma_{\nu'} = -n_{\nu'}\sqrt{h'}d^3\zeta$, our earlier result that $\mathcal{P}_{\mu'} = -\sqrt{h'}n_{\nu'}T^{\nu'}_{\mu'}$, and the definition $\xi^{\alpha} = -\sigma^{\alpha}$, we see that the above definitions for momentum and spin coincide with Dixon's definitions in Eqs. (1.2).

The equations of motion for the effective particle, as derived in Sec. II, must be generalized to account for the fact that the index *I* now includes the continuous labels ζ^{i} . The variation of the action, from Eq. (2.5), becomes

$$\delta S = \int_{s_0}^{s_1} \mathrm{d}s \left(P_{\alpha} \Delta \dot{X}^{\alpha} + \frac{\partial L}{\partial R_{\mu\nu\rho\sigma}} \nabla_{\alpha} R_{\mu\nu\rho\sigma} \delta X^{\alpha} + \cdots \right. \\ \left. + \int_{\mathscr{S}} \mathrm{d}^3 \zeta \left[\pi_{\alpha} \Delta \dot{\xi}^{\alpha} + \frac{\delta L}{\delta \xi^{\alpha}} \Delta \xi^{\alpha} + \frac{\delta L}{\delta \xi^{\alpha}_{,i}} \Delta \xi^{\alpha}_{,i} \right] \right). \tag{3.27}$$

After integration by parts, we obtain the generalization of Eq. (2.9):

$$\delta S = \int_{s_0}^{s_1} \mathrm{d}s \left(-\mathring{P}_{\alpha} + \frac{\partial L}{\partial R_{\mu\nu\rho\sigma}} \nabla_{\alpha} R_{\mu\nu\rho\sigma} + \cdots \right. \\ \left. + R^{\mu}_{\nu\alpha\sigma} \dot{X}^{\sigma} \int_{\mathscr{S}} \mathrm{d}^3 \zeta \pi_{\mu} \xi^{\nu} \right) \delta X^{\alpha} \\ \left. + \int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta \left(-\mathring{\pi}_{\alpha} + \frac{\delta L}{\delta \xi^{\alpha}} - \frac{\mathcal{D}}{\mathcal{D} \zeta^i} \left(\frac{\delta L}{\delta \xi^{\alpha}_{,i}} \right) \right) \Delta \xi^{\alpha} \\ \left. + \int_{s_0}^{s_1} \mathrm{d}s \int_{\partial \mathscr{S}} \mathrm{d}^2 \zeta \left(\eta_i \frac{\delta L}{\delta \xi^{\alpha}_{,i}} \right) \Delta \xi^{\alpha}.$$
(3.28)

The equations of motion obtained by extremizing *S* with respect to $\xi^{\alpha}(\zeta)$ are

$$\overset{\,\,{}_{\,\,}}{\pi}_{\alpha} = \frac{\delta L}{\delta \xi^{\alpha}} - \frac{\mathcal{D}}{\mathcal{D} \zeta^{i}} \left(\frac{\delta L}{\delta \xi^{\alpha}_{,i}} \right) \quad (\zeta \in \mathscr{S}), \tag{3.29a}$$

$$\eta_i \frac{\delta L}{\delta \xi_{i,}^{\alpha}} = 0 \quad (\zeta \in \partial \mathscr{S}).$$
(3.29b)

These equations generalize the effective particle Eq. (2.10b). They are equivalent to Eqs. (3.17).

The equation of motion that comes from extremization of *S* with respect to the fiducial particle world line X^{α} is

$$\overset{\circ}{P}_{\alpha} = -R_{\alpha\beta}{}^{\rho\sigma}\dot{X}^{\beta}\int_{\mathscr{S}} \mathrm{d}^{3}\zeta\xi_{\rho}\pi_{\sigma} + \frac{\partial L}{\partial R_{\mu\nu\rho\sigma}}\nabla_{\alpha}R_{\mu\nu\rho\sigma} + \cdots.$$
(3.30)

This equation generalizes the effective particle Eq. (2.10a). It is equivalent to Eq. (3.18). As shown before, Eq. (3.30) is redundant; it can be derived from Eqs. (3.29).

We have shown that the continuous body, described by the action (3.14), can be interpreted as an effective single particle with action (3.22). This result still holds when the Lagrangian density depends on higher-order spatial derivatives of $X^{\mu'}$. In that case, the formulas for converting derivatives of $X^{\mu'}$ into derivatives of ξ^{α} and bitensors are more complicated. Nevertheless, it is still true that the final Lagrangian density can be written in terms of a $X^{\mu}(s)$, $\xi^{\alpha}(s,\zeta)$ and their derivatives, in combinations that depend on the geometry in a neighborhood of the fiducal world line. In terms of a series expansion, the dependence on geometry appears as the Riemann tensor and its derivatives evaluated at $X^{\mu}(s)$.

We can also generalize the action (3.14) by allowing for further dependence on X' through some exterior fields. This would not change the main result—we simply use the parallel propagator to express these fields in terms of tensors defined along the fiducial world line.

IV. GAUGE CONDITIONS

In this section we assume the action (3.3) for the continuous body is reparametrization invariant

(RP-invariant). A reparametrization $\tilde{s} = F(s, \zeta)$ consists in replacing the coordinates $X^{\mu'}$ with $\tilde{X}^{\mu'}$ such that $\tilde{X}^{\mu'}(\tilde{s}, \zeta) = X^{\mu'}(s, \zeta)$. In this case, the action can be written in the form (3.4) for an elastic body [21].

A natural way to remove the gauge freedom for the continuous body is to choose the "proper time gauge" $\dot{X}^{\mu'}\dot{X}_{\mu'} = -1$. With this condition, the separation *ds* between neighboring constant *s* surfaces coincides with the proper time interval measured along each of the world lines.

The effective particle action (3.22) inherits RP-invariance from the continuous body (3.3). Recall the change of variables from $X^{\mu'}$ to X^{μ} and ξ^{μ} , defined by the exponential map, Eq. (3.13). RP-invariance consists in replacing the vector $\xi^{\alpha}(s, \zeta)$ with $\tilde{\xi}^{\alpha}(\tilde{s}, \zeta)$ such that

$$\tilde{X}'(\tilde{s},\zeta) = \exp\left(X(\tilde{s}), \tilde{\xi}(\tilde{s},\zeta)\right). \tag{4.1}$$

Note that the reparametrization $\tilde{s} = F(s, \zeta)$ only changes the parametrization of the particle world lines. It does not affect the parametrization of the fiducial world line. See Fig. 1.

The fiducial world line can be chosen arbitrarily. This freedom appears as a gauge symmetry for the effective particle action (3.22), in addition to the RP-invariance. This gauge symmetry can be identified by varying the action,

$$\delta S = \int \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta \left(\frac{\delta S}{\delta X^{\mu'}} \right) \delta X^{\mu'}, \qquad (4.2)$$

where $\delta X^{\mu'}$ depends on δX^{α} and $\delta \xi^{\alpha}$ through the exponential map (3.13). Using the results of Appendix B, we find $\delta X^{\mu'} = K^{\mu'}{}_{\alpha} \delta X^{\alpha} + H^{\mu'}{}_{\alpha} \Delta \xi^{\alpha}$. This shows that the action is invariant, $\delta S = 0$, for any variation satisfying

$$\Delta\xi^{\beta} = \sigma^{\beta}{}_{\mu'}K^{\mu'}{}_{\alpha}\delta X^{\alpha}. \tag{4.3}$$

This invariance holds independent of the equations of motion.

For the effective particle, the reparametrization invariance (4.1) and fiducial world line invariance (4.3) are independent symmetries, each requiring their own conditions for gauge fixing. RP-invariance is characterized by 1 real function of 4 real parameters. World line invariance is characterized by 4 real functions of 1 real parameter.

There are many ways to fix RP-invariance for the effective particle. We will consider two different conditions, namely, the "proper time" gauge and the "normal" gauge. For the fiducial world line invariance, it is generally convenient to impose $\dot{X}^{\mu}\dot{X}_{\mu} = -1$. This partially fixes the gauge by setting the fiducial world line parameter equal to proper time. The remaining freedom in the fiducial world line is removed by imposing a center of mass condition. We show that for a specific definition of the center of mass, the center of mass condition is related to a spin supplementary condition.

The proper time gauge for an effective particle is an approximation to the conditions $\dot{X}^{\mu'}\dot{X}_{\mu'} = -1$, where $X^{\mu'}$ is defined in terms of X^{α} and ξ^{α} by the exponential map. In our examples in the next sections, we will take this approximation to third order in ϵ , where we suppose $|\xi|/\ell < \epsilon$ for any relevant length scale ℓ . For example, ℓ would include the radius of curvature $\sim 1/\sqrt{|R|}$. This condition allows for convergence of series expansions in ξ . We also impose $|\xi|/\alpha < \epsilon$. This second condition requires the relative motions of the particles to be much less than the speed of light (so that vibrational energy is not comparable to the total mass-energy).



FIG. 1. Each figure shows the fiducial world line X(s) and the world line for a generic particle $X'(s, \zeta)$ in the body. The dots on the fiducial world line indicate the parameter values for X(s). In the left figure, the particle world lines are parametrized by s; in the right figure, the particle world lines are parametrized by \tilde{s} . The spacetime point $X'(1, \zeta) = \exp(X(1), \xi(1, \zeta))$ in the left figure coincides with the spacetime point $\tilde{X}'(2, \zeta) = \exp(X(2), \tilde{\xi}(2, \zeta))$ in the right figure. The dashed line in each figure is the geodesic that defines the exponential map.

Using Eqs. (3.19) and (B7), we have

$$g^{\alpha}{}_{\mu'}\dot{X}^{\mu'} = \left(\delta^{\alpha}_{\beta} - \frac{1}{2}R^{\alpha}{}_{\xi\beta\xi} - \frac{1}{6}R^{\alpha}{}_{\xi\beta\xi;\xi}\right)\dot{X}^{\beta} + \left(\delta^{\alpha}_{\beta} - \frac{1}{6}R^{\alpha}{}_{\xi\beta\xi}\right)\dot{\xi}^{\alpha} + \mathcal{O}(\epsilon^{4}) = \dot{X}^{\alpha} + \dot{\xi}^{\alpha} - \frac{1}{2}R^{\alpha}{}_{\xi\dot{X}\xi} - \frac{1}{6}R^{\alpha}{}_{\xi\dot{\xi}\xi\xi} - \frac{1}{6}R^{\alpha}{}_{\xi\dot{X}\xi;\xi} + \mathcal{O}(\epsilon^{4}),$$
(4.4)

which gives

$$\dot{X}^{\mu'}\dot{X}_{\mu'} = \dot{X}_{\alpha}\dot{X}^{\alpha} + 2\dot{X}_{\alpha}\dot{\xi}^{\alpha} + \dot{\xi}^{2} - R_{\dot{X}\xi\dot{X}\xi} - \frac{4}{3}R_{\dot{X}\xi\ddot{\xi}\xi} - \frac{1}{3}R_{\dot{X}\xi\ddot{\xi}\xi\xi} - \frac{1}{3}R_{\dot{X}\xi\dot{\xi}\xi\xi} + \mathcal{O}(\epsilon^{4}).$$
(4.5)

The occurrence of \dot{X} and ξ as indices on the Riemann curvature tensor indicates contraction of those indices with the specified vectors. We now partially fix the fiducial world line by setting $\dot{X}_{\alpha}\dot{X}^{\alpha} = -1$. Then the proper time gauge condition $\dot{X}^{\mu'}\dot{X}_{\mu'} = -1$ yields

$$2\dot{X}_{\alpha}\overset{\alpha}{\xi}^{\alpha} + \overset{\alpha}{\xi}^{2} - R_{\dot{X}\xi\dot{\chi}\xi} - \frac{4}{3}R_{\dot{X}\xi\dot{\xi}\xi} - \frac{1}{3}R_{\dot{X}\xi\dot{\chi}\xi;\xi} + \mathcal{O}(\epsilon^{4}) = 0,$$
(4.6)

to third order in ϵ .

An alternative to the proper time gauge is the "normal gauge", in which we simply suppose the vectors $\xi^{\alpha}(s, \zeta)$ inhabit the subspace normal to some timelike vector field $V^{\alpha}(s)$ defined along the world line, that is,

$$V_{\alpha}(s)\xi^{\alpha}(s,\zeta) = 0. \tag{4.7}$$

The vector field can be a defined in various ways; for example, V^{α} might equal four-velocity U^{α} or the momentum P^{α} .

With $\dot{X}^{\alpha}\dot{X}_{\alpha} = -1$, three degrees of freedom remain to be fixed for the fiducial world line. Two options are a "center of mass" condition and or a "spin supplementary" condition $V_{\alpha}S^{\alpha\beta} = 0$. We will consider a natural formulation of the center of mass condition which is equivalent to a spin supplementary condition up to quadrupole order, though further multipole corrections exist.

The center of mass at parameter time *s* can be defined as the Frechet-Karcher mean of the surface $\Sigma(s)$. Specifically, X(s) is the center of mass of $\Sigma(s)$ if it minimizes the function [22]

$$f(y) \equiv \int_{\mathscr{S}} d^{3}\zeta w(\zeta) \sigma(y, X'(s, \zeta)), \qquad (4.8)$$

where $w(\zeta)$ can be any density defined on \mathscr{S} . Essentially, this definition tells us that the center of mass minimizes a weighted average of the squared distance between itself and every other point on the surface. It may be the case that the point x = X(s) that minimizes f(y) does not lie on the surface $\Sigma(s)$.

The most useful form of the center of mass condition is found by setting $\partial_{\alpha} f = 0$ at the minimum. With the relation (B3) this yields $\Xi_{\alpha} = 0$, where

$$\Xi_{\alpha} \equiv \int_{\mathscr{S}} d^{3} \zeta w \xi_{\alpha} \tag{4.9}$$

defines the center of mass

One natural choice for the density weight is $w = \sqrt{h'n^{\mu'}n^{\nu'}T_{\mu'\nu'}}$, where $n^{\mu'}$ is the unit vector field normal to the surface $\Sigma(s)$ of constant *s* and *h'* is the determinant of the induced metric on $\Sigma(s)$. This *w* is the energy density relative to a fleet of observers at rest in $\Sigma(s)$. If we choose the normal gauge, then $\Sigma(s)$ is (a subset of) the surface of geodesics passing through X(s) which are normal to V_{α} at X(s). This gauge choice has been considered by Costa and Natario [5], but using different definitions of P_{α} and $S_{\alpha\beta}$ than those used in this paper. They showed that with the normal gauge (with their definitions of spin and momentum), the center of mass condition $\Xi_{\alpha} = 0$ with $w = -\sqrt{h'}Z^{0,\mu'}n^{\nu'}T_{\mu'\nu'}$ is exactly equivalent to a spin supplementary condition $V_{\alpha}S^{\alpha\beta} = 0$. We produce a similar result for our definitions of P_{α} , $S_{\alpha\beta}$ and *w*.

The quickest way to derive the result is by using the series expansion of $\overline{H}^{\alpha}{}_{\mu'}n^{\mu'}$. If we differentiate the normal gauge condition (4.7) with respect to ζ^{i} , we see that it satisfies the property $V_{\alpha}\xi^{\alpha}{}_{,i} = 0$. We also have $n_{\mu'}X^{\mu'}{}_{,i} = 0$ everywhere on $\Sigma(s)$. Using the identity $\overline{H}^{\alpha}{}_{\mu'}H^{\nu'}{}_{\alpha} = \delta^{\mu'}{}_{\nu'}$ and the result (3.19b), we can write $n_{\nu'}\delta^{\nu'}_{\mu'}X^{\mu'}{}_{,i} = 0$ as

$$(H^{\nu'}{}_{\alpha}n_{\nu'})(\overset{-1}{H^{\alpha}}{}_{\mu'}X^{\mu'}_{,i}) = H^{\nu'}{}_{\alpha}n_{\nu'}\xi^{\alpha}_{,i} = 0.$$
(4.10)

This means that $H^{\mu'}{}_{\alpha}n_{\mu'}$ is parallel to V_{α} . Next, set $H^{\mu'}{}_{\alpha}n_{\mu'} = AV_{\alpha}$ and write

$$g^{\mu'\nu'}n_{\mu'}n_{\nu'} = A^2 (H^{\alpha}_{\ \mu'}H^{\beta}_{\ \nu'}g^{\mu'\nu'})V_{\alpha}V_{\beta}.$$
 (4.11)

Using the series expansion (B6) and $n^{\mu'}n_{\mu'} = -1$ we have

$$A = \left(1 - \frac{1}{3}R_{V\xi V\xi} - \frac{1}{6}\nabla_{\xi}R_{V\xi V\xi} + \mathcal{O}(\epsilon^{4})\right)^{-\frac{1}{2}}$$

= $1 + \frac{1}{6}R_{V\xi V\xi} + \frac{1}{12}\nabla_{\xi}R_{V\xi V\xi} + \mathcal{O}(\epsilon^{4}).$ (4.12)

Now we can calculate

S. P. LOOMIS and J. DAVID BROWN

$$\overset{-1}{H^{\alpha}}_{\mu'}n^{\mu'} = \overset{-1}{H^{\alpha}}_{\mu'}g^{\mu'\nu'}n_{\nu'} = \overset{-1}{H^{\alpha}}_{\mu'}g^{\mu'\nu'}(\overset{-1}{H^{\beta}}_{\nu'}H^{\lambda'}{}_{\beta})n_{\lambda'}.$$
 (4.13)

Using the results above for $H^{\lambda'}{}_{\beta}n_{\lambda'}$ and the expansion (B6b) from Appendix B, we have

$$\begin{split} \stackrel{-1}{H^{\alpha}}_{\mu'} n^{\mu'} &= \left(g^{\alpha\beta} + \frac{1}{3} R^{\alpha}{}_{\xi}{}^{\beta}{}_{\xi} + \frac{1}{6} \nabla_{\xi} R^{\alpha}{}_{\xi}{}^{\beta}{}_{\xi} + \mathcal{O}(\epsilon^{4}) \right) \\ &\times \left(1 + \frac{1}{6} R_{V\xi V\xi} + \frac{1}{12} \nabla_{\xi} R_{V\xi V\xi} + \mathcal{O}(\epsilon^{4}) \right) V_{\beta} \\ &= V^{\alpha} + \frac{1}{3} \psi^{\alpha\beta} \left(R_{\beta\xi V\xi} + \frac{1}{2} \nabla_{\xi} R_{\beta\xi V\xi} \right) + \mathcal{O}(\epsilon^{4}). \end{split}$$

$$(4.14)$$

where $\psi^{\alpha\beta} \equiv g^{\alpha\beta} + \frac{1}{2} V^{\alpha} V^{\beta}$.

Now set $w = \sqrt{h'}n^{\mu'}n^{\nu'}T_{\mu'\nu'} = -n^{\mu'}\mathcal{P}_{\mu'}$, where the second equality follows from the results of Sec. III, to obtain

$$\int_{\mathscr{S}} d^{3}\zeta w \xi_{\alpha} = -\int_{\mathscr{S}} d^{3}\zeta n^{\mu'} H^{\beta}{}_{\mu'} H^{\nu'}{}_{\beta} \mathcal{P}_{\nu'} \xi_{\alpha}$$
$$= -2 \int_{\mathscr{S}} d^{3}\zeta H^{\beta}{}_{\mu'} n^{\mu'} \xi_{[\alpha} H^{\nu'}{}_{\beta]} \mathcal{P}_{\nu'}. \quad (4.15)$$

Here we utilize the result $H^{\beta}{}_{\mu'}n^{\mu'}\xi_{\beta} = n^{\mu'}\sigma^{\beta}{}_{\mu'}\sigma_{\beta} = n^{\mu'}\sigma_{\mu'}=0$, derived from the relations in Appendix B and the fact that $\sigma_{\mu'}$ is tangent to $\Sigma(s)$ and therefore orthogonal to $n^{\mu'}$. Now replacing $H^{\beta}{}_{\mu'}n^{\mu'}$ with the expansion in Eq. (4.14) and using the definition of spin from Eqs. (3.24b) and (3.26), we find that the center of mass satisfies

$$\Xi_{\alpha} \equiv \int_{\mathscr{S}} \mathrm{d}^{3} \zeta w \xi_{\alpha} = V^{\beta} S_{\beta \alpha} - \frac{2}{3} \psi^{\beta \gamma} V^{\delta} R_{\gamma \rho \delta \sigma} O^{\rho \sigma}{}_{\alpha \beta} + \mathcal{O}(\epsilon^{4}).$$

$$(4.16)$$

Here,

$$O^{\rho\sigma}{}_{\alpha\beta} \equiv \int_{\mathscr{S}} \mathrm{d}^{3}\zeta\xi^{\rho}\xi^{\sigma}\xi_{[\alpha}H^{\mu'}{}_{\beta]}\mathcal{P}_{\mu'} \qquad (4.17)$$

is an octupole moment in the sense that it is an integral over a density field with three position vectors. It is not clear to us how this term relates to the gravitational or reduced multipoles. In any case, Eq. (4.16) shows that the center of mass condition $\Xi_{\alpha} = 0$ with appropriately chosen weight w is equivalent to a spin supplementary condition up to second order in ϵ , with an octupole correction at third order.

PHYSICAL REVIEW D 95, 044025 (2017)

V. EXAMPLE: DUST

An extremely simple example of a continuous body is a cloud of dust. This is a special case of an elastic body Eq. (3.4) with the Lagrangian density $\mathcal{L} = -\alpha\rho$, where $\rho(\zeta, f_{ij}) = \rho(\zeta)$ is independent of the fleet metric f_{ij} . Thus, for the dust cloud, there is no interaction energy between particles.

With the definition $\alpha = \sqrt{-\dot{X}^{\mu'}\dot{X}_{\mu'}}$, the dust cloud action is

$$S[X'] = -\int_{s_0}^{s_1} \mathrm{d}s \int_{\mathscr{S}} \mathrm{d}^3 \zeta \rho \sqrt{-\dot{X}^{\mu'} \dot{X}_{\mu'}}.$$
 (5.1)

The equations of motion are given in (3.10), with primes placed on the spacetime indices. In this case the stress density $S^i_{\mu'} \equiv \partial \mathcal{L} / \partial X^{\mu'}_{,i}$ vanishes, so the equations of motion reduce to $\mathring{\mathcal{P}}_{\mu'} = 0$. The momentum density is $\mathcal{P}_{\mu'} \equiv \partial \mathcal{L} / \partial \dot{X}^{\mu'} = \rho U_{\mu'}$, where $U_{\mu'}(s,\zeta) \equiv \dot{X}_{\mu'} / \sqrt{-\dot{X}^{\mu'}\dot{X}_{\mu'}}$ is the four-velocity of the particle with label ζ^i . Therefore the equations of motion for the dust cloud imply $\mathring{U}_{\mu'} = 0$; as expected, each dust particle follows a geodesic.

We now use the exponential map to write the action in terms of a fiducial world line X^{μ} and vectors ξ^{μ} . From Eq. (4.5) we have

$$\sqrt{-\dot{X}^{\mu'}\dot{X}_{\mu'}} = \sqrt{-\dot{X}^2} \left\{ 1 - U \cdot \overset{*}{\xi} - \frac{1}{2} (1 + U \cdot \overset{*}{\xi}) \right. \\ \left. \times \left(f_{\alpha\beta} \overset{*}{\xi^{\alpha}} \overset{*}{\xi^{\beta}} - R_{U\xi U\xi} \right) + \frac{2}{3} R_{U\xi \overset{*}{\xi\xi\xi}} \right. \\ \left. + \frac{1}{6} R_{U\xi U\xi;\xi} + \mathcal{O}(\epsilon^4) \right\}$$
(5.2)

where we have used the abbreviations $U^{\alpha} \equiv \dot{X}^{\alpha}/\sqrt{-\dot{X}^2}$ and $\overset{*}{\xi^{\alpha}} \equiv \overset{\circ}{\xi}^{\alpha}/\sqrt{-\dot{X}^2}$. The action is given to third order in ϵ by inserting (5.2) into (5.1).

The canonical momenta conjugate to X^{α} and ξ^{α} are

$$P_{\alpha} = \int_{\mathscr{S}} \mathrm{d}^{3}\zeta \rho \left\{ U_{\alpha} + \frac{1}{2} (f_{\beta\sigma} \xi^{\beta} \xi^{\sigma} - R_{U\xi U\xi}) + (1 + U \cdot \xi^{*}) f_{\alpha\beta} \xi^{\beta} - R_{\alpha\xi U\xi} + \mathcal{O}(\epsilon^{3}) \right\}$$
(5.3)

and

$$\pi_{\alpha} = \rho \left\{ U_{\alpha} + \frac{1}{2} U_{\alpha} (f_{\beta\sigma} \xi^{\beta} \xi^{\sigma} - R_{U\xi U\xi}) + (1 + U \cdot \xi) f_{\alpha\beta} \xi^{\beta} - \frac{2}{3} R_{U\xi \alpha\xi} + \mathcal{O}(\epsilon^{3}) \right\}, \quad (5.4)$$

respectively. The equations of motion (2.10) are

CONTINUOUS BODY DYNAMICS AND THE MATHISSON- ...

$$\overset{\circ}{P}_{\alpha} = \int_{\mathscr{S}} \mathrm{d}^{3} \zeta \rho \sqrt{-\dot{X}^{2}} \left\{ (1 + U \cdot \overset{*}{\xi}) R_{\alpha U U \xi} + R_{\alpha U \xi \xi} - \frac{1}{2} R_{U \xi U \xi; \alpha} + \mathcal{O}(\epsilon^{3}) \right\}$$
(5.5)

and

$${}^{\circ}_{\pi}{}_{\alpha} = \rho \sqrt{-\dot{X}^{2}} \bigg\{ (1 + U \cdot \overset{*}{\xi}) R_{\alpha U U \xi} + \frac{4}{3} R_{\alpha (U \xi) \xi} + \frac{1}{3} R_{\alpha U U \xi; \xi} - \frac{1}{6} R_{U \xi U \xi; \alpha} + \mathcal{O}(\epsilon^{3}) \bigg\}.$$
(5.6)

Equation (5.5) follows from Eq. (5.6). To show this, we multiply Eq. (5.6) by $\overline{H}^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta}$ and integrate over the matter space \mathscr{S} . We can make use of the relations

$$\overset{-1}{H^{\alpha}}_{\mu'}K^{\mu'}{}_{\beta} = \sigma^{\alpha}{}_{\beta} = \delta^{\alpha}_{\beta} - \frac{1}{3}R^{\alpha}{}_{\xi\beta\xi} + \mathcal{O}(\epsilon^3) \qquad (5.7)$$

which follow from Eqs. (B5) and (B7) of Appendix B. The calculation leading from (5.6) to (5.5) uses the first and second Bianchi identities.

The spin for the dust cloud, from the definition (2.11), is

$$S_{\alpha\beta} = 2 \int_{\mathscr{S}} d^{3}\zeta \rho \left\{ \xi_{[\alpha} U_{\beta]} + \frac{1}{2} \xi_{[\alpha} U_{\beta]} (f_{\sigma\rho} \xi^{\sigma} \xi^{\rho} - R_{U\xi U\xi}) \right. \\ \left. + (1 + U \cdot \xi^{*}) \xi_{[\alpha} f_{\beta]\sigma} \xi^{\sigma} - \frac{2}{3} \xi_{[\alpha} R_{\beta]\xi U\xi} + \mathcal{O}(\epsilon^{4}) \right\}.$$

$$(5.8)$$

One can show by direct calculation that the spin $S_{\alpha\beta}$ and the momentum P_{α} from (5.3) satisfy the MPD equations (2.15).

We now apply gauge conditions to analyze geodesic deviation. First, we use proper time to parametrize the fiducual world line, so that $\sqrt{-\dot{X}^2} = 1$. With this choice we have $U^{\alpha} = \dot{X}^{\alpha}$ and $\dot{\xi}^{\sigma} = \dot{\xi}^{\alpha}$. Applying the proper time gauge (4.6), the momentum (5.4) becomes

$$\pi_{\alpha} = \rho \left\{ \dot{X}_{\alpha} + \overset{\circ}{\xi}_{\alpha} - \frac{2}{3} R_{\dot{X}\xi\alpha\xi} + \mathcal{O}(\epsilon^3) \right\}.$$
(5.9)

The equation of motion (5.6) reduces to

$$\overset{\circ}{\pi}_{\alpha} = \rho \left\{ R_{\alpha \dot{X} \dot{X} \xi} + \frac{4}{3} R_{\alpha (\dot{X} \overset{\circ}{\xi}) \xi} + \frac{1}{3} R_{\alpha \dot{X} \dot{X} \xi; \xi} - \frac{1}{6} R_{\dot{X} \xi \dot{X} \xi; \alpha} + \mathcal{O}(\epsilon^3) \right\}.$$
(5.10)

Finally, we can fix the fiducial world line to be a geodesic; this implies $\mathring{U}^{\alpha} = \mathring{X}^{\alpha} = 0$. The equation of motion (5.10) together with the definition (5.9) yields

$$\begin{aligned} \overset{\circ\circ}{\xi}_{\alpha}^{\circ} &-\frac{2}{3}R_{\alpha\xi\dot{X}\xi;\dot{X}} + \frac{4}{3}R_{\alpha(\xi\xi)\dot{X}} \\ &= R_{\alpha\dot{X}\dot{X}\xi} + \frac{4}{3}R_{\alpha(\dot{X}\xi)\xi} - \frac{1}{6}R_{\dot{X}\xi\dot{X}\xi;\alpha} \\ &+ \frac{1}{3}R_{\alpha\dot{X}\dot{X}\xi;\xi} + \mathcal{O}(\epsilon^{3}). \end{aligned}$$
(5.11)

Applying the first and second Bianchi identities simplifies this to

$$\overset{\circ\circ}{\xi}_{\alpha}^{\circ} = R_{\alpha \dot{X} \dot{X} \xi} + 2R_{\alpha \dot{\xi} \dot{X} \xi} - \nabla_{(\xi} R_{\dot{X}) \alpha \dot{X} \xi} + \mathcal{O}(\epsilon^3) \quad (5.12)$$

which matches previous derivations of the geodesic deviation equations to second order [23,24].

Geodesic deviation is often studied in terms of the *n*th order Jacobi fields J_n^{α} , each of which is associated with a finite differential equation linear in J_n^{α} and only including lower-order Jacobi fields. We can generalize to our three-dimensional congruence of geodesics, representing Jacobi fields with spatial derivatives of the vector ξ^{α} evaluated at ζ_0^i :

$$J^{\alpha}_{i_1\cdots i_n} \equiv \xi^{\alpha}_{,i_1\cdots i_n}(s,\zeta_0). \tag{5.13}$$

The equations of motion for the first- and second-order Jacobi fields can be computed from the first and second spatial derivatives of (5.12), respectively, evaluated at ζ_0^i . This yields

$$\widetilde{J}_{i}^{\alpha} = R^{\alpha}{}_{\dot{X}\dot{X}\beta}J_{i}^{\beta}, \qquad (5.14a)$$

$$\begin{split} \overset{\circ}{J}_{ij}^{\alpha} &= R^{\alpha}{}_{\dot{X}\dot{X}\beta}J_{ij}^{\beta} + 2\nabla_{(\beta}R^{\alpha}{}_{\dot{X})\dot{X}\gamma}J_{(i}^{\beta}J_{j)}^{\gamma} \\ &+ 4R^{\alpha}{}_{\dot{\beta}\dot{X}\gamma}\overset{\circ}{J}_{(i}^{\beta}J_{j)}^{\gamma} \end{split}$$
(5.14b)

which are the three-dimensional analogues of the traditional geodesic deviation equation and Bazanski's equation [25], respectively. These results are exact (requiring no further expansion in ϵ) because all $\mathcal{O}(\epsilon^3)$ terms vanish when their first and second spatial derivatives are evaluated at ζ_0^i .

Alternatively, instead of applying the proper time gauge and fixing the fiducial world line to be a geodesic, we can choose the normal gauge $U_{\alpha}\xi^{\alpha} = 0$ and the center-of-mass condition $\Xi_{\alpha} = 0$. For dust, the center of mass with density weight $w = \sqrt{h'}n^{\mu'}n^{\nu'}T_{\mu'\nu'}$ is

$$\Xi_{\alpha} = \int_{\mathscr{S}} \mathrm{d}^{3} \zeta \gamma \rho \xi_{\alpha} \tag{5.15}$$

where $\gamma = -n_{\mu'}U^{\mu'}$ is the relativistic gamma factor between the dust particles and the observers at rest in $\Sigma(s)$. The series expansion for γ can be computed using Eqs. (3.19a), (5.7), and the results for $H^{\mu'}{}_{\alpha}n_{\mu'}$ from Sec. IV. To second order, we find

$$\gamma = 1 + \frac{1}{2} f_{\alpha\beta} \xi^{\ast} \xi^{\ast} \xi^{\beta} + \mathcal{O}(\epsilon^3).$$
 (5.16)

Under these gauge conditions, the spin becomes

$$S_{\alpha\beta} = 2 \int_{\mathscr{S}} \mathrm{d}^{3}\zeta \rho \left\{ (1 + U \cdot \overset{*}{\xi}) \xi_{[\alpha} f_{\beta]\gamma} \overset{*}{\xi^{\gamma}} - \frac{1}{2} R_{U\xi U\xi} \xi_{[\alpha} U_{\beta]} - \frac{2}{3} \xi_{[\alpha} R_{\beta]\xi U\xi} + \mathcal{O}(\epsilon^{4}) \right\}.$$
(5.17)

To order ϵ^2 , the spin has the familiar form for the angular momentum of a system about its center of mass; roughly, $S_{\alpha\beta}$ is a sum over particles of the "cross product" between the position vector ξ_{α} and the relative momentum vector $d^3\zeta\rho f_{\beta\gamma}\xi^{\gamma}$ (projected orthogonal to the four-velocity U_{α} of the fiducial world line).

Taking the inner product of $S_{\alpha\beta}$ with U^{β} yields

$$U^{\beta}S_{\beta\alpha} - \frac{1}{6} \int_{\mathscr{S}} \mathrm{d}^{3}\zeta \rho \xi_{\alpha} R_{U\xi U\xi} + \mathcal{O}(\epsilon^{4}) = 0.$$
 (5.18)

This is precisely the center of mass condition $\Xi_{\alpha} = 0$ with Ξ_{α} written in terms of spin as in Eq. (4.16). It has the form of a spin supplementary condition with octupole correction.

VI. DISCUSSION

In Sec. II we generalized the result of Steinhoff [13], to show that "effective particle" actions of the form (2.1) always yield the MPD equations. These actions make no explicit reference to a body frame or angular velocity, and the spin is defined directly in terms of the internal structure of the effective particle.

We define continuous bodies in terms of a congruence of world lines in Sec. III, and show how a continuous body can be expressed as an effective particle. The momentum and spin for this effective particle are exactly equivalent to Dixon's definitions for the momentum and spin of extended bodies.

The MPD equations are often considered to be "incomplete", requiring the addition of a spin supplementary condition. By starting with the action principle for a continuous body, written as an effective particle, we see clearly that the fiducial particle world line is a gauge degree of freedom. The equations of motion, and the MPD equations in particular, are "complete" in the sense of providing an unambiguous description of the evolution of the physical system. They are incomplete only in the sense that gauge conditions are needed to specify the evolution of the gauge variables. In Sec. IV we defined the center of mass of the continuous body and fixed the effective particle world line to coincide with the center of mass. With the chosen center of mass density weight *w*, we showed that this gauge condition can be written as a spin-supplementary condition with higher-order corrections.

Note that the action (3.3) depends on the position X^{μ} through the metric tensor, which is treated as a fixed field. Treating the metric as a dynamical variable can provide a new direction for the study of gravitational self-force.

ACKNOWLEDGMENTS

We would like to offer our gratitude to Justin Vines for sharing his notes and for helpful conversations. This research was supported by the Provost's Professional Experience Program at NCSU.

APPENDIX A: COVARIANT VARIATIONS AND THE EXPONENTIAL MAP

Consider a tensor ψ_I that lives in the tangent space at a point *x* in a manifold \mathcal{M} . The variation $\delta \psi_I = \psi'_I - \psi_I$ is the difference between ψ_I and another tensor, ψ'_I , which is (in some sense) close to ψ_I . If ψ'_I lives in a nearby tangent space at point *x'*, then $\delta \psi_I$ is not a tensor. We define the difference between ψ'_I and ψ_I as a tensor by parallel transporting ψ'_I to the tangent space at *x*, then subtracting. This difference is $\Delta \psi_I = \delta \psi_I + \Gamma^{\sigma}_{\alpha \rho} \delta x^{\alpha} G^{\rho}{}_{\sigma} \psi_I$, where $\delta x = x' - x$. This leads us to define the covariant variation

$$\Delta \equiv \delta + \Gamma^{\sigma}_{\alpha\rho} \delta x^{\alpha} G^{\rho}{}_{\sigma} \tag{A1}$$

acting on tensors. Here, $G^{\rho}{}_{\sigma}$ is an operator acting on tensor indices. Specifically, $G^{\alpha}{}_{\beta}$ are representations of the Lie algebra of the general linear group GL(4) acting on the tensor space [15]. In particular, for contravariant and covariant vectors ψ^{μ} and ψ_{μ} , we have

$$G^{\alpha}{}_{\beta}\psi^{\mu} = \delta^{\mu}_{\beta}\psi^{\alpha}, \qquad (A2a)$$

$$\mathbf{G}^{\alpha}{}_{\beta}\psi_{\mu} = -\delta^{\alpha}_{\mu}\psi_{\beta}. \tag{A2b}$$

The G's satisfy the commutation relations

$$[\mathbf{G}^{\alpha}{}_{\beta},\mathbf{G}^{\mu}{}_{\nu}] = \delta^{\mu}_{\beta}\mathbf{G}^{\alpha}{}_{\nu} - \delta^{\alpha}_{\nu}\mathbf{G}^{\mu}{}_{\beta} \tag{A3}$$

for the Lie algebra of GL(4).

For a given point $x \in \mathcal{M}$ and a vector $\xi \in T_x \mathcal{M}$, let $\gamma_{x,\xi}(u)$ be the affinely parametrized geodesic satisfying $\gamma_{x,\xi}(0) = x$ and $\dot{\gamma}_{x,\xi}(0) = \xi$. Define the exponential map as $\exp(x,\xi) = \gamma_{x,\xi}(1)$. In component form, we abbreviate $x^{\mu'} = \exp^{\mu'}(x,\xi)$, where $x^{\mu'}$ are the coordinates of the point x'. Primed indices are also used to denote tensors in the tangent space at x'.

The variation of the exponential map is

$$\delta x^{\mu'} = \frac{\partial x^{\mu'}}{\partial x^{\alpha}} \delta x^{\alpha} + \frac{\partial x^{\mu'}}{\partial \xi^{\alpha}} \delta \xi^{\alpha}.$$
 (A4)

As discussed above, $\delta \xi^{\alpha}$ is not a tensor whenever the variation of the base point *x* is nonzero. Using the definition (A1), and Eq. (A2), we have

$$\delta x^{\mu'} = \left(\frac{\partial x^{\mu'}}{\partial x^{\alpha}} - \frac{\partial x^{\mu'}}{\partial \xi^{\sigma}} \Gamma^{\sigma}_{\alpha\rho} \xi^{\rho}\right) \delta x^{\alpha} + \frac{\partial x^{\mu'}}{\partial \xi^{\alpha}} \Delta \xi^{\alpha}.$$
 (A5)

The coefficients of δx^{α} and $\Delta \xi^{\alpha}$ are

$$K^{\mu'}{}_{\alpha} \equiv \frac{\partial x^{\mu'}}{\partial x^{\alpha}} - \Gamma^{\gamma}{}_{\alpha\beta}\xi^{\beta}\frac{\partial x^{\mu'}}{\partial \xi^{\gamma}}, \qquad (A6a)$$

$$H^{\mu'}{}_{\alpha} \equiv \frac{\partial x^{\mu'}}{\partial \xi^{\alpha}}.$$
 (A6b)

These are bitensors, meaning that they depend on two points x and x', and they have indices that exist in the tangent spaces of both points.

Using this notation, the variation of the exponential map becomes

$$\delta x^{\mu'} = K^{\mu'}{}_{\alpha} \delta x^{\alpha} + H^{\mu'}{}_{\alpha} \Delta \xi^{\alpha}. \tag{A7}$$

We use this result in the main text to compute results (3.19a) and (3.19b).

The definitions of $H^{\mu'}{}_{\alpha}$ and $K^{\mu'}{}_{\alpha}$ as the vertical and horizontal derivatives of the exponential function were, to our knowledge, first given in [23], though Dixon employs similar definitions in [9]. *H* and *K* are often called the *Jacobi propagators* and were first discovered in the context of Synge's world function σ [6].

Next, consider a tensor function $T(x, \psi_I)$ on the tensor bundle. That is, $T(x, \psi_I)$ depends on the point $x \in \mathcal{M}$ and tensor arguments ψ_I . Both *T* and ψ_I exist in the tangent space of *x*. The variation of *T* is

$$\delta T = \frac{\partial T}{\partial x^{\alpha}} \delta x^{\alpha} + \frac{\partial T}{\partial \psi_I} \delta \psi_I.$$
 (A8)

When δx^{α} is not zero, neither δT nor $\delta \psi_I$ are tensors. Using the covariant variation (A1), we have

$$\Delta T = \left(\frac{\partial T}{\partial x^{\alpha}} + \Gamma^{\sigma}_{\alpha\rho} \mathbf{G}^{\rho}{}_{\sigma}T - \frac{\partial T}{\partial \psi_{I}} \Gamma^{\sigma}_{\alpha\rho} \mathbf{G}^{\rho}{}_{\sigma}\psi_{I}\right) \delta x^{\alpha} + \frac{\partial T}{\partial \psi_{I}} \Delta \psi_{I}.$$
(A9)

Each of the factors on the right-hand side of this equation is covariant. Note that the first two terms in parenthesis give the traditional definition of the covariant derivative, PHYSICAL REVIEW D 95, 044025 (2017)

$$\nabla_{\alpha}T \equiv \frac{\partial T}{\partial x^{\alpha}} + \Gamma^{\sigma}_{\alpha\rho} \mathbf{G}^{\rho}{}_{\sigma}T.$$
 (A10)

When T depends on ψ_I , $\nabla_{\alpha}T$ is not covariant. Covariance requires the third term in parenthesis above, which arises from the parallel transport of ψ as the base point x is varied.

It would be a misnomer to call ∇_{α} the "covariant derivative": we shall opt here to call it by its other traditional name, the Levi-Civita derivative. The covariant derivative is defined by

$$\frac{\mathcal{D}T}{\mathcal{D}x^{\alpha}} \equiv \nabla_{\alpha}T - \frac{\partial T}{\partial \psi_{I}}\Gamma^{\sigma}_{\alpha\rho}\mathbf{G}^{\rho}{}_{\sigma}\psi_{I}.$$
(A11)

With this notation, the covariant variation of T becomes

$$\Delta T = \frac{\mathcal{D}T}{\mathcal{D}x^{\alpha}} \delta x^{\alpha} + \frac{\partial T}{\partial \psi_I} \Delta \psi_I.$$
 (A12)

We also use the shorthand notation $\mathcal{D}_{\alpha}T = \mathcal{D}T/\mathcal{D}x^{\alpha}$. It is worth noting that when ψ_I is the contravariant vector ξ^{α} , the covariant derivative \mathcal{D}_{α} corresponds to the vertical derivative $\nabla_{*\alpha}$ defined by Dixon and $\partial/\partial\psi_I = \partial/\partial\xi^{\alpha}$ corresponds to the horizontal derivative $\nabla_{\alpha*}$ [8,26]. The notation $\mathcal{D}T/\mathcal{D}x^{\alpha}$ for the covariant partial derivative is adopted from Vines [19].

For the case where *T* has no explicit dependence on *x* [that is, $T = T(\psi_I)$], Eq. (A8) becomes $\delta T = (\partial T / \partial \psi_I) \delta \psi_I$. We can derive a "chain rule" for the generators G^{ρ}_{σ} by considering the passive coordinate transformations $\tilde{x}^{\alpha} = x^{\alpha} + \delta x^{\alpha}$. Note that, in this context, δx^{α} denotes the difference between coordinate values, using two different coordinate systems, at a single point in the manifold \mathcal{M} . Elsewhere in this appendix, δx^{α} denotes the difference between coordinates values, using a single coordinate system, for two distinct points of \mathcal{M} .

Under the passive coordinate transformation, T and ψ_I transform as

$$\delta T = -\partial_{\alpha} \delta x^{\beta} G^{\alpha}{}_{\beta} T, \qquad (A13a)$$

$$\delta \psi_I = -\partial_\alpha \delta x^\beta \mathbf{G}^\alpha{}_\beta \psi_I. \tag{A13b}$$

Using these results in the relation $\delta T = (\partial T / \partial \psi_I) \delta \psi_I$ and noting that $\partial_\alpha \delta x^\beta$ is arbitrary, we have

$$\mathbf{G}^{\alpha}{}_{\beta}T = \frac{\partial T}{\partial \psi_I} \mathbf{G}^{\alpha}{}_{\beta}\psi_I. \tag{A14}$$

In this paper we often consider the case $T = T(\phi_A, \psi_I)$, where $\phi_A = \phi_A(x)$ is a field defined on the manifold, and $\psi_I(s)$ is a tensor function defined solely along a world line $x^{\alpha} = X^{\alpha}(s)$. In this case the results (A9) and (A14) become

$$\Delta T = \frac{\partial T}{\partial \phi_A} \Delta \phi_A + \frac{\partial T}{\partial \psi_I} \Delta \psi_I, \qquad (A15a)$$

 Δ

$$\mathbf{G}^{\alpha}{}_{\beta}T = \frac{\partial T}{\partial \phi_A} \mathbf{G}^{\alpha}{}_{\beta}\phi_A + \frac{\partial T}{\partial \psi_I} \mathbf{G}^{\alpha}{}_{\beta}\psi_I, \qquad (A15b)$$

respectively.

The results (A15) are applied in the main text to the effective particle Lagrangian, which is a spacetime scalar, by evaluating these relations at the spacetime point x = X(s) on the particle world line. In particular, the variation of the action Eq. (2.5) and the property (2.13) of the Lagrangian are obtained from Eqs. (A15). The results (A15) are also applied to the Lagrangian density for the continuous body by evaluating them at the spacetime point $x = X(s, \zeta)$. See for example the variation of the action in Eq. (3.6) and the calculation leading to the stress-energy tensor (3.12).

Now suppose that $T_1(x, \psi_I)$ is a tensor function on the tensor bundle defined by $T_1(x, \psi_I) \equiv T_2(\phi_A(x, \psi_I), \psi_I)$, where $\phi_A(x, \psi)$ are also functions on the tensor bundle. By applying Eq. (A15a) to T_2 and Eq. (A12) to T_1 and ϕ_A , we find the chain rules

$$\frac{\mathcal{D}T_1}{\mathcal{D}x^{\alpha}} = \frac{\partial T_2}{\partial \phi_A} \frac{\mathcal{D}\phi_A}{\mathcal{D}x^{\alpha}}$$
(A16a)

$$\frac{\partial T_1}{\partial \psi_I} = \frac{\partial T_2}{\partial \phi_A} \frac{\partial \phi_A}{\partial \psi_I} + \frac{\partial T_2}{\partial \psi_I}.$$
 (A16b)

These results are used in Appendix B.

Finally, suppose we have two independent variations Δ_1 and Δ_2 acting on a general tensor $T(x, \psi_I)$. The covariant variation Δ_1 corresponds to a change in x^{α} by $\delta_1 x^{\alpha}$ and Δ_2 corresponds to the change $\delta_2 x^{\alpha}$. Explicitly, we have $\Delta_1 =$ $\delta_1 + \Gamma^{\sigma}_{\alpha\rho} \delta_1 x^{\alpha}$ and $\Delta_2 = \delta_2 + \Gamma^{\sigma}_{\alpha\rho} \delta_2 x^{\alpha}$. In general these variations will not commute. Rather,

$$\begin{split} [\Delta_1, \Delta_2] &= [\delta_1 + \Gamma^{\sigma}_{\alpha\rho} \delta_1 x^{\alpha} \mathbf{G}^{\rho}{}_{\sigma}, \Delta_2] \\ &= \delta_1 \Delta_2 + \Gamma^{\sigma}_{\alpha\rho} \delta_1 x^{\alpha} \mathbf{G}^{\rho}{}_{\sigma} \Delta_2 - (1 \leftrightarrow 2) \quad (A17) \end{split}$$

where $(1 \leftrightarrow 2)$ denotes the previous terms with indices 1 and 2 exchanged. Note that Δ_2 is a scalar operator—it carries no tensor indices—so it commutes with the generator G^{ρ}_{σ} . Thus, we have

$$\begin{aligned} \mathbf{G}^{\rho}{}_{\sigma}\Delta_{2}T &= \Delta_{2}(\mathbf{G}^{\rho}{}_{\sigma}T) \\ &= \delta_{2}(\mathbf{G}^{\rho}{}_{\sigma}T) + \Gamma^{\nu}{}_{\beta\mu}\delta_{2}x^{\beta}\mathbf{G}^{\mu}{}_{\nu}\mathbf{G}^{\rho}{}_{\sigma}T. \end{aligned} \tag{A18}$$

With this result it is straightforward to obtain

$$\begin{aligned} {}_{1}, \Delta_{2}] &= \delta_{1}\delta_{2} + \delta_{1}(\Gamma^{\nu}_{\beta\mu}\delta_{2}x^{\beta}G^{\mu}{}_{\nu}) \\ &+ \Gamma^{\sigma}_{\alpha\rho}\Gamma^{\nu}_{\beta\mu}\delta_{1}x^{\alpha}\delta_{2}x^{\beta}G^{\mu}{}_{\nu}G^{\rho}_{\sigma} - (1 \leftrightarrow 2). \end{aligned}$$
 (A19)

(The variation δ_1 in the second term on the right acts only on the factor in parenthesis.) Terms such as $\delta_1 \Gamma^{\nu}_{\beta\mu}$ yield derivatives of Christoffel symbols. The terms $G^{\mu}_{\nu} G^{\rho}_{\sigma}$ are simplified using the commutation relations (A3). The final result is

$$\begin{aligned} [\Delta_1, \Delta_2] &= [\delta_1, \delta_2] + \Gamma^{\nu}_{\alpha\mu} ([\delta_1, \delta_2] x^{\alpha}) G^{\mu}{}_{\nu} \\ &+ R^{\sigma}{}_{\rho\mu\nu} \delta_1 x^{\mu} \delta_2 x^{\nu} G^{\rho}{}_{\sigma}, \end{aligned} \tag{A20}$$

where $R^{\sigma}_{\rho\mu\nu}$ is the Riemann tensor. Note that the first two terms on the right-hand side are the covariant extension of the operator $[\delta_1, \delta_2]$.

Now suppose x is a point on a world line, x = X(s), so the tensors T, ϕ_A and ψ_I depend on the parameter s. Let $\delta_2 = \delta s(d/ds)$, where δs is infinitesimal. Then $\delta_2 x^{\alpha} = \dot{X}^{\alpha} \delta s$ and $\Delta_2 = \delta s(D/Ds)$, where D/Ds is the covariant derivative along the world line. We can also drop the subscripts from δ_1 and Δ_1 . The result (A20) now gives the commutator of a general covariant variation Δ with the covariant derivative D/Ds. Since δ and d/ds commute, we have

$$\left[\Delta, \frac{\mathrm{D}}{\mathrm{D}s}\right] = R^{\sigma}{}_{\rho\alpha\beta}\delta X^{\alpha}\dot{X}^{\beta}\mathrm{G}^{\rho}{}_{\sigma}.$$
 (A21)

This result is used in the variation of the Lagrangian in Sec. II.

APPENDIX B: BITENSORS AND SYNGE'S WORLD FUNCTION

Synge's world function is closely related to the exponential function. Given two points *x* and *x'*, take the geodesic $\gamma_{x,x'}(u)$ affinely parametrized so that $\gamma_{x,x'}(0) = x$ and $\gamma_{x,x'}(1) = x'$. Then Synge's world function is defined as [27]

$$\sigma(x,x') \equiv \frac{1}{2} \int_0^1 \mathrm{d}u g_{\alpha\beta}(\gamma_{x,x'}(u)) \frac{\mathrm{d}\gamma^{\alpha}_{x,x'}}{\mathrm{d}u} \frac{\mathrm{d}\gamma^{\beta}_{x,x'}}{\mathrm{d}u}, \quad (B1)$$

which is one-half the squared distance of the geodesic connecting x and x'. Obviously, this function is well-defined only when x and x' are close enough that they are connected by one geodesic. In this paper we consider the approximation of small bodies and assume that this is the case for any spacelike-separated x and x' in the world tube \mathcal{W} of our body.

When taking derivatives of σ , we omit semicolons following the standard notation, and as before use primed indices for derivatives with respect to x' and unprimed indices for x. It can be seen that Synge's world function satisfies the formulae [28]

$$\sigma^{\mu'}(x, x') = \dot{\gamma}^{\mu'}_{x, x'}(1),$$
 (B2b)

$$\sigma^{\mu'}\sigma_{\mu'} = \sigma^{\alpha}\sigma_{\alpha} = 2\sigma, \tag{B2c}$$

$$\sigma^{\mu'}_{\alpha}\sigma_{\mu'}=\sigma^{\beta}_{\alpha}\sigma_{\beta}=\sigma_{\alpha}, \hspace{1cm} (\text{B2d})$$

$$\sigma_{\mu'}^{\nu'}\sigma_{\nu'} = \sigma_{\mu'}^{\alpha}\sigma_{\alpha} = \sigma_{\mu'}.$$
 (B2e)

Based on (B2a) and the definition of the exponential map, we see that $x'(x,\xi)$ and $\sigma(x,x')$ are related by

$$\xi^{\alpha} = -\sigma^{\alpha}(x, x'). \tag{B3}$$

We can treat each side of this equation as a tensor function of x^{α} and ξ^{α} by writing $\xi^{\alpha} = -\sigma^{\alpha}(x, x'(x, \xi))$. This allows us to apply the covariant variation (A12) and get

$$\Delta\xi^{\alpha} = -(\sigma^{\alpha}{}_{\beta} + \sigma^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta})\delta x^{\beta} - \sigma^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta}\Delta\xi^{\beta}$$
(B4)

where we have used the chain rule (A16). Matching coefficients gives $-\sigma^{\alpha}_{\mu'}H^{\mu'}{}_{\beta} = \delta^{\alpha}_{\beta}$ and $\sigma^{\alpha}{}_{\beta} + \sigma^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta} = 0$, which imply

$$K^{\mu'}{}_{\alpha} = -\overset{-1}{\sigma}^{\mu'}{}_{\beta}\sigma^{\beta}{}_{\alpha}, \qquad (B5a)$$

$$H^{\mu'}{}_{\alpha} = -\overset{-1}{\sigma}^{\mu'}{}_{\alpha}. \tag{B5b}$$

The symbol -1 over the matrices indicates matrix inversion. Equations (B5b) and (B5a) are Dixon's original definitions for *H* and *K* [6].

We can use the Jacobi propagator H and its inverse to transfer indices on bitensors into a single tangent space, so that we can sensibly define series expansions for bitensors. In Riemann normal coordinates, the components of $\xi^{\alpha}(x, x')$ are (by definition) equal to the coordinates of the point x'. Thus, in Riemann normal coordinates, we have $H^{\mu'}{}_{\alpha} = \delta^{\mu'}_{\alpha}$. (This equality is by components, and is obviously not a tensor equation.) As such, the coefficients

of a series expansion mediated by H and its inverse can be found from the Taylor expansion of the same tensor field in Riemann normal coordinates. For example, in this paper we use the following expansion for the metric tensor and its inverse:

$$H^{\mu'}{}_{\alpha}H^{\nu'}{}_{\beta}g_{\mu'\nu'}(x') = g_{\alpha\beta}(x) - \frac{1}{3}R_{\alpha\xi\beta\xi}(x) - \frac{1}{6}R_{\alpha\xi\beta\xi;\xi}(x) + \mathcal{O}(\epsilon^4), \qquad (B6a)$$

$$\begin{split} \stackrel{-1}{H^{\alpha}}_{\mu'} \stackrel{-1}{H^{\beta}}_{\nu'} g^{\mu'\nu'}(x') &= g^{\alpha\beta}(x) + \frac{1}{3} R^{\alpha}{}_{\xi}{}^{\beta}{}_{\xi}(x) \\ &+ \frac{1}{6} R^{\alpha}{}_{\xi}{}^{\beta}{}_{\xi;\xi}(x) + \mathcal{O}(\epsilon^{4}). \end{split} \tag{B6b}$$

These follow from the familiar expansions for the metric and its inverse in Riemann normal coordinates [29]. Here, ξ appears as an index in the Riemann curvature tensor, indicating contraction of that index with ξ^{α} . The coefficients in the *H*-series expansion of a tensor field are the *Veblen extensions* of that tensor field [30].

Another important bitensor is the parallel propagator $g^{\mu'}{}_{\alpha}(x, x')$. This bitensor parallel propagates contravariant indices from $T_x \mathcal{M}$ to $T_{x'} \mathcal{M}$ along the geodesic $\gamma_{x,x'}$. Because the metric is parallel propagated, we have $g^{\alpha\beta}(x)g^{\mu'}{}_{\alpha}g^{\nu'}{}_{\beta} = g^{\mu'\nu'}(x')$. It follows that $g^{\alpha}{}_{\mu'}$ parallel propagates covariant indices from $T_x \mathcal{M}$ to $T_{x'} \mathcal{M}$. We also have the relations $g^{\alpha}{}_{\mu'}g^{\mu'}{}_{\beta} = \delta^{\alpha}{}_{\beta}$ and $g^{\mu'}{}_{\alpha}g^{\alpha}{}_{\nu'} = \delta^{\mu'}{}_{\nu'}$, so that $g^{\alpha}{}_{\mu'}$ and $g^{\mu'}{}_{\alpha}$ can inversely propagate contravariant and covariant indices from $T_x \mathcal{M}$ to $T_x \mathcal{M}$, respectively.

We can also use the parallel propagator to define series expansions for bitensors. In this paper we use the expansions [19]

$$g^{\alpha}{}_{\mu'}H^{\mu'}{}_{\beta} = \delta^{\alpha}_{\beta} - \frac{1}{6}R^{\alpha}{}_{\xi\beta\xi} - \frac{1}{12}R^{\alpha}{}_{\xi\beta\xi;\xi} + \mathcal{O}(\epsilon^4), \quad (B7a)$$

$$g^{\alpha}{}_{\mu'}K^{\mu'}{}_{\beta} = \delta^{\alpha}_{\beta} - \frac{1}{2}R^{\alpha}{}_{\xi\beta\xi} - \frac{1}{6}R^{\alpha}{}_{\xi\beta\xi;\xi} + \mathcal{O}(\epsilon^4) \qquad (B7b)$$

for the Jacobi propagators.

- M. Mathisson, Republication of: New mechanics of material systems, Gen. Relativ. Gravit. 42, 1011 (2010).
- [2] A. Papapetrou, Spinning test particles in general relativity I, Proc. R. Soc. A 209, 248 (1951).
- [3] F. Pirani, Republication of: On the physical significance of the Riemann tensor, Gen. Relativ. Gravit. **41**, 1215 (2009).
- [4] K. Kyrian and O. Semerak, Spinning test particles in a Kerr field II, Mon. Not. R. Astron. Soc. 382, 1922 (2007).
- [5] L. F. O. Costa and J. Natário, Center of mass, spin supplementary conditions, and the momentum of spinning particles, Fundamental Theories of Physics **179**, 215 (2015).

- S. P. LOOMIS and J. DAVID BROWN
- [6] W.G. Dixon, Dynamics of extended bodies in general relativity. I. Momentum and angular momentum, Proc. R. Soc. A **314**, 499 (1970).
- [7] W.G. Dixon, Dynamics of extended bodies in general relativity. II. Moments of the charge-current vector, Proc. R. Soc. A **319**, 509 (1970).
- [8] W. G. Dixon, Dynamics of extended bodies in general relativity. III. Equations of motion, Proc. Trans. R. Soc. Lond. A 277, 59 (1974).
- [9] W. G. Dixon, The new mechanics of Myron Mathisson and its subsequent development, Fundam. Theor. Phys. **179**, 1 (2015).
- [10] I. Bailey and W. Israel, Lagrangian dynamics of spinning particles and polarized media in general relativity, Commun. Math. Phys. 42, 65 (1975).
- [11] A. J. Hanson and T. Regge, The relativistic spherical top, Ann. Phys. (N.Y.) 87, 498 (1974).
- [12] R. Porto, Post-Newtonian corrections to the motion of spinning bodies in nonrelativistic general relativity, Phys. Rev. D 73, 104031 (2006).
- [13] J. Steinhoff, Spin and quadrupole contributions to the motion of astrophysical binaries, Fundamental Theories of Physics 179, 615 (2015).
- [14] H. Fuchs, General Lagrangian approach to the equations of motion for test particles with internal structure, Astron. Nachr. 307, 281 (1986).
- [15] B. DeWitt, in *Relativity, Groups, and Topology, Les Houches 1963* (Gordon and Breach, New York, 1964).
- [16] B. Carter and H. Quintana, Foundations of general relativistic high-pressure elasticity theory, Proc. R. Soc. A 331, 57 (1972).

- [17] J. Kijowski, A. Smólski, and A. Górnika, Hamiltonian theory of self-gravitating perfect fluid and a method of effective deparametrization of Einstein's theory of gravitation, Phys. Rev. D 41, 1875 (1990).
- [18] J. D. Brown and D. Marolf, On relativistic material reference systems, Phys. Rev. D 53, 1835 (1996).
- [19] J. Vines (to be published).
- [20] B. DeWitt, in *Gravitation: An Introduction to Current Research*, edited by L. Witten (Wiley, New York, 1962).
- [21] J. D. Brown and S. P. Loomis (to be published).
- [22] H. Karcher, Riemannian center of mass and mollifier smoothing, Commun. Pure Appl. Math. 30, 509 (1977).
- [23] J. Vines, Geodesic deviation at higher orders via covariant bitensors, Gen. Relativ. Gravit. 47, 59 (2015).
- [24] A. N. Aleksandrov and K. A. Piragas, Geodesic structure, Theor. Math. Phys. 38, 48 (1979).
- [25] S. L. Bazanski, Kinematics of relative motion of test particles in general relativity, Annales de L'Institut Henri Poincare Section Physique Theorique 27, 115 (1977).
- [26] W. Dixon, in *Isolated Gravitating Systems in General Relativity*, edited by J. Ehlers (North-Holland Publishing Company, Amsterdam, 1979).
- [27] J. L. Synge, *Relativity: The General Theory* (North-Holland Publishing Company, Amsterdam, 1960).
- [28] E. Poisson, A. Pound, and I. Vega, The motion of point particles in curved spacetime, Living Rev. Relativ. 14, 7 (2011).
- [29] L. Brewin, Riemann normal coordinate expansions using Cadabra, Classical Quantum Gravity 26, 175017 (2009).
- [30] O. Veblen and T. Y. Thomas, The geometry of paths, Trans. Am. Math. Soc. 25, 551 (1923).