

Light axion-like dark matter must be present during inflation

Luca Visinelli*

*The Oskar Klein Centre for Cosmoparticle Physics, Department of Physics, Stockholm University, AlbaNova, 10691 Stockholm, Sweden; Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, 10691 Stockholm, Sweden and University of Helsinki,**P.O. Box 64, Helsinki FI-00014 Finland*

(Received 29 March 2017; published 27 July 2017)

Axion-like particles (ALPs) might constitute the totality of the cold dark matter (CDM) observed. The parameter space of ALPs depends on the mass of the particle m and on the energy scale of inflation H_I , the latter being bound by the nondetection of primordial gravitational waves. We show that the bound on H_I implies the existence of a mass scale $\bar{m}_\chi = 10 \text{ neV} - 0.5 \text{ peV}$, depending on the ALP susceptibility χ , such that the energy density of ALPs of mass smaller than \bar{m}_χ is too low to explain the present CDM budget, if the ALP field has originated after the end of inflation. This bound affects ultra-light axions (ULAs), which have recently regained popularity as CDM candidates. Light ($m < m_\chi$) ALPs can then be CDM candidates only if the ALP field has already originated during the inflationary period, in which case the parameter space is constrained by the nondetection of axion isocurvature fluctuations. We comment on the effects on these bounds from additional physics beyond the standard model, besides ALPs.

DOI: [10.1103/PhysRevD.96.023013](https://doi.org/10.1103/PhysRevD.96.023013)**I. INTRODUCTION**

In the era of precision cosmology, the cold dark matter (CDM) budget in our Universe has been established at about 84% of the total matter in the Universe, yet its composition remains unknown. Among the proposed hypothetical particles which could address this fundamental question is the QCD axion [1,2], the quantum of the axion field arising from the spontaneous breaking of a U(1) symmetry conjectured by Peccei and Quinn (PQ [3,4]) to solve the strong-CP problem in quantum chromodynamics (QCD). The symmetry breaking occurs at a yet unknown energy scale f_a , which is constrained by measurements to be much larger than the electroweak energy scale [5]. The mass of the QCD axion at zero temperature m_0 is related to the axion energy scale f_a by $m_0 f_a = \Lambda_a^2$, where the energy scale Λ_a is related to the QCD parameter Λ_{QCD} . Realistic “invisible” axion models introduce new particles that further extend the standard model: examples include the coupling of the axion to heavy quarks [6,7] or to a Higgs doublet [8,9].

The history and the properties of axions produced in the early Universe depend on the relative magnitude of the energy scale f_a compared to the inflation energy scale H_I [10–19]. In fact, if $f_a > H_I/2\pi$, the breaking of the U(1)_{PQ} symmetry occurs before reheating begins and axions must be present during inflation, while, if $f_a < H_I/2\pi$, the axion field originates after the end of inflation. Measurements of the CMB properties constrain the parameter space of the axion, including the scale of inflation H_I and axion isocurvature fluctuations. See Refs. [20–27] for reviews of the QCD axion.

Besides the QCD axion, other axion-like particles (ALPs) arise from various ultraviolet completion models, in which additional U(1) symmetries which are spontaneously broken are introduced, as well as some other underlying physics. In fact, although the ALP mass might share a common origin with the QCD axion, it is possible for these particle not to be related to the dynamics of the gauge fields whatsoever. Examples include “accidental” axions [28–34] and axions from string theory [19,35–42]. See also Ref. [43] for the effects of wormholes to the QCD axion potential. In all these scenarios, two energy scales emerge: the symmetry-breaking scale Λ and the ALP decay constant f . Similarly to the QCD axion, the ALP field acquires a mass $m \sim \Lambda^2/f$, so that, contrarily to the QCD axion, the mass m and the energy scale f can be treated as independent parameters. An interesting proposed ALP is the ultra-light axion (ULA), of mass $m_{\text{ULA}} \approx 10^{-22} \text{ eV}$ [44–55]. Such a light axion, recently revised in Refs. [56,57], would have a wavelength of astrophysical scale $\lambda \sim 1 \text{ kpc}$ and could possibly address some controversies arising when treating small scales in the standard Λ CDM cosmology, namely the missing satellites and the cusp-core problems (see Ref. [58] for a review).

ALPs from global and accidental U(1) symmetries share a common cosmological history with the QCD axion and spectate inflation whenever $f > H_I/2\pi$. One of the main results of the present paper is to show that, in the opposite regime $f < H_I/2\pi$, the observational constraint on H_I coming from the Planck mission leads to a lower bound on the ALP mass, $m \gtrsim \bar{m}_\chi$, for some limiting mass \bar{m}_χ whose value depends on the ALP susceptibility χ . We find a numerical value $\bar{m}_\chi = 10 \text{ neV} - 0.5 \text{ peV}$, depending on the value of χ . This means that, if the CDM is discovered to be entirely composed of an ALP of mass $m < \bar{m}_\chi$, e.g. ULAs,

*luca.visinelli@fysik.su.se

such particles must be already present during inflation. Instead, if an ALP is discovered with $m > \bar{m}_\chi$, both cosmological origins are possible. We also show that, when $f > H_I/2\pi$ and the U(1) symmetry is never restored afterwards, the nondetection of axion isocurvature fluctuations by the Planck mission leads to an upper bound on the scale of inflation H_I , regardless of the ALP mass. Although this second result is quite straightforward to derive, it has never been stressed in the past literature.

The paper is organized as follows. In Sec. II we review the temperature dependence of the QCD axion mass, the ALP parameter space, and we derive the lower bound on the ALP mass. In Sec. III we show results for the ALP parameter space, assuming either a cosine or a harmonic potential. In Sec. IV, we discuss some exceptions to the computation used coming from the effects of some physics beyond the standard model, including the modification to the effective number of degrees of freedom, nonstandard cosmologies, or entropy dilution. Conclusions are drawn in Sec. V.

II. ALPS AND INFLATION

A. Reviewing the temperature dependence of the QCD axion mass

The QCD axion mass originates from nonperturbative effects during the QCD phase transition. At zero temperature, the axion gets a mass m_0 from mixing with the neutral pion [1],

$$m_0 = \frac{\Lambda_a^2}{f_a} = \frac{\sqrt{z} m_\pi f_\pi}{1+z} \frac{m_\pi f_\pi}{f_a}, \quad (1)$$

where $z = m_u/m_d$ is the ratio of the masses of the up and down quarks, m_π and f_π are respectively the mass and the energy scale of the pion, and f_a is the QCD axion energy scale. The energy scale Λ_a is proportional to the QCD scale Λ_{QCD} , so that the axion mass is tied to the underlying QCD theory. Using $z = 0.48(5)$, $m_\pi = 132$ MeV, and $f_\pi = 92.3$ MeV, the authors in Ref. [59] obtain $\Lambda_a = 75.5$ MeV, a value slightly smaller than what obtained in other work. For example, Ref. [24] obtains $\Lambda_a = 78$ MeV within the framework of the ‘‘interacting instanton liquid model’’, fixing the QCD scale to $\Lambda_{\text{QCD}} = 400$ MeV. Recently, more refined computations on the QCD lattice have become accessible [60–62].

When temperature-dependent effects become important, the QCD axion mass acquires a complicated dependence on the plasma temperature [63,64]. Here, we model such dependence as [24,65,66]

$$m_a(T) = \begin{cases} \frac{\alpha^2 \Lambda_{\text{QCD}}^2}{f_a} \left(\frac{\Lambda_{\text{QCD}}}{T}\right)^{\chi/2}, & \text{for } T \geq T_{\text{eff}}, \\ m_0, & \text{for } T < T_{\text{eff}}, \end{cases} \quad (2)$$

where χ is the QCD axion susceptibility and α is a numerical factor. At present, there is no general consensus on the

numerical value of the susceptibility, which depends on the particle content of the embedding theory [67,68], as well as the computational technique used [24,60,65]. Reference [24] obtains $\chi = 6.68$ and $\alpha = (1.68 \times 10^{-7})^{1/4} \approx 0.02$ while the methods in Refs. [15,63,64,69] predict $\chi = 8$ and

$$\alpha = \frac{\Lambda_a}{\Lambda_{\text{QCD}}} C^{1/2} \left(\frac{\Lambda_{\text{QCD}}}{200 \text{ MeV}}\right)^{1/4} \approx 0.03\text{--}0.05, \quad (3)$$

where $C = 0.018$, see Eq. (4) in Ref. [69]. In addition, we have introduced the temperature scale $T_{\text{eff}} = \Lambda_{\text{QCD}}(\alpha\Lambda_{\text{QCD}}/\Lambda_a)^{4/\chi}$ at which the two expressions in Eq. (2) match. This allows us to rewrite Eq. (2) as $m_a(T) = m_0 G(T)$, with the function

$$G(T) = \begin{cases} \left(\frac{T_{\text{eff}}}{T}\right)^{\chi/2}, & \text{for } T \geq T_{\text{eff}}, \\ 1, & \text{for } T < T_{\text{eff}}. \end{cases} \quad (4)$$

B. Observational constraints

The QCD axion, and more generally ALPs, are suitable CDM candidates in some region of the parameter space, provided that these particles are produced nonthermally. In the following, we assume that the totality of the observed CDM budget is in the form of ALPs of mass m . This is equivalent to demanding that the energy density in ALPs, here ρ_A , is equal to the present CDM energy density ρ_{CDM} . We write this requirement as

$$\Omega_A h^2 = \Omega_{\text{CDM}} h^2 = 0.1197 \pm 0.0022 \quad \text{at 68\% Confidence Level (CL)}, \quad (5)$$

where $\Omega_A = \rho_A/\rho_{\text{crit}}$ and $\Omega_{\text{CDM}} = \rho_{\text{CDM}}/\rho_{\text{crit}}$ are, respectively, the energy densities in ALPs and in the observed CDM [70], both given in units of the critical density $\rho_{\text{crit}} = 3H_0^2 M_{\text{Pl}}^2/8\pi$, with the Planck mass $M_{\text{Pl}} = 1.221 \times 10^{19}$ GeV and where h is the Hubble constant H_0 in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

Besides its mass, energy scale, and initial value of the misalignment angle, the ALP energy budget depends on the Hubble expansion rate H_I at the end of inflation, which is constrained from measurements on the scalar power spectrum $\Delta_{\mathcal{R}}^2(k_0)$ and the tensor-to-scalar ratio r_{k_0} at the pivotal scale k_0 as [71,72]

$$H_I < \frac{M_{\text{Pl}}}{4} \sqrt{\pi \Delta_{\mathcal{R}}^2(k_0) r_{k_0}} \sim 7 \times 10^{13} \text{ GeV}. \quad (6)$$

The numerical value of the bound has been computed by using the measurements at the wave number $k_0 = 0.05 \text{ Mpc}^{-1}$ [73–77]

$$\Delta_{\mathcal{R}}^2(k_0) = (2.215_{-0.079}^{+0.032}) \times 10^{-9}, \quad \text{at 68\% CL}, \quad (7)$$

$$r_{k_0} < 0.07, \quad \text{at 95\% CL.} \quad (8)$$

We finally comment on isocurvature perturbations. Quantum fluctuations imprint into all massless scalar field a present during inflation, with variance [78,79]

$$\langle |\delta a^2| \rangle = \left(\frac{H_I}{2\pi} \right)^2. \quad (9)$$

Primordial quantum fluctuations later develop into isocurvature perturbations [80], which modify the number density of axions, since the gauge invariant entropy perturbation is nonzero [81–83],

$$\mathcal{S}_a = \frac{\delta(n_a/s)}{n_a/s} \neq 0, \quad (10)$$

where s is the comoving entropy and n_a the axion number density. If all of the CDM is in axions, then we define [69,84,85]

$$\Delta_{S,A}^2 \equiv \langle |\mathcal{S}_a|^2 \rangle = \Delta_{\mathcal{R}}^2(k_0) \frac{\beta}{1-\beta}, \quad (11)$$

where the parameter β is constrained from Planck [73,74] at the scale $k_0 = 0.05 \text{ Mpc}^{-1}$ as

$$\beta \lesssim 0.037, \quad \text{at 95\% CL,} \quad (12)$$

independently on the ALP mass.

C. Constraining the ALP mass

We now consider the parameter space of ALPs produced through the vacuum realignment mechanism (VRM) [86–88], as revised in Appendix. Although, in principle, other mechanisms in addition to the VRM like the decay of topological defects produced at the PQ phase transition through the Kibble mechanism [89] and the decay of parent particles into ALPs might sensibly contribute to the present abundance of cold ALPs, we do not consider them here.

Similarly to what obtained for axions, we represent the ALP mass as $m(T) = mG(T)$, where m is a new parameter and $G(T)$ is given in Eq. (4). The ALP susceptibility χ might take any real non-negative value and is left here as a free parameter. An infinite susceptibility corresponds to the ALP mass abruptly jumping from zero to the value m at temperature T_{eff} ; any finite value of χ results in a smoother transition. ALPs from string theory or arising from accidental symmetries have $\chi = 0$. The ALP energy scale f is related to the ALP mass by $f = \Lambda^2/m$, where Λ is a new energy scale specified by an underlying theory. Finally, we write $T_{\text{eff}} = c\Lambda$, for some constant value c .

We review the non-thermal production of a cosmological population of ALPs from the misalignment mechanism in the Appendix, assuming that ALPs move in the potential

$$V(\theta) = f^2 m^2(T)(1 - \cos \theta), \quad (13)$$

where $\theta = a/f$ and a is the ALP field. We assume that, when the ALP field originates, the initial value of the misalignment angle is θ_i . The present value of the ALP energy density obtained from the misalignment mechanism is given in Eq. (A17),

$$\rho_A = \frac{\Lambda^4 G(T_1) g_S(T_0)}{2 g_S(T_1)} \left(\frac{T_0}{T_1} \right)^3 \langle \theta_i^2 \rangle, \quad (14)$$

where $\langle \theta_i^2 \rangle$ is the initial value of the misalignment angle squared, averaged over our Hubble volume, while the effective number of relativistic (“ R ”) and entropy (“ S ”) degrees of freedom are defined as [79]

$$g_R(T) = \sum_i \frac{15g_i}{\pi^4} \left(\frac{T_i}{T} \right)^4 \times \int_0^{+\infty} \frac{\sqrt{x^2 + y_i^2}}{\exp(\sqrt{x^2 + y_i^2}) + (-1)^{Q_i^f}} dx, \quad (15)$$

$$g_S(T) = \frac{3}{4} \sum_i \frac{15g_i}{\pi^4} \left(\frac{T_i}{T} \right)^3 \int_0^{+\infty} \frac{x^2 \sqrt{x^2 + y_i^2}}{\exp(\sqrt{x^2 + y_i^2}) + (-1)^{Q_i^f}} \times \left(1 + \frac{x^2}{3(x^2 + y_i^2)} \right) dx. \quad (16)$$

In the expressions above, T is the temperature of the plasma, and the sum runs over the i species considered, each with temperature T_i , mass m_i , $y_i \equiv m_i/T_i$, and $Q_i^f = 1$ ($Q_i^f = 0$) if i is a fermion (boson). Instead of computing the integrals in Eqs. (15)–(16), we have considered the parametrization in Refs. [24,90], where the effective number of degrees of freedom are approximated with a series of step functions, for temperatures up to $O(100 \text{ GeV})$.

In Eq. (14), we have introduced the initial value of the misalignment angle θ_i , which is the ALP field in units of f , and angle brackets define the average over all possible values of θ_i . In this scenario, θ_i takes different values within our Hubble horizon, so

$$\langle \theta_i^2 \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \theta_i^2 F(\theta_i) d\theta_i, \quad (17)$$

where the weighting function $F(\theta_i)$ has been thoroughly discussed in the literature [18,57,66,91–94]. Here, we take [57]

$$F(\theta_i) = \ln \left[\frac{e}{1 - (\theta_i/\pi)^4} \right], \quad (18)$$

which gives $\sqrt{\langle \theta_i^2 \rangle} = 2.45$.

Coherent oscillations in the ALP field begin at temperature T_1 given by $3H(T_1) = m$, see Eq. (A2) below, and the Hubble rate during radiation domination is

$$H(T) = \mathcal{A}(T) \frac{T^2}{3M_{\text{Pl}}}, \quad \mathcal{A}(T) = \sqrt{\frac{4\pi^3}{5} g_*(T)}. \quad (19)$$

The temperature T_1 at which the coherent oscillations in the ALP field begin is

$$T_1 = T_{\text{eff}} \begin{cases} \left(\frac{\hat{f}}{f}\right)^{\frac{2}{4+\gamma}}, & \text{for } f \leq \hat{f}, \\ \left(\frac{\hat{f}}{f}\right)^{\frac{1}{2}}, & \text{for } f > \hat{f}. \end{cases} \quad (20)$$

where we have defined the axion energy scale

$$\hat{f} \equiv \frac{M_{\text{Pl}}}{c^2 \mathcal{A}(T_1)}. \quad (21)$$

Inserting Eq. (20) into Eq. (14), we obtain the present ALP energy density as

$$\rho_A = \hat{\rho}_A \langle \theta_i^2 \rangle \left(\frac{m}{\hat{f}}\right)^{1/2} \begin{cases} \left(\frac{\hat{f}}{f}\right)^{\frac{16+3\gamma}{2(4+\gamma)}}, & \text{for } f \leq \hat{f}, \\ \left(\frac{\hat{f}}{f}\right)^2, & \text{for } f > \hat{f}, \end{cases} \quad (22)$$

where we have defined

$$\hat{\rho}_A = \frac{g_{*S}(T_0)}{g_{*S}(T_1)} \frac{\hat{f}}{2} \left(\frac{T_0}{c}\right)^3. \quad (23)$$

If the ALP field originates after inflation, the energy density is a function of the mass m and the ALP energy scale f only, but it does not depend on θ_i which is averaged out. Equating ρ_A in Eq. (22) with the CDM energy density $\rho_{\text{CDM}} = \Omega_{\text{CDM}} \rho_{\text{crit}}$ gives

$$f = \hat{f} \begin{cases} \left(\frac{\rho_{\text{CDM}}}{\hat{\rho}_A \langle \theta_i^2 \rangle}\right)^{\frac{8+2\gamma}{16+3\gamma}} \left(\frac{\hat{f}}{m}\right)^{\frac{4+\gamma}{16+3\gamma}}, & \text{for } f \leq \hat{f}, \\ \left(\frac{\rho_{\text{CDM}}}{\hat{\rho}_A \langle \theta_i^2 \rangle}\right)^{\frac{1}{2}} \left(\frac{\hat{f}}{m}\right)^{\frac{1}{4}}, & \text{for } f > \hat{f}. \end{cases} \quad (24)$$

For any value of m , Eq. (24) expresses the value of f for which the ALP explains the observed CDM budget.

We show that lighter ALPs cannot make the totality of the CDM when produced after the end of inflation. In facts, the region where $f < H_I/2\pi$ (which implies $f < \hat{f}$) is constrained by the bound on H_I expressed in Eq. (6), which leads to the lower bound on the ALP mass,

$$m \geq \bar{m}_\chi \equiv \hat{f} \left[\frac{64\pi}{\Delta_{\mathcal{R}}^2(k_0) r_{k_0}} \left(\frac{\hat{f}}{M_{\text{Pl}}}\right)^2 \right]^{\frac{16+3\gamma}{8+2\gamma}} \left(\frac{\rho_{\text{CDM}}}{\hat{\rho}_A \langle \theta_i^2 \rangle}\right)^2. \quad (25)$$

The numerical value of \bar{m}_χ depends on the susceptibility χ and on the value of the constant c in the model. Setting $c = 1$, we obtain the limiting cases $\bar{m}_0 = 10 \text{ neV}$ and

$\bar{m}_\infty = 0.5 \text{ peV}$. Axion theories where $m < \bar{m}_\chi$ must embed the axion production in the inflationary mechanism, as we discuss below. We remark that the bound in Eq. (25) only applies if the ALP field originated after the end of inflation, $f < H_I/2\pi$, and if the ALP field has originated from the breaking of a U(1) symmetry. In these scenarios, a Hubble volume contains a multitude of patches where the axion field has a different, random value. These patches are bound by topological defects which could decay and leave to an additional component of the cold ALP energy density. The inclusion of nonrelativistic ALPs from the decay of topological defects would increment their number density, potentially reducing the value of \bar{m}_χ by a couple of orders of magnitude. Here, we do not consider such contribution. Notice that the result in Eq. (25) does not depend on the value of Λ .

D. ALPs and inflation

ALPs of mass smaller than \bar{m}_χ can still be regarded as CDM candidates, although the related U(1) symmetry must have broken during the inflationary period, with the ALP energy scale satisfying $f > H_I/2\pi$. The cosmological properties of such ALPs would greatly differ from those described in the region $f < H_I/2\pi$, in particular no defects are present and a unique value of θ_i is singled out by the inflationary period within our Hubble volume. For example, consider the case of an ULA of mass $m_{\text{ULA}} = 10^{-22} \text{ eV}$, which is the mass scale proposed to solve some small-scale galactic problems [44–55] and recently has been vigorously reconsidered as a possible CDM candidate [56]. Since the mass scale m_{ULA} falls well within the limit excluded by Eq. (25), ULAs must have been produced during inflation to be the CDM, with a precise relation between the initial misalignment angle and the energy scale given by Eq. (24) with $\langle \theta_i^2 \rangle$ replaced by $\theta_i^2 F(\theta_i)$. The replacements account for the fact




FIG. 1. The energy scale f as a function of the ALP mass m , Eq. (24). We have chosen the initial misalignment angle $\theta_i = 1$ and different values of the susceptibility: $\chi = 0$ (blue solid line), $\chi = 8$ (green dotted line), $\chi = +\infty$ (red dashed line).

that the angle average $\langle \theta_i^2 \rangle$ singles out a uniform value for θ_i over the entire Hubble volume. In this scenario, we expect that the initial misalignment angle is of the order of one, with smaller values of θ_i still possible albeit disfavored. In Fig. 1, we show the value for f given in Eq. (24), as a function of the ALP mass m , for the value $\theta_i = 1$ and for different values of the ALP susceptibility: $\chi = 0$ (blue solid line), $\chi = 8$ (green dotted line), $\chi = +\infty$ (red dashed line). Values of f of the order of the GUT scale $f \sim 10^{15}$ GeV are expected for $m \sim 10^{-17} - 10^{-13}$ eV, while the ULA mass $m_{\text{ULA}} \sim 10^{-22}$ eV gives $f \sim 10^{17}$ GeV [56]. For higher values of the ALP mass, the spread among f for different values of χ widens.

III. FRAMING THE ALP PARAMETER SPACE

A. Cosine potential

We apply the expression for axion isocurvature fluctuations in Eq. (11) to the ALP scenario, to obtain [95]

$$\Delta_{S,A}^2 = \left(\frac{\partial \ln \rho_A}{\partial \theta_i} \right)^2 \langle \delta \theta_i^2 \rangle = \left(\frac{H_I \mathcal{F}(\theta_i)}{\pi \theta_i f} \right)^2, \quad (26)$$

where in the last step we have used Eq. (9), and where we defined the function

$$\mathcal{F}(x) = 1 + \frac{x F'(x)}{2F(x)}. \quad (27)$$

Results on the various bounds on the ALP parameter space are summarized in Fig. 2. Since we do not consider the contribution from the decay of topological defects, the parameter space of CDM ALPs depends on six quantities, f , θ_i , H_I , m , c , and χ . We show how the parameter space modifies when considering different values of the ALP mass: $m = 10^{-20}$ eV (top left), $m = 10^{-10}$ eV (top right), $m = 10^{-5}$ eV (bottom left), and $m = 10^{-3}$ eV (bottom right). For each panel, the line $f = H_I/2\pi$ separates the region where the axion is present during inflation (top-left) from the region where the axion field originates after inflation (bottom-right), for a fixed value $c = 1$. This line has to be thought as a qualitative bound between the two scenarios we will describe, since the exact details depend on the inflationary model, the preheating-reheating scenarios, and axion particle physics. The horizontal line labeled ‘‘ALP CDM’’ corresponds to the




FIG. 2. The ALP parameter space for different values of the ALP mass. We have assumed the ALP potential in Eq. (13). Top left: $m = 10^{-20}$ eV. Top right: $m = 10^{-10}$ eV. Bottom left: $m = 10^{-5}$ eV. Bottom right: $m = 10^{-3}$ eV. We have shown results for different values of the susceptibility: $\chi = 0$ (solid blue), $\chi = 8$ (dotted green), $\chi = +\infty$ (dashed red). The yellow region is excluded by CDM overproduction, $\rho_A > \rho_{\text{CDM}}$. The region labeled ‘‘Axion isocurvature fluctuations’’ is excluded below the curve shown. For clarity, we have shaded in yellow the region below the lowest of the three curves only. Horizontal lines show the values of f for which the ALP is the CDM for each ALP mass and for each value of χ . The green region is accessible, however the ALP is a subdominant CDM component, $\rho_A < \rho_{\text{CDM}}$. The bound labeled ‘‘Tensor modes’’ is derived from the nondetection of primordial gravitational waves, see Eq. (6).




FIG. 3. Same as Fig. 2, but for a harmonic potential in Eq. (30).

requirement that the primordial ALP condensate has started behaving like CDM at matter-radiation equality (See Ref. [96] for details),

$$f \gtrsim \frac{53 \text{ TeV}}{\pi} \sqrt{\frac{\text{eV}}{m}}. \quad (28)$$

We first discuss the scenario where $f > H_I/2\pi$. The region is bound by the nondetection of axion isocurvature fluctuations, obtained from Eq. (12) with the requirement that $\rho_A = \rho_{\text{CDM}}$. We plot the bound for three different values of the susceptibility: $\chi = 0$ (blue solid line), $\chi = 8$ (green dotted line), $\chi = +\infty$ (red dashed line). For clarity, we shade in yellow the region below the minimum of the three curves although we have to bear in mind that the whole parameter space below a curve of fixed χ has been ruled out. The change in the slope corresponds to the argument of the anharmonicity function $F(\theta_i)$ approaching π . For each value of χ , the horizontal lines in the allowed parameter space show the “natural” value of f for which $\rho_A = \rho_{\text{CDM}}$ and $\theta_i = 1$, as shown in Fig. 1. For $m = 10^{-20}$ eV, the natural value of the axion energy scale is of the order of $f \sim 10^{16}$ GeV, corresponding to the “ALP miracle” discussed in Ref. [56]. For smaller values of the ALP mass, the natural value of f lowers, and the spread among different values of χ widens, as shown in Fig. 1. The bound from isocurvature fluctuations steepens when θ_i decreases, and it is vertical for $\theta_i \ll 1$ and for $\chi = 0$, or for $f > \bar{f}$. We reformulate this constraint as an upper

bound on H_I for a given ALP theory, which is obtained by combining Eqs. (11), (24), and (26) as

$$H_I \leq \pi \hat{f} \left(\frac{\hat{f}}{m} \right)^{\frac{1}{4}} \sqrt{\frac{\rho_{\text{CDM}}}{\hat{\rho}_A} \frac{\beta}{1-\beta} \Delta_{\mathcal{R}}^2(k_0)} = 10^7 \text{ GeV} \sqrt[4]{\frac{\text{eV}}{m}}. \quad (29)$$

Isocurvature bounds have been used in the string axiverse realization discussed in Ref. [19], neglecting the dependence on the susceptibility and the anharmonic corrections in the potential. The presence of axion isocurvatures in the CMB, whose constrain on the power spectrum leads to Eq. (29), relies on the fact that the PQ symmetry has never been restored after the end of inflation. Caveats that allow to evade the bound from isocurvature fluctuations in Eq. (29) include the presence of more than one ALP [95] or by identifying the inflaton with the radial component of the PQ field [97]. This latter technique has been embedded into the SMASH model [98] where, for a decay scale $f \lesssim 4 \times 10^{16}$ GeV, the PQ symmetry is restored immediately after the end of inflation and isocurvature modes are absent, so that the bound in Eq. (29) does not apply.

In the second scenario $f < H_I/2\pi$, the axion is not present during inflation. In this scenario, a horizontal line gives the value of f for which the ALP is the CDM for a given value of the susceptibility. ALPs with an energy scale smaller than this value are a subdominant CDM component (green region, $\rho_A < \rho_{\text{CDM}}$), while values above

are excluded (yellow region, $\rho_A > \rho_{\text{CDM}}$). The constraint in Eq. (25) applies in this region of the parameter space, for some values of the ALP mass. For $m = 10^{-20}$ eV, which lies below the critical value \bar{m}_χ in Eq. (25), we always have $\rho_A < \rho_{\text{CDM}}$, so the region $f < H_I/2\pi$ is shaded with green. Larger values of the ALP mass allow for $\rho_A = \rho_{\text{CDM}}$ for some values of f and χ , avoiding the constraint in Eq. (25).

B. Harmonic potential

In Fig. 2, we have shown the parameter space of ALPs moving in the cosine ALP potential in Eq. (13), including the nonharmonic corrections through the function $F(\theta_i)$ in Eq. (18). However, the ALP potential can greatly differ from what expressed in Eq. (13). For example, in the presence of a monodromy [99–101], the degeneracy among the minima of the cosine potential is lifted by a quadratic potential, which might dominate the axion CDM potential [102]. We repeat the computation in the previous section for a harmonic potential, by switching off the nonharmonic corrections, setting $F(\theta_i) = 1$, considering the ALP moving in the quadratic potential

$$V_H(\theta) = \frac{1}{2}f^2m^2(T)\theta^2. \quad (30)$$

Inserting Eq. (22) into Eq. (26) for a harmonic potential to eliminate θ_i leads to a relation between H_I and f ,

$$f = \hat{f} \left[\left(\frac{\pi \hat{f}}{H_I} \right)^2 \frac{\rho_{\text{CDM}} \beta \Delta_{\mathcal{R}}^2(k_0)}{\hat{\rho}_A (1 - \beta)} \right]^{\frac{8+2\chi}{8+\chi}}. \quad (31)$$

We show results for the parameter space thus obtained in Fig. 3. Notice that the upper left panel ($m = 10^{-20}$ eV) qualitatively reproduces the results recently obtained in Ref. [57] when the anharmonic corrections are neglected in the isocurvature modes. Equation (29) describes the vertical blue line at the boundary of the region excluded by the nonobservation of isocurvature fluctuations.

IV. EFFECTS OF ADDITIONAL PHYSICS BEYOND THE STANDARD MODEL

Additional new physics might sensibly alter the axion parameter space presented in Fig. 2. Besides the QCD axion and other ALPs, examples of new physics not currently described within the framework of the standard model include additional particles whose presence modifies the effective number of degrees of freedom, or heavy scalar fields that might have dominated the Universe before the onset of radiation domination. We discuss some of the issues in the following. We focus on the case in which the axion mass is independent of temperature, since results can be easily generalized.

A. Effective number of degrees of freedom

The existence of particles that are still to be discovered would alter the effective number of relativistic and entropy degrees of freedom for temperatures larger than $T \gtrsim O(100)$ GeV. For example, the maximum number of effective relativistic degrees of freedom is 106.75 in the standard model, and 228.75 in the minimal supersymmetric standard model [79]. Setting $3H(T) = m$, with H given in Eq. (19) and $T = 1$ TeV, we obtain that corrections to $g_R(T)$ from physics beyond the standard model become important when $m \gtrsim 10^{-4}$ eV. We thus neglect these contributions when deriving the results in Sec. II C.

B. Nonstandard cosmological history

The content of the Universe for temperatures larger than $T_{\text{RH}} \gtrsim 4$ MeV is currently unknown, with lower bound being obtained from the requirement that the big-bang nucleosynthesis is achieved in a radiation-dominated cosmology [103–107]. However, for higher temperatures, the expansion rate of the Universe could have been dominated by some unknown form of energy, with an equation of state that differs from the one describing a relativistic fluid. A popular example is the early domination of a massive scalar field ϕ , emerging as a by-product of the decay of the inflaton field. In the following, we refer to this modified cosmology as being ϕ -dominated. The effect of a non-standard cosmological history might vary the present value of the axion energy density by orders of magnitude [93], depending on the equation of state for the fluid that dominates the expansion and the presence of an entropy dilution fact. In a nutshell, in a ϕ -dominated Universe the ALP begins to oscillate at a temperature T_1 that is different from what obtained in the standard picture, because of a different relation between temperature and time in the modified cosmology. Assuming that the equation of state of the ϕ field in the modified cosmology is $p = w\rho$ ($w = 1/3$ for radiation), for times t larger than the moment t_{RH} at which the Universe transitions from ϕ domination to radiation domination, the Hubble rate is

$$H = \frac{2}{3(w+1)t} = H_{\text{RH}} \left(\frac{T}{T_{\text{RH}}} \right)^{\frac{3(w+1)}{2}}, \quad (32)$$

where the last expression is valid only if the entropy density $s = g_S T^3$ in a comoving volume is conserved, we have neglected the contribution from the entropy degrees of freedom, and

$$H_{\text{RH}} = H(t_{\text{RH}}) = \mathcal{A}(T_{\text{RH}}) \frac{T_{\text{RH}}^2}{3M_{\text{Pl}}}. \quad (33)$$

We consider the temperature dependence of the ALP mass as $m(T) = m(T_{\text{eff}}/T)^{\chi/2}$, while the constant ALP mass case is obtained by setting $\chi = 0$. An early ϕ domination

modifies the temperature at which coherent oscillations begin, Eq. (20), as

$$T_1 = T_{\text{eff}} \left(\frac{\hat{f}_{\text{RH}}}{f} \right)^{\frac{2}{3(w+1)+\chi}} \left(\frac{T_{\text{RH}}}{T_{\text{eff}}} \right)^{\frac{3w-1}{3(w+1)+\chi}}. \quad (34)$$

where $\hat{f}_{\text{RH}} \equiv M_{\text{Pl}}/c^2 \mathcal{A}(T_{\text{RH}}) \approx \hat{f}$. The new value of T_1 modifies the present energy density, given by Eq. (14) when it is assumed entropy conservation from the onset of axion oscillations. The ALP energy density is

$$\rho_A = \hat{\rho}_A \left(\frac{m}{\hat{f}} \right)^{1/2} \langle \theta_i^2 \rangle \left(\frac{c\hat{\Lambda}}{T_{\text{RH}}} \right)^{\frac{(6+\chi)(3w-1)}{6(w+1)+2\chi}} \left(\frac{f}{\hat{f}} \right)^{\frac{2(16+3\chi)+(3w-1)(8+\chi)}{4(3(w+1)+\chi)}}, \quad (35)$$

where $\hat{\rho}_A$ has been defined in Eq. (23). Notice that, setting $w = 1/3$, we obtain the energy density given in the first line in Eq. (22). The axion energy scale for which the ALP is the CDM particle reads

$$f = \hat{f} \left(\frac{\rho_{\text{CDM}}}{\hat{\rho}_A \langle \theta_i^2 \rangle} \sqrt{\frac{\hat{f}}{m}} \right)^{\frac{4[3(w+1)+\chi]}{2(16+3\chi)+(3w-1)(8+\chi)}} \left(\frac{T_{\text{RH}}}{c\hat{\Lambda}} \right)^{\frac{2(6+\chi)(3w-1)}{2(16+3\chi)+(3w-1)(8+\chi)}}. \quad (36)$$

For a generic cosmological mode, the constraint in Eq. (25) for the region $f < H_1/2\pi$ modifies as

$$m \geq \hat{f} \left[\frac{64\pi}{\Delta_{\mathcal{R}}^2(k_0) r_{k_0}} \left(\frac{\hat{f}}{M_{\text{Pl}}} \right)^2 \right]^{\frac{2(16+3\chi)+(3w-1)(8+\chi)}{2[3w(8+\chi)+\chi]}} \times \left(\frac{\rho_{\text{CDM}}}{\hat{\rho}_A \langle \theta_i^2 \rangle} \right)^{\frac{4[3(w+1)+\chi]}{3w(8+\chi)+\chi}} \left(\frac{T_{\text{RH}}}{c\hat{f}} \right)^{\frac{2(6+\chi)(3w-1)}{3w(8+\chi)+\chi}}. \quad (37)$$

The latter expression depends on two additional parameters w and T_{RH} , and gives the result already obtained in Eq. (25) for $w = 1/3$.

For $w < 1/3$, Eq. (37) can be restated as a lower bound on the reheating temperature, valid when assuming that the ALPs considered make up the totality of the CDM observed and that coherent oscillations in the field began after inflation, in a ϕ -dominated cosmology. In the case of an early matter-dominated cosmology $w = 0$, the bound in Eq. (37) can be restated as a bound on the reheating temperature as

$$T_{\text{RH}} \geq c\hat{f} \left(\frac{\hat{f}}{m} \right)^{\frac{\chi}{2(6+\chi)}} \left[\frac{64\pi}{\Delta_{\mathcal{R}}^2(k_0) r_{k_0}} \left(\frac{\hat{f}}{M_{\text{Pl}}} \right)^2 \right]^{\frac{24+5\chi}{4(6+\chi)}} \left(\frac{\rho_{\text{CDM}}}{\hat{\rho}_A \langle \theta_i^2 \rangle} \right)^{\frac{6+2\chi}{6+\chi}}. \quad (38)$$

If the mass is not affected by nonperturbative effects and $\chi = 0$, like for accidental ALPs, the expression above becomes independent on m and yields the bound $T_{\text{RH}} \gtrsim 3 \text{ GeV}$, which is about three orders of magnitude

more stringent than what obtained in Refs. [103–107] using BBN considerations. We nevertheless stress that the bound in Eq. (38) can be easily evaded, given the strong assumptions under which it has been derived.

C. Dilution factor

Some scenarios predict a violation in the conservation of the total entropy in a comoving volume, sa^3 , due for example to the decay into lighter degrees of freedom of the ϕ field that dominates the Universe at that time. This is the case, for example, of a low-temperature reheating (LTR) stage [87,108–110], in which the Universe is dominated by a massive, decaying moduli field. In this situation, the relation between the scale factor and temperature changes from the simple relation $g_s^{1/3} T \sim 1/a$ to a generic relation $aT^\delta \sim \text{const}$, where δ is a new constant in the model. For example, $\delta = 8/3$ in the LTR scenario [111]. A different parametrization consists in assuming that a certain amount of entropy γ is produced during the decaying stage [108,112,113]. Either way, the effect of entropy dilution reduces the present energy density of axions in Eq. (22) by a factor γ , and the bound on the ALP mass in Eq. (25) is lowered. In general, we obtain the ALP energy density to be diluted by a factor $\rho_A \rightarrow \rho_A/\gamma$. If γ were to be independent on the ALP mass, we would get a reduction by $\bar{m}_\chi \rightarrow \bar{m}_\chi/\gamma^2$.

We compute the dilution factor in the LTR scenario as

$$\begin{aligned} \gamma &= \frac{g_S(T_{\text{RH}}) a_R^3 T_R^3}{g_S(T_1) a_1^3 T_1^3} = \frac{g_S(T_{\text{RH}})}{g_S(T_1)} \left(\frac{T_1}{T_{\text{RH}}} \right)^{3(\delta-1)} \\ &= \frac{g_S(T_{\text{RH}})}{g_S(T_1)} \left(\frac{T_{\text{eff}}}{T_{\text{RH}}} \right)^{3(\delta-1)} \left(\frac{\hat{f}}{f} \right)^{\frac{6(\delta-1)}{4+\chi}}, \end{aligned} \quad (39)$$

where in the last step we have used the expression for T_1 in Eq. (20) for the case $f \leq \hat{f}$. Since we expect oscillations to begin in the ϕ -dominated scenario, for which $T_1 > T_{\text{RH}}$, demanding $\delta > 1$ indeed leads to a dilution that is larger than one. For example, using $T_{\text{RH}} = 4 \text{ MeV}$ and $m = 10^{-5} \text{ eV}$ with $\delta = 8/3$ and $\chi = 0$, we obtain $\gamma \approx 10^{20}$. This large discrepancy with respect to the standard cosmology scenario has been used in Ref. [93] to dilute the energy density of the cosmological QCD axion, obtaining results that sensibly differ from the standard picture. Taking the expression for ρ_A in Eq. (22), we rephrase the bound in Eq. (25) when the dilution in Eq. (39) is added as

$$\begin{aligned} m &\geq \left(\frac{g_S(T_1)}{g_S(T_{\text{RH}})} \frac{g_R(T_1)}{g_R(T_0)} \frac{H_0^2}{T_0^3} \frac{48\Omega_{\text{CDM}}}{\langle \theta_i^2 \rangle \Delta_{\mathcal{R}}^2(k_0) r_{k_0}} \right)^{\frac{2}{3\delta-2}} \\ &\times \left(\frac{\mathcal{A}(T_1)}{M_{\text{Pl}}} \right)^{\frac{3(\delta-2)}{3\delta-2}} T_{\text{RH}}^{\frac{6(\delta-1)}{3\delta-2}} \sim 10^{-13} \text{ eV}. \end{aligned} \quad (40)$$

We have treated separately the effects due to the modified expansion rate and dilution to obtain the bounds in Eqs. (37)

and (40). A consistent derivation within a modified cosmology (say, LTR), has to consistently take into account both effects.

V. CONCLUSION

The present energy density of ALPs depends on both its mass m and the energy scale f . In general, these parameters can be tuned so that $\rho_A = \rho_{\text{CDM}}$. However, in models where the ALP field originates after inflation, we have shown in Sec. II C that the bound on the scale of inflation H_I from the nondetection of primordial gravitational waves leads to a minimum value of the ALP mass \bar{m}_χ below which the tuning of m and f is no longer possible. An ALP with mass $m < \bar{m}_\chi$ can still be a CDM candidate if it spectates inflation. In this latter scenario, the scale of inflation H_I is bound by the ALP mass through Eq. (29) which, although used in other work [13–19], has never been explicitly derived before. We have shown how these results affect the parameter space of the ALP for different values of the mass and of the susceptibility in Fig. 2 (cosine potential) and Fig. 3 (harmonic potential). Finally, we have commented on how results are affected by the presence of additional physics beyond the standard model, focusing on the modification of the effective number of degrees of freedom, nonstandard inflation and post-inflation cosmologies, and entropy dilution.

ACKNOWLEDGMENTS

The author would like to thank the anonymous referee for the careful read and the helpful suggestions, which led to a substantial improvement of the manuscript with respect to its original version, and Javier Redondo (U. Zaragoza) for the useful discussion. The author acknowledges support by Katherine Freese through a grant from the Swedish Research Council (Contract No. 638-2013-8993).

APPENDIX: REVIEW OF THE VACUUM REALIGNMENT MECHANISM

1. Equation of motion for the axion field

The ALP field originates from the breaking of the PQ symmetry at a temperature of the order of f . The equation of motion for the angular variable of the ALP field at any time is

$$\ddot{\theta} + 3H\dot{\theta} - \frac{\bar{\nabla}^2}{R^2}\theta + m^2(T)\sin\theta = 0, \quad (\text{A1})$$

where θ is the ALP field in units of f , $\bar{\nabla}$ is the Laplacian operator with respect to the physical coordinates \bar{x} , and R is the scale factor. To derive Eq. (A1), we have considered the simplest possible ALP potential $V(\theta) = f^2 m^2(T)(1 - \cos\theta)$. The mass term in the equation of motion becomes important when the Hubble rate is comparable to the axion mass,

$$H(T_1) = 3m(T_1), \quad (\text{A2})$$

whose solution gives the temperature T_1 when coherent oscillations begin. Setting the scale factor and the Hubble rate at temperature T_1 respectively as R_1 and H_1 , we rescale time t and scale factor R as $t \rightarrow H_1 t$ and $R \rightarrow R/R_1$, so that Eq. (A1) reads

$$\ddot{\theta} + 3H\dot{\theta} - \frac{\nabla^2}{R^2}\theta + 9g^2\sin\theta = 0, \quad (\text{A3})$$

where the Laplacian operator is written with respect to the comoving spatial coordinates $x = H_1 R_1 \bar{x}$ and $g = G(T)/G(T_1)$. We work in a radiation-dominated cosmology, where time and scale factor are related by $R \propto t^{1/2}$. Setting $\theta = \psi/R$, Eq. (A3) reads

$$\psi'' - \nabla^2\psi + 9g^2R^3\sin\left(\frac{\psi}{R}\right) = 0, \quad (\text{A4})$$

where a prime indicates a derivation with respect to R . Equation (A4) coincides with the results in Ref. [114], where the conformal time η is used as the independent variable in place of the scale factor R .

Taking the Fourier transform of the axion field as

$$\psi(\mathbf{x}) = \int e^{-iq\mathbf{x}}\psi(q), \quad (\text{A5})$$

we rewrite Eq. (A4) as

$$\psi'' + q^2\psi + 9g^2R^3\sin\left(\frac{\psi}{R}\right) = 0. \quad (\text{A6})$$

Equation (A6) expresses the full equation of motion for the axion field in the variable R , conveniently written to be solved numerically.

2. Approximate solutions of the equation of motion

Analytic solutions to Eq. (A6) can be obtained in the limiting regime $\psi/R \ll 1$, where Eq. (A6) reads

$$\psi'' + \kappa^2(R)\psi = 0, \quad (\text{A7})$$

with the wave number $\kappa^2(R) = q^2 + 9g^2R^2$. An approximate solution of Eq. (A7), valid in the adiabatic regime in which higher derivatives are neglected, is [114,115]

$$\psi = \psi_0(R) \exp\left(i \int^R \kappa(R') dR'\right), \quad (\text{A8})$$

where the amplitude ψ_0 is given by

$$|\psi_0(R)|^2 \kappa(R) = \text{const.} \quad (\text{A9})$$

Each term appearing in $\kappa(R)$ is the leading term in a particular regime of the evolution of the axion field. We analyze these approximate behavior in depths in the following.

- (i) Solution at early times, outside the horizon

At early times $t \sim R^2 \lesssim t_1$ prior to the onset of axion oscillations, the mass term in Eq. (A7) can be neglected since $m(T) \ll m(T_1)$. Defining the physical wavelength $\lambda = R/q$, we distinguish two different regimes in this approximation, corresponding to the evolution of the modes outside the horizon ($\lambda \gtrsim t$) or inside the horizon ($\lambda \lesssim t$). In the first case $\lambda \gtrsim t$, Eq. (A7) at early times reduces to $\psi'' = 0$, with solution ($\psi = R\phi$)

$$\phi(q, t) = \phi_1(q) + \frac{\phi_2(q)}{R} = \phi_1(q) + \frac{\phi_2(q)}{t^{1/2}}, \quad (\text{A10})$$

the first solution being a constant in time $\phi_1(q)$, while the second solution dropping to zero. The axion field for modes larger than the horizon is “frozen by causality” [115].

- (ii) Solution at early times, inside the horizon

Equation (A7) for modes that evolve inside the horizon $\lambda \lesssim t$ reduces to

$$\psi'' + q^2\psi = 0, \quad (\text{A11})$$

whose solution in a closed form, obtained through Eq. (A8) and $\phi = \psi/R$, reads

$$\phi \propto R^{-1} \exp(iqR). \quad (\text{A12})$$

The dependence of the amplitude $|\phi| \sim 1/R$ in Eq. (A12) is crucial, since it shows that the axion number density scales as cold matter,

$$n_A(q, t) \sim \frac{|\phi|^2}{\lambda} \sim \frac{1}{R^3}. \quad (\text{A13})$$

- (iii) Solution for the zero mode at the onset of oscillations

An approximate solution of Eq. (A7) for the zero-momentum mode $q = 0$, valid after the onset of axion oscillations when $t \sim t_1$, is obtained by setting

$$\kappa(R) \approx 3gR, \quad (\text{A14})$$

so that the adiabatic solution for ψ in Eq. (A8) in this slowly oscillating regime gives the axion number density

$$n_A^{\text{mis}}(R) = \frac{1}{2} m(R) f^2 \frac{|\psi|^2}{R^2} = n_1^{\text{mis}} \left(\frac{R}{R_1} \right)^{-3}, \quad (\text{A15})$$

where n_1^{mis} is the number density of axions from the misalignment mechanism at temperature T_1 ,

$$n_1^{\text{mis}} = \frac{1}{2} m(T_1) f^2 F(\theta_i) \theta_i^2, \quad (\text{A16})$$

and $F(\theta_i)$ is a function that accounts for neglecting the nonharmonic higher-order terms in the Taylor expansion of the sine function, see Eq. (18). Equation (A15) shows that the axion number density of the zero modes after the onset of axion oscillations scales as cold matter, with R^{-3} . The present ALP energy density is found by conservation of the comoving axion number density,

$$\rho_A = mn_1^{\text{mis}} \frac{s(T_0)}{s(T_1)} = mn_1^{\text{mis}} \frac{g_{*s}(T_0)}{g_{*s}(T_1)} \left(\frac{T_0}{T_1} \right)^3, \quad (\text{A17})$$

where $s(T)$ is the entropy density and $g_{*s}(T)$ is the number of degrees of freedom at temperature T .

-
- [1] S. Weinberg, *Phys. Rev. Lett.* **40**, 223 (1978).
 [2] F. Wilczek, *Phys. Rev. Lett.* **40**, 279 (1978).
 [3] R. D. Peccei and H. R. Quinn, *Phys. Rev. Lett.* **38**, 1440 (1977).
 [4] R. D. Peccei and H. R. Quinn, *Phys. Rev. D* **16**, 1791 (1977).
 [5] G. G. Raffelt, in *Axions: Theory, Cosmology, and Experimental Searches*, edited by M. Kuster, G. Raffelt, and B. Beltrán (Springer Berlin Heidelberg, Berlin, Heidelberg, 2008), p. 51.
 [6] J. E. Kim, *Phys. Rev. Lett.* **43**, 103 (1979).
 [7] M. Shifman, A. Vainshtein, and V. Zakharov, *Nucl. Phys.* **B166**, 493 (1980).
 [8] M. Dine, W. Fischler, and M. Srednicki, *Phys. Lett.* **104B**, 199 (1981).
 [9] A. R. Zhitnitsky, *Sov. J. Nucl. Phys.* **31**, 260 (1980).
 [10] A. D. Linde, *Phys. Lett. B* **201**, 437 (1988).
 [11] A. D. Linde, *Phys. Lett. B* **259**, 38 (1991).
 [12] M. S. Turner and F. Wilczek, *Phys. Rev. Lett.* **66**, 5 (1991).
 [13] F. Wilczek, [arXiv:hep-ph/0408167](https://arxiv.org/abs/hep-ph/0408167).
 [14] M. Tegmark, A. Aguirre, M. Rees, and F. Wilczek, *Phys. Rev. D* **73**, 023505 (2006).

- [15] M. P. Hertzberg, M. Tegmark, and F. Wilczek, *Phys. Rev. D* **78**, 083507 (2008).
- [16] B. Freivogel, *J. Cosmol. Astropart. Phys.* **03** (2010) 021.
- [17] K. J. Mack, *J. Cosmol. Astropart. Phys.* **07** (2011) 021.
- [18] L. Visinelli and P. Gondolo, *Phys. Rev. D* **80**, 035024 (2009).
- [19] B. S. Acharya, K. Bobkov, and P. Kumar, *J. High Energy Phys.* **11** (2010) 105.
- [20] G. G. Raffelt, in *Dark matter in cosmology, clocks and test of fundamental laws. Proceedings, 30th Rencontres de Moriond, 15th Moriond Workshop, Villars sur Ollon, Switzerland, 1995* (Editions Frontieres, Gif-Sur-Yvette, 1995), p. 159, [arXiv:hep-ph/9502358](https://arxiv.org/abs/hep-ph/9502358).
- [21] G. G. Raffelt, *Quantum theories and renormalization group in gravity and cosmology: Proceedings, 2nd International Conference, IRGAC 2006, Barcelona, Spain, 2006*, *J. Phys. A* **40**, 6607 (2007).
- [22] P. Sikivie, *Axions: Theory, cosmology, and experimental searches. Proceedings, 1st Joint ILIAS-CERN-CAST axion training, Geneva, Switzerland, 2005, Lect. Notes Phys.* **741**, 19 (2008).
- [23] J. E. Kim and G. Carosi, *Rev. Mod. Phys.* **82**, 557 (2010).
- [24] O. Wantz and E. P. S. Shellard, *Phys. Rev. D* **82**, 123508 (2010).
- [25] M. Kawasaki and K. Nakayama, *Annu. Rev. Nucl. Part. Sci.* **63**, 69 (2013).
- [26] D. J. E. Marsh, *Phys. Rep.* **643**, 1 (2016).
- [27] J. E. Kim, [arXiv:1703.03114](https://arxiv.org/abs/1703.03114).
- [28] K.-S. Choi, I.-W. Kim, and J. E. Kim, *J. High Energy Phys.* **03** (2007) 116.
- [29] K.-S. Choi, H. P. Nilles, S. Ramos-Sanchez, and P. K. S. Vaudrevange, *Phys. Lett. B* **675**, 381 (2009).
- [30] A. G. Dias, A. C. B. Machado, C. C. Nishi, A. Ringwald, and P. Vaudrevange, *J. High Energy Phys.* **06** (2014) 037.
- [31] T. Higaki and F. Takahashi, *J. High Energy Phys.* **07** (2014) 074.
- [32] J. E. Kim and D. J. E. Marsh, *Phys. Rev. D* **93**, 025027 (2016).
- [33] M. Redi and R. Sato, *J. High Energy Phys.* **05** (2016) 104.
- [34] L. Di Luzio, E. Nardi, and L. Ubaldi, *Phys. Rev. Lett.* **119**, 011801 (2017).
- [35] P. Svrcek and E. Witten, *J. High Energy Phys.* **06** (2006) 051.
- [36] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper, and J. March-Russell, *Phys. Rev. D* **81**, 123530 (2010).
- [37] M. Dine, G. Festuccia, J. Kehayias, and W. Wu, *J. High Energy Phys.* **01** (2011) 012.
- [38] A. Ringwald, *Proceedings, 18th International Symposium on Particles, Strings and Cosmology (PASCOS 2012): Merida, Yucatan, Mexico, J. Phys. Conf. Ser.* **485**, 012013 (2014).
- [39] J. E. Kim, *Phys. Rev. Lett.* **111**, 031801 (2013).
- [40] T. C. Bachlechner, C. Long, and L. McAllister, *J. High Energy Phys.* **12** (2015) 042.
- [41] J. Halverson, C. Long, and P. Nath, [arXiv:1703.07779](https://arxiv.org/abs/1703.07779).
- [42] M. J. Stott, D. J. E. Marsh, C. Pongkitivanichkul, L. C. Price, and B. S. Acharya, [arXiv:1706.03236](https://arxiv.org/abs/1706.03236).
- [43] R. Alonso and A. Urbano, [arXiv:1706.07415](https://arxiv.org/abs/1706.07415).
- [44] M. Baldeschi, G. Gelmini, and R. Ruffini, *Phys. Lett. B* **122**, 221 (1983).
- [45] M. Membrado, A. F. Pacheco, and J. Sañudo, *Phys. Rev. A* **39**, 4207 (1989).
- [46] W. H. Press, B. S. Ryden, and D. N. Spergel, *Phys. Rev. Lett.* **64**, 1084 (1990).
- [47] S.-J. Sin, *Phys. Rev. D* **50**, 3650 (1994).
- [48] S. U. Ji and S. J. Sin, *Phys. Rev. D* **50**, 3655 (1994).
- [49] J.-w. Lee and I. G. Koh, *Phys. Rev. D* **53**, 2236 (1996).
- [50] F. S. Guzmán and T. Matos, *Classical Quantum Gravity* **17**, L9 (2000).
- [51] V. Sahni and L. Wang, *Phys. Rev. D* **62**, 103517 (2000).
- [52] P. J. E. Peebles, *Astrophys. J.* **534**, L127 (2000).
- [53] J. Goodman, *New Astron.* **5**, 103 (2000).
- [54] T. Matos and L. A. Ureña-López, *Classical Quantum Gravity* **17**, L75 (2000).
- [55] W. Hu, R. Barkana, and A. Gruzinov, *Phys. Rev. Lett.* **85**, 1158 (2000).
- [56] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, *Phys. Rev. D* **95**, 043541 (2017).
- [57] A. Diez-Tejedor and D. J. E. Marsh, [arXiv:1702.02116](https://arxiv.org/abs/1702.02116).
- [58] D. H. Weinberg, J. S. Bullock, F. Governato, R. Kuzio de Naray, and A. H. G. Peter, *Sackler Colloquium: Dark Matter Universe: On the Threshold of Discovery Irvine, USA, 2012, Proc. Nat. Acad. Sci.* **112**, 12249 (2014).
- [59] G. Grilli di Cortona, E. Hardy, J. Pardo Vega, and G. Villadoro, *J. High Energy Phys.* **01** (2016) 034.
- [60] S. Borsanyi, M. Dierigl, Z. Fodor, S. D. Katz, S. W. Mages, D. Nogradi, J. Redondo, A. Ringwald, and K. K. Szabo, *Phys. Lett. B* **752**, 175 (2016).
- [61] S. Borsanyi *et al.*, *Nature (London)* **539**, 69 (2016).
- [62] P. Petreczky, H.-P. Schadler, and S. Sharma, *Phys. Lett. B* **762**, 498 (2016).
- [63] D. J. Gross, R. D. Pisarski, and L. G. Yaffe, *Rev. Mod. Phys.* **53**, 43 (1981).
- [64] P. Fox, A. Pierce, and S. D. Thomas, [arXiv:hep-th/0409059](https://arxiv.org/abs/hep-th/0409059).
- [65] M. S. Turner, *Phys. Rev. D* **33**, 889 (1986).
- [66] K. J. Bae, J.-H. Huh, and J. E. Kim, *J. Cosmol. Astropart. Phys.* **09** (2008) 005.
- [67] H. Davoudiasl, *Phys. Lett. B* **646**, 172 (2007).
- [68] H. Davoudiasl and C. W. Murphy, *Phys. Rev. Lett.* **118**, 141801 (2017).
- [69] M. Beltran, J. Garcia-Bellido, and J. Lesgourgues, *Phys. Rev. D* **75**, 103507 (2007).
- [70] P. A. R. Ade *et al.* (Planck), *Astron. Astrophys.* **594**, A13 (2016).
- [71] D. Lyth, *Phys. Lett.* **147B**, 403 (1984).
- [72] D. H. Lyth and E. D. Stewart, *Phys. Lett. B* **283**, 189 (1992).
- [73] P. A. R. Ade *et al.* (Planck), *Astron. Astrophys.* **571**, A16 (2014).
- [74] P. A. R. Ade *et al.* (Planck), *Astron. Astrophys.* **571**, A22 (2014).
- [75] D. Barkats *et al.* (BICEP1), *Astrophys. J.* **783**, 67 (2014).
- [76] P. A. R. Ade *et al.* (BICEP2, Planck), *Phys. Rev. Lett.* **114**, 101301 (2015).
- [77] P. A. R. Ade *et al.* (BICEP2, Keck Array), *Phys. Rev. Lett.* **116**, 031302 (2016).
- [78] D. H. Lyth, *Phys. Lett. B* **236**, 408 (1990).
- [79] E. W. Kolb and M. S. Turner, *The Early Universe*, (Addison-Wesley, Redwood City, CA, 1990).

- [80] T. Kobayashi, R. Kurematsu, and F. Takahashi, *J. Cosmol. Astropart. Phys.* **09** (2013) 032.
- [81] M. Axenides, R. Brandenberger, and M. Turner, *Phys. Lett.* **126B**, 178 (1983).
- [82] A. D. Linde, *Phys. Lett.* **158B**, 375 (1985).
- [83] D. Seckel and M. S. Turner, *Phys. Rev. D* **32**, 3178 (1985).
- [84] P. Crotty, J. Garcia-Bellido, J. Lesgourgues, and A. Riazuelo, *Phys. Rev. Lett.* **91**, 171301 (2003).
- [85] M. Beltran, J. Garcia-Bellido, J. Lesgourgues, A. R. Liddle, and A. Slosar, *Phys. Rev. D* **71**, 063532 (2005).
- [86] L. F. Abbott and P. Sikivie, *Phys. Lett.* **120B**, 133 (1983).
- [87] M. Dine and W. Fischler, *Phys. Lett.* **120B**, 137 (1983).
- [88] J. Preskill, M. B. Wise, and F. Wilczek, *Phys. Lett.* **120B**, 127 (1983).
- [89] T. W. B. Kibble, *J. Phys. A* **9**, 1387 (1976).
- [90] T. S. Coleman and M. Roos, *Phys. Rev. D* **68**, 027702 (2003).
- [91] D. H. Lyth, *Phys. Rev. D* **45**, 3394 (1992).
- [92] K. Strobl and T. J. Weiler, *Phys. Rev. D* **50**, 7690 (1994).
- [93] L. Visinelli and P. Gondolo, *Phys. Rev. D* **81**, 063508 (2010).
- [94] P. Gondolo and L. Visinelli, *Phys. Rev. Lett.* **113**, 011802 (2014).
- [95] N. Kitajima and F. Takahashi, *J. Cosmol. Astropart. Phys.* **01** (2015) 032.
- [96] P. Arias, D. Cadamuro, M. Goodsell, J. Jaeckel, J. Redondo, and A. Ringwald, *J. Cosmol. Astropart. Phys.* **06** (2012) 013.
- [97] M. Fairbairn, R. Hogan, and D. J. E. Marsh, *Phys. Rev. D* **91**, 023509 (2015).
- [98] G. Ballesteros, J. Redondo, A. Ringwald, and C. Tamarit, [arXiv:1610.01639](https://arxiv.org/abs/1610.01639).
- [99] E. Silverstein and A. Westphal, *Phys. Rev. D* **78**, 106003 (2008).
- [100] L. McAllister, E. Silverstein, and A. Westphal, *Phys. Rev. D* **82**, 046003 (2010).
- [101] N. Kaloper and L. Sorbo, *Phys. Rev. Lett.* **102**, 121301 (2009).
- [102] J. Jaeckel, V. M. Mehta, and L. T. Witkowski, *J. Cosmol. Astropart. Phys.* **01** (2017) 036.
- [103] M. Kawasaki, K. Kohri, and N. Sugiyama, *Phys. Rev. Lett.* **82**, 4168 (1999).
- [104] M. Kawasaki, K. Kohri, and N. Sugiyama, *Phys. Rev. D* **62**, 023506 (2000).
- [105] S. Hannestad, *Phys. Rev. D* **70**, 043506 (2004).
- [106] K. Ichikawa, M. Kawasaki, and F. Takahashi, *Phys. Rev. D* **72**, 043522 (2005).
- [107] F. D. Bernardis, L. Pagano, and A. Melchiorri, *Astropart. Phys.* **30**, 192 (2008).
- [108] P. J. Steinhardt and M. S. Turner, *Phys. Lett.* **129B**, 51 (1983).
- [109] M. S. Turner, *Phys. Rev. D* **28**, 1243 (1983).
- [110] R. J. Scherrer and M. S. Turner, *Phys. Rev. D* **31**, 681 (1985).
- [111] G. F. Giudice, E. W. Kolb, and A. Riotto, *Phys. Rev. D* **64**, 023508 (2001).
- [112] G. Lazarides, C. Panagiotakopoulos, and Q. Shafi, *Phys. Lett. B* **192**, 323 (1987).
- [113] G. Lazarides, R. K. Schaefer, D. Seckel, and Q. Shafi, *Nucl. Phys.* **B346**, 193 (1990).
- [114] E. W. Kolb and I. I. Tkachev, *Phys. Rev. D* **49**, 5040 (1994).
- [115] P. Sikivie, in *Axions: Theory, Cosmology, and Experimental Searches*, edited by M. Kuster, G. Raffelt, and B. Beltrán (Springer Berlin Heidelberg, Berlin, Heidelberg, 2008), p. 19.