

New constraints on dark matter production during kinationKayla Redmond^{*} and Adrienne L. Erickcek[†]*Department of Physics and Astronomy, University of North Carolina at Chapel Hill,
Phillips Hall CB3255, Chapel Hill, North Carolina 27599, USA*

(Received 27 March 2017; published 15 August 2017)

Our ignorance of the period between the end of inflation and the beginning of big bang nucleosynthesis limits our understanding of the origins and evolution of dark matter. One possibility is that the Universe's energy density was dominated by a fast-rolling scalar field while the radiation bath was hot enough to thermally produce dark matter. We investigate the evolution of the dark matter density and derive analytic expressions for the dark matter relic abundance generated during such a period of kination. Kination scenarios in which dark matter does not reach thermal equilibrium require $\langle\sigma v\rangle < 2.7 \times 10^{-38} \text{ cm}^3 \text{ s}^{-1}$ to generate the observed dark matter density while allowing the Universe to become radiation dominated by a temperature of 3 MeV. Kination scenarios in which dark matter does reach thermal equilibrium require $\langle\sigma v\rangle > 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ in order to generate the observed dark matter abundance. We use observations of dwarf spheroidal galaxies by the Fermi Gamma-Ray Telescope and observations of the Galactic Center by the High Energy Stereoscopic System to constrain these kination scenarios. Combining the unitarity constraint on $\langle\sigma v\rangle$ with these observational constraints sets a lower limit on the temperature at which the Universe can become radiation dominated following a period of kination if $\langle\sigma v\rangle > 3 \times 10^{-31} \text{ cm}^3 \text{ s}^{-1}$. This lower limit is between 0.05 and 1 GeV, depending on the dark matter annihilation channel.

DOI: [10.1103/PhysRevD.96.043511](https://doi.org/10.1103/PhysRevD.96.043511)**I. INTRODUCTION**

The expansion history of the Universe before big bang nucleosynthesis (BBN) is uncertain. The fact that the primordial curvature perturbation spectrum is almost scale invariant strongly suggests that shortly after the big bang, the Universe experienced a period of inflation [1–3]. The energy scale of inflation is not known, but it is generally assumed to be greater than 10^{10} GeV. The successful BBN prediction of the abundances of light elements only requires that the Universe be radiation dominated at a temperature of 3 MeV [4–8]. Thus, there is a gap in the cosmological record between the theorized energy scale of inflation and 3 MeV.

In the simplest model, inflation is powered by a scalar field defined as the inflaton, and the Universe becomes radiation dominated when the inflaton decays into relativistic particles [9–11]. Another possibility is that a different scalar field dominates the Universe after inflation [12]. If either of these scalar fields oscillates around the minimum of their potential, it behaves like a pressureless fluid and the Universe would be in an early-matter-dominated era [12–23]. An alternative scenario is that a fast-rolling scalar field (a kination) dominates the energy density of the Universe prior to the onset of radiation domination. When the kination's energy density is dominant, the Universe is said to be in a period of kination [24–26]. Kination was initially proposed as an inflationary model that does not

require the complete conversion of the false vacuum energy into radiation to initiate the onset of radiation domination [24]. Kination also facilitates baryogenesis; if the electro-weak phase transition occurs during kination, then baryogenesis is possible during a second-order phase transition [25]. Finally, if the kination's potential energy becomes dominant at very late times, it can accelerate the expansion of the Universe and mimic the effects of a cosmological constant [26–30].

The uncertainties in the thermal history of the Universe prior to BBN limit our understanding of the origins of dark matter [12,15,16,20,23,31–34]. We study the effects of kination on the thermal production of dark matter. We derive analytic expressions for the dark matter relic abundance generated during kination and confirm that our analytic results match the numeric solutions to the Boltzmann equation. Our relic abundance expressions depend on the dark matter mass m_χ , the velocity-averaged dark matter annihilation cross section $\langle\sigma v\rangle$, and the temperature at which the Universe becomes radiation dominated, T_{RH} . Our analytic expressions allow us to solve for the $\langle\sigma v\rangle$ values that will generate the observed dark matter abundance. We determine that in order to achieve the observed dark matter abundance, kination models in which dark matter reaches thermal equilibrium require $\langle\sigma v\rangle$ values that would underproduce dark matter during radiation domination. In contrast, kination models in which dark matter does not reach thermal equilibrium require $\langle\sigma v\rangle$ values that would overproduce dark matter during radiation domination. Using the most recent constraints on m_χ and $\langle\sigma v\rangle$ from Fermi-LAT PASS-8 observations of dwarf spheroidal galaxies [35] and

^{*}kayla.jaye@unc.edu[†]erickcek@physics.unc.edu

High Energy Stereoscopic System (H.E.S.S.) observations of the Galactic center [36], we constrain T_{RH} for kination scenarios where dark matter reaches thermal equilibrium.

Prior investigations of dark matter production during kination have focused on specific kination potentials. References [37–41] investigated how the relic abundance of dark matter is affected if the kination has an exponential potential, while Refs. [42,43] studied kination models where the kination has an inverse power-law potential. While these prior works did place constraints on dark matter parameters, those constraints were dependent on the specified kination potential. Our relic abundance expressions are independent of the kination potential, and our constraints on m_χ , T_{RH} , and $\langle\sigma v\rangle$ are applicable to all kination scenarios in which dark matter is a thermal relic. Furthermore, improvements in the observational constraints on $\langle\sigma v\rangle$ over the past six years allow us to place tighter constraints than previous works. We determine that kination scenarios in which dark matter reaches thermal equilibrium have a minimum allowed reheat temperature between 0.05 and 1 GeV, depending on the dark matter annihilation channel.

In Sec. II A, we discuss the evolution equations that govern the thermal production of dark matter during kination. In Secs. II B and II C, we present analytic derivations of the dark matter relic abundance for dark matter that does and does not reach thermal equilibrium. In Sec. III, we use observational data from Fermi-LAT and H.E.S.S. to constrain m_χ , T_{RH} , and $\langle\sigma v\rangle$. In Sec. IV, we summarize our results. Natural units ($\hbar = c = k_B = 1$) are used throughout this work.

II. THERMAL DARK MATTER DURING KINATION

A. Kination cosmology

The scenario we consider consists of a fast-rolling scalar field (the kination) that dominates the energy density of the Universe prior to BBN. The kination’s energy density is dominated by its kinetic energy, meaning that the kination’s energy density equals its pressure and that the equation of state parameter is $w = 1$. Therefore, the kination’s energy density scales as a^{-6} , where a is the scale factor, and will eventually become subdominant to radiation, whose energy density scales as a^{-4} . Reheating is defined as the point at which the radiation energy density becomes the dominant component of the Universe. It is important to note, however, that during kination, the temperature of the radiation bath is higher than the temperature at reheating. Therefore, it is possible to thermally produce dark matter prior to the onset of radiation domination.

We consider three energy density components during kination: dark matter, radiation, and the kination. The evolution of these energy densities are governed by three free parameters: the dark matter mass m_χ , the reheat

temperature T_{RH} , and the velocity-averaged dark matter annihilation cross section $\langle\sigma v\rangle$. Throughout this work, we assume s -wave dark matter annihilation. In kination scenarios, T_{RH} is the temperature at which the radiation energy density equals the kination energy density. We assume that the kination does not decay nor interact with dark matter or radiation (see Ref. [39] for an analysis of decaying kination cosmologies). Radiation and dark matter on the other hand are thermally coupled via pair production and annihilation. Therefore, the equations for the energy density of the scalar field ρ_ϕ , the radiation energy density ρ_r , and the dark matter number density n_χ are

$$\frac{d}{dt}\rho_\phi = -6H\rho_\phi, \quad (1a)$$

$$\frac{d}{dt}n_\chi = -3Hn_\chi - \langle\sigma v\rangle(n_\chi^2 - n_{\chi,\text{eq}}^2), \quad (1b)$$

$$\frac{d}{dt}\rho_r = -4H\rho_r + \langle\sigma v\rangle E_\chi(n_\chi^2 - n_{\chi,\text{eq}}^2), \quad (1c)$$

where $\langle E_\chi\rangle = \rho_\chi/n_\chi$ is the average energy of a dark matter particle and $n_{\chi,\text{eq}}$ is the number density of dark matter particles in thermal equilibrium.¹ For a dark matter particle with mass m_χ and internal degrees of freedom g_χ within a thermal bath of temperature T ,

$$n_{\chi,\text{eq}} = \frac{g_\chi}{2\pi^2} \int_{m_\chi}^{\infty} \frac{\sqrt{E^2 - m_\chi^2}}{e^{E/T} + 1} E dE. \quad (2)$$

When evaluating the average energy of a dark matter particle, we make the approximation that $\langle E_\chi\rangle \simeq \sqrt{m_\chi^2 + (3.151T)^2}$, which matches ρ_χ/n_χ to within 10%.

Figure 1 shows the evolution of the kination, radiation, and dark matter densities obtained by numerically solving Eq. (1). Initially, the kination’s energy density is dominant, but since it scales away more quickly than the radiation energy density, it eventually becomes subdominant. Figure 1 shows that the radiation energy density scales as a^{-4} and is unaffected by dark matter annihilation or pair production. Since ρ_r and ρ_ϕ evolve independently of n_χ , we can solve for their evolution analytically. We then use these solutions to numerically solve Eq. (1b) and calculate the dark matter relic abundance.

To accurately describe the evolution of ρ_r , we need to take into account the energy injection that occurs when Standard Model particles become nonrelativistic. When a particle species becomes nonrelativistic, its entropy is transferred to the remaining relativistic particles. Entropy is conserved during kination; therefore, the universal entropy

¹Throughout this work, we assume that the dark matter is composed of Majorana particles, and therefore $\chi = \bar{\chi}$.




FIG. 1. The density evolution of the kinaton, radiation, and dark matter. In this figure, $m_\chi = 10^4$ GeV, and reheating occurs when $a/a_I = 2.7 \times 10^4$; the reheat temperature is 2 GeV. The two solid curves show the evolution of ρ_χ for the two values of $\langle\sigma v\rangle$ that produce the observed dark matter density: $\Omega_\chi h^2 = 0.12$ [44]. The top solid curve corresponds to the freeze-out scenario with $\langle\sigma v\rangle = 7.5 \times 10^{-25} \text{ cm}^3 \text{ s}^{-1}$, whereas the bottom solid curve corresponds to the freeze-in scenario with $\langle\sigma v\rangle = 6.7 \times 10^{-46} \text{ cm}^3 \text{ s}^{-1}$. The dotted line shows the equilibrium dark matter density, $\rho_{\chi,\text{eq}} = \langle E_\chi \rangle n_{\chi,\text{eq}}$.

sa^3 must remain constant, where s is the entropy density: $s \equiv (2\pi^2/45)T^3 g_{*s}(T)$, and $g_{*s}(T)$ is the effective number of degrees of freedom that contribute to the entropy density. Due to the conservation of entropy, radiation cools during kination according to the same proportionality as during radiation domination: $T \propto g_{*s}(T)^{-1/3} a^{-1}$.

To evaluate the temperature of the radiation bath, we set a maximum temperature of T_{MAX} at which $\rho_\chi = 0$. We set $T_{\text{MAX}} = 8m_\chi$ to ensure that if the dark matter is capable of reaching thermal equilibrium, it will have adequate time to do so. If the dark matter cannot reach thermal equilibrium, setting $T_{\text{MAX}} = 8m_\chi$ ensures there will be enough time for maximal pair production. Therefore, the dark matter relic abundance will not be sensitive to T_{MAX} . Using T_{MAX} , we construct an expression for the temperature evolution during kination that accounts for changes in $g_{*s}(T)$:

$$T = T_{\text{MAX}} \left[\frac{g_{*s}(T_{\text{MAX}})}{g_{*s}(T)} \right]^{1/3} \frac{a_I}{a}, \quad (3)$$

where a_I is the scale factor value when $T = T_{\text{MAX}}$.

The final step in evaluating ρ_r is to connect Eq. (3) and the definition of ρ_r . The radiation energy density is $\rho_r \equiv (\pi^2/30)g_*(T)T^4$, where $g_*(T)$ is the number of relativistic degrees of freedom at temperature T . Using this definition of ρ_r and Eq. (3), we see that the evolution of ρ_r during kination is

$$\rho_r = \frac{\pi^2}{30} g_*(T) T_{\text{MAX}}^4 \left[\frac{g_{*s}(T_{\text{MAX}})}{g_{*s}(T)} \right]^{4/3} \left(\frac{a_I}{a} \right)^4. \quad (4)$$




FIG. 2. The evolution of the comoving dark matter number density and equilibrium number density with $T_{\text{RH}} = 20$ GeV, $m_\chi = 3000$ GeV, and $\langle\sigma v\rangle = 10^{-32} \text{ cm}^3 \text{ s}^{-1}$. The vertical line represents the point of reheating at $a_{\text{RH}}/a_I = 800$. The solid horizontal line shows the comoving number density at the point of freeze-out solved from $H(T_F) = \langle\sigma v\rangle n_{\chi,\text{eq}}$. This figure demonstrates that dark matter annihilations after freeze-out significantly decrease the dark matter number density.

Next, we analytically solve for ρ_ϕ . Solving Eq. (1a) yields $\rho_\phi = \rho_{\phi,I}(a_I/a)^6$, where $\rho_{\phi,I}$ is ρ_ϕ when $a = a_I$. By defining a_{RH} as the scale factor value at the onset of radiation domination we see that ρ_ϕ evaluated at reheating equals $\rho_{\phi,I}(a_I/a_{\text{RH}})^6$. Using Eq. (4), we can evaluate ρ_r at reheating. Considering that at reheating $\rho_\phi = \rho_r$, this implies that

$$\rho_{\phi,I} = \frac{\pi^2}{30} g_*(T_{\text{RH}}) T_{\text{MAX}}^4 \left[\frac{g_{*s}(T_{\text{MAX}})}{g_{*s}(T_{\text{RH}})} \right]^{4/3} \left(\frac{a_{\text{RH}}}{a_I} \right)^2. \quad (5)$$

Using Eq. (3) to relate a_{RH} to T_{RH} , we obtain the evolution of ρ_ϕ during kination:

$$\rho_\phi = \frac{\pi^2}{30} \left[\frac{T_{\text{MAX}}^3}{T_{\text{RH}}} \right]^2 \left[\frac{g_{*s}(T_{\text{MAX}})}{g_{*s}(T_{\text{RH}})} \right]^2 g_*(T_{\text{RH}}) \left(\frac{a_I}{a} \right)^6. \quad (6)$$

Now that we have obtained expressions for $T(a)$, $\rho_r(a)$ and $\rho_\phi(a)$, we have the necessary components to numerically solve Eq. (1b) for $n_\chi(a)$, as shown in Fig. 2. Figure 3 shows the dark matter relic abundance as a function of $\langle\sigma v\rangle$ for several values of T_{RH} and m_χ . For small $\langle\sigma v\rangle$ values, the dark matter cannot reach thermal equilibrium, and Fig. 3 shows that as $\langle\sigma v\rangle$ increases the dark matter relic abundance increases. Once $\langle\sigma v\rangle$ becomes large enough, pair production will bring n_χ up to its thermal equilibrium value. If dark matter reaches thermal equilibrium, we see from Fig. 3 that as $\langle\sigma v\rangle$ increases, the dark matter relic abundance decreases. In the following sections, we derive analytic expressions for the dark matter relic abundance generated during kination and analyze how the relic abundance is influenced by T_{RH} .




FIG. 3. The observed dark matter abundance $\Omega_\chi h^2$ as a function of the dark matter velocity-averaged annihilation cross section $\langle\sigma v\rangle$. Dark matter freezes in at small $\langle\sigma v\rangle$ where $\Omega_\chi h^2 \propto \langle\sigma v\rangle$, whereas dark matter freezes out at large $\langle\sigma v\rangle$ where $\Omega_\chi h^2 \propto \langle\sigma v\rangle^{-1}$. In the left panel we see that decreasing m_χ decreases $\Omega_\chi h^2$ for both cases. In the right panel we see that decreasing T_{RH} decreases $\Omega_\chi h^2$ for the freeze-in case but increases $\Omega_\chi h^2$ for the freeze-out case. In both panels the solid curves represent the numerical solutions to Eq. (1), while the symbols represent the analytic approximations represented by Eqs. (16) and (21). The solid black line represents the Planck measurement for the observed dark matter abundance, $\Omega_\chi h^2 = 0.12$ [44].

B. Freeze-out

If $\langle\sigma v\rangle$ is sufficiently large, then pair production brings dark matter into thermal equilibrium: $n_\chi = n_{\chi,\text{eq}}$, as defined in Eq. (2). Once $H \simeq \langle\sigma v\rangle n_{\chi,\text{eq}}$, the dark matter deviates from equilibrium and “freezes out.” If dark matter freezes out during radiation domination, nearly all dark matter annihilations cease at freeze-out. However, if dark matter freezes out during kination, we see from Fig. 2 that we need to take dark matter annihilations between the time of freeze-out and reheating into account to get an accurate relic abundance.

To analytically solve for the evolution of the dark matter number density between freeze-out and reheating, we define a dimensionless comoving number density $Y \equiv n_\chi (a/a_I)^3 T_{\text{RH}}^{-3}$. Equation (1b) is rewritten as

$$\frac{dY}{da} = \langle\sigma v\rangle \frac{T_{\text{RH}}^3 a_I^3}{H a^4} (Y_{\text{eq}}^2 - Y^2). \quad (7)$$

After the dark matter freezes out, $Y^2 \gg Y_{\text{eq}}^2$. Since during kination $H = H(a_I)[a_I/a]^3$, we simplify Eq. (7) to

$$\frac{dY}{da} = \frac{-\lambda_{\text{KD}}}{a} Y^2, \quad (8)$$

where $\lambda_{\text{KD}} = T_{\text{RH}}^3 \langle\sigma v\rangle / H(a_I)$. Integrating Eq. (8) from freeze-out to reheating yields

$$\frac{1}{Y_{\text{F}}} - \frac{1}{Y_{\text{RH}}} = -\lambda_{\text{KD}} \ln \frac{a_{\text{RH}}}{a_{\text{F}}}, \quad (9)$$

where Y_{F} and Y_{RH} are the comoving dark matter number densities at freeze-out and reheating. Therefore, if freeze-out

occurs during kination, the dark matter comoving number density experiences a logarithmic decrease between freeze-out and reheating.

To evaluate the current dark matter density we need to reevaluate Eq. (7) during radiation domination and solve for Y at some late time. During radiation domination $H = H(a_{\text{RH}})[a_{\text{RH}}/a]^2$, and by defining $\lambda_{\text{RD}} = [T_{\text{RH}}^3 \langle\sigma v\rangle / H(a_{\text{RH}})] \times [a_I^3 / a_{\text{RH}}^2]$, Eq. (7) simplifies to

$$\frac{dY}{da} = \frac{-\lambda_{\text{RD}}}{a^2} Y^2. \quad (10)$$

Solving Eq. (10) from reheating to a very late time yields

$$\frac{1}{Y_{\text{RH}}} - \frac{1}{Y_{\text{LT}}} = -\lambda_{\text{RD}} \left(\frac{1}{a_{\text{RH}}} \right), \quad (11)$$

where Y_{LT} is the comoving dark matter number density at some late time ($a = a_{\text{LT}}$). To obtain Eq. (11), we use the fact that $a_{\text{LT}} \gg a_{\text{RH}}$. Therefore, if dark matter freezes out during kination, Y experiences a logarithmic decrease between freeze-out and reheating, after which Y approaches a constant value.

Utilizing Y_{RH} from Eq. (9) and rewriting Eq. (11) in terms of the dark matter number density yields

$$n_{\chi,\text{LT}} = \left[\frac{\langle\sigma v\rangle a_{\text{LT}}^3}{H(a_I) a_I^3} \left(\ln \left[\frac{a_{\text{RH}}}{a_{\text{F}}} \right] + 1 \right) + \frac{a_{\text{LT}}^3}{n_{\chi,\text{F}} a_{\text{F}}^3} \right]^{-1}. \quad (12)$$

We wish to express Eq. (12) in terms of our free parameters m_χ , T_{RH} , and $\langle\sigma v\rangle$. We can express $H(a_I)$ and a_{RH} in terms of T_{RH} and T_{MAX} using Eqs. (3) and (5). In addition, since $n_{\chi,\text{F}} \simeq n_{\chi,\text{eq}}$, $n_{\chi,\text{F}} \simeq H(T_{\text{F}}) / \langle\sigma v\rangle$. If freeze-out occurs

during kination, $H^2 \approx (8\pi G/3)\rho_\phi$; combining this with Eqs. (3) and (6) allows us to solve for $H(T_F)$:

$$H(T_F) = \left(\frac{4\pi^3}{45}\right)^{1/2} \left(\frac{T_F^3}{m_{\text{pl}} T_{\text{RH}}}\right) \left(\frac{g_{*sF}}{g_{*s\text{RH}}}\right) g_{*s\text{RH}}^{1/2}, \quad (13)$$

where $g_{*s\text{RH}} = g_{*s}(T_{\text{RH}})$ and $g_{*s\text{RH}} = g_{*s}(T_{\text{RH}})$. These relations allow us to rewrite Eq. (12) as

$$n_{\chi,\text{LT}} = \left(\frac{4\pi^3}{45}\right)^{1/2} \frac{T_{\text{LT}}^3 g_{*s\text{LT}} g_{*s\text{RH}}^{1/2}}{\langle\sigma v\rangle m_{\text{pl}} T_{\text{RH}} g_{*s\text{RH}}} \times \left(\ln \left[\frac{T_F}{T_{\text{RH}}} \left(\frac{g_{*sF}}{g_{*s\text{RH}}}\right)^{1/3} \right] + 2 \right)^{-1}, \quad (14)$$

where T_F is obtained by numerically solving $H(T_F) = \langle\sigma v\rangle n_{\chi,\text{eq}}$. For kination scenarios, m_χ/T_F is roughly between 20 and 30.

After $a_{\text{LT}}, n_\chi \propto a^{-3}$, which allows us to relate the dark matter density at a_{LT} to today:

$$\rho_{\chi,0} = \rho_{\chi,\text{LT}} \left(\frac{a_{\text{LT}}}{a_0}\right)^3 = \rho_{\chi,\text{LT}} \left(\frac{T_0}{T_{\text{LT}}}\right)^3 \left(\frac{g_{*s0}}{g_{*s\text{LT}}}\right), \quad (15)$$

where T_0 is the radiation temperature today and $g_{*s0} = 3.91$. Bringing all of the previous components together and scaling our analytic expression by a factor of 1.22, thereby ensuring that it matches the numeric solution of Eq. (1b) within 20% for $m_\chi/T_{\text{RH}} > 100$, we obtain an analytic expression for the freeze-out dark matter relic abundance:

$$\Omega_\chi h^2 = 6.06 \left(\frac{3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}}{\langle\sigma v\rangle}\right) \frac{g_{*s\text{RH}}^{1/2}}{g_{*s\text{RH}}} \left(\frac{m_\chi/T_{\text{RH}}}{150}\right) \times \left(\ln \left[\frac{T_F/T_{\text{RH}}}{10} \left(\frac{g_{*sF}}{g_{*s\text{RH}}}\right)^{1/3} \right] + 4.3 \right)^{-1}. \quad (16)$$

Equation (16) indicates that decreasing T_{RH} increases the relic abundance. Decreasing T_{RH} requires increasing the kination energy density, which increases the Hubble parameter during kination. Since $n_{\chi,F} \approx H(T_F)/\langle\sigma v\rangle$, increasing the Hubble parameter increases the dark matter number density at freeze-out and thus increases the relic abundance. Furthermore, in our calculation of Y_{RH} we showed that dark matter annihilations do not cease during kination. As a result, Eq. (16) includes an inverse logarithmic term that depends on the ratio T_F/T_{RH} .

Figure 3 shows the dark matter relic abundance as a function of $\langle\sigma v\rangle$ for several values of T_{RH} and m_χ . In Fig. 3 we see that for sufficiently large $\langle\sigma v\rangle$ the freeze-out dark matter relic abundances from Eq. (16), represented by the circular symbols, match the numeric solutions to Eq. (1b), represented by the curves. We can solve for the minimum $\langle\sigma v\rangle$ that will result in the dark matter reaching thermal equilibrium. If dark matter freezes out, $H(T_F) = \langle\sigma v\rangle n_{\chi,\text{eq}}$,

and we can rewrite this equation in terms of a new variable x , where $x \equiv m_\chi/T_F$. Assuming that dark matter is nonrelativistic, $H(T_F) = \langle\sigma v\rangle n_{\chi,\text{eq}}$ can be rewritten as $x^{-3/2} e^x g_{*s}(m_\chi/x) = \text{const} \times \langle\sigma v\rangle$. The left-hand side of this equation has a minimum value near $x \sim 1.5$ which implies that there is a minimum $\langle\sigma v\rangle$ for which a solution will exist. This lower bound on $\langle\sigma v\rangle$ is

$$\langle\sigma v\rangle > 9.37 \times 10^{-33} \text{ cm}^3 \text{ s}^{-1} \left(\frac{2}{g_\chi}\right) \times \left(\frac{3 \text{ MeV}}{T_{\text{RH}}}\right) \left(\frac{g_{*s\text{RH}}^{1/2}}{g_{*s\text{RH}}}\right) g_{*s}(m_\chi/1.5). \quad (17)$$

The horizontal line in Fig. 3 represents the Planck measurement of the observed dark matter abundance. To reproduce the observed dark matter abundance, freeze-out cases during kination require larger $\langle\sigma v\rangle$ than that required if freeze-out occurs during radiation domination. This comes from the fact that at a given temperature the Hubble parameter during kination is always higher than it is during radiation domination, which causes freeze-out to occur earlier. In order to compensate for the earlier freeze-out and reproduce the observed dark matter abundance, freeze-out scenarios during kination require $\langle\sigma v\rangle > 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$. Since this lower bound on $\langle\sigma v\rangle$ is more stringent than Eq. (17), Eq. (16) is applicable to all freeze-out scenarios that generate the observed dark matter abundance.

C. Freeze-in

For cross sections that violate Eq. (17), dark matter pair production is not sufficient to bring the dark matter into thermal equilibrium. Once pair production ceases, the dark matter ‘‘freezes in’’ and the comoving dark matter number density remains constant. The first step in determining the freeze-in dark matter relic abundance is to calculate the comoving dark matter number density when pair production ceases.

In freeze-in scenarios, the dark matter number density does not reach thermal equilibrium, so $n_\chi \ll n_{\chi,\text{eq}}$. The dimensionless comoving dark matter equilibrium number density is $Y_{\text{eq}} \equiv n_{\chi,\text{eq}} T_{\text{RH}}^{-3} (a/a_I)^3$, and for freeze-in scenarios $Y_{\text{eq}} \gg Y$. Therefore, Eq. (7) reduces to

$$\frac{dY}{da} = \frac{\langle\sigma v\rangle T_{\text{RH}}^3}{H(a_I) a} Y_{\text{eq}}^2 \quad (18)$$

for freeze-in scenarios during kination.

Equation (18) implies that dY/da diverges as $a \rightarrow 0$ if $\langle\sigma v\rangle$ is independent of temperature. The same divergence occurs if the Universe is radiation dominated during dark matter production, and it would make the freeze-in abundance of dark matter dependent on T_{MAX} . Previous analyses of the freeze-in process avoided this sensitivity to




FIG. 4. The evolution of dY/da given $T_{\text{RH}} = 1 \text{ GeV}$, $m_\chi = 5 \times 10^4 \text{ GeV}$, and $\langle\sigma v\rangle_s = 10^{-47} \text{ cm}^3 \text{ s}^{-1}$. The vertical line represents the scale factor at which pair production peaks, defined as a_* . For kination scenarios where dark matter freezes in and $T_{\text{MAX}} = 8m_\chi$, $a_*/a_I \approx 8$.

high-energy physics by assuming that $\langle\sigma v\rangle \propto 1/T^2$ for relativistic particles [45,46]. We take the same approach and set $\langle\sigma v\rangle = \langle\sigma v\rangle_s (m_\chi/T)^2$ for $T > m_\chi$ and $\langle\sigma v\rangle = \langle\sigma v\rangle_s$ for $T < m_\chi$, where $\langle\sigma v\rangle_s$ is the s -wave dark matter annihilation cross section for massive particles. With this scaling, $dY/da \rightarrow 0$ as $a \rightarrow 0$, and the production of dark matter is finite during kination even if $T_{\text{MAX}} \rightarrow \infty$. Figure 4 shows that dY/da increases until $a = a_*$, which we define as the scale factor value at which pair production peaks. The temperature at which pair production peaks is $T_* = m_\chi$. Therefore, dY/da reaches its maximum when $a_*/a_I = 8[g_{*s}(T_{\text{MAX}})/g_{*s}(T_*)]^{1/3}$.

Figure 4 also indicates that the integral of dY/da converges; Y will approach a constant value as pair production becomes less and less efficient. However, Eq. (18) is only valid during kination, so it will only provide an accurate dark matter density if nearly all the pair production occurs prior to reheating. Truncating the integration of dY/da at a_{pp} , where $T(a_{\text{pp}}) = m_\chi/3.9$, reduces the value of Y by less than 1% compared to integrating dY/da out to $a = \infty$. Therefore, pair production has effectively halted when $T < m_\chi/3.9$, and we can use Eq. (18) to compute the relic abundance of dark matter provided that $T_{\text{RH}} < m_\chi/3.9$. Integrating Eq. (18) from 0 to a_{pp} , while taking into account the fact that $\langle\sigma v\rangle$ changes from $\langle\sigma v\rangle_s (m_\chi/T)^2$ to $\langle\sigma v\rangle_s$ at $T = m_\chi$, gives

$$Y_{\text{pp}} = 3.1 \times 10^{-4} \left(\frac{T_{\text{MAX}}/T_{\text{RH}}}{150} \right)^3 \left(\frac{T_{\text{RH}}}{5 \text{ GeV}} \right) \left(\frac{g_\chi}{2} \right)^2 \times \left(\frac{\langle\sigma v\rangle_s}{10^{-45} \text{ cm}^3 \text{ s}^{-1}} \right) \left(\frac{g_{*s\text{RH}}}{g_{*s\text{RH}}^{1/2}} \right) \left(\frac{g_{*s\text{MAX}}}{g_{*s}^2(m_\chi)} \right). \quad (19)$$

After pair production ends, Y remains nearly constant, and thus $Y_{\text{pp}} = Y_{\text{RH}}$. Therefore, the dark matter density at reheating can be written as

$$\rho_{\chi,\text{RH}} = m_\chi Y_{\text{pp}} T_{\text{RH}}^3 (a_I/a_{\text{RH}})^3. \quad (20)$$

Equation (3) indicates that $(a_I/a_{\text{RH}})^3 \propto T_{\text{MAX}}^{-3} g_{*s\text{MAX}}^{-1}$. As a result, inserting Eq. (19) into Eq. (20) seems to indicate that $\rho_{\chi,\text{RH}}$ is independent of T_{MAX} , but this is not generically true. When integrating Eq. (18) from $a = 0$ to a_{pp} to obtain Eq. (19), we effectively integrated from $T = \infty$ to $T = m_\chi/3.9$. However, integrating instead from $T = 8m_\chi$ to $T = m_\chi/3.9$ does not significantly change the result. Therefore, if $T_{\text{MAX}} \geq 8m_\chi$, the freeze-in dark matter abundance does not depend on T_{MAX} . Conversely, if $T_{\text{MAX}} < 8m_\chi$, $\rho_{\chi,\text{RH}}$ will decrease as T_{MAX} decreases because maximal pair production is not reached.

After reheating, $n_\chi \propto a^{-3}$, and we can use Eq. (15) to evolve ρ_χ from reheating to today. Combining the previous expressions and scaling our analytic expression by a factor of 0.95 to match the numeric solution of Eq. (1b) provides an analytic expression for the freeze-in dark matter relic abundance:

$$\Omega_\chi h^2 = 1.08 \left(\frac{m_\chi}{1 \text{ GeV}} \right) \left(\frac{T_{\text{RH}}}{100 \text{ GeV}} \right) \left(\frac{g_\chi}{2} \right)^2 \times \left(\frac{\langle\sigma v\rangle_s}{10^{-45} \text{ cm}^3 \text{ s}^{-1}} \right) \left(\frac{g_{*s\text{RH}}}{g_{*s\text{RH}}^{1/2}} \right) g_{*s}^{-2}(m_\chi). \quad (21)$$

Equation (21) indicates that increasing T_{RH} leads to a larger relic abundance. To understand how the freeze-in dark matter relic abundance relates to T_{RH} we need to investigate how n_χ relates to the Hubble parameter. The connection between the Hubble parameter and n_χ stems from the cooling rate dT/dt . During kination $T \propto a^{-1}$, and therefore $dT/dt = -HT$. Rewriting dn/dt as a function of temperature yields $dn/dt = (dT/dt)(dn/dT) = f(T)$, where $f(T)$ is the right-hand side of Eq. (1b). This allows us to express dn/dT as

$$\frac{dn}{dT} = \frac{-f(T)}{HT}, \quad (22)$$

which implies that $n \propto 1/H$. Since increasing T_{RH} decreases H during kination, it also decreases the cooling rate, leaving more time for pair production and thereby increasing the dark matter number density.

In order to reach the observed dark matter abundance, scenarios in which dark matter freezes in during kination require larger $\langle\sigma v\rangle_s$ values than if freeze-in occurs during radiation domination. For example, given a dark matter mass of 100 GeV, a freeze-in scenario during radiation domination requires $\langle\sigma v\rangle_s = 10^{-47} \text{ cm}^3 \text{ s}^{-1}$ in order for $\Omega_\chi h^2 = 0.12$ [44,47]. For the same m_χ , a freeze-in scenario during kination with $T_{\text{RH}} = 0.033 \text{ GeV}$ requires $\langle\sigma v\rangle_s = 10^{-41} \text{ cm}^3 \text{ s}^{-1}$. Freeze-in scenarios during kination require larger $\langle\sigma v\rangle_s$ values to generate the observed dark matter abundance because the increased

cooling rate during kination leaves less time for pair production.

In Fig. 3, we see that for sufficiently small cross sections the dark matter relic abundances from Eq. (21), represented by the star symbols, match the numeric solutions to Eq. (1b), represented by the curves. We have already discussed how freeze-in requires the dark matter particles to never reach thermal equilibrium. For this to hold true, the dark matter number density at the end of pair production must be less than the dark matter equilibrium number density at the peak of pair production: $Y(T_{\text{PP}}) < Y_{\text{eq}}(T_*)$. To solve for the largest $\langle\sigma v\rangle_s$ that will not result in dark matter reaching thermal equilibrium, we approximate the equilibrium number density as being nonrelativistic:

$$Y_{\text{eq}}(a) = g_\chi \left(\frac{a}{a_I}\right)^3 T_{\text{RH}}^{-3} \left(\frac{m_\chi T}{2\pi}\right)^{3/2} e^{-m_\chi/T}. \quad (23)$$

Evaluating Eq. (23) at a_* and equating that to Eq. (19) gives us the cross sections that will result in dark matter freezing in during kination:

$$\langle\sigma v\rangle_s \leq 8.5 \times 10^{-34} \text{ cm}^3 \text{ s}^{-1} \left(\frac{g_{*\text{RH}}}{g_{*\text{sRH}}}\right)^{1/2} \times g_{*\text{s}}(m_\chi) \left(\frac{2}{g_\chi}\right) \left(\frac{3 \text{ MeV}}{T_{\text{RH}}}\right). \quad (24)$$

Figure 3 demonstrates that there is a range of $\langle\sigma v\rangle_s$ values for each reheat temperature where neither a freeze-out nor freeze-in scenario will result in the observed dark matter abundance. The left panel of Fig. 3 shows that, at a fixed reheat temperature, decreasing m_χ increases the freeze-in cross section and decreases the freeze-out cross section that generates the observed dark matter abundance. However, once $m_\chi \lesssim 3T_{\text{RH}}$, dark matter will no longer freeze in during kination, and as discussed in Sec. II B, $\langle\sigma v\rangle > 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ is required to generate the observed dark matter abundance if dark matter freezes out during kination.

The right panel of Fig. 3 shows that decreasing T_{RH} increases the cross section that generates the observed dark matter abundance via the freeze-in mechanism. Therefore, setting $T_{\text{RH}} = 3 \text{ MeV}$ gives an upper bound on the cross sections that can generate the observed dark matter abundance in freeze-in scenarios:

$$\langle\sigma v\rangle_s < 2.66 \times 10^{-38} \text{ cm}^3 \text{ s}^{-1} \left(\frac{0.175 \text{ GeV}}{m_\chi}\right). \quad (25)$$

This maximal cross section is calculated using $g_{*\text{s}}(m_\chi) = g_{*\text{s}}(0.175 \text{ GeV})$. When deriving Eq. (21) we assumed that $g_{*\text{s}}(T)$ was approximately constant during pair production, which implies that $\Omega_\chi h^2 \propto g_{*\text{s}}^{-2}(m_\chi)$. If dark matter freezes in during kination and m_χ is less than 0.17 GeV, then the QCD phase transition occurs before the

peak of pair production. At the QCD phase transition $g_{*\text{s}}(T)$ sharply decreases, resulting in an increase in the relic abundance as given by Eq. (21). To compensate for the increased relic abundance, freeze-in scenarios with $m_\chi \leq 0.17 \text{ GeV}$ require cross sections smaller than the one calculated in Eq. (25). Therefore, the largest cross section that can generate the observed dark matter abundance in freeze-in scenarios is $2.7 \times 10^{-38} \text{ cm}^3 \text{ s}^{-1}$.

The relic abundances from Eq. (21) are within 20% of the solutions to Eq. (1b) for $m_\chi > 0.17 \text{ GeV}$ and $T_{\text{RH}} < m_\chi/3.9$. If $m_\chi \leq 0.17 \text{ GeV}$, we need to take into consideration the evolution of $g_{*\text{s}}(T)$ to accurately calculate the relic abundance. Allowing for the evolution of $g_{*\text{s}}(T)$, Y_{PP} is rewritten as

$$Y_{\text{PP}} = \left(\frac{45}{4\pi^3}\right)^{1/2} \frac{\langle\sigma v\rangle_s m_{\text{pl}} T_{\text{MAX}}^3}{T_{\text{RH}}^2 g_{*\text{RH}}^{1/2} m_\chi^6} g_{*\text{sMAX}} g_{*\text{sRH}} \times \left(\int_0^1 n_{\chi,\text{eq}}^2 x^7 g_{*\text{s}}^{-2}(m_\chi/x) dx + \int_1^{x_{\text{PP}}} n_{\chi,\text{eq}}^2 x^5 g_{*\text{s}}^{-2}(m_\chi/x) dx \right). \quad (26)$$

Using this expression for Y_{PP} and a scaling factor of 0.4, we construct a modified relic abundance expression that takes into account the evolution of $g_{*\text{s}}(T)$. For freeze-in scenarios with $m_\chi \leq 0.17 \text{ GeV}$ and $T_{\text{RH}} < m_\chi/3.9$, this updated expression for Y_{PP} brings the analytic relic abundance solutions to within 25% of the numeric solution to Eq. (1b).

III. CONSTRAINTS ON KINATION

To constrain kination cosmologies, we first solve Eqs. (16) and (21) for all combinations of the variables m_χ , T_{RH} , and $\langle\sigma v\rangle$ that produce the observed dark matter abundance of $\Omega_\chi h^2 = 0.12$ [44]. We set the minimum allowed reheat temperature to 3 MeV to ensure that the period of kination does not alter the cosmic microwave background or the abundances of light elements [4–8].² Next, we compare our allowed parameters to current constraints on m_χ and $\langle\sigma v\rangle$ from Fermi-LAT and H.E.S.S. Specifically, we use the Fermi-LAT PASS-8 constraints from observations of dwarf spheroidals [35] and H.E.S.S. constraints from observations of the Galactic center [36]. The Fermi-LAT data cover dark matter masses ranging from $2 \text{ GeV} \leq m_\chi \leq 10^4 \text{ GeV}$, while the H.E.S.S. data cover dark matter masses ranging from $125 \text{ GeV} \leq m_\chi \leq 7 \times 10^4 \text{ GeV}$.

Figure 5 shows the allowed parameter space for m_χ and $\langle\sigma v\rangle_s$ for scenarios in which dark matter freezes in during kination. To ensure that freeze-in occurs before the onset

²These constraints on the reheat temperature were derived assuming that the radiation-dominated era was preceded by an early-matter-dominated era, but we expect that similar constraints would apply to kination.

of radiation domination, we have restricted ourselves to $m_\chi/3.9 > T_{\text{RH}}$. This restriction comes from the fact that the temperature at which pair production effectively ceases is $T_{\text{PP}} = m_\chi/3.9$. Since the minimum reheat temperature is 3 MeV, we require that $m_\chi > 0.012$ GeV to ensure that freeze-in occurs before radiation domination. From Eq. (25) and Fig. 5, we see that scenarios in which dark matter freezes in during kination require $\langle\sigma v\rangle < 2.7 \times 10^{-38} \text{ cm}^3 \text{ s}^{-1}$.

Figure 6 shows the allowed parameter space for m_χ and $\langle\sigma v\rangle$ for scenarios in which dark matter freezes out during kination. To obtain the observed dark matter abundance, freeze-out scenarios during kination must have an annihilation cross section greater than $3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$. As discussed in Sec. II B, freeze-out occurs earlier during kination than during radiation domination, and to compensate, freeze-out scenarios during kination require larger annihilation cross sections to generate the same dark matter density.

In Fig. 6 we include the Fermi-LAT [35] and H.E.S.S. [36] constraints for dark matter that annihilates via the $b\bar{b}$ channel. The Fermi-LAT bounds cover a range of dark matter masses from the mass of the bottom quark to a mass of 10^4 GeV. The H.E.S.S. bounds add additional constraints to dark matter masses ranging from 200 to 7×10^4 GeV. Dark matter annihilation cross sections above the observational bounds are ruled out as these signals would have already been observed. Figure 6 also includes the partial-wave unitarity bound, which requires $\langle\sigma v\rangle \lesssim 1/m_\chi^2$ [48–50]. We see from Fig. 6 that the unitarity bound rules out all kination scenarios with $\langle\sigma v\rangle$ values larger than $4.5 \times 10^{-23} \text{ cm}^3 \text{ s}^{-1}$. In addition, if dark matter annihilates via the $b\bar{b}$ channel, Fermi-LAT and H.E.S.S. observations constrain $\langle\sigma v\rangle$ to be less than $2 \times 10^{-25} \text{ cm}^3 \text{ s}^{-1}$ and T_{RH} to be greater than 1 GeV.




FIG. 5. Allowed freeze-in parameter space for m_χ and $\langle\sigma v\rangle_s$. Equation (21) is not applicable to scenarios with $m_\chi/3.9 < T_{\text{RH}}$ because more than 1% of pair production occurs during radiation domination. Dark matter produced via freeze-in requires very small annihilation cross sections in order to reach the observed dark matter abundance. These small annihilation cross sections are not constrainable with current astrophysical observations.

Figure 7 shows the Fermi-LAT, H.E.S.S., and unitarity constraints on m_χ and T_{RH} for scenarios in which dark matter freezes out during kination for various annihilation channels. For every value of m_χ and T_{RH} we calculate the $\langle\sigma v\rangle$ that will produce the observed dark matter abundance via freeze-out using Eq. (16). If the calculated $\langle\sigma v\rangle$ is above the Fermi-LAT or H.E.S.S. constraints, then that scenario is ruled out. The ruled-out area below $T_{\text{RH}} = 3$ MeV represents the fact that, in order to produce the correct abundance of light elements, reheating must occur before a temperature of ~ 3 MeV. The solid black line represents when $m_\chi = 100T_{\text{RH}}$. As discussed in Sec. II B, Eq. (16) is accurate for $T_{\text{RH}} < m_\chi/100$. As T_{RH} increases beyond $m_\chi/100$, numerical tests with $m_\chi > 17$ GeV indicate that the $\langle\sigma v\rangle$ value that yields the observed dark matter abundance rapidly decreases to $3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ as freeze-out occurs closer to radiation domination. Therefore, we make the conservative assumption that $\langle\sigma v\rangle = 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ will give the observed dark matter abundance if $m_\chi > 17$ GeV and $T_{\text{RH}} > m_\chi/100$.

For $m_\chi < 17$ GeV, numerical tests show that Eq. (16) remains accurate for reheat temperatures slightly higher than $m_\chi/100$ if reheating occurs after the QCD phase transition. The QCD phase transition causes a sharp decrease in g_* when $T=0.17$ GeV, and since $T_{\text{RH}} > 3$ MeV, $g_{*,\text{SRH}} = g_{*,\text{RH}}$ and $H \propto g_{*,\text{SRH}}^{-1/2}$ during kination. Consequently, the Hubble parameter at a given temperature during kination sharply increases as T_{RH} goes below 0.17 GeV, which causes




FIG. 6. Allowed freeze-out parameter space for m_χ and $\langle\sigma v\rangle$. To obtain the observed dark matter abundance, scenarios with annihilation cross sections smaller than $3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ require freeze-out to occur during radiation domination. The short dashed line represents the H.E.S.S. [36] constraints for annihilation in the $b\bar{b}$ channel, and the medium dashed line is the Fermi-LAT [35] constraints, also for the $b\bar{b}$ annihilation channel. The long dashed line is the unitarity bound: $\langle\sigma v\rangle \lesssim 1/m_\chi^2$. All of the kination scenarios above the Fermi-LAT and H.E.S.S. constraint lines are ruled out as dark matter annihilations would have already been detected by the corresponding observations.




FIG. 7. Constraints on m_χ and T_{RH} for dark matter produced via the freeze-out mechanism. These panels represent the following annihilation channels: $b\bar{b}$ (top left), e^+e^- (top right), $\mu^+\mu^-$ (middle left), $\tau^+\tau^-$ (middle right), $u\bar{u}$ (bottom left), and W^+W^- (bottom right). The solid black line represents when $m_\chi = 100T_{RH}$. Below this line it is possible to reproduce the observed dark matter abundance if dark matter freezes out during kination. The shaded regions are those that are constrained by Fermi-LAT, H.E.S.S., unitarity, and BBN.

freeze-out to occur earlier. Therefore, Eq. (16) is applicable for scenarios with T_{RH} slightly higher than $m_\chi/100$ if $m_\chi < 17$ GeV because freeze-out still occurs during kination. For most annihilation channels, scenarios that generate the observed relic abundance with $m_\chi < 17$ GeV are ruled out by Fermi-LAT constraints. The exception is dark matter annihilating via $\mu^+\mu^-$. Figure 7 shows that, for dark matter annihilating via $\mu^+\mu^-$, Fermi-LAT constraints rule out all scenarios with $m_\chi \lesssim 8$ GeV. In addition,

if m_χ is between 8 and 17 GeV, Fermi-LAT constraints rule out scenarios with $T_{RH} \lesssim 0.17$ GeV. In these scenarios, freeze-out occurs during kination even though T_{RH} may be higher than $m_\chi/100$. As T_{RH} increases beyond 0.17 GeV, numerical tests with $8 \text{ GeV} < m_\chi < 17 \text{ GeV}$ indicate that the $\langle\sigma v\rangle$ value required to obtain the observed dark matter abundance rapidly decreases to $3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ as freeze-out occurs closer to radiation domination.

The resulting constraints on m_χ and T_{RH} are contingent on dark matter reaching thermal equilibrium during kination. Equation (17) indicates that decreasing T_{RH} and increasing $g_{*s}(m_\chi/1.5)$ increases the minimum value of $\langle\sigma v\rangle$ that results in dark matter reaching thermal equilibrium. Therefore, solving Eq. (17) with the minimum reheat temperature of 3 MeV and $g_{*s}(m_\chi/1.5) = 100$ shows that, if $\langle\sigma v\rangle > 3 \times 10^{-31} \text{ cm}^3 \text{ s}^{-1}$, dark matter will freeze out during kination regardless of T_{RH} or m_χ .

The Fermi-LAT and unitarity constraints establish an allowed mass range for each annihilation channel. The unitarity bound on $\langle\sigma v\rangle$ places an upper bound on the allowed dark matter mass of $1.9 \times 10^4 \text{ GeV}$ for all annihilation channels. The lower bound on the dark matter mass comes from the Fermi-LAT observations and is between 8 and 160 GeV, depending on the annihilation channel. As T_{RH} decreases, the range of viable masses decreases. The addition of the H.E.S.S. constraints restrict dark matter annihilating via $\tau^+\tau^-$ to have a mass around either 250 or 9000 GeV. For dark matter masses between 470 and 2500 GeV, the H.E.S.S. observations constrain $\langle\sigma v\rangle$ to be less than $3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ for dark matter annihilating via $\tau^+\tau^-$, thereby ruling out all scenarios where dark matter freezes out during kination or radiation domination.

Figure 7 also shows that with the Fermi-LAT, H.E.S.S., and unitarity constraints we can place lower limits on T_{RH} if dark matter reaches thermal equilibrium during kination. For example, we can rule out kination scenarios with reheat temperatures below 0.05 GeV for dark matter annihilating via the e^+e^- or $\mu^+\mu^-$ annihilation channel. We are also able to rule out kination scenarios with reheat temperatures below 0.6 GeV for the $\tau^+\tau^-$ and $u\bar{u}$ annihilation channels as well as reheat temperatures below 1 GeV for the $b\bar{b}$ and W^+W^- annihilation channels. In addition, kination scenarios where dark matter annihilates via the $b\bar{b}$, $\tau^+\tau^-$, or W^+W^- annihilation channel require T_{RH} be very close to T_{F} , which implies that these kination scenarios are on the verge of being ruled out.

Throughout this work, we assumed that dark matter consisted of one particle species. If dark matter consists of multiple particle species, then it is possible that only a fraction of the dark matter is thermally produced during kination. To determine what effect this has on the T_{RH} constraints shown in Fig. 7, we neglect the $\ln(T_{\text{F}}/T_{\text{RH}})$ term in Eq. (16) and make the rough estimate that for dark matter freezing out during kination, $\Omega_\chi h^2 \propto m_\chi / (\langle\sigma v\rangle T_{\text{RH}})$. If only a fraction of the dark matter consists of a thermal relic, such that $\Omega_\chi h^2 = f \Omega_{\text{dm}} h^2$, then to scale the relic abundance by a factor of f for a fixed dark matter mass requires scaling the product of $\langle\sigma v\rangle$ and T_{RH} by a factor of $1/f$. In addition, since the Fermi-LAT and H.E.S.S. constraints are obtained using the dark matter annihilation rate $\Gamma = \langle\sigma v\rangle \rho_\chi^2 / m_\chi^2$, altering ρ_χ will subsequently reduce the annihilation rate by a factor of f^2 and raise the maximum

allowed annihilation cross section $\langle\sigma v\rangle_{\text{max}}$ by a factor of $1/f^2$. Therefore, the minimum allowed reheat temperature $T_{\text{RH,min}} \propto \langle\sigma v\rangle T_{\text{RH}} / \langle\sigma v\rangle_{\text{max}} \propto f^{-1}/f^{-2} \propto f$. For example, for dark matter annihilating via W^+W^- , the minimum allowed reheat temperature is 1 GeV if dark matter consists of a single particle species. If only a fraction of dark matter is thermally produced during kination and $\Omega_\chi h^2 = 0.05$, then $f = 0.42$ and the new minimum allowed reheat temperature is roughly 0.34 GeV.

IV. CONCLUSION

Our uncertainty regarding the expansion history of the Universe between the end of inflation and the beginning of BBN allows for the possibility that within this period there was an intermittent era of kination. In this paper we have investigated the effects that a period of kination has on the thermal production of dark matter. Previous studies on this topic have required the use of specific kination potentials [38–40,42,43]. Our analysis is independent of the kination potential, and therefore our constraints on m_χ , T_{RH} , and $\langle\sigma v\rangle$ are applicable to all kination models assuming that dark matter consists of one particle species that undergoes s -wave annihilation. In addition to numerically solving for the dark matter relic abundance, we have also derived analytic relic abundance equations for freeze-out and freeze-in kination scenarios.

Our scenarios are determined by three parameters: the dark matter mass m_χ , the reheat temperature T_{RH} , and the dark matter annihilation cross section $\langle\sigma v\rangle$. In deriving relic abundance equations for freeze-out [Eq. (16)] and freeze-in [Eq. (21)] scenarios, we have deduced physical relationships between our parameters and the dark matter relic abundance. For example, at a given temperature, the Hubble parameter during kination is higher than that during radiation domination. Therefore, freeze-out occurs earlier during kination, which increases the relic abundance. In order to compensate for this larger relic abundance, freeze-out scenarios require larger-than-canonical $\langle\sigma v\rangle$ values in order to increase the annihilation rate and subsequently decrease the dark matter abundance to the observed value. If on the other hand, dark matter freezes in during kination, an increase in T_{RH} will increase the relic abundance. Increasing T_{RH} decreases ρ_ϕ , which decreases the Hubble parameter. Decreasing the Hubble parameter decreases the cooling rate, leaving more time for pair production and thereby increasing the relic abundance. To compensate for the increased cooling rate and bring the dark matter abundance into agreement with the observed value, freeze-in scenarios during kination require larger $\langle\sigma v\rangle$ values compared to during radiation domination. Overall, to reach the observed dark matter abundance, freeze-out scenarios during kination require $\langle\sigma v\rangle$ values that would underproduce dark matter during radiation domination, whereas freeze-in scenarios require $\langle\sigma v\rangle$ values that would overproduce dark matter during radiation domination.

Therefore, the possibility that dark matter was thermally produced during kination significantly widens the field of potential dark matter candidates. In particular, thermally produced winos and Higgsinos, which generally have $\langle\sigma v\rangle > 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$, could constitute all the dark matter if they freeze out during a period of kination.

Our analytic relic abundance equations allow us to efficiently determine the dark matter parameter space that would result in the observed dark matter abundance. To ensure that freeze-out occurs before reheating and that reheating occurs at a temperature above 3 MeV, $\langle\sigma v\rangle$ values between 2.7×10^{-38} and $3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ are forbidden for all dark matter masses. Using the PASS 8 Fermi-LAT observations of dwarf spheroidal galaxies, H.E.S.S. observations of the Galactic center, and the unitarity bound on $\langle\sigma v\rangle$ we further constrain kination models. The observational bounds and subsequent constraints only apply to freeze-out scenarios because the required $\langle\sigma v\rangle$ values for freeze-in scenarios are below observational thresholds. From the unitarity constraint, we were able to rule out all kination scenarios with $\langle\sigma v\rangle$ greater than $4.5 \times 10^{-23} \text{ cm}^3 \text{ s}^{-1}$. These constraints also allowed us to rule out kination scenarios with reheat temperatures below 0.05 GeV for dark matter annihilating via the e^+e^- or $\mu^+\mu^-$ annihilation channel. Similarly, we ruled out kination scenarios with reheat temperatures below 0.6 GeV for the $\tau^+\tau^-$ and $u\bar{u}$ annihilation channels as well as reheat temperatures below 1 GeV for the $b\bar{b}$ and W^+W^- annihilation channels. Since these new bounds on T_{RH} are below the electroweak phase transition, kination could facilitate baryogenesis [25]. These bounds on T_{RH} are contingent on dark matter freezing out during kination. We have shown that, if $\langle\sigma v\rangle > 3 \times 10^{-31} \text{ cm}^3 \text{ s}^{-1}$, dark matter will freeze out during kination regardless of T_{RH} or m_χ . We also note that we only consider s -wave dark matter annihilation. If we consider a p -wave process, the annihilation rate in the Galaxy would be suppressed relative to

the annihilation rate at freeze-out and our bounds would no longer apply.

We have shown that scenarios in which dark matter is thermally produced during kination are not ruled out by current observational limits on the dark matter annihilation cross section. In these scenarios, $\langle\sigma v\rangle > 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ is required to generate the observed dark matter abundance via a freeze-out process. The observed dark matter abundance can be generated by a freeze-in process if $\langle\sigma v\rangle < 2.7 \times 10^{-38} \text{ cm}^3 \text{ s}^{-1}$. Therefore, our uncertainty regarding the pre-BBN expansion history prevents us from knowing the dark matter annihilation cross section that yields the current dark matter abundance; there exists a degeneracy between the allowed values of $\langle\sigma v\rangle$ and T_{RH} that cannot be eliminated by relic abundance calculations alone. One possible approach to breaking this degeneracy involves studying the evolution of the dark matter perturbations during kination. The evolution of perturbations during kination will impact the small-scale matter power spectrum. Therefore, studying how the small-scale matter power spectrum and small-scale structure formation are affected by kination may provide a means to further constrain kination scenarios and reduce the range of viable annihilation cross sections.

ACKNOWLEDGMENTS

We thank Carisa Miller for her comments on our manuscript. This work was supported by National Science Foundation Grant No. PHY-1417446.

Note added.—While we were finishing this paper, Ref. [51] appeared on the arXiv. This paper also considers dark matter production during kination, and it similarly identifies the logarithmic decrease of the dark matter number density between freeze-out and reheating.

-
- [1] A. H. Guth, *Phys. Rev. D* **23**, 347 (1981).
 - [2] A. D. Linde, *Phys. Lett. B* **108**, 389 (1982).
 - [3] A. Albrecht and P. J. Steinhardt, *Phys. Rev. Lett.* **48**, 1220 (1982).
 - [4] M. Kawasaki, K. Kohri, and N. Sugiyama, *Phys. Rev. Lett.* **82**, 4168 (1999).
 - [5] M. Kawasaki, K. Kohri, and N. Sugiyama, *Phys. Rev. D* **62**, 023506 (2000).
 - [6] S. Hannestad, *Phys. Rev. D* **70**, 043506 (2004).
 - [7] K. Ichikawa, M. Kawasaki, and F. Takahashi, *Phys. Rev. D* **72**, 043522 (2005).
 - [8] K. Ichikawa, M. Kawasaki, and F. Takahashi, *J. Cosmol. Astropart. Phys.* **05** (2007) 007.
 - [9] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Phys. Rev. Lett.* **73**, 3195 (1994).
 - [10] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Phys. Rev. D* **56**, 3258 (1997).
 - [11] R. Allahverdi, R. Brandenberger, F.-Y. Cyr-Racine, and A. Mazumdar, *Annu. Rev. Nucl. Part. Sci.* **60**, 27 (2010).
 - [12] G. Kane, K. Sinha, and S. Watson, *Int. J. Mod. Phys. D* **24**, 1530022 (2015).
 - [13] M. S. Turner, *Phys. Rev. D* **28**, 1243 (1983).
 - [14] D. J. H. Chung, E. W. Kolb, and A. Riotto, *Phys. Rev. D* **60**, 063504 (1999).
 - [15] M. Kamionkowski and M. S. Turner, *Phys. Rev. D* **42**, 3310 (1990).

- [16] G. F. Giudice, E. W. Kolb, and A. Riotto, *Phys. Rev. D* **64**, 023508 (2001).
- [17] N. Fornengo, A. Riotto, and S. Scopel, *Phys. Rev. D* **67**, 023514 (2003).
- [18] C. Pallis, *Astropart. Phys.* **21**, 689 (2004).
- [19] G. Gelmini, P. Gondolo, A. Soldatenko, and C. E. Yaguna, *Phys. Rev. D* **74**, 083514 (2006).
- [20] G. B. Gelmini and P. Gondolo, *Phys. Rev. D* **74**, 023510 (2006).
- [21] M. Drewes, *J. Cosmol. Astropart. Phys.* **11** (2014) 020.
- [22] G. L. Kane, P. Kumar, B. D. Nelson, and B. Zheng, *Phys. Rev. D* **93**, 063527 (2016).
- [23] A. L. Erickcek, *Phys. Rev. D* **92**, 103505 (2015).
- [24] B. Spokoiny, *Phys. Lett. B* **315**, 40 (1993).
- [25] M. Joyce, *Phys. Rev. D* **55**, 1875 (1997).
- [26] P. G. Ferreira and M. Joyce, *Phys. Rev. D* **58**, 023503 (1998).
- [27] P. J. E. Peebles and A. Vilenkin, *Phys. Rev. D* **59**, 063505 (1999).
- [28] K. Dimopoulos and J. W. F. Valle, *Astropart. Phys.* **18**, 287 (2002).
- [29] K. Dimopoulos, *Phys. Rev. D* **68**, 123506 (2003).
- [30] D. J. H. Chung, L. L. Everett, K. Kong, and K. T. Matchev, *J. High Energy Phys.* **10** (2007) 016.
- [31] M. Drees, H. Imminiyaz, and M. Kakizaki, *Phys. Rev. D* **73**, 123502 (2006).
- [32] D. Grin, T. L. Smith, and M. Kamionkowski, *Phys. Rev. D* **77**, 085020 (2008).
- [33] S. Watson, *Adv. Ser. Dir. High Energy Phys.* **21**, 305 (2010).
- [34] A. Beniwal, M. Lewicki, J. D. Wells, M. White, and A. G. Williams, [arXiv:1702.06124](https://arxiv.org/abs/1702.06124).
- [35] M. Ackermann *et al.* (Fermi-LAT Collaboration), *Phys. Rev. Lett.* **115**, 231301 (2015).
- [36] V. Lefranc and E. Moulin (H.E.S.S. Collaboration), *Proc. Sci.*, ICRC2015 (2016) 1208 [[arXiv:1509.04123](https://arxiv.org/abs/1509.04123)].
- [37] S. Profumo and P. Ullio, *J. Cosmol. Astropart. Phys.* **11** (2003) 006.
- [38] C. Pallis, *J. Cosmol. Astropart. Phys.* **10** (2005) 015.
- [39] C. Pallis, *Nucl. Phys.* **B751**, 129 (2006).
- [40] C. Pallis, in *Proceedings of the 6th International Workshop on the identification of dark matter (IDM 2006): Rhodes, Greece, 2006* (World scientific, Singapore, 2006), pp. 602–608.
- [41] M. E. Gomez, S. Lola, C. Pallis, and J. Rodriguez-Quintero, *J. Cosmol. Astropart. Phys.* **01** (2009) 027.
- [42] S. Lola, C. Pallis, and E. Tzelati, *J. Cosmol. Astropart. Phys.* **11** (2009) 017.
- [43] C. Pallis, *Nucl. Phys.* **B831**, 217 (2010).
- [44] P. A. R. Ade *et al.* (Planck Collaboration), *Astron. Astrophys.* **594**, A13 (2016).
- [45] L. J. Hall, K. Jedamzik, J. March-Russell, and S. M. West, *J. High Energy Phys.* **03** (2010) 080.
- [46] C. E. Yaguna, *J. High Energy Phys.* **08** (2011) 060.
- [47] P. S. B. Dev, A. Mazumdar, and S. Qutub, *Front. Phys.* **2**, 26 (2014).
- [48] K. Griest and M. Kamionkowski, *Phys. Rev. Lett.* **64**, 615 (1990).
- [49] L. Hui, *Phys. Rev. Lett.* **86**, 3467 (2001).
- [50] W.-L. Guo and X. Zhang, *Phys. Rev. D* **79**, 115023 (2009).
- [51] F. D’Eramo, N. Fernandez, and S. Profumo, *J. Cosmol. Astropart. Phys.* **05** (2017) 012.