

# Probing $Z - Z'$ mixing with ATLAS and CMS resonant diboson production data at the LHC at $\sqrt{s} = 13$ TeV

P. Osland\*

*Department of Physics and Technology, University of Bergen, Postboks 7803, N-5020 Bergen, Norway*

A. A. Pankov†

*The Abdus Salam ICTP Affiliated Centre, Technical University of Gomel, 246746 Gomel, Belarus and Institute for Nuclear Problems, Belarusian State University, 220030 Minsk, Belarus*

A. V. Tsytrinov‡

*The Abdus Salam ICTP Affiliated Centre, Technical University of Gomel, 246746 Gomel, Belarus*

(Received 14 July 2017; published 27 September 2017)

The study of electroweak boson pair production provides a powerful test of the spontaneously broken gauge symmetry of the standard model (SM) and can be used to search for new phenomena beyond the SM. Extra neutral vector bosons  $Z'$  decaying to charged gauge vector boson pairs  $W^+W^-$  are predicted in many scenarios of new physics, including models with an extended gauge sector. The diboson production allows us to place stringent constraints on the  $Z$ - $Z'$  mixing factor  $\xi$  and  $Z'$  mass,  $M_{Z'}$ . We present the  $Z'$  exclusion region in the  $\xi - M_{Z'}$  plane for the first time by using data comprised of  $pp$  collisions at  $\sqrt{s} = 13$  TeV and recorded by the ATLAS and CMS detectors at the CERN LHC, with integrated luminosities of 36.1 and 35.9 fb<sup>-1</sup>, respectively. The exclusion region has been significantly extended compared to that obtained from the previous analysis performed with Tevatron data as well as with LHC data collected at 7 and 8 TeV. Also, we found that these constraints on the  $Z$ - $Z'$  mixing factor are more severe than those derived from the global analysis of electroweak data. Further improvement on the constraining of this mixing can be achieved from the analysis of data to be collected at higher luminosity.

DOI: 10.1103/PhysRevD.96.055040

## I. INTRODUCTION

Many new physics (NP) scenarios beyond the SM [1], including superstring and left-right-symmetric models, predict the existence of new neutral and charged gauge bosons, which might be light enough to be accessible at current and/or future colliders [2]. The search for these new neutral  $Z'$  and charged  $W'$  gauge bosons is an important aspect of the experimental physics program of high-energy colliders. In this note we concentrate on the former one.

Present limits from direct production at the LHC and virtual effects at Large Electron-Positron Collider (LEP), through interference or mixing with the  $Z$  boson, imply that any new  $Z'$  boson is rather heavy and mixes very little with the  $Z$  boson. Depending on the considered theoretical model,  $Z'$  masses of the order of 4.5 TeV [3,4] and  $Z$ - $Z'$  mixing angles at the level of a few per mil are excluded [5] (see also [6,7]). The mixing angle is strongly constrained by very high-precision experiments at LEP and the SLAC Linear Collider (SLC) [8]. They include measurements from the  $Z$  line shape, from the leptonic branching ratios normalized to the total hadronic  $Z$  decay width as well as from leptonic forward-backward asymmetries. A  $Z'$  boson,

if lighter than about 5 TeV, could be discovered at the LHC [9,10] with  $\sqrt{s} = 14$  TeV in the Drell-Yan (DY) process  $pp \rightarrow Z' \rightarrow \ell^+\ell^- + X$  with  $\ell = e, \mu$ .

After the discovery of a  $Z'$  boson at the LHC via the DY process, some diagnostics of its couplings and  $Z$ - $Z'$  mixing needs to be performed in order to identify the underlying theoretical framework. In this paper we investigate the implications of the ATLAS [11] and CMS [12] data in the diboson channel

$$pp \rightarrow W^+W^- + X \quad (1)$$

to probe the  $Z'$  boson that arises, e.g., in a popular model with extended gauge sector proposed in [13]. The analysis is based on  $pp$  collision data at a center-of-mass energy  $\sqrt{s} = 13$  TeV, collected by the ATLAS (36.1 fb<sup>-1</sup>) and CMS (35.9 fb<sup>-1</sup>) experiments at the LHC. In particular, the data is used to probe the  $Z$ - $Z'$  mixing. In ATLAS  $W^+W^-$  events are reconstructed via their semileptonic decays of the  $W$ 's where one  $W$  boson decays into a charged lepton ( $l = e, \mu$ ) and a neutrino, and the other into two jets [11], while in CMS  $W$  bosons decay hadronically with two reconstructed jets (dijet channel) [12].

The  $W^\pm$  boson pair production process (1) is important for studying the electroweak gauge symmetry. General properties of the weak gauge bosons are closely related to

\*Per.Osland@uib.no

†pankov@ictp.it

‡tsytrin@rambler.ru

electroweak symmetry breaking and the structure of the gauge sector, like the existence and structure of trilinear couplings. In addition, the diboson decay modes of the  $Z'$  probe the gauge coupling strength between the new and the standard-model gauge bosons [14–17]. Furthermore, the coupling strength strongly influences the decay branching ratios and the natural widths of such a new gauge boson. Thus, detailed examination of the process (1) will both test the gauge sector of the SM with high accuracy and shed light on NP that may appear beyond the SM. Here, we examine the feasibility of observing a  $Z'$  boson in the  $W^\pm$  pair production process at the LHC, which in contrast to the DY process is not the principal discovery channel, but can help to understand the origin of new gauge bosons.

Direct searches for a heavy  $WW$  resonance have been performed at the Tevatron by both the CDF and D0 collaborations. The D0 collaboration explored diboson resonant production up to  $\mathcal{O}(700 \text{ GeV})$  using the  $\ell\nu\ell'\nu'$  and  $\ell\nu jj$  final states [18]. The CDF collaboration also searched for resonant  $WW$  production in the  $\ell\nu jj$  final state, resulting in a lower limit on the mass of  $Z'$  and  $W'$  bosons [6], excluding masses up to  $\mathcal{O}(900 \text{ GeV})$ , depending on the mixing.

The direct  $WW$  resonance search by the ATLAS and CMS Collaborations using, respectively, semileptonic  $\ell\nu jj$  and hadronic final-state events in  $pp$  collision data at 13 TeV set mass limits of  $\mathcal{O}(3 \text{ TeV})^1$  on such resonances [11,12] (see also [19–21]).

In this work, we derive bounds on a possible new neutral spin-1 resonance ( $Z'$ ) from the available ATLAS and CMS data on  $W^+W^-$  pair production [11,12]. We present results as constraints on the relevant  $Z$ - $Z'$  mixing factor introduced in Sec. II and on the  $M_{Z'}$  mass.

The paper is organized as follows. In Sec. II, we briefly review a model involving an additional  $Z'$  boson and emphasize the role of  $Z$ - $Z'$  mixing in the process (1). We give expressions for basic observables (cross sections) of the process under consideration at parton and hadron levels. We also discuss achievable constraints on  $Z'$  model parameters from different experiments underlining the role of the LHC in substantially improving the current limits on the  $Z$ - $Z'$  mixing. Section III presents some concluding remarks.

## II. CROSS-SECTION AND CONSTRAINTS ON $Z$ - $Z'$ MIXING

There are many theoretical models which predict a  $Z'$  with mass possibly in the TeV range. We will consider a NP model where  $Z'$ 's interact with light quarks and charged gauge bosons via their mixing with the SM  $Z$  assuming that the  $Z'$  couplings exhibit the same Lorentz structure as those of the SM. In particular, in the present analysis we will focus on a gauge boson of the “sequential standard model”

(SSM). In the simple reference model described in [13,22], the couplings of the  $Z'$  boson to fermions (quarks, leptons) and  $W$  bosons are a direct transcription of the corresponding standard-model couplings. Note that such a  $Z'$  boson is not expected in the context of gauge theories unless it has additional couplings to exotic fermions. However, it serves as a useful reference case when comparing constraints from various sources. It could also play the role of an excited state of the ordinary  $Z$  in models of compositeness or with extra dimensions at the weak scale.

In many extended gauge models, while the couplings to fermions are not much different from those of the SM, the  $Z'WW$  coupling is substantially suppressed with respect to that of the SM. In fact, in an extended gauge model the standard-model trilinear gauge boson coupling strength,  $g_{WWZ}(= \cot\theta_W)$ , is replaced by  $g_{WWZ} \rightarrow \xi \cdot g_{WWZ}$ , where  $\xi = \mathcal{C} \cdot (M_W/M_{Z'})^2$  is the mixing factor and  $\mathcal{C}$  the coupling strength scaling factor [23]. We will set cross section limits on such  $Z'_{\text{SSM}}$  as a function of the mass  $M_{Z'}$  and  $\xi$ . One should note that most  $Z'$  search results report mass limits along the  $\xi = (M_W/M_{Z'})^2$  line ( $\mathcal{C} = 1$  is referred to as “reference model”) and we have also done so for comparison.

The differential cross section for  $Z'$  production in the process (1) from initial quark-antiquark states can be written as

$$\frac{d\sigma^{Z'}}{dM dy dz} = K \frac{2M}{s} \sum_q [f_{q|P_1}(\xi_1) f_{\bar{q}|P_2}(\xi_2) + f_{\bar{q}|P_1}(\xi_1) f_{q|P_2}(\xi_2)] \frac{d\hat{\sigma}_{q\bar{q}}^{Z'}}{dz}. \quad (2)$$

Here,  $s$  denotes the proton-proton center-of-mass energy squared,  $z \equiv \cos\theta$ , with  $\theta$  the  $W^-$ -boson–quark angle in the  $W^+W^-$  center-of-mass frame and  $y$  is the diboson rapidity. Furthermore,  $f_{q|P_1}(\xi_1, M)$  and  $f_{\bar{q}|P_2}(\xi_2, M)$  are parton distribution functions for the protons  $P_1$  and  $P_2$ , respectively, with  $\xi_{1,2} = (M/\sqrt{s}) \exp(\pm y)$  the parton fractional momenta. Finally,  $d\hat{\sigma}_{q\bar{q}}^{Z'}/dz$  are the partonic differential cross sections. In (2), the  $K$  factor accounts for higher-order QCD contributions. For numerical computation, we use CTEQ-6L1 parton distributions [24]. Our estimates will be at the Born level, hence the factorization scale  $\mu_F$  enters solely through the parton distribution functions, as the parton-level cross-section at this order does not depend on  $\mu_F$ . As regards the scale dependence of the parton distributions we choose for the factorization scale the  $W^+W^-$  invariant mass,  $\mu_F^2 = M^2 = \hat{s}$ , with  $\hat{s} = \xi_1 \xi_2 s$  the parton subprocess c.m. energy squared. The obtained constraints presented in the following are not significantly modified when  $\mu_F$  is varied from  $\mu_F/2$  to  $2\mu_F$ .

The cross-section for the narrow  $Z'$  state production and subsequent decay into a  $W^+W^-$  pair needed in order to estimate the expected number of  $Z'$  events,  $N^{Z'}$ , is derived

<sup>1</sup>The quoted limit assumes  $g_{WWZ}/g_{WWZ} = (M_W/M_{Z'})^2$ .

from (2) by integrating the right-hand side over  $z$ , over the rapidity of the  $W^\pm$ -pair  $y$  and invariant mass  $M$  around the resonance peak ( $M_R - \Delta M/2$ ,  $M_R + \Delta M/2$ ):

$$\sigma^{Z'}(pp \rightarrow W^+W^- + X) = \int_{M_R - \Delta M/2}^{M_R + \Delta M/2} dM \int_{-Y}^Y dy \times \int_{-z_{\text{cut}}}^{z_{\text{cut}}} dz \frac{d\sigma^{Z'}}{dM dy dz}, \quad (3)$$

where the phase space can be found, e.g., in [16]. Using Eq. (3), the number of signal events for a narrow  $Z'$  resonance state can be written as follows

$$N^{Z'} = \mathcal{L} \cdot \varepsilon \cdot \sigma^{Z'}(pp \rightarrow W^+W^- + X) \equiv \mathcal{L} \cdot \varepsilon \cdot A_{WW} \cdot \sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow W^+W^-). \quad (4)$$

Here,  $\mathcal{L}$  denotes the integrated luminosity, and the overall kinematic and geometric acceptance times trigger, reconstruction and selection efficiencies,  $A_{WW} \times \varepsilon$ , is defined as the number of signal events passing the full event selection divided by the number of generated events [25,26]. Finally,  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow W^+W^-)$  is the (theoretical) total production cross-section times branching ratio extrapolated to the total phase space.

The differential cross-section for the processes  $q\bar{q} \rightarrow Z'_{\text{SSM}} \rightarrow W^+W^-$ , averaged over quark colors, can be written as [16]

$$\frac{d\hat{\sigma}_{q\bar{q}}^{Z'}}{d\cos\theta} = \frac{1}{3} \frac{\pi\alpha^2 \cot^2\theta_W}{16} (v_f^2 + a_f^2) \frac{\hat{s}}{(\hat{s} - M_{Z'}^2)^2 + M_{Z'}^2 \Gamma_{Z'}^2} \times \xi^2 \beta_W^3 \left( \frac{\hat{s}^2}{M_W^4} \sin^2\theta + 4 \frac{\hat{s}}{M_W^2} (4 - \sin^2\theta) + 12 \sin^2\theta \right), \quad (5)$$

where  $v_f = (T_{3,f} - 2Q_f s_W^2)/(2s_W c_W)$ ,  $a_f = T_{3,f}/(2s_W c_W)$ . Finally,  $M_{Z'}$  and  $\Gamma_{Z'}$  denote the mass and total width of the  $Z'$  boson.

In the calculation of the total width  $\Gamma_{Z'}$  we included the following channels:  $Z' \rightarrow f\bar{f}$ ,  $W^+W^-$ , and  $ZH$  [27], where  $H$  is the SM Higgs boson and  $f$  are the SM fermions ( $f = l, \nu, q$ ). The total width  $\Gamma_{Z'}$  of the  $Z'$  boson can be written as follows:

$$\Gamma_{Z'} = \sum_f \Gamma_{Z'}^{ff} + \Gamma_{Z'}^{WW} + \Gamma_{Z'}^{ZH}. \quad (6)$$

The presence of the two last decay channels, which are often neglected, is due to  $Z$ - $Z'$  mixing. However for large  $Z'$  masses there is an enhancement that cancels the suppression due to the tiny  $Z$ - $Z'$  mixing parameter  $\xi$  [28]. Notice that for all  $M_{Z'}$  values of interest for LHC the width of the  $Z'_{\text{SSM}}$  boson is considerably smaller than the mass resolution  $\Delta M$ .

The expression for the partial width of the  $Z' \rightarrow W^+W^-$  decay channel can be written as [13]:

$$\Gamma_{Z'}^{WW} = \frac{\alpha}{48} \cot^2\theta_W M_{Z'} \left( \frac{M_{Z'}}{M_W} \right)^4 \left( 1 - 4 \frac{M_W^2}{M_{Z'}^2} \right)^{3/2} \times \left[ 1 + 20 \left( \frac{M_W}{M_{Z'}} \right)^2 + 12 \left( \frac{M_W}{M_{Z'}} \right)^4 \right] \xi^2. \quad (7)$$

The dominant term in the second line of Eq. (5), for  $M^2 \gg M_W^2$ , is proportional to  $(M/M_W)^4 \sin^2\theta$  and corresponds to the production of longitudinally polarized  $W$ 's,  $Z' \rightarrow W_L^+ W_L^-$ . This strong dependence on the invariant mass results in a very steep growth of the cross section with energy and therefore a substantial increase of the cross section sensitivity to  $Z$ - $Z'$  mixing at high  $M$ . In its turn, for a fixed mixing factor  $\xi$  and at large  $M_{Z'}$  where  $\Gamma_{Z'}^{WW}$  dominates over  $\sum_f \Gamma_{Z'}^{ff}$  and  $\Gamma_{Z'}^{ZH}$  the total width increases very rapidly with the mass  $M_{Z'}$  because of the quintic dependence on the  $Z'$  mass of the  $W^+W^-$  mode as shown in Eq. (7) [13]. In this case, the  $W^+W^-$  mode becomes dominant and  $\text{Br}(Z' \rightarrow W^+W^-) \rightarrow 1$ , while the fermionic decay channels are increasingly suppressed. While the equivalence theorem [29,30] might suggest a value for  $\text{Br}(Z' \rightarrow ZH)$  comparable to  $\text{Br}(Z' \rightarrow W^+W^-)$ , we note that this is inhibited by the vanishing  $\text{SU}(2)$  structure constant for the  $ZZZ$  coupling.

Further contributions of decays involving Higgs and/or gauge bosons and supersymmetric partners (including sfermions), which are not accounted for in (6), could increase  $\Gamma_{Z'}$  by a model-dependent amount, as large as 50% [14]. In this case,  $\Gamma_{Z'}$  would be larger, with a corresponding suppression in the branching ratio to  $W^\pm$ , and the bounds from LHC (and their ability for observing the  $Z$ - $Z'$  mixing effect) would be weaker.

In our early study [15], to gain some approximate understanding of the acceptances for signal and background at different values of the invariant mass  $M$  of the  $W^+W^-$  pair and  $Z'$  mass, we performed a simple study as follows. To estimate the  $2\sigma$  constraints on the  $Z$ - $Z'$  mixing parameter at the LHC, we compared the events due to a  $Z'$  signal to the events from the SM background in a 3% interval around the relevant values of the diboson  $W^+W^-$  invariant mass. This should be compatible with the expected energy resolution and with the fact that  $\Gamma_{Z'}/M_{Z'} \sim 3.0$ – $3.5\%$ . We then required the signal events to be at least a  $2\sigma$  fluctuation over the expected background, and in any case more than 3. This rough statistical analysis, as a preliminary stage, was enough to get an approximate answer to the questions we wanted to address.

Here, we are making a more careful analysis, employing the most recent measurements of diboson processes provided by the experimental collaborations ATLAS and CMS, which have control of the information needed to perform it in a more accurate way. In particular, for  $Z'_{\text{SSM}}$




FIG. 1. Observed and expected 95% C.L. upper limits on the production cross section times the branching fraction for  $Z' \rightarrow W^+W^-$  as a function of  $Z'$  mass,  $M_{Z'}$ . Theoretical production cross sections  $\sigma \times \text{Br}(Z' \rightarrow W^+W^-)$  for  $Z'_{\text{SSM}}$  and reference model are calculated from PYTHIA 6.409 [31] with a  $K$ -factor of 1.9, and given by dash-dotted curves. Labels attached to the curves for the  $Z'_{\text{SSM}}$  cross section correspond to the considered mixing factor  $\xi$ . Upper panel: ATLAS data for  $36.1 \text{ fb}^{-1}$  [11], lower panel: CMS data for  $35.9 \text{ fb}^{-1}$  [12].

we compute the LHC  $Z'$  production cross-section multiplied by the branching ratio into two  $W$  bosons,  $\sigma(pp \rightarrow Z') \times \text{Br}(Z' \rightarrow W^+W^-)$ , as a function of two parameters ( $M_{Z'}$ ,  $\xi$ ), and compare it with the limits established by the ATLAS and CMS experiments.

ATLAS [11] and CMS [12] analyzed the  $W^+W^-$  production in process (1) through the semileptonic and hadronic final states, respectively. Our strategy in the

present analysis is to use the SM backgrounds that have been carefully evaluated by the experimental collaborations and we simulate only the  $Z'$  signal. Figure 1 shows the observed and expected 95% C.L. upper limits on the production cross-section times the branching fraction for  $Z' \rightarrow W^+W^-$  as a function of  $Z'$  mass,  $M_{Z'}$ . The data analyzed comprises pp collisions at  $\sqrt{s} = 13 \text{ TeV}$ , recorded by the ATLAS ( $36.1 \text{ fb}^{-1}$ ) and CMS




FIG. 2.  $Z'$  exclusion regions in the two-dimensional plane of  $(M_{Z'}, \xi)$  obtained from CDF (Tevatron) [6], precision electroweak (EW) data [5] and preliminary LHC data as analyzed here. The vertical dot-dashed line corresponds to the  $Z'_{\text{SSM}}$  mass constraints obtained from the DY process at the LHC [3,4]. Left panel: ATLAS data for  $36.1 \text{ fb}^{-1}$  [11], right panel: CMS data for  $35.9 \text{ fb}^{-1}$  [12]. Exclusion plots with  $100 \text{ fb}^{-1}$  of data correspond to an extrapolation of the expected sensitivity.

( $35.9 \text{ fb}^{-1}$ ) detectors at the LHC [11,12]. The inner (green) and outer (yellow) bands around the expected limits represent  $\pm 1\sigma$  and  $\pm 2\sigma$  uncertainties, respectively. Also shown are theoretical production cross sections  $\sigma \times \text{Br}(Z' \rightarrow W^+W^-)$  for  $Z'_{\text{SSM}}$ , calculated from PYTHIA 6.409 [31] adapted for such kind of analysis.<sup>2</sup> Higher-order QCD corrections for the SM and  $Z'$  boson cases were estimated using a  $K$ -factor, for which we adopt a mass-independent value of 1.9 [32–34] (however, note that also a lower value close to unity has been given [35]). These theoretical curves for the cross sections, in descending order, correspond to values of the  $Z$ - $Z'$  mixing factor  $\xi$  from 0.01 to 0.0005. The intersection points of the expected (and observed) upper limits on the production cross-section with these theoretical cross-sections for various  $\xi$  give the corresponding lower bounds on  $(M_{Z'}, \xi)$  displayed in Fig. 2. The line with the attached label “Reference model” indicates PYTHIA defaults (except for the above-mentioned  $K$ -factor) which is commonly used for mass exclusion regions. We found that the expected (observed) exclusion limits are  $M_{Z'} < 3.7(3.8) \text{ TeV}$  (ATLAS) and  $M_{Z'} < 3.2(3.5) \text{ TeV}$  (CMS).

<sup>2</sup>Contributions from  $WW$  fusion are not taken into account, since  $\text{Br}(Z' \rightarrow W^+W^-)$  is restricted to the value of 2% for  $M_{Z'} \leq 3.8 \text{ TeV}$  in the parameter region of interest [green and yellow bands shown in Fig. 1(a)], and actually below 10% up to  $M_{Z'} = 5 \text{ TeV}$ . Notice that the reference model predicts  $\text{Br}(Z' \rightarrow W^+W^-) \approx 2\%$ .

In Fig. 2, we collect limits on the  $Z'$  parameters, starting with the Tevatron studies of diboson  $W^+W^-$  pair production [6]. The limits on  $\xi$  and  $M_{Z'}$  at the Tevatron assume that no decay channels into exotic fermions or superpartners are open to the  $Z'$ . Otherwise, the limits would be moderately weaker. Interestingly, Fig. 2 shows that at heavy  $Z'$  masses, the limits on  $\xi$  obtained from the ATLAS and CMS diboson resonance production data at the LHC at 13 TeV are stronger than those derived from the global analysis of the precision electroweak data [5].

Also, here we have extrapolated the experimental sensitivity curves for higher expected luminosity downwards by a factor of  $1/\sqrt{D}$  where  $D$  is the ratio of the expected integrated luminosity of  $100 \text{ fb}^{-1}$  that will be collected at Run 2 by 2018 to the current integrated luminosities of  $36.1 \text{ fb}^{-1}$  and  $35.9 \text{ fb}^{-1}$ . It is clear that further improvement on the constraining of this mixing can be achieved from the analysis of such data.

### III. CONCLUDING REMARKS

If a new  $Z'$  boson exists in the mass range  $\sim 4\text{--}5 \text{ TeV}$ , its discovery is possible at the LHC in the Drell–Yan channel. Moreover, the detection of the  $Z' \rightarrow W^+W^-$  mode is eminently possible and would give valuable information on the  $Z$ - $Z'$  mixing. This paper presents an analysis of  $Z$ - $Z'$  mixing in the process of  $W$  pair production. The analysis is based on preliminary pp collision data at a center-of-mass energy  $\sqrt{s} = 13 \text{ TeV}$ ,

collected by the ATLAS and CMS experiments at the LHC. We analyze the popular  $Z'_{SSM}$  model and determine limits on its mass,  $M_{Z'}$ , as well as on the  $Z$ - $Z'$  mixing (angle) factor,  $\xi$ . We present the  $Z'$  exclusion region in the  $\xi - M_{Z'}$  plane for the first time by using these data. The exclusion limits represent a large improvement over previously published results obtained at the Tevatron, and also over precision electroweak data and results obtained from proton-proton collisions at  $\sqrt{s} = 7$  and 8 TeV. These are the most stringent exclusion limits to date on the  $\xi - M_{Z'}$  plane. Further improvement on the

constraining of this mixing can be achieved from the analysis of data which will be collected at higher luminosity in the near future at Run 2 of the LHC.

## ACKNOWLEDGMENTS

This research has been partially supported by the Abdus Salam ICTP (TRIL Programme) and the Belarusian Republican Foundation for Fundamental Research. The work of P. O. has been supported by the Research Council of Norway.

- 
- [1] C. Patrignani *et al.* (Particle Data Group), Review of particle physics, *Chin. Phys. C* **40**, 100001 (2016).
- [2] P. Langacker, The physics of heavy  $Z'$  gauge bosons, *Rev. Mod. Phys.* **81**, 1199 (2009).
- [3] M. Aaboud *et al.* (ATLAS Collaboration), Search for new high-mass phenomena in the dilepton final state using 36.1 fb<sup>-1</sup> of proton-proton collision data at  $\sqrt{s} = 13$  TeV with the ATLAS detector, [arXiv:1707.02424](https://arxiv.org/abs/1707.02424).
- [4] (CMS Collaboration), Report No. CMS-PAS-EXO-15-005.
- [5] J. Erler, P. Langacker, S. Munir, and E. Rojas, Improved constraints on Z-prime bosons from electroweak precision data, *J. High Energy Phys.* **08** (2009) 017.
- [6] T. Aaltonen *et al.* (CDF Collaboration), Search for  $WW$  and  $WZ$  Resonances Decaying to Electron, Missing  $E_T$ , and Two Jets in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV, *Phys. Rev. Lett.* **104**, 241801 (2010).
- [7] V. V. Andreev and A. A. Pankov, *Yad. Fiz.* **75**, 67 (2012) [Constraints on the  $Z - Z'$  mixing angle from data measured for the process  $e^+e^- \rightarrow W^+W^-$  at the LEP2 collider, *Phys. At. Nucl.* **75**, 76 (2012)].
- [8] S. Schael *et al.* (ALEPH and DELPHI and L3 and OPAL and SLD Collaborations and LEP Electroweak Working Group and SLD Electroweak Group and SLD Heavy Flavour Group), Precision electroweak measurements on the  $Z$  resonance, *Phys. Rep.* **427**, 257 (2006).
- [9] M. Dittmar, A. S. Nicollerat, and A. Djouadi, Z-prime studies at the LHC: An update, *Phys. Lett. B* **583**, 111 (2004).
- [10] S. Godfrey and T. Martin,  $Z'$  discovery reach at future hadron colliders: A Snowmass white paper, [arXiv:1309.1688](https://arxiv.org/abs/1309.1688).
- [11] The ATLAS Collaboration, Report No. ATLAS-CONF-2017-051.
- [12] The CMS Collaboration, Report No. CMS-PAS-B2G-17-001.
- [13] G. Altarelli, B. Mele, and M. Ruiz-Altaba, Searching for new heavy vector bosons in  $p\bar{p}$  colliders, *Z. Phys. C* **45**, 109 (1989); Erratum, *Z. Phys. C* **47**, 676(E) (1990).
- [14] A. A. Pankov and N. Paver, Probing  $Z$ - $Z'$  mixing at future  $e^+e^-$  colliders, *Phys. Rev. D* **48**, 63 (1993).
- [15] V. V. Andreev, G. Moortgat-Pick, P. Osland, A. A. Pankov, and N. Paver, Discriminating  $Z'$  from anomalous trilinear gauge coupling signatures in  $e^+e^- \rightarrow W^+W^-$  at ILC with polarized beams, *Eur. Phys. J. C* **72**, 2147 (2012).
- [16] V. V. Andreev, P. Osland, and A. A. Pankov, Precise determination of  $Z$ - $Z'$  mixing at the CERN LHC, *Phys. Rev. D* **90**, 055025 (2014).
- [17] V. V. Andreev, A. A. Pankov, and V. A. Bednyakov, Effects of  $ZZ'$  mixing in double W-bosons production at the large hadron collider, *Yad. Fiz.* **78**, 775 (2015) *Phys. At. Nucl.* **78**, 725 (2015).
- [18] V. M. Abazov *et al.* (D0 Collaboration), Search for Resonant  $WW$  and  $WZ$  Production in  $p\bar{p}$  Collisions at  $s = 1.96$  TeV, *Phys. Rev. Lett.* **107**, 011801 (2011).
- [19] A. M. Sirunyan *et al.* (CMS Collaboration), Search for massive resonances decaying into  $WW$ ,  $WZ$  or  $ZZ$  bosons in proton-proton collisions at  $\sqrt{s} = 13$  TeV, *J. High Energy Phys.* **03** (2017) 162.
- [20] The ATLAS Collaboration, Report No. ATLAS-CONF-2015-073.
- [21] CMS Collaboration, Report No. CMS-PAS-B2G-16-021.
- [22] D. Benchechroun, C. Driouichi, and A. Hoummada, Search for  $Z' \rightarrow WW$  at LHC with the ATLAS detector, *Eur. Phys. J. direct A* **3**, 1 (2001).
- [23] P. Langacker and M. x. Luo, Constraints on additional  $Z$  bosons, *Phys. Rev. D* **45**, 278 (1992).
- [24] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky, and W. K. Tung, New generation of parton distributions with uncertainties from global QCD analysis, *J. High Energy Phys.* **07** (2002) 012.
- [25] G. Aad *et al.* (ATLAS Collaboration), Search for new phenomena in the  $WW$  to  $\ell\nu\ell'\nu'$  final state in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *Phys. Lett. B* **718**, 860 (2013).
- [26] G. Aad *et al.* (ATLAS Collaboration), Measurement of  $W^+W^-$  production in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector and limits on anomalous  $WWZ$  and  $WW\gamma$  couplings, *Phys. Rev. D* **87**, 112001 (2013); Erratum, *Phys. Rev. D* **88**, 079906 (2013).
- [27] V. Barger, P. Langacker, and H. S. Lee, Six-Lepton Z-Prime Resonance at the LHC, *Phys. Rev. Lett.* **103**, 251802 (2009).

- [28] E. Salvioni, G. Villadoro, and F. Zwirner, Minimal Z-prime models: Present bounds and early LHC reach, *J. High Energy Phys.* **11** (2009) 068.
- [29] J. M. Cornwall, D. N. Levin, and G. Tiktopoulos, Derivation of gauge invariance from high-energy unitarity bounds on the matrix, *Phys. Rev. D* **10**, 1145 (1974); Erratum, *Phys. Rev. D* **11**, 972(E) (1975).
- [30] B. W. Lee, C. Quigg, and H. B. Thacker, The Strength of Weak Interactions at Very High-Energies and the Higgs Boson Mass, *Phys. Rev. Lett.* **38**, 883 (1977).
- [31] T. Sjostrand, S. Mrenna, and P. Z. Skands, PYTHIA 6.4 Physics and Manual, *J. High Energy Phys.* **05** (2006) 026.
- [32] S. Frixione, A next-to-leading order calculation of the cross-section for the production of  $W^+W^-$  pairs in hadronic collisions, *Nucl. Phys.* **B410**, 280 (1993).
- [33] N. Agarwal, V. Ravindran, V. K. Tiwari, and A. Tripathi, Next-to-leading order QCD corrections to  $W^+W^-$  production at the LHC in Randall Sundrum model, *Phys. Lett. B* **690**, 390 (2010).
- [34] T. Gehrmann, M. Grazzini, S. Kallweit, P. Maierhöfer, A. von Manteuffel, S. Pozzorini, D. Rathlev, and L. Tancredi,  $W^+W^-$  Production at Hadron Colliders in Next to Next to Leading Order QCD, *Phys. Rev. Lett.* **113**, 212001 (2014).
- [35] Y. M. Bai, L. Guo, X. Z. Li, W. G. Ma, and R. Y. Zhang, Revisiting the large extra dimension effects on  $W$ -pair production at the LHC in next-to-leading order QCD, *Phys. Rev. D* **85**, 016008 (2012).