

Inflation in an external four-form gauge field

Jeff Greensite

*Physics and Astronomy Department, San Francisco State University,
San Francisco, California 94132, USA*

(Received 15 May 2017; published 14 September 2017)

We consider the possibility that the Universe, viewed as a three-brane, originated in a region of strong external field strength due to a four-form gauge field in the bulk. It is shown that in a scenario of this kind inflation is generic for a wide range of initial conditions. This is true even for small field inflation with a simple quadratic inflaton potential, as well as for Higgs potentials with the initial field well away from the local maximum, not necessarily starting from rest. The power spectrum, spectral index, and r parameter help to constrain parameters in this scenario, and $r < 0.1$ favors Higgs potentials.

DOI: [10.1103/PhysRevD.96.063512](https://doi.org/10.1103/PhysRevD.96.063512)

I. INTRODUCTION

The view that our Universe should be viewed as a three-brane in higher dimensions has been advocated, in various forms, by many authors, e.g. [1–7]. But three branes couple in a natural way to an Abelian four-form gauge field in the bulk. If the embedding coordinates of the brane are denoted $\phi^a(x)$, and the brane is charged with respect to the gauge field, then the corresponding interaction term is

$$S_A = \frac{q_0}{4!} \int d^4x A_{abcd}[\phi(x)] \epsilon^{\alpha\beta\gamma\delta} \partial_\alpha \phi^a \partial_\beta \phi^b \partial_\gamma \phi^c \partial_\delta \phi^d. \quad (1)$$

It is interesting to ask what effect a strong external four-form gauge field might have had on the dynamics of the early Universe, if the Universe were exposed to such a field at that early time.

Of course such a question can only be answered in the context of a specific model. The model I will consider here was proposed in [8]. It consists of S_A plus the usual action of the standard model fields. In addition, there is the Einstein-Hilbert action, where the metric depends on the embedding, and an inflaton action. What is a little non-standard is that the inflaton fields are regarded as a subset of the $D + 1 + N$ embedding coordinates, while the induced metric $g_{\mu\nu}$ is taken to depend only on the remaining embedding coordinates. Explicitly,

$$S = S_{\text{SM}} + S_A + \frac{1}{16\pi G} \int d^4x \sqrt{-g} R - \sigma^4 \int d^4x \sqrt{-g} \left(\frac{1}{2} g^{\mu\nu} \partial_\mu \phi^s \partial_\nu \phi^s + U(\phi) \right), \quad (2)$$

where S_{SM} is the action of standard model (and possibly beyond-standard-model) fields, and

$$g_{\mu\nu} = \partial_\mu \phi^A \eta_{AB} \partial_\nu \phi^B, \quad A, B = 0, 1, \dots, D \quad (3)$$

is the induced metric of a three-brane in a $(D + 1)$ -dimensional Minkowski space. The constant σ has the dimensions of mass. The remaining N coordinates

$$\{\phi^s, s = D + 1, \dots, D + 1 + N\} \quad (4)$$

are identified with inflaton fields. We adopt the convention that upper case Latin indices run from 0 to D , indices r, s run from $D + 1$ to $D + N$, and all other lower case Latin indices run from 0 to $D + N$. It is convenient to define

$$\varphi^s = \sigma^2 \phi^s \quad \text{and} \quad V(\varphi) = \sigma^4 U(\phi), \quad (5)$$

so that the inflaton field and the potential have the conventional dimensions. We will consider in detail two specific examples, namely a simple quadratic potential

$$V(\varphi) = \frac{1}{2} m^2 \varphi^s \varphi^s, \quad (6)$$

and a Higgs potential

$$V(\varphi) = \lambda (\varphi^s \varphi^s - m^2)^2, \quad (7)$$

where, in the latter case, the inflaton starts out at $\varphi^s \varphi^s < m^2$.

In Sec. II below we review, following Ref. [8], the equations of motion of this system, and then specialize to the simplest nontrivial case of a two-component inflaton and a constant external field strength, leading to early-universe dynamical equations for the inflaton zero mode and the metric scale factor. In Sec. III these equations are simplified to something analogous to slow-roll equations, although in contrast to a slow roll down a potential hill the dynamics results in a kind of spiral motion in inflaton field space towards the minimum of the inflaton potential. What is going on is that while the inflaton potential tends to pull the inflaton field towards the minimum of the potential, the external field provides a counterbalancing

velocity-dependent force analogous to a $\mathbf{v} \times \mathbf{B}$ force, orthogonal to gravitational friction, away from the minimum. It is shown that inflation in this model does not require fine-tuning of either couplings or initial conditions, even in the small-field case, as illustrated by numerical solutions in Sec. IV. Scalar field perturbations are considered in Sec. V, and the power spectrum, spectral index n_s , and r parameter are expressed in terms of the parameters of the model and the initial state. It has been inferred from the Planck data that the tensor-to-scalar ratio is $r < 0.1$. This fact favors the Higgs potential over the quadratic potential in the external field scenario, although both are inflationary at small fields. Other aspects of this scenario are discussed in Secs. VI through VIII, with conclusions in Sec. IX.

It should be mentioned that ours is not the only inflationary scenario that has been proposed involving a kind of orbital or spiral motion. There is, e.g. the ‘‘Spinflation’’ picture [9] which considers the angular motion of a braneworld in the extended throat of a Calabi-Yau manifold, and a ‘‘Spiral Inflation’’ scenario [10] which introduces a potential which is nonperiodic in an angular variable in inflaton field space. These scenarios are very different from the one presented here, where there is a flat metric in the bulk, inflaton potentials of a standard form, and an external four-form gauge field, but they do share the idea that orbital motion of some kind may be an important component in the inflationary process.

II. EQUATIONS OF MOTION

Variation of the action (2) with respect to the embedding coordinates ϕ^A leads to equations of motion

$$2\eta_{AB}\partial_\mu(E^{\mu\nu}\partial_\nu\phi^B) - \frac{q_0}{4!}F_{ABCD}\epsilon^{\alpha\beta\gamma\delta}\partial_\alpha\phi^a\partial_\beta\phi^b\partial_\gamma\phi^c\partial_\delta\phi^d = 0, \quad (8)$$

where

$$E^{\mu\nu} \equiv \frac{1}{2}\sqrt{-g}\left\{-\frac{1}{8\pi G}G^{\mu\nu} + T^{\mu\nu}\right\}, \quad (9)$$

with $G^{\mu\nu}, T^{\mu\nu}$ the Einstein and stress-energy tensors of $S - S_A$. Variation of the action with respect to the inflaton fields φ^s leads to the equations of motion

$$\partial_\mu(\sqrt{-g}g^{\mu\nu}\partial_\nu\varphi^s) - \sqrt{-g}\frac{\partial V}{\partial\varphi^s} + \frac{q_0}{4!\sigma^2}F_{sabcd}\epsilon^{\alpha\beta\gamma\delta}\partial_\alpha\phi^a\partial_\beta\phi^b\partial_\gamma\phi^c\partial_\delta\phi^d = 0, \quad (10)$$

where F_{abcd} is the field strength

$$F_{fabcd} = \frac{\partial A_{abcd}}{\partial\phi^f} - \frac{\partial A_{fbcd}}{\partial\phi^a} + \frac{\partial A_{facd}}{\partial\phi^b} - \frac{\partial A_{fabd}}{\partial\phi^c} + \frac{\partial A_{fabc}}{\partial\phi^d} \quad (11)$$

corresponding to the four-form gauge field. These equations are of course supplemented by the usual equations of motion of the standard model fields, which we ignore for now.

For the purposes of simplified cosmology it is sufficient to assume that the induced metric on the three-brane has the usual Friedmann-Lemaître form with scale factor $a(t)$, which requires a minimum of five coordinates ϕ^A in the bulk. For zero spatial curvature, we can choose the embedding [11,12]

$$\begin{aligned} \phi^0 &= \frac{1}{2}\left\{a(t) + \int^t \frac{dt'}{da/dt'} + a(t)r^2\right\} \\ \phi^1 &= a(t)r\cos(\theta) \\ \phi^2 &= a(t)r\sin(\theta)\cos(\chi) \\ \phi^3 &= a(t)r\sin(\theta)\sin(\chi) \\ \phi^4 &= \frac{1}{2}\left\{a(t) - \int^t \frac{dt'}{da/dt'} - a(t)r^2\right\}, \end{aligned} \quad (12)$$

and the remaining inflaton coordinates are numbered ϕ^5, ϕ^6 .

Next we suppose that there is a constant field strength in the bulk oriented orthogonal to the inflaton plane, with the gauge field taken to have components

$$\begin{aligned} A_{5123}[\phi] &= -\frac{1}{2}B\phi^6, \\ A_{6123}[\phi] &= \frac{1}{2}B\phi^5. \end{aligned} \quad (13)$$

The four-form gauge field A_{abcd} is antisymmetric under permutations of indices, but apart from (13) and components obtained from (13) by permutation, it is assumed that all other components vanish.

For an initial discussion of inflation in this scenario, we make the usual simplifying assumptions of spatial homogeneity and isotropy, taking, in particular,

$$\phi^{5,6}(x, y, z, t) = \phi^{5,6}(t), \quad (14)$$

and $\phi^a = 0$ for $a > 6$. In conjunction with (13), this has the consequence that

$$F_{ABCD}\epsilon^{\alpha\beta\gamma\delta}\partial_\alpha\phi^a\partial_\beta\phi^b\partial_\gamma\phi^c\partial_\delta\phi^d = 0. \quad (15)$$

This is because two of the indices $abcd$ must be 5 and 6, so the expression necessarily includes at least one space derivative of φ^s , which vanishes according to (14). Then the

equation of motion (8) is satisfied by $E^{\mu\nu} = 0$, which are the standard Einstein field equations. For a Friedmann-Lemaître metric, disregarding the other standard model fields, the Einstein equations are just the conventional expressions for the $a(t)$ scale factor coupled to a pair of scalar fields,

$$\begin{aligned}\frac{\dot{a}^2}{a^2} &= \frac{8\pi G}{3} \left(\frac{1}{2} \partial_t \varphi^s \partial_t \varphi^s + V(\varphi) \right), \\ \frac{\ddot{a}}{a} &= \frac{8\pi G}{3} (-\partial_t \varphi^s \partial_t \varphi^s + V(\varphi)).\end{aligned}\quad (16)$$

The equations of motion for the φ^s , however, involve the field strength

$$\begin{aligned}\partial_t^2 \varphi^5 - qB \partial_t \varphi^6 + 3 \frac{\dot{a}}{a} \partial_t \varphi^5 + \frac{\partial V}{\partial \varphi^5} &= 0, \\ \partial_t^2 \varphi^6 + qB \partial_t \varphi^5 + 3 \frac{\dot{a}}{a} \partial_t \varphi^6 + \frac{\partial V}{\partial \varphi^6} &= 0,\end{aligned}\quad (17)$$

where $q = q_0/\sigma^4$. It is not hard to verify the consistency of (16) and (17).

III. THE SLOW SPIRAL

Numerical solutions of (16) and (17), a sample of which are presented in the next section, show that for a very wide range of initial conditions $\varphi^s(0)$, $\partial_t \varphi^s(0)$, field strength qB , and parameters m , λ in the potential, the evolution of $\varphi^s(t)$ rapidly settles into a spiral in the $\varphi^5 - \varphi^6$ plane, drifting slowly towards the origin in the case of the quadratic potential, and towards $\varphi^2 = m^2$ in the case of the Higgs potential. It is the drift towards the minimum, rather than the magnitude of $\partial_t \varphi$ itself, which is “slow” in this scenario.

The spiral motion in the $\varphi^{5,6}$ plane, with a gradual drift to the minimum, is best understood as a balance of forces in a plane. Define

$$\boldsymbol{\varphi} \equiv \begin{bmatrix} \varphi^5 \\ \varphi^6 \\ 0 \end{bmatrix}, \quad \mathbf{B} \equiv \begin{bmatrix} 0 \\ 0 \\ B \end{bmatrix}, \quad \boldsymbol{\nabla} \equiv \begin{bmatrix} \partial/\partial \varphi^5 \\ \partial/\partial \varphi^6 \\ 0 \end{bmatrix}, \quad (18)$$

and $H = \dot{a}/a$ as usual. Then the inflaton equations of motion can be written as

$$\partial_t^2 \boldsymbol{\varphi} + q\mathbf{B} \times \partial_t \boldsymbol{\varphi} + 3H \partial_t \boldsymbol{\varphi} + \boldsymbol{\nabla} V = 0, \quad (19)$$

which can be thought of as the equations of motion of a particle in a plane. There are three forces on this “particle”: a force $-\boldsymbol{\nabla} V$ towards the minimum of the potential, a gravitational drag force in a direction opposite to the velocity $\partial_t \boldsymbol{\varphi}$, and a “Lorentz force” in the plane which




FIG. 1. “Force” diagrams for the slow spiral approximation, where the spiral is counterclockwise inward towards the origin for the quadratic potential (a), and clockwise outward towards $\rho = m$ for the Higgs potential (b). The approximation is that $\partial_t^2 \boldsymbol{\varphi}$ towards the origin (solid dot) is simply the centripetal acceleration required for circular motion in the $\boldsymbol{\varphi}$ plane, neglecting any acceleration beyond that.

is perpendicular to the velocity. In the standard slow-roll approximation with a single inflaton field, the second time derivative of the inflaton field is neglected. In our case the slow roll is in the radial direction, so the approximation is that the second time derivative can be equated to the centripetal acceleration required for circular motion. Referring to Fig. 1 and defining

$$\begin{aligned}\rho &\equiv |\boldsymbol{\varphi}|, & v &\equiv |\partial_t \boldsymbol{\varphi}|, & V'(\rho) &= \frac{\partial V}{\partial \rho} \\ \beta &= \begin{cases} 1 & \text{Higgs potential} \\ -1 & \text{quadratic potential} \end{cases},\end{aligned}\quad (20)$$

the equations of motion in this “slow spiral” approximation are

$$V' + \beta(qB \cos \theta + 3H \sin \theta)v = \frac{v^2 \cos^2 \theta}{\rho}, \quad (21)$$

$$qBv \sin \theta - 3Hv \cos \theta = 0. \quad (22)$$

Solving (21) for v , we have

$$v = \frac{\rho\beta}{2\cos^2\theta} \left[(qB \cos\theta + 3H \sin\theta) - \sqrt{(qB \cos\theta + 3H \sin\theta)^2 + 4V'\cos^2\theta/\rho} \right], \quad (23)$$

and from (22)

$$\tan\theta = \frac{3H}{qB}. \quad (24)$$

For the cases that we will be concerned with,

$$\theta \ll 1, \quad qB \gg H, \quad (qB)^2 \gg V'/\rho, \quad (25)$$

so that (23) simplifies to

$$v \approx -\beta \frac{V'}{qB}. \quad (26)$$

In polar coordinates, where

$$\varphi^5 = \rho \cos\alpha, \quad \varphi^6 = \rho \sin\alpha, \quad (27)$$

the equations of motion are

$$\frac{d\rho}{dt} = -v \sin\theta \approx -3H \frac{V'}{(qB)^2}, \quad (28)$$

$$\frac{d\alpha}{dt} = \beta \frac{v \cos\theta}{\rho} \approx -\frac{V'}{qB\rho}. \quad (29)$$

Assuming that the fractional variation of H is negligible in a period $v/(2\pi\rho)$, the solution of these equations is a spiral in the φ plane towards the minimum of the potential, i.e. inwards towards $\rho = 0$ for the quadratic potential or, if $\rho < m$ initially, outwards to $\rho = m$ for the Higgs potential.

IV. NUMERICAL SOLUTIONS

Next we compare the slow spiral equations to numerical solutions of (16) and (17), which make no assumptions about force balance or the time-dependence of H . In a conventional treatment with $qB = 0$, slow roll inflation in a quadratic potential requires a strong initial field ρ_0 above the Planck scale, $\rho_0 > M_P$, while small field inflation in a Higgs potential calls for an initial ρ_0 fine tuned to be extremely close to the local maximum, with v_0 very close to zero. However, only taking $qB \gg m$, the equations of motion in the external field scenario predict inflationary behavior in either potential without these special conditions.

We begin with the quadratic potential (6). Inflationary behavior is generic, requiring only that $qB \gg m$ and $\rho_0 > m$. For purposes of illustration we choose parameters to be motivated (for the Higgs potential) in the next section as follows:




FIG. 2. Numerical solution of the equations of motion (16) and (17) in a quadratic potential with initial $\rho_0 = 3m, v_0 = 0$, over a period of 2×10^6 Planck times.

$$m = 0.0083, \quad qB = 0.22 \quad (30)$$

in Planck units. For the initial condition, we take

$$\rho_0 = 3m, \quad v_0 = 0. \quad (31)$$

The trajectory in the space of $\varphi_5, \varphi_6, V(\rho)$, obtained from these initial conditions by solving (16) and (17) numerically for Planck times $0 \leq t_P \leq 2 \times 10^6$, is displayed in Fig. 2. There is some initial transient behavior which is shown in Fig. 3. This behavior can be understood qualitatively: From rest, the charged “particle” tends to fall towards the minimum of the potential, but as it falls there is a force orthogonal to the motion due to the external B field. The combination of forces induces the transient oscillatory behavior seen in Fig. 3, which dies out after $t_P = 2000$ Planck times, i.e. the transient behavior is a very small fraction of the trajectory, and dies out rapidly.




FIG. 3. The initial trajectory of the previous figure, projected into the $\varphi_5 - \varphi_6$ plane, which illustrates the initial transient behavior that dies out after roughly 2000 Planck times.





FIG. 4. Same as Fig. 2, but with a random initial velocity v_0 . The transient behavior is clearly seen in this example.

If the Universe starts out at nonzero ρ_0 , then there is no particular reason that it also starts out at rest, so we may also consider initial conditions with some random choice of v_0 . A solution of this kind, with $(\partial_t \varphi_5, \partial_t \varphi_6) = (8.4, 15.6) \times 10^{-4}$ in Planck units, and all other parameters

as above, is shown in Fig. 4, over a period of 2×10^6 Planck times. The transient behavior is much more noticeable in this case, but again it dies out after about 10,000 Planck times and the evolution settles into the slow spiral described in the previous section. The behavior of $H = \dot{a}/a$ is shown in Fig. 5(a), and we compare, in Figs. 5(b) and 5(c), the behavior of the computed $\theta(t)$, $v(t)$ to the predictions of the slow spiral equations, (24) and (26), respectively. The slow spiral equations clearly agree very well with the numerical solutions for $t_p > 10,000$, i.e. after the transient behavior has died out. The number N of e -foldings of course depends on the initial conditions and the assumed start of reheating. For the trajectory shown in these figures, $N = 483$.

Inflation is also generic for the Higgs potential when $qB \gg m$ and $m > \rho_0$, without much tuning beyond those conditions, for any coupling λ . As an illustration we carry out the computation for the Higgs potential with the same parameters $m = 0.0083$, $qB = 0.22$ as above, and an arbitrary choice of coupling $\lambda = 1$. The initial configuration is taken to be at $\rho_0 = 0.2m$, with a random choice of initial



(a) $H = \dot{a}/a$, quadratic potential

(b) $\theta(t)$, quadratic potential

(c) $v(t)$, quadratic potential

FIG. 5. For the example of the quadratic potential with a random initial v_0 : (a) The expansion rate $H = \dot{a}/a$. (b) Comparison of the computed $\theta(t)$ in the $\varphi_5 - \varphi_6$ plane with the slow spiral prediction (24). (c) Comparison of the speed $v(t)$ in the $\varphi_5 - \varphi_6$ plane with the slow spiral prediction (26).




FIG. 6. Numerical solution of the equations of motion (16) and (17) in a Higgs potential with initial $\rho_0 = 0.2m$ and random v_0 , over a period of 1×10^6 Planck times. The initial transient behavior is shown in green.

velocity $(\partial_t \varphi_5, \partial_t \varphi_6) = (-1.37, 1.50) \times 10^{-4}$ in Planck units. The result, for a duration of 1×10^6 Planck times, is shown in Fig. 6, with the transient and later behaviors shown in different colors. Again the transient behavior dies out beyond 10,000 Planck times. The Hubble variable


$H = \dot{a}/a$ vs time, is shown in Fig. 7(a), and the computed $\theta(t)$, $v(t)$, compared to the slow spiral predictions (24) and (26), are shown in Figs. 7(b) and 7(c), respectively. Once again it is clear that the slow spiral is an attractor, at least under the stated conditions $qB \gg m$ and $m > \rho_0$. For the evolution shown, where the trajectory is computed up to $\rho_{\text{final}} = \frac{7}{8}m$, the number of e -foldings is $N = 104$.

V. FLUCTUATION SPECTRUM

We now consider the equations of motion to first order in fluctuations around the homogeneous isotropic background. Denote

$$\varphi^s(x, t) = \varphi_0^s(t) + \delta\varphi^s(x, t), \quad (32)$$

where $\varphi_0^s(t), a(t)$ are solutions of the spatially homogeneous equations (16) and (17). In a standard slow-roll scenario with a single inflaton field, it is possible to choose a gauge (essentially a choice of time coordinate) such that $\delta\varphi = 0$. The transformed inflaton field is



(a) $H = \dot{a}/a$, Higgs potential

(b) $\theta(t)$, Higgs potential

(c) $v(t)$, Higgs potential

FIG. 7. For the example of the Higgs potential with a random initial v_0 : (a) The expansion rate $H = \dot{a}/a$. (b) Comparison of the computed $\theta(t)$ in the $\varphi_5 - \varphi_6$ plane with the slow spiral prediction (24). (c) Comparison of the speed $v(t)$ in the $\varphi_5 - \varphi_6$ plane with the slow spiral prediction (26).

$$\varphi'(x', t') = \varphi(x, t), \quad (33) \quad \text{where}$$

where $x' = x$ and

$$t'(x, t) = \varphi_0^{-1}[\varphi(x, t)]. \quad (34)$$

Since $t'(x, t)$ is, by definition, the time at which $\varphi_0(t') = \varphi(x, t)$, it is clear that $\varphi'(x', t') = \varphi_0(t')$ and therefore, by construction, $\delta\varphi(x, t) = 0$ in this (comoving) gauge.

With two inflaton components, it is not possible to transform to $\delta\varphi = 0$ exactly. However, in the slow spiral regime it is possible to come close to this, under certain conditions. The reason one cannot find a time coordinate such that $\varphi'(x', t') = \varphi_0(t')$ is simply because $\varphi_0(t)$ defines a linelike object, i.e. a spiral, while $\varphi(x, t)$ lies in a plane. But one can still choose $t'(x, t)$ to be such that the modulus of

$$\delta\varphi(x, t') \equiv \varphi(x, t) - \varphi_0(t') \quad (35)$$

is minimized. In this “minimal” gauge $|\delta\varphi(x', t')|$ is the distance in the φ plane between $(\varphi^5(x, t), \varphi^6(x, t))$ and the nearest point on the spiral. This distance can be no greater than half the distance between neighboring arcs of the spiral. From Eqs. (28) and (29), this upper bound on the minimal distance is

$$\Delta\rho = \frac{1}{2} \left| \frac{d\rho}{d\alpha} \right| 2\pi = \frac{3\pi H}{qB} \rho. \quad (36)$$

So in minimal gauge the magnitude of $\delta\varphi$ is as small as it can be, with a fractional deviation $\delta\rho/\rho$ of order H/qB . If the Hubble expansion rate H is very much smaller than qB , then in this gauge the fluctuations away from $\varphi_0(t)$ can be neglected by comparison to the $\delta\phi^A$, which give rise to fluctuations in the metric. We will assume that this $H \ll qB$ condition is satisfied, and that in an expansion of the action $S = S_0[\varphi_0, \phi_0^A] + \delta S$ to second order in the fluctuations the terms involving $\delta\varphi$ can be ignored relative to terms quadratic in $\delta\phi^A$.

At this point we can simply follow the approach of Maldacena [13], who derives the power spectrum in a comoving gauge where the fluctuations of a single inflaton field vanish. The only modification is to replace the single inflaton field expression $(\partial_t \varphi_0)^2$ by $v^2 = \partial_t \varphi_0^s \partial_t \varphi_0^s$. Of course, in quantizing perturbations around the Friedmann-Lemaître metric one must keep in mind that $g_{\mu\nu}$ is the induced metric (3). However, in a minimal gauge with $\delta\varphi \approx 0$, the fluctuation δS in the action depends on the fluctuations of $\phi^A = \phi_0^A + \delta\phi^A$ only through the induced metric, and therefore we have

$$\begin{aligned} Z &= \int D\delta\phi^C \exp[-(S_0 + \delta S[\partial_\mu \phi^A \eta_{AB} \partial_\nu \phi^B])] \\ &= \int D\delta\phi^C \int Dg_{\alpha\beta} \delta[g_{\mu\nu} - \partial_\mu \phi^A \eta_{AB} \partial_\nu \phi^B] e^{-(S_0 + \delta S[g_{\mu\nu}])} \\ &= \int Dg_{\alpha\beta} M[g] e^{-(S_0 + \delta S[g_{\mu\nu}])}, \end{aligned} \quad (37)$$

$$M[g] = \int D\delta\phi^C \delta[g_{\mu\nu} - \partial_\mu \phi^A \eta_{AB} \partial_\nu \phi^B]. \quad (38)$$

The measure factor $M[g]$ will not contribute at the quadratic order in metric fluctuations considered here. Then, following [13], we obtain the familiar result for the dimensionless power spectrum of the scalar curvature fluctuations

$$P_R(k) = \frac{1}{4\pi^2} \frac{H^4}{v^2}. \quad (39)$$

For the spectral index n_s , with $N(t) = \int dt H$ the number of e -foldings, we have [14]

$$\begin{aligned} n_s - 1 &= \frac{d \ln P_R}{d \ln k} \\ &\approx \left(2 \frac{d \ln H^2}{dN} - \frac{d \ln v^2}{dN} \right) \left(1 - \frac{1}{2} \frac{d \ln H^2}{dN} \right) \\ &= \frac{1}{H} \left(2 \frac{d \ln H^2}{dt} - \frac{d \ln v^2}{dt} \right) \left(1 - \frac{1}{2H} \frac{d \ln H^2}{dt} \right), \end{aligned} \quad (40)$$

and for the r parameter

$$r = \frac{P_{grav}}{P_R} = \frac{\frac{2}{\pi^2} \frac{H^2}{M_p^2}}{\frac{1}{4\pi^2} \frac{H^4}{v^2}} = 8 \frac{v^2}{M_p^2 H^2}, \quad (41)$$

where M_p is the Planck mass. Now we apply the slow spiral equations (26) and (28), using also the approximation to the Friedmann equation $H^2 \approx V(\rho)/3M_p$. Then we have

$$\begin{aligned} P_R^{1/2} &= \frac{1}{6\pi M_p^2} \frac{qB V(\rho)}{|V'(\rho)|} \\ n_s - 1 &= -\frac{6}{(qB)^2} \left(\frac{V^2}{V} - V'' \right) \left(1 + \frac{3}{2} \frac{1}{(qB)^2} \frac{V^2}{V} \right) \\ r &= \frac{24}{(qB)^2} \frac{V^2}{V}, \end{aligned} \quad (42)$$

and from the Planck 2015 [15] and earlier data it is known that

$$P_R^{1/2} \approx 4.6 \times 10^{-5} \quad n_s - 1 \approx -0.04 \quad r < 0.1 \quad (43)$$

at the pivot scale $k = 0.05 \text{ Mpc}^{-1}$.

A. Quadratic potential

For the quadratic potential $V(\rho) = \frac{1}{2} m^2 \rho^2$ these expressions become

$$P_R^{1/2} = \frac{1}{12\pi M_P^2} qB \rho, \quad n_s - 1 = -\frac{6m^2}{(qB)^2}, \quad r = 48 \left(\frac{m}{qB} \right)^2. \quad (44)$$

But from the Planck data on the spectral index, $n_s - 1 \approx -0.04$, these equations imply that $r \approx 0.32$, which is incompatible with the Planck data for r . On these grounds, the quadratic potential is ruled out.

B. Higgs potential

For the Higgs potential $V(\rho) = \lambda(\rho^2 - m^2)^2$ the corresponding expressions are

$$\begin{aligned} P_R^{1/2} &= \frac{1}{24\pi M_P^2} \frac{qB}{\rho} |\rho^2 - m^2| \\ n_s - 1 &\approx -\frac{24}{(qB)^2} \lambda(\rho^2 + m^2) \\ r &= 384 \frac{\lambda \rho^2}{(qB)^2}. \end{aligned} \quad (45)$$

The coupling λ can be entirely eliminated from these equations by writing

$$m = m_0 \lambda^{-1/4}, \quad qB = (qB)_0 \lambda^{1/4}, \quad \rho = \alpha m, \quad (46)$$

and then the constants m_0 , $(qB)_0$, and α at the pivot scale are determined, given P_R , n_s and r . Unfortunately, we only have an upper bound for r , so the best one can do so far is to determine μ , b , α as a function of $r < 0.1$. For sufficiently small r we find the approximate solution

$$\begin{aligned} m &\approx \sqrt{24\pi} \left(\frac{P_R}{384} \right)^{1/4} \left(\frac{r}{\lambda} \right)^{1/4} M_P \\ \rho_* &\approx \sqrt{\frac{24r}{384(1-n_s)}} m \\ &= 24 \sqrt{\frac{\pi}{384(1-n_s)}} \left(\frac{P_R}{384} \right)^{1/4} \frac{r^{3/4}}{\lambda^{1/4}} M_P \\ qB &\approx 24 \sqrt{\frac{\pi}{(1-n_s)}} \left(\frac{P_R}{384} \right)^{1/4} r^{1/4} \lambda^{1/4} M_P, \end{aligned} \quad (47)$$

where ρ_* is ρ at the pivot scale. To get some numerical feeling for this, suppose r is at the experimental upper limit $r = 0.1$ and $\lambda = 1$. Then, using the data (43), we find from (45)

$$\rho_* = 3.6 \times 10^{-3} M_P, \quad m = 8.3 \times 10^{-3} M_P, \quad qB = 0.22 M_P. \quad (48)$$

We can also compute the running of the spectral index

$$\begin{aligned} \frac{dn_s}{d \ln k} &= \frac{dn_s}{dN} \frac{dN}{d \ln k} = \frac{1}{H} \frac{dn_s}{dt} (1 + \epsilon) \approx -\frac{1}{H} \frac{96\lambda}{(qB)^2} \rho \frac{d\rho}{dt} \\ &= 288\lambda^2 \frac{\rho^2(\rho^2 - m^2)}{(qB)^4} \end{aligned} \quad (49)$$

which, evaluated at $\rho = \rho_*$ in our $r = 0.1$ example, is

$$\frac{dn_s}{d \ln k} = -0.9 \times 10^{-4}. \quad (50)$$

Note that, from (47), this value is λ -independent, and can be compared to the Planck 2015 [15] estimate of

$$\frac{dn_s}{d \ln k} = -(3.3 \pm 7.4) \times 10^{-3}. \quad (51)$$

VI. e-FOLDINGS IN THE HIGGS SLOW SPIRAL

We have seen that after a brief transient, the solution of the homogenous equations of motion (16) and (17) settles into a trajectory which is well described by the slow spiral equations of motion derived in Sec. III. We have also seen, in the previous section, that the constants m and qB are determined, for the Higgs potential, from the CMB data for P_R , n_s , r , and the Higgs coupling λ . Moreover, m and qB depend on simple fractional powers of λ , r , shown in (47). From this power dependence, and the slow spiral equations of motion, it is quite easy to see that the number N of e -foldings is independent of λ and r . For the Higgs potential, with the minimum at $\rho = m$, this number is determined by the known values of P_R , n_s , and the (as yet unknown) fractions of ρ/m at the beginning and end of inflation.

Let m_0 , $(qB)_0$ denote the values of m , qB shown in (48), which were determined from the Planck data assuming $\lambda = 1$, $r = 0.1$. Then from (47)

$$m = m_0 \lambda^{-1/4} \left(\frac{r}{0.1} \right)^{1/4}, \quad qB = (qB)_0 \lambda^{1/4} \left(\frac{r}{0.1} \right)^{1/4} \quad (52)$$

and also denote

$$\rho = R \lambda^{-1/4} \left(\frac{r}{0.1} \right)^{1/4}. \quad (53)$$

Substituting these expressions into the slow spiral equations of motion, we find

$$\begin{aligned} H^2 &= \frac{1}{3M_P} (R^2 - m_0^2)^2 \left(\frac{r}{0.1} \right) = H_0^2 \left(\frac{r}{0.1} \right) \\ v &= \frac{4R(R^2 - m_0^2)}{(qB)_0} \left(\frac{r}{0.1} \right)^{1/2} = v_0 \left(\frac{r}{0.1} \right)^{1/2} \\ \frac{dR}{dt} &= -3H_0 \frac{4R(R^2 - m_0^2)}{(qB)_0^2} \left(\frac{r}{0.1} \right). \end{aligned} \quad (54)$$

Then the number of e -foldings from the beginning of inflation at $t = 0, R = R_0$ to the end of inflation at $t = t_f, R = R_f$ is

$$\begin{aligned} N &= \int_0^{t_f} dt H = \int_{R_0}^{R_f} dR \left(\frac{dR}{dt} \right)^{-1} H \\ &= \frac{(qB)_0^2}{12} \int_{R_0}^{R_f} \frac{dR}{R(m_0^2 - R^2)} \\ &= \frac{(qB)_0^2}{24m_0^2} \log \left[\frac{R_f^2(m_0^2 - R_0^2)}{R_0^2(m_0^2 - R_f^2)} \right]. \end{aligned} \quad (55)$$

Note that both λ and r have dropped out of this expression. Since $(qB)_0$ and m_0 are determined from the Planck data, the number of e -foldings depends entirely on the fractions

$$f_0 = \frac{\rho_{\text{initial}}}{m} = \frac{R_0}{m_0}, \quad f_f = \frac{\rho_{\text{final}}}{m} = \frac{R_f}{m_0}, \quad (56)$$

in terms of which

$$\begin{aligned} N &= \frac{(qB)_0^2}{24m_0^2} \log \left[\frac{f_f^2(1 - f_0^2)}{f_0^2(1 - f_f^2)} \right] \\ &= 29.3 \log \left[\frac{f_f^2(1 - f_0^2)}{f_0^2(1 - f_f^2)} \right]. \end{aligned} \quad (57)$$

Although the number of e -foldings is λ, r -independent, the period of inflation does have an r dependence, increasing as r decreases like $1/\sqrt{r}$ as follows:

$$\begin{aligned} t_f &= \int_{R_0}^{R_f} dR \left(\frac{dR}{dt} \right)^{-1} \\ &= \frac{(qB)_0^2}{4\sqrt{3}} M_P \left(\frac{r}{0.1} \right)^{-1/2} \int_{R_0}^{R_f} \frac{dR}{R(m_0^2 - R^2)} \\ &= \frac{(qB)_0^2}{8\sqrt{3}} \frac{M_P}{m_0^4} \left(\frac{r}{0.1} \right)^{-1/2} \left\{ \log \left[\frac{R_f^2(m_0^2 - R_0^2)}{R_0^2(m_0^2 - R_f^2)} \right] m \right. \\ &\quad \left. - \frac{m_0^2(R_f^2 - R_0^2)}{(m_0^2 - R_0^2)(m_0^2 - R_f^2)} \right\}. \end{aligned} \quad (58)$$

Finally, although we do not know the initial and ending points $\rho_{\text{initial}}, \rho_{\text{final}}$, it is clear that ρ_{initial} must be less than ρ at the pivot momentum. From (47)

$$\rho_* = R_* \lambda^{-1/4} \left(\frac{r}{0.1} \right)^{3/4} \quad (59)$$

where $R_* = 3.6 \times 10^{-3} M_P$ is the value of ρ_* shown in (48) for $\lambda = 1, r = 0.1$. Then defining, for $r \leq 0.1$,

$$f_* = \frac{\rho_*}{m} = \frac{R_*}{m_0} \left(\frac{r}{0.1} \right)^{1/2} = 0.43 \left(\frac{r}{0.1} \right)^{1/2}, \quad (60)$$




FIG. 8. The number of e -foldings N , in the Higgs potential case, as a function of the inflaton field $\rho = |\varphi|$ at the start and end of inflation. The latter are displayed as fractions $f_0 = \rho_{\text{initial}}/m$ and $f_f = \rho_{\text{final}}/m$ of the minimum of the Higgs potential at $\rho = m$.

it is necessary that $f_0 < f_*$. The number of e -foldings N vs the fractions f_0, f_f are displayed in Fig. 8.

VII. EXCITATIONS

It is worth mentioning, for the sake of completeness, some intriguing effects of the external four-form gauge field if the field strength persists into the late universe. It is found, when the inflaton field is quantized in the presence of the constant background four-form field strength, that there is an analogy to ordinary Landau levels, and the spectrum of the quantized field is

$$\begin{aligned} E &= \sum_k \left\{ \sqrt{k^2 + \frac{1}{4} q^2 B^2 + m^2(n_1(k) + n_2(k))} \right. \\ &\quad \left. + \frac{1}{2} qB(n_1(k) - n_2(k)) \right\} + E_0, \end{aligned} \quad (61)$$

where $n_1(k), n_2(k)$ are occupation numbers, E_0 is the ground-state energy, and the sum runs over momenta with nonzero occupation numbers. Excitations with definite energy and momentum satisfy one of two dispersion relations

$$\begin{aligned} E_1(k) &= \sqrt{k^2 + M^2 + m^2} + M, \quad \text{and} \\ E_2(k) &= \sqrt{k^2 + M^2 + m^2} - M, \end{aligned} \quad (62)$$

where $M = \frac{1}{2} qB$. It can be shown that these two types of heavy/light excitations propagate like ordinary massive particles with inertial mass $M' = \sqrt{M^2 + m^2}$, i.e. with group velocity $v = k/\sqrt{k^2 + M^2 + m^2}$, but interact gravitationally with gravitational masses $M' \pm M$. Thus, there is

an apparent violation of the Principle of Equivalence, due to the interaction with the external field strength of the four-form gauge field. For details, cf. [8].

VIII. MANY BRANEWORLDS

Having introduced a braneworld scenario in which the braneworld can interact with an external four-form Abelian gauge field, one can of course formulate the obvious generalization: a higher-dimensional universe containing many braneworlds, interacting with one another via the four-form gauge field, which itself has dynamics and wave propagation in the bulk. The action would be

$$\begin{aligned}
 S = & \int \prod_a d\phi^a F_{bcdef}[\phi] F^{bcdef}[\phi] \\
 & + \sum_n \left\{ \frac{q_0}{4!} \int d\phi^a \wedge d\phi^b \wedge d\phi^c \wedge d\phi^d A_{abcd}[\phi(x_n)] \right. \\
 & + S_{infl}[\{\phi^s(x_n)\}, \sqrt{g(x_n)}] + S_{EH}[g_{\mu\nu}(x_n)] \\
 & \left. + S_{SM}[\{\Phi(x_n)\}, \sqrt{g(x_n)}] \right\}, \quad (63)
 \end{aligned}$$

where the sum is over braneworlds, $\phi^a(x_n)$ are the coordinates of the n th braneworld in the bulk, S_{infl} , S_{EH} , S_{SM} are the inflaton, Einstein-Hilbert, and standard model actions, with $\{\Phi(x_n)\}$ the set of standard model fields living on the n th braneworld, and

$g_{\mu\nu}(x_n) = \partial_\mu \phi^A(x_n) \eta_{AB} \partial_\nu \phi^B(x_n)$ the induced metric of the n th braneworld. At the quantum level one could even speculate, going well beyond the conjecture raised in this article, that our $D = 4$ dimensional Universe was created in a strong external gauge field via pair production of a three brane-antibrane pair.

We are content to make these speculations, but will not pursue them here. A consistent formulation would probably require making some connection to string theory.

IX. CONCLUSIONS

We have pointed out that if the Universe is to be thought of as a three-brane propagating in a higher-dimensional space, then it is natural for that three-brane to couple to a four-form gauge field in the bulk. If the very early Universe were exposed to a strong external field strength due to the four-form gauge field, then it seems that the resulting ‘‘slow spiral’’ of the inflaton field would solve one of the main problems associated with small field inflation in a Higgs (or other hilltop) potential, namely, the need for fine-tuning the initial value of the inflaton field and its time derivative. We reserve questions regarding reheating and non-Gaussianity in this scenario for later investigation.

ACKNOWLEDGMENTS

This research is supported by the U.S. Department of Energy under Grant No. DE-SC0013682.

-
- [1] T. Regge and C. Teitelboim, General Relativity a la String: A Progress Report, in *Marcel Grossmann Meeting on the Recent Progress of the Fundamentals of General Relativity*, edited by R. Ruffini (North-Holland, Amsterdam, 1977).
- [2] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, *Phys. Rev. D* **59**, 086004 (1999).
- [3] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 4690 (1999).
- [4] M. Alishahiha, E. Silverstein, and D. Tong, *Phys. Rev. D* **70**, 123505 (2004).
- [5] S. H. Henry Tye, *Lect. Notes Phys.* **737**, 949 (2008).
- [6] G. R. Dvali, G. Gabadadze, and M. Porrati, *Phys. Lett. B* **485**, 208 (2000).
- [7] M. Pavsic, *Phys. Lett. A* **283**, 8 (2001). M. Pavsic and V. Tapia, arXiv:gr-qc/0010045.
- [8] J. Greensite, *Phys. Rev. D* **95**, 023506 (2017).
- [9] D. A. Easson, R. Gregory, D. F. Mota, G. Tasinato, and I. Zavala, *J. Cosmol. Astropart. Phys.* **02** (2008) 010.
- [10] I.-S. Yang, *Phys. Rev. D* **85**, 123532 (2012).
- [11] J. Rosen, *Rev. Mod. Phys.* **37**, 204 (1965).
- [12] M. Lachize-Rey, *Astron. Astrophys.* **364**, 894 (2000).
- [13] J. M. Maldacena, *J. High Energy Phys.* **05** (2003) 013.
- [14] D. Baumann, *Physics of the Large and the Small: TASI 2009*, edited by C. Csaki and S. Dodelson (World Scientific, Singapore, 2011).
- [15] P. A. R. Ade *et al.* (Planck Collaboration), *Astron. Astrophys.* **594**, A20 (2016).