# Mass hierarchy and CP-phase sensitivity of ORCA using the Fermilab neutrino beam

Ushak Rahaman<sup>\*</sup>

Department of Physics, Indian Institute of Technology Bombay, Powai, Mumbai 400076, India

Soebur Razzaque<sup>†</sup>

Department of Physics, University of Johannesburg, PO Box 524, Auckland Park 2006, South Africa (Received 17 March 2017; published 27 October 2017)

We explore neutrino mass hierarchy determination and *CP*-phase measurement using the multimegaton water Cherenkov detector KM3NeT-Oscillation Research with Cosmics in the Abyss in the Mediterranean sea receiving neutrino beam from the Fermilab Long Baseline Neutrino Facility over a 6900 km baseline. We find that with the proposed beam luminosity of  $1.2 \times 10^{21}$  proton on target per year, it will be possible to determine mass hierarchy at  $\gtrsim 4\sigma$  confidence level within 1 year in the neutrino mode alone. A combined 1 year in neutrino and 1 year in antineutrino mode can determine hierarchy at  $\gtrsim 6\sigma$  confidence level. We also find that a nonzero *CP* phase can be detected with up to ~1.8 $\sigma$  significance after 10 years of data taking. We explore degeneracy of neutrino oscillation parameters and uncertainties in detection efficiencies affecting the results.

DOI: 10.1103/PhysRevD.96.073007

# I. INTRODUCTION

Determination of neutrino mass hierarchy and measurement of the CP phase have been widely discussed in recent years using multimegaton water/ice Cherenkov detectors and with atmospheric neutrino flux [1–12]. Such measurements are based on neutrino propagation over long baseline, through the mantle and core of the earth, for which oscillations in vacuum and in matter including Mikheev-Smirnov-Wolfenstein (MSW) resonance [13,14] and parametric enhancement effects [15,16] are relevant. Determining the mass hierarchy and measuring the CPphase are among the foremost objectives of neutrino physics as well as particle physics.

While atmospheric neutrinos can provide large statistics of events in multimegaton scale water/ice Cherenkov detectors with few GeV threshold, such as the proposed "Precision IceCube Next Generation Upgrade" (PINGU) [6] and Oscillation Research with Cosmics in the Abyss (ORCA) option of KM3NeT [8], there are a number of uncertainties that reduce the significance to hierarchy determination and CP-phase measurement. The uncertainties directly related to atmospheric neutrinos are flux normalizations and spectral shapes, presence of both neutrinos and antineutrinos in the flux, presence of both  $\nu_{e}$  and  $\nu_{\mu}$  flavors in the flux, etc. Careful analyses have been carried out to minimize these uncertainties with dedicated detector-specific simulations, resulting  $\approx 3\sigma$  significance for hierarchy determination within  $\approx 3-4$  years of data taking [17,18]. Sensitivity to the *CP*-phase  $\delta$  will be rather poor unless further upgrades in future are considered [7], giving  $\sim 1$  megaton effective mass at sub-GeV energies.

Long baseline accelerator experiments such as T2K and NO $\nu$ A currently provide no significant constraint on hierarchy, with only a weak preference to normal mass hierarchy [19,20]. Sensitivities to *CP* phase are similarly poor, with 90% C.L. exclusion of the  $\delta = (0.15, 0.83)\pi$  region for normal hierarchy and  $\delta = (-0.08, 1.09)\pi$  region for inverted hierarchy by T2K [19]. NO $\nu$ A excluded the  $\delta = (0.1, 0.5)\pi$  region for inverted hierarchy at 90% C.L. most recently by using  $\nu_e$  appearance data [21]. Future accelerator experiments such as "Deep Underground Neutrino Experiment" (DUNE) using the neutrino beam provided by Long Baseline Neutrino Facility (LBNF) at the Fermilab are expected to determine hierarchy and possibly find evidence of a nonzero *CP* phase [22].

A combination of multimegaton scale water/ice Cherenkov detector at the South Pole receiving accelerator neutrino beam from the Northern hemisphere has been explored recently to do oscillation physics [23,24]. In particular, sensitivities to mass hierarchy and *CP* phase were explored by choosing a specific example from the CERN to PINGU over a baseline of 11810 km [24]. Possibilities of exploring mass hierarchy in experimental setups with beam from CERN (Fermilab) to Lake Baikal (KM3NeT) [25] and from CERN to Super-Kamiokande [26] have been explored as well. Other, shorter, baselines have also been explored using neutrino beam and huge atmospheric detectors to determine mass hierarchy [27,28].

In this paper we explore neutrino mass hierarchy and *CP*-phase sensitivities of the KM3NeT-ORCA detector in the Mediterranean sea, receiving LBNF neutrino beam over a baseline of 6900 km. We use proposed characteristics of

ushak@phy.iitb.ac.in

srazzaque@uj.ac.za

the detector from simulations [8] such as energy resolution and efficiency to specific interaction channels. Similarly we use neutrino and antineutrino beam characteristics of the LBNF beam [22]. We investigate hierarchy sensitivity with marginalizations over all relevant oscillation parameters. We also investigate degeneracy of the oscillation parameters to measure the *CP* phase. We address particle flavor (electrons and muons) misidentification in the detector affecting our results.

We discuss briefly long baseline oscillation relevant for Fermilab to ORCA in Sec. II. In Sec. III we discuss the LBNF beam and detector characteristics as well as details of simulations. We present and discuss our results in Sec. IV and draw conclusions in Sec. V.

# **II. LONG BASELINE OSCILLATION**

The probability  $P_{\mu e}$  of  $\nu_{\mu}$  oscillating into  $\nu_{e}$  and the  $\nu_{\mu}$ survival probability  $P_{\mu\mu}$  are functions of mixing angles  $\theta_{12}$ ,  $\theta_{13}$ ,  $\theta_{23}$ , mass-squared differences  $\Delta m_{31}^2$  and  $\Delta m_{21}^2$  for three flavors and the *CP*-violating phase  $\delta$ . The baseline for the LBNF beam from the Fermilab to ORCA site, off the coast of Toulon, France [8] is L = 6900 km. Neutrinos along this trajectory do not pass through the core of the earth and a quasiconstant density approximation is reasonable for calculation of oscillation probabilities. Sensitivity to the mass hierarchy comes mostly at energies  $\gtrsim 2$  GeV and one can ignore 1–2 mixing and mass splitting in this energy range. The corresponding neutrino oscillation probabilities for normal mass hierarchy (NH) are [2,29,30]

$$P_{\mu e} = \sin^2 \theta_{23} P_A,$$
  

$$P_{\mu \mu} = 1 - \frac{1}{2} \sin^2 2\theta_{23} - \sin^4 \theta_{23} P_A$$
  

$$+ \frac{1}{2} \sin^2 2\theta_{23} \sqrt{1 - P_A} \cos \phi_X,$$
 (1)

where, for constant density approximation,

$$P_{A} = \sin^{2}2\theta_{13}^{m}\sin^{2}\frac{\phi_{31}^{m}}{2}$$

$$\phi_{X} = \tan^{-1}\left(\cos 2\theta_{13}^{m}\tan\frac{\phi_{31}^{m}}{2}\right)$$

$$+\frac{\Delta m_{31}^{2}L}{4E_{\nu}}\left(1+\frac{2VE_{\nu}}{\Delta m_{31}^{2}}\right)$$
(2)

and

$$\phi_{31}^{m} = \frac{\Delta m_{31}^{2} L}{2E_{\nu}} \sqrt{\left(\cos 2\theta_{13} - \frac{2VE_{\nu}}{\Delta m_{31}^{2}}\right)^{2} + \sin^{2}2\theta_{13}}$$
$$\cos 2\theta_{13}^{m} = \frac{\cos 2\theta_{13} - \frac{2VE_{\nu}}{\Delta m_{31}^{2}}}{\sqrt{\left(\cos 2\theta_{13} - \frac{2VE_{\nu}}{\Delta m_{31}^{2}}\right)^{2} + \sin^{2}2\theta_{13}}}$$
(3)

are the oscillation phase and mixing angle in matter, respectively. The matter potential  $V = \sqrt{2}G_F N_e$ , with  $G_F$  and  $N_e$  being the Fermi constant and average electron number density along the trajectory, respectively.

Oscillation probabilities for antineutrinos  $\bar{P}_{\mu e}$  and  $\bar{P}_{\mu \mu}$  can be written by substituting  $V \rightarrow -V$  in the above equations. It can be seen from Eqs. (3) and (2) that for antineutrinos  $\cos 2\theta_{13}^m \approx 1$  and correspondingly  $P_A(V \rightarrow -V) \approx 0$ . Therefore the antineutrino probability  $\bar{P}_{\mu e}$  in matter is suppressed. Also, for antineutrinos  $\phi_X(V \rightarrow -V) \approx \phi_{31} = \Delta m_{31}^2 L/2E_{\nu}$  and the probability  $\bar{P}_{\mu\mu}$  assumes the vacuum oscillation probability.

For inverted mass hierarchy (IH) the oscillation probabilities for the neutrino and antineutrinos are exchanged in this approximation of zero 1–2 mixing such that  $P_{\alpha\beta}^{\text{IH}} = \bar{P}_{\alpha\beta}^{\text{NH}}$  and  $\bar{P}_{\alpha\beta}^{\text{IH}} = P_{\alpha\beta}^{\text{NH}}$  [2].

Our numerical calculation of probabilities, using the preliminary reference Earth model [31], agree very well with above expectation for NH and IH for neutrinos and antineutrinos, for the same *CP*-phase values. We have used the best fit values of the neutrino oscillation parameters from Refs. [32,33] for numerical calculation.

In Fig. 1, we have plotted  $P_{\mu e}$  (left panels) and  $\bar{P}_{\mu e}$  (right panels) for NH and IH and for three values of  $\delta$ . The upper and lower panels correspond to the lower octant (LO) and higher octant (HO) of  $\theta_{23}$ , respectively. We have used a baseline of 6900 km (distance between Fermilab and the ORCA detector in the Mediterranean sea). The dominant feature of the probabilities in these plots is the 1-3 resonance at an energy  $E_{\nu}^{\rm R} = \Delta m_{31}^2 \cos 2\theta_{13}/2V \approx 6.6 \text{ GeV}$  for NH (IH) in case of  $P_{\mu e}$  ( $\bar{P}_{\mu e}$ ). The sensitivity to hierarchy mostly comes from this feature in this energy region. The effect of octant for  $\theta_{23}$  is just a shift in the overall probability values since those are proportional to  $\sin^2 \theta_{23}$  as in Eq. (1). It is clear from Fig. 1 that unlike other long baseline experiments such as NO $\nu$ A and T2K, no hierarchy- $\delta$  degeneracy [34,35] exists for this baseline and the probabilities for NH is well separated from IH.

We have plotted the survival probabilities  $P_{\mu\mu}$  (left panels) and  $\bar{P}_{\mu\mu}$  (right panels) in Fig. 2. As discussed for Eqs. (1)–(3), the probabilities  $\bar{P}_{\mu\mu}^{\rm NH} = P_{\mu\mu}^{\rm IH}$  are consistent with vacuum oscillation, which are different from the probabilities  $\bar{P}_{\mu\mu}^{\rm IH} = P_{\mu\mu}^{\rm NH}$ . These differences give sensitivity to hierarchy in the muon neutrino channel. Again, the upper and lower panels in Fig. 2 correspond to LO and HO of  $\theta_{23}$  as in Fig. 1. But the difference between the LO and HO for  $P_{\mu\mu}$  is opposite than for  $P_{\mu e}$ , as evidenced from Eq. (1).

To understand the *CP* dependence of the probabilities one cannot ignore the 1-2 mixing, however, as was done in Eq. (1). The 1-2 oscillation phase at low energies and close to the 1-2 resonance energy in constant density medium is [7]

$$\phi_{21}^{m} = \frac{\Delta m_{21}^{2} L}{2E_{\nu}} \sqrt{\left(\cos 2\theta_{12} - \frac{2VE_{\nu}}{\Delta m_{21}^{2}}\right)^{2} + \sin^{2}2\theta_{12}}.$$
 (4)



FIG. 1. Probabilities  $P_{\mu e}$  ( $\bar{P}_{\mu e}$ ) in the left (right) panels as functions of energy for different hierarchies and  $\delta$  values for a baseline of 6900 km. The upper (lower) panels show the probabilities for  $\sin^2\theta_{23} = 0.44$  (0.584).

The *CP*-dependent parts of the probabilities, after averaging over fast oscillations driven by the 1-3 oscillation phase are [7]

$$\langle P_{e\mu}^{\delta} \rangle = \frac{J_{\theta}}{2} \left[ \cos \delta \cos 2\theta_{12}^{m} \sin^{2} \frac{\phi_{21}^{m}}{2} + \frac{1}{2} \sin \delta \sin \phi_{21}^{m} \right],$$

$$\langle P_{\mu\mu}^{\delta} \rangle = -\frac{J_{\theta}}{2} \cos \delta \sin^{2} \frac{\phi_{21}^{m}}{2} \cos 2\theta_{12}^{m},$$

$$(5)$$

where

$$J_{\theta} \equiv \sin 2\theta_{23} \sin 2\theta_{12}^m \sin 2\theta_{13}^m \cos \theta_{13}^m \tag{6}$$

is the Jarlskog invariant in matter. Note that  $P_{\mu e}^{\delta} = P_{e\mu}^{-\delta}$  and the probabilities for antineutrinos can be found by changing  $V \rightarrow -V$  and  $\delta \rightarrow -\delta$  in Eq. (5). For energies above the 1–2 resonance,  $\cos 2\theta_{12}^m \approx -1$ .

Equation (5) describes the general behavior of the *CP* dependence of the numerically calculated probabilities in Figs. 1 and 2 rather well. A systematic shift in  $P_{\mu e}$  with  $\delta$ 



FIG. 2. The same as Fig. 1 but for  $P_{\mu\mu}$  and  $\bar{P}_{\mu\mu}$ .

over a large energy range in Fig. 1 is due to the *CP*-odd second term of the first line in Eq. (5) containing  $\sin \delta$ , as discussed in details in Ref. [7]. The main contribution to the *CP* sensitivity comes from this term in  $P_{\mu e}$ . The effect of *CP* in the case of  $P_{\mu\mu}$  is opposite to that of  $P_{\mu e}$  and is *CP* even because of the  $\cos \delta$  term. Therefore in Fig. 2 the  $\delta = 90^{\circ}$  and  $-90^{\circ}$  probabilities are the same.

From the above probability plots, we can conclude that since the probabilities for NH and IH are quite well separated, it is possible to get a good hierarchy sensitivity for the proposed Fermilab beam to the ORCA detector in the Mediterranean sea. The effect of *CP* on the other hand is rather mild at the probability level and will be difficult to measure.

Next we investigate the effect of marginalization over  $\theta_{23}$ on the hierarchy sensitivity from the probability plots. Figure 3 shows the probabilities for different  $\theta_{23}$  values in the LO and HO, i.e.,  $\sin^2 \theta_{23} = 0.440$  and 0.584, respectively. Both the  $P_{\mu e}$  and  $P_{\mu \mu}$  probabilities for different octants and hierarchy combinations are relatively well separated from each other. Thus the octant degeneracy cannot mimic a NH probability for IH and vice versa, and the sensitivity to hierarchy is not lost.



FIG. 3. Probabilities  $P_{\mu e}$  ( $P_{\mu \mu}$ ) in the upper (lower) panels for different octants of  $\theta_{23}$  and hierarchy combinations but with the same *CP* phase. The left (right) panels show the probabilities for the neutrino (antineutrino). We have used a baseline of 6900 km.

Finally, in Fig. 4 we have plotted the probability curves for LO, NH, and different  $\delta$  and  $\sin^2 2\theta_{13}$  combinations for both oscillation and survival cases to investigate the effect of marginalization over  $\sin^2 2\theta_{13}$  on the *CP*-phase sensitivity. In the case of  $P_{\mu e}$ , the probability for  $\sin^2 2\theta_{13} =$ 0.085 and  $\delta = -90^\circ$  is the same as the probability for  $\sin^2 2\theta_{13} = 0.085$  and  $\delta = 0$  and close to all other  $\sin^2 2\theta_{13} =$ 0.085 and  $\delta = 90^\circ$  are very similar to that for  $\sin^2 2\theta_{13} =$ 0.091 and  $\delta = 0$  and close to probabilities with other  $\delta$  and  $\sin^2 2\theta_{13} = 0.091$  combinations. In the case of survival probabilities, curves with all  $\sin^2 2\theta_{13}$  and  $\delta$  combinations are very close to each other. Therefore marginalization over  $\sin^2 2\theta_{13}$  is expected to reduce sensitivity to the *CP* phase.

# III. BEAM AND DETECTOR CHARACTERISTICS AND SIMULATION DETAILS

To do our simulations, we have used the proposed LBNF beam, designed for the DUNE program [22]. The beam is bended 7.2° westward and 5.8° downward along the far



FIG. 4. Probabilities  $P_{\mu e}$  ( $P_{\mu\mu}$ ) in upper (lower) panels for LO-NH and different  $\delta -\sin^2 2\theta_{13}$  combinations. The left (right) panels show the probabilities for the neutrino (antineutrino). We have used a baseline of 6900 km.

detector at the Sanford Underground Research Facility (SURF) 1300 km away [36]. For the location of the KM3NeT-ORCA detector, the beam will need to be bent 32.8° downward. For the ORCA detector, the beam line needs to be turned. In this case DUNE and ORCA cannot share the same beam, but can use the same flux and normalization. We have used 120 GeV beam of 1.2 MW power, corresponding to  $1.2 \times 10^{21}$  protons on target per year. The dimensions of the muon decay path are 250 m×4m and the beam peaks at around 2.5 GeV. The  $\nu_{\mu}$  beam is contaminated by  $\nu_{e}$ ,  $\bar{\nu}_{\mu}$ , and  $\bar{\nu}_{e}$ .

We also use seven energy bins in the range 1–7 GeV to calculate event rates. The bin widths are 0.27, 0.73, 0.75, 1 GeV, 1.25, 1.3, and 0.7 GeV, respectively. The typical number of signal electron (muon) neutrino events in the 1–7 GeV energy range is  $\approx 140/\text{yr}$  ( $\approx 630/\text{yr}$ ). For detector characteristics, we followed specification published in the KM3NeT letter of intention [8]. The mass of the detector is 3.6 Mton. The signal and background efficiencies for different interactions were calculated by taking the ratio of the effective volume, given in Fig. 60 of Ref. [8] with the

total detector volume of  $3.6 \times 10^6 \text{ m}^3$  in the relevant energy range. Note that this effective volume is at the trigger level and may change with different quality cuts, although the ratios may not change. With this caveat in mind we have calculated efficiencies from the abovementioned figure for different interactions inside the detector for each of the seven energy bins in the 1–7 GeV range. These interactions are  $e^-$ ,  $e^+$ ,  $\mu^-$ , and  $\mu^+$  events as well as  $\nu_{\mu}$  and  $\bar{\nu}_{\mu}$  neutral current events.

Signals for the oscillation channel come from the electrons generated by the  $\nu_{\mu} \rightarrow \nu_{e}$  oscillation and backgrounds come from muon misidentification, beam contamination, and neutral current pion interactions. Similarly, for the survival channel, signals come from muons due to the  $\nu_{\mu} \rightarrow \nu_{\mu}$  survival and backgrounds come from beam contamination and electron misidentification. We assumed a nominal value of 30% for the misidentification factor [8] and then changed it to 10% to see the effect on the *CP* phase sensitivity, and to 10% and 50% to see the effect on mass hierarchy sensitivity.

To calculate simulated experimental event rates and theoretical event rates, and then to calculate  $\Delta \chi^2$  from those, we have used the software GLoBES [37,38]. We used the best fit values of the neutrino oscillation parameters from Refs. [32,33]. For LO, the best fit value of  $\sin^2 \theta_{23}$  is 0.44 and for HO, it is 0.584. The best fit value of  $\sin^2 \theta_{13}$  is 0.085 and that of  $\sin^2 \theta_{12}$  is 0.31. For the solar mass-squared difference we have used  $\Delta m_{21}^2 = 7.49 \times 10^{-5} \text{ eV}^2$ , and for the atmospheric mass-squared difference, we have used  $\Delta m_{\mu\mu}^2 = 2.52 \times 10^{-3} \text{ eV}^2$ , where [39]

$$\Delta m_{\mu\mu}^{2} = \sin^{2}\theta_{12}\Delta m_{31}^{2} + \cos^{2}\theta_{12}\Delta m_{32}^{2} + \cos\delta\sin 2\theta_{12}\sin\theta_{13}\tan\theta_{12}\Delta m_{21}^{2}.$$
 (7)

To generate theoretical event rates, we vary the test values of  $\sin^2 2\theta_{13}$  and  $\Delta m^2_{\mu\mu}$  in their  $2\sigma$  range with 3.5% and 3% uncertainty, respectively. We also vary the test values of  $\sin^2 \theta_{23}$  in its  $3\sigma$  range [0.35, 0.65] in steps of 0.01. For the test values of  $\delta$ , we vary it in the full range of [-180°, 180°] in steps of 2.5°. We used Gaussian priors for  $\sin^2 2\theta_{23}$ ,  $\sin^2 2\theta_{13}$ , and  $|\Delta m^2_{\mu\mu}|$ .

Automatic bin-based energy smearing is done for generated events with a Gaussian smearing function in GLoBES, defined as

$$R^{c}(E, E') = \frac{1}{\sigma(E)\sqrt{2\pi}} e^{-\frac{(E-E')^{2}}{2\sigma^{2}(E)}},$$
(8)

where E' is the reconstructed neutrino energy. The corresponding energy resolution function is given by

$$\sigma(E) = \alpha E + \beta \sqrt{E + \gamma}, \tag{9}$$

where we have used  $\alpha = \beta = 0$  and  $\gamma = 0.4$  for the 1–7 GeV energy range relevant for the Fermilab-LBNF beam.

For systematics, We have used signal normalization and signal calibration errors in energy as 5% and 0.01%, respectively, for the  $\nu_e$  appearance events. For the same  $\nu_e$  appearance events the background normalization and calibration errors are 10% and 0.01%, respectively. For the  $\nu_{\mu}$  events the signal normalization and calibration errors in energy are 2.5% and 0.01% while the background normalization and calibration errors are 10% and 0.01%, respectively. Use of systematics in GLoBES has been discussed in detail in Refs. [37,38].

#### **IV. RESULTS AND DISCUSSION**

The results from our study are shown in the mass hierarchy and *CP*-phase sensitivity plots in this section. We have taken into account various uncertainties in the oscillation parameters as well as uncertainties in the event reconstruction at the detector. The square root of the  $\Delta \chi^2$ values reported here is the significance in number of  $\sigma$  or confidence level (C.L.) for one parameter.

#### A. Mass hierarchy sensitivity

The KM3NeT-ORCA has very good sensitivity to neutrino mass hierarchy, thanks to its large mass at an energy range where matter effect is significant for the baseline we have considered. As mentioned earlier, the effect of *CP*-phase uncertainty on determination of mass hierarchy is rather mild in this energy range.

In general, for true NH, the hierarchy sensitivity at  $\delta = 90^{\circ}$  is larger than  $\delta = -90^{\circ}$ . This is because of a larger event rate for  $\delta = 90^{\circ}$ , following larger probability (see Fig. 5) that arises from the *CP*-odd term in  $\langle P_{e\mu}^{\delta} \rangle$  in Eq. (5), resulting in smaller statistical uncertainty. The opposite is true for the case when the true hierarchy is IH. These can be seen immediately from our hierarchy sensitivity plots.

In Fig. 5, we have plotted the mass hierarchy sensitivity of the experimental setup described in Sec. III, after 1 year of neutrino run only (solid lines) and after  $\nu$  and  $\bar{\nu}$  runs of 1 year each, respectively (dashed lines). From the sensitivity curves we can see that it is possible to determine mass hierarchy for 1-year  $\nu$  run only (solid lines) at more than  $4\sigma$ significance, if  $\theta_{23}$  lies in the LO and the true hierarchy is either NH (upper-left panel) or IH (upper-right panel). The same is true when  $\theta_{23}$  is in the HO (lower panels). The sensitivity, however, is higher when  $\theta_{23}$  is in the HO and the NH is true (lower-left panel) and when  $\theta_{23}$  is in the LO and the IH is true (upper-right panel) as compared to the same octant but opposite hierarchy cases.

We also see in Fig. 5 that if an initial 1-year  $\nu$  run is followed by a 1-year  $\bar{\nu}$  run (dashed lines), it will be possible to determine the mass hierarchy at  $\gtrsim 6\sigma$  C.L. for all octants



FIG. 5. Mass hierarchy discovery potential after 1-year  $\nu$  run only (solid lines) and after 1-year  $\nu$  and 1-year  $\bar{\nu}$  run (dashed lines). The upper (lower) panels are for LO (HO) of  $\theta_{23}$  and the left (right) panels are for the assumed NH (IH) as the true hierarchy. Marginalizations have been done over  $\sin^2 2\theta_{13}$ ,  $\sin^2 \theta_{23}$ , and  $|\Delta m^2_{\mu\mu}|$ . We used a particle misidentification factor of 30%.

of  $\theta_{23}$  and true hierarchy combinations. Note that we have marginalized over the oscillation parameters  $\sin^2 2\theta_{13}$ ,  $\sin^2 \theta_{23}$ , and  $|\Delta m^2_{\mu\mu}|$  to compute the sensitivity curves.

In Fig. 6 we have plotted the hierarchy discovery sensitivity after 1-year  $\nu$  and 1-year  $\bar{\nu}$  run, each for the  $\nu_{\mu} \rightarrow \nu_{e}$  appearance/oscillation channel only (solid lines) and the same for the  $\nu_{\mu} \rightarrow \nu_{\mu}$  disappearance/survival channel only (dashed lines). Note that hierarchy sensitivity is dominated by the oscillation channel  $\nu_{\mu} \rightarrow \nu_{e}$ ; see Eq. (1). Even if the  $\nu_{\mu} \rightarrow \nu_{\mu}$  survival channel is



FIG. 6. Mass hierarchy discovery potential for  $\nu_{\mu} \rightarrow \nu_{e}$  oscillation channel only (appearance—solid lines) and for  $\nu_{\mu} \rightarrow \nu_{\mu}$  survival channel only (disappearance—dashed lines), for 1-year  $\nu$  and 1-year  $\bar{\nu}$  run. The octant of  $\theta_{23}$ , assumed true hierarchy, marginalizations, and particle misidentification information are the same as in Fig. 5.



FIG. 7. Mass hierarchy discovery potential after 1-year  $\nu$  and 1-year  $\bar{\nu}$  run but for particle misidentification factor 10% (solid lines) and 50% (dashed lines). The octants of  $\theta_{23}$ , assumed true hierarchy, and marginalization information are the same as in Fig. 5.



FIG. 8. *CP*-violation discovery potential after 10 years of data taking. Marginalizations have been done over  $\sin^2 2\theta_{13}$ ,  $\sin^2 \theta_{23}$ ,  $|\Delta m^2_{\mu\mu}|$ , and mass hierarchy. We used a particle misidentification factor of 30%.



FIG. 9. Same as Fig. 8 but marginalization over  $|\Delta m_{uu}^2|$  only.

ignored, it is possible to determine the hierarchy at  $\gtrsim 4\sigma$  for all octants of  $\theta_{23}$  and true hierarchy. On the other hand, the hierarchy can only be established at  $\approx 3\sigma$  level in the  $\nu_{\mu} \rightarrow \nu_{\mu}$  channel only (dashed lines). The relatively flat feature (*CP* insensitive) of the dashed curves arises from the smallness of the *CP*-even term of  $\langle P_{\mu\mu} \rangle$  in Eq. (5).

In Fig. 7 we have explored the effects of electron-muon misidentification factors on mass hierarchy discovery. The solid (dashed) lines correspond to the case when the electron-muon misidentification factor is 10% (50%). It can be seen that when LO-NH or HO-IH is the true hierarchy-octant combination, it is possible to determine hierarchy at  $\gtrsim 8\sigma$  C.L. for 10% misidentification. For HO-NH combination the significance is  $\gtrsim 10\sigma$ . Therefore the hierarchy discovery potential for this experimental setup increases significantly if the electron and muon identification capability improves. On the other hand, it can be seen that the hierarchy discovery potential degrades drastically if the muon-electron identification factor is 50%, wrong hierarchy can be excluded at  $\approx 6\sigma$  C.L. for



FIG. 10. Same as Fig. 8 but with 10% particle misidentification.

all octant and true hierarchy combinations, which is still quite good.

#### B. CP-phase sensitivity

We have studied sensitivity of ORCA, using the Fermilab-LBNF beam, to measure a nonzero *CP*-phase value both in the case of an unknown mass hierarchy and a known mass hierarchy. As mentioned earlier, the neutrino beam peaks at an energy  $\approx 2.5$  GeV and the most sensitivity to the *CP* phase at the probability level is at lower energy. Furthermore the effective mass of the ORCA detector is also rather small below  $\sim 1-3$  GeV. Both of these factors reduce sensitivity to the *CP* phase. Therefore one must consider a much longer run time.

In Fig. 8 we have shown the *CP*-phase sensitivity plots after the 10-year run in the  $\nu$  mode (solid lines) as well as 5-year in the  $\nu$  and 5-year in the  $\bar{\nu}$  mode run (dashed lines) as functions of the true  $\delta$ . We have marginalized over the test values of the oscillation parameters  $\sin^2 2\theta_{13}$ ,  $\sin^2 \theta_{23}$ ,  $|\Delta m^2_{\mu\mu}|$ , and mass hierarchy. Both the muon and electron channels have been used and the electron-muon misidentification factor is 30% in these plots. Note that in the case

# USHAK RAHAMAN and SOEBUR RAZZAQUE

that true hierarchy is NH, the 10-year  $\nu$  run gives the most sensitivity to the *CP* phase (left panels—solid lines) because of a higher event rate, resulting from higher probability, and hence smaller statistical uncertainty in the resonance channel. The same is not true for the IH because of suppressed probability for neutrinos in this case. Furthermore, a 5-year  $\nu$  and 5-year  $\bar{\nu}$  run (dashed lines) does not improve the sensitivity to the *CP* phase. This is partly because of a smaller cross section for  $\bar{\nu}$ , which reduces the overall event rate, even though  $\bar{P}_{\mu e}$  is enhanced for  $\bar{\nu}$  in case of IH.

We also see in Fig. 8 that when NH is the true hierarchy, the sensitivity to a nonzero *CP* phase is up to  $\approx 1.5\sigma$  when  $\theta_{23}$  is either in the LO or HO and for true  $\delta = \pm 90^{\circ}$  (left panels—solid lines). But for IH, the sensitivity is  $< 1.2\sigma$  for all values of the true *CP* and for either octants of  $\theta_{23}$  (right panels).

In Fig. 9 we have plotted sensitivity to the *CP* phase in a case when the mass hierarchy is known (e.g., from an early run of the discussed experimental setup) and the values of  $\theta_{13}$  and  $\theta_{23}$  are accurately measured from other experiments. Note that removing marginalization from  $\sin^2 2\theta_{13}$ ,  $\sin^2 \theta_{23}$ , and hierarchy improves significance in general but not by a large value. This is again due to relatively high (~1–3 GeV) threshold energy for the proposed ORCA detector where the effect of the *CP* phase on the oscillation probabilities is low.

We have explored the sensitivity to the *CP* phase affected by the electron-muon misidentification in Fig. 10. We used a 10% misidentification factor and in general the sensitivity improves; in particular, if the true hierarchy is NH, a nonzero *CP* phase can be measured for  $-120^{\circ} \leq \delta \leq -60^{\circ}$  and for  $60^{\circ} \leq \delta \leq 120^{\circ}$  at  $\geq 1\sigma$  C.L. Marginalizations over  $\sin^2 2\theta_{13}$ ,  $\sin^2 \theta_{23}$ ,  $|\Delta m_{\mu\mu}^2|$  and mass hierarchy have been taken into account in this case.

### V. CONCLUSIONS

We have explored a possibility to determine the neutrino mass hierarchy and *CP* phase with the proposed KM3NeT-ORCA detector in the Mediterranean sea receiving a neutrino beam from the Fermilab-LBNF, over a baseline of 6900 km. We used publicly available detector characteristics and neutrino flux to simulate events and compute sensitivities using the GLoBES software.

Detailed efficiencies for different interaction channels in the detector have been taken into account as well as smearing of the events in energy. We marginalized over uncertainties of the oscillation parameters and also took into account realistic particle misidentification factor in the detector.

The mass hierarchy can be determined with  $\gtrsim 4\sigma$  significance when the true hierarchy is either normal or inverted, and for either octant of  $\theta_{23}$  within 1 year of neutrino run. A higher significance can be achieved with 1-year neutrino and 1-year antineutrino run. For this case  $\gtrsim 4\sigma$  significance can be reached in the electron neutrino channel only.

The *CP*-phase value can only be measured in the  $\pm (60^{\circ}-120^{\circ})$  range at  $\approx 1\sigma$  C.L. for 10-year neutrino run in the case that the true mass hierarchy is normal. Reducing particle misidentification in the detector can help (up to  $\sim 1.8\sigma$  for 10% misidentification) but will not increase the sensitivity significantly.

# ACKNOWLEDGMENTS

S. R. was funded by a National Research Foundation (South Africa) Grant CPRR 2014 No. 87823, which allowed a visit by U. R. to the University of Johannesburg where most of this work was completed.

- O. Mena, I. Mocioiu, and S. Razzaque, Phys. Rev. D 78, 093003 (2008).
- [2] E. K. Akhmedov, S. Razzaque, and A. Yu. Smirnov, J. High Energy Phys. 02 (2013) 082; 07 (2013) 26(E).
- [3] W. Winter, Phys. Rev. D 88, 013013 (2013).
- [4] M. Ribordy and A. Y. Smirnov, Phys. Rev. D 87, 113007 (2013).
- [5] D. Franco, C. Jollet, A. Kouchner, V. Kulikovskiy, A. Meregaglia, S. Perasso, T. Pradier, A. Tonazzo, and V. Van Elewyck, J. High Energy Phys. 04 (2013) 008.
- [6] M. G. Aartsen et al. (IceCube PINGU), arXiv:1401.2046.
- [7] S. Razzaque and A. Yu. Smirnov, J. High Energy Phys. 05 (2015) 139.
- [8] S. Adrian-Martinez *et al.* (KM3Net), J. Phys. G 43, 084001 (2016).

- [9] S.-F. Ge, K. Hagiwara, and C. Rott, J. High Energy Phys. 06 (2014) 150.
- [10] S.-F. Ge and K. Hagiwara, J. High Energy Phys. 09 (2014) 024.
- [11] F. Capozzi, E. Lisi, and A. Marrone, Phys. Rev. D 91, 073011 (2015).
- [12] S. Adrin-Martnez et al., J. High Energy Phys. 05 (2017) 008.
- [13] L. Wolfenstein, Phys. Rev. D 17, 2369 (1978).
- [14] S. P. Mikheev and A. Yu. Smirnov, Yad. Fiz. 42, 1441 (1985) [Sov. J. Nucl. Phys. 42, 913 (1985)].
- [15] E. K. Akhmedov, Nucl. Phys. B538, 25 (1999).
- [16] Q. Y. Liu, S. P. Mikheyev, and A. Yu. Smirnov, Phys. Lett. B 440, 319 (1998).
- [17] A. Kouchner, J. Phys. Conf. Ser. 718, 062030 (2016).

MASS HIERARCHY AND CP-PHASE SENSITIVITY OF ...

- [18] M. G. Aartsen et al. (IceCube), J. Phys. G 44, 054006 (2017).
- [19] K. Abe et al. (T2K), Phys. Rev. D 91, 072010 (2015).
- [20] P. Adamson et al. (NOvA), Phys. Rev. D 93, 051104 (2016).
- [21] P. Adamson *et al.* (NOvA), Phys. Rev. Lett. **116**, 151806 (2016).
- [22] R. Acciarri et al. (DUNE), arXiv:1601.05471.
- [23] D. Fargion, D. D'Armiento, P. Desiati, and P. Paggi, Astrophys. J. 758, 3 (2012).
- [24] J. Tang and W. Winter, J. High Energy Phys. 02 (2012) 028.
- [25] S. K. Agarwalla and P. Hernandez, J. High Energy Phys. 10 (2012) 086.
- [26] C. Lujan-Peschard, G. Pagliaroli, and F. Vissani, Eur. Phys. J. C 73, 2439 (2013).
- [27] J. Brunner, arXiv:1304.6230.
- [28] C. Vallee, in 18th International Workshop on Neutrino Factories and Future Neutrino Facilities Search (NuFact16), Quy Nhon, Vietnam, August 21–27, 2016 (2016).
- [29] E. K. Akhmedov, A. Dighe, P. Lipari, and A. Y. Smirnov, Nucl. Phys. B542, 3 (1999).

- [30] E. K. Akhmedov, M. Maltoni, and A. Yu. Smirnov, J. High Energy Phys. 06 (2008) 072.
- [31] A. M. Dziewonski and D. L. Anderson, Phys. Earth Planet. Inter. 25, 297 (1981).
- [32] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, and T. Schwetz, J. High Energy Phys. 01 (2017) 087.
- [33] http://www.nu-fit.org/?q=node/45.M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado, and T. Schwetz, J. High Energy Phys. 12 (2012) 123.
- [34] O. Mena and S. J. Parke, Phys. Rev. D 70, 093011 (2004).
- [35] S. Prakash, S. K. Raut, and S. U. Sankar, Phys. Rev. D 86, 033012 (2012).
- [36] J. Strait et al. (DUNE), arXiv:1601.05823.
- [37] P. Huber, M. Lindner, and W. Winter, Comput. Phys. Commun. 167, 195 (2005).
- [38] P. Huber, J. Kopp, M. Lindner, M. Rolinec, and W. Winter, Comput. Phys. Commun. **177**, 432 (2007).
- [39] H. Nunokawa, S. J. Parke, and R. Zukanovich Funchal, Phys. Rev. D 72, 013009 (2005).