

Damping of gravitational waves by matterGordon Baym,^{1,2} Subodh P. Patil,² and C. J. Pethick^{2,3}¹*Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, Illinois 61801-3080, USA*²*The Niels Bohr International Academy, The Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*³*NORDITA, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden*

(Received 3 August 2017; published 16 October 2017)

We develop a unified description, via the Boltzmann equation, of damping of gravitational waves by matter, incorporating collisions. We identify two physically distinct damping mechanisms—collisional and Landau damping. We first consider damping in flat spacetime, and then generalize the results to allow for cosmological expansion. In the first regime, maximal collisional damping of a gravitational wave, independent of the details of the collisions in the matter is, as we show, significant only when its wavelength is comparable to the size of the horizon. Thus damping by intergalactic or interstellar matter for all but primordial gravitational radiation can be neglected. Although collisions in matter lead to a shear viscosity, they also act to erase anisotropic stresses, thus suppressing the damping of gravitational waves. Damping of primordial gravitational waves remains possible. We generalize Weinberg’s calculation of gravitational wave damping, now including collisions and particles of finite mass, and interpret the collisionless limit in terms of Landau damping. While Landau damping of gravitational waves cannot occur in flat spacetime, the expansion of the universe allows such damping by spreading the frequency of a gravitational wave of given wave vector.

DOI: [10.1103/PhysRevD.96.084033](https://doi.org/10.1103/PhysRevD.96.084033)**I. INTRODUCTION**

The opening of a new window on the universe through the ongoing observations of gravitational waves [1] underlines the importance of reexamining how they propagate through the matter in the universe, and asking what gravitational wave measurements can teach one about this matter. Half a century ago, Hawking showed that if matter could be treated in the hydrodynamic limit the damping rate of a gravitational wave would be $\gamma = 16\pi G\eta$, where G is Newton’s gravitational constant and η the viscosity of the matter [2,3]. Using this result, Goswami *et al.* [4] argued that gravitational wave observations could be used to constrain the viscosity of dark matter between the source and Earth. But, as Hawking first pointed out, there are in general too few collisions in matter for hydrodynamics to be valid, and the damping would be less than the hydrodynamic result. Reference [5] estimated damping in the almost collisionless limit by investigating the response of individual particles to a gravitational wave and found that the damping rate of the wave by nonrelativistic particles is

$$\gamma \sim \frac{Gnm}{\omega^2} \left(\frac{\bar{v}}{c} \right)^2 \frac{1}{\tau}. \quad (1)$$

Here ω is the frequency of the wave, n the particle density, m the particle mass, \bar{v} the typical particle velocity, and τ the particle-particle collision time; the damping is $\sim 1/(\omega\tau)^2$ smaller than the viscous result.

In addition to damping by collisions in matter, gravitational waves can also be attenuated by Landau damping, in which particles surf the gravitational wave and extract its energy, first proposed for gravitational waves in Ref. [6]. This effect was originally investigated in the context of plasma physics [7], then in galactic dynamics [8], and later in quantum chromodynamic plasmas [9]. In a static flat universe massive particles cannot produce Landau damping since they move more slowly than a gravitational wave. In an expanding universe, however, Landau damping becomes possible, as we show, since the expansion in the presence of matter effectively spreads the frequency of a gravitational wave. Indeed the damping of cosmological gravitational waves by noninteracting neutrinos, as examined by Weinberg [10]¹ and expanded upon in Refs. [12–15], can in fact be understood in terms of Landau damping, as we indicate below.

Our aim in this paper is to present a unified treatment of the damping of gravitational waves by matter, for arbitrary collision rates, thus encompassing the hydrodynamic and nearly collisionless limits studied earlier, as well as cosmological expansion. We begin, in Sec. II, by considering a weak gravitational wave propagating through a dilute gas of colliding particles of arbitrary mass in an otherwise flat spacetime, and calculate, in Sec. III, the response of the matter to the wave using the Boltzmann

¹Similar effects were previously studied in the context of cosmic microwave anisotropies by Misner in [11].

equation. For simplicity we work in the relaxation or collision time approximation.

As we show, in Sec. IV, the maximum damping of the amplitude of a gravitational wave with frequency ω is of order $1/(\omega\tau_H)$ where τ_H is the age of the universe. Thus collisional damping by matter of gravitational waves generated by astrophysical sources cannot provide useful information about the nature of matter in the universe. Furthermore, damping by dense environments surrounded localized sources of gravitational radiation is, as we estimate, insignificant. After a general discussion of Landau damping in Sec. VI, we generalize the Boltzmann equation results in Sec. VII to describe collisional damping by particles of arbitrary mass in the presence of an expanding cosmological background.

II. STATIC SPACETIME

Initially, we do not include the expansion of the universe, and consider rather the Minkowski space metric with a gravitational wave superimposed:

$$ds^2 = -dt^2 + g_{ij}dx^i dx^j, \quad (2)$$

where

$$g_{ij} = \delta_{ij} + h_{ij}(\vec{r}, t), \quad (3)$$

with h_{ij} the weak metric perturbation caused by a gravitational wave. We work with h_{ij} in the transverse-traceless gauge, and generally set $c = 1$.

The effects on a gravitational wave passing through matter are given in terms of the gravitational wave equation in the transverse traceless gauge

$$\partial_\mu \partial^\mu h_{ij} = \left(-\frac{\partial^2}{\partial t^2} + \nabla^2 \right) h_{ij} = -16\pi G \pi_{ij}, \quad (4)$$

where π_{ij} is the transverse traceless part of the matter stress tensor, $T_{ij,M}$, defined by

$$\pi_{ij}^i \equiv T_{j,M}^i - \frac{\delta_j^i}{3} \sum_{k=1}^3 T_{k,M}^k. \quad (5)$$

In equilibrium, $\frac{1}{3} \sum_{k=1}^3 T_{k,M}^k$ is simply the pressure P of the matter.

The effect of a gravitational wave on a particle is given in terms of the dispersion relation,

$$p_\mu p_\nu g^{\mu\nu} + m^2 = 0, \quad (6)$$

which in the present case implies that the particle energy ϵ is given by

$$\epsilon^2 = g^{ij} p_i p_j + m^2. \quad (7)$$

Thus a weak gravitational wave changes the particle dispersion relation from $\epsilon_0 = \sqrt{p^2 + m^2}$ to

$$\epsilon = \epsilon_0 + \delta\epsilon. \quad (8)$$

To first order in h_{ij}

$$\delta\epsilon = \frac{1}{2} h^{ij} \frac{p_i p_j}{\epsilon_0} = -\frac{1}{2} h_{ij} \frac{p_i p_j}{\epsilon_0}, \quad (9)$$

since to this order, $h^{ij} = -h_{ij}$.

III. BOLTZMANN EQUATION

We treat the matter as a dilute gas and calculate π_{ij} from the Boltzmann equation for the matter. We first write the nonlinear Boltzmann equation for the particle distribution function $f(r^i, p_j)$ as a function of the particle positions and canonical momenta,

$$\left(\frac{\partial}{\partial t} + \vec{\nabla}_p \epsilon \cdot \vec{\nabla}_r - \vec{\nabla}_r \epsilon \cdot \vec{\nabla}_p \right) f = \mathcal{C}, \quad (10)$$

where \mathcal{C} is the collision term. Here position gradients are taken with respect to r^i , and momentum gradients with respect to p_i . This form of the equation is valid for relativistic as well as nonrelativistic particles.

The conservation laws of energy and momentum are found by taking the moments of Eq. (10) with respect to p_i and ϵ ; assuming that collisions conserve the total energy and momentum of the particles, we find (as in standard Fermi liquid theory)

$$\frac{\partial}{\partial t} \int_p \epsilon f + \nabla_i \int_p \epsilon v^i f = \int_p \frac{\partial \epsilon}{\partial t} f, \quad (11)$$

and

$$\frac{\partial}{\partial t} \int_p p_i f + \nabla_j T_{i,M}^j = - \int_p (\nabla_{r_i} \epsilon) f, \quad (12)$$

where $\int_p \equiv g \int d^3 p / (2\pi)^3$, with g the number of internal states, e.g., spin, and

$$T_{i,M}^j = \int_p p_i v^j f = g^{jk} \int_p \frac{p_i p_k}{\epsilon} f \quad (13)$$

is the matter stress tensor.

With Eq. (9) the right side of Eq. (11) becomes

$$\int_p \frac{\partial \epsilon}{\partial t} f = \frac{1}{2} \frac{\partial h^{ij}}{\partial t} \int_p \frac{p_i p_j}{\epsilon} f = \frac{1}{2} \frac{\partial h^{ij}}{\partial t} T_{ij,M}, \quad (14)$$

so that the change in energy of the matter is given by

$$\frac{\partial E}{\partial t} = \frac{1}{2} \int d^3 r \frac{\partial h^{ij}}{\partial t} \pi_{ij}. \quad (15)$$

Only the transverse-traceless part of the stress tensor, Eq. (5), enters Eq. (14).

Note that, for a wave of the form $H(z - ct)$, say, the right side of Eq. (12) becomes

$$-\int_p \frac{\partial \epsilon}{\partial z} f = \frac{1}{2} \frac{\partial h^{ij}}{\partial t} \pi_{ij}, \quad (16)$$

indicating that as momentum q is transferred from the gravitational wave, energy q is also transferred. Since π^{ij} is itself at least of first order in h_{ij} the energy and momentum transfers are second order and higher in the amplitude of the gravitational wave.

We turn now to calculating the transverse-traceless part of the matter stress tensor, Eq. (13); to linear order in h_{ij} ,

$$\delta T_{ij,M} = \int_p \frac{p_i p_j}{\epsilon_0} \left[\delta f - \frac{\delta \epsilon}{\epsilon_0} f_0 \right] \quad (17)$$

where $\delta f = f - f_0$, with f_0 the distribution function in the absence of h_{ij} . The $\delta \epsilon$ term arises from the dependence of ϵ in the denominator on h_{ij} , Eq. (9). Subtracting out the trace, we find, after using the vanishing trace of h_{ij} and writing the equilibrium pressure of the matter as $\int_p (p^2/3\epsilon_0) f_0$, that

$$\pi_{ij} = \int_p \frac{p_i p_j}{\epsilon_0} \left[\delta f - \frac{\delta \epsilon}{\epsilon_0} f_0 \right] + h^{ij} \int_p \frac{p^2}{3\epsilon_0} f_0. \quad (18)$$

The second term of this expression is manifestly traceless; the trace of the first term vanishes since the integrations over both δf and $\delta \epsilon$, being symmetric in angles, vanish.

The latter two terms that contain f_0 can be simply combined, with an integration by parts using the transverse-traceless structure of h_{ij} , into a term proportional to $\partial f_0 / \partial \epsilon$ so that Eq. (18) becomes,

$$\pi_{ij} = \int_p \frac{p_i p_j}{\epsilon_0} \left[\delta f - \delta \epsilon \frac{\partial f_0}{\partial \epsilon} \right]. \quad (19)$$

This combination of terms falls out naturally, as we shall see, from the Boltzmann equation.

Collisions between the particles, prior to freeze-out, tend to bring the distribution function into a local equilibrium in the presence of h_{ij} :

$$f \rightarrow f_h = \frac{1}{e^{\beta(\epsilon - \mu)} \mp 1}, \quad (20)$$

where ϵ , given by Eq. (7), depends on h_{ij} ; β is the inverse temperature, and μ the particle chemical potential. Note that to first order in h_{ij} ,

$$f_h = f_0 + \delta \epsilon \frac{\partial f}{\partial \epsilon}. \quad (21)$$

For simplicity we employ a collision time approximation. Since the only disturbances relevant here involve spherical harmonics of degree greater than one, we can write the collision term as

$$C = -\frac{f - f_h}{\tau} = -\frac{1}{\tau} \left(\delta f - \delta \epsilon \frac{\partial f}{\partial \epsilon} \right), \quad (22)$$

where τ is the collision time, and $\delta f - \delta \epsilon \partial f / \partial \epsilon$ is the deviation of the distribution from local equilibrium; the additional terms commonly introduced to ensure conservation of particle number and total momentum (which involve spherical harmonics of degree zero and one) are not relevant [7]. The linearized Boltzmann equation then reduces to

$$\left(\frac{\partial}{\partial t} + \frac{1}{\tau} + \vec{v} \cdot \vec{\nabla}_{\vec{r}} \right) \delta f = \left(\vec{v} \cdot \nabla_r \delta \epsilon + \frac{1}{\tau} \delta \epsilon \right) \frac{\partial f_0}{\partial \epsilon}. \quad (23)$$

With $h_{ij}(\vec{r}, t) = e^{i(\vec{q} \cdot \vec{r} - \omega t)} h_{ij}$, and Fourier transforming in space and time we find the solution of Eq. (23),

$$\delta f = \frac{\partial f_0}{\partial \epsilon} \left(\frac{-\vec{q} \cdot \vec{v} + i/\tau}{\omega - \vec{q} \cdot \vec{v} + i/\tau} \right) \delta \epsilon. \quad (24)$$

The deviation from local equilibrium is thus given by

$$\delta f - \frac{\partial f_0}{\partial \epsilon} \delta \epsilon = - \left(\frac{\omega}{\omega - \vec{q} \cdot \vec{v} + i/\tau} \right) \frac{\partial f_0}{\partial \epsilon} \delta \epsilon. \quad (25)$$

We then find the general result,

$$\pi_{ij}(q, \omega) = \int_p \frac{p_i p_j}{\epsilon_0} \delta \epsilon \left(\frac{\omega}{\vec{q} \cdot \vec{v} - \omega - i/\tau} \right) \frac{\partial f_0}{\partial \epsilon}. \quad (26)$$

Fourier transformed back to time, the stress tensor is

$$\pi_{ij}(q, t) = - \int_p \frac{p_i p_j}{\epsilon_0} \int_{-\infty}^t dt' e^{-(iq \cdot v + 1/\tau)(t-t')} \frac{\partial f_0}{\partial \epsilon} \dot{\delta \epsilon}(t'). \quad (27)$$

The response can be written in the more general form

$$\pi_{ij}(q, \omega) = -\omega h_{ij} \int \frac{d\omega'}{2\pi} \frac{A(q, \omega')}{\omega - \omega' + i\xi}, \quad (28)$$

where the spectral function is

$$A(q, \omega) = -2 \int_p \left(\frac{p_x p_y}{\epsilon_0} \right)^2 \frac{1/\tau}{(\omega - \vec{q} \cdot \vec{v})^2 + 1/\tau^2} \frac{\partial f_0}{\partial \epsilon}, \quad (29)$$

and ξ is a positive infinitesimal. In the collisionless limit, $1/\tau \rightarrow 0$, and for relativistic particles,

$$A(q, \omega) = \pi \rho \langle (1 - \xi^2) \delta(\omega - q\xi) \rangle, \quad (30)$$

where ρ is the energy density of the excitations, and the angular brackets denote the average over $\xi \equiv \hat{q} \cdot \hat{v}$. Fourier transformed back to time,

$$\pi_{ij}(q, t) = - \int_{-\infty}^t dt' \int \frac{d\omega}{2\pi} e^{-i\omega(t-t')} A(q, \omega) \dot{h}_{ij}(t'). \quad (31)$$

The damping of gravitational waves is governed by the imaginary part of the response,

$$\Im \left(\frac{\pi_{ij}}{h_{ij}} \right) = -\omega \int_p \left(\frac{p_i p_j}{\epsilon} \right)^2 \frac{1/\tau}{(\omega - \vec{q} \cdot \vec{v})^2 + 1/\tau^2} \frac{\partial f_0}{\partial \epsilon}. \quad (32)$$

In the collision-dominated regime, $\tau \ll 1/\omega$; doing the angular averages in the integrals we have

$$\begin{aligned} \pi_{ij} &= i\tau \omega \int_p \frac{p_i p_j}{\epsilon_0} \delta\epsilon \frac{\partial f_0}{\partial \epsilon} = -\frac{i\tau \omega}{15} \int_p \frac{p^4}{\epsilon_0^2} \frac{\partial f_0}{\partial \epsilon} h_{ij} \\ &= -\eta \dot{h}_{ij}. \end{aligned} \quad (33)$$

The viscosity calculated in the relaxation time approximation is

$$\eta = - \int_p \left(\frac{p_i p_j}{\epsilon_0} \right)^2 \frac{\partial f_0}{\partial \epsilon_0} \tau, \quad (34)$$

with $i \neq j$. In this limit the damping rate of a gravitational wave, from Eq. (4), is $16\pi G\eta$, in agreement with earlier hydrodynamic treatments [2,3].

For nonrelativistic matter, $T \ll mc^2$, the $\vec{q} \cdot \vec{v}$ in the denominator of A can be neglected, and we have

$$\pi_{ij} \simeq \frac{\omega}{\omega + i/\tau} P h_{ij}. \quad (35)$$

From the imaginary part of Eq. (4), the dispersion relation of gravitational waves is

$$\omega \simeq q + \frac{8\pi G}{\omega} \frac{\pi_{ij}}{h_{ij}}, \quad (36)$$

so that the damping of a wave is given by

$$\Im \omega = \frac{8\pi G}{\omega} \Im \left(\frac{\pi_{ij}}{h_{ij}} \right). \quad (37)$$

For fully relativistic matter, in the nearly collisionless limit, to first order in $1/\tau\omega$,

$$\pi_{ij} = -P \left(1 - \frac{2i}{\tau\omega} \right) h_{ij}, \quad (38)$$

while in the collision-dominated regime, π_{ij} is given by Eq. (33).

IV. MAXIMUM COLLISIONAL DAMPING

As one can see from Eq. (32), $\Im(\pi_{xy}/h_{xy})$ has its maximum magnitude for $\omega\tau \sim 1$. The possible damping is thus limited by

$$\text{Max}(|\Im \omega|) \lesssim -\frac{8\pi G}{\omega} \int_p \frac{p^4}{15\epsilon_0^2} \frac{\partial f_0}{\partial \epsilon} \leq \frac{8\pi GP}{\omega}, \quad (39)$$

where P is the total local pressure of the matter under consideration, which gives rise to the damping. It is instructive to write this bound in terms of the expansion rate of the universe, defined by

$$\frac{1}{\tau_H^2} = \frac{8\pi G}{3} \bar{\rho} = \left(\frac{\dot{a}}{a} \right)^2, \quad (40)$$

where a is the cosmological scale parameter and $\bar{\rho}$ the mean mass density of the universe. Since the mean pressure obeys $\bar{P} \leq \bar{\rho}/3$, we find

$$\text{Max}(|\Im \omega|) \lesssim \frac{P}{\bar{P}} \frac{1}{\omega \tau_H^2}. \quad (41)$$

A wave traversing matter will average the local pressure, and thus we can conclude,

$$\text{Max}(|\Im \omega|) \lesssim \frac{1}{\omega \tau_H^2}, \quad (42)$$

indicating that damping of a gravitational wave by matter in the universe can only be significant for a wave of frequency of order $1/\tau_H$. This bound includes all contributions from dark matter particles as well [4].

To express the result (42) in another way, the collisional damping of a gravitational wave within the characteristic expansion time of the universe is of order $1/\omega\tau_H$. For $\omega \sim 10^3 \text{ s}^{-1}$, as in the recent gravitational wave detections [1], and $\tau_H \sim 10^{18} \text{ s}$, this ratio is $\sim 10^{-21}$. Collisional damping in intergalactic or interstellar matter of gravitational waves produced by astrophysical sources is not useful to determine the nature of matter in the universe. This result is valid for any particle-like form of dark matter, including that in a possible shadow universe [16] or matter that only interacts with gravitationally suppressed interactions [17]. Furthermore, collisional damping in locally high dense environments, e.g., in the neighborhood of

mergers of black holes or neutron stars, is also negligible for gravitational waves produced by astrophysical sources, as we argue in the next section. On the other hand for primordial gravitational waves with $\omega \sim 10^{-16} - 10^{-15}$, one has $1/\omega\tau_H \sim 10^{-3} - 10^{-2}$, an effect that could play a role in interpretation of future precision measurements of the spectrum of primordial gravitational radiation.²

As we discuss in the following sections, Landau damping cannot occur in a flat spacetime. Even in an expanding universe, the Landau damping rate is $\sim 1/\omega^2 t_H^3$, so that the total damping within the expansion time of the universe is $\sim 1/(\omega t_H)^2$, a factor $1/\omega t_H$ smaller than the maximum collision damping.

One can write the contribution to the damping from a particular component, s , of the matter, e.g., neutrinos or dark matter, in the form

$$|\Im\omega|_s \equiv n_s \sigma_{\text{GW},s}, \quad (43)$$

where $\sigma_{\text{GW},s}$ is the graviton scattering cross section on particles of species s , and n_s is the number density of species s . The ratio of the cross section to the Planck length, ℓ_{Pl} , squared is essentially bounded above by

$$\frac{\sigma_{\text{GW},s}}{\ell_{\text{Pl}}^2} \lesssim \frac{\langle pv \rangle_s}{\omega} \quad (44)$$

where $\langle pv \rangle_s$ is the mean product of the particle momentum and velocity of species s , which is of order the temperature for a species in thermal equilibrium, or the temperature at which the species froze out. Thus in general,

$$\frac{\sigma_{\text{GW},i}}{\ell_{\text{Pl}}^2} \lesssim \frac{T}{\omega}, \quad (45)$$

with the above understanding of T .

V. MAXIMAL COLLISIONAL DAMPING IN DENSE ENVIRONMENTS

We look now at the damping of gravitational wave produced by binary astrophysical sources as the waves pass through the dense medium surrounding the sources. Collisional damping is limited by $|\Im\omega| < \gamma_{\text{max}}$, where from Eq. (39),

$$\gamma_{\text{max}} \sim \frac{1}{M_{\text{pl}}^2} \frac{P}{\omega} = \frac{w}{M_{\text{pl}}^2} \frac{\rho}{\omega}; \quad (46)$$

we have introduced the Planck mass $M_{\text{pl}}^{-2} = 8\pi G$ and written the relation between the pressure and the energy density by the equation of state parameter $w = P/\rho$.

²We thank Vicky Kalogera and Chris Pankow for this observation.

Assuming the gravitational wave source to a binary system inside a region surrounded by matter with a given density profile with equation of state parameter w , we find collisional damping along a line of sight to be significant if

$$\int_0^R d\gamma_{\text{max}} \sim 1 \quad (47)$$

where R is a physical radius enclosing the ambient matter. A reasonable first estimate is simply to associate the integral with the characteristic size R_c of the dense region:

$$\int_0^R d\gamma_{\text{max}} \sim \frac{w}{M_{\text{pl}}^2} \frac{\rho}{\omega} R_c \sim \frac{w}{M_{\text{pl}}^2} \frac{M}{R_c^2 \omega} \quad (48)$$

where M is the total mass in the region with characteristic size R_c . To go beyond this estimate, one could take the density profile from detailed calculations, e.g. the profile of a typical dark matter halo (determined via phenomenological models that fit N-body simulations such as the Navarro-Frenk-White or Einasto density profiles [18]), and find numerical factors that little affect the conclusion. In “natural” units, one solar mass $M_\odot \sim 10^{66}$ eV, $1 \text{ Hz} \sim 10^{-15}$ eV, $M_{\text{pl}}^{-1} = \ell_{\text{Pl}} \sim 10^{-35}$ m, and $1 \text{ kpc} \sim 10^{19}$ m, one has

$$\int_0^R d\gamma_{\text{max}} \sim w \frac{10^{-27}}{(R_c/\text{kpc})^2} \left(\frac{M}{M_\odot} \right) \frac{1}{(\nu/\text{Hz})}, \quad (49)$$

where $\nu = \omega/2\pi$. We first consider a typical galactic halo surrounding a binary system source of the gravitational wave. Here, typically $M \sim 10^{12} M_\odot$, $R_c \sim 100 \text{ kpc}$, so that

$$\int_0^R d\gamma_{\text{max}} \sim w \frac{10^{-19}}{(\nu/\text{Hz})} \quad (50)$$

which is feeble for all astrophysical sources within the halo.

We next consider the ambient region surrounding a binary system similar to that which gave rise to GW150914—containing a dense distribution of (not necessarily dark) matter. If the ambient matter has a mass comparable to that of the binary system localized within some region $R_c \gg R_s$, the Schwarzschild radius associated with the total mass of the binary system, we find that at the lowest frequencies of the binary system, the factor $M/M_\odot \times (\nu/\text{Hz})^{-1}$ is $\mathcal{O}(1)$; furthermore for the expected nonrelativistic low pressure surrounding matter, $w \sim c_s^2$, the square of the adiabatic sound velocity. Thus

$$\int_0^R d\gamma_{\text{max}} \sim c_s^2 \frac{10^{11}}{(R_c/\text{m})^2}. \quad (51)$$

which can only be of order unity if the ambient matter is localized to within a radius $R_c \lesssim c_s \times 300 \text{ km}$ around the binary system, which even for mildly nonrelativistic ambient matter [e.g. $c_s \sim \mathcal{O}(10^{-1})$] would require a mass

comparable to that of the binary system to be crammed into a region comparable to the Schwarzschild radius of the final merged black hole ($\sim 70 \text{ km} \sim R_s$); such a high density is contrary to our initial assumption that $R_c \gg R_s$. Requiring that this matter be distributed within a region an order of magnitude larger than the binary system yields $\int_0^R dr \gamma_{\text{max}} \sim c_s^2 \ll 1$. We conclude that a distribution of nonrelativistic matter of high density surrounding the source of gravitational radiation is not capable of significantly damping gravitational radiation.

More realistically, one is in general far from the condition of maximal collisional damping, that the collision rate τ^{-1} , be comparable to the frequency of the gravitational wave. Maximal collisional damping is a highly unlikely prospect even for relativistic matter jets and lobes close to the merger of neutron star/black hole binary systems [19,20]. To see this we write roughly, $\tau^{-1} = n\sigma v$, where n is the density of particles, σ is a particle-particle scattering cross section, and v a mean particle velocity. In terms of the mass, M , and characteristic radius, R_c , of the dense environment,

$$\frac{1}{\tau} \sim \frac{M}{M_\odot} \frac{10^{35}}{(R_c/\text{m})^3} \left(\frac{\sigma}{\text{fm}^2} \right) \frac{v}{c} \text{s}^{-1}, \quad (52)$$

Clearly, for a typical nuclear or particle physics cross section, the above is much larger than the typical frequency of an astrophysical binary system by many orders of magnitude, so that $\omega\tau \ll 1$.

VI. LANDAU DAMPING: GENERAL CONSIDERATIONS

In flat space in the collisionless limit ($\tau \rightarrow \infty$) Eq. (32) reduces to

$$\Im \left(\frac{\pi_{ij}}{h_{ij}} \right) = \pi\omega \int_p \left(\frac{p_i p_j}{\epsilon_0} \right)^2 \delta(\omega - \vec{q} \cdot \vec{v}) \frac{\partial f_0}{\partial \epsilon}, \quad (53)$$

a result describing Landau damping, the decay of the mode into a single particle-hole pair.³ The particle-hole excitations are spacelike. For a gravitational wave, $\omega = q$, the integral vanishes except possibly for massless particles moving in the same direction, say \hat{z} , as the gravitational wave. However, for such particles, the factor $p_i^2 p_j^2 \rightarrow p_x^2 p_y^2$ vanishes; Landau damping is forbidden in the absence of cosmological expansion. Following general remarks on Landau damping in this section we show in the following section how the collisionless damping process described by

³In the language of quantum mechanics, the damping may be regarded as the creation, with amplitude $\propto 1/(\omega - \vec{q} \cdot \vec{v}_s)$, of a virtual single particle-hole pair, which subsequently decays into two real particle-hole pairs. Equation (32) can be understood, when $1/\tau \rightarrow 0$, as this amplitude squared, summed over all momenta.

Weinberg [10] can be understood as a generalization of Landau damping, driven by the expansion of the universe.

In an expanding universe, the gravitational wave energy changes during expansion, i.e., the frequency of the gravitational wave is not constant, since the expansion absorbs energy from the wave. This energy loss is different from Landau damping by the matter traversed by the wave. When the phase velocity of the wave is different from the group velocity of the excitations in the matter, energy in flat spacetime is pumped to and fro between the wave and the matter, but the net rate of transfer is zero because the energy transferred in one half-cycle of the wave is exactly cancelled by the loss in the other half-cycle. In an expanding universe, however, the cancellation is incomplete.

We recall the energy loss caused by expansion. A weak gravitational wave of period small compared with the age of the universe behaves in the absence of matter as $h_{ij} = \chi(u)e^{-iqu}$, where u is conformal time, related to coordinate time by $du = dt/a(t)$; as we see in the next section, $\chi(u) \propto 1/a(u)$. This structure is expected on the basis of simple arguments: the energy of a gravitational wave packet is proportional to the energy density in the wave packet times the volume of the packet. The energy density of the wave varies as $g^{ii}(\partial\chi/\partial x^i)^2 \sim a^{-4}$ and the volume of the packet varies as a^3 , so the total energy decreases as $1/a$. This result also agrees with simple redshift arguments: The energy of a massless particle varies as $1+z \sim 1/a$ owing to the expansion of the universe, and thus the energy density measured in locally Minkowskian spacetime ($ds^2 = -dt^2 + (d\vec{r})^2$) varies as $1/(1+z)^4$. For example, the redshift of the first LIGO event GW150914 was $z = 0.09^{+0.03}_{-0.04}$ [1], leading to an energy density reduction by a factor $\approx 1 - 1/(1.09)^4 \approx 30\%$ from cosmological expansion. By comparison, even were the intervening matter collisionless, Landau damping of the wave would be totally negligible, $\sim 1/(\omega t_H)^2$.

VII. GRAVITATIONAL WAVE DAMPING WITH COSMOLOGICAL EXPANSION

We turn now to relate the gravitational radiation damping derived by Weinberg [10] to the calculations above, and to Landau damping in collisionless plasmas, driven by the expansion of the universe. We first generalize the treatment of [10] to allow for massive particles and collisions, working in conformal time, u , related to coordinate time by $dt = a du$, where a is the scale parameter of the expansion. The metric in the presence of expansion and a gravity wave is given by

$$ds^2 = a(u)^2 [-du^2 + (\delta_{ij} + h_{ij}) dx^i dx^j]. \quad (54)$$

Owing to the explicit a^2 , upper and lower components of vectors are related by $x_\mu = a^2 x^\mu$ to zeroth order in h .

In addition, the energy of a particle in the metric (54) is given, in the absence of h_{ij} , by

$$\epsilon_0^2 = p_i^2 + a^2 m^2. \quad (55)$$

We study, following [10], the evolution of the coupled gravitational wave—matter system, after an initial time u_0 at which the matter distribution function is given as f_0 , essentially the f_h in Eq. (20). Since f_0 includes the metric perturbations $h_{ij}(u_0)$ at that time, the additional perturbations of the energy that modify the distribution function by $\delta f(u)$ at later time depend only on the deviation from $h_{ij}(u_0)$, that is,

$$\delta\epsilon = -\frac{p_i p_j}{2\epsilon_0} [h_{ij}(u) - h_{ij}(u_0)]. \quad (56)$$

The Boltzmann equation in conformal time, Fourier transformed in space [cf. Eq. (23)] is

$$\left(\frac{\partial}{\partial u} + \frac{1}{\tau_c} + i\vec{q} \cdot \vec{v}\right) \delta f = \frac{\partial f_0}{\partial \epsilon} \left(\frac{1}{\tau_c} + i\vec{q} \cdot \vec{v}\right) \delta\epsilon, \quad (57)$$

The particle velocity, v , the distribution function f_0 , and the conformal collision time $\tau_c = \tau/a$, are directly dependent on the scale factor. (More generally τ will depend on the cosmological epoch and thus contain further dependence on the scale factor, a question we do not pursue here.) In particular

$$v^i = \partial\epsilon_0/\partial p_i = \frac{p_i}{\sqrt{p_j p_j + m^2 a(u)^2}}. \quad (58)$$

Similarly, an equilibrium distribution function,

$$f_0 = \frac{1}{e^{\epsilon_0/T_0(u)} \mp 1} \quad (59)$$

(with \mp for bosons or fermions) depends on a through the term $m^2 a^2$ in ϵ , and $T_0(u)/a(u)$ is the temperature of the dark matter. For massless particles, T_0 is constant. In addition τ_c is a function of the ambient density along the trajectory of the gravitational wave, and so in general depends on time through its dependence on the scale factor, as well as through the evolving particle distributions. Equation (57) has the general solution

$$\delta f(u, p) = \int_{u_0}^u du' \frac{\partial e^{-\Phi(u, u')}}{\partial u'} \frac{\partial f_0}{\partial \epsilon}(u') \delta\epsilon(u'); \quad (60)$$

where we write

$$\begin{aligned} \Phi(u, u') &\equiv \int_{u'}^u du'' \left(\frac{1}{\tau_c(u'')} + i\vec{q} \cdot \vec{v}(u'') \right) \\ &= \ell(u, u') + i\vec{q} \cdot \hat{p}s(u, u'), \end{aligned} \quad (61)$$

in terms of

$$\ell(u, u') = \int_{u'}^u \frac{du''}{\tau_c(u'')}, \quad (62)$$

and

$$s(u, u') = q \int_{u'}^u du'' v(u''), \quad (63)$$

which is the displacement of the particle in the interval u' to u times the wave vector. The deviation from local equilibrium in (65) is therefore

$$\delta f - \frac{\partial f}{\partial \epsilon} \delta\epsilon = - \int_{u_0}^u du' e^{-\Phi(u, u')} \frac{\partial}{\partial u'} \left[\frac{\partial f_0}{\partial \epsilon} \delta\epsilon(u') \right]. \quad (64)$$

On a cosmological background,

$$\pi_{ij} = \int_p \frac{p_i p_j}{\sqrt{-g}\epsilon_0} \left[\delta f - \frac{\partial f}{\partial \epsilon} \delta\epsilon \right], \quad (65)$$

the generalization of Eq. (19), where in the absence of a gravitational wave, $\sqrt{-g} = a^4$. With Eq. (64), we then have

$$\pi_{ij}(q, u) = - \int_p \frac{p_i p_j}{a(u)^4 \epsilon_0} \int_{u_0}^u du' e^{-\Phi(u, u')} \frac{\partial}{\partial u'} \left[\frac{\partial f_0}{\partial \epsilon} \delta\epsilon(u') \right]. \quad (66)$$

This equation is the direct generalization of Eq. (28) to an expanding spacetime.

Using $\epsilon d\epsilon = p dp$ we have

$$\begin{aligned} \pi_{ij} &= \frac{1}{a(u)^4} \int_p \frac{p_i p_j p_k p_l}{2p\epsilon_0(u)} \int_{u_0}^u du' e^{-\Phi(u, u')} \\ &\quad \times \frac{\partial}{\partial u'} \left[\frac{\partial f_0}{\partial p} (h_{kl}(u') - h_{kl}(u_0)) \right]. \end{aligned} \quad (67)$$

Since $k \neq l$ the angular average above has the form $(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{jk})K(s)$ where in terms of spherical Bessel functions $K(s) = j_2(s)/s^2$, and explicitly,

$$\begin{aligned} K(s) &= \int \frac{d\Omega}{4\pi} e^{-i\zeta s} (1 - \zeta^2)^2 \sin^2 \varphi \cos^2 \varphi \\ &= -\frac{\sin s}{s^3} - 3\frac{\cos s}{s^4} + 3\frac{\sin s}{s^5}, \end{aligned} \quad (68)$$

with $\zeta = \cos \theta$; thus

$$\pi_{ij}(q, u) = \frac{1}{a(u)^4} \int_p \frac{p^3}{\epsilon_0(u)} \int_{u_0}^u du' e^{-\ell(u, u')} K(s) \times \frac{\partial}{\partial u'} \left[\frac{\partial f_0}{\partial p} (h_{ij}(u') - h_{ij}(u_0)) \right]. \quad (69)$$

In the massless limit, we integrate the momentum derivative by parts, using $\epsilon d\epsilon = p dp$, and noting that $s \rightarrow q(u - u')$, to obtain

$$\pi_{ij} = -4\bar{\rho} \int_{u_0}^u du' e^{-\ell(u, u')} K(q(u - u')) h'_{ij}(u'), \quad (70)$$

where the prime denotes d/du , and

$$\bar{\rho} = \frac{1}{a^4} \int_p p f_0 \quad (71)$$

is the mass density of the matter. Away from the massless limit, generalizing Ref. [10], we find extra contributions from the p dependence of s in K , an effect that was investigated in the context of collisionless massive neutrinos and axions in [15], where it was found that the damping is reduced when including masses. We see from Eq. (69) or (70), that, as expected, the net effect of collisional interactions is to efficiently erase anisotropic stresses, and hence limit their ability to damp gravitational waves.

Since astrophysical sources of gravitational waves have characteristic frequencies much greater than the inverse Hubble scale at late times, we can expand the time dependence of the mode functions in powers of $a'/(aq)$. For such a wave, the spatial Fourier component \vec{q} obeys the equation of motion,

$$h''_{ij} + 2\frac{a'}{a} h'_{ij} + q^2 h_{ij} = 16\pi G a^2 \pi_{ij}. \quad (72)$$

The solution for $h_{ij}(q, u)$ in the absence of matter is, to lowest order in $a'/(aq)$,

$$h_{ij}(q, u) \propto \frac{e^{-iqu}}{a(u)} \quad (73)$$

(during radiation domination, this result is exact). We assume that the gravitational wave is in the form of a wave packet for which

$$h_{ij}(\vec{r}, u) = \int \frac{d^3 q}{(2\pi)^3} \frac{e^{-iqu}}{a(u)} F(q), \quad (74)$$

where $F(q)$ is localized about a wave vector \vec{q}_0 . We consider the absorption by a region of matter much smaller than the horizon size.

To see the mechanism of Landau damping in the absence of collisions, we calculate the damping of the wave directly

in terms of the energy transfer to the matter, writing, from Eq. (15), using conformal time

$$\frac{\partial E}{\partial u} = -\frac{1}{2} \int d^3 r h'_{ij}(\vec{r}, u) \pi_{ij}(\vec{r}, u) = -\frac{1}{2} \Re \int \frac{d^3 q}{(2\pi)^3} h'_{ij}{}^*(q, u) \pi_{ij}(q, u). \quad (75)$$

We work in the massless limit, in order to illustrate the physics with the fewest complications. Then f_0 does not depend on a , and $\Phi(u, u') \rightarrow iq\zeta(u - u')$, and one has,

$$\begin{aligned} \frac{\partial E}{\partial u} &= -\frac{1}{4a(u)^4} \Re \int \frac{d^3 q}{(2\pi)^3} h'_{ij}{}^*(q, u) \int_p \frac{p_i p_j p_k p_l}{p^2} \\ &\quad \times \int_{u_0}^u du' e^{-iq\zeta(u-u')} \frac{\partial f_0}{\partial p} h'_{kl}(q, u') \\ &= -\frac{1}{16a(u)^4} \Re \int \frac{d^3 q}{(2\pi)^3} h'_{ij}{}^*(q, u) \int_p p^2 \frac{\partial f_0}{\partial p} \\ &\quad \times (1 - \zeta^2)^2 \int_{u_0}^u du' e^{-iq\zeta(u-u')} h'_{ij}(q, u'). \end{aligned} \quad (76)$$

From Eq. (73), we see that $h'_{ij}(u) = -(iq + \mathcal{H}(u))h_{ij}(u)$, where $\mathcal{H} \equiv a'(u)/a(u)$. The explicit $\mathcal{H}(u)$, which is small relative to the q term and for an astrophysical gravitational wave produces only a small correction to the Landau damping, can be neglected. Over the time span of a gravitational wave packet transversing a given region of matter, the scale factor $a(u')$ in $h_{ij}(u')$ can be expanded as $a(u') = a(u) + a'(u)(u' - u) \simeq a(u)e^{\mathcal{H}(u'-u)}$. Thus the u' integral can be written as

$$\begin{aligned} &\frac{-iq e^{-iqu} F(q)}{a(u)} \int_{u_0}^u du' e^{(iq(1-\zeta) + \mathcal{H}(u))(u-u')} \\ &\simeq \frac{1}{\mathcal{H} + iq(1-\zeta)} (h'_{ij}(q, u) - e^{-iq\zeta(u-u_0)} h'_{ij}(q, u_0)). \end{aligned} \quad (77)$$

Since the characteristic frequencies are large compared with $1/(u - u_0)$ the phase factor in the final term will average to zero inside the q and ζ integrals in Eq. (76). We find then

$$\frac{\partial E}{\partial u} = \frac{\bar{\rho}}{4} \int \frac{d^3 q}{(2\pi)^3} |h'_{ij}(q, u)|^2 \int_{-1}^1 \frac{d\zeta}{2} \frac{\mathcal{H}(1 - \zeta^2)^2}{\mathcal{H}^2 + q^2(1 - \zeta)^2}. \quad (78)$$

We see here how expansion of the universe introduces a spread in frequencies $\sim \pm \mathcal{H}$ about q , thus allowing Landau damping; in the absence of expansion, $\mathcal{H} = 0$, and Landau damping vanishes.

To lowest order in \mathcal{H}/q the integral is simply $4/3$, so that

$$\frac{\partial E}{\partial u} = \frac{\bar{\rho}}{3} \frac{a'}{a} \int \frac{d^3 q}{(2\pi)^3} \frac{|h'_{ij}(q, u)|^2}{q^2}. \quad (79)$$

The energy density of the gravitational wave is

$$E_{gw} = \int \frac{d^3 q}{(2\pi)^3} \frac{|h'_{ij}(q, t)|^2}{32\pi G}, \quad (80)$$

so that for a wave packet centered about a frequency \bar{q}

$$\frac{\partial E}{\partial u} = \frac{32\pi G \bar{\rho}}{3\bar{q}^2} \mathcal{H} E_{gw}. \quad (81)$$

Finally we note that $8\pi G \bar{\rho}/3 \sim (a'/a^2)^2$ and thus

$$\frac{\partial E}{\partial u} \sim \frac{4\mathcal{H}^3}{(a\bar{q})^2} E_{gw}. \quad (82)$$

The characteristic absorption time via Landau damping is thus $\sim \omega^2 t_H^3$ (with $\omega = \bar{q}$), which is thoroughly negligible. The corresponding fractional change in energy over an expansion time of the universe is $\sim 1/(\omega t_H)^2$.

VIII. CONCLUDING REMARKS

In this paper we have laid out a framework for evaluating the damping of gravitational radiation by matter with arbitrary mass particles and collision strengths. By considering the damping of gravitational waves in both flat spacetime and in an expanding universe, we identify two distinct mechanisms through which damping can occur—the first in which collisions produce the damping, and the second, via Landau damping.

When the expansion of spacetime can be neglected, the damping of a wave of a given frequency, proportional to the *relaxation rate* $1/\tau$ in the collisionless regime ($\omega\tau \gg 1$) and to the *collision time*, τ , in the hydrodynamic regime ($\omega\tau \ll 1$), is maximal when $\omega\tau \simeq 1$. For the frequencies to which LIGO is sensitive and for plausible models of dark

matter, calculations of damping based on hydrodynamical considerations are gross overestimates, and we conclude that it is impossible from current observations of gravitational waves to put useful bounds on the properties of dark matter. Landau damping in this case is not possible because particles have velocities less than c . As we estimate in Sec. V, collisional damping of gravitational waves of frequencies produced by astrophysical binary systems, propagating through dense local environments, is also insignificant.

Collisionless damping is possible in an expanding universe since the frequency of the gravitational wave and the energies of particles depend on time. Damping of gravitational waves by free-streaming relativistic particles as discussed by Weinberg [10], may, as we have shown, be regarded as a generalization of Landau damping; we have also generalized Weinberg's formalism to allow for collisions in the matter. We note in passing that one can straightforwardly extend the present framework to incorporate scenarios of nonthermal dark matter, since it was not essential to assume a specific functional form for the distribution, see, e.g., Eq. (20).

In the future we will apply the present framework to study damping of stochastic gravitational waves produced after inflation, e.g., in (first order) phase transitions in the early universe, during matter domination where the matter consists of ultralight dark matter. Such scenarios are similar to damping by neutrinos during radiation domination [10,13,15], and might describe damping by axions [21–23].

ACKNOWLEDGMENTS

We are grateful to Diego Blas, Stu Shapiro and Subir Sarkar for very helpful remarks. The research of author G. B. was supported in part by NSF Grants No. PHY1305891 and No. PHY1714042. He is grateful to the Aspen Center for Physics, supported in part by NSF Grants No. PHY1066292 and No. PHY1607611, and the Niels Bohr International Academy where parts of this research were carried out. Author S.P. is supported by funds from Danmarks Grundforskningsfond under Grant No. 1041811001.

-
- [1] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), *Phys. Rev. Lett.* **116**, 061102 (2016); **116**, 241103 (2016); **118**, 221101 (2017); **119**, 141101 (2017).
 - [2] S. W. Hawking, *Astrophys. J.* **145**, 544 (1966).
 - [3] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (Wiley, New York, 1972), Ch. X.
 - [4] G. Goswami, G. K. Chakravarty, S. Mohanty, and A. R. Prasanna, *Phys. Rev. D* **95**, 103509 (2017).

- [5] A. P. Lightman, W. H. Press, R. H. Price, and S. A. Teukolsky, *Problem Book in Relativity and Gravitation* (Princeton University Press, Princeton, NJ, 1975), Problem 18.15.
- [6] S. Gayer and C. F. Kennel, *Phys. Rev. D* **19**, 1070 (1979).
- [7] A. A. Abrikosov and I. M. Khalatnikov, *Rep. Prog. Phys.* **22**, 329 (1959), Eq. (10.1).
- [8] D. Lynden-Bell, *Mon. Not. R. Astron. Soc.* **124**, 279 (1962).
- [9] G. Baym, H. Monien, C. J. Pethick, and D. G. Ravenhall, *Phys. Rev. Lett.* **64**, 1867 (1990).

- [10] S. Weinberg, [Phys. Rev. D](#) **69**, 023503 (2004).
- [11] C. W. Misner, [Phys. Rev. Lett.](#) **19**, 533 (1967).
- [12] J. R. Pritchard and M. Kamionkowski, [Ann. Phys. \(Amsterdam\)](#) **318**, 2 (2005).
- [13] T. Watanabe and E. Komatsu, [Phys. Rev. D](#) **73**, 123515 (2006).
- [14] B. A. Stefanek and W. W. Repko, [Phys. Rev. D](#) **88**, 083536 (2013).
- [15] J. B. Dent, L. M. Krauss, S. Sabharwal, and T. Vachaspati, [Phys. Rev. D](#) **88**, 084008 (2013).
- [16] K. Nishijima and M. H. Saffouri, [Phys. Rev. Lett.](#) **14**, 205 (1965).
- [17] M. Garny, M. Sandora, and M. S. Sloth, [Phys. Rev. Lett.](#) **116**, 101302 (2016).
- [18] A. W. Graham, D. Merritt, B. Moore, J. Diemand, and B. Terzic, [Astron. J.](#) **132**, 2685 (2006).
- [19] V. Paschalidis, M. Ruiz, and S. L. Shapiro, [Astrophys. J. Lett.](#) **806**, L14 (2015).
- [20] M. Ruiz, R. Lang, V. Paschalidis, and S. L. Shapiro, [Astrophys. J. Lett.](#) **824**, L6 (2016).
- [21] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, [Phys. Rev. D](#) **95**, 043541 (2017).
- [22] N. Banik and P. Sikivie, in *Universal Themes of Bose-Einstein Condensation*, edited by D. Snoke, N. Proukakis, and P. Littlewood (Cambridge University Press, Cambridge, England, 2017).
- [23] D. J. E. Marsh, [Phys. Rep.](#) **643**, 1 (2016).