Ś

Critical flavor number of the Thirring model in three dimensions

Björn H. Wellegehausen

Institut für Theoretische Physik, Justus-Liebig-Universität Giessen, 35392 Giessen, Germany and Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, 07743 Jena, Germany

Daniel Schmidt[†] and Andreas Wipf[‡]

Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, 07743 Jena, Germany (Received 10 August 2017; published 9 November 2017)

The Thirring model is a four-fermion theory with a current-current interaction and U(2N) chiral symmetry. It is closely related to three-dimensional OED and other models used to describe properties of graphene. In addition, it serves as a toy model to study chiral symmetry breaking. In the limit of flavor number $N \rightarrow 1/2$ it is equivalent to the Gross-Neveu model, which shows a parity-breaking discrete phase transition. The model was already studied with different methods, including Dyson-Schwinger equations, functional renormalization group methods, and lattice simulations. Most studies agree that there is a phase transition from a symmetric phase to a spontaneously broken phase for a small number of fermion flavors, but no symmetry breaking for large N. But there is no consensus on the critical flavor number $N^{\rm cr}$ above which there is no phase transition anymore and on further details of the critical behavior. Values of N found in the literature vary between 2 and 7. All earlier lattice studies were performed with staggered fermions. Thus it is questionable if in the continuum limit the lattice model recovers the internal symmetries of the continuum model. We present new results from lattice Monte Carlo simulations of the Thirring model with SLAC fermions which exactly implement *all* internal symmetries of the continuum model even at finite lattice spacing. If we reformulate the model in an irreducible representation of the Clifford algebra, we find, in contradiction to earlier results, that the behavior for even and odd flavor numbers is very different: for even flavor numbers, chiral and parity symmetry are always unbroken; for odd flavor numbers, parity symmetry is spontaneously broken below the critical flavor number $N_{ir}^{cr} = 9$, while chiral symmetry is still unbroken.

DOI: 10.1103/PhysRevD.96.094504

I. INTRODUCTION

The Thirring model [1] is a fermionic quantum field theory with a current-current interaction. While it was originally studied in two spacetime dimensions, a lot of recent works were concerned with its three-dimensional version with a varying number of N flavors. This model is renormalizable in a 1/N-expansion for 2 < d < 4 [2–5]. Its Lagrangian in Euclidean spacetime is given by

$$\mathcal{L} = \bar{\Psi}_a \mathrm{i} \Gamma^\mu \partial_\mu \Psi_a - \frac{g^2}{2N} (\bar{\Psi}_a \Gamma_\mu \Psi_a)^2, \quad a = 1, \dots, N, \quad (1)$$

with summation over fermion flavors. In three dimensions, an irreducible representation of the Clifford algebra is two-dimensional, but we will start with a reducible representation here and take the well-known Γ_{μ} matrices of the four-dimensional theory with four-component spinor fields Ψ_a . This is motivated by a strong similarity to

three-dimensional QED [5–8], which is often used to model electronic properties of materials like graphene [9,10] or high-temperature superconductors [11,12].

The Thirring model is also interesting on its own, because it has a large continuous chiral symmetry. It is believed that spontaneous breaking of this symmetry can happen with the pattern

$$U(2N) \to U(N) \otimes U(N),$$
 (2)

but only for a small number of flavors. A critical flavor number N^{cr} should exist, such that chiral symmetry breaking (csb) only occurs for $N < N^{cr}$. The main focus of our work is to find the value of N^{cr} for the reducible Thirring model.

While there is a broad agreement on this general behavior in the literature, the predictions for N^{cr} vary to some extent. Numerous works have employed a large-N expansion: Different studies have used Dyson-Schwinger equations (DSEs), with the first [13] reporting $N^{cr} \approx 3.24$. Later work [6,14] found $N^{cr} \approx 4.32$ in the limit $g^2 \rightarrow \infty$. By constructing an effective potential by an inversion method, Kondo [15] reported $N^{cr} = 2$ for infinite Thirring coupling.

bjoern.wellegehausen@uni-jena.de

d.schmidt@uni-jena.de

^{*}wipf@tpi.uni-jena.de

Additionally, these works reported relations between the chiral condensate, N, and g^2 that are qualitatively not in agreement with each other.

A recent extensive study [16,17] of four-fermion theories with functional renormalization group methods spotted a structure with three interacting fixed points in the plane spanned by Thirring and Gross-Neveu coupling. The fixed point governing the critical behavior of the Thirring model is only on the axis of pure Thirring interaction for $N \to \infty$, while it is off the axis for any finite N. On the contrary, for small N this fixed point is dominated by another fourfermion interaction, showing dynamical generation of a fermion mass. This is not the case for large N. Balancing this competition, the authors found $N^{cr} \approx 5.1(7)$.

Regarding lattice field theory, many simulations with staggered fermions and a small mass have been performed so far. With a setup using the standard hybrid Monte Carlo (HMC) algorithm, simulations are only possible with an integer number of flavors. Since for staggered fermions each lattice flavor corresponds to two continuum flavors [18], only simulations with even N are possible. The first results [19] reported a change of the chiral behavior for $2 < N^{cr} < 6$. Another series of publications [20–22] used the same algorithm with a slightly different action and found chiral symmetry breaking with a second-order phase transition for N = 2 and N = 4, while the transition for N = 6 is different. The authors claimed that in the latter case there is a first-order transition with coexisting symmetric and broken phases. They concluded that N^{cr} is between 4 and 6. Simulations with a hybrid molecular dynamics algorithm were performed in Ref. [23], allowing also for odd and noninteger values of N. The authors presented a phase diagram in the (N, g^2) plane with a critical line. Along this line, the critical exponents and the order of the phase transition change from a second-order transition at N = 4 to a first-order transition at N = 6. For N = 5 the simulations did not lead to a conclusive result.

In a more recent lattice study [24] with this setup, simulations in the limit $g^2 \rightarrow \infty$ were performed, in order to compare the results with those from DSE approaches. To ensure transversality of the vacuum polarization tensor for large *N* on the lattice, a renormalization of the coupling was necessary. The renormalized coupling at leading order in 1/N is

$$g_{\rm R}^2 = \frac{g^2}{1 - g^2 J(m)}$$
 with $J(m) \to \frac{2}{3}$, (3)

where the integral J(m) was given in Ref. [21]. Obviously, g_R^2 becomes negative if the bare coupling g^2 is larger than 3/2. Thus the strong-coupling limit is reached for finite bare coupling. For stronger bare couplings, an unphysical phase is present. All these works found a nonmonotonic behavior of the chiral condensate, and it is argued that its maximum corresponds to the point where

the renormalized coupling becomes negative, although the coupling at the maximum does not match the value of 3/2. Looking at the maximal value of the chiral condensate for different N at small masses, the authors concluded that $N^{\rm cr} = 6.6(1)$.

The fermion bag approach was applied by Chandrasekharan [25] to study the Thirring model with a single staggered flavor, corresponding to N = 2, and to obtain critical exponents for the csb phase transition [26,27]. This was the first lattice work in the chiral limit m = 0. But since staggered fermions do not preserve all internal symmetries it is not clear if the correct symmetrybreaking pattern (2) is recovered in the continuum limit. In a subsequent work the authors [28] observed that their lattice version of the Thirring model has the same symmetry and critical exponents as the Gross-Neveu model. This seems to contradict our knowledge about the continuum models. More recently, the first study with domain-wall fermions was performed [29]. Contrary to the older works, no remnant of csb was found in the extrapolation $m \to 0$ for N = 2. Very recent preliminary results [30] for N = 1 showed no csb either.

In the present work we follow an alternative route and simulate the Thirring model with chiral SLAC fermions [31,32]. It is well known that these fermions should not be used in lattice gauge theories [33], but they have been successfully applied to simulate supersymmetric Yukawa models [34-38], where the renormalizability of lattice perturbation theory with SLAC fermions up to one-loop was also established. With these fermions, the step scaling function of the nonlinear O(3) sigma model has been calculated to high accuracy on moderately large lattices; see Ref. [39]. Thus there are good reasons to believe that they work well for four-fermion theories. SLAC fermions admit the exact internal symmetry at any finite lattice spacing. Nevertheless, it was difficult to establish csb at any N in previous investigations [40,41]. Including an explicit breaking, a physical phase transition was observed, but it seems to merge with the artifact transition explained around Eq. (3), leaving no reliable trace of csb when performing the limit to the massless Thirring model. An attempt to use Fierz identities to reformulate the Thirring model showed a very strong sign problem preventing HMC simulations [41]. The present work will present an approach to circumvent these problems and provide a definite answer about the existence and values of the critical flavor number. We shall see that there is no symmetry breaking in all models with integer N. This is in complete agreement with the absence of bilinear condensation in three-dimensional QED with an even number of massless irreducible flavors [42]. In contrast, we present numerical evidence that all models with half-integer $N \leq N^{\rm cr} = 9/2$ show a symmetry breaking.

To ease the computations, we also consider the Thirring model in the irreducible representation, where the chiral symmetry is merely a flavor symmetry. Then, dynamical generation of a fermion mass is associated to a spontaneous breaking of parity. This was studied by DSE in the large-*N* limit, where a dynamically generated mass was found [43], implying $N^{cr} \rightarrow \infty$. In Refs. [44,45] a different parity-breaking pattern emerged when Fierz identities were used to compute the effective potential: a dynamical mass generation for two and three irreducible flavors was seen, whereas the potential became unbounded from below for $N \rightarrow \infty$. Similarly, in the functional Schrödinger picture no symmetry breaking in the large-*N* limit was found, while it appeared when higher-order corrections in 1/N were included [46].

With the help of an auxiliary vector field v_{μ} , the Lagrangian (1) can be written in the equivalent form

$$\mathcal{L} = \bar{\Psi}_a \mathbf{i} \Gamma^\mu D_\mu \Psi_a + \lambda v_\mu v^\mu, \qquad \lambda = \frac{N}{2g^2}, \qquad (4)$$

with the "covariant derivative" $D_{\mu} = \partial_{\mu} - iv_{\mu}$. In an adapted base the reducible matrices Γ_{μ} and reducible spinors take the form

$$\Gamma_{\mu} = \sigma_3 \otimes \gamma_{\mu}, \qquad \Psi_a = \begin{pmatrix} \Psi_{1,a} \\ \Psi_{2,a} \end{pmatrix}, \qquad \bar{\Psi}_a = (\bar{\psi}_{1,a}, \bar{\psi}_{2,a}), \tag{5}$$

where the two-dimensional γ_{μ} form an irreducible representation of the Clifford algebra in three dimensions. For massless fermions the overall sign of $\not D = \gamma^{\mu}(\partial_{\mu} - iv_{\mu})$ in

is irrelevant and the model with *N* reducible flavors is equivalent to the model with $N_{ir} = 2N$ irreducible flavors, for which the second term in Eq. (6) has a positive sign. For massive fermions this is no longer true: when integrating out irreducible fermions a parity-violating (imaginary) Chern-Simons-like term is generated.¹ In contrast, the fermionic determinant is real for reducible massive fermions and we do not expect parity breaking in models with reducible fermions.

This paper is organised as follows. In Sec. II, we summarize the symmetries of the Thirring model in the reducible and an irreducible representation. The relation between both formulations is further discussed. Then, in Sec. III we use a Fierz identity to rewrite the Lagrangian for irreducible fermions. The main derivation of the effective potential for local condensates can be found in Sec. IV, while its symmetries are studied in Sec. V. We also give the

explicit forms of the potential for $N_{\rm ir} = 1$ and 2. Next, we study the effective potential in the strong-coupling limit in Sec. VI, before we present our main results from numerical simulations in Sec. VII. Finally, a discussion of our findings and a comparison with previous results can be found in Sec. VIII.

II. CONTINUUM SYMMETRIES

First we discuss the internal symmetries of the Thirring model in three dimensions. The results look different for N fermions in the four-dimensional reducible representation of the Lorentz group and N_{ir} fermions in a two-dimensional irreducible representation. The flavor numbers are related as $N_{ir} = 2N$.

In the reducible representation Ψ_a is a four-component Dirac spinor, where a = 1, ..., N labels the flavors. The theory possesses a U(2N) chiral symmetry generated by

$$T = T_{\rm f} \otimes \{\mathbb{1}, \Gamma_4, \Gamma_5, \mathrm{i}\Gamma_4\Gamma_5\},\tag{7}$$

where $T_f \in \mathfrak{u}(N)$ generates rotations in flavor space. A Dirac spinor transforms under the chiral and flavor rotations as

$$\begin{split} \Psi &\to e^{\mathrm{i}T_{\mathrm{f}}^{a}\otimes(t_{1}^{a}\mathbb{1}+t_{2}^{a}\Gamma_{4}+t_{3}^{a}\Gamma_{5}+\mathrm{i}t_{4}^{a}\Gamma_{4}\Gamma_{5})}\Psi, \\ \bar{\Psi} &\to \bar{\Psi}e^{-\mathrm{i}T_{\mathrm{f}}^{a}\otimes(t_{1}^{a}\mathbb{1}-t_{2}^{a}\Gamma_{4}-t_{3}^{a}\Gamma_{5}+\mathrm{i}t_{4}^{a}\Gamma_{4}\Gamma_{5})}. \end{split}$$
(8)

Furthermore, the theory is invariant under a discrete \mathbb{Z}_2 parity transformation

$$\Psi(x) \to i\Gamma_1\Gamma_4\Psi(x') \quad \text{with} \quad x' = (x_0, -x_1, x_2), \quad (9)$$

where alternative formulations using Γ_5 instead of Γ_4 are possible. A detailed discussion of both continuous and discrete symmetries can be found in Refs. [13,16]. In total, the global symmetry group is $U(2N) \otimes \mathbb{Z}_2$. Now, we can define a parity-even chiral condensate $\Sigma = \bar{\Psi}\Psi$ and a parity-odd condensate $\Sigma_{45} = \bar{\Psi}i\Gamma_4\Gamma_5\Psi$. The chiral condensate Σ is an order parameter for spontaneous breaking of the continuous chiral symmetry according to

$$U(2N) \otimes \mathbb{Z}_2 \xrightarrow{\Sigma} U(N) \otimes U(N) \otimes \mathbb{Z}_2,$$
 (10)

while the parity condensate Σ_{45} serves as an order parameter for discrete parity breaking,

$$U(2N) \otimes \mathbb{Z}_2 \stackrel{\Sigma_{45}}{\to} U(2N). \tag{11}$$

For calculations it is often more convenient to reformulate the Thirring model in an irreducible representation. A useful reduction is given by

¹More accurately, an imaginary Chern-Simons term is generated in the limit $m \to \infty$. For small *m* the imaginary part of the effective action is proportional to the η invariant.

$$\Gamma_{\mu} = \sigma_3 \otimes \gamma_{\mu}, \qquad \Gamma_4 = \sigma_2 \otimes \mathbb{I},$$

$$\Gamma_5 = \sigma_1 \otimes \mathbb{I} \quad \text{and} \quad i\Gamma_4\Gamma_5 = \sigma_3 \otimes \mathbb{I}. \qquad (12)$$

In order to obtain a *standard* kinetic term in the irreducible representation, we decompose the N four-component Ψ_a as

$$\Psi_a = \begin{pmatrix} \Psi_{1,a} \\ \Psi_{2,a} \end{pmatrix} \quad \text{and} \quad \bar{\Psi}_a = (\bar{\psi}_{1,a}, -\bar{\psi}_{2,a}), \quad (13)$$

where $\psi_{i,a}$ are two-component spinors in a fixed irreducible representation. With this decomposition, the Lagrangian is given by

$$\mathcal{L} = \bar{\psi}_{\alpha} i \partial \!\!\!/ \psi_{\alpha} - \frac{g^2}{2N_f} (\bar{\psi}_{\alpha} \gamma_{\mu} \psi_{\alpha})^2, \qquad (14)$$

where the irreducible flavor index $\alpha = (i, a)$ assumes $2N = N_{ir}$ different values. Note the difference to the decomposition (5). The condensates are

$$\Sigma = \sum_{a=1}^{N} \left(\bar{\psi}_{1,a} \psi_{1,a} - \bar{\psi}_{2,a} \psi_{2,a} \right) \quad \text{and} \tag{15}$$

$$\pi = \Sigma_{45} = \sum_{\alpha=1}^{N_{\rm ir}} \bar{\psi}_{\alpha} \psi_{\alpha}.$$
 (16)

The parity condensate in the reducible representation π turns into a *chiral condensate* in the irreducible representation, while for even N_{ir} the former chiral condensate now has a flavor-staggered structure. This already indicates that in the irreducible representation the behavior is different for even and odd flavor numbers N_{ir} . We will further investigate this difference in the next sections.

The continuous chiral symmetry of the reducible Thirring model becomes a pure flavor symmetry in the irreducible representation, i.e., the theory is invariant under $U(N_{\rm ir})$ flavor transformations given by

$$\psi \to U\psi$$
 and $\bar{\psi} \to \bar{\psi}U^{\dagger}$ (17)

with a unitary matrix $U = \exp(it_iT_i)$ acting in flavor space only. Here, the T_i are generators of the algebra $\mathfrak{u}(N_{ir})$. Due to its relation to the chiral transformations of the fourcomponent spinors, this flavor symmetry will still be called *chiral symmetry* in the following, although the concept of chirality does not exist in odd dimensions.

The parity transformation of the four-component spinors turns into a combination of flavor rotations and a parity transformation in the irreducible representation, which is given by

$$\psi(x) \to \gamma_1 \psi(x')$$
 and $\bar{\psi}(x) \to -\bar{\psi}(x')\gamma_1$. (18)

The *irreducible chiral condensate* π is invariant under chiral transformations (i.e., two-component flavor rotations) while it breaks the discrete \mathbb{Z}_2 parity symmetry.

III. FIERZ IDENTITIES

To integrate over the fermionic fields in the functional integral, it is useful to "linearize" the four-fermion term by a Hubbard-Stratonovich transformation, which transforms the Lagrangian (14) into the equivalent form

$$\mathcal{L} = \bar{\psi}(\mathbf{i}\partial \!\!\!/ + \gamma_{\mu}v_{\mu})\psi + \lambda v_{\mu}^2, \tag{19}$$

with $\lambda = N/2g^2$ and three real scalar fields v_{μ} . After integration over the ψ_{α} we obtain the effective action

$$S_{\rm eff} = -N_{\rm ir} \ln \det \left(i \partial \!\!\!/ + \gamma_{\mu} v_{\mu} \right) + \lambda \int d^3 x v_{\mu}(x)^2. \quad (20)$$

Note that this action is not gauge invariant and v_{μ} should not be viewed as gauge potential.²

A technical problem arises if one discretizes this formulation on a (hypercubic) lattice to perform Monte Carlo simulations. Since v_{μ} is invariant under chiral transformations there is no natural order parameter for chiral symmetry breaking in the massless theory. But the advantage of the vector formulation is that except for $N_{ir} = 1$ it is free of a fermion sign problem. This is obvious for even N_{ir} since $i\partial + \gamma_{\mu}v_{\mu}$ is Hermitian. For odd N_{ir} we never observed a negative sign in our Monte Carlo simulations except for $N_{ir} = 1$.

We can circumvent the technical problem by applying a Fierz transformation to the four-fermion interaction

$$(\bar{\psi}_{\alpha}\gamma_{\mu}\psi_{\alpha})^{2} = -2(\bar{\psi}_{\alpha}\psi_{\beta})(\bar{\psi}_{\beta}\psi_{\alpha}) - (\bar{\psi}_{\alpha}\psi_{\alpha})^{2}.$$
 (21)

This identity was also applied in, for example, Refs. [44,45]. The transformed Lagrangian reads

$$\mathcal{L} = \bar{\psi} i \partial \!\!\!/ \psi + \frac{g^2}{N_f} (\bar{\psi}_{\alpha} \psi_{\beta}) (\bar{\psi}_{\beta} \psi_{\alpha}) + \frac{g^2}{2N_f} (\bar{\psi} \psi)^2.$$
(22)

Now, by applying the Hubbard-Stratonovich transformation we can reproduce the four-fermion terms with the help of a matrix-valued field,

$$\mathcal{L} = \bar{\psi} \left(i \partial \!\!\!/ + iT + \frac{i}{2} tr T \right) \psi + \frac{\lambda}{2} tr T^2 + \frac{\lambda}{4} (tr T)^2, \quad (23)$$

where $T = T^{\dagger}$ is a generic $\mathfrak{u}(N_{ir})$ -algebra valued field, i.e.,

$$T = t_i T_i \quad \text{with} \quad i = 1 \dots N_{\text{ir}}^2. \tag{24}$$

²It could be promoted to a gauge potential after introducing a Stückelberg field [6].

Under a chiral transformation the components of T transform according to

$$T \to UTU^{\dagger}, \qquad U \in U(N_{\rm ir}), \tag{25}$$

such that their expectation values serve as order parameters for chiral symmetry breaking. They are related to fermionic condensates by DSEs. Unfortunately, this formulation of the model suffers from a severe sign problem on the lattice [41].

IV. EFFECTIVE POTENTIAL

The vector formulation (19) and matrix formulation (23) each have their own advantages and disadvantages. The former can (except for $N_{ir} = 1$) be simulated without a sign problem, but information about chiral symmetry is not directly accessible. On the other hand, in the matrix formulation we have direct access to order parameters for csb, but there is a strong sign problem that prevents reliable simulations. We proceed with an analytical treatment in the matrix formulation and calculate the resulting expectation values in the vector formulation. We begin with splitting the matrix field in Eq. (23) as

$$T(x) = T^{c}(x) + T^{\perp}(x),$$
 (26)

where the first term is in the Cartan subalgebra of $\mathfrak{u}(N_{ir})$ and the second is in its orthogonal complement. We shall introduce a dual-variables formulation in Sec. IVA and afterwards present a calculation of the (constraint) effective potential

$$V_{\text{eff}}(T^{\text{c}}) = -\ln \int \mathcal{D}T \mathcal{D}\phi e^{-S_{\text{eff}}(T,\phi)} \delta(T^{\text{c}} - T^{\text{c}}(x_0))$$
$$= -\ln \sum_{n=0}^{2N_{\text{ir}}} \sum_{i=1}^{N_{\text{ir}}} a_{n,i}(t_i)^n, \qquad T^{\text{c}} = t_i H^i,$$

in which the constraint fixes the field at an arbitrary point x_0 to the prescribed value T^c in Sec. IV B. The matrix field $T(x_0)$ is Hermitian and can be diagonalized by a global chiral transformation. Hence it is sufficient to calculate the effective potential for a field in the Cartan subalgebra with generators H^i . In Sec. IV C we will relate the coefficients $a_{n,i}$ to expectation values of observables $O_{n,i}$ in the vector field formulation,

$$a_{n,i} = \langle O_{n,i} \rangle_v . \tag{27}$$

This allows us to employ Monte Carlo simulations in the vector field formulation to calculate the effective potential defined in the matrix formulation.

A. The partition function in the dual variables formulation

First we reformulate the partition function of the Thirring model with massive fermions in terms of discrete *spin* (or dual) variables. Then we take derivatives with respect to the mass to relate observables in the dual formulation to powers of fermionic bilinears. Actually, we introduce an *x*-dependent and diagonal fermion mass matrix

$$M(x) = \text{diag}(m_1(x), \dots, m_{N_{\text{ir}}}(x)).$$
(28)

The partition function for the massive model is now given by (an integral over spacetime is assumed in the exponent)

$$Z(\lambda, M) = \left(\frac{\pi}{\lambda}\right)^{\frac{V}{2}} \int \mathcal{D}T \, Z_F[T, M] e^{-\frac{\lambda}{2} \mathrm{tr} T^2 - \frac{\lambda}{4} (\mathrm{tr} T)^2}, \quad (29)$$

where the fermionic partition function $Z_F[T, M]$ is given by the fermionic integral

$$Z_F[T, M] = \int \mathcal{D}\bar{\psi}\mathcal{D}\psi e^{-\bar{\psi}\mathrm{i}\not\partial\psi} e^{-\bar{\psi}\mathrm{i}H\psi}.$$
 (30)

Here, the shifted field $H = T + \frac{1}{2}trT + M$ was introduced. With spinor index *i* and flavor indices α and β the product expansion of the second exponent leads to

$$Z_F[T,M] = \int \mathcal{D}\bar{\psi}\mathcal{D}\psi e^{-\bar{\psi}i\not\partial\!\!\!/\psi} \prod_{xi\alpha\beta} (1 - \bar{\psi}^{\alpha}_{xi}\mathrm{i}H^{\alpha\beta}_x\psi^{\beta}_{xi}), \quad (31)$$

where we used that ψ_{xi}^{α} and $\bar{\psi}_{xi}^{\alpha}$ are Grassmann variables. A similar expansion is met in the fermion bag approach, where one expands the integrand directly in powers of the four-fermion term [27]. After integrating over the auxiliary field the two expansions yield the same results, although the intermediate expressions are different. A related expansion is also encountered in attempts to dualize gauge theories [47].

At this point we introduce a spin field $k_{xi}^{\alpha\beta} \in \{0,1\} = \mathbb{Z}_2$, rearrange the weight function as a sum over configurations of the *k*-field and perform the integration over the fermions to get

$$Z_F[T, M] = \sum_{\{k_{xi}^{\alpha\beta}\}} (-\mathbf{i})^k \det(\mathbf{i}\partial [k]) \prod_{x\alpha\beta} (H_x^{\alpha\beta})^{k_x^{\alpha\beta}}, \quad (32)$$

where we introduced the abbreviations

$$k_x^{\alpha\beta} = \sum_i k_{xi}^{\alpha\beta} = k_{x0}^{\alpha\beta} + k_{x1}^{\alpha\beta}, \qquad k = \sum_{xi\alpha\beta} k_{xi}^{\alpha\beta} \quad (33)$$

and $\partial [k]$ is the matrix in which the $\{xi\alpha\}$ th row and $\{xi\beta\}$ th column of ∂ are removed whenever $k_{xi}^{\alpha\beta} = 1$. A similar expansion in terms of minors of a fermionic matrix was

recently presented within a transfer matrix approach in Ref. [48]. It is important to note that in the minor expansion there are constraints on the spin field $k_{xi}^{\alpha\beta}$ in order to get nonvanishing contributions to the weight function: due to $\psi_{xi}^{\alpha}\psi_{xi}^{\alpha} = \bar{\psi}_{xi}^{\alpha}\bar{\psi}_{xi}^{\alpha} = 0$ for fixed *x*, *i*, the sum over rows and columns of the matrix $k^{\alpha\beta}$ has to be zero or one,

$$\sum_{\alpha} k_{xi}^{\alpha\beta} \in \{0,1\} \quad \text{and} \quad \sum_{\beta} k_{xi}^{\alpha\beta} \in \{0,1\}.$$
(34)

Summing over the spinor index *i* leads to the following local constraints on the elements of the matrix $k_x = (k_x^{\alpha\beta})$ in Eq. (33):

$$\sum_{\alpha} k_x^{\alpha\beta} \in \{0, 1, 2\} \text{ and } \sum_{\beta} k_x^{\alpha\beta} \in \{0, 1, 2\}.$$
(35)

We can summarize these constraints in a local constraints function:

$$\delta_{\text{constr}}(k_x) = \begin{cases} 1 & \text{all local constraints are fulfilled}, \\ 0 & \text{else.} \end{cases}$$
(36)

Inserting Eq. (32) into the partition function (29), we observe that all contributions, with the exception of the minors, are given by a product over the lattice sites. After rescaling the variables $\tilde{T} = \sqrt{\lambda}T$ and $\tilde{M} = \sqrt{\lambda}M$, the partition function reads

$$Z(\lambda, \tilde{M}) = C \sum_{\{k_{xi}^{\alpha\beta}\}} \lambda^{-\frac{k}{2}} \det\left(\mathscr{D}[k]\right)$$
$$\cdot \prod_{x} \delta_{\text{constr}}(k_{x}) W_{\text{loc}}(k_{x}, \tilde{M}(x)), \quad (37)$$

with an overall factor $C = (\pi/\lambda)^{\gamma/2}$, where $\gamma = V(N_{\rm ir}^2 + 1)$. The local weight function is defined for any k_x by

$$W_{\rm loc}(k,\tilde{M}) = \int \prod_{i=1}^{N_{\rm tr}^2} \left(\frac{\mathrm{d}\tilde{t}_i}{\sqrt{\pi}}\right) e^{-\frac{1}{2}\mathrm{tr}\tilde{T}^2 - \frac{1}{4}(\mathrm{tr}\tilde{T})^2} \\ \cdot \prod_{\alpha\beta} \left(\tilde{T}^{\alpha\beta} + \frac{1}{2}\mathrm{tr}\tilde{T}\delta^{\alpha\beta} + \tilde{M}^{\alpha\beta}\right)^{k^{\alpha\beta}}.$$
 (38)

In the following we drop the tilde above variables again to simplify our notation. Because of Eq. (35) the exponent $k^{\alpha\beta}$ only takes the values 0,1, and 2. The integration variables are the expansion coefficients in $T = t_i T_i$.

The integration over the non-Cartan fields is performed in Appendix A and leads to the final form of the local weight function

$$W_{\rm loc}(k,M) = w_{\rm o}(k_{\perp})W_{{\bf p}(k)}(M), \qquad (39)$$

where $\mathbf{p}(k) = (k^{11}, k^{22}, ...)$ is the N_{ir} -component vector that collects the diagonal entries of $k^{\alpha\beta}$. Recall that these entries can assume the values 0, 1, or 2. In addition, the first factor $w_o(k_{\perp})$ is a non-negative integer depending on the nondiagonal entries of the matrix $(k^{\alpha\beta})$. The explicit form of w_o is given in Appendix A. The second factor is an integral over the Cartan subalgebra,

$$W_{\mathbf{p}}(M) = \int \prod_{i=1}^{N_{\mathrm{tr}}} \left(\frac{\mathrm{d}t_i}{\sqrt{\pi}}\right) e^{-tAt} \\ \cdot \prod_{\alpha} \left(T^{\alpha\alpha} + \frac{1}{2}\mathrm{tr}T + m_{\alpha}\right)^{p_{\alpha}}.$$
 (40)

The symmetric matrix A in the exponent can be written as

$$A_{ij} = \frac{1}{2}\operatorname{tr}(H_i H_j) + \frac{1}{4}\operatorname{tr} H_i \operatorname{tr} H_j \tag{41}$$

and the integration is only over the Cartan subalgebra with generators H_i . Although the final result does not depend on a specific choice of the generators, we use the generators given in Appendix A for the explicit calculations. W_p is a polynomial in m_{α} of degree p_{α} . The explicit form of $W_{\text{loc}}(k, M)$ is also given in Appendix A.

In this form it looks like the partition function can be simulated with a fermion-bag-like algorithm. Unfortunately the minors induce a severe sign problem. So far we could solve this problem for $N_{\rm ir} = 1$, where we applied the fermion bag algorithm to the model with chiral SLAC fermions [49]. In the following we refer to the integer number k in Eq. (33) as the lattice filling factor, because it counts how many fermions take part in the local interaction. Due to Pauli blocking it can only take the values $k = 0, ..., 2VN_{ir}$. In the strong-coupling limit $(\lambda \rightarrow 0)$ we find $k = 2VN_{ir}$ on every configuration such that every lattice site is occupied by the maximal number of fermions. Here we expect strong lattice artifacts due to saturation effects. In the weak-coupling limit $(\lambda \to \infty)$ we have k = 0 and the theory reduces to a theory of free fermions on the lattice.

B. Effective potential

In this section we derive an effective theory for the local chiral condensates $\langle \bar{\psi}_{\alpha} \psi_{\alpha} \rangle$ (no sum), or equivalently the expectation values of scalar fields in the Cartan subalgebra, that are related to the condensates by DSEs. Therefore, we constrain the local mass matrix defined in Eq. (28) to

$$M = \operatorname{diag}(m_1, \dots, m_{N_{\operatorname{ir}}})\delta_{x, x_0} \tag{42}$$

with a fixed lattice point x_0 and determine the dependence of the partition function on the parameters m_a . This allows us to relate contributions to the partition function $Z(\lambda, M)$ in the dual formulation to expectation values of chirally invariant local observables in the original vector formulation by differentiating with respect to these local mass parameters. Formally, the *M*-dependent partition function is given by

$$Z(\lambda, M) = \sum_{\mathcal{P}} \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} a_{\mathbf{p}} W_{\mathbf{p}}(M), \qquad (43)$$

where the first sum extends over the $(N_{\rm ir} + 1)(N_{\rm ir} + 2)/2$ triplets

$$\mathcal{P} = (P_0, P_1, P_2). \tag{44}$$

The elements of \mathcal{P} obey

$$P_k \in \{0, 1, ..., N_{ir}\}$$
 and $\sum_{k=0}^{2} P_k = N_{ir}.$ (45)

The second sum in Eq. (43) extends over the

$$\operatorname{Vol}(\mathcal{S}_{\mathcal{P}}) = \frac{N_{\mathrm{ir}}!}{P_0! P_1! P_2!}$$
(46)

permutations $S_{\mathcal{P}}$ of the N_{ir} -tuple **p** with P_0 elements equal to zero, P_1 elements equal to one, and P_2 elements equal to two. All together, the double sum in Eq. (43) consists of $3^{N_{ir}}$ terms. The coefficients $a_{\mathbf{p}}$ depend on λ , the volume, and further details of the lattice formulation such as, for example, the choice of lattice fermion derivative.

When all flavors have the same mass $(m_{\alpha} = m)$, the coefficients $a_{\mathbf{p}}$ do not depend on a specific permutation (permutation of flavors) but only on the three numbers P_k and we can write the partition function as

$$Z(\lambda, m) = \sum_{\mathcal{P}} a_{\mathcal{P}} \cdot \operatorname{Vol}(\mathcal{S}_{\mathcal{P}}) W_{\mathcal{P}}(m), \qquad (47)$$

where

$$W_{\mathcal{P}}(m) = W_{\mathbf{p}_{\text{sort}}}(M)|_{M(m_{\alpha} \to m)}$$

with $\mathbf{p}_{\text{sort}} = (\underbrace{0, \dots, 0}_{P_0 \text{ times}}, \underbrace{1, \dots, 1}_{P_1 \text{ times}}, \underbrace{2, \dots, 2}_{P_2 \text{ times}})$ (48)

is the weight of a particular representative in an orbit of the permutation group.

The constraint effective potential in the limit of a constant mass in flavor space is then given by the negative logarithm of the distribution function, i.e.,

$$V_{\text{eff}}(\lambda, T, m) = \frac{1}{2} \text{tr}(T^{c})^{2} + \frac{1}{4} (\text{tr}T^{c})^{2}$$
$$-\ln\left(\sum_{\mathcal{P}} a_{\mathcal{P}} \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} \prod_{\alpha=1}^{N_{\text{ir}}} \left(T^{\alpha \alpha} + \frac{1}{2} \text{tr}T + m\right)^{p_{\alpha}}\right).$$
(49)

Here, we have to sum over all permutations again, because no integration over the diagonal scalar fields is performed in the effective potential.

C. Relation to observables

In order to relate the coefficients $a_{\mathcal{P}}$ to expectation values, we take derivatives of the partition function (43) with respect to the local masses m_{α} and afterwards set $m_{\alpha} = m$. With the definition of moments of the Gaussian weights

$$W_{\mathbf{p},\mathbf{q}}(m) = \prod_{\alpha=1}^{N_{\mathrm{ir}}} \left(\frac{\partial^{q_{\alpha}}}{\partial m_{\alpha}^{q_{\alpha}}} \right) W_{\mathbf{p}}(M)|_{m_{\alpha} \to m}$$
(50)

we can write the partition function and its derivatives with respect to the local masses as

$$Z_{\mathbf{q}}(\lambda, m) = \sum_{\mathcal{P}} a_{\mathcal{P}} \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} W_{\mathbf{p}, \mathbf{q}}(m).$$
(51)

Since $W_{\mathbf{p}}(M)$ is a polynomial of degree p_{α} in m_{α} we only get a nonzero result in Eq. (50) when $q_{\alpha} \leq p_{\alpha}$. In the limit of a constant mass in flavor space, the expectation values $Z_{\mathbf{q}}$ do not depend on the ordering of the flavors, but only on the number Q_0 of zeroth, Q_1 of first, and Q_2 of second derivatives. Denoting the triple $\{Q_0, Q_1, Q_2\}$ by Q, we can write

$$\sigma_{\mathcal{Q}} = \frac{1}{\operatorname{Vol}(\mathcal{S}_{\mathcal{Q}})} \sum_{\mathbf{q} \in \mathcal{S}_{\mathcal{Q}}} Z_{\mathbf{q}}(\lambda, m).$$
(52)

The index sets Q and P are identical, with the constraints that their three entries sum to N_{ir} . The sum in Eq. (52) is over the Vol $(S_Q) = N_{ir}!/(Q_0!Q_1!Q_2!)$ permutations S_Q of **q**. The derivatives of the partition sum are directly related to expectation values via

$$\sigma_{\mathcal{Q}} = \lambda^{-\frac{|\mathbf{q}|}{2}} Z \cdot \left\langle \prod_{\alpha} (\bar{\psi}_{\alpha} \psi_{\alpha})^{q_{\alpha}} \right\rangle.$$
 (53)

Here, the right-hand side can be computed, up to the arbitrary normalization factor $Z = \sigma_{N_{ir},0,0}$, with conventional simulations of the Thirring model in the vector field formulation. Inserting Eq. (51) into Eq. (52), we obtain

$$\sigma_{\mathcal{Q}} = \sum_{\mathcal{P}} a_{\mathcal{P}} \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} \frac{1}{\operatorname{Vol}(\mathcal{S}_{\mathcal{P}})} \sum_{\mathbf{q} \in \mathcal{S}_{\mathcal{Q}}} W_{\mathbf{p},\mathbf{q}}(m).$$
(54)

We can also write this equation as a matrix equation,

$$\sigma_{\mathcal{Q}} = K_{\mathcal{QP}} a_{\mathcal{P}} \quad \text{with}$$

$$K_{\mathcal{QP}} = \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} \frac{1}{\operatorname{Vol}(\mathcal{S}_{\mathcal{Q}})} \sum_{\mathbf{q} \in \mathcal{S}_{\mathcal{Q}}} W_{\mathbf{p},\mathbf{q}}(m), \quad (55)$$

where *K* is a square matrix that can always be represented in an upper triangular form (due to $q_{\alpha} \le p_{\alpha}$) with nonzero diagonal elements. Therefore a unique solution for the coefficients $a_{\mathcal{P}}$ is given by

$$\vec{a} = K^{-1}\vec{\sigma}.\tag{56}$$

For symmetry reasons, the double sum for the matrix coefficients simplifies to

$$K_{\mathcal{QP}} = \frac{\operatorname{Vol}(\mathcal{S}_{\mathcal{P}})}{\operatorname{Vol}(\mathcal{S}_{\mathcal{Q}})} \sum_{\mathbf{q} \in \mathcal{S}_{\mathcal{Q}}} W_{\mathbf{p}(\mathcal{P}),\mathbf{q}}(m),$$
(57)

where $\mathbf{p}(\mathcal{P})$ is one representative in the equivalence class defined by \mathcal{P} . In this way, we can uniquely relate the coefficients \vec{a} to expectation values $\vec{\sigma}$ of the Thirring model. Alternatively, the coefficients \vec{a} can be calculated directly with a fermion bag simulation of the partition function (37).

V. SYMMETRIES OF THE EFFECTIVE POTENTIAL

A suitable order parameter for chiral symmetry breaking is the position of the global minimum of the effective potential. We can simplify the discussion of the potential, if we locally apply a chiral transformation such that the local condensates are in the Cartan subalgebra of $u(N_{ir})$. As part of the remaining symmetry of the Cartan subalgebra, it is possible to exchange flavors and the sign of the condensate, without changing physics. Therefore, minima of the effective potential can only occur in the directions

$$T_{\min}^{c} = \frac{2x}{N_{ir}} \begin{pmatrix} \pm 1 & \\ & \ddots & \\ & & \pm 1 \end{pmatrix} \in \mathbb{Z}_{2}^{N_{ir}}, \qquad (58)$$

where $x \sim (\bar{\psi}\psi)_{\alpha}$ is a free parameter. These directions where also spotted by the simulation results for the full effective potential. Once a direction is fixed we are left with the problem of finding the minimizing x. Physically equivalent solutions are related by a reflection $x \rightarrow -x$ or a permutation of flavors. The latter is given by the action of the Weyl group of $U(N_{\rm ir})$. We conclude that physically distinct solutions are characterized by the trace of $T_{\rm min}^c$,

$$tr T_{\min}^{c} = \frac{2x}{N_{ir}} n \quad \text{with}$$

$$n = \begin{cases} 0, 2, 4, \dots, N_{ir} & N_{ir} \text{ even,} \\ 1, 3, 5, \dots, N_{ir} & N_{ir} \text{ odd.} \end{cases}$$
(59)

This leads to $n_{sol} = \left[\frac{1}{2}N_{ir}\right] + 1$ different solutions for the potential. Every solution gives rise to a different breaking pattern of chiral symmetry, leaving different subgroups intact. A nonvanishing expectation value in direction *n* breaks the symmetry down to

$$U(N_{\rm ir}) \rightarrow U(n_+) \otimes U(n_-), \qquad n_{\pm} = \frac{N_{\rm ir} \pm n}{2}.$$
 (60)

A symmetric breaking with $n_{+} = n_{-}$ is only possible for even N_{ir} with n = 0. This is the proposed breaking of the reducible Thirring model. For the solution with $n = N_{ir}$ and $x \neq 0$, only parity symmetry is broken. Therefore, the solution with n = 0 is called a *Thirring*-like breaking while the solution with $n = N_{ir}$ is called a *Gross-Neveu*like breaking.

Along the different directions, the effective potential in the massless case is

$$V_{\text{eff}}(x,n) = \frac{2N_{\text{ir}} + n^2}{N_{\text{ir}}^2} x^2 - \ln\left(\sum_{\mathcal{P}} a_{\mathcal{P}} \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} (n+2)^{|\mathbf{p}|_{<}} (n-2)^{|\mathbf{p}|_{>}} \left(\frac{x}{N_{\text{ir}}}\right)^{|\mathbf{p}|}\right),$$
(61)

where the exponents are given by the (partial) sums,

$$|\mathbf{p}|_{<} = \sum_{\alpha \le n_{+}} p_{\alpha}, \quad |\mathbf{p}|_{>} = \sum_{\alpha > n_{+}} p_{\alpha}, \quad |\mathbf{p}| = \sum_{\alpha = 1}^{N_{\text{ir}}} p_{\alpha}. \quad (62)$$

In the massive case, $n \pm 2$ in Eq. (61) is replaced by $n \pm 2 + mN_{ir}$. Evaluating the above potential in the Thirring-like direction, we get

$$V_{\text{eff}}^{\text{Th}}(x) = \frac{2x^2}{N_{\text{ir}}} - \ln\left(\sum_{\mathcal{P}} a_{\mathcal{P}} \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} (-1)^{|\mathbf{p}|_{>}} \left(\frac{2x}{N_{\text{ir}}}\right)^{|\mathbf{p}|}\right), \quad (63)$$

and in the Gross-Neveu-like direction

$$V_{\text{eff}}^{\text{GN}}(x) = \frac{N_{\text{ir}} + 2}{N_{\text{ir}}} x^2 - \ln\left(\sum_{\mathcal{P}} a_{\mathcal{P}} \sum_{\mathbf{p} \in \mathcal{S}_{\mathcal{P}}} \left(\frac{N_{\text{ir}} + 2}{N_{\text{ir}}} x\right)^{|\mathbf{p}|}\right).$$
(64)

An important quantity to determine chiral symmetry breaking is the curvature κ of the effective potential at the origin, CRITICAL FLAVOR NUMBER OF THE THIRRING MODEL ...

$$\kappa(n) = \frac{d^2 V_{\text{eff}}(x, n)}{dx^2} \bigg|_{x=0}.$$
(65)

Possible phase transitions are expected to be second order. Thus, we conclude that chiral symmetry is unbroken if all curvatures $\kappa(n)$ are positive, while it is spontaneously broken if at least one of the $\kappa(n)$ is negative.

A. Effective theory for $N_{\rm ir} = 1$

For one irreducible flavor the Thirring model is equivalent to the Gross-Neveu model. This follows from the Fierz identity (21) for one flavor. For massless fermions the integral (40) vanishes for $\mathbf{p} = (1)$ such that only $\mathbf{p} = (0)$ and $\mathbf{p} = (2)$ remain. Likewise only multiderivatives with $\mathbf{q} = (0)$ and $\mathbf{q} = (2)$ appear. The *K*-matrix is then given by

$$K = \begin{pmatrix} 1 & \frac{3}{2} \\ 0 & 2 \end{pmatrix}. \tag{66}$$

The explicit form of the matrix depends on the normalization of the generators H_i given in Appendix A. With the arbitrary normalization $\sigma_{1,0,0} = 1$, we find for the coefficients \vec{a}

$$a_{1,0,0} = 1 - \frac{3}{4}\sigma_{0,0,1}$$
 and $a_{0,0,1} = \frac{1}{2}\sigma_{0,0,1}$, (67)

and obtain for the effective potential

$$V_{\rm eff}(x) = 3x^2 - \ln\left(1 - \frac{3}{4}\sigma_{0,0,1} + \frac{9}{2}\sigma_{0,0,1}x^2\right).$$
 (68)

The curvature of the potential at the origin is then

$$\kappa = 18 \frac{\sigma_{0,0,1} - 4/9}{\sigma_{0,0,1} - 4/3}, \qquad \sigma_{0,0,1} = \frac{1}{\lambda} \langle (\bar{\psi}\psi)^2 \rangle_{m=0}.$$
(69)

At the critical inverse coupling λ_c the curvature vanishes and we get $\sigma_{0,0,1}(\lambda_c) = 4/9$. The condensate $\langle (\bar{\psi}\psi)^2 \rangle$ is later calculated within the strong-coupling expansion since Monte Carlo simulations in the vector formulation face a severe sign problem.

The fermion bag approach, which is free of a sign problem and directly yields the coefficients \vec{a} , will be discussed in a follow-up paper [49].

B. Effective theory for $N_{\rm ir} = 2$

For two massless irreducible flavors the Thirring model is equivalent to the massless reducible model with N = 1. For a vanishing *m* the nonvanishing **p** configurations are $\mathbf{p} = \{(0,0), (1,1), (2,0), (0,2), (2,2)\}$. The configurations $\mathbf{p} = (0,2)$ and $\mathbf{p} = (2,0)$ are equivalent under flavor exchange and the corresponding coefficients are

TABLE I. Configurations and weights for $N_{\rm ir} = 2$ in the massless limit.

Ι	а	р	$W_{(0,0)}$	$W_{(1,1)}$	$W_{(2,0)}$	$W_{(0,2)}$	$W_{(2,2)}$
1	a _{2,0,0}	(0, 0)	1	0	0	0	0
2	<i>a</i> _{0,2,0}	(1, 1)	$\frac{1}{2}$	1	0	0	0
3	$a_{1,0,1}$	(2, 0)	$\frac{3}{2}$	0	2	0	0
3	$a_{1,0,1}$	(0, 2)	$\frac{3}{2}$	0	0	2	0
4	<i>a</i> _{0,0,2}	(2, 2)	$\frac{11}{4}$	2	3	3	4

the same. The weights are listed in Table I. The *K*-matrix and its inverse are then given by

$$K = \begin{pmatrix} 1 & \frac{1}{2} & 3 & \frac{11}{4} \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 2 & 3 \\ 0 & 0 & 0 & 4 \end{pmatrix},$$
$$K^{-1} = \begin{pmatrix} 1 & -\frac{1}{2} & -\frac{3}{2} & \frac{11}{16} \\ 0 & 1 & 0 & -\frac{1}{2} \\ 0 & 0 & \frac{1}{2} & -\frac{3}{8} \\ 0 & 0 & 0 & \frac{1}{4} \end{pmatrix}.$$
 (70)

We obtain with the normalization $\sigma_{2,0,0} = 1$

$$a_{2,0,0} = 1 - \frac{1}{2}\sigma_{0,2,0} - \frac{3}{2}\sigma_{1,0,1} + \frac{11}{16}\sigma_{0,0,2},$$

$$a_{0,2,0} = \sigma_{0,2,0} - \frac{1}{2}\sigma_{0,0,2},$$

$$a_{1,0,1} = \frac{1}{2}\sigma_{1,0,1} - \frac{3}{8}\sigma_{0,0,2},$$

$$a_{0,0,2} = \frac{1}{4}\sigma_{0,0,2}.$$
(71)

The Gross-Neveu-type potential is given by

$$V_{\rm eff}^{\rm GN}(x) = 2x^2 - \ln\left(a_{2,0,0} + 4a_{0,2,0}x^2 + 8a_{1,0,1}x^2 + 16a_{0,0,2}x^4\right)$$
(72)

and the Thirring-type potential is given by

$$V_{\text{eff}}^{\text{Th}}(x) = x^2 - \ln \left(a_{2,0,0} - a_{0,2,0} x^2 + 2a_{1,0,1} x^2 + a_{0,0,2} x^4 \right).$$
(73)

In the following sections, the observables σ are calculated first in the strong-coupling expansion and after that with Monte Carlo simulations in the vector formulation of the Thirring model which for $N_{\rm ir} \ge 2$ has no sign problem. In this section we compute the effective potential in the strong-coupling expansion. The lattice partition function in the presence of fermion sources is given by

$$Z[\eta,\bar{\eta}] = \int \mathcal{D}v \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-\sum_{x} (\lambda v^{2} - \bar{\psi}(i\not\!\!\!d + \not\!\!\!)\psi - \bar{\psi}\eta - \bar{\eta}\psi)}$$

$$= K \left[\frac{\delta}{\delta\eta}, \frac{\delta}{\delta\bar{\eta}}\right] \int \mathcal{D}v \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{-\sum_{x} (\lambda v^{2} - \bar{\psi}\not\!\!/\psi - \bar{\psi}\eta - \bar{\eta}\psi)},$$

(74)

where the sum is over all lattice points x and where we already use dimensionless fields and inverse coupling λ . The kinetic operator is given by

$$K\left[\frac{\delta}{\delta\eta},\frac{\delta}{\delta\bar{\eta}}\right] = e^{-\sum_{x,y}\frac{\delta}{\delta\eta_x}\mathbf{i}\not\theta_{xy}\frac{\delta}{\delta\eta_y}} \tag{75}$$

and ∂_{xy} is a lattice-regularized derivative operator. With a rescaling of the sources according to $\chi = \lambda^{1/4} \eta$ we shift the explicit λ dependence to the kinetic operator. Furthermore, derivatives with respect to the sources χ exactly reproduce the expectation values (52) needed in the effective potential. After performing the integration over the vector field and the fermions (details are given in Appendix B), we obtain our master equation for the strong-coupling expansion:

$$Z[\chi,\bar{\chi}] = K\left[\frac{\delta}{\delta\chi}, \frac{\delta}{\delta\bar{\chi}}, \lambda\right] \prod_{x} \sum_{k=0}^{N_{\rm ir}} \frac{\Gamma(\frac{3}{2} + N_{\rm ir} - k)}{\Gamma(2k+2)} ((\bar{\chi}_{x}\gamma_{\mu}\chi_{x})^{2})^{k}.$$
(76)

In the following we are only interested in local fermionic observables. Therefore, *after* applying the kinetic operator K we set the source χ at all points x with the exception of x_0 to zero. With the definitions

$$K^{(n)} = (-1)^n \frac{\lambda^{n/2}}{n!} \left(\sum_{x,y} \frac{\delta}{\delta \chi_x} i \partial_{xy} \frac{\delta}{\delta \bar{\chi}_y} \right)^n,$$

$$F^{(k)}(x) = \frac{\Gamma(\frac{3}{2} + N_{\rm ir} - k)}{\Gamma(2k + 2)} ((\bar{\chi}_x \gamma_\mu \chi_x)^2)^k \tag{77}$$

we can write the partition function as

$$Z[\chi_{x_0}, \bar{\chi}_{x_0}] = \sum_n K^{(n)} \prod_x \sum_k F^{(k)}(x)|_{\chi_{x \neq x_0} = 0}.$$
 (78)

In Appendix B we show, that in the infinite-volume limit and up to any finite order n in the expansion of the kinetic operator, the partition function has the form

$$Z[\chi_{x_0}, \bar{\chi}_{x_0}] = C(\lambda) \sum_{k} F^{(k)}(x_0)$$

= $C(\lambda) \sum_{k=0}^{N_{\rm ir}} \frac{\Gamma(\frac{3}{2} + N_{\rm ir} - k)}{\Gamma(2k+2)} ((\bar{\chi}_{x_0} \gamma_{\mu} \chi_{x_0})^2)^k,$ (79)

where $C(\lambda)$ is an unknown function that cancels in expectation values. Then the solution for the expectation values is

$$\sigma_{Q} = \lambda^{-Q_{2}-Q_{1}/2} \prod_{\alpha=1}^{Q_{1}} \partial_{m_{\alpha}} \prod_{\beta=Q_{1}+1}^{Q_{1}+Q_{2}} \partial_{m_{\beta}}^{2} \ln Z(M)|_{M=0}$$
$$= \frac{\Gamma(\frac{3}{2}+N_{\rm ir}-Q_{2})}{\Gamma(\frac{3}{2}+N_{\rm ir})} \delta_{Q_{1},0}.$$
(80)

For the vector interaction we obtain

$$\frac{1}{4\lambda} \langle (\bar{\psi} \gamma_{\mu} \psi)^2 \rangle = N_{\rm ir} \tag{81}$$

and it follows that the normalized lattice filling factor takes its maximal value $\langle k_{\text{norm}} \rangle = 1$. In conclusion, within the strong-coupling expansion we are not able to leave the strong-coupling phase where strong lattice artifacts dominate due to complete Pauli blocking on every lattice site.

A. Results for the effective potential

With the observables from the strong-coupling expansion we can calculate the effective potential in the lattice artifact phase [which was discussed around Eq. (3)] in the infinite-volume limit. The results for various flavor numbers N_{ir} are shown in Fig. 1. For $N_{ir} = 1$, the Gross-Neveu-type potential reads

$$V_{\rm eff}^{\rm GN}(x) = 2x^2 + \ln\left(\frac{1}{4x^2}\right) = -2\ln(2x) + 2x^2.$$
 (82)

It has a global minimum at $x = \pm 1/\sqrt{2}$ and parity symmetry is spontaneously broken. Its curvature at the origin diverges.

For odd flavor numbers parity symmetry is always broken at strong coupling with a minimum of the potential in the Gross-Neveu-like direction. With increasing N_{ir} the curvature at the origin decreases. Extrapolating the curvature of the Gross-Neveu-type potential to $N_{ir} = \infty$ predicts a broken symmetry in this limit. At strong lattice coupling there is no critical flavor number and parity symmetry is always broken for odd N_{ir} . For $N_{ir} = 7$ we observe that the potential with n = 1 has a positive curvature while this is not the case for smaller N_{ir} , indicating that the breaking of parity symmetry becomes weaker for larger flavor numbers. CRITICAL FLAVOR NUMBER OF THE THIRRING MODEL ...

PHYSICAL REVIEW D 96, 094504 (2017)



FIG. 1. Effective potentials in the strong-coupling expansion for $N_{\rm ir} = 1, 2, 3, 4, 5, 7$ with different values of n.

For $N_{\rm ir} = 2$, the potentials at strong coupling read

$$V_{\text{eff}}^{\text{GN}}(x) = 2x^2 - \ln(8x^4 - 4x^2 + 1) + \ln\left(\frac{15}{8}\right)$$
$$= \ln\left(\frac{15}{8}\right) + 6x^2 - \frac{32x^6}{3} + O(x^8),$$
$$V_{\text{eff}}^{\text{Th}}(x) = x^2 - \ln(x^4 + 2x^2 + 2) + \ln\left(\frac{15}{4}\right)$$
$$= \ln\left(\frac{15}{8}\right) + \frac{x^6}{6} + O(x^8). \tag{83}$$

The curvature in the Gross-Neveu direction is positive. In the Thirring-like direction the leading power is x^6 with a positive coefficient such that the minimum of the potential is at x = 0. Hence chiral symmetry is unbroken. For larger even N_{ir} the curvature in all directions is positive. In conclusion, chiral and parity symmetry is unbroken for any even N_{ir} and parity symmetry is broken for any odd N_{ir} in the strong-coupling limit. In the next section we will investigate with Monte Carlo simulations whether these results hold outside the lattice artifact phase and in the continuum limit.

VII. SIMULATION RESULTS

The simulations have been performed with the lattice action

$$S(\lambda) = N_{\rm ir} \left(\frac{\lambda}{2} \sum_{x} v_{\mu}^2(x) - \ln \det(i \not D) \right)$$
(84)

for fermions in the irreducible representation of the Clifford algebra with $\gamma_{\mu} = \sigma_{\mu}$. Note, that we have rescaled $\lambda \rightarrow N_{ir}\lambda$ for our simulations. We use the SLAC derivative for fermions with a Hermitian Dirac operator,

$$\mathbf{i} \mathbf{D} = \sigma_{\mu} (\mathbf{i} \partial_{\mu}^{\mathrm{SLAC}} + v_{\mu}), \tag{85}$$

because it preserves the continuum $U(N_{ir})$ chiral symmetry and the discrete parity symmetry exactly even at finite lattice spacing. Note that Wilson fermions in the irreducible representation also preserve the continuum chiral symmetry but break parity. In order to implement antiperiodic boundary conditions for the fermions in the time direction, we simulate on lattices with volumes $V = L \times (L - 1)^2$ with even L. Most of our simulations have been carried out on lattices with L = 8, 12, 16, and 20 with statistics of 1000 to 10 000 configurations. For all flavor numbers we use a rational HMC algorithm with

$$(\det(\not D \not D^{\dagger})^{N_{\rm ir}/2N_{\rm PF}})^{N_{\rm PF}},\tag{86}$$

where the number of pseudofermions is $N_{\rm PF} = 2N_{\rm ir}$. In order to calculate the expectation values of powers of the condensate, we use $N_{\rm est} = 200 \times N_{\rm ir}$ stochastic estimators for the fermion propagator on every Monte Carlo configuration.

The fermion determinant det(iD) is real but not necessarily positive. Therefore we do not have a sign problem for even flavor numbers. Furthermore, simulations on smaller lattices, where we can compute the fermion determinant numerically, showed that we only have a sign problem for $N_{ir} = 1$. In this case, the sign problem can be solved with a fermion-bag-inspired algorithm and



FIG. 2. Effective potentials for $N_{\rm ir} = 4$ and $\lambda = 0.118$ along the different directions labeled by n = 0, 2, and 4 on a lattice with L = 16.

appropriate resummations of certain weights. More details on the sign problem will be published in Ref. [49]. For $N_{\rm ir} = 2$ to $N_{\rm ir} = 11$ we determined the observables σ_Q in Eq. (53) via Monte Carlo simulations in the vector formulation and calculated the effective potential with the formalism described in the previous sections.

The potentials for $N_{\rm ir} = 4$ and corresponding values of *n* on a $16 \times 15 \times 15$ lattice are depicted in Fig. 2 for $\lambda = 0.118$. Statistical errors are always obtained with a jackknife procedure and are indicated by the width of the curves. It turns out that for every value of λ the minimum of the potential is always at the origin x = 0. Therefore we conclude that there is no spontaneous chiral or parity symmetry breaking for $N_{\rm ir} = 4$, at least on the lattice with L = 16.

For $N_{\rm ir} = 5$ the potentials are depicted in Fig. 3 for two values of λ . For $\lambda = 0.102$ the potential has two global minima in the Gross-Neveu direction at $x \approx \pm 0.22$ while for the larger value $\lambda = 0.118$ the minimum of the potential is at x = 0. This suggests that for $N_{\rm ir} = 5$ parity is broken at strong coupling (small λ). On larger lattices this result still holds true.

To check, that our conclusions are not blurred by lattice artifacts we investigated the transition from the strong to weak coupling regime more carefully and determined the critical coupling where the lattice theory shows a transition from an artificial lattice phase at strong coupling to a continuum phase at weak coupling for all simulated N_{ir} . In the lattice artifact phase and in the infinite-volume limit, observables should only trivially depend on λ . An important quantity to investigate here is the first derivative of the partition function with respect to λ . It is connected to the normalized fermion filling factor in the dual variables approach,

$$\langle k_{\text{norm}} \rangle = \frac{\langle k \rangle}{2VN_{\text{ir}}} \in [0, 1],$$
 (87)



FIG. 3. Effective potentials for $N_{\rm ir} = 5$ at $\lambda = 0.102$ (upper panel) and $\lambda = 0.118$ (lower panel) for the different directions n = 1, 3, and 5 on a lattice with L = 16.

where k has been defined in Eq. (33), and therefore is an interesting quantity to investigate lattice artifacts. The relation is

$$\frac{\lambda}{2N_{\rm ir}V}\frac{d\ln Z(\lambda)}{d\lambda} = C + \langle k_{\rm norm} \rangle(\lambda), \tag{88}$$

where the constant *C* only depends on the flavor number. At the transition from the lattice-artifact phase to the physical weak-coupling phase, we expect a jump or a peak in the first derivative $\partial_{\lambda} \langle k_{\text{norm}} \rangle$. The critical value λ^* is then obtained as the position of the jump (or peak) in the infinite-volume limit. The results for $\langle k_{\text{norm}} \rangle$ are shown in Fig. 4 for different N_{ir} and volumes. Below λ^* , the expectation value $\langle k_{\text{norm}} \rangle$ depends only weakly on the volume and increases towards its strong-coupling value $\langle k_{\text{norm}} \rangle = 1$ with increasing lattice volume. The curves for different volumes intersect close to the critical λ^* . In the physical phase $\langle k_{\text{norm}} \rangle$ decreases with increasing lattice volume.

The variation of $\langle k_{\text{norm}} \rangle$ with λ for $N_{\text{ir}} = 9$ is depicted in Fig. 5. It stays finite in the infinite-volume limit and develops a jump at the critical coupling. We observe that for larger flavor numbers the curves for L = 16 and L = 20



FIG. 4. Lattice filling factor $\langle k_{norm} \rangle$ for different N_{ir} and lattices volumes. Larger volumes are indicated by a darker color shade.



FIG. 5. Derivative of the lattice filling factor $\langle k_{\text{norm}} \rangle$ for $N_{\text{ir}} = 9$ and different lattice volumes.

lie almost on top of each other, indicating that finite-volume effects are already small on these relatively small lattices. Even for smaller N_{ir} , finite-volume effects are small on the larger lattices. Therefore, we identify the critical coupling on our largest lattice as the infinite-volume coupling λ^* . The relatively small finite-size effects are an additional advantage of the SLAC derivative that approximates the continuum derivative for a fixed number of lattice points much better than the naive central derivative used for Wilson or staggered fermions [35]. The results for the

critical λ^* for all $N_{\rm ir}$ between 1 and 11 are displayed in Table II. The results for $N_{\rm ir} = 1$ with a lattice size L = 8were obtained with a fermion bag algorithm directly calculating the coefficients $a_{\mathcal{P}}$ of the effective potential. The lower curve in the phase diagram in Fig. 8 shows the phase boundary, separating the strong-coupling latticeartifact regime from the physical weak-coupling regime. We see that with increasing flavor number the critical value λ^* decreases monotonically.

After having localized the transition point between the artifact and physical phases, we calculate the curvature κ of the effective potential at the origin as a function of λ and compare the critical value λ_c , at which the curvature vanishes, to the critical value λ^* of the artifact transition. For even $N_{\rm ir}$ we show κ for both the Gross-Neveu-like breaking as well as the Thirring-like breaking. The results for $N_{ir} \in \{2, 4\}$ are depicted in Fig. 6, together with the results from the strong-coupling expansion (solid lines). The grey bars show the allowed critical values λ^* . For $N_{\rm ir} = 2$ the curvature in the Thirring direction vanishes at strong coupling and increases at the transition from the strong-coupling lattice-artifact phase to the weak-coupling physical phase. The potential in the Gross-Neveu direction is always positive. Therefore, we conclude that there is no chiral symmetry breaking for $N_{\rm ir} = 2$, which corresponds to one reducible flavor. For $N_{\rm ir} = 4$ the curvature for all directions is always positive and there is clearly no chiral symmetry breaking. We also checked this for larger even numbers of flavors with the same result: chiral symmetry is always unbroken for even flavor numbers. This implies that there is no spontaneous symmetry breaking for all reducible Thirring models. This is one important conclusion of our work which conflicts with earlier findings but agrees with very recent simulations [30,41,42].

For odd flavor numbers there is no Thirring-like potential. Furthermore, we checked that the minimum of the full potential is either at the origin or in the Gross-Neveu direction. Therefore we show the curvature in the Gross-Neveu direction in Fig. 7 for $N_{ir} = 3, 5, 7, 9$, and 11 on the 16×15^2 lattice. The critical inverse coupling λ_c is defined by vanishing curvature and values are shown in Table III. Again we compare the coupling λ_c to the critical value of the strong-coupling transition λ^* . For $N_{ir} = 3, 5,$ and 7 we observe that the parity phase transition at λ_c lies within the physical phase, i.e., $\lambda_c > \lambda^*$ and we conclude that parity

TABLE II. Critical λ^* for different flavor numbers and lattice volumes. For larger lattices there are small finite-size effects. Simulations for $N_{ir} = 1$ were done with a fermion bag algorithm.

N _{ir}	1	2	3	4	5	7	9	11
$\lambda^*(L=8)$	0.35(1)	0.223(6)	0.158(4)	0.122(4)	0.098(2)	0.073(2)	0.058(2)	0.048(2)
$\lambda^*(L=12)$		0.214(4)	0.149(4)	0.114(3)	0.094(3)	0.068(2)	0.054(2)	0.046(2)
$\lambda^*(L=16)$		0.208(4)	0.146(4)	0.112(3)	0.091(2)	0.067(1)	0.054(1)	0.045(1)
$\lambda^*(L=20)$						0.066(1)	0.053(1)	0.045(1)



FIG. 6. Curvature of the effective potential at the origin for $N_{\rm ir} = 2$ (top) and $N_{\rm ir} = 4$ (bottom) compared to the critical coupling λ^* (grey bar). The width of the grey bar indicates the statistical error of λ^* .



FIG. 7. Curvature of the effective potential for different odd flavor numbers N_{ir} on the lattice with L = 16 (darker shade) and L = 20 (lighter shade). The colored bars and their widths denote the corresponding transition from the lattice-artifact phase to the physical phase at λ^* including the statistical error.

TABLE III. Critical inverse coupling λ_c on a lattice with L = 16 and L = 20 separating the parity-broken from the parity-symmetric phase.

N _{ir}	1	3	5	7	9	11
$\overline{\lambda_{\rm c}(L=16)}$	0.39(1)	0.172(2)	0.110(4)	0.077(1)	0.054(2)	
$\lambda_{\rm c}(L=20)$				0.074(2)	0.051(2)	

symmetry is spontaneously broken for these flavor numbers. For $N_{\rm ir} = 11$ the curvature is always positive and therefore parity symmetry is always unbroken. For $N_{\rm ir} = 9$ both critical couplings coincide within error bars and it is still unclear whether parity symmetry is spontaneously broken or not.

The upper curve in Fig. 8 shows the linear interpolation between the critical values λ_c for the Thirring model with odd flavor numbers. We conclude that the critical flavor number for parity breaking is $N_{\rm ir}^{\rm cr} = 9$. The minimum $x_{\rm min}$ of the effective potential is an order parameter for the breaking of parity symmetry and therefore related to a parity condensate π . For a second-order phase transition, π should decrease continuously to zero with increasing coupling λ . In Fig. 9 we show the condensate for different odd flavor numbers in the physical phase on a lattice with L = 16. For $N_{ir} \in \{3, 5, 7\}$ it decreases monotonically to zero while for $N_{\rm ir} = 9$ the condensate approaches zero at vanishing *renormalized* inverse coupling $\lambda_{\rm R} = \lambda - \lambda^*$. This observation is consistent with the scenario proposed in Refs. [6,14,15] where parity at the critical flavor number is only broken at vanishing inverse coupling.

The length dimension of the physical inverse coupling is $[\lambda_{phys}] = -1$ and therefore the dimensionless lattice coupling is related to a physical coupling by $a\lambda_{phys} = \lambda$. In the cases where we did not find a second-order phase



FIG. 8. Phase diagram for odd flavor numbers. The lower curve separates the strong-coupling lattice-artifact phase from the physical weak-coupling phase and is known for odd and even flavor numbers. The upper curve for odd flavor numbers shows the physical phase transition associated with breaking of the discrete parity symmetry.



FIG. 9. Parity condensate $\pi \sim x_{\min}$ for odd flavor numbers in the physical phase as a function of the renormalized inverse coupling $\lambda_{\rm R} = \lambda - \lambda^*$.

transition, we can perform the continuum limit by $\lambda_{\rm R} \rightarrow 0$ for a fixed physical inverse coupling within the physical phase. For odd $N_{\rm ir} < N_{\rm ir}^{\rm cr}$, where we spotted a second-order phase transition, we build the continuum limit by $\lambda \rightarrow \lambda_c$, corresponding to a non-Gaussian fixed point. Both limits coincide for $N_{\rm ir} = N_{\rm ir}^{\rm cr}$.

VIII. CONCLUSIONS AND DISCUSSION

Our main observation is that the irreducible Thirring model with an odd number of irreducible flavors behaves differently compared to the model with an even number of flavors. For massless fermions the latter class is equivalent to the class of well-studied reducible models with $N = 2N_{\rm ir}$. While for odd flavor numbers we find a critical flavor number $N_{\rm ir}^{\rm cr} = 9$ below which the Thirring model shows spontaneous parity breaking, for even flavor numbers, neither parity nor chiral symmetry is broken. This implies that *no spontaneous breaking of chiral symmetry exists for all reducible models*, which are usually discussed in the literature. Our earlier, more straightforward simulations with SLAC fermions already pointed to this result [40,41]. Furthermore, our conclusions are also consistent with recent simulations with domain-wall fermions [29,30].

Regarding the older lattice simulations with staggered fermions [19–24], including the fermion bag formulation [27], their results seem not to be valid for the reducible Thirring model, likely because their lattice formulation does not have the correct symmetry. To see this more clearly we should recall that for massive fermions the reducible models are no longer equivalent to the irreducible models with $N_{ir} = 2N$ flavors. For reducible massive models log det($i\not D + im$) is real and does not contain any imaginary Chern-Simons-type term as it does for all irreducible models are not equal to the zero-mass limits of the irreducible models are not equal to the zero-mass limits of the irreducible models with $N_{ir} = 2N$. Thus any lattice simulation (or any

other regularization) which needs a fine-tuning to reach the chiral limit may yield erroneous results. It may very well happen that instead of the massless reducible model one simulates an irreducible model at small masses. This could be a partial explanation of why the earlier prediction $8 < N_{\rm irr}^{\rm cr} < 12$ is consistent with ours, but only for odd flavor numbers in the irreducible representation.

Also note that most of the previous analytical studies focused on csb in the reducible representation assuming conservation of the reducible parity (9) [6,14,15,43] or did not distinguish between irreducible and reducible models [13,46]. For example, the authors of Ref. [13] used DSEs to investigate mass generation either from parity or chiral symmetry breaking and found $N^{cr} \approx 12.97/D$, where *D* is the dimension of the representation of the Clifford algebra. These earlier results are not consistent with ours and other ongoing simulations with chiral fermions [29,30], where csb is not present at all.

Not much emphasis was put on parity breaking for odd flavor numbers in the irreducible representation, but most studies found a cancellation of the Chern-Simons terms for even N_{ir} [6,15,43]. As discussed above, this is a delicate issue and the answer depends on the order of limits $\lim_{V\to\infty}$ and $\lim_{m\to 0}$. We obtained our results for zero masses in a finite volume, in which case the fermion determinant is real and no (imaginary) Chern-Simons term can be generated. If one considers the Thirring model on \mathbb{R}^3 , such a term can show up for even N_{ir} [44,45,50]. It also can show up if one uses a regularization which breaks chiral symmetry explicitly.

To obtain our novel results it was essential to employ chiral fermions. But with massless chiral fermions it seems impossible to calculate the chiral condensates directly [40]. The main ingredient to circumvent this difficulty was to use both the vector and the matrix formulations of the Thirring models. By introducing auxiliary (local) masses we could relate expansion coefficients for effective potentials of a massless model in the matrix formulation to expectation values of condensates in the vector formulation of the same model. The actual proof and explicit mapping from coefficients to condensates are based on a reformulation of the matrix models in terms of dual spin variables $k_{xi}^{\alpha\beta}$. They are introduced to represent the result of the integration over the fermionic variables.

Our analytic results hold for other types of chiral fermions. Actually, at present we replace SLAC fermions by overlap fermions in our simulation code to calculate the condensates related to the coefficients of effective potentials.³ We expect to find results comparable to SLAC fermions and in particular a similar value for $N_{\rm ir}^{\rm cr}$.

We already mentioned that the irreducible one-flavor model (which is equivalent to the irreducible one-flavor Gross-Neveu model) has a severe sign problem. We could show that in the dual formulation there are subtle

³In collaboration with Rajamani Narayanan.

cancellations of terms such that the sign problem actually goes away [49]. It would be interesting to show that a similar fermion-bag-type algorithm without sign problems exists for multiflavor Thirring models.

ACKNOWLEDGMENTS

We are grateful to Shailesh Chandrasekharan, Holger Gies, Lukas Janssen, Simon Hands, Rajamani Narayanan, and Urs Wenger for helpful discussions and comments. This work was supported by the Helmholtz International Center for FAIR within the LOEWE initiative of the State of Hesse. D. S. was supported by the graduate school GRK 1523/2. Simulations were performed on the LOEWE-CSC at the University of Frankfurt and on the HPC cluster at the University of Jena.

APPENDIX A: FERMION BAG

Here we present further details concerning the functional integral in the dual formulation introduced in Sec. IVA. We summarize relevant results up to the point where simulations with a fermion bag algorithm are possible. For that purpose it is convenient to use the Lagrangian

$$\mathcal{L} = \bar{\psi}(\mathrm{i}\partial \!\!\!/ + \mathrm{i}T + \mathrm{i}\phi)\psi + \frac{\lambda}{2}\mathrm{tr}T^2 + \lambda\phi^2, \qquad (\mathrm{A1})$$

with an additional scalar field ϕ in place of the equivalent Lagrangian (23). In the formulation with a scalar field the integrals that occur over the Hermitian matrix *T* are more readily calculated. The equivalence of the two formulations can be seen after splitting *T* into its trace-free contribution and a multiple of the identity. Then one observes that the integration over tr*T* in the formulation (23) and over tr*T* and ϕ in the formulation (A1) yield the same results, up to an overall factor $\propto \lambda^{V/2}$. The transition to dual variables is the same as for the Lagrangian without scalar fields, with the only difference being that the matrix $(H_x^{\alpha\beta})$ in the interaction term is now

$$H_x^{\alpha\beta} = T_x^{\alpha\beta} + \phi_x \delta^{\alpha\beta} + M_x^{\alpha\beta}, \qquad (A2)$$

and this gives rise to a slightly different local weight and an additional integration over ϕ . Instead of Eq. (38) one obtains

$$W_{\rm loc}(k,M) = \int \prod_{i=1}^{N_{\rm ir}^2} \left(\frac{\mathrm{d}t_i}{\sqrt{\pi}}\right) \int \mathrm{d}\phi \mathrm{e}^{-\frac{1}{2}\mathrm{tr}T^2 - \phi^2} \\ \cdot \prod_{\alpha\beta} (T^{\alpha\beta} + \phi\delta^{\alpha\beta} + M^{\alpha\beta})^{k^{\alpha\beta}}, \qquad (A3)$$

up to an irrelevant overall factor $\propto \sqrt{\lambda}$. Since *T* is Hermitian, the exponential function factorizes as follows:

PHYSICAL REVIEW D 96, 094504 (2017)

$$e^{-\frac{1}{2}trT^2} = \prod_{\alpha < \beta} e^{-|T^{\alpha\beta}|^2} \prod_{\alpha} e^{-(T^{\alpha\alpha})^2}.$$
 (A4)

It implies the following factorization of the local weight:

$$W_{\rm loc}(k,M) = \int \mathrm{d}\phi \mathrm{e}^{-\phi^2} W_{\mathbf{p}(k)}(\phi,M) w_{\rm o}(k_{\perp}), \qquad (A5)$$

where the integral over the off-diagonal matrix elements produces the *M*- and ϕ -independent factor

$$w_{\rm o}(k_{\perp}) = \prod_{\alpha < \beta} W_{\perp}(k^{\alpha\beta}, k^{\beta\alpha}). \tag{A6}$$

The function W_{\perp} is determined by a complex Gaussian integral

$$W_{\perp}(k,k') = \int \frac{\mathrm{d}z \mathrm{d}\bar{z}}{\pi} \mathrm{e}^{-|z|^2} z^k (z^*)^{k'} = k! \delta_{kk'}, \quad (A7)$$

leading to the local constraints $k^{\alpha\beta} = k^{\beta\alpha}$. Thus we obtain

$$w_{\rm o}(k_{\perp}) = \prod_{\alpha < \beta} (k^{\alpha \beta})! \tag{A8}$$

Recall that the entries of the symmetric matrix $(k^{\alpha\beta})$ must obey the local constraints in Eq. (35).

The integral over the Cartan variables leads to a term similar to Eq. (40) and given by

$$W_{\mathbf{p}(k)}(\phi, M) = \int \prod_{i=1}^{N_{\rm ir}} \left(\frac{\mathrm{d}t_i}{\sqrt{\pi}}\right) e^{-tAt} \cdot \prod_{\alpha} (T^{\alpha\alpha} + \phi + m_{\alpha})^{p_{\alpha}(k)},$$
(A9)

with $p_{\alpha}(k) = k^{\alpha \alpha} \in \{0, 1, 2\}$. Note that the diagonal matrix elements $T^{\alpha \alpha}$ are linear functions in the integration variables t_i . The symmetric positive matrix *A* has matrix elements

$$A_{ij} = \frac{1}{2} \operatorname{tr}(H_i H_j) \tag{A10}$$

and differs slightly from the matrix (41) in the formulation without a scalar field ϕ . The normalization of the generators H_i is such that

$$\operatorname{tr}(H_i H_j) = \begin{pmatrix} \frac{4}{N_{ir}+2} & 0 & 0 & 0 & \dots & 0\\ 0 & 2 & -1 & 0 & \dots & 0\\ 0 & -1 & 2 & -1 & \ddots & \vdots\\ \vdots & \ddots & \ddots & \ddots & \ddots & 0\\ 0 & \dots & 0 & -1 & 2 & -1\\ 0 & \dots & 0 & 0 & -1 & 2 \end{pmatrix}. \quad (A11)$$

CRITICAL FLAVOR NUMBER OF THE THIRRING MODEL ...

The final integration over the variables t_i yields the factor

$$W_{\mathbf{p}}(\phi, M) = \prod_{\alpha: p_a = 1} (\phi + m_a) \prod_{\alpha: p_a = 2} (1 + (\phi + m_a)^2).$$
(A12)

In the limit of $m \to 0$, the ϕ integral (A5) vanishes for odd P_1 , and for even P_1 it is given by the confluent hypergeometric function,

$$W_{\rm loc}(k) = w_{\rm o}(k_{\perp})\Gamma\left(\frac{1+P_1}{2}\right)U\left(\frac{1+P_1}{2}, \frac{3+P_1}{2}+P_2, 1\right),$$
(A13)

where, as in the main body of the text, the number P_k with $k \in \{0, 1, 2\}$ counts the number of indices p_α with $p_\alpha = k$. In this form, the local weights are suitable for simulations with a fermion bag algorithm. Unfortunately, this formulation does not solve the sign problem that was introduced by the Fierz transformation.

APPENDIX B: STRONG-COUPLING EXPANSION

The partition function in the vector formulation of the Thirring model with fermionic sources is given by Eq. (74). Here we perform the integration over the vector field v_{μ} and the fermions. After rescaling the vector field according to $\tilde{v}_{\mu} = \sqrt{\lambda}v_{\mu}$ (and afterwards dropping the tilde) the integration over the fermions yields

$$Z[\eta,\bar{\eta}] = (\lambda)^{-(3/2+N_{\rm ir})V} K\left[\frac{\delta}{\delta\eta},\frac{\delta}{\delta\bar{\eta}}\right] \\ \times \int \mathcal{D}v(\det p_x')^{N_{\rm ir}} e^{-\sum_x v_x^2} e^{\sum_x \sqrt{\lambda}\,\bar{\eta}_x \frac{1}{x_x'}\eta_x}.$$
 (B1)

The integral over the vector field factorizes and we can expand in powers of the fermionic bilinear in the exponent,

$$Z[\eta,\bar{\eta}] = (\lambda)^{-(3/2+N_{\rm ir})V} K\left[\frac{\delta}{\delta\eta},\frac{\delta}{\delta\bar{\eta}}\right] \\ \times \int \mathcal{D}v e^{-\sum_{x} (v_{x}^{2}+N_{\rm ir}\ln v_{x}^{2})} e^{\sqrt{\lambda}\sum_{x}\bar{\eta}_{x}\oint_{x}\eta_{x}/v_{x}^{2}} \\ = (\lambda)^{-(\frac{3}{2}+N_{\rm ir})V} K\left[\frac{\delta}{\delta\eta},\frac{\delta}{\delta\bar{\eta}}\right] \prod_{x} \sum_{k=0}^{N_{\rm ir}} I_{k}(\eta_{x},\bar{\eta}_{x}), \qquad (B2)$$

where I_k is the three-dimensional one-site integral

$$I_k(\eta,\bar{\eta}) = \frac{\lambda^k}{(2k)!} \int d^3 v e^{-v^2} (v^2)^{N_{\rm ir}-2k} (\bar{\eta} \not= \eta)^{2k}.$$
(B3)

Here, we already used that only even powers of $p \neq contribute$ to the integrals over v_{μ} . In spherical coordinates $v = r\hat{v}$, the integration over the radial direction is simple and we obtain

$$I_{k}(\eta,\bar{\eta}) = \frac{\lambda^{k}}{(2k)!} \int dr d^{2} \hat{v} e^{-r^{2}} r^{2(N_{\rm ir}+1-k)} (\bar{\eta} \hat{p} \eta)^{2k}$$
$$= \frac{\lambda^{k} \Gamma(\frac{3}{2} + N_{\rm ir} - k)}{2(2k)!} \int_{S^{2}} d^{2} \hat{v} (\bar{\eta} \hat{p} \eta)^{2k}. \tag{B4}$$

The remaining integral can be calculated from the generating function

$$z(j) = \int d^2 \hat{v} e^{\sum j_{\mu} \hat{v}_{\mu}} = 4\pi \frac{\sinh|j|}{|j|}, \qquad j \in \mathbb{R}^3, \qquad (B5)$$

by taking derivatives. This way we find

$$\int d^2 \hat{v} (\bar{\eta} \gamma_\mu \eta \hat{v}_\mu)^{2k} = \left(\bar{\eta} \gamma_\mu \eta \frac{\partial}{\partial j_\mu} \right)^{2k} z(j)|_{j=0}$$
$$= \frac{4\pi}{2k+1} ((\bar{\eta} \gamma_\mu \eta)^2)^k.$$
(B6)

Hence, the integral over the vector field yields

$$I_{k}(\eta,\bar{\eta}) = 2\pi \frac{\lambda^{k} \Gamma(\frac{3}{2} + N_{\rm ir} - k)}{\Gamma(2k+2)} ((\bar{\eta}\gamma_{\mu}\eta)^{2})^{k}, \quad (B7)$$

such that the final expression for the partition function after integration over vector and fermion fields is given by Eq. (76).

In the main body of the paper, just after Eq. (78), we remark that in the infinite-volume limit the partition function only has a trivial dependence on the inverse coupling λ and that local expectation values do not depend on λ at all. In order to show this, we write the *n*th order in the corresponding expansion as a sum over the configurations $\check{k} = \{k_x | x \neq x_0\}$, where k_x is the order of the function $F^{(k_x)}(x)$. We obtain for the partition function

$$Z^{(n)}[\chi_{x_0}, \bar{\chi}_{x_0}] = K^{(n)} \prod_{x} \sum_{k_x} F^{(k_x)}(x)|_{\chi_{x \neq x_0} = 0}$$

= $\sum_{\tilde{k}} K^{(n)} \prod_{x \neq x_0} F^{(k_x)}(x) \sum_{k_{x_0}} F^{(k_{x_0})}(x_0)|_{\chi_{x \neq x_0} = 0},$
(B8)

where a particular point $x = x_0$ was singled out, because we later differentiate the partition function with respect to the source at this point. The operator $K^{(n)}$ contains (n, n)derivatives with respect to $(\chi, \bar{\chi})$ at all lattice points while the function $F^{(k)}(x)$ contains (2k, 2k) fermion sources $(\chi, \bar{\chi})$ at the lattice point *x*. Symbolically, we introduce the operator $D^{(i,i)}$, that collects (i, i) derivatives together with the 2i sums over the lattice points. This allows us to write the partition function as

$$Z^{(n)}[\chi_{x_0}, \bar{\chi}_{x_0}] = \sum_{i=1}^n A^{(i)} B^{(i)}, \tag{B9}$$

with the functions A and B defined as

$$A^{(i)} = \left(D^{(i,i)} \sum_{\tilde{k}} \prod_{x \neq x_0} F^{(k_x)}(x) \right) \Big|_{\chi=0},$$

$$B^{(i)} = D^{(n-i,n-i)} \sum_{k_{x_0}} F^{(k_{x_0})}(x_0).$$
 (B10)

To investigate the volume dependence of the *A* functions we act with *i* derivatives on the *F* functions and afterwards set the sources to zero. Only terms with $2\sum k_x = i$ yield a nonvanishing contribution to the partition function. Furthermore, we need more than one lattice point, because the massless inverse fermion propagator vanishes for x = y (this is true for SLAC fermions on lattices with even *L*). Therefore, the number n_x of lattice points with sources in

the product of the *F* function is $n_x = 2, ..., i/2$. For the first lattice point, we have V - 1 possibilities, for the second lattice point we have V - 2, etc. Thus the volume dependence of $A^{(i)}$ is

$$A^{(i)} \sim \sum_{n_x=2}^{i/2} a_{n_x} \binom{V-1}{n_x} \xrightarrow{V \gg i} \sum_{n_x=2}^{i/2} a_{n_x} \frac{V^{n_x}}{n_x!}.$$
 (B11)

We conclude that the dominant contribution to the partition function in the infinite-volume limit and for a fixed order of the expansion is

$$Z^{(n)}[\chi_{x_0}, \bar{\chi}_{x_0}] = A^{(n)}B^{(n)} \sim C(\lambda) \sum_k F^{(k)}(x_0).$$
(B12)

This leads to the form of the partition function given in Eq. (79).

- [1] W. E. Thirring, Ann. Phys. (N.Y.) 3, 91 (1958).
- [2] G. Parisi, Nucl. Phys. B100, 368 (1975).
- [3] K. Gawedzki and A. Kupiainen, Phys. Rev. Lett. 55, 363 (1985).
- [4] B. Rosenstein, B. Warr, and S. H. Park, Phys. Rep. 205, 59 (1991).
- [5] S. Hands, Phys. Rev. D 51, 5816 (1995).
- [6] T. Itoh, Y. Kim, M. Sugiura, and K. Yamawaki, Prog. Theor. Phys. 93, 417 (1995).
- [7] V. P. Gusynin and P. K. Pyatkovskiy, Phys. Rev. D 94 (2016).
- [8] A. V. Kotikov and S. Teber, Phys. Rev. D 94 (2016).
- [9] G. W. Semenoff, Phys. Rev. Lett. 53, 2449 (1984).
- [10] S. Hands and C. Strouthos, Phys. Rev. B 78 (2008).
- [11] I. F. Herbut, Phys. Rev. B 66, 094504 (2002).
- [12] M. Franz, Z. Tesanovic, and O. Vafek, Phys. Rev. B 66, 054535 (2002).
- [13] M. Gomes, R. S. Mendes, R. F. Ribeiro, and A. J. da Silva, Phys. Rev. D 43, 3516 (1991).
- [14] M. Sugiura, Prog. Theor. Phys. 97, 311 (1997).
- [15] K.-I. Kondo, Nucl. Phys. B450, 251 (1995).
- [16] H. Gies and L. Janssen, Phys. Rev. D 82 (2010).
- [17] L. Janssen and H. Gies, Phys. Rev. D 86 (2012).
- [18] C. Burden and A. N. Burkitt, Eur. Phys. Lett. 3, 545 (1987).
- [19] S. Kim and Y. Kim, arXiv:hep-lat/9605021.
- [20] L. Del Debbio and S. Hands, Phys. Lett. B **373**, 171 (1996).
- [21] L. Del Debbio, S. J. Hands, and J. C. Mehegan, Nucl. Phys. B502, 269 (1997).
- [22] L. Del Debbio and S. J. Hands, Nucl. Phys. B552, 339 (1999).
- [23] S. Hands and B. Lucini, Phys. Lett. B 461, 263 (1999).
- [24] S. Christofi, S. Hands, and C. Strouthos, Phys. Rev. D 75, 101701 (2007).

- [25] S. Chandrasekharan, Phys. Rev. D 82 (2010).
- [26] S. Chandrasekharan and A. Li, *Proc. Sci.*, LATTICE2011 (2011) 058, arXiv:1111.5276.
- [27] S. Chandrasekharan and A. Li, Phys. Rev. Lett. 108, 140404 (2012).
- [28] S. Chandrasekharan and A. Li, Phys. Rev. D 88, 021701 (2013).
- [29] S. Hands, J. High Energy Phys. 11 (2016) 015.
- [30] S. Hands, arXiv:1708.07686.
- [31] S. D. Drell, M. Weinstein, and S. Yankielowicz, Phys. Rev. D 14, 1627 (1976).
- [32] S. D. Drell, M. Weinstein, and S. Yankielowicz, Phys. Rev. D 14, 487 (1976).
- [33] L. H. Karsten and J. Smit, Phys. Lett. B 85, 100 (1979).
- [34] A. Kirchberg, J. D. Länge, and A. Wipf, Ann. Phys. (Amsterdam) 316, 357 (2005).
- [35] G. Bergner, T. Kästner, S. Uhlmann, and A. Wipf, Ann. Phys. (Amsterdam) 323, 946 (2008).
- [36] T. Kästner, G. Bergner, S. Uhlmann, A. Wipf, and C. Wozar, Phys. Rev. D 78, 095001 (2008).
- [37] G. Bergner, J. High Energy Phys. 01 (2010) 024.
- [38] G. Bergner, F. Bruckmann, and J. M. Pawlowski, Phys. Rev. D 79, 115007 (2009).
- [39] R. Flore, D. Körner, A. Wipf, and C. Wozar, J. High Energy Phys. 11 (2012) 159.
- [40] D. Schmidt, B. Wellegehausen, and A. Wipf, Proc. Sci., LATTICE2015 (2015) 050, arXiv:1511.00522.
- [41] D. Schmidt, B. Wellegehausen, and A. Wipf, *Proc. Sci.*, LATTICE2016 (2016) 247, arXiv:1611.00275.
- [42] N. Karthik and R. Narayanan, Phys. Rev. D 93, 045020 (2016).
- [43] D. K. Hong and S. H. Park, Phys. Rev. D 49, 5507 (1994).

- [44] Y. M. Ahn, B. K. Chung, J.-M. Chung, and Q.-H. Park, arXiv:hep-th/9404181.
- [45] Y. M. Ahn and Q. H. Park, J. Korean Phys. Soc. 33, 655 (1998).
- [46] S. J. Hyun, G. H. Lee, and J. H. Yee, Phys. Rev. D 50, 6542 (1994).
- [47] C. Gattringer and C. Marchis, Nucl. Phys. B916, 627 (2017).
- [48] U. Wenger, G. Bergner, and H. Liu, *Proc. Sci.*, LATTICE2016 (2016) 395, arXiv:1612.04291.
- [49] D. Schmidt, B. Wellegehausen, and A. Wipf (to be published).
- [50] G. L. Rossini and F. A. Schaposnik, Phys. Lett. B 338, 465 (1994).