HAWC observations strongly favor pulsar interpretations of the cosmic-ray positron excess

Dan Hooper,^{1,2,3,*} Ilias Cholis,^{4,†} Tim Linden,^{5,‡} and Ke Fang^{6,7,§}

¹Fermi National Accelerator Laboratory, Center for Particle Astrophysics, Batavia, Illinois 60510, USA

²University of Chicago, Department of Astronomy and Astrophysics, Chicago, Illinois 60637, USA

³University of Chicago, Kavli Institute for Cosmological Physics, Chicago, Illinois 60637, USA

⁴Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, Maryland 21218, USA

⁵Ohio State University, Center for Cosmology and AstroParticle Physics (CCAPP),

Columbus, Ohio 43210, USA

⁶University_of Maryland, Department of Astronomy, College Park, Maryland 20742, USA

¹Joint Space-Science Institute, College Park, Maryland 20742, USA

(Received 12 July 2017; published 17 November 2017)

Recent measurements of the Geminga and B0656 + 14 pulsars by the gamma-ray telescope HAWC (along with earlier measurements by Milagro) indicate that these objects generate significant fluxes of very high-energy electrons. In this paper, we use the very high-energy gamma-ray intensity and spectrum of these pulsars to calculate and constrain their expected contributions to the local cosmic-ray positron spectrum. Among models that are capable of reproducing the observed characteristics of the gamma-ray emission, we find that pulsars invariably produce a flux of high-energy positrons that is similar in spectrum and magnitude to the positron fraction measured by PAMELA and AMS-02. In light of this result, we conclude that it is very likely that pulsars provide the dominant contribution to the long perplexing cosmic-ray positron excess.

DOI: 10.1103/PhysRevD.96.103013

I. INTRODUCTION

Measurements of the cosmic-ray positron fraction by the PAMELA [1] and AMS-02 [2] experiments (as well as HEAT [3], AMS-01 [4], and Fermi [5]) have identified an excess of high-energy positrons relative to the standard predictions for secondary production in the interstellar medium (ISM). This result indicates that significant quantities of ~0.01–1 TeV positrons are produced within the local volume (the surrounding ~kpc) of the Milky Way. The origin of these particles has been debated, and possible sources include nearby and young pulsars [6–12], as well as annihilating dark matter particles [12–25]. This excess could also plausibly arise in nearby supernova remnants, which may be able to produce and accelerate secondary positrons before they escape into the ISM [26–32].

From the catalog of known pulsars, Geminga (PSR J0633 + 1746) and B0656 + 14 (PSR J0659 + 1414, thought to be associated with the Monogem supernova remnant) are particularly attractive candidates to account for the observed positron excess. These pulsars are each

relatively young (370,000 and 110,000 yr, respectively) and are located within a few hundred parsecs of the solar system $(250^{+230}_{-80} \text{ and } 280^{+30}_{-30} \text{ pc}, \text{ respectively [33]})$. Much of what we know about these and other pulsars is based on gamma-ray observations. Geminga, in particular, is an extremely bright source of GeV-scale emission, thought to originate from the pulsar itself [34-37]. In contrast, observations of Geminga at very high energies reveal a significantly extended morphology. In particular, the Milagro Collaboration has reported the detection of gamma-ray emission at ~35 TeV with a flux of $(3.77\pm1.07)\times10^{-16}~TeV^{-1}~cm^{-2}\,s^{-1}$ and an extended profile with a full-width at half maximum of $2.6^{+0.7}_{-0.9}$ deg [38]. Very recently, the HAWC Collaboration has presented their measurements of the Geminga pulsar, confirming its angular extension (~2° radius) and reporting a flux of $(4.87 \pm 0.68 \pm 2.4) \times 10^{-14} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ at 7 TeV, with a local spectral index of $-2.23 \pm 0.08 \pm 0.2$ [39] (see also Refs. [40-42]). HAWC has also reported the detection of very high-energy emission from B0656 + 14, with a similar degree of spatial extension and with a flux and spectral index at 7 TeV of $(2.30\pm$ 0.73 ± 1.2 × 10⁻¹⁴ TeV⁻¹ cm⁻²s⁻¹ and -2.03 ± 0.14 ± 0.2, respectively [39].

These observations by HAWC and Milagro allow us to conclude that these pulsars, in fact, deposit a significant fraction of their total spin-down power into high-energy leptons. Furthermore, we will show in this paper that the flux of leptons required to explain these observations is

dhooper@fnal.gov

ORCID: http://orcid.org/0000-0001-8837-4127 [†]icholis1@jhu.edu

ORCID: http://orcid.org/0000-0002-3805-6478 [‡]linden.70@osu.edu

ORCID: http://orcid.org/0000-0001-9888-0971 [§]kefang@umd.edu

ORCID: http://orcid.org/0000-0002-5387-8138

HOOPER, CHOLIS, LINDEN, and FANG

roughly equal to that required for the Geminga and B0656 + 14 pulsars to produce an order 1 fraction of the positron excess observed by PAMELA and AMS-02. The spectrum, morphology, and intensity of the gamma-ray emission measured by HAWC and Milagro leave us with little choice but to conclude that nearby pulsars are likely to be the dominant source of the observed cosmic-ray positrons.

II. INVERSE COMPTON SCATTERING OF VERY HIGH-ENERGY ELECTRONS AND POSITRONS NEAR PULSARS

The gamma-ray emission observed from Geminga and B0656 + 14 by HAWC and Milagro is almost certainly generated through the inverse Compton scattering of very high-energy leptons. The angular extension of this signal rules out other scenarios, with the possible exception of pion production. A pion production origin, however, would require an unrealistically large quantity ($\gtrsim 10^{46}$ erg) of $\mathcal{O}(10^2)$ TeV protons to be confined to the region surrounding Geminga for $\gtrsim 10^5$ yr. Such a scenario can also be constrained to some degree by the lack of TeV neutrinos detected by the IceCube experiment [43].

To study the diffusion and energy losses of electrons and positrons produced in nearby pulsars, we utilize the standard propagation equation,

$$\frac{\partial}{\partial t} \frac{dn_e}{dE_e}(E_e, r, t)
= \vec{\nabla} \cdot \left[D(E_e) \vec{\nabla} \frac{dn_e}{dE_e}(E_e, r, t) - \vec{v}_c \frac{dn_e}{dE_e}(E_e, r, t) \right]
+ \frac{\partial}{\partial E_e} \left[\frac{dE_e}{dt}(r) \frac{dn_e}{dE_e}(E_e, r, t) \right] + \delta(r) Q(E_e, t), \quad (2.1)$$

where dn_e/dE_e is the differential number density of electrons/positrons at a distance *r* from the pulsar, *D* is the diffusion coefficient, \vec{v}_c is the convection velocity (which we take to be directed outward from the pulsar), and the source term *Q* describes the spectrum and time profile of electrons/positrons injected into the ISM. Energy losses are dominated by a combination of inverse Compton and synchrotron losses and are given by [44]

$$-\frac{dE_e}{dt}(r) = \sum_i \frac{4}{3} \sigma_T \rho_i(r) S_i(E_e) \left(\frac{E_e}{m_e}\right)^2 + \frac{4}{3} \sigma_T \rho_{\text{mag}}(r) \left(\frac{E_e}{m_e}\right)^2, \qquad (2.2)$$

$$\equiv b(E_e, r) \left(\frac{E_e}{\text{GeV}}\right)^2, \qquad (2.3)$$

where σ_T is the Thomson cross section and

$$b(r) \approx 1.02 \times 10^{-16} \text{ GeV/s}$$
$$\times \left[\sum_{i} \frac{\rho_i(r)}{\text{eV/cm}^3} S_i(E_e) + 0.224 \left(\frac{B(r)}{3 \,\mu\text{G}} \right)^2 \right]. \quad (2.4)$$

The sum in this expression is carried out over the various components of the radiation backgrounds, consisting of the cosmic microwave background (CMB), infrared emission (IR), starlight (star), and ultraviolet emission (UV). Throughout our analysis, we adopt the following parameters: $\rho_{\rm CMB} = 0.260 \text{ eV/cm}^3$, $\rho_{\rm IR} = 0.60 \text{ eV/cm}^3$, $\rho_{\rm star} = 0.60 \text{ eV/cm}^3$, $\rho_{\rm UV} = 0.10 \text{ eV/cm}^3$, and $\rho_{\rm mag} = 0.224 \text{ eV/cm}^3$ (corresponding to $B = 3 \mu$ G) and $T_{\rm CMB} = 2.7 \text{ K}$, $T_{\rm IR} = 20 \text{ K}$, $T_{\rm star} = 5000 \text{ K}$, and $T_{\rm UV} = 20,000 \text{ K}$. For low to moderate electron energies, these parameters correspond to a value of $b \approx 1.8 \times 10^{-16} \text{ GeV/s}$. At very high energies ($E_e \gtrsim m_e^2/2T$), however, inverse Compton scattering is further suppressed by the following factor [45]:

$$S_i(E_e) \approx \frac{45m_e^2/64\pi^2 T_i^2}{(45m_e^2/64\pi^2 T_i^2) + (E_e^2/m_e^2)}.$$
 (2.5)

To solve Eq. (2.2), we calculate the distribution of the electrons and positrons that were emitted a time t ago and then sum the contributions produced over different periods of time. Considering an injected spectrum of the form $Q(E_e, t) = \delta(t)Q_0E^{-\alpha}\exp(-E_e/E_c)$, the solution to Eq. (2.2) (neglecting convection) is given by

$$\frac{dn_{e}}{dE_{e}}(E_{e}, r, t) = \frac{Q_{0}E_{0}^{2-\alpha}}{8\pi^{3/2}E_{e}^{2}L_{dif}^{3}(E_{e}, t)} \times \exp\left[\frac{-E_{0}}{E_{c}}\right] \exp\left[\frac{-r^{2}}{4L_{dif}^{2}(E_{e}, t)}\right], \quad (2.6)$$

where E_0 is the initial energy of an electron that has an energy of E_e after a time *t*. At low to moderate energies, for which $S_i(E_e) \approx 1$, this quantity is given by $E_0 \approx E_e/(1 - E_e bt)$. Including the effects of Klein-Nishina suppression, E_0 is determined by solving $\int_{E_0}^{E_e} dE/b(E)E^2 = t$. The diffusion length scale is given by

$$L_{\rm dif}(E_e, t) \equiv \left[\int_{E_0}^{E_e} \frac{D(E')}{-dE_e/dt(E')} dE' \right]^{1/2}.$$
 (2.7)

Throughout this study, we adopt a parametrization of $D(E_e) = D_0 E_e^{\delta}$ for the diffusion coefficient.

To account for the time profile of the electrons and positrons injected from a given pulsar, we adopt a function proportional to the spin-down power (the rate at which the pulsar loses rotational kinetic energy through magnetic dipole braking) [46], HAWC OBSERVATIONS STRONGLY FAVOR PULSAR ...

$$\dot{E} = -\frac{8\pi^4 B^2 R^6}{3c^3 P(t)^4}$$

\$\approx 1.0 \times 10^{35} erg/s
\$\times \left(\frac{B}{1.6 \times 10^{12} \text{ G}} \right)^2 \left(\frac{R}{15 \text{ km}} \right)^6 \left(\frac{0.23 \text{s}}{P(t)} \right)^4, (2.8)\$

where B is the strength of the magnetic field at the surface of the neutron star, R is the radius of the neutron star, and the rotational period evolves as follows,

$$P(t) = P_0 \left(1 + \frac{t}{\tau} \right)^{1/2},$$
 (2.9)

where P_0 is the initial period and τ is the spin-down time scale:

$$\tau = \frac{3c^3 I P_0^2}{4\pi^2 B^2 R^6}$$

$$\approx 9.1 \times 10^3 \text{ yr} \left(\frac{1.6 \times 10^{12} \text{ G}}{B}\right)^2 \left(\frac{M}{1.4 M_{\odot}}\right) \left(\frac{15 \text{ km}}{R}\right)^4$$

$$\times \left(\frac{P_0}{0.040 \text{ sec}}\right)^2. \tag{2.10}$$

In Eqs. (2.8) and (2.10), we adopted benchmark values for the neutron star's magnetic field, radius, and mass, chosen to match Geminga's observed period and its rate of change.

At energies within the range measured by MILAGRO and HAWC, inverse Compton scattering yields photons with energies not very far below that of the incident electrons and positrons, $E_{\gamma} \sim E_e$. Adopting this approximation, the angular profile of gamma rays generated through inverse Compton scattering is given by

$$\begin{split} \Phi_{\gamma}(E_{\gamma} = E_{e}, \psi) \propto \int \dot{E} dt \int_{\log} \frac{dn_{e}}{dE_{e}} (E_{e}, r, t) \rho_{\rm rad}(r) dl \\ \propto \int \frac{\tau^{2}}{(t+\tau)^{2}} dt \int_{\log} e^{-r^{2}/4L_{\rm dif}^{2}(E_{e}, t)} \rho_{\rm rad}(r) dl, \end{split}$$

$$(2.11)$$

where ψ is the angle observed away from the pulsar, and $r^2 = l^2 + d^2 - 2ld \cos \psi$, where *d* is the distance between the pulsar and the observer. If we adopt a uniform distribution of radiation in the vicinity of the pulsar, this reduces to a profile of the form $\Phi_{\gamma}(\psi) \propto \exp[-d^2 \sin^2 \psi / 4L_{\text{dif}}^2(E_e, t)]$. Observations of Geminga by both Miligro and HAWC indicate that the very high-energy gamma-ray emission from this source is extended over a region of a few degrees across the sky. This in turn requires a diffusion length given by $L_{\text{dif}}(E_e) \approx (250 \text{ pc}) \sin(0.5 \times 2.6^\circ) / 2(\ln(2))^{1/2} \approx 2.6 \text{ pc}$. In contrast, adopting parameters appropriate for the ISM $(D_0 \approx 2 \times 10^{28} \text{ cm}^2/\text{s}, \delta \approx 0.4, b = 1.8 \times 10^{-16} \text{ GeV/s})$, we find



FIG. 1. The angular distribution of the flux of inverse Compton emission (per solid angle) from 35 TeV electrons (corresponding to photons in the approximate energy range measured by Milagro and HAWC) from the Geminga pulsar. Here, we have adopted diffusion and energy loss parameters which correspond to the conditions found in the ISM, $D_0 = 2 \times 10^{28} \text{ cm}^2/\text{s}$, $\delta = 0.4$, and $b = 1.8 \times 10^{-16} \text{ GeV/s}$, and spectral parameters as given by $\alpha = 1.5$, $E_c = 100 \text{ TeV}$. The solid black line represents the angular profile predicted assuming an isotropic radiation distribution, whereas the dashed blue line (visible in the upper left corner) also includes a contribution from a population of radiation which is distributed according to an r^{-2} profile, normalized to the total spin-down power of Geminga. In either case, the predicted profile is dramatically broader than the ~2° extension reported by both Milagro and HAWC.

$$L_{\rm dif}(E_e, t) \simeq 200 \ {\rm pc} \left(\frac{35 \ {\rm TeV}}{E_e}\right)^{0.3} (1 - (1 - E_e b t)^{0.6})^{1/2}.$$

(2.12)

Assuming conditions for cosmic-ray transport that are similar to those found in the ISM, this calculation shows that we should have expected the inverse Compton emission observed at very high energies to be extended over a scale of ~60°, dramatically more than the ~2° extension reported by both Milagro and HAWC (see Fig. 1).

To resolve this puzzle, one might be tempted to consider the possibility that the pulsar is surrounded by a dense radiation field, which intensifies the resulting inverse Compton emission from the surrounding parsecs. The problem with this scenario, however, is that there is not nearly enough power available to generate the required density of radiation. More quantitatively, for an r^{-2} profile of radiation to exceed the energy density of the CMB at a distance of 1 pc from the pulsar would require an amount of energy equivalent to more than ten times the total spindown power of Geminga. Adopting an extreme benchmark in which 100% of Geminga's energy budget is transferred into radiation, the profile of inverse Compton emission is altered only very modestly, by less than 10% at $\psi = 1^{\circ}$ (see Fig. 1). Based on these considerations, it does not appear that local concentrations of radiation play a significant role in explaining the angular extent of the gamma-ray emission observed from Geminga or B0656 + 14.

A more likely solution is that the conditions that dictate cosmic-ray diffusion around Geminga and B0656 + 14 are very different from those found elsewhere in the ISM, leading energetic leptons to escape from the surrounding regions much more slowly. In order to accommodate the observed extension in this way, we require that these particles only diffuse a distance of a few parsecs before losing most of their energy. For an energy loss time of 5000 yr (corresponding to $E_e = 35$ TeV and $b = 1.8 \times 10^{-16}$ GeV/s), this requires a diffusion coefficient of $D \sim 10^{27}$ cm²/s (see Fig. 2). Although this is significantly smaller than that found in the bulk of the ISM, we note that it is similar to that predicted for standard Bohmian diffusion, $D_{\text{Bohm}} = r_L c/3 \approx 1.2 \times 10^{27}$ cm²/s $\times (E_e/35 \text{ TeV})(\mu G/B)$.

If pulsars such as Geminga are typically surrounded by a region with inefficient diffusion ($D \ll D_{\rm ISM}$), the volume of such regions must be fairly small to avoid conflicting with secondary-to-primary ratios in the cosmic-ray spectrum as measured at Earth. In particular, if such regions have a typical radius of $r_{\rm region}$, then such regions will occupy roughly the following fraction of the volume of the Milky Way's disk,



FIG. 2. The angular distribution of the flux of inverse Compton emission (per solid angle) from 35 TeV electrons from the Geminga pulsar, for different choices of the diffusion parameters, D_0 , δ [such that $D = D_0(E_e/\text{GeV})^{\delta}$]. We have adopted an energy loss parameter of $b = 1.8 \times 10^{-16}$ GeV/s and spectral parameters of $\alpha = 1.5$ and $E_c = 100$ TeV. To accommodate the ~2° extension reported by Milagro and HAWC, Geminga must be enclosed within an environment with a diffusion coefficient that is ~10³ times smaller than that corresponding to the typical conditions of the ISM.

$$f \sim \frac{N_{\text{region}} \times \frac{2n}{3} r_{\text{region}}^{2}}{\pi R_{\text{MW}}^{2} \times 2z_{\text{MW}}}$$
$$\sim 0.03 \times \left(\frac{r_{\text{region}}}{50 \text{ pc}}\right)^{3} \left(\frac{\dot{N}_{\text{SN}}}{0.03 \text{ yr}^{-1}}\right) \left(\frac{\tau_{\text{region}}}{10^{6} \text{ yr}}\right)$$
$$\times \left(\frac{20 \text{ kpc}}{R_{\text{MW}}}\right)^{2} \left(\frac{200 \text{ pc}}{z_{\text{MW}}}\right), \qquad (2.13)$$

where $\dot{N}_{\rm SN}$ is the rate at which new pulsars appear in the Galaxy, $\tau_{\rm region}$ is the length of time that such regions persist, and $N_{\rm region} = \dot{N}_{\rm SN} \times \tau_{\rm region}$ is the number of such regions present at a given time. The quantities $R_{\rm MW}$ and $z_{\rm MW}$ denote the radius and half-width of the Galaxy's cylindrical disk. Combined with Milagro and HAWC observations of Geminga and B0656 + 14, these considerations suggest 5 pc $\lesssim r_{\rm region} \lesssim 50$ pc, for which there will be little impact on the observed secondary-to-primary ratios (other than the positron fraction).

In Fig. 3, we plot the gamma-ray spectrum from the region surrounding Geminga, compared to the measurements by HAWC and Milagro (at 7 and 35 TeV, respectively). In performing this calculation, we utilized the full differential cross section for inverse Compton scattering [44]. In the left (right) frame, we have adopted a spectrum of injected electrons with an index of $\alpha = 1.5$ (1.9) and in each case selected a value of E_c that provides the best fit to the combination of these two measurements. In this part of our calculation, we have parametrized the transport of the cosmic rays out of the region surrounding Geminga with an escape time, t_{esc} , which is intended to account for phenomena such as convective winds or energy-independent diffusion [7]. In particular, we show results for two values of the escape time ($t_{\rm esc} = 1.8 \times 10^4$ and $t_{\rm esc} = 1.8 \times 10^5$ yr), which if interpreted in terms of convection correspond to $v_c = 554 \text{ km/s} \times (r_{\text{region}}/10 \text{ pc})$ and $v_c = 55.4 \text{ km/s} \times (r_{\text{region}}/10 \text{ pc})$ $(r_{\rm region}/10 \text{ pc})$, respectively. In these four cases, the best fits were found for $E_c = 44$ TeV ($\alpha = 1.5$, $t_{esc} = 1.8 \times$ 10⁵ yr), $E_c = 35$ TeV ($\alpha = 1.5, t_{\rm esc} = 1.8 \times 10^4$ yr), $E_c =$ 67 TeV ($\alpha = 1.9$, $t_{esc} = 1.8 \times 10^5$ yr), and $E_c = 49$ TeV $(\alpha = 1.9, t_{\rm esc} = 1.8 \times 10^4 \text{ yr}).$

In addition to their flux measurement, the HAWC Collaboration has also reported a value of $-2.23 \pm 0.08 \pm 0.2$ (statistical and systematic uncertainties, respectively) for Geminga's spectral slope at 7 TeV. Among the models shown in Fig 3, those with $t_{\rm esc} = 1.8 \times 10^5$ yr predict spectral slopes at 7 TeV of -2.47 ($\alpha = 1.5$) or -2.59 ($\alpha = 1.9$). Such values are in some tension with those reported by HAWC. In contrast, for those models with a shorter escape time, $t_{\rm esc} = 1.8 \times 10^4$ yr, we instead predict a spectral slope of -2.23 ($\alpha = 1.5$) or -2.32 ($\alpha = 1.9$), in excellent agreement with HAWC's measurement. This favors scenarios in which convection or energy-independent diffusion plays a very important role in the transport of high-energy leptons, especially among those



FIG. 3. The gamma-ray spectrum from the region surrounding Geminga, compared to measurements by HAWC and Milagro (shown at 7 and 35 TeV, respectively). In the left (right) frame, we adopt a spectrum of injected electrons with an index of $\alpha = 1.5$ (1.9) and have in each case selected a value of E_c that provides the best fit to the combination of these two measurements. The blue-dashed (solid green) curves correspond to a case in which the escape time from the region surrounding Geminga is $1.8 \times 10^5 (1.8 \times 10^4)$ yr. As discussed in the text, the case of $t_{esc} \approx 1.8 \times 10^5$ yr is disfavored by the value of the spectral index reported by the HAWC Collaboration.

with energies below ~10 TeV. We lastly note that, among this range of models, between 7.2% and 29% of Geminga's total current spin-down power is being deposited into electron-positron pairs with $E_e > 10$ GeV.

III. IMPLICATIONS FOR THE COSMIC-RAY POSITRON EXCESS

Although the angular extent of the emission observed from Geminga and B0656 + 14 by HAWC and Milagro indicates that very high-energy ($E_e \gtrsim 10$ TeV) leptons are effectively trapped within a few parsecs of these sources, the same fate need not be experienced by lower-energy electrons and positrons. In particular, even a modest degree of convection (i.e. the streaming of particles away from the source at a constant velocity) can remove sub-TeV leptons from the region before they lose a substantial fraction of their energy, while higher-energy leptons lose the vast majority of their energy to inverse Compton scattering before escaping the same region [7].

To address this more quantitatively, we can compare the time scale for energy losses, $\tau_{\rm loss} = 1/E_e b$, to that for the escape from the region, $\tau_{\rm escape} = r_{\rm region}/v_c$. For convective winds with a velocity of v_c , particles with $E_e \ll v_c/br_{\rm region}$ are left largely unaffected, while those with $E_e \gg v_c/br_{\rm region}$ lose the majority of their energy before escaping. In the left frame of Fig. 4, we plot the fraction of energy that an electron loses before escaping a region of radius 10 pc for several values of v_c . It is expected that future observations by the Cherenkov Telescope Array (CTA) will provide an important test of this transition by measuring the intensity and angular extent of the ~0.1–10 TeV emission from Geminga and B0656 + 14.



FIG. 4. Left: The fraction of energy that is lost by an electron over the time that is required to move via convection a distance of 10 pc. Right: The mean fraction of energy that is lost by an electron over the time required to move via either convection ($v_c = 100 \text{ km/s}$) or energy-independent diffusion ($D_0 = 10^{26} \text{ cm}^2/\text{s}$) a distance of 10 pc. For comparison, in each frame, we show the (gamma-ray) energies at which HAWC and Milagro have reported fluxes.



FIG. 5. The cosmic-ray positron spectrum as measured by AMS-02, compared to the predictions from standard secondary production in the ISM (solid black) and including a contribution from the Geminga pulsar. In each case, we have chosen the normalization and spectral shape to match that of the very high-energy gamma-ray emission measured by HAWC and Milagro (see Fig. 3). In the left (right) frame, we adopt a spectrum of injected electrons with an index of $\alpha = 1.5$ (1.9) and have in each case selected a value of E_c that provides the best fit to the combination of the HAWC and Milagro measurements. The solid green curves correspond to a case with $t_{esc} = 1.8 \times 10^4$ yr. For comparison, we also show as blue-dashed curves the result with $t_{esc} = 1.8 \times 10^5$ yr, although this case is disfavored by the spectral index reported by the HAWC Collaboration.

Although one can describe the escape of sub-TeV from the regions surrounding Geminga and B0656 + 14 in terms of convection, we emphasize that other physical means are also possible. For example, if we simply introduce a diffusion coefficient with no energy dependence ($\delta = 0$), we effectively mimic the effects of convection (see the right frame of Fig. 4). We remain agnostic as to the physical means by which sub-TeV leptons escape the regions surrounding these sources and encourage the reader to interpret convection within this model as a proxy for whatever mechanism expels particles from the low-diffusion region, as quantified in the previous paragraph.

In Fig. 5, we plot the cosmic-ray positron fraction as measured by AMS-02, compared to the predictions from the Geminga pulsar, using the same choices of parameters as adopted in Fig. 3. In each frame, the solid black curve denotes the contribution from standard secondary production in the ISM, while the other curves include both this contribution and that from Geminga. We remind the reader that those models with $t_{\rm esc} = 1.8 \times 10^5$ yr (dotted blue curves) do not lead to a spectral index compatible with the measurement of HAWC and thus should be viewed as a poor fit to the data. We have not presented the contribution to the cosmic-ray electron spectrum from Geminga, as we do not expect this individual source to dominate the spectrum in the energy range considered here. In our estimation, it remains likely that sources such as supernova remnants significantly contribute to the observed spectrum of cosmic-ray electrons.

The positron fraction presented in Fig. 5 includes a distinctive feature at 400–500 GeV, which is a result of energy losses. More specifically, a positron with an infinite energy will be reduced over a time *t* to an energy of $E_e \simeq (bt)^{-1}$, which for t = 370,000 yr (the age of

Geminga) yields a final energy of \sim 500 GeV. Any positrons from Geminga above this energy were injected at later times and thus have not cooled to the same extent.

The main lesson from the results shown in Fig. 5 is that when the spectral shape and overall normalization of Geminga are fixed to reproduce the results of HAWC and Milagro, this pulsar is found to generate a nonnegligible portion of the observed positron fraction. That being said, the overall size of this contribution to the cosmic-ray positron flux can vary by a factor of order unity depending on the precise values of the escape time, t_{esc} , and spin-down time scale, τ [see Eq. (2.10)] that are adopted. The impact of the convection velocity is clearly evident in Fig. 5. The resulting positron flux scales approximately as τ^{-1} (we have adopted a value of $\tau = 9.1 \times 10^3$ yr). Furthermore, the time profile of a pulsar's emission could plausibly depart to some extent from that predicted from standard magnetic dipole braking [46], potentially altering the normalization of the positron flux predicted here, as well as the inferred age of the pulsar.

Finally, in Fig. 6, we plot the contributions to the positron fraction from the Geminga and B0656 + 14 pulsars, as well as the average contribution from those pulsars located more than 500 pc away from the Solar System. For each source, we have adopted $\alpha = 1.9$, $E_c = 49$ TeV, and $t_{\rm esc} = 1.8 \times 10^4$ yr and normalized their contributions with $\tau = 4.3 \times 10^3$ yr and adopted a total birth rate of two pulsars per century in the Milky Way.¹ For other details regarding the calculation of the contribution from distant pulsars, we direct the reader to Ref. [6]. In

¹We produce nearly identical results if we instead adopt our default value for $\tau \simeq 9.1 \times 10^3$ and a somewhat lower value for the escape time, $t_{\rm esc} = 8.4 \times 10^3$ yr.



FIG. 6. As in Fig. 5, but showing contributions from Geminga, B0656 + 14, and all pulsars more than 0.5 kpc from the Solar System. For each source, we adopted $\alpha = 1.9$, $E_c = 49$ TeV, and $t_{\rm esc} = 1.8 \times 10^4$ yr and normalized their contributions with $\tau = 4.3 \times 10^3$ yr, adopting a total birth rate of two pulsars per century in the Milky Way. While we expect many of these parameters to vary from pulsar to pulsar, making a detailed prediction of this kind difficult and possibly unreliable, this calculation provides significant support for the conclusion that a sizable fraction of the observed positron excess originates from pulsars.

reality, we expect many of these parameters to vary from pulsar to pulsar, making a detailed prediction of this kind difficult and possibly unreliable. That being said, this exercise clearly demonstrates that in light of the measurements by HAWC and Milagro, it appears very likely that a sizable fraction of the observed positron excess originates from this class of sources. In addition, we note that it was recently shown that the stochastic acceleration of cosmicray secondaries in supernova remnants is also likely to contribute to the local positron flux [32].

IV. SUMMARY AND CONCLUSIONS

In this paper, we have made use of measurements by the very high-energy gamma-ray telescopes HAWC and Milagro to better understand and constrain the injection of high-energy electrons and positrons from the nearby pulsars Geminga and B0656 + 14. The angular extension of

the \gtrsim TeV gamma-ray emission observed from these pulsars indicates that very high-energy leptons are effectively trapped within the surrounding several parsecs around these sources. Furthermore, their very high-energy gamma-ray spectra indicate that lower-energy leptons are able to escape more easily, suggesting the presence of features such as strong convective winds or energy-independent diffusion.

In models that are able to reproduce the characteristics of the gamma-ray emission reported by HAWC and Milagro, these pulsars invariably provide a significant contribution to the local cosmic-ray positron spectrum, and thus to the measured positron fraction. Although it is not yet possible to precisely predict the normalization of the positron flux from these sources, these results show that Geminga and B0656 + 14 are expected to generate a significant fraction of the observed high-energy positron flux. If we make the entirely reasonable assumption that other pulsars in the Milky Way behave in a fashion similar to Geminga and B0656 + 14, we find that it is very likely that pulsars are responsible for much, if not the entirety, of the reported positron excess.

An important test of this conclusion will come from future Imaging Atmospheric Cherenkov Telescopes (IACTs), such as the CTA. Although existing IACTs have not yet reported any significant detection of TeV-scale emission from Geminga or B0656 + 14 [47,48], next generation telescopes will be far better suited to detect emission that is extended over the angular scales reported by Milagro and HAWC. Such a measurement is expected to be able to confirm the basic features of the model presented here and enable us to produce a more detailed determination of the spectrum of electrons and positrons that are injected from this class of sources.

ACKNOWLEDGMENTS

D. H. is supported by the U.S. Department of Energy under Contract No. DE-FG02-13ER41958. Fermilab is operated by Fermi Research Alliance, LLC, under Contract No. DE- AC02-07CH11359 with the U.S. Department of Energy. I. C. acknowledges support from NASA Grant No. NNX15AB18G and from the Simons Foundation. T. L. acknowledges support from NSF Grant No. PHY-1404311.

- O. Adriani *et al.* (PAMELA Collaboration), PAMELA Results on the Cosmic-Ray Antiproton Flux from 60 MeV to 180 GeV in Kinetic Energy, Phys. Rev. Lett. 105, 121101 (2010).
- [2] M. Aguilar *et al.* (AMS Collaboration), First Result from the Alpha Magnetic Spectrometer on the International Space

Station: Precision Measurement of the Positron Fraction in Primary Cosmic Rays of 0.5350 GeV, Phys. Rev. Lett. **110**, 141102 (2013).

[3] S. W. Barwick *et al.* (HEAT Collaboration), Measurements of the cosmic ray positron fraction from 1-GeV to 50-GeV, Astrophys. J. **482**, L191 (1997).

- [4] M. Aguilar *et al.* (AMS 01 Collaboration), Cosmic-ray positron fraction measurement from 1 to 30-GeV with AMS-01, Phys. Lett. B 646, 145 (2007).
- [5] M. Ackermann *et al.* (Fermi-LAT Collaboration), Measurement of Separate Cosmic-Ray Electron and Positron Spectra with the Fermi Large Area Telescope, Phys. Rev. Lett. **108**, 011103 (2012).
- [6] D. Hooper, P. Blasi, and P. D. Serpico, Pulsars as the sources of high energy cosmic ray positrons, J. Cosmol. Astropart. Phys. 01 (2009) 025.
- [7] H. Yuksel, M. D. Kistler, and T. Stanev, TeV Gamma Rays from Geminga and the Origin of the GeV Positron Excess, Phys. Rev. Lett. **103**, 051101 (2009).
- [8] S. Profumo, Dissecting cosmic-ray electron-positron data with Occam's Razor: the role of known Pulsars, Central Eur. J. Phys. 10, 1 (2011).
- [9] D. Malyshev, I. Cholis, and J. Gelfand, Pulsars versus dark matter interpretation of ATIC/PAMELA, Phys. Rev. D 80, 063005 (2009).
- [10] D. Grasso *et al.* (Fermi-LAT Collaboration), On possible interpretations of the high energy electron-positron spectrum measured by the Fermi Large Area Telescope, Astropart. Phys. **32**, 140 (2009).
- [11] T. Linden and S. Profumo, Probing the pulsar origin of the anomalous positron fraction with AMS-02 and Atmospheric Cherenkov Telescopes, Astrophys. J. 772, 18 (2013).
- [12] I. Cholis and D. Hooper, Dark matter and pulsar origins of the rising cosmic ray positron fraction in light of new data from AMS, Phys. Rev. D 88, 023013 (2013).
- [13] L. Bergstrom, T. Bringmann, and J. Edsjo, New Positron Spectral Features from Supersymmetric Dark Matter—A Way to Explain the PAMELA Data?, Phys. Rev. D 78, 103520 (2008).
- [14] M. Cirelli and A. Strumia, Minimal dark matter predictions and the PAMELA positron excess, *Proc. Sci.*, IDM2008 (2008) 089.
- [15] I. Cholis, L. Goodenough, D. Hooper, M. Simet, and N. Weiner, High energy positrons from annihilating dark matter, Phys. Rev. D 80, 123511 (2009).
- [16] M. Cirelli, M. Kadastik, M. Raidal, and A. Strumia, Modelindependent implications of the e + -, anti-proton cosmic ray spectra on properties of Dark Matter, Nucl. Phys. B813, 1 (2009).
- [17] A. E. Nelson and C. Spitzer, Slightly non-minimal dark matter in PAMELA and ATIC, J. High Energy Phys. 10 (2010) 066.
- [18] N. Arkani-Hamed, D. P. Finkbeiner, T. R. Slatyer, and N. Weiner, A theory of dark matter, Phys. Rev. D 79, 015014 (2009).
- [19] I. Cholis, D. P. Finkbeiner, L. Goodenough, and N. Weiner, The PAMELA positron excess from annihilations into a light boson, J. Cosmol. Astropart. Phys. 12 (2009) 007.
- [20] I. Cholis, G. Dobler, D. P. Finkbeiner, L. Goodenough, and N. Weiner, The case for a 700 + GeV WIMP: Cosmic ray spectra from ATIC and PAMELA, Phys. Rev. D 80, 123518 (2009).
- [21] R. Harnik and G. D. Kribs, An effective theory of Dirac dark matter, Phys. Rev. D 79, 095007 (2009).
- [22] P. J. Fox and E. Poppitz, Leptophilic dark matter, Phys. Rev. D 79, 083528 (2009).

- [23] M. Pospelov and A. Ritz, Astrophysical signatures of secluded dark matter, Phys. Lett. B 671, 391 (2009).
- [24] J. D. March-Russell and S. M. West, WIMPonium and boost factors for indirect dark matter detection, Phys. Lett. B 676, 133 (2009).
- [25] S. Chang and L. Goodenough, Charge asymmetric cosmic ray signals from dark natter decay, Phys. Rev. D 84, 023524 (2011).
- [26] P. Blasi, The Origin of the Positron Excess in Cosmic Rays, Phys. Rev. Lett. 103, 051104 (2009).
- [27] P. Mertsch and S. Sarkar, Testing Astrophysical Models for the PAMELA Positron Excess with Cosmic Ray Nuclei, Phys. Rev. Lett. 103, 081104 (2009).
- [28] M. Ahlers, P. Mertsch, and S. Sarkar, On cosmic ray acceleration in supernova remnants and the FERMI/ PAMELA data, Phys. Rev. D 80, 123017 (2009).
- [29] I. Cholis and D. Hooper, Constraining the origin of the rising cosmic ray positron fraction with the boron-to-carbon ratio, Phys. Rev. D 89, 043013 (2014).
- [30] M. Kachelriess, S. Ostapchenko, and R. Tomas, Antimatter production in supernova remnants, Astrophys. J. 733, 119 (2011).
- [31] M. Kachelrie and S. Ostapchenko, *B/C* ratio and the PA-MELA positron excess, Phys. Rev. D 87, 047301 (2013).
- [32] I. Cholis, D. Hooper, and T. Linden, Evidence for the stochastic acceleration of secondary antiprotons by supernova remnants, Phys. Rev. D 95, 123007 (2017).
- [33] J. P. W. Verbiest, J. M. Weisberg, A. A. Chael, K. J. Lee, and D. R. Lorimer, On pulsar distance measurements and their uncertainties, Astrophys. J. 755, 39 (2012).
- [34] F. Acero *et al.*, Constraints on the galactic population of TeV pulsar wind nebulae using Fermi Large Area Telescope observations, Astrophys. J. **773**, 77 (2013).
- [35] A. A. Abdo *et al.*, Fermi-LAT observations of the Geminga pulsar, Astrophys. J. **720**, 272 (2010).
- [36] M. Razzano, D. Dumora, and F. Gargano, High-energy gamma-ray observations of Geminga with the Fermi Large Area Telescope, arXiv:0912.5442.
- [37] M. Ackermann *et al.* (Fermi-LAT Collaboration), Fermi-LAT search for pulsar wind nebulae around gamma-ray pulsars, Astrophys. J. **726**, 35 (2011).
- [38] A. A. Abdo *et al.*, Milagro observations of multi-TeV emission from galactic sources in the Fermi bright source list, Astrophys. J. **700**, L127 (2009).
- [39] A. U. Abeysekara *et al.*, The 2HWC HAWC Observatory Gamma Ray Catalog, Astrophys. J. **843**, 40 (2017).
- [40] B. M. Baughman and J. Wood (HAWC Collaboration), TeV Gamma-Ray Emission Observed from Geminga with HAWC, arXiv:1508.03497.
- [41] A. Carramiñana, First year results of the High Altitude Water Cherenkov observatory, J. Phys. Conf. Ser. 761, 012034 (2016).
- [42] J. Pretz (HAWC Collaboration), Highlights from the high altitude water Cherenkov observatory, *Proc. Sci.*, ICRC2015, (2016) 025.
- [43] M. G. Aartsen *et al.* (IceCube Collaboration), Searches for extended and point-like neutrino sources with four years of IceCube data, Astrophys. J. **796**, 109 (2014).
- [44] G. R. Blumenthal and R. J. Gould, Bremsstrahlung, synchrotron radiation, and Compton scattering of high-energy

HAWC OBSERVATIONS STRONGLY FAVOR PULSAR ...

electrons traversing dilute gases, Rev. Mod. Phys. 42, 237 (1970).

- [45] M. S. Longair, *High Energy Astrophysics*, 3rd ed. (Cambridge University Press, New York, 2011).
- [46] B. M. Gaensler and P. O. Slane, The evolution and structure of pulsar wind nebulae, Annu. Rev. Astron. Astrophys. 44, 17 (2006).
- [47] M. L. Ahnen *et al.*, Search for VHE gamma-ray emission from Geminga pulsar and nebula with the MAGIC telescopes, Astron. Astrophys. **591**, A138 (2016).
- [48] A. Flinders (VERITAS Collaboration), VERITAS observations of the Geminga supernova remnant, *Proc. Sci.*, ICRC2015, (2016) 795.