

Linear sigma model for multiflavor gauge theories

Y. Meurice

Department of Physics and Astronomy, The University of Iowa, Iowa City, Iowa 52242, USA
(Received 29 September 2017; published 11 December 2017)

We consider a linear sigma model describing $2N_f^2$ bosons (σ , \mathbf{a}_0 , η' and $\boldsymbol{\pi}$) as an approximate effective theory for a $SU(3)$ local gauge theory with N_f Dirac fermions in the fundamental representation. The model has a renormalizable $U(N_f)_L \otimes U(N_f)_R$ invariant part, which has an approximate $O(2N_f^2)$ symmetry, and two additional terms, one describing the effects of a $SU(N_f)_V$ invariant mass term and the other the effects of the axial anomaly. We calculate the spectrum for arbitrary N_f . Using preliminary and published lattice results from the LatKMI collaboration, we found combinations of the masses that vary slowly with the explicit chiral symmetry breaking and N_f . This suggests that the anomaly term plays a leading role in the mass spectrum and that simple formulas such as $M_\sigma^2 \simeq (2/N_f - C_\sigma)M_{\eta'}^2$ should apply in the chiral limit. Lattice measurements of $M_{\eta'}^2$ and of approximate constants such as C_σ could help in locating the boundary of the conformal window. We show that our calculation can be adapted for arbitrary representations of the gauge group and in particular to the minimal model with two sextets, where similar patterns are likely to apply.

DOI: 10.1103/PhysRevD.96.114507

I. INTRODUCTION

The linear version of the sigma models introduced by Gell-Mann and Levy [1] has played an influential role [2] in the establishment of the standard model. In today's usage [3,4], the nonlinear versions not involving the σ -particle [$f_0(500)$] are favored in quantum chromodynamics (QCD) based low-energy calculations. However, when dealing with the explicit breaking of the axial $U(1)_A$ symmetry, linear models are used to describe the η' [5–8]. In addition to the η' and $\boldsymbol{\pi}$, the linear models involve the σ and the \mathbf{a}_0 [0^+ isovectors, $a_0(980)$]. In QCD, the σ and the \mathbf{a}_0 are correctly considered as “heavy” particles compared to the “light” pions which—unlike their 0^+ counterparts—become massless in the chiral limit. However if enough light flavors are added, the separation between light and heavy changes, and the possibility of having a light σ is quite attractive from the low-energy point of view.

The spectrum of multiflavor gauge theories has been vigorously investigated in the context of finding hypothetical strongly interacting particles responsible for the formation of the Brout-Englert-Higgs particle and the electroweak symmetry breaking. For recent reviews of the physics motivations and the literature on this subject, we recommend Refs. [9–11]. The estimation of the mass of flavor singlets using lattice simulations involves disconnected diagrams and is computationally expensive. There are only a few available results. For instance, the estimation of the η' in QCD has only been achieved recently [12–14]. Similarly, there are only a few results available in the multiflavor case.

Recently, light σ masses were found for $SU(3)$ gauge theories with eight [15–18] and twelve [17,19] fundamental flavors and also for two sextets [20]. In addition, preliminary results [21,22] concerning the mass of the η' were announced

at recent conferences. With this information, we investigate the possibility that the explicit breaking of the axial $U(1)_A$ symmetry, which depends in a distinct way on N_f , plays an important role in the determination of the spectrum and the boundary of the conformal window where a nontrivial infrared fixed point is present [23–25]. As the models mentioned above have a low-energy behavior significantly different from QCD, it is very desirable from the point of view of model building to have a simple effective description of this behavior.

In this paper, we consider a generalization of the models discussed for $N_f = 2$ [7], and $N_f = 3$ [5,6,8] in the context of QCD. This is a linear sigma model describing $2N_f^2$ bosons (σ , \mathbf{a}_0 , η' and $\boldsymbol{\pi}$), using the QCD terminology. We use it here as an approximate effective theory for a $SU(3)$ local gauge theory with N_f Dirac fermions in the fundamental representation. The Lagrangian has a renormalizable $U(N_f)_L \otimes U(N_f)_R$ invariant part and two additional terms, one representing a $SU(N_f)_V$ invariant mass term and the other the axial anomaly.

A $SU(N_f)_V$ invariant effective theory for pions is discussed in [26] for arbitrary N_f . There has been a recent interest to include the σ in dilatonic effective theories [27–29]. We briefly motivate the inclusion of the “complex partners,” the η' and \mathbf{a}_0 . In the case of $N_f = 2$, the term that we use to describe the axial anomaly effect is simply a mass term with alternate signs,

$$V_a|_{N_f=2} \propto (\eta'^2 - \sigma^2 + \mathbf{a}_0^2 - \boldsymbol{\pi}^2). \quad (1)$$

The $N_f = 2$ model was used by 't Hooft [7] to explain the role that the instantons play in the spectrum because if we

replace the effective bosonic degrees of freedom in (1) by their quark content ($\mathbf{a}_0 \sim \bar{\psi}\tau\psi$ etc.), we recognize a term of the 't Hooft determinant [30]. In the following, we discuss the generalization for an arbitrary number of flavors N_f . As we see, the axial anomaly term is essential to get a light σ .

The main goal of the article is to show that linear combinations of ratios of meson masses given in Eq. (26) vary slowly with N_f and the quark mass. The slow variations are shown in Figs. 1 and 2. The linear coefficients depend only on N_f and are thus model independent. These empirical results suggest that in the chiral limit,

$$M_{mes.}^2 \simeq (2/N_f - C_{mes.})M_{\eta'}^2, \quad (2)$$

where the subscript *mes.* refers to a σ or \mathbf{a}_0 meson and $C_{mes.}$ is positive, significantly smaller than 1 and approximately N_f independent.

This leads to a simple picture where mesons which are considered heavy for $N_f = 3$ become light and possibly massless for larger N_f . This picture does not depend crucially on the details of the linear model we used to guess the slow variations in Eq. (26). The $C_{mes.}$ can be expressed in terms of the parameters and couplings entering in the simple linear sigma model used in this article; however refinements are needed to get a more complete description of the spectrum. Recent work on linear sigma models [31–34] discusses variations of the form of the anomaly term, inclusion of tetraquarks and vector mesons. In addition, the inclusion of finite-temperature effects [35,36] could help clarify the nature of the transition for $N_f = 8$ [37]. It would be interesting to calculate accurately the $C_{mes.}$ and test the hypothesis that they are meson independent and approximately equal to $2/N_{fc}$, where N_{fc} is the smallest value of N_f in the conformal window.

The linear sigma model is presented in Sec. II. The tree level spectrum is calculated for arbitrary N_f in Sec. III.




FIG. 1. R_σ for $N_f = 8$ (diamonds) and 12 (upside-down triangles) and R_{a_0} for $N_f = 8$ (squares) and 12 (triangles), versus am_f .




FIG. 2. R_σ for $N_f = 8$ (diamonds) and 12 (upside-down triangles) and R_{a_0} for $N_f = 8$ (squares) and 12 (triangles), versus $M_\pi/M_{\eta'}$.

In Sec. IV, we introduce dimensionless quantities involving the masses. Preliminary [22] and published [17] lattice results indicate that they vary slowly with the explicit chiral symmetry breaking and N_f . This provides approximate mass formulas that, if properly refined by future lattice calculations may help identify instabilities for large enough N_f and pinpoint the boundary of the conformal window. In Sec. V we show how to extend our results for fermions in an arbitrary representation of the gauge group. In the conclusions, we discuss possible improvements and the relevance of the results for future lattice calculations.

II. THE MODEL

Following Refs. [5–8], we consider a $N_f \times N_f$ matrix of effective fields ϕ_{ij} having the same quantum numbers as $\bar{\psi}_{Rj}\psi_{Li}$ with the summation over the color indices being implicit. Under a general transformation of $U(N_f)_L \otimes U(N_f)_R$, we have

$$\phi \rightarrow U_L \phi U_R^\dagger. \quad (3)$$

We now use a basis of $N_f \times N_f$ Hermitian matrices Γ^α such that

$$\text{Tr}(\Gamma^\alpha \Gamma^\beta) = (1/2)\delta^{\alpha\beta}, \quad (4)$$

to express ϕ in terms of N_f^2 scalars (0^+ in J^P notation), denoted S_α , and N_f^2 pseudoscalars (0^-), denoted P_α ,

$$\phi = (S_\alpha + iP_\alpha)\Gamma^\alpha, \quad (5)$$

with a summation over $\alpha = 0, 1, \dots, N_f^2 - 1$. We use the convention that $\Gamma_0 = 1/\sqrt{2N_f}$ while the remaining $N_f^2 - 1$ matrices are traceless.

We introduce the diagonal subgroup $U(N_f)_V$ defined by the elements of $U(N_f)_L \otimes U(N_f)_R$ such that $U_L = U_R$. Using Eqs. (5) and (3) we see that under $U(N_f)_V$, S_0 and P_0 are singlets denoted σ and η' respectively while the remaining components transform like the adjoint representation and are denoted \mathbf{a}_0 and $\boldsymbol{\pi}$ respectively.

We consider the effective Lagrangian

$$\mathcal{L} = \text{Tr} \partial_\mu \phi \partial^\mu \phi^\dagger - V \quad (6)$$

with the potential split into three parts

$$V = V_0 + V_a + V_m, \quad (7)$$

that we now proceed to define and discuss. The first term is the most general $U(N_f)_L \otimes U(N_f)_R$ invariant renormalizable expression,

$$V_0 \equiv -\mu^2 \text{Tr}(\phi^\dagger \phi) + (1/2)(\lambda_\sigma - \lambda_{a0})(\text{Tr}(\phi^\dagger \phi))^2 + (N_f/2)\lambda_{a0} \text{Tr}((\phi^\dagger \phi)^2). \quad (8)$$

The use of $\lambda_\sigma - \lambda_{a0}$ becomes clear when we write the mass formulas. The stability of V_0 is discussed in Sec. IV. Note that the first two terms and the kinetic term have a larger group of symmetry $O(2N_f^2)$. The second term

$$V_a \equiv -2(2N_f)^{N_f/2-2} X (\det \phi + \det \phi^\dagger), \quad (9)$$

is invariant under $SU(N_f)_L \otimes SU(N_f)_R$ but breaks the axial $U(1)_A$. It takes into account the effect of the axial anomaly for the fundamental representation. The generalization to arbitrary representations is discussed in Sec. V. The prefactor $2(2N_f)^{N_f/2-2}$ is chosen in order to make the expression of the spectrum as simple as possible. The parameter X has a mass dimension $4 - N_f$. Related effective descriptions of the breaking of the $U(1)_A$ can be found in the literature [38–41].

Finally the third term represents the effect of the mass term which is the same for the N_f flavors,

$$V_m \equiv -(b/\sqrt{2N_f})(\text{Tr} \phi + \text{Tr} \phi^\dagger) = -b\sigma. \quad (10)$$

It is invariant under $SU(N)_V$.

In the following, we assume that chiral symmetry is spontaneously broken by a $SU(N_f)$ invariant vacuum expectation value (vev),

$$\langle \phi_{ij} \rangle = v \delta_{ij} / \sqrt{2N_f}. \quad (11)$$

This amounts to saying that $\langle \sigma \rangle = v$ while the other vevs are 0. We impose that

$$\partial V / \partial \phi|_{\langle \phi \rangle} = 0. \quad (12)$$

Thanks to the simple form of the vevs in Eq. (11), these N_f^2 equations reduce to a single one,

$$-\mu^2 v + (1/2)\lambda_\sigma v^3 - (X/N_f)v^{N_f-1} = b. \quad (13)$$

III. THE SPECTRUM

We can now calculate the tree level spectrum. The normalization (4) implies that the kinetic term in Eq. (6) is canonical,

$$\text{Tr} \partial_\mu \phi \partial^\mu \phi^\dagger = (1/2) \sum_\alpha (\partial_\mu S_\alpha \partial^\mu S_\alpha + \partial_\mu P_\alpha \partial^\mu P_\alpha). \quad (14)$$

The mass of the fields is then obtained as the second derivatives in an obvious way (as for free Klein-Gordon fields). In addition, the unbroken $SU(N_f)_V$ symmetry simplifies the formulas which can be expressed in terms of four masses,

$$\begin{aligned} \partial^2 V / \partial S_0 \partial S_0|_{\langle \phi \rangle} &= M_\sigma^2, \\ \partial^2 V / \partial S_i \partial S_j|_{\langle \phi \rangle} &= \delta_{ij} M_{a0}^2, \\ \partial^2 V / \partial P_0 \partial P_0|_{\langle \phi \rangle} &= M_{\eta'}^2, \\ \partial^2 V / \partial P_i \partial P_j|_{\langle \phi \rangle} &= \delta_{ij} M_\pi^2. \end{aligned} \quad (15)$$

By convention, in the four above equations, the latin indices run from 1 to $N_f^2 - 1$ (isovector indices in the $N_f = 2$ language). The notation M_σ^2 does not mean that this quantity is automatically positive. A negative value could indicate an instability. Since $\langle \phi \rangle$ is proportional to the identity, the second derivatives can be easily calculated at the assumed vevs. For instance, the derivative of the determinant involves the inverse which can then be easily evaluated. This would not be the case for an arbitrary breaking where we would need to use f and d symbols [8]. For the pions, we have

$$M_\pi^2 = -\mu^2 + (1/2)\lambda_\sigma v^2 - (X/N_f)v^{N_f-2}. \quad (16)$$

Using the minimization condition (13), this can be recast in the form

$$M_\pi^2 v = b. \quad (17)$$

In other words in the absence of the explicit mass breaking ($b = 0$), we have the familiar result $M_\pi^2 v = 0$ and $v \neq 0$ implies that in this chiral limit, the pions are exactly massless Nambu-Goldstone bosons. The vev v is related to the pion decay constant in the following way:

$$f_\pi = \sqrt{2/N_f} v. \quad (18)$$

This result can be obtained by considering a $U(N_f)_A$ transformation,

$$\phi \rightarrow U\phi U \simeq \phi + i\omega_\alpha \{\Gamma^\alpha, \phi\}, \quad (19)$$

and showing that the Noether current satisfies the PCAC relation

$$\partial^\mu J_\mu^\alpha = b\sqrt{2/N_f}P^\alpha. \quad (20)$$

We used the convention that

$$\langle \Omega J_\mu^\alpha(x) | P^\beta(p) \rangle = i\delta^{\alpha\beta} f_\pi p_\mu e^{-ipx}. \quad (21)$$

The other results for the spectrum can be written in a compact way,

$$\begin{aligned} M_{\eta'}^2 - M_\pi^2 &= Xv^{N_f-2}, \\ M_\sigma^2 - M_\pi^2 &= \lambda_\sigma v^2 - (1 - 2/N_f)Xv^{N_f-2}, \\ M_{a_0}^2 - M_\pi^2 &= \lambda_{a_0}v^2 + (2/N_f)Xv^{N_f-2}. \end{aligned} \quad (22)$$

One can check that these results agree with the corresponding results [8] for $N_f = 3$ in the $SU(3)_V$ limit. In the chiral limit ($b = 0$), Eq. (22) reduce to

$$\begin{aligned} M_\sigma^2 &= \lambda_\sigma v^2 - (1 - 2/N_f)M_{\eta'}^2, \\ M_{a_0}^2 &= \lambda_{a_0}v^2 + (2/N_f)M_{\eta'}^2. \end{aligned} \quad (23)$$

The sign of the interaction in the anomaly term V_a has been chosen in such a way that $M_{\eta'}^2 \geq M_\pi^2$ as in QCD. This feature persists in more general situations. This implies that for $N_f \geq 3$ the mass of the σ has two contributions of opposite sign. In order to have $M_\sigma \simeq M_\pi$, the two contributions should either be small separately or cancel each other. We see that the second possibility seems to be realized in a certain number of situations.

Notice that if b , λ_{a_0} and X are set to 0, $M_{\eta'} = M_{a_0} = 0$ and the η' and \mathbf{a}_0 could be interpreted as N_f^2 additional Nambu-Goldstone bosons. This is because in this limit, the effective Lagrangian has an $O(2N_f^2)$ symmetry and the vev of the σ breaks it down to $O(2N_f^2 - 1)$ resulting in a total of $2N_f^2 - 1$ Nambu-Goldstone bosons.

In the next section, we explain that (1) it is legitimate to consider the $\lambda_{a_0}v^2$ contribution as small, and (2) the $M_{\eta'}^2$ is a significant contribution partially suppressed by the $1/N_f$ factor in the $M_{a_0}^2$ formula. In that sense, it seems legitimate to treat the \mathbf{a}_0 as light particles in some region of the (m_f, N_f) plane. We should add that the chiral limit obtained from a linear fit of the data for M_{a_0} (see Fig. 28 of Ref. [17]) gives a value significantly smaller than masses quoted with a finite m_f . For instance, at $am_f = 0.03$, we have $aM_{a_0} \simeq 0.46$,

while the chiral extrapolation is $aM_{a_0} \simeq 0.16$. In other words the slope in the aM_{a_0} versus am_f graph is rather large (about 10).

IV. DIMENSIONLESS RATIOS

In order to allow comparisons of numerical results at different lattice spacings, we introduce the dimensionless ratios,

$$R_\sigma \equiv \lambda_\sigma v^2 / M_{\eta'}^2, \quad (24)$$

and

$$R_{a_0} \equiv \lambda_{a_0} v^2 / M_{\eta'}^2. \quad (25)$$

We want to test the idea that these quantities vary slowly with the explicit breaking of chiral symmetry (due to the mass of the fermions m_f) and N_f . If this is the case, the mass formulas have a simple approximate form which could provide a nice intuitive picture. To make things completely clear, the ratios should be understood as functions of the spectroscopic data, namely

$$\begin{aligned} R_\sigma &= (M_\sigma^2 - M_\pi^2) / M_{\eta'}^2 + (1 - 2/N_f)(1 - M_\pi^2 / M_{\eta'}^2), \\ R_{a_0} &= (M_{a_0}^2 - M_\pi^2) / M_{\eta'}^2 - (2/N_f)(1 - M_\pi^2 / M_{\eta'}^2). \end{aligned} \quad (26)$$

Using the preliminary results presented at Lattice 2017 [22] and the published results of Ref. [17] we obtained Fig. 1. As explained in the appendix, these results have been obtained by using the largest volume available for each mass. The error bars do not take into account the finite volume effects and could be significantly larger if we had taken them into account. Nevertheless, Fig. 1 is consistent with the idea of slowly varying ratios. We propose the order of magnitude estimates,

$$R_\sigma \simeq 0.7 + \delta_\sigma(m_f, N_f), \quad (27)$$

and

$$R_{a_0} \simeq -0.1 + \delta_{a_0}(m_f, N_f), \quad (28)$$

where the $\delta(m_f, N_f)$ describe slow variations that need to be studied carefully in the future. The spectroscopic data are often given in terms of m_f in lattice spacing units, but we could as well parametrize the residuals in terms of another symmetry breaking parameter, for instance M_π^2 . Another possibility is to use the dimensionless ratio $M_\pi / M_{\eta'}$ which does not involve the lattice spacing and may be more suitable to compare results at different N_f . This is done in Fig. 2.

Figure 1 shows that for small m_f , we have negative values of R_{a_0} . Can λ_{a_0} be negative? The stability of the effective potential requires that V_0 defined in (8) stays positive for large values of the field ϕ . Using the symmetry

$U(N_f)_L \otimes U(N_f)_R$ of V_0 , we can diagonalize ϕ . $\phi^\dagger \phi$ is then a diagonal matrix with positive diagonal terms $|\alpha_i|^2$ and the sum of the two quartic terms of V_0 remain positive when $\lambda_{a0} = -|\lambda_{a0}|$ is negative provided that

$$(1/2)(\lambda_\sigma + |\lambda_{a0}|) \left(\sum_{i=1}^{N_f} |\alpha_i|^2 \right)^2 \geq (N_f/2) |\lambda_{a0}| \sum_{i=1}^{N_f} |\alpha_i|^4. \quad (29)$$

This inequality should remain valid for any choice of α_i . Considering the case where only one $|\alpha_i|$ becomes arbitrarily large, we get the requirement

$$\lambda_\sigma \geq (N_f - 1) |\lambda_{a0}|. \quad (30)$$

Using the inequality

$$\left(\sum_{i=1}^{N_f} |\alpha_i|^2 \right)^2 \geq \sum_{i=1}^{N_f} |\alpha_i|^4, \quad (31)$$

we see that the requirement (30) is also a sufficient condition in general.

Using the estimates of Eqs. (27) and (28) in the chiral limit, we obtain the simple approximate picture,

$$(M_\sigma/M_{\eta'})^2 \approx 2/N_f - 0.3 + \delta_\sigma(0, N_f), \quad (32)$$

$$(M_{a_0}/M_{\eta'})^2 \approx 2/N_f - 0.1 + \delta_{a_0}(0, N_f). \quad (33)$$

The stability bound $M_\sigma^2 \geq 0$ could provide an estimate of N_{fc} . The idea of having $R_{a_0} < 0$ is attractive because it implies additional stability bounds on N_f coming from $M_{a_0}^2 \geq 0$. For instance, if we use the rough chiral limit estimates $R_\sigma \approx 0.8$ and $R_{a_0} \approx -0.2$, suggested by Fig. 1, we obtain the same $N_{fc} \approx 10$ from (32) with $\delta_\sigma(0, N_f) = 0.1$, and (33) with $\delta_{a_0}(0, N_f) = -0.1$, respectively. In addition, the stability bound of Eq. (30) implies

$$N_f \leq 1 + R_\sigma/|R_{a_0}|. \quad (34)$$

With our chiral limit estimates this corresponds to $N_{fc} \approx 5$. These numbers should not be taken too seriously. They are just meant to illustrate the fact that a careful study of the residuals $\delta(m_f, N_f)$ might help to pinpoint the boundary of the conformal window.

Figure 1 suggests that we could try to find functional relations that are approximately N_f independent, at least for sufficiently massive theories. For the common mass $am_f = 0.04$, the ratios are almost identical despite significantly different meson masses (see Table I in the appendix). Results at smaller am_f would be interesting to see if the chiral limits are very different for $N_f = 12$ and $N_f = 8$, as expected if they are on opposite sides of the conformal window. Since am_f depend explicitly on the lattice spacing

a , we have also plotted the ratios versus the dimensionless ratio $M_\pi/M_{\eta'}$ in Fig. 2. This may be a better way to proceed with the functional relations idea. We also have one data point available [22] for $N_f = 4$ at much smaller volume and $am_f = 0.01$ providing $R_\sigma \sim 0.55$ which is close to 0.7. More data for this case as well as $N_f = 6$ and 10 would be very desirable to study the residual functions δ .

V. MODIFICATION FOR HIGHER REPRESENTATIONS

If the microscopic theory is defined by fermions in higher dimensional representations, for instance $SU(3)$ sextets (the twice symmetric representation), we need to modify the axial anomaly term as

$$V_a(K) = -2(2N_f)^{N_f/2-2} X((\det \phi)^K + (\det \phi^\dagger)^K), \quad (35)$$

with K being the number of zero modes in an instanton background. It also appears in the one-loop coefficient of the Callan-Symanzik beta function. It can be written as $K = 2T[R]$ where $T[R]$ is the trace normalization of the representation R (see Ref. [42] for a lattice discussion). For instance, for $SU(3)$ sextets, $K = 5$. In general, $T[R]$ can be calculated in terms of the Casimir operator of the representation $C_2(R)$ using the relation $T[R] = d(R)C_2(R)/d(\text{Adjoint})$. The coupling X has now a dimension $4 - KN_f$.

The minimization equation (13), M_π^2 and $M_{a_0}^2$, can be generalized by changing

$$X \rightarrow \tilde{X} \equiv XK(v/\sqrt{2N_f})^{N_f(K-1)} \quad (36)$$

into the equations for the fundamental representation derived above. For $M_{\eta'}^2$ and M_σ^2 , in addition to this substitution we need to add a term $(K-1)\tilde{X}$ with a positive sign for η' and a negative sign for σ . In the ‘‘minimal’’ case of $N_f = 2$ $SU(3)$ sextets studied in Refs. [20,43–46],

$$\begin{aligned} M_{\eta'}^2 - M_\pi^2 &= (25/256)Xv^8, \\ M_\sigma^2 - M_\pi^2 &= \lambda_\sigma v^2 - (5/64)Xv^8, \\ M_{a_0}^2 - M_\pi^2 &= \lambda_{a_0} v^2 + (5/256)Xv^8. \end{aligned} \quad (37)$$

Again we see that the M_σ^2 receives contributions of opposite signs, making $M_\sigma^2 < M_\pi^2$ observed in Ref. [20] possible in our model. In the chiral limit ($b = 0$), Eq. (37) reduce to

$$\begin{aligned} M_\sigma^2 &= \lambda_\sigma v^2 - (4/5)M_{\eta'}^2, \\ M_{a_0}^2 &= \lambda_{a_0} v^2 + (1/5)M_{\eta'}^2. \end{aligned} \quad (38)$$

Calculating $M_{\eta'}^2$ for this model would be quite interesting.

VI. CONCLUSIONS

In summary, we have adapted a linear sigma model used in the context of the study of the QCD axial anomaly, to an arbitrary number N_f of equal mass fermions. Equations (22) provide simple formulas for the tree level spectrum. Using preliminary [22] and published [17] results, we found combinations of the masses that appear to vary slowly with the explicit chiral symmetry breaking and N_f . If confirmed by new numerical calculations at different masses and N_f , this would imply that the axial anomaly term plays a leading role in the mass spectrum and is essential to get a light σ . The measurement of $M_{\eta'}^2$ is a crucial ingredient to locate the boundary of the conformal window.

The ratios in Figs. 1 and 2 suggest looking for approximately N_f -independent relationships. It is expected [10] that for m_f large enough, the fermions decouple and the low-energy theory is confining. Consequently, it is plausible that the massive theories at different N_f have effective theories that can be smoothly connected. However, if $N_f = 12$ and $N_f = 8$ are on opposite sides of the conformal window, the massless limit of their effective theory should reveal important differences. Investigating the massless limit of $N_f = 12$ within the framework proposed here should be quite interesting. Larger masses should also be investigated. If R_σ stays approximately flat when m_f increases, one would expect that the σ remains the lightest state. On the lattice, at sufficiently large mass and coupling, the line of first order transition has an end point where one expects a second order phase transition and a light scalar in its vicinity [47–49]. If this is a lattice artifact or something

that could have a counterpart in the continuum is an open question that is worth investigating.

Our main results are model independent and suggest that instabilities appear for N_f large enough. More work needs to be done in order to pinpoint a precise value for the critical value N_{fc} . In addition to using more accurate results expected from ongoing and future simulations, one could also consider more refined sigma models. For instance, we modeled the anomaly effect with a simple determinant term. Variations of this form including powers, logarithms and multiplication by traces are discussed in Refs. [6,31,34]. We have also neglected the vector mesons and the mixing with tetraquark states which are considered in Refs. [31–33]. If used together with numerical spectroscopic results, this could lead to global fits and a better determination of N_{fc} . Another interesting variation of the calculation performed here would be to relax the equality of the masses of the N_f flavors, for instance by splitting 12 flavors into four light and eight heavy flavors [50]. This work is in progress.

ACKNOWLEDGMENTS

We thank Y. Aoki, C. Bernard, T. DeGrand, D. Negradi, E. Rinaldi, D. Schaich, B. Svetitsky and O. Witzel for useful conversations and suggestions. We thank E. Rinaldi for providing the preliminary η' masses [22] with error bars. This research was supported in part by the Department of Energy (DOE) under Award No. DE-SC0010113. This work started while attending the program ‘‘Lattice gauge theory for the LHC and beyond’’ at the Kavli Institute for Theoretical Physics, which is supported by the National Science Foundation under Grant No. PHY11-25915.

APPENDIX: DATA FOR FIGS. 1 AND 2

In this appendix, we explain how we selected the data used in Figs. 1 and 2. We used Tables XVII, XXI, XXII, XXIII, XXVII, XXVIII and XXIX of [17] for $N_f = 8$ and Table XXXVIII (with $\beta = 4$) for $N_f = 12$. We always used the largest volume available. For instance, $36^3 \times 48$ for the four masses with $am_f = 0.02$ and $N_f = 8$. So different masses may have different volumes. The volume effects are typically a bit smaller than the quoted systematic and statistical errors but not negligible. The rest of the data come from the graphs of Ref. [22] available on the Lattice 2017 link of Ref. [51]. This is collected in Table I. After our article was submitted, the data presented in Ref. [22] appeared in a preprint [52]. In addition, a linear sigma model (without an anomaly term) discussed for $N_f = 8$ at the same conference appeared as a preprint [53].

TABLE I. LatKMI data used for the graph.

N_f	am_f	aM_π^2	aM_σ^2	$aM_{a_0}^2$	$aM_{\eta'}^2$
8	0.012	0.164(1)	0.151(15)	0.279(10)	0.875(55)
8	0.015	0.186(1)	0.162(23)	0.310 (10)	0.954(63)
8	0.020	0.221(1)	0.190(17)	0.365(11)	0.956(49)
8	0.030	0.281(2)	0.282(27)	0.480(39)	0.945(69)
8	0.040	0.335(2)	0.365(43)	0.567(23)	0.977(42)
12	0.040	0.2718(7)	0.24(1)	0.3820(21)	0.815(43)
12	0.050	0.3186(4)	0.28(2)	0.4469(25)	0.850(44)
12	0.060	0.3629(3)	0.30(2)	0.5032(36)	1.026(59)

- [1] M. Gell-Mann and M. Levy, *Nuovo Cimento* **16**, 705 (1960).
- [2] B. Lee, in *Chiral Dynamics*, Documents on Modern Physics (Gordon & Breach Science Publishers Ltd, Philadelphia, USA, 1972).
- [3] S. Scherer and M. R. Schindler, arXiv:hep-ph/0505265.
- [4] J. Bijnens, *Prog. Part. Nucl. Phys.* **58**, 521 (2007).
- [5] J. Schechter and Y. Ueda, *Phys. Rev. D* **3**, 2874 (1971).
- [6] C. Rosenzweig, J. Schechter, and C. G. Trahern, *Phys. Rev. D* **21**, 3388 (1980).
- [7] G. 't Hooft, *Phys. Rep.* **142**, 357 (1986).
- [8] Y. Meurice, *Mod. Phys. Lett. A* **02**, 699 (1987).
- [9] B. Svetitsky, in *35th International Symposium on Lattice Field Theory (Lattice 2017), Granada, Spain, 2017* (2017) [arXiv:1708.04840].
- [10] T. DeGrand, *Rev. Mod. Phys.* **88**, 015001 (2016).
- [11] D. Negradi and A. Patella, *Int. J. Mod. Phys. A* **31**, 1643003 (2016).
- [12] N. H. Christ, C. Dawson, T. Izubuchi, C. Jung, Q. Liu, R. D. Mawhinney, C. T. Sachrajda, A. Soni, and R. Zhou, *Phys. Rev. Lett.* **105**, 241601 (2010).
- [13] E. B. Gregory, A. C. Irving, C. M. Richards, and C. McNeile (UKQCD Collaboration), *Phys. Rev. D* **86**, 014504 (2012).
- [14] K. Ottnad, C. Urbach, and F. Zimmermann (OTM Collaboration), *Nucl. Phys.* **B896**, 470 (2015).
- [15] Y. Aoki *et al.* (LatKMI Collaboration), *Phys. Rev. D* **89**, 111502 (2014).
- [16] T. Appelquist, R. C. Brower, G. T. Fleming, A. Hasenfratz, X. Y. Jin, J. Kiskis, E. T. Neil, J. C. Osborn, C. Rebbi, E. Rinaldi, D. Schaich, P. Vranas, E. Weinberg, and O. Witzel (Lattice Strong Dynamics Collaboration), *Phys. Rev. D* **93**, 114514 (2016).
- [17] Y. Aoki *et al.* (LatKMI Collaboration), *Phys. Rev. D* **96**, 014508 (2017).
- [18] A. D. Gasbarro and G. T. Fleming *Proc. Sci.*, LATTICE2016 (2017) 242 [arXiv:1702.00480].
- [19] Y. Aoki, T. Aoyama, M. Kurachi, T. Maskawa, K.-i. Nagai, H. Ohki, E. Rinaldi, A. Shibata, K. Yamawaki, and T. Yamazaki (LatKMI Collaboration), *Phys. Rev. Lett.* **111**, 162001 (2013).
- [20] Z. Fodor, K. Holland, J. Kuti, S. Mondal, D. Negradi, and C. H. Wong *Proc. Sci.*, LATTICE2014 (2015) 244.
- [21] Y. Aoki, in *Lattice 2016*.
- [22] E. Rinaldi, in *Lattice 2017*.
- [23] T. Banks and A. Zaks, *Nucl. Phys.* **B196**, 189 (1982).
- [24] F. Sannino and K. Tuominen, *Phys. Rev. D* **71**, 051901 (2005).
- [25] D. D. Dietrich and F. Sannino, *Phys. Rev. D* **75**, 085018 (2007).
- [26] J. Bijnens and J. Lu, *J. High Energy Phys.* **11** (2009) 116.
- [27] M. Golterman and Y. Shamir, *Phys. Rev. D* **94**, 054502 (2016).
- [28] M. Golterman and Y. Shamir *Proc. Sci.*, LATTICE2016 (2016) 205 [arXiv:1610.01752].
- [29] T. Appelquist, J. Ingoldby, and M. Piai, *J. High Energy Phys.* **07** (2017) 035.
- [30] G. 't Hooft, *Phys. Rev. D* **14**, 3432 (1976).
- [31] A. H. Fariborz, R. Jora, and J. Schechter, *Phys. Rev. D* **77**, 094004 (2008).
- [32] D. Parganlija, F. Giacosa, and D. H. Rischke, *Phys. Rev. D* **82**, 054024 (2010).
- [33] D. Parganlija, P. Kovács, G. Wolf, F. Giacosa, and D. H. Rischke, *Phys. Rev. D* **87**, 014011 (2013).
- [34] P. Kovács and G. Wolf, in *Proceedings of the 5th Workshop on Excited QCD 2013, Sarajevo, Bosnia-Herzegovina, 2013*; *Acta Phys. Pol.* **6**, 853 (2013).
- [35] P. Kovács and Z. Szép, *Phys. Rev. D* **75**, 025015 (2007).
- [36] P. Kovács, Z. Szép, and G. Wolf, *Phys. Rev. D* **93**, 114014 (2016).
- [37] A. Deuzeman, M. P. Lombardo, and E. Pallante, *Phys. Lett. B* **670**, 41 (2008).
- [38] P. Di Vecchia and G. Veneziano, *Nucl. Phys.* **B171**, 253 (1980).
- [39] K. Kawarabayashi and N. Ohta, *Nucl. Phys.* **B175**, 477 (1980).
- [40] N. Ohta, *Prog. Theor. Phys.* **66**, 1408 (1981); **67**, 993(E) (1982).
- [41] K. Kawarabayashi and N. Ohta, *Prog. Theor. Phys.* **66**, 1789 (1981).
- [42] Z. Fodor, K. Holland, J. Kuti, D. Negradi, and C. Schroeder, *J. High Energy Phys.* **08** (2009) 084.
- [43] D. D. Dietrich, F. Sannino, and K. Tuominen, *Phys. Rev. D* **72**, 055001 (2005).
- [44] T. DeGrand, Y. Shamir, and B. Svetitsky, *Phys. Rev. D* **79**, 034501 (2009).
- [45] T. DeGrand, Y. Shamir, and B. Svetitsky, *Phys. Rev. D* **87**, 074507 (2013).
- [46] Z. Fodor, K. Holland, J. Kuti, D. Negradi, C. Schroeder, and C. H. Wong, *Phys. Lett. B* **718**, 657 (2012).
- [47] X.-Y. Jin and R. D. Mawhinney, in *Proceedings of the KMI-GCOE Workshop on Strong Coupling Gauge Theories in the LHC Perspective (SCGT 12), Nagoya, Japan, 2012* (World Scientific, Singapore, 2014), pp. 96–102.
- [48] Z. Gelzer, Y. Liu, Y. Meurice, and D. K. Sinclair, *Proc. Sci.*, LATTICE2013 (2014) 078 [arXiv:1312.3906].
- [49] Z. Gelzer, Y. Liu, and Y. Meurice *Proc. Sci.*, LATTICE2014 (2014) 255 [arXiv:1411.3360].
- [50] R. C. Brower, A. Hasenfratz, C. Rebbi, E. Weinberg, and O. Witzel, *Phys. Rev. D* **93**, 075028 (2016).
- [51] Contributions to Lattice 2017, See <https://makondo.ugr.es/event/0/session/96/contribution/350/material/slides/0.pdf>.
- [52] Y. Aoki *et al.*, in *35th International Symposium on Lattice Field Theory (Lattice 2017), Granada, Spain, 2017* (2017) [arXiv:1710.06549].
- [53] A. Gasbarro, in *35th International Symposium on Lattice Field Theory (Lattice 2017), Granada, Spain 2017* (2017) [arXiv:1710.08545].