

Inflation and high-scale supersymmetry with an EeV gravitino

 Emilian Dudas,^{1,*} Tony Gherghetta,^{2,†} Yann Mambrini,^{3,‡} and Keith A. Olive^{2,4,§}
¹*CPhT, Ecole Polytechnique, 91128 Palaiseau Cedex, France*
²*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*
³*Laboratoire de Physique Théorique Université Paris-Sud, F-91405 Orsay, France*
⁴*William I. Fine Theoretical Physics Institute, University of Minnesota, Minneapolis, Minnesota 55455, USA*

(Received 28 October 2017; published 28 December 2017)

We consider inflation and supersymmetry breaking in the context of a minimal model of supersymmetry in which the only “low” energy remnant of supersymmetry is the gravitino with a mass of order an EeV. In this theory, the supersymmetry breaking scale is above the inflaton mass, $m \approx 3 \times 10^{13}$ GeV, as are all sfermion and gaugino masses. In particular, for a no-scale formulation of Starobinsky-like inflation using the volume modulus T , we show that inflation can be accommodated even when the supersymmetry breaking scale is very large. Reheating is driven through a gravitational coupling to the two Higgs doublets and is enhanced by the large μ -parameter. This leads to gravitino cold dark matter where the mass is constrained to be in the range $0.1 \text{ EeV} \lesssim m_{3/2} \lesssim 1000 \text{ EeV}$.

DOI: 10.1103/PhysRevD.96.115032

I. INTRODUCTION

While the Higgs boson was discovered at the LHC [1,2] and is consistent with predictions of low energy supersymmetry (SUSY), [3–7], so far supersymmetry has not been seen experimentally [8]. Whether supersymmetry is waiting around the corner, or is broken at some high scale (intermediate or above) is currently unknown. If indeed supersymmetry is broken above the inflationary scale, it may well be that the only remnant of supersymmetry at low energies is the gravitino which may yet play the role of dark matter [9,10].¹

Below the Planck and grand unified theory (GUT) scales, it would appear that there is an intermediate scale (between the GUT scale and electroweak scale) associated with inflation. For the sake of definiteness, let us consider the Starobinsky model of inflation as an example [12–14]. The inflaton potential can be written as

$$V(t) = \frac{3}{4} m^2 (1 - e^{-\sqrt{\frac{2}{3}} t})^2, \quad (1)$$

where t is the canonically normalized inflaton field. The inflaton mass scale, m , can be determined by the amplitude of density fluctuations [15],

$$A_s = \frac{3m^2}{8\pi^2} \sinh^4(t_*/\sqrt{6}) = 2.1 \times 10^{-9}, \quad (2)$$

where $t_* \approx 5.35$ corresponds to 55 e-folds of inflation. Solving for m in (2), we have $m = 1.2 \times 10^{-5} M_P \approx 3 \times 10^{13}$ GeV, where $M_P = 1/\sqrt{8\pi G_N} \approx 2.4 \times 10^{18}$ GeV.

Interestingly, the mass scale around 10^{13} GeV, may also correspond to an intermediate scale gauge group whose breaking may yield a large Majorana mass for right-handed neutrinos, $M_R \approx m$, appropriate for the see-saw mechanism [16].

Here, we consider the possibility that the supersymmetry breaking scale is also of order the inflaton mass, m . An example is provided by supersymmetry breaking via a Polonyi sector which is achieved with a superpotential of the form [17]

$$W_P = \tilde{m}^2 (Z + b), \quad (3)$$

where Z is the chiral superfield responsible for breaking supersymmetry with auxiliary field component $F \equiv \tilde{m}^2$. Of course if \tilde{m} in (3) is of order the inflaton mass, m , then the masses of the entire supersymmetric spectrum would be of order the intermediate scale and clearly out of reach of any accelerator search. However, the gravitino mass,

$$m_{3/2} = \frac{\tilde{m}^2}{\sqrt{3} M_P}, \quad (4)$$

would be significantly lighter and could still provide for the dark matter in the universe [9,10]. In fact, to avoid over production of gravitinos through the decay of the inflaton to R -parity = -1 matter fields (which subsequently decay to gravitinos), it was argued [10], that the sparticle spectrum should lie above the inflaton mass, thus providing a lower limit to the supersymmetry breaking scale and hence a

*Emilian.Dudas@cph.t.polytechnique.fr

†tgher@umn.edu

‡yann.mambrini@th.u-psud.fr

§olive@physics.umn.edu

¹In this case supersymmetry is nonlinearly realized at lower energies [11].

lower limit to the gravitino mass of $m_{3/2} > 0.2$ EeV. Only an EeV gravitino mass is left behind.

The EeV gravitino as a dark matter candidate is produced after inflation in the process of reheating. The common mechanism [18–32], for producing a single gravitino in thermal scattering processes has a cross section which is temperature independent and scales as $m_{\text{SUSY}}^2/M_P^2 m_{3/2}^2$, where m_{SUSY} is a typical sparticle mass. The rate therefore is roughly $\Gamma \sim T^3 m_{\text{SUSY}}^2/M_P^2 m_{3/2}^2$, where we have assumed predominantly Goldstino production in the limit $m_{3/2} \ll m_{\text{SUSY}}$. If sparticle production is kinematically forbidden, single gravitino production (which must be accompanied by a massive sparticle (gluino), is not operative. Instead, the rate for gravitino production during reheating is suppressed, as only processes which produce two gravitinos are allowed. This cross section is temperature dependent and scales as $\langle \sigma v \rangle \propto T^6/F^4$, so that the rate is roughly $\Gamma \sim T^9/F^4$. In this case, the final gravitino abundance scales as $n_{3/2}/n_\gamma \sim \Gamma/H \sim T^7 M_P/F^4$ evaluated at the reheating temperature, in contrast to the abundance for single gravitino production, $n_{3/2}/n_\gamma \sim \Gamma/H \sim T m_{\text{SUSY}}^2/M_P m_{3/2}^2$. For reheating temperatures of order 10^{10} GeV, the gravitino abundance matches the CMB determined cold dark matter density [15]. It is also possible that the inflaton can decay to two gravitinos, but this is more model dependent and we return to this possibility in Sec. III D.

The phenomenology of this high scale supersymmetric model is simple. The (not so) low energy spectrum consists of the gravitino and perhaps the scalars associated with the chiral superfield, Z . However, as we discuss in Sec. III A, we expect that these are also hierarchically more massive than the gravitino. As has been shown recently [33,34], even with a spectrum as massive as discussed here, a 125 GeV Higgs mass can still be attained if $\tan \beta$ is either small (close to 1) or large (above 60).

The clear drawback of such a model is its testability. In fact, the model in its simplest and most minimal form predicts no signatures in either accelerator searches, or direct and indirect searches for dark matter. Of course if supersymmetry is actually discovered at the LHC, then this model can be ruled out. In a modest extension of the model with R-parity violation in the lepton-Higgs sector, the gravitino becomes unstable, though still suitably long lived. The detection of very high energy neutrinos or photons at HAWC or the Pierre Auger Observatory would be a signature of this model. Another possible signature may come from the observation of non-Gaussianities in the CMB due to scalars with masses near the Hubble scale during inflation [35].

The paper is organized as follows. In Sec. II, we discuss exemplary inflation models with high scale supersymmetry breaking. We focus on models based on no-scale supergravity [36,37] which lead to Starobinsky-like potentials [38–51]. In particular, models which allow for high scale

supersymmetry breaking with stabilized fields [38,39,41,43,45,47] without spoiling the inflationary properties of the potential. In Sec. III, we discuss the phenomenological aspects of the model.² We begin in Sec. III A with a model for gaugino and scalar masses. The model utilizes the strong stabilization of the Polonyi field [45,54–61] used to generate large gaugino masses. Scalar masses are then obtained through threshold corrections as in gaugino mediation [62] or a more strongly-coupled mediation mechanism [63,64]. The requirement that all sparticle masses lie above the inflaton mass will set a constraint on the stabilization scale. The effects of supersymmetry breaking on inflation is then discussed in Sec. III B. In Sec. III C we determine the conditions under which we can obtain a Higgs mass of 125 GeV, as well as preserving the stability of the Higgs vacuum. The requirements for gravitino dark matter are outlined in Sec. III D. Our concluding remarks are given in Sec. IV.

II. INFLATION AND SUPERSYMMETRY BREAKING

There are many ways to proceed in constructing a model of inflation which incorporates supersymmetry breaking. While it would be an overstatement to say that recent Planck results [15] on the CMB spectrum parameters, n_s and r , corresponding to the tilt of the scalar perturbation spectrum and the scalar to tensor ratio, respectively predict Starobinsky-like inflation models, it is clear that these models are for now in very good agreement with Planck results. In particular, we will use formulations of the Starobinsky model based on no-scale supergravity.

A. No-scale supergravity and Starobinsky-like inflation

The Kähler potential in the context of no-scale supergravity can be written in the form [36,37],

$$K = -3 \ln \left(T + \bar{T} - \frac{1}{3} \sum_i |\phi_i|^2 \right), \quad (5)$$

where T is a volume modulus and the ϕ_i include all matter fields and possibly a supersymmetry breaking Polonyi-like field, Z [17]. The inflaton may be identified with either T or a matter-like field, ϕ . Equivalently, we may use a set of field redefinitions [65] and write

$$K = -3 \ln \left(1 - \sum_i \frac{|y_i|^2}{3} \right), \quad (6)$$

where the y_i include all matter fields, moduli and the inflaton. For now, let us ignore the matter fields, and concentrate on a two-field model. There are at least two

²For alternative approaches to inflation in high scale supersymmetric models see [52,53].

independent families [41] of superpotentials which lead to Starobinsky-like inflation. In the first, T is a modulus and ϕ is the inflaton with a Wess-Zumino (WZ) superpotential written as [38]

$$W = m \left(\frac{\phi^2}{2} - \frac{\phi^3}{3\sqrt{3}} \right) \quad \text{WZ}, \quad (7)$$

or in the symmetric basis

$$W = m \left[\frac{y_1^2}{2} \left(1 + \frac{y_2}{\sqrt{3}} \right) - \frac{y_1^3}{3\sqrt{3}} \right] \quad \text{WZ}, \quad (8)$$

which is a WZ model for the inflaton y_1 with an interaction term $y_1^2 y_2$. In both bases, when ϕ (y_1) is redefined to a field x with a canonical kinetic term, the potential is exactly of the form of the Starobinsky potential (1) (with t identified as x), assuming that some dynamics stabilizes and fixes T (y_2): $\langle T \rangle = \langle T^* \rangle = 1/2$ ($\langle y_2 \rangle = 0$). One way to accomplish this is by adding a quartic term in the Kähler potential [41,66].

The second family of models first formulated as an R^2 extension to supergravity by Cecotti [67] can be written as

$$W = \sqrt{3}m\phi(T - 1/2) \quad \text{C}, \quad (9)$$

or

$$W = m y_1 y_2 (1 + y_2/\sqrt{3}) \quad \text{C}, \quad (10)$$

In this case, the inflaton is associated with T (y_2) and it must be assumed that ϕ (y_1) is stabilized at the origin [39,41]. Again, when T (y_2) is normalized to give a proper kinetic term, we get the Starobinsky potential shown in Eq. (1) [39].

In either case (WZ or C), the mass parameter m is related to the inflaton mass, and is set by the amplitude of density fluctuations measured in the CMB through Eq. (2).

B. Effects of supersymmetry breaking on inflation

Supersymmetry breaking can be accomplished in various ways, but in many of these, there are constraints on the SUSY breaking scale due to its effect on the inflationary potential. In general, SUSY breaking perturbs the potential, but these effects may be small, if the supersymmetry breaking scale, $\tilde{m} \ll m$. Indeed, this was one of the initial motivations behind supersymmetric formulations of inflation [68].

The simplest possibility we can consider is adding a constant, w_0 to the superpotential. In most low energy models of SUSY phenomenology, we would relate w_0 to the weak scale through $w_0 = \tilde{m} M_P^2$ and the gravitino mass is just

$$m_{3/2} = \frac{w_0}{(T + \bar{T})^{3/2}} = \tilde{m}, \quad (11)$$

with $T + \bar{T} = 1$, in Planck units. However, in this case low energy SUSY breaking parameters such as soft scalar masses, m_0 , trilinear A -terms and the bi-linear B_0 are all proportional to $m_{3/2}$ ($m_0 = 0$ for untwisted matter fields) [45,69]. The gaugino mass in this case is

$$M_{1/2} = \left| \frac{1}{2} e^{G/2} \frac{\tilde{f}_T}{\text{Ref}} (G^{-1})_T^T G^T \right| = \left| \frac{1}{2} w_0 \frac{\tilde{f}_T}{\text{Ref}} \right|, \quad (12)$$

where $f_{\alpha\beta} = f \delta_{\alpha\beta}$ is the gauge kinetic function. For $T + \bar{T} = 1$, it is unlikely that we get a hierarchy $m_{3/2} \ll M_{1/2}$. Moreover, we would like to relate the supersymmetry breaking scale to the inflationary scale $\tilde{m} = m$. However, *a priori*, we can set w_0 to be either $\tilde{m} M_P^2$, or $\tilde{m}^2 M_P$, or \tilde{m}^3 , giving $m_{3/2} = m$, $m^2/M_P \approx 0.4$ EeV, or $m^3/M_P^2 \approx 5$ TeV (though the latter may be of phenomenological interest at the LHC).

In [50], a linear term $a^2 \phi$ for the inflaton in the WZ model given in (7) was proposed, making the association between the inflaton and Polonyi field. For small a , the theory works quite well, and thus predicts a small (weak scale) gravitino mass. The inflationary capability of the theory breaks down when $a \gtrsim 5 \times 10^{-5}$ corresponding to an upper limit on the gravitino mass of $m_{3/2} = a^4/2m \lesssim 10^6$ GeV.

Next we can consider adding a strongly stabilized (twisted) Polonyi field to the WZ model,

$$w_0 \rightarrow \tilde{m}^2(z + b)(T + 1/2)^p, \quad (13)$$

with

$$K \supset z\bar{z} - \frac{(z\bar{z})^2}{\Lambda_z^2}. \quad (14)$$

The factor $(T + 1/2)^p$ is needed to avoid a deSitter vacuum with weak scale energy density and we will take $p = 3$ as an example here [45]. Choosing $b \approx 1/\sqrt{3}$ gives a minimum with zero vacuum energy at $\langle z \rangle \approx \Lambda_z^2/\sqrt{12}$. The mass of the Polonyi field is now hierarchically larger than the gravitino mass

$$m_z^2 = \frac{12m_{3/2}^2 M_P^2}{\Lambda_z^2}. \quad (15)$$

Once again, for $\tilde{m}^2 \ll m M_P$, this works quite well so long as Λ_z is not too small ($\Lambda_z \gtrsim 2(\tilde{m}^2/m M_P)^{1/3}$). Increasing \tilde{m} , leads to the formation of a new minimum at large field values (of the canonically normalized inflaton), which quickly becomes the global minimum. In Fig. 1, we




FIG. 1. Projections of the effective inflationary potential for the model (7) with the Polonyi sector ((13) and (14)), for $p = 3$. Here $\langle T \rangle = 1/2$, $\langle z \rangle \simeq \Lambda_z^2/\sqrt{12}$ and $b \simeq 1/\sqrt{3}$, and we use the nominal value $\Lambda_z = 10^{-2}$. Shown is the potential for different choices of $\tilde{m}^2/m = 10^{-8}, 10^{-7}, 2 \times 10^{-7}, 5 \times 10^{-7}$ (in Planck units) in black, blue, green, and red.

show the potential for fixed $p = 3$, $\Lambda_z = 10^{-2}$, and several values of \tilde{m} . For $\tilde{m}^2/m < 10^{-8}M_P$, the potential is indistinguishable from that shown as 10^{-8} . For $m \approx 10^{-5}M_P$, we see that this model works fine for weak scale supersymmetry breaking, and for scales as large as $\tilde{m}^2/M_P \lesssim 10^{-12}M_P \sim 3$ PeV, corresponding to a gravitino mass of $m_{3/2} \simeq \tilde{m}^2/\sqrt{3}M_P \lesssim 1.7$ PeV. In particular, $m_{3/2} = 0.2$ EeV would correspond to $\tilde{m}^2/m \approx 10^{-5}M_P$ which would badly spoil the inflationary potential.

While the WZ models are perfectly acceptable for low scale supersymmetry breaking, our objective here is high scale breaking and thus we turn our attention to the case C, for the superpotential given by (9). To achieve supersymmetry breaking and generate a finite gravitino mass, we can again add a constant, w_0 to the superpotential. In this case, if $w_0 \ll m$, the minimum is shifted slightly to [45]

$$\langle T \rangle = \frac{1}{2} - \frac{w_0^2}{m^2}, \quad \langle \phi \rangle = \sqrt{3} \frac{w_0}{m}, \quad (16)$$

but the vacuum energy density is necessarily negative, $V_0 = -3\langle e^G \rangle = -3m^2w_0^2/(m^2 - 3w_0^2) < 0$.

However, adding an untwisted Polonyi field, so that the Kähler potential becomes

$$K = -3 \ln \left(T + \bar{T} - \frac{1}{3} \sum_i |\phi_i|^2 - \frac{1}{3} |z|^2 + \frac{|z|^4}{\Lambda_z^2} \right), \quad (17)$$

with the superpotential given in (13) with $p = 0$ leaves the Starobinsky potential (now a function of T) unchanged, save for a shift in the minimum to

$$\begin{aligned} \langle T \rangle &\simeq \frac{1}{2} + \frac{1}{3} \left(\frac{\tilde{m}^2}{mM_P} \right)^2, & \langle \phi \rangle &\simeq \frac{\tilde{m}^2}{m}, \\ \langle z \rangle &\simeq \frac{\Lambda_z^2}{6\sqrt{3}}, & b &\simeq \frac{1}{\sqrt{3}} \left(1 - \frac{1}{6} \left(\frac{\tilde{m}^2}{mM_P} \right)^2 \right), \end{aligned} \quad (18)$$

when $\tilde{m}^2/(mM_P), \Lambda_z/M_P \ll 1$. The mass of z is $\sqrt{3}$ times larger than the twisted Polonyi mass given in Eq. (15).

Alternatively, one can add a twisted Polonyi field with Kähler potential

$$K = -3 \ln \left(T + \bar{T} - \frac{1}{3} \sum_i |\phi_i|^2 \right) + |z|^2 - \frac{|z|^4}{\Lambda_z^2}, \quad (19)$$

and the same superpotential (9). This also leaves the Starobinsky potential unchanged, with a similar shift in the minimum to [45]

$$\begin{aligned} \langle T \rangle &\simeq \frac{1}{2} + \frac{2}{3} \left(\frac{\tilde{m}^2}{mM_P} \right)^2, & \langle \phi \rangle &\simeq \frac{\tilde{m}^2}{m}, \\ \langle z \rangle &\simeq \frac{\Lambda_z^2}{2\sqrt{3}}, & b &\simeq \frac{1}{\sqrt{3}} \left(1 - \frac{1}{2} \left(\frac{\tilde{m}^2}{mM_P} \right)^2 \right), \end{aligned} \quad (20)$$

Unlike the WZ case discussed above, the inflationary potential maintains its form even for large \tilde{m} , and arbitrarily small Λ_z . For example, in Fig. 2, we show the inflationary potential with $\tilde{m}^2 = 0.9mM_P$ and $\Lambda_z = 10^{-3}M_P$ both the twisted (solid) and untwisted (dashed) Polonyi models. Here $t = \sqrt{3/2} \ln(2T)$ is the canonically normalized inflaton. In the figure, $\langle z \rangle$ and b have been fixed at the approximate values given in (18) and (20), respectively (higher order terms in (18) cannot be neglected).




FIG. 2. Projections of the effective inflationary potential for the model (9) with the Polonyi sector [(13) and (17)], with $p = 0$. We use the nominal values $\Lambda_z = 10^{-3}$ with $\tilde{m}^2/m = 0.9$. The values $\langle z \rangle$, b , and $\langle \phi \rangle$ are given approximately by (18) (shown by the dashed curve), and by (20) (shown by the solid curve).

The gravitino mass for small Λ_z can be written as

$$m_{3/2} = m \frac{5\tilde{m}^6 + 6\tilde{m}^2 m^2 M_P^2}{2(3m^2 M_P^2 + \tilde{m}^4)^{3/2}} \quad \text{untwisted,} \quad (21)$$

$$m_{3/2} = m \frac{4\tilde{m}^6 + 2\tilde{m}^2 m^3 M_P}{2\sqrt{3}(m^2 M_P^2 + \tilde{m}^4)^{3/2}} \quad \text{twisted,} \quad (22)$$

which in the limit of small \tilde{m}^2/mM_P for both cases gives the expected result $m_{3/2} = \tilde{m}^2/\sqrt{3}M_P$. Indeed, for $\tilde{m}^2 = 0.9mM_P$, we obtain $m_{3/2} \sim 10^{13}$ GeV. It is indeed rather surprising that even for a large Polonyi mass scale, the inflationary dynamics are little affected. This is only true for case C given by Eq. (9). Thus we are free to make the simple choice of $\tilde{m} = m \approx 10^{-5}M_P$. In this case, the gravitino mass is

$$m_{3/2} = \frac{m^2}{\sqrt{3}M_P} \approx 0.2 \text{ EeV.} \quad (23)$$

Furthermore, it was shown in the first reference of [47] that one also needs to impose $m_{3/2} < H$ in order to keep perturbative control of the Kähler potential. In what follows, we will restrict our attention to the twisted Polonyi model.

To avoid the production of the Polonyi field during reheating, we can derive an upper limit on Λ_z from the requirement that $m_z = \sqrt{12}m_{3/2}M_P/\Lambda_z > m$. This limit is shown by the blue dashed line in Fig. 3. Another upper limit on Λ_z is obtained by requiring that the branching ratio of inflaton decays to gravitinos does not lead to an excess abundance of gravitinos (discussed in more detail in Sec. III D). This constraint is shown by the negatively sloped pink dot-dashed line. Acceptable parameters lie below both lines (blue dashed and pink dot-dashed) Finally, we also have a lower bound on Λ_z , stemming from our effective correction to the Kähler potential which imposes $F < \langle z \rangle < \Lambda_z^2$ or $1/2 \log(m_{3/2}/M_P) < \log(\Lambda_z/M_P)$. This lower bound is shown by the green dotted line in Fig. 3, and all values of $10^{-5} < m_{3/2}/m < 10^{-1}$ are allowed so long as Λ_z lies in the pale shaded region.

III. THE PARTICLE SPECTRUM

A. Gaugino and scalar masses

We next discuss ways to generate gaugino and scalar masses via perturbative mediations of supersymmetry breaking, and then we will turn to more strongly-coupled mediation mechanisms.

One perturbative possibility for mediation of supersymmetry breaking is via gaugino mediation [62]. New physics at a messenger mass scale M generates a coupling of the supersymmetry breaking field, $Z = z + \theta^2 F$ (ignoring fermionic components) to the gauge fields




FIG. 3. Bounds on the stabilization parameter Λ_z as a function of the gravitino mass. The blue dashed line shows the upper limit on Λ_z from the requirement that the mass of the Polonyi field lies above the inflaton mass. The pink dot-dashed line (negatively sloped) is an upper limit on Λ_z derived from an upper limit on the branching ratio of inflaton decays to gravitinos. The green dotted line shows the lower limit on Λ_z , assuming $F < \Lambda_z^2$. The shaded region is allowed by all constraints. Note, however, that there are lower bounds on the gravitino mass given in Eqs. (27) and (30).

$$\int d^2\theta \left(f_0 + f_1 \frac{Z}{M} \right) \text{tr} W^\alpha W_\alpha, \quad (24)$$

where at leading order $f_0 = 1/(4g^2)$. This will give rise to gaugino masses of order

$$M_{1/2} \sim g^2 f_1 \frac{F}{M}. \quad (25)$$

Scalar masses are then generated by SM loop corrections and will be of order

$$m_0^2 \sim \frac{g^2}{16\pi^2} M_{1/2}^2, \quad (26)$$

where g is a SM gauge coupling. Usually there is a log enhancement of the scalar masses due to the running between the mediation scale M and the scale of the superpartners. In our case however, since the superpartners are very heavy there is not much running, so numerically we can use as an order of magnitude estimate $m_0 \sim (\alpha/g)M_{1/2} \sim 0.06M_{1/2}$. In order for the effective field theory to be well-defined (higher-dimensional operators are negligible), one needs to impose $F < M^2$. We also forbid inflaton decays into superpartners, which roughly

requires $m_0 > 3 \times 10^{13}$ GeV. Under such assumptions with $\sqrt{F} \sim M$, one obtains $\sqrt{F} > 5 \times 10^{14}$ GeV and therefore the minimum value of the gravitino mass in such a scenario is

$$m_{3/2} > 6 \times 10^{10} \text{ GeV} = 60 \text{ EeV}. \quad (27)$$

It seems difficult to decrease the gravitino mass much below 100 EeV with known perturbative mediations of supersymmetry breaking. For example, in a model based on gauge mediation, scalar masses would be expected to be of order the gaugino masses such that $m_0 \sim M_{1/2} \sim (g^2/16\pi^2)F/M$ (hence there is no relative loop suppression). The lower limit on $m_{3/2}$ is then increased to $m_{3/2} > 2 \times 10^{13}$ GeV, which is essentially incompatible with the required upper limit, $m_{3/2} < m$.

In order to decrease the viable values of the gravitino mass, some strong coupling effects seem to be needed, like for example in holographic models of supersymmetry breaking of the type described in [63], or general gauge mediation [64]. In more generic terms, this means a mediation mechanism with no loop suppression in the generation of visible sector soft masses, so that we obtain $m_0 \sim M_{1/2} \sim F/M$. Thus we can start with the same gauge kinetic function as in (24), namely

$$f_{\alpha\beta} = \left(f_0 + f_1 \frac{Z}{M} \right) \delta_{\alpha\beta}, \quad (28)$$

and generate squark/slepton/Higgs soft masses through operators of the type

$$\int d^4\theta \frac{Z^\dagger Z}{M^2} Q^\dagger Q, \quad (29)$$

where Q denotes a generic MSSM chiral superfield. In this case, we recover the limit

$$m_{3/2} > 0.2 \text{ EeV}, \quad (30)$$

and corresponds to the bound derived in [10]. Perturbativity of the correction to gauge couplings $f_1 \frac{\langle z \rangle}{M} < f_0$, together with the requirement, $M_{1/2} > m$ leads to the new lower limit

$$m_{3/2} > \frac{4\langle z \rangle}{\sqrt{3}} \frac{m}{M_P}. \quad (31)$$

However, this constraint is easily satisfied, once the other constraints, such as $F \leq M^2$ are taken into account.

Note that at the lower gravitino mass limit (30) we obtain the bound $g^2 f_1 \gtrsim 1$. Since mediation is strongly coupled, this is not really surprising. In both holographic models and general gauge mediation setups, additional states of mass,

M and heavier are expected. In order to not perturb our single-field inflation framework, the masses of these states should be above the Hubble scale H during inflation,³ which implies generically $M > H$. This condition is satisfied by the range of gravitino masses in Fig. 3 and is saturated at the lower bound.

Finally, we comment on the partial wave unitarity limit arising from the scattering of two gluons into two gravitinos [70]. For gaugino mediation, tree-level unitarity is violated at a scale $\approx 17/(g^2 f_1)M$, which for $g^2 f_1 \lesssim 17$ is above the messenger scale M (where new degrees of freedom should appear), and therefore compatible with the constraint arising from the gravitino mass limit (30).

B. Constraints on the scale of supersymmetry breaking from reheating

Reheating proceeds by coupling the inflaton to the MSSM sector. Since all superpartners are above the inflaton mass and reheating temperature, reheating produces predominantly SM particles (the abundance of gravitinos is discussed in Sec. III D). Radiative corrections with MSSM fields in loops correct the inflaton potential. In low-energy supersymmetry such corrections are tiny, since they are proportional to the scale of supersymmetry breaking [68]. In our case with high-scale supersymmetry breaking, there may be large radiative corrections that can spoil flatness of the inflaton potential. Such constraints can put upper limits on the superpartner masses and therefore on the gravitino mass.

For example, a direct coupling (through the gauge kinetic function) of the inflaton, t to gauge fields, $f \ni h_1 t/M_P$, would induce quadratic and quartic corrections of magnitude

$$\delta m^2 \sim \frac{h_1^2}{16\pi^2} M_{1/2}^2, \quad (32)$$

$$\delta \lambda \sim \frac{h_1^4}{16\pi^2} \frac{M_{1/2}^2}{M_P^2}, \quad (33)$$

which both place nontrivial bounds on the coupling h_1 . For reheating dominated decays to gauge bosons, this can be translated into a limit on the reheating temperature and eventually the gravitino abundance.

As we discuss in more detail in Sec. III D, reheating in this model proceeds via the gravitational coupling of the inflaton to two Higgs bosons. The coupling of the inflaton field, t to MSSM fields was derived in [45], and the relevant bosonic coupling is

³It is also possible that all additional scalars obtain Hubble scale masses during inflation, therefore avoiding this condition.

$$\begin{aligned} \mathcal{L}_{\text{eff}} \ni & \frac{\text{Re}T}{\sqrt{3}}(n_I + n_L - 3)W^{IL}\bar{W}_{LJ}\Phi_I\bar{\Phi}^J, \\ & \sim \mu^2 e\sqrt{\frac{3}{2}}(|h_u|^2 + |h_d|^2), \end{aligned} \quad (34)$$

where $n_{I,J}$ are modular weights⁴ of the superfields $\Phi_{I,J}$ and should be taken to be equal to one for untwisted Higgs fields. The coupling for the Higgs fields is then $\mu^2/\sqrt{3}M_P$, where μ is the MSSM Higgs mixing mass, which is now expected to be of order the scalar masses. The quadratic and quartic corrections in Eqs. (32) and (33) are found with the replacement $h_1 \rightarrow \mu^2/mM_P$. Requiring $\delta m^2 \ll m^2$ and $\delta\lambda \ll 10^{-14}$ sets a rough bound on $\mu/m \lesssim 10^2$ which we will see below is satisfied when μ is adjusted to give the correct gravitino relic density. However a more model-independent statement is that reheating sets constraints on inflaton couplings to MSSM fields. Once we fix such couplings, the requirement that quantum corrections do not spoil flatness of the inflaton potential generically set upper bounds on superpartner masses. Such bounds are dependent on the inflationary model and details of reheating, but they typically indicate that the scale of supersymmetry breaking should not be too much higher than the inflationary mass scale.

C. The Higgs mass and vacuum stability

The fact that the sfermion, gaugino and Higgsino masses are above 3×10^{13} GeV, leads to important implications for the Higgs boson mass and vacuum stability. It is well known that to obtain the 125 GeV Higgs mass for $\tan\beta \lesssim 50$, the maximum supersymmetry breaking scale is approximately 10^{10} GeV. However in Ref. [33], it was noted that the supersymmetry breaking scale can be increased to the GUT scale ($\sim 10^{16}$ GeV), for very large values of $\tan\beta \sim 200$. This assumes a degenerate superpartner spectrum (at \tilde{m}), with a bottom superpotential Yukawa coupling, $\hat{y}_b = y_b/\cos\beta$ such that $\hat{\alpha}_b = \hat{y}_b^2/(4\pi) \sim 0.5$. Even though this coupling is perturbative, a very close Landau pole develops at $\Lambda \sim 10\tilde{m}$.

Instead, Ref. [34] considered nondegenerate superpartner masses, and showed that GUT scale masses can in fact be accommodated for much smaller values of $\tan\beta$. In particular, assuming $\tilde{m} = m_{\tilde{Q}_{L,3}} = m_{\tilde{\tau}_R} = 10^{16}$ GeV and $\tan\beta = 1$, a scan of gaugino and first/second generation sfermion masses in the range $[\tilde{m}, 100\tilde{m}]$, and Higgsino masses in the range $[\frac{\tilde{m}}{100}, \tilde{m}]$ gives rise to the required threshold corrections of the Higgs quartic coupling.

The analysis in Ref. [34] suggests that our superpartner spectrum can give rise to similar threshold corrections needed for a 125 GeV Higgs mass. For concreteness we will consider a gaugino-mediated spectrum, and expect

⁴Note that the definition of modular weights in our paper is opposite in sign with respect to the standard convention.

similar qualitative features for a spectrum generated by a more strongly-coupled mediation mechanism. Identifying the right-handed stau mass with the scale $\tilde{m} = m_{\tilde{\tau}_R} \sim 5 \times 10^{13}$ GeV, will then determine the size of the remaining sfermion and gaugino masses. Assuming that the gaugino masses are generated at a scale $m < \mu_0 \leq 10^{16}$ GeV, the sfermion masses are then approximately given by

$$m_{\tilde{Q}}^2 \simeq \frac{1}{16\pi^2} \left(\frac{32}{3} g_3^2 + 6g_2^2 + \frac{2}{15} g_1^2 \right) M_{1/2}^2 \log \frac{\mu_0}{\tilde{m}}, \quad (35)$$

$$m_{\tilde{u}}^2 \simeq \frac{1}{16\pi^2} \left(\frac{32}{3} g_3^2 + \frac{32}{15} g_1^2 \right) M_{1/2}^2 \log \frac{\mu_0}{\tilde{m}}, \quad (36)$$

$$m_{\tilde{d}}^2 \simeq \frac{1}{16\pi^2} \left(\frac{32}{3} g_3^2 + \frac{8}{15} g_1^2 \right) M_{1/2}^2 \log \frac{\mu_0}{\tilde{m}}, \quad (37)$$

$$m_{\tilde{L}}^2 \simeq \frac{1}{16\pi^2} \left(6g_2^2 + \frac{6}{5} g_1^2 \right) M_{1/2}^2 \log \frac{\mu_0}{\tilde{m}}, \quad (38)$$

$$m_{\tilde{e}}^2 \simeq \frac{1}{16\pi^2} \left(\frac{24}{5} g_1^2 \right) M_{1/2}^2 \log \frac{\mu_0}{\tilde{m}}. \quad (39)$$

Other soft parameters such as A_t , $B\mu$ and $m_{H_u,d}^2$ are also generated radiatively. Using this approximate spectrum we can then compute the one-loop threshold corrections as given in Refs. [34,71]. The result is shown in Fig. 4. The contribution to the Higgs quartic coupling from our superpartner spectrum overlaps with the $\pm 3\sigma$ band of the λ_H coupling provided that $0.75 \lesssim \tan\beta \lesssim 1.34$.

However, note that the one-loop threshold correction matches at a *negative* value of the Higgs quartic coupling. This occurs because of our somewhat compressed




FIG. 4. The Higgs quartic coupling, λ_H at the matching scale $\tilde{m} = 5 \times 10^{13}$ GeV, as a function of $\tan\beta$. The solid blue line is the difference between the SM and tree-level SUSY values, with the dashed lines indicating the $\pm 3\sigma$ contours due to the error in the top-quark Yukawa coupling $y_t(m_t)$, where $m_t = 173.1 \pm 0.7$ GeV [72]. The green line shows the one-loop threshold correction arising from our superpartner spectrum.

spectrum, $m_{\tilde{Q}}/\tilde{m} \sim 1.9$, $M_{1/2}/\tilde{m} \sim 2$, $\mu/\tilde{m} \sim 1.5$, and the fact that the gaugino/Higgsinos contribute negatively, while the positive stop contribution is suppressed by mixing near $\tan\beta \sim 1$. A more detailed determination of the superpartner spectrum that gives different mass ratios may lead to a positive correction and then the matching could occur for positive values of λ_H .

Nevertheless even with a positive one-loop threshold correction from our superpartner spectrum, the value of the top quark Yukawa coupling, $y_t(m_t)$ would need to be near its -3σ extreme value that is allowed by the large uncertainty in the top-quark mass measurement. Otherwise one needs to rely on other threshold effects to stabilize the Higgs potential below the scale \tilde{m} . For instance this could be due to the inflaton coupling to the Higgs, $y_I \sim \frac{1}{\sqrt{3}} \frac{\mu^2}{m M_p} \sim 10^{-3}$. However, assuming that the supersymmetric and soft contributions to the inflaton mass are of the same order, this coupling causes a shift [73] in the Higgs quartic coupling by an amount given by $\delta\lambda_H \approx y_I^2 \sin^2 2\beta \leq y_I^2 \sim 10^{-6}$, which is negligibly small.

Alternatively, a heavy scalar singlet with a quartic coupling to the Higgs, could be introduced that is related to the generation of the neutrino masses [74].⁵ If this correction causes the Higgs quartic to be large and positive at the SUSY scale \tilde{m} , then a negative one-loop threshold correction would not be a problem, since it could then be absorbed by the SUSY tree-level contribution for sufficiently large $\tan\beta$. Nonetheless this does require an extra tuning in the model in order that this scalar singlet remains light (at an intermediate scale).

Another potential concern is that the required $\tan\beta$ values in Fig. 4 are near one, and normally in the MSSM this would cause a Landau pole in the top quark Yukawa coupling to appear below the GUT scale. However since our sparticle spectrum is quite heavy, the top quark Yukawa coupling is reduced by a factor of two at the scale $\tilde{m} \approx 5 \times 10^{13}$ GeV. The matching condition for the top Yukawa coupling $y_t = \hat{y}_t \sin\beta$, where \hat{y}_t is the superpotential Yukawa coupling, then allows for a lower value of $\tan\beta$, with a corresponding larger value of \hat{y}_t . Since there is relatively little running in our high-scale SUSY model above the scale \tilde{m} , the larger \hat{y}_t value can remain perturbative below the GUT scale.

Furthermore, to radiatively break electroweak symmetry requires that at some scale, $\frac{dm_{H_u}^2}{dt} = 0$, or $\hat{y}_t m_{\tilde{Q}} \sim g_2 M_2$. In the gaugino-mediated model this condition occurs when $\tan\beta \sim 0.5$, which is incompatible with the range required in Fig. 4.⁶ However if there were a new positive

contribution to the Higgs quartic coupling then it may be possible to also achieve radiative electroweak symmetry breaking. For instance, starting the running above the GUT scale 10^{16} GeV would give different sfermion mass ratios to make $\delta\lambda_H$ positive and/or allow $m_{H_u}^2$ to run negative before \tilde{m} . The details of these possibilities are beyond the scope of this paper and will be left for future work.

D. Dark matter

One of the main motivations of our high-scale SUSY model is its ability to account for the dark matter in the form of gravitinos with masses $m_{3/2} \gtrsim 0.2$ EeV. Because the supersymmetric particle spectrum lies above the inflaton mass, the dominant mechanism for gravitino production becomes SM + SM \rightarrow 2 gravitinos with longitudinal polarizations [9,10] or the decay of the inflaton directly to gravitinos depending on the reheating temperature.

The gravitino production rate was derived in [9]

$$R = n^2 \langle \sigma v \rangle \approx 21.65 \times \frac{T^{12}}{F^4}, \quad (40)$$

where n is the number density of incoming states. This temperature dependence can be understood as follows: one uses $n \propto T^3$, and we expect the gravitino production cross section to scale as $\langle \sigma v \rangle \propto T^6/F^4$. From the rate $R(T)$, we can determine that $\Gamma \sim R/n \sim T^9/M_P^4 m_{3/2}^4$ (assuming $m_{3/2} \ll \tilde{m}$) leading to a gravitino abundance $n_{3/2}/n_\gamma \sim \Gamma/H \sim T^7/M_P^3 m_{3/2}^4$ evaluated at $T = T_{RH}$ or

$$\Omega_{3/2} h^2 \approx 0.11 \left(\frac{0.1 \text{ EeV}}{m_{3/2}} \right)^3 \left(\frac{T_{RH}}{2.0 \times 10^{10} \text{ GeV}} \right)^7, \quad (41)$$

assuming instantaneous decay and thermalization. Thus, thermal production of gravitinos with $m_{3/2} > 0.2$ EeV would require $T_{RH} > 3 \times 10^{10}$ GeV.

It is known however, that the reheating process is not instantaneous, and that the temperature of the Universe during inflaton decay can exceed T_{RH} by orders of magnitude [32,76] up to a value T_{\max} . Due to the strong temperature dependence of the gravitino production cross section, there will be significant production of gravitinos at T_{\max} , which is not fully diluted by the entropy produced in subsequent decays. The final gravitino abundance in this case (with $\sigma \propto T^6$) relative to the instantaneous approximation is [77]

$$r_{3/2} = \frac{56}{5} \ln \left(\frac{T_{\max}}{T_{RH}} \right), \quad (42)$$

where

$$T_{\max} \approx 0.5 \left(\frac{m}{\Gamma_T} \right)^{1/4} T_{RH}, \quad (43)$$

⁵In [75], an additional $SU(2)_L$ triplet was introduced to insure stability and aid in keeping the Higgs mass at 125 GeV.

⁶In addition, requiring that there is no color-breaking minimum deeper than the electroweak minimum [71], leads to $\tan\beta \gtrsim 0.6$ for our spectrum.

for inflationary models of this type, where Γ_T is the total inflaton decay rate.

For the inflationary model discussed above, there are many possible decay channels all of which are Planck suppressed. The decay channel given in Eq. (34) is ordinarily (with weak scale supersymmetry breaking) negligible as $\mu^2/(mM_P) \ll 1$. However, in our case, since $\mu > m$, this is actually the dominant inflaton decay mode $t \rightarrow H_{u,d}H^{*d,u}$ which ultimately corresponds to a decay of $t \rightarrow hh$ where h is the SM Higgs boson. The decay rate to two Higgs bosons is [45]

$$\Gamma_{2h} = \frac{\mu^4}{384\pi m M_P^2} \sin^2 2\beta, \quad (44)$$

where an additional factor of 1/16 has been included in writing $H_u^0 = h/\sqrt{2} \sin \alpha$, and $H_d^0 = h/\sqrt{2} \cos \alpha$, and noting that $\alpha = \beta$ in the high scale SUSY limit.⁷

If we define an effective Yukawa-like coupling, $y_I = \mu^2/(4\sqrt{3}mM_P)$, such that $\Gamma_{2h} = \frac{y_I^2}{8\pi} m$, we can express the reheating temperature in terms of y_I [10,30,32]

$$T_{RH} = \left(\frac{10}{g_s}\right)^{1/4} \left(\frac{2\Gamma_{2h}M_P}{\pi c}\right)^{1/2} = 0.5 \frac{y_I}{2\pi} (mM_P)^{1/2}, \quad (45)$$

where g_s is the effective number of light degrees of freedom, in this case set by the standard model, $g_s = 427/4$ and $c \approx 1.2$ is a constant. We can then reexpress the relic abundance (41) as

$$\begin{aligned} \Omega_{3/2} h^2 &\simeq 0.11 r_{3/2} \left(\frac{0.1 \text{ EeV}}{m_{3/2}}\right)^3 \left(\frac{m}{3 \times 10^{13} \text{ GeV}}\right)^{7/2} \\ &\times \left(\frac{y_I}{2.9 \times 10^{-5}}\right)^7, \\ &= 0.11 r_{3/2} \left(\frac{0.1 \text{ EeV}}{m_{3/2}}\right)^3 \left(\frac{3 \times 10^{13} \text{ GeV}}{m}\right)^{7/2} \\ &\times \left(\frac{\mu}{1.2 \times 10^{14} \text{ GeV}}\right)^{14}, \end{aligned} \quad (46)$$

where we have included the enhancement factor $r_{3/2}$ from Eq. (42). The enhancement factor depends on $\ln \mu$, and for the range of μ values considered here (roughly $10^{14} - 10^{15}$ GeV), $r_{3/2}$ varies very little and we take it as a constant $r_{3/2} = 25$.

The value of μ needed to obtain the correct relic density of gravitinos is shown by the solid line in Fig. 5 using

⁷We note that parametric resonance effects such as those studied in [78] are not effective in this model. The adiabatic condition $\dot{m}_h/m_h^2 > 1$ is at best satisfied during the first inflaton oscillation, where m_h is the t -dependent Higgs mass. One can easily check that $\dot{m}_h/m_h^2 \sim Am/\mu M_P$, where $A \lesssim M_P$ is the amplitude of inflaton oscillations and we require $m < \mu$.




FIG. 5. The value of μ relative to the inflaton mass needed to obtain the correct relic density of gravitinos thermally through reheating (solid line) as a function of the gravitino mass. Also shown (dashed lines) are the values of μ needed to obtain the correct relic density of gravitinos through inflaton decays for a given value of $\log \Lambda_z/M_P$ as labeled.

Eq. (46). It is rather amazing that independent of the supersymmetric particle spectrum discussed above, the value of μ needed for the correct abundance of gravitinos is in the range of roughly 3-30 times the inflaton mass. This is exactly where one might expect the Higgsino mass to lie given our spectrum of heavy scalars and gauginos.

It is also possible that μ takes values below the solid line in Fig. 5. In that case, the abundance of gravitinos is below the needed relic density of dark matter (by the same token, values of μ above the solid line are excluded as they yield a relic density in excess of the observed one). Nevertheless, it is still possible to recover the correct relic density through inflaton decay to gravitinos. The gravitino abundance produced by inflaton decay for a given branching fraction to gravitinos, $B_{3/2} = \Gamma_{3/2}/\Gamma_{2h}$, was computed in [10]

$$\begin{aligned} \Omega_{3/2}^{\text{decay}} h^2 &= 0.11 \left(\frac{B_{3/2}}{1.3 \times 10^{-13}}\right) \left(\frac{y_I}{2.9 \times 10^{-5}}\right) \\ &\times \left(\frac{m_{3/2}}{0.1 \text{ EeV}}\right) \left(\frac{3 \times 10^{13} \text{ GeV}}{m}\right)^{1/2}. \end{aligned} \quad (47)$$

The decay of the inflaton to two gravitinos was computed in [45] with

$$\Gamma_{3/2} = \left(\frac{\Lambda_z}{M_P}\right)^4 \frac{3m_{3/2}^2 m}{256\pi M_P^2}, \quad (48)$$

so that

$$B_{3/2} = \frac{9}{2} \left(\frac{\Lambda_z}{M_P} \right)^4 \left(\frac{m_{3/2}}{m} \right)^2 \left(\frac{m}{\mu} \right)^4. \quad (49)$$

Using Eq. (48), inputting $B_{3/2}$ and y_I , we obtain the correct relic density of gravitinos along the sloped dashed and dotted lines for different values of Λ_z/M_P as labeled.

As one can see from Fig. 5, the value of the abundance of gravitinos from the decay of the inflaton is strongly dependent on the value of Λ_z as this scale controls the branching ratio $B_{3/2}$. As a consequence, we can derive an upper limit to Λ_z

$$\frac{\Lambda_z}{M_P} \leq 2.4 \times 10^{-4} \left(\frac{m}{m_{3/2}} \right)^{9/14}, \quad (50)$$

which is shown in Fig. 3 by the negatively sloped pink dot-dashed line. This upper limit in turn imposes an upper limit to the gravitino mass $m_{3/2} \lesssim 0.1m$.

IV. CONCLUSIONS

It may be that supersymmetry is not physically realized at energy scales accessible to the LHC. The hierarchy problem and naturalness biased our expectations that the supersymmetry mass scale was at or near the weak scale making experimental discovery all but inevitable. Such is not (yet) the case, and the mass scale of the supersymmetric spectrum remains unknown. It is therefore plausible to consider the possibility that nearly the entire supersymmetric spectrum lies at very high energies. If it is above the inflationary scale it is quite possible that the rich spectrum of supersymmetric partners were never produced in the early universe after inflationary reheating.

The exception could be the gravitino whose mass may remain below the inflaton mass. In this case, gravitinos could be produced (in pairs) during reheating [9,10] and because of the strong sensitivity to temperature, have enhanced production at the start of reheating when the temperature of the radiation plasma is above the reheating scale [77]. For a sufficiently high reheating temperature ($T_{RH} \gtrsim 10^{10}$ GeV), thermally produced gravitinos would have the correct relic density to account for the observed cold dark matter in the Universe.

In this paper, we have constructed a working model incorporating both inflation and supersymmetry breaking which leads to the heavy supersymmetric spectrum with gravitino dark matter. Our starting point is no-scale supergravity [36,37]. In the family of models formulated in Ref. [67], the inflaton is associated with the volume modulus, T , and the scalar potential is identical to that derived in the Starobinsky model [12]. Supersymmetry

breaking is achieved by adding a strongly stabilized Polonyi field which preserves the potential for inflation. Reheating occurs through the gravitational coupling of the inflaton to the standard model Higgs scalars. Because supersymmetry breaking occurs at a high scale, the μ parameter is large (larger than the inflaton mass) and the dominant decay channel is to two Higgs bosons. Depending on the value of μ , gravitinos may be produced through reheating with the correct relic density over a wide range of gravitino masses. For smaller values of μ (but still larger than the inflaton mass), the correct relic density may be obtained via the decays of the inflaton directly to gravitinos. All constraints are satisfied for gravitino masses in the range $0.1 \text{ EeV} \lesssim m_{3/2} \lesssim 1000 \text{ EeV}$.

While we have shown that the Higgs mass in this class of models can be compatible with the experimental value when $\tan\beta \approx 1$, we leave open for future study the questions of vacuum stability, and radiative electroweak symmetry breaking—two attractive features normally associated with supersymmetric models. Both stability and symmetry breaking can be achieved without supersymmetry via an intermediate scale such as in SO(10) grand unification [79], and therefore suggests that in models of high scale supersymmetry breaking, grand unification plays a crucial role in determining the Higgs mass.

Finally, the minimal setup of our model predicts no signatures in either collider or direct/indirect dark matter searches. Instead, scalars with masses near the Hubble scale during inflation could lead to non-Gaussianities in the CMB that may eventually be observed. Alternatively, by introducing R-parity violation in the lepton-Higgs sector, the gravitino can become unstable with a suitably long-lived decay. The detection of the decay products, such as very high energy neutrinos or photons at HAWC or the Pierre Auger Observatory, would then be a possible sign of high scale supersymmetry with an EeV gravitino, and thus these types of signatures are worthy of further study.

ACKNOWLEDGMENTS

We would like to thank Sebastian Ellis and Marco Peloso for helpful discussions. This work was supported by the France-US PICS No. 06482. Y. M. acknowledges partial support from the European Union Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie: RISE InvisiblesPlus (Grant No. 690575), the ITN Elusives (Grant No. 674896), and the ERC advanced grants Higgs@LHC. E. D. acknowledges partial support from the ANR Black-dS-String. The work of T. G. and K. A. O. was supported in part by the DOE Grant No. DE-SC0011842 at the University of Minnesota.

- [1] G. Aad *et al.* (ATLAS Collaboration), Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012); S. Chatrchyan *et al.* (CMS Collaboration), Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
- [2] G. Aad *et al.* (ATLAS and CMS Collaborations), Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments, *Phys. Rev. Lett.* **114**, 191803 (2015).
- [3] J. R. Ellis, G. Ridolfi, and F. Zwirner, Radiative corrections to the masses of supersymmetric Higgs bosons, *Phys. Lett. B* **257**, 83 (1991); On radiative corrections to supersymmetric Higgs boson masses and their implications for LEP searches, *Phys. Lett. B* **262**, 477 (1991); Y. Okada, M. Yamaguchi, and T. Yanagida, Upper bound of the lightest Higgs boson mass in the minimal supersymmetric standard model, *Prog. Theor. Phys.* **85**, 1 (1991); A. Yamada, Radiative corrections to the Higgs masses in the minimal supersymmetric standard model, *Phys. Lett. B* **263**, 233 (1991); Howard E. Haber and Ralf Hempfling, Can the Mass of the Lightest Higgs Boson of the Minimal Supersymmetric Model be Larger than m_z ?, *Phys. Rev. Lett.* **66**, 1815 (1991); M. Drees and M. M. Nojiri, One-loop corrections to the Higgs sector in minimal supergravity models, *Phys. Rev. D* **45**, 2482 (1992); P. H. Chankowski, S. Pokorski, and J. Rosiek, Charged and neutral supersymmetric Higgs boson masses. Complete one-loop analysis, *Phys. Lett. B* **274**, 191 (1992); One-loop corrections to the supersymmetric Higgs boson couplings and LEP phenomenology, *Phys. Lett. B* **286**, 307 (1992).
- [4] G. Degrandi, P. Slavich, and F. Zwirner, On the neutral Higgs boson masses in the MSSM for arbitrary stop mixing, *Nucl. Phys. B* **611**, 403 (2001); A. Brignole, G. Degrandi, P. Slavich, and F. Zwirner, On the $O(\alpha_t^2)$ two-loop corrections to the neutral Higgs boson masses in the MSSM, *Nucl. Phys. B* **631**, 195 (2002); A. Brignole, G. Degrandi, P. Slavich, and F. Zwirner, On the two-loop sbottom corrections to the neutral Higgs boson masses in the MSSM, *Nucl. Phys. B* **643**, 79 (2002).
- [5] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich, and G. Weiglein, Towards high-precision predictions for the MSSM Higgs sector, *Eur. Phys. J. C* **28**, 133 (2003).
- [6] B. C. Allanach, A. Djouadi, J. L. Kneur, W. Porod, and P. Slavich, Precise determination of the neutral Higgs boson masses in the MSSM, *J. High Energy Phys.* **09** (2004) 044.
- [7] J. R. Ellis, D. V. Nanopoulos, K. A. Olive, and Y. Santoso, On the Higgs mass in the CMSSM, *Phys. Lett. B* **633**, 583 (2006).
- [8] G. Aad *et al.* (ATLAS Collaboration), Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton-proton collision data, *J. High Energy Phys.* **09** (2014) 176; G. Aad *et al.* (ATLAS Collaboration), Summary of the searches for squarks and gluinos using $\sqrt{s} = 8$ TeV pp collisions with the ATLAS experiment at the LHC, *J. High Energy Phys.* **10** (2015) 054; S. Chatrchyan *et al.* (CMS Collaboration), Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 8$ TeV, *J. High Energy Phys.* **06** (2014) 055.
- [9] K. Benakli, Y. Chen, E. Dudas, and Y. Mambrini, A minimal model of gravitino dark matter, *Phys. Rev. D* **95**, 095002 (2017).
- [10] E. Dudas, Y. Mambrini, and K. Olive, Case for an EeV Gravitino, *Phys. Rev. Lett.* **119**, 051801 (2017).
- [11] D. V. Volkov and V. P. Akulov, Is the neutrino a Goldstone particle?, *Phys. Lett.* **46B**, 109 (1973); For applications of nonlinear supersymmetry to inflation, see e.g. I. Antoniadis, E. Dudas, S. Ferrara, and A. Sagnotti, The Volkov–Akulov–Starobinsky supergravity, *Phys. Lett. B* **733**, 32 (2014); S. Ferrara, R. Kallosh, and A. Linde, Cosmology with nilpotent superfields, *J. High Energy Phys.* **10** (2014) 143; G. Dall’Agata and F. Zwirner, On sgoldstino-less supergravity models of inflation, *J. High Energy Phys.* **12** (2014) 172.
- [12] A. A. Starobinsky, A new type of isotropic cosmological models without singularity, *Phys. Lett.* **91B**, 99 (1980).
- [13] V. F. Mukhanov and G. V. Chibisov, Pis’ma Zh. Eksp. Teor. Fiz. **33**, 549 (1981) [Quantum fluctuation and nonsingular universe. (In Russian), *JETP Lett.* **33**, 532 (1981)].
- [14] A. A. Starobinsky, The perturbation spectrum evolving from a nonsingular initially de-Sitter cosmology and the microwave background anisotropy, *Sov. Astron. Lett.* **9**, 302 (1983).
- [15] P. A. R. Ade *et al.* (Planck Collaboration), Planck 2015 results. XIII. Cosmological parameters, *Astron. Astrophys.* **594**, A13 (2016); Planck 2015 results. XX. Constraints on inflation, *Astron. Astrophys.* **594**, A20 (2016).
- [16] P. Minkowski, $\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?, *Phys. Lett.* **67B**, 421 (1977); T. Yanagida, Horizontal symmetry and masses of neutrinos, *Conf. Proc. C* **7902131**, 95 (1979); M. Gell-Mann, P. Ramond, and R. Slansky, Complex spinors and unified theories, *Conf. Proc. C* **790927**, 315 (1979); S. L. Glashow, The future of elementary particle physics, *NATO Sci. Ser. B* **59**, 687 (1980); R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Violation, *Phys. Rev. Lett.* **44**, 912 (1980); R. N. Mohapatra and G. Senjanovic, Neutrino masses and mixings in gauge models with spontaneous parity violation, *Phys. Rev. D* **23**, 165 (1981); J. Schechter and J. W. F. Valle, Neutrino masses in $SU(2) \times U(1)$ theories, *Phys. Rev. D* **22**, 2227 (1980); J. Schechter and J. W. F. Valle, Neutrino decay and spontaneous violation of lepton number, *Phys. Rev. D* **25**, 774 (1982).
- [17] J. Polonyi, Budapest Report No. KFKI-1977-93, 1977.
- [18] D. V. Nanopoulos, K. A. Olive, and M. Srednicki, After primordial inflation, *Phys. Lett.* **127B**, 30 (1983).
- [19] J. Ellis, J. Hagelin, D. Nanopoulos, K. Olive, and M. Srednicki, Supersymmetric relics from the big bang, *Nucl. Phys. B* **238**, 453 (1984).
- [20] M. Y. Khlopov and A. D. Linde, Is it easy to save the gravitino?, *Phys. Lett.* **138B**, 265 (1984).
- [21] J. R. Ellis, J. E. Kim, and D. V. Nanopoulos, Cosmological gravitino regeneration and decay, *Phys. Lett.* **145B**, 181 (1984).
- [22] R. Juszkiewicz, J. Silk, and A. Stebbins, Constraints on cosmologically regenerated gravitinos, *Phys. Lett.* **158B**, 463 (1985).

- [23] T. Moroi, H. Murayama, and M. Yamaguchi, Cosmological constraints on the light stable gravitino, *Phys. Lett. B* **303**, 289 (1993).
- [24] M. Kawasaki and T. Moroi, Gravitino production in the inflationary universe and the effects on big bang nucleosynthesis, *Prog. Theor. Phys.* **93**, 879 (1995).
- [25] T. Moroi, Effects of the gravitino on the inflationary universe, [arXiv:hep-ph/9503210](https://arxiv.org/abs/hep-ph/9503210).
- [26] J. R. Ellis, D. V. Nanopoulos, K. A. Olive, and S. J. Rey, On the thermal regeneration rate for light gravitinos in the early universe, *Astropart. Phys.* **4**, 371 (1996).
- [27] G. F. Giudice, A. Riotto, and I. Tkachev, Thermal and nonthermal production of gravitinos in the early universe, *J. High Energy Phys.* **11** (1999) 036.
- [28] M. Bolz, A. Brandenburg, and W. Buchmuller, Thermal production of gravitinos, *Nucl. Phys.* **B606**, 518 (2001); Erratum, *Nucl. Phys.* **B790**, 336 (2008).
- [29] J. Pradler and F. D. Steffen, Thermal gravitino production and collider tests of leptogenesis, *Phys. Rev. D* **75**, 023509 (2007).
- [30] J. Pradler and F. D. Steffen, Constraints on the reheating temperature in gravitino dark matter scenarios, *Phys. Lett. B* **648**, 224 (2007).
- [31] V. S. Rychkov and A. Strumia, Thermal production of gravitinos, *Phys. Rev. D* **75**, 075011 (2007).
- [32] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, K. A. Olive, and M. Peloso, Post-inflationary gravitino production revisited, *J. Cosmol. Astropart. Phys.* **03** (2016) 008.
- [33] J. P. Vega and G. Villadoro, SusyHD: Higgs mass determination in supersymmetry, *J. High Energy Phys.* **07** (2015) 159.
- [34] S. A. R. Ellis and J. D. Wells, High-scale supersymmetry, the Higgs mass and gauge unification, *Phys. Rev. D* **96**, 055024 (2017).
- [35] N. Arkani-Hamed and J. Maldacena, Cosmological collider physics, [arXiv:1503.08043](https://arxiv.org/abs/1503.08043).
- [36] E. Cremmer, S. Ferrara, C. Kounnas, and D. V. Nanopoulos, Naturally vanishing cosmological constant in $N = 1$ supergravity, *Phys. Lett.* **133B** (1983) 61; J. R. Ellis, A. B. Lahanas, D. V. Nanopoulos, and K. Tamvakis, No-scale supersymmetric standard model, *Phys. Lett.* **134B** (1984) 429.
- [37] A. B. Lahanas and D. V. Nanopoulos, The road to no scale supergravity, *Phys. Rep.* **145**, 1 (1987).
- [38] J. Ellis, D. V. Nanopoulos, and K. A. Olive, No-Scale Supergravity Realization of the Starobinsky Model of Inflation, *Phys. Rev. Lett.* **111**, 111301 (2013).
- [39] R. Kallosh and A. Linde, Superconformal generalizations of the Starobinsky model, *J. Cosmol. Astropart. Phys.* **06** (2013) 028.
- [40] F. Farakos, A. Kehagias, and A. Riotto, On the Starobinsky model of inflation from supergravity, *Nucl. Phys.* **B876**, 187 (2013).
- [41] J. Ellis, D. V. Nanopoulos, and K. A. Olive, Starobinsky-like inflationary models as avatars of no-scale supergravity, *J. Cosmol. Astropart. Phys.* **10** (2013) 009.
- [42] J. Ellis, D. V. Nanopoulos, and K. A. Olive, A no-scale framework for sub-Planckian physics, *Phys. Rev. D* **89**, 043502 (2014).
- [43] I. Antoniadis, E. Dudas, S. Ferrara, and A. Sagnotti, The Volkov-Akulov-Starobinsky supergravity, *Phys. Lett. B* **733**, 32 (2014).
- [44] D. Roest, M. Scalisi, and I. Zavala, Kahler potentials for Planck inflation, *J. Cosmol. Astropart. Phys.* **11** (2013) 007; S. Ferrara, R. Kallosh, A. Linde, and M. Porrati, Minimal supergravity models of inflation, *Phys. Rev. D* **88** (2013) 085038; W. Buchmüller, V. Domcke, and C. Wieck, No-scale D-term inflation with stabilized moduli, *Phys. Lett. B* **730**, 155 (2014); C. Pallis, Linking Starobinsky-type inflation in no-scale supergravity to MSSM, *J. Cosmol. Astropart. Phys.* **04** (2014) 024; C. Pallis, Induced-gravity inflation in no-scale supergravity and beyond, *J. Cosmol. Astropart. Phys.* **08** (2014) 057; S. Ferrara, A. Kehagias, and A. Riotto, The imaginary Starobinsky model, *Fortsch. Phys.* **62**, 573 (2014); S. Ferrara, A. Kehagias, and A. Riotto, The imaginary Starobinsky model and higher curvature corrections, *Fortsch. Phys.* **63**, 2 (2015); R. Kallosh, A. Linde, B. Vercnocke, and W. Chemissany, Is imaginary Starobinsky model real?, *J. Cosmol. Astropart. Phys.* **07** (2014) 053; K. Hamaguchi, T. Moroi, and T. Terada, Complexified Starobinsky inflation in supergravity in the light of recent BICEP2 result, *Phys. Lett. B* **733**, 305 (2014); J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, Resurrecting quadratic inflation in no-scale supergravity in light of BICEP2, *J. Cosmol. Astropart. Phys.* **05** (2014) 037; J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, A no-scale inflationary model to fit them all, *J. Cosmol. Astropart. Phys.* **08** (2014) 044; J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, Two-field analysis of no-scale supergravity inflation, *J. Cosmol. Astropart. Phys.* **01** (2015) 010; W. Buchmuller, E. Dudas, L. Heurtier, A. Westphal, C. Wieck, and M. W. Winkler, Challenges for large-field inflation and moduli stabilization, *J. High Energy Phys.* **04** (2015) 058; C. Kounnas, D. Lust, and N. Toumbas, R^2 inflation from scale invariant supergravity and anomaly free superstrings with fluxes, *Fortsch. Phys.* **63**, 12 (2015); T. Terada, Y. Watanabe, Y. Yamada, and J. Yokoyama, Reheating processes after Starobinsky inflation in old-minimal supergravity, *J. High Energy Phys.* **02** (2015) 105; A. B. Lahanas and K. Tamvakis, Inflation in no-scale supergravity, *Phys. Rev. D* **91**, 085001 (2015); I. Dalianis and F. Farakos, On the initial conditions for inflation with plateau potentials: the $R + R^2$ (super)gravity case, *J. Cosmol. Astropart. Phys.* **07** (2015) 044; D. Roest and M. Scalisi, Cosmological attractors from -scale supergravity, *Phys. Rev. D* **92**, 043525 (2015); I. Garg and S. Mohanty, No scale SUGRA SO(10) derived Starobinsky model of inflation, *Phys. Lett. B* **751**, 7 (2015); M. Scalisi, Cosmological α -attractors and de Sitter landscape, *J. High Energy Phys.* **12** (2015) 134; S. Ferrara, A. Kehagias, and M. Porrati, \mathcal{R}^2 supergravity, *J. High Energy Phys.* **08** (2015) 001; G. D. Diamandis, A. B. Lahanas, and K. Tamvakis, Towards a formulation of $f(R)$ supergravity, *Phys. Rev. D* **92**, 105023 (2015); C. Pallis and N. Toumbas, Starobinsky inflation: From non-SUSY to SUGRA realizations, *Adv. High Energy Phys.* **2017**, 6759267 (2017); T. Kobayashi, O. Seto, and T. H. Tatsuishi, Pole inflation and attractors in supergravity, [arXiv:1703.09960](https://arxiv.org/abs/1703.09960).
- [45] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, Phenomenological aspects of no-scale inflation models, *J. Cosmol. Astropart. Phys.* **10** (2015) 003.

- [46] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, Calculations of inflaton decays and reheating: With applications to no-scale inflation models, *J. Cosmol. Astropart. Phys.* **07** (2015) 050.
- [47] E. Dudas and C. Wieck, Moduli backreaction and supersymmetry breaking in string-inspired inflation models, *J. High Energy Phys.* **10** (2015) 062; W. Buchmuller, E. Dudas, L. Heurtier, A. Westphal, C. Wieck, and M. W. Winkler, Challenges for large-field inflation and moduli stabilization, *J. High Energy Phys.* **04** (2015) 058; W. Buchmuller, E. Dudas, L. Heurtier, and C. Wieck, Large-field inflation and supersymmetry breaking, *J. High Energy Phys.* **09** (2014) 053.
- [48] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, and K. A. Olive, No-scale inflation, *Classical Quantum Gravity* **33**, 094001 (2016).
- [49] J. Ellis, M. A. G. Garcia, N. Nagata, D. V. Nanopoulos, and K. A. Olive, Starobinsky-like inflation and neutrino masses in a no-scale SO(10) model, *J. Cosmol. Astropart. Phys.* **11** (2016) 018.
- [50] M. C. Romao and S. F. King, Starobinsky-like inflation in no-scale supergravity Wess-Zumino model with Polonyi term, *J. High Energy Phys.* **07** (2017) 033.
- [51] J. Ellis, M. A. G. Garcia, N. Nagata, D. V. Nanopoulos, and K. A. Olive, Starobinsky-like inflation, supercosmology and neutrino masses in no-scale flipped SU(5), *J. Cosmol. Astropart. Phys.* **07** (2017) 006.
- [52] K. Schmitz and T. T. Yanagida, Dynamical supersymmetry breaking and late-time R symmetry breaking as the origin of cosmic inflation, *Phys. Rev. D* **94**, 074021 (2016).
- [53] V. Domcke and K. Schmitz, Unified model of D-term inflation, *Phys. Rev. D* **95**, 075020 (2017).
- [54] M. Dine, R. Kitano, A. Morisse, and Y. Shirman, Moduli decays and gravitinos, *Phys. Rev. D* **73**, 123518 (2006); R. Kitano, Gravitational gauge mediation, *Phys. Lett. B* **641**, 203 (2006); R. Kallosh and A. D. Linde, O'kklt, *J. High Energy Phys.* **02** (2007) 002; H. Abe, T. Higaki, and T. Kobayashi, More about F-term uplifting, *Phys. Rev. D* **76**, 105003 (2007); J. Fan, M. Reece, and L.-T. Wang, Mitigating moduli messes in low-scale SUSY breaking, *J. High Energy Phys.* **09** (2011) 126.
- [55] E. Dudas, C. Papineau, and S. Pokorski, Moduli stabilization and uplifting with dynamically generated F-terms, *J. High Energy Phys.* **02** (2007) 028; H. Abe, T. Higaki, T. Kobayashi, and Y. Omura, Moduli stabilization, F-term uplifting and soft supersymmetry breaking terms, *Phys. Rev. D* **75**, 025019 (2007).
- [56] R. Kallosh, A. Linde, K. A. Olive, and T. Rube, Chaotic inflation and supersymmetry breaking, *Phys. Rev. D* **84**, 083519 (2011); A. Linde, Y. Mambrini, and K. A. Olive, Supersymmetry breaking due to moduli stabilization in string theory, *Phys. Rev. D* **85**, 066005 (2012).
- [57] E. Dudas, A. Linde, Y. Mambrini, A. Mustafayev, and K. A. Olive, Strong moduli stabilization and phenomenology, *Eur. Phys. J. C* **73**, 2268 (2013).
- [58] J. L. Evans, M. Ibe, K. A. Olive, and T. T. Yanagida, Universality in pure gravity mediation, *Eur. Phys. J. C* **73**, 2468 (2013); J. L. Evans, K. A. Olive, M. Ibe, and T. T. Yanagida, Non-universalities in pure gravity mediation, *Eur. Phys. J. C* **73**, 2611 (2013).
- [59] K. Nakayama, F. Takahashi, and T. T. Yanagida, Eluding the gravitino overproduction in inflaton decay, *Phys. Lett. B* **718**, 526 (2012).
- [60] M. A. G. Garcia and K. A. Olive, Affleck-Dine baryogenesis and inflation in supergravity with strongly stabilized moduli, *J. Cosmol. Astropart. Phys.* **09** (2013) 007.
- [61] J. L. Evans, M. A. G. Garcia, and K. A. Olive, The moduli and gravitino (non)-problems in models with strongly stabilized moduli, *J. Cosmol. Astropart. Phys.* **03** (2014) 022.
- [62] Z. Chacko, M. A. Luty, A. E. Nelson, and E. Ponton, Gaugino mediated supersymmetry breaking, *J. High Energy Phys.* **01** (2000) 003; D. E. Kaplan, G. D. Kribs, and M. Schmaltz, Supersymmetry breaking through transparent extra dimensions, *Phys. Rev. D* **62**, 035010 (2000).
- [63] T. Gherghetta and A. Pomarol, A warped supersymmetric standard model, *Nucl. Phys.* **B602**, 3 (2001); S. Abel and T. Gherghetta, A slice of AdS₅ as the large N limit of Seiberg duality, *J. High Energy Phys.* **12** (2010) 091.
- [64] P. Meade, N. Seiberg, and D. Shih, General gauge mediation, *Prog. Theor. Phys. Suppl.* **177**, 143 (2009); M. Buican, P. Meade, N. Seiberg, and D. Shih, Exploring general gauge mediation, *J. High Energy Phys.* **03** (2009) 016.
- [65] J. R. Ellis, C. Kounnas, and D. V. Nanopoulos, No scale supersymmetric GUTs, *Nucl. Phys.* **B247**, 373 (1984).
- [66] J. R. Ellis, C. Kounnas, and D. V. Nanopoulos, No scale supergravity models with a Planck mass gravitino, *Phys. Lett.* **143B**, 410 (1984).
- [67] S. Cecotti, Higher derivative supergravity is equivalent to standard supergravity coupled to matter. I., *Phys. Lett. B* **190**, 86 (1987).
- [68] J. R. Ellis, D. V. Nanopoulos, K. A. Olive, and K. Tamvakis, Cosmological inflation cries out for supersymmetry, *Phys. Lett.* **118B**, 335 (1982); Fluctuations in a supersymmetric inflationary universe, *Phys. Lett.* **120B**, 331 (1983); Primordial supersymmetric inflation, *Nucl. Phys.* **B221**, 524 (1983); K. Nakayama and F. Takahashi, Low-scale supersymmetry from inflation, *J. Cosmol. Astropart. Phys.* **10** (2011) 033.
- [69] K. Choi, A. Falkowski, H. P. Nilles, and M. Olechowski, Soft supersymmetry breaking in KKL_T flux compactification, *Nucl. Phys.* **B718**, 113 (2005); O. Lebedev, H. P. Nilles, and M. Ratz, De Sitter vacua from matter superpotentials, *Phys. Lett. B* **636**, 126 (2006); O. Lebedev, V. Lowen, Y. Mambrini, H. P. Nilles, and M. Ratz, Metastable vacua in flux compactifications and their phenomenology, *J. High Energy Phys.* **02** (2007) 063.
- [70] T. Bhattacharya and P. Roy, Unitarity limit on the gaugino-gravitino mass ratio, *Phys. Lett. B* **206**, 655 (1988); Tree unitarity in broken supergravity. 1. Single gravitino amplitude, *Nucl. Phys.* **B328**, 469 (1989).
- [71] E. Bagnaschi, G. F. Giudice, P. Slavich, and A. Strumia, Higgs mass and unnatural supersymmetry, *J. High Energy Phys.* **09** (2014) 092.
- [72] G. Degrandi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori, and A. Strumia, Higgs mass and vacuum stability in the standard model at NNLO, *J. High Energy Phys.* **08** (2012) 098.
- [73] G. F. Giudice and A. Strumia, Probing high-scale and split supersymmetry with Higgs mass measurements, *Nucl. Phys.* **B858**, 63 (2012).

- [74] J. Elias-Miro, J.R. Espinosa, G.F. Giudice, H.M. Lee, and A. Strumia, Stabilization of the electroweak vacuum by a scalar threshold effect, *J. High Energy Phys.* **06** (2012) 031.
- [75] C. Liu and Z. h. Zhao, θ_{13} and the Higgs mass from high scale supersymmetry, *Commun. Theor. Phys.* **59**, 467 (2013).
- [76] D. J. H. Chung, E. W. Kolb, and A. Riotto, Production of massive particles during reheating, *Phys. Rev. D* **60**, 063504 (1999); G. F. Giudice, E. W. Kolb, and A. Riotto, Largest temperature of the radiation era and its cosmological implications, *Phys. Rev. D* **64** (2001) 023508; E. W. Kolb, A. Notari, and A. Riotto, On the reheating stage after inflation, *Phys. Rev. D* **68**, 123505 (2003).
- [77] M. A. G. Garcia, Y. Mambrini, K. A. Olive, and M. Peloso, Enhancement of the dark matter abundance before reheating: Applications to gravitino dark matter, *Phys. Rev. D* **96**, 103510 (2017).
- [78] L. Kofman, A. D. Linde, and A. A. Starobinsky, Towards the theory of reheating after inflation, *Phys. Rev. D* **56**, 3258 (1997).
- [79] Y. Mambrini, N. Nagata, K. A. Olive, and J. Zheng, Vacuum stability and radiative electroweak symmetry breaking in an SO(10) dark matter model, *Phys. Rev. D* **93**, 111703 (2016).