

## Black holes as quantum gravity condensates

Daniele Oriti,<sup>1,\*</sup> Daniele Pranzetti,<sup>2,†</sup> and Lorenzo Sindoni<sup>3,‡</sup>

<sup>1</sup>*Max Planck Institute for Gravitational Physics (AEI), Am Mühlenberg 1, D-14476 Golm, Germany*

<sup>2</sup>*Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario N2L 2Y5, Canada*

<sup>3</sup>*InveniaLabs, Parkside Place, Parkside, CBI 1HQ Cambridge, United Kingdom*



(Received 15 January 2018; published 19 March 2018)

We model spherically symmetric black holes within the group field theory formalism for quantum gravity via generalized condensate states, involving sums over arbitrarily refined graphs (dual to three-dimensional triangulations). The construction relies heavily on both the combinatorial tools of random tensor models and the quantum geometric data of loop quantum gravity, both part of the group field theory formalism. Armed with the detailed microscopic structure, we compute the entropy associated with the black hole horizon, which turns out to be equivalently the Boltzmann entropy of its microscopic degrees of freedom and the entanglement entropy between the inside and outside regions. We recover the area law under very general conditions, as well as the Bekenstein-Hawking formula. The result is also shown to be generically independent of any specific value of the Immirzi parameter.

DOI: [10.1103/PhysRevD.97.066017](https://doi.org/10.1103/PhysRevD.97.066017)

### I. INTRODUCTION

The physics of black holes is a crucial part of the current frontier of theoretical physics. Beside their importance for theoretical (and observational) astrophysics [1,2], they offer a precious laboratory for testing the foundations of our theories of fundamental interactions and of all other physical systems, and their mutual (in)compatibility: general relativity, quantum mechanics (and its offspring, quantum information), and statistical mechanics. On the one hand, black hole thermodynamics [3–7] (see also [8] for a recent review and references covering the various developments) remains to date a surprising set of (theoretical) facts, still in search for a microscopic (statistical) understanding. It constantly challenges our assumed foundations of physical theories, at least when framed within semiclassical physics: locality, conservation of information, the equivalence principle, Lorentz invariance, quantum monogamy, causality, to name only a few of the principles that have been suggested to be somehow in conflict with it. On the other hand, together with the cosmology of the very early Universe, black holes (and their thermodynamics) are especially important for quantum gravity research, representing the main concrete challenge and the first testing ground at the nonperturbative level. Quantum gravity is supposed to complete the very definition of what black holes are, by providing the new physics replacing their central curvature singularity, and to identify

the microscopic degrees of freedom whose statistical mechanics is ultimately responsible for their macroscopic thermodynamical properties.

In this paper, we tackle the issue of defining quantum states representing (spherically symmetric) black holes, and in particular their horizon degrees of freedom, within a full quantum gravity formalism, and to compute their macroscopic entropy from first principles, within the same formalism. The quantum gravity framework we use is group field theory (GFT) [9,10], closely related to random tensor models [11] and to loop quantum gravity (LQG) [12,13]. To clarify what is the new contribution we give here to this topic, it is in fact useful to relate and compare our work to that done so far within loop quantum gravity, forming by now an extensive literature.

In the canonical LQG approach to black hole entropy calculation, the main strategy to the computation of black hole entropy has been based on classical symmetry reduction. The starting point for modeling the black hole horizon is the local definition provided by the notion of isolated horizon (IH) [14,15] (see [16,17] for reviews), which enters as a specific set of boundary conditions. Their implication is that a spherically symmetric isolated horizon appears at any  $r_0$  for which the following classical geometrical relation holds [18,19],

$$C_{IH} \equiv F^i(A) + \frac{\pi}{\mathcal{A}_{IH}} (1 - \gamma^2) \Sigma^i = 0. \quad (1)$$

In the expression above,  $\mathcal{A}_{IH}$  is the area of the isolated horizon,  $F^i(A)$  the field strength of the Ashtekar–Barbero connection  $A_a^i = \Gamma_a^i + \gamma K_a^i$  on the IH, and  $\Sigma^i \equiv \epsilon^i{}_{jk} \Sigma^{jk}$  the

\*daniele.oriti@aei.mpg.de

†dpranzetti@perimeterinstitute.ca

‡lorenzo.sindoni@invenialabs.co.uk

2-form densitized triad pulled back on the horizon. Spherically symmetric geometries constrained by such boundary conditions are then taken to *define* the classical degrees of freedom of the black hole system, which is then quantized.

In the quantum theory, the IH condition (1) induces a relation between the flux associated to a link coming from the bulk and ending on the horizon and the holonomy on the horizon around the given link. This is imposed as an operatorial constraint equation on the tensor product of the bulk Hilbert space, quantized through standard LQG techniques, and the boundary Hilbert space, constructed by relying on Chern-Simons theory formalism, which also defines the boundary dynamics. As a result, fluctuations of geometrical operators coming from the bulk get coupled to those living on the boundary, through identification of quantum numbers of the two Hilbert spaces, and Chern-Simons curvature excitations (punctures) get identified with quanta of space degrees of freedom. The microcanonical counting of these degrees of freedom à la Boltzmann in the semiclassical limit of large IH area yields a leading term for the entropy linear in  $\mathcal{A}_{IH}$  together with a subleading logarithmic term [20–29]. Extension of the SU(2)-invariant formulation of isolated horizons to the distorted and the rotating cases has been achieved in [30,31].

The realization that, in this classically reduced context, the relevant degrees of freedom are associated to punctures on the horizon (described by a simple Chern-Simons theory) combined with the general key result obtained in the full LQG theory that area operator has discrete spectrum, with spin networks as eigenstates and eigenvalues carried by the links of their supporting graphs, has motivated another well-explored strategy (in fact, the first one to be followed [32–34]). This was the construction of several toy models within LQG, i.e., simple spin network states incorporating enough features of the mentioned description of quantized isolated horizons to have a chance to capture interesting black hole physics. Most of such toy models consist of spin-network states based on a single fixed graph, interpreted as having a number of nodes inside a black hole horizon (encoding the bulk degrees of freedom), but often limited to a single intertwiner state, and a number of links crossing it, providing for the horizon degrees of freedom.

Both the mentioned strategies have produced very important and interesting results, and will certainly contribute to the complete understanding of the physics of quantum black holes in quantum gravity. The reliability of the results obtained within a symmetry reduced treatment is notoriously questionable, however, and the limitations of both strategies are apparent, motivating the search for a more complete description of quantum black holes within the full quantum gravity formalism.

Moreover, in all the LQG-based constructions so far, the numerical value of the Barbero-Immirzi parameter needs to

be fixed to recover exactly the coefficient 1/4 in the Bekenstein-Hawking entropy area formula. The long-standing issue, extensively debated in the literature [35–43], is whether this necessity might be signalling an incompleteness in the identification of the microscopic degrees of freedom counted in the LQG entropy calculation, or some other limitation in the usual constructions. On the basis of the semiclassical nature of the Bekenstein-Hawking formula and the expectation that the Barbero-Immirzi parameter should play no important role in the classical description of gravity, a natural option to remove this undesired feature of the entropy calculation is that other (noninternal and so far neglected) degrees of freedom should be taken into account. A recent proposal within the LQG framework, discussed in [44], uses new boundary degrees of freedom representing information channels between gravitational subsystems. However, the implications of the discovery of these new degrees of freedom for the black hole entropy calculation have not been investigated in detail yet.

In this paper, we concentrate on an alternative proposal for the fundamental boundary (and bulk) degrees of freedom, working within the full quantum gravity formalism and using a construction that gives us access to a continuum description of a black hole quantum geometry. More precisely, we want to use the construction [45] of GFT condensate states, generalizing those used in a cosmological context [46–49], and representing continuum spherically symmetric geometries within the GFT formalism. Leveraging on the structure of the Fock representation of GFTs, it is possible to define a quantum black hole horizon in the full, nontruncated, theory and then compute its entropy. A short report of some of these results appeared in [50]. Here we provide a much more detailed presentation of those calculations, as well as a more general entropy counting which extends to a wider and possibly more physically relevant class of generalized condensates for a spherically symmetric black hole.

We describe in some detail the construction of the class of states that our analysis relies on, and highlight their convenient features in the following; we note up front, however, the main limitation of our construction. Implementing the dynamics for generic GFT condensates is not as easy as in the case of those used for cosmological applications [46–49]. Therefore we treat these states as kinematical trial states, under the hypothesis that they can represent some reasonable approximation of realistic states, i.e., solutions of the full quantum dynamics, at least in some regime.

This remains a hypothesis, but it is supported by three considerations. First, these states naturally include some form of homogeneity (specifically, wave function homogeneity, to be clarified in the following) which already restricts the possible shape of the states in combination with the combinatorial restrictions that allow one to assign them

a clear topological interpretation (in particular, as encoding spherical symmetry). This sort of restrictions is expected to apply also to an exact state (i.e., solving the equations of motion of the theory) with spherical symmetry, or some slightly less local variant (e.g. involving vertices along tangential loops), resulting from constraints on curvature observables. Second, very importantly (and for the first time, to the best of our knowledge), the quantum states we use already include a sum over a family of triangulations, which is an inevitable consequence of the interacting nature of the GFT equations of motion. It is not obvious that the superposition of different graphs realized in physical states is of the same melonic nature that we are using, even if the dominance of melonic diagrams is a recurrent property of GFT (and random tensor) models [51]; however, the superposition that we are using explores a whole family of triangulations obtained by refinement, which means that, while differing in some crucial aspects, we might be still close to the exact state (indeed, for general wave functions, states associated to different graphs are *not* orthogonal [10]). While the calculation of the overlap between our condensate states and an exact spherically symmetric state (assuming this notion makes sense in the full theory) is nontrivial, we can expect that this overlap is nonzero.<sup>1</sup> Third, these particular states implement a nonobvious macroscopic property, i.e., holography, in a form compatible with a nonperturbative full quantum gravity regime. This means that they contain some of the structural properties necessary to match the dynamics of the classical theory.

Still, it remains true that our analysis stays at the kinematical level (as in all the previous LQG literature on the subject). We use, however, a possible proxy for the dynamics of the theory, because we impose a condition of maximization of the entropy that one could expect to be satisfied by solutions of the quantum dynamics and, in particular, by black hole configurations. The (maximal) entropy we compute for our states is shown to be interpretable both as a Boltzmann entropy of horizon degrees of freedom and as an entanglement entropy across the same horizon, to scale with the mean area of the black hole horizon, and to match (under additional assumptions, amounting to semiclassicality conditions) the Bekenstein-Hawking result.

We depart from the canonical LQG approach to black hole entropy in three main ways: (i) the horizon quantum state is defined including a sum over triangulations, including very refined ones, admitting in this way an interpretation in terms of a continuum geometry, whose information is encoded in a single collective variable, i.e., the condensate wave function; crucially, this basic aspect of the construction also implies that our states encode in their

very definition a coarse-graining of microscopic degrees of freedom allowing one to control them via a limited number of variables—this point is also relevant for their (lack of) dynamical character: being the result of a coarse graining, we should not expect them to be exact solutions of the microscopic quantum dynamics; (ii) the interior bulk degrees of freedom are not removed (or drastically reduced) by hand through the introduction of a single intertwiner model; space-time does not end at the horizon in our construction, but both an interior and an exterior bulk are included in the quantum state; (iii) our construction relies uniquely on structures and techniques proper of the GFT formalism, i.e., both boundary and bulk degrees of freedom are described in a unique, consistent way, removing ambiguities present in the canonical LQG approach, where spin-network states are coupled to a Chern-Simons theory on the boundary.<sup>2</sup>

Most importantly, with all the limitations of our work, and the inevitable approximations and assumptions, we work from beginning to end within the full quantum gravity formalism, without any preliminary classical symmetry reduction and with realistic quantum states of the full theory. These points of departure from the standard treatment are also at the origin of a remarkable feature of the entropy calculation we present in this paper: by replacing the area operator eigenstates for the quantum isolated horizon by condensate states, the result of the entropy calculation yields the semiclassical Bekenstein-Hawking entropy area law with no explicit dependence on the value of the Barbero-Immirzi parameter.

## II. SHELL CONDENSATE STATE: GROUP ELEMENT REPRESENTATION

We start by presenting in some detail the construction of our quantum black hole states in the GFT formalism. We do so, in this and in the next sections, in two equivalent representations of the Hilbert (Fock) space of the theory. This serves also the purpose of showing the generality of our construction, and of our results. We take a sort of engineering approach, by building up our quantum states piece by piece, starting from the fundamental building blocks provided by the GFT formalism, i.e., GFT quanta corresponding to individual spin-network vertices, in turn dual to fundamental three-dimensional simplices. Specifically, we construct spherically symmetric configurations of quantum space as glueings (along a radial direction) of homogeneous spherical shells.

As pointed out above, in this work we build upon the construction of a continuum quantum geometry representing a spherically symmetric shell performed in [45], to

<sup>1</sup>Its precise value would be a quantitative measure of the goodness of the approximation used.

<sup>2</sup>See however [52,53] for a more uniform treatment of bulk and boundary degrees of freedom through techniques developed in the context of 2 + 1 LQG [54–58].

which we refer for notation and basic notions.<sup>3</sup> Their construction can be easily described: we start with a seed state for a given shell, containing few quanta, and then we act upon it with a series of refinement operators in order to increase the number of fundamental blocks (four-valent vertices), preserving the initial topology. The same wave function is associated to each new fundamental block, enforcing homogeneity and symmetry.

Therefore, the GFT condensate state for a given shell is formed by an infinite superposition of graphs, each with a certain number of 4 vertices connected together. A color  $t = \{B, W\}$  is associated to each 4-vertex and four  $SU(2)$  group elements (denoted by the letter  $g$  variously decorated) are assigned to the links departing from a given 4-vertex and labeled by a number  $I = \{1, 2, 3, 4\}$ . A shell is formed of three parts: an outer boundary, an inner boundary, and a bulk in between. In order to keep track of these three parts of each shell, we add a color  $s = \{+, 0, -\}$  to the vertex wave function, labeling respectively these three parts, so that we can specify the region to which a given vertex of a shell belongs.

Following the conventions of [45] (see also the Appendix), the action of the refinement operators on the initial seed state is such that each boundary of a given shell is formed by open radial links all with the same color, with outer and inner boundaries having different colors. The next step is to glue shells together. In order to do this, while still being able to distinguish different shells, we need to introduce a single extra label  $r \in \mathbb{N}$ , also associated to the shell wave function, which can be interpreted as an effective radial coordinate. Therefore, the field operator associated to the fundamental building block  $v$  reads

$$\begin{aligned}\hat{\sigma}_{r,t^v s^v}(h_I^v) &= \int dg_I^v \sigma_{r,s^v}(h_I^v g_I^v) \hat{\varphi}_{t^v}(g_I^v), \\ \hat{\sigma}_{r,t^v s^v}^\dagger(h_I^v) &= \int dg_I^v \overline{\sigma_{r,s^v}(h_I^v g_I^v)} \hat{\varphi}_{t^v}^\dagger(g_I^v),\end{aligned}\quad (2)$$

satisfying the commutation relations

$$\begin{aligned}[\hat{\sigma}_{r,t^v s^v}(h_I^v), \hat{\sigma}_{r',t'^w s'^w}(h_I'^w)] &= \delta_{r,r'} \delta_{t^v,t'^w} \delta_{s^v,s'^w} \Delta_L(h_I^v, h_I'^w) \\ &\equiv \delta_{r,r'} \delta_{t^v,t'^w} \delta_{s^v,s'^w} \int_{SU(2)} d\gamma \\ &\quad \times \prod_{I=1}^4 \delta(\gamma h_I^v (h_I'^w)^{-1}),\end{aligned}\quad (3)$$


where the rhs guarantees the left gauge invariance of the vertex wave function, namely

$$\hat{\sigma}_{r,t^v s^v}(h_I^v) = \hat{\sigma}_{r,t^v s^v}(\gamma h_I^v), \quad \forall \gamma \in SU(2). \quad (4)$$


<sup>3</sup>For the reader's convenience, we have reported the basic notion in the Appendix.

Moreover, the  $\delta_{r,r'}$  implies that operators associated to different shells commute with each other. The above field operators are constructed out of the fundamental GFT field operators by convolution with the condensate wave function  $\sigma$ ; they thus create/annihilate GFT quanta, all associated with such a wave function. This association with a unique wave function for all the quanta forming a given shell is what we call wave function homogeneity, which puts these states in correspondence with homogeneous (continuum) spatial geometries, and also what characterizes the same states as GFT condensates. The condensate quanta are then glued to one another, for a given radial parameter, to form the three-dimensional triangulations constituting the shell, with quantum correlations encoding spatial topology [59,60].

In order to form a full space foliation we glue all the radial links belonging to the outer boundary of a given shell  $r$  with the radial links belonging to the inner boundary of the shell  $r+1$ . For the gluing to be done consistently, the two boundaries have to have the same number and color of radial links. Two glued shells can be graphically represented as



The use of bipartite colored graphs in order to be able to encode the information about the spatial topology suggests that it is most convenient to adopt a construction of the seed state in terms of melonic graphs, and of the associated refinement operators in terms of dipole (or melonic) moves. Explicitly, the seed state for a given shell  $r$  is graphically represented as



and, in terms of field operators, the seed state is given by



$$|\tau\rangle = \int (dg)^{10} \hat{\sigma}_{r,B+}^\dagger(e, g_2, g_3, g_4) \hat{\sigma}_{r,W+}^\dagger(e, g'_2, g_3, g_4) \hat{\sigma}_{r,B0}^\dagger(g''_1, g'_2, g''_3, g'_4) \\ \times \hat{\sigma}_{r,W0}^\dagger(g''_1, g_2, g''_3, g'_4) \hat{\sigma}_{r,B-}^\dagger(g'''_1, g'_2, g'_3, e) \hat{\sigma}_{r,W-}^\dagger(g'''_1, g'_2, g'_3, e) |0\rangle, \quad (7)$$

where we have arbitrarily assigned the color 1 to the radial links of the boundary + and the color 4 to the boundary -, and, for the moment, we have set the gluing group elements  $h$ 's associated to both sets of radial links to the identity.

There are three refinement operators for each shell: Two refine the boundaries and one the bulk vertices. The complete set of refinement operators has been studied in [45]. Here we just concentrate on one of the two boundaries, namely the + one. The construction for the other one follows the same logic. The action of the operator for the refinement of white vertices has a simple graphical representation,

$$\widehat{\mathcal{M}}_{r,W+} : \begin{array}{c} |1 \\ \circ \\ \hline 2 \quad 4 \\ \hline 3 \end{array} \rightarrow \begin{array}{c} 1'' \quad | \quad 1' \\ \circ \quad 4' \quad 4' \quad \bullet \\ \hline 2 \quad 2' \quad 2' \quad 2 \\ \hline 3' \quad 3' \quad 3 \quad |1 \\ \circ \end{array}, \quad (8)$$

with the one for black vertices having a similar structure (notice the colors of the edge in the loop, however),

$$\widehat{\mathcal{M}}_{r,B+} : \begin{array}{c} |1 \\ \bullet \\ \hline 4 \quad 2 \\ \hline 3 \end{array} \rightarrow \begin{array}{c} 1'' \quad | \quad 1' \\ \bullet \quad 2' \quad 2' \quad \circ \\ \hline 4 \quad 4' \quad 4' \quad 4 \\ \hline 3' \quad 3' \quad 3 \quad |1 \\ \bullet \end{array}. \quad (9)$$

These two moves are the ones involving the minimum number of vertices, keeping fixed the topology. In terms of the group fields, the two move operators read

$$\widehat{\mathcal{M}}_{r,W+} \equiv \int dk_2 dk_3 dk_4 dh_{4'} dh_{2'} dh_{3'} \\ \times \hat{\sigma}_{W-}^\dagger(e, k_2, h_{3'}, h_{4'}) \hat{\sigma}_{B-}^\dagger(e, h_{2'}, h_{3'}, h_{4'}) \hat{\sigma}_{W-}^\dagger(e, h_{2'}, k_3, k_4) \hat{\sigma}_{W-}(e, k_2, k_3, k_4) \quad (10)$$

and

$$\widehat{\mathcal{M}}_{r,B+} \equiv \int dk_2 dk_3 dk_4 dh_{4'} dh_{2'} dh_{3'} \\ \times \hat{\sigma}_{r,B+}^\dagger(e, h_{2'}, h_{3'}, k_4) \hat{\sigma}_{r,W+}^\dagger(e, h_{2'}, h_{3'}, h_{4'}) \hat{\sigma}_{r,B+}^\dagger(e, k_2, k_3, h_{4'}) \hat{\sigma}_{r,B+}(e, k_2, k_3, k_4). \quad (11)$$

These are graph topology-preserving operators and their actions (9) and (8) can be straightforwardly verified by computing their commutators with, respectively,  $\hat{\sigma}_{r,B+}^\dagger(e, g_2, g_3, g_4)$ ,  $\hat{\sigma}_{r,W+}^\dagger(e, g_2, g_3, g_4)$ . In a similar fashion, we can build refinement operators for vertices belonging to the other two components of the shell. We can thus arbitrarily refine the seed state (7) by repeated action of the operators  $\widehat{\mathcal{M}}_{r,ts}$  in order to implement the sum over triangulations while preserving the desired topology and the key feature of wave function homogeneity. In fact, the refinement moves are implemented through operators built out of field operators (2) dressed with the same wave function as the seed state; in this way, the geometric information of the finer shell states is still encoded in the same small number of parameters. The state of a given shell  $r$  can then be written as

$$|\Psi_r\rangle = F_r(\widehat{\mathcal{M}}_{r,Bs}, \widehat{\mathcal{M}}_{r,Ws})|\tau\rangle, \quad (12)$$

where, at this stage,  $F_r$  is a generic function of the refinement operators associated to the given shell  $r$ .

This completes the definition of the spherically symmetric quantum states we use in the following, for describing the microstructure of quantum black holes.

### A. The area operator

The physical properties encoded in our quantum states have to be extracted by computing suitable operators. While the combinatorial aspects of the GFT formalism, shared with random tensor models, were crucial in the definition of our quantum states, alongside its second quantization tools, the quantum geometric aspects, shared with loop quantum

gravity (and simplicial approaches), become prominent in the physical interpretation of the same states and in the identification of interesting operators.

An important geometric operator for our purposes is the area operator. Following the prescription in [45], a second quantized version of a shell boundaries area operator is defined by

$$\begin{aligned}\hat{\mathbb{A}}_{Jr,s} &= \sum_{t=B,W} \hat{\mathbb{A}}_{Jr,ts} \\ &\equiv \kappa \sum_{t=B,W} \int dh_1^v \hat{\sigma}_{r,ts}^\dagger(h_1^v) \sqrt{E_J^i E_J^j \delta_{ij}} \triangleright \hat{\sigma}_{r,ts}(h_1^v),\end{aligned}\quad (13)$$

where  $\kappa = 8\pi\gamma\ell_p^2$ , introducing the dependence on the Barbero-Immirzi parameter  $\gamma$ . In the expression above the label  $s$  takes values  $\{+, -\}$ , according to which boundary of the shell  $r$  we want to compute the area of, and the index  $J$  matches the number associated to the radial links dual to the given boundary.

The action of the operator (13) is computed using the definition

$$E_J^i \triangleright f(g_I) := \lim_{\epsilon \rightarrow 0} \frac{d}{d\epsilon} f(g_1, \dots, e^{-i\epsilon \tau^i} g_J, \dots, g_4) \quad (14)$$

for a given function  $f: SU(2)^4 \rightarrow \mathbb{C}$ . It is immediate to see that the expectation value of the area operator (13) on a shell boundary state can be written as a function of the number of quanta and a single vertex expectation value,

$$\begin{aligned}\langle \hat{\mathbb{A}}_{Jr,s} \rangle &= \kappa \langle \hat{n}_{r,s} \rangle \int dh_1^v dg_1^v \sigma_{r,s}(h_1^v g_1^v) \\ &\quad \times \sqrt{E_J^i E_J^j \delta_{ij}} \triangleright \sigma_{r,s}(h_1^v g_1^v) \equiv \langle \hat{n}_{r,s} \rangle a_{Jr,s},\end{aligned}\quad (15)$$

where we have defined  $a_{Jr,s}$  as the expectation value of the area operator on a single radial link  $J$  in the boundary  $s$  of the shell  $r$ , and  $\hat{n}_{r,s}$  is the number operator given by

$$\hat{n}_{r,s} = \sum_{t=B,W} \hat{n}_{r,ts} = \sum_{t=B,W} \int dh_1^v \hat{\sigma}_{r,ts}^\dagger(h_1^v) \hat{\sigma}_{r,ts}(h_1^v). \quad (16)$$

Notice that, due to the definition of the seed state and the refinement operators, after each refinement action the graphs are such that

$$n_{r,Bs} = n_{r,Ws} = \frac{n_{r,s}}{2} \quad (17)$$

always holds, where  $n \equiv \langle \hat{n} \rangle$ .

An analog one-body operator can be constructed for the volume. Also in this case, the factorization property (15) holds. The structure of these expectation values should come as natural given the structure of the wave function, boiling down to dimensional considerations (areas are extensive quantities) and to the fact that a single wave function has been used.

### III. SHELL CONDENSATE STATE: SPIN REPRESENTATION

Let us now introduce a dual spin representation for generalized GFT condensates, and a different example of condensate states constructed by the same scheme but relying on this dual representation. This provides a useful computational toolkit and, at the same time, it allows us to circumvent the issue of non-normalizability in the kinematical Hilbert space (the Fock space) of the condensate states constructed out of the field operators (2). In practice, when working in the group representation, and for the condensate states described in the previous section, a regularization scheme is generally required in order to obtain finite expectation values for geometric operators like (15). On the other hand, when working in the spin representation at fixed spins, and constructing adapted condensate states using the same scheme, this issue does not arise. Beside these technical advantages, we detail this dual construction for two additional reasons. First, it shows the generality of our construction and results; in fact, starting from these two basic definitions of condensate states, one can envisage new definitions combining or interpolating between the two basic ones (which play a role of possible bases for linear combinations possessing similar properties). Second, this dual spin-based construction is the one producing quantum GFT condensate states that are the closest to the quantum states customarily used in loop quantum gravity as models of quantum black holes, based on eigenstates of the area operator and thus labeled by fixed spins on the links puncturing the black hole horizon. This facilitates comparison of our results with the LQG literature.

The spin representation follows from a straightforward Peter-Weyl decomposition of the vertex wave function, with the field operators (2) now becoming

$$\begin{aligned}\hat{\sigma}_{r,ts}(h_I) &= \int dg_I \sigma_{r,s}(h_I g_I) \hat{\phi}_t(g_I) = \int dg_I \sum_{\{j\}, l_R, l_L} \sigma_{r,s}^{j_1 \dots j_4 l_L l_R} \iota_{m_1 m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l_L} \iota_{n_1 n_2 n_3 n_4}^{j_1 j_2 j_3 j_4 l_R} \prod_{I=1}^4 D_{m_I o_I}^{j_I}(h_I) D_{o_I n_I}^{j_I}(g_I) \hat{\phi}_t(g_I) \\ &= \sum_{\{j\}, l_R, l_L} \sigma^{j_1 \dots j_4 l_L l_R} \iota_{m_1 m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l_L} \hat{a}_{(r,ts) o_1 \dots o_4}^{j_1 \dots j_4 l_R} \prod_{I=1}^4 D_{m_I o_I}^{j_I}(h_I),\end{aligned}\quad (18)$$

where we defined the following new field operators

$$\hat{a}_{(r,ts)o_1\dots o_4}^{j_1\dots j_4 l_R} = \sigma_{r,s} t_{n_1 n_2 n_3 n_4}^{j_1 j_2 j_3 j_4 l_R} \int dg_I \prod_{I=1}^4 D_{o_I n_I}^{j_I} (g_I) \hat{\varphi}_I (g_I), \quad (19)$$

and we have assumed the factorization property

$$\sigma_{r,s}^{j_1\dots j_4 l_R} = \sigma_{r,s} \sigma^{j_1\dots j_4 l_R}. \quad (20)$$

One might wonder whether such a restriction is plausible or not. Let us notice that this particular choice means that different layers differ more because of their volume (controlled essentially by the number of quanta, determined by  $|\sigma_{r,s}|^2$ ) rather than their intrinsic geometry (type of curvature, for instance). This situation is very natural in spherical symmetry, as it is by its very definition in the continuum setting: the geometry of different two-dimensional spheres singled out by the isometry group differs only by a rescaling determined by the radial coordinate, the rest being constrained by the isometry group itself.

From the commutation relation of the basic field operators

$$\begin{aligned} [\hat{\varphi}_I (g_I), \hat{\varphi}_I^\dagger (g_I')] &= \delta_{I,I'} \Delta_R (g, g') \\ &\equiv \delta_{I,I'} \int_{SU(2)} d\gamma \prod_{I=1}^4 \delta(g_I \gamma (g_I')^{-1}) \end{aligned} \quad (21)$$

and the orthonormality relations between the Wigner matrices

$$\int dg D_{mn}^j (g) \overline{D_{m'n'}^j (g)} = \frac{1}{d_j} \delta_{j,j'} \delta_{m,m'} \delta_{n,n'}, \quad (22)$$

it is immediate to see that the new field operators (19) satisfy

$$\widehat{\mathcal{M}}_{(r,B+)o_1 o_1' o_1''}^{j_1 j_2 j_3 j_4 l_R} = \frac{1}{|\sigma_{r,s}|^2} \left( \prod_{I=1}^4 d_{j_I} \right) \sum_{\{m\}} \hat{a}_{(r,B+)o_1 m_2 m_3 m_4}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,W+)o_1' m_2' m_3' m_4'}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,B+)o_1'' m_2'' m_3'' m_4''}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,B+)o_1'' m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l_R}, \quad (27)$$

where we notice the absence of the sign flips in the last term, due to the fact that we have  $\hat{a}$  instead of  $\hat{a}^\dagger$ .

By means of (23), we can now verify that the refinement operators above realize the move depicted in (9), namely

$$[\widehat{\mathcal{M}}_{(r,B+)o_1 o_1' o_1''}^{j_1 j_2 j_3 j_4 l_R}, \hat{a}_{(r,B+)n_1 n_2 n_3 n_4}^{\dagger j_1' j_2' j_3' j_4' l_R}] = \sum_{\{m\}} \hat{a}_{(r,B+)o_1 m_2 n_3 n_4}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,W+)o_1' m_2' m_3' m_4'}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,B+)o_1'' m_2'' m_3'' m_4''}^{\dagger j_1 j_2 j_3 j_4 l_R} \delta_{o_1'' n_1} \delta_{l_R, l_R'} \prod_I \delta_{j_I, j_I'}. \quad (28)$$

Similarly, we have

$$\widehat{\mathcal{M}}_{(r,W+)o_1 o_1' o_1''}^{j_1 j_2 j_3 j_4 l_R} = \frac{1}{|\sigma_{r,s}|^2} \left( \prod_{I=1}^4 d_{j_I} \right) \sum_{\{m\}} \hat{a}_{(r,W+)o_1 m_2 m_3 m_4}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,B+)o_1' m_2' m_3' m_4'}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,W+)o_1'' m_2'' m_3'' m_4''}^{\dagger j_1 j_2 j_3 j_4 l_R} \hat{a}_{(r,W+)o_1'' m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l_R}, \quad (29)$$

and a direct calculation as in the black case shows that the operator above implements the action (8).

$$\begin{aligned} &[\hat{a}_{(r,ts)o_1\dots o_4}^{j_1\dots j_4 l_R}, \hat{a}_{(r',t's')o_1'\dots o_4'}^{\dagger j_1'\dots j_4' l_R'}] \\ &= |\sigma_{r,s}|^2 \delta_{r,r'} \delta_{t,t'} \delta_{s,s'} \delta_{l_R, l_R'} n(j_1, j_2, j_3, j_4, l_R) \\ &\quad \times \prod_{I=1}^4 \frac{1}{d_{j_I}} \delta_{o_I o_I'} \delta_{j_I j_I'}, \end{aligned} \quad (23)$$

where

$$\delta^{l, l'} n(j_1, j_2, j_3, j_4, l) = \sum_{\{m\}} t_{m_1 m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l} t_{m_1 m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l'} \quad (24)$$

is the intertwiners normalization factor. With the convention

$$t_{m_1 m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l} = \sum_{m, m'} C_{m_1 m_2 m}^{j_1 j_2 l} C_{m_3 m_4 m'}^{j_3 j_4 l'} C_{mm'0}^{l l'}, \quad (25)$$

where the  $C$ 's are the Clebsch-Gordan coefficients, the normalization factor  $n(j_1, j_2, j_3, j_4, l)$  is equal to 1 and therefore we omit it from now on.

The wave function of the new field operators then reads

$$\begin{aligned} a_{(r,s)o_1\dots o_4}^{j_1\dots j_4 l_R} (g_I) &\equiv \langle g_I | \hat{a}_{(r,ts)o_1\dots o_4}^{\dagger j_1\dots j_4 l_R} | 0 \rangle \\ &= \sigma_{r,s} t_{n_1 n_2 n_3 n_4}^{j_1 j_2 j_3 j_4 l_R} \prod_{I=1}^4 D_{o_I n_I}^{j_I} (g_I). \end{aligned} \quad (26)$$

## A. Refinement operators

We now write the refinement operators  $\widehat{\mathcal{M}}_{r, B_s}, \widehat{\mathcal{M}}_{r, W_s}$  in terms of the spin representation field operators (19). We concentrate on the operators for the outer boundary  $s = +$ . Refinement operators for the other boundaries can be constructed straightforwardly in a similar fashion. Assuming again that the boundary radial links have label 1, we define

### B. Area expectation value

Let us compute the area expectation value on a single-vertex state created by the field operator (19). Consistently with our previous convention, the vertex radial link dual to the boundary of interest has color 1. Using (15) and (26), we get

$$\begin{aligned}
 a_{1r,s} &= \kappa \int dg_I a_{(r,s)o_1\dots o_4}^{j_1\dots j_4 l_R}(g_I) \sqrt{E_1^i E_1^j \delta_{ij}} \triangleright \overline{a_{(r,s)o_1\dots o_4}^{j_1\dots j_4 l_R}(g_I)} \\
 &= \kappa \int dg_I |\sigma_{r,s}|^2 t_{m_1 m_2 m_3 m_4}^{j_1 j_2 j_3 j_4 l_R} t_{n_1 n_2 n_3 n_4}^{j_1 j_2 j_3 j_4 l_R} \sqrt{j_1(j_1+1)} \\
 &\quad \times \prod_{I=1}^4 D_{o_I m_I}^{j_I}(g_I) \overline{D_{o_I n_I}^{j_I}(g_I)} \\
 &= \kappa |\sigma_{r,s}|^2 \sqrt{j_1(j_1+1)} \prod_{I=1}^4 \frac{1}{d_{j_I}}. \tag{30}
 \end{aligned}$$

As anticipated above, the only dependence on the vertex wave function is through the normalization factor  $|\sigma_{r,s}|^2$ . As we have already remarked, this might seem suspicious but it is to be expected if we take seriously the idea that these states have leaves which support the same symmetry group.

### IV. SHELL DENSITY MATRIX

Given the single shell state (12), we can construct a pure state associated to a full three-dimensional spatial foliation through a shell-gluing procedure. This can be obtained through the definition of a full foliation refinement operator built out of the single-shell ones. At this kinematical stage, there is considerable freedom left in such a definition, with the only constraint coming from the requirement of spatial topology preservation. More precisely, the refinement and gluing operations have to be carried out in such a way that the outer boundary of a given shell  $r$  always contains the same number of vertices as the inner boundary of the shell  $r+1$  it is glued to, so that no open links are created. It should be remarked, however, that the structure of the refinement operators associated to the three different components of a given shell is such that these can act independently on one another. Therefore, without further

inputs coming from the dynamics, the most generic complete-foliation state can be written as a product of single-shell states, namely

$$|\Psi\rangle = \prod_r |\Psi_r\rangle, \tag{31}$$

upon which the constraint on the synchronization of nearby shell boundaries refinement is applied (thus removing the factorization into shells and introducing correlations).

From the pure state (31) we can obtain the density matrix

$$\hat{\rho} = |\Psi\rangle\langle\Psi| \tag{32}$$

of the full three-dimensional spatial foliation.

### A. Reduced density matrix for a single shell

We now want to compute the reduced density matrix associated to the outer boundary of a single shell  $r$  by tracing over the rest of the bulk state. This reduced density matrix is the central object for our entropy calculation.

In general, the boundary component of a given shell state (12) contains a superpositions of all graphs that can be obtained by all possible combinations of strings of refinement operators associated to the boundary. The coefficients of such superposition are determined by the specific form of the function  $F_r$ . We come back to this important point in a moment. For now, in order to understand the entanglement structure between different shells, let us start by simply considering a given graph  $A$  associated to the shell  $r$  outer boundary and the graph  $B$  of the inner boundary of the shell  $r+1$  right outside. The result of this simple example can be easily generalized to the rest of the graph. Therefore, if we assume that both graphs are formed by  $n$  vertices (in order to be properly glued they must have the same number of building blocks), where again we take the connecting radial links of color 1, we can write the wave function of these two components as

$$\psi(g_I^{A_1}, \dots, g_I^{A_n}, g_I^{B_1}, \dots, g_I^{B_n}) = \prod_{i=1}^n a_{A_i m_i^1 \dots m_i^4}^{j_1 \dots j_4 l_R}(g_I^{A_i}) a_{B_i n_i^1 \dots n_i^4}^{j_1 \dots j_4 l_R}(g_I^{B_i}) \delta_{m_i^1, -n_i^1} t_{m_i^2, -n_i^2}^{j_2(i)} \prod_{J=2}^4 \delta_{m_i^J, -n_i^J} t_{n_i^J, -n_i^J}^{j_J(i)}, \tag{33}$$

where the  $\delta$ 's are used to keep track of the connectivity of the whole graph  $A \cup B$ , with the notation  $t_J^m(i)$  ( $t_J^j(i)$ ) indicating the target vertex in the graph  $A$  ( $B$ ) of the edge of color  $J$  departing from the vertex  $i$  in the graph  $A$  ( $B$ ), and similarly for  $t_J^m(i)$  with the target vertex in the graph  $B$  (instead of  $A$ ), encoding the connectivity between the two boundaries through the radial links of color 1. We can thus write the total density matrix as



$$\rho^{(n)}(g_I^{A_1}, \dots, g_I^{A_n}, g_I^{B_1}, \dots, g_I^{B_n}; g_I^{A_1}, \dots, g_I^{A_n}, g_I^{B_1}, \dots, g_I^{B_n}) \quad (34)$$

$$\begin{aligned} &= C \prod_{i=1}^n a_{A_i m_1^i \dots m_4^i}^{j_1 \dots j_4 L_R}(g_I^{A_i}) a_{B_i n_1^i \dots n_4^i}^{j_1 \dots j_4 L_R}(g_I^{B_i}) \delta_{m_1^i, -n_1^{i'}} \prod_{J=2}^4 \delta_{m_J^i, -m_J^{i'}} \delta_{n_J^i, -n_J^{i'}} \\ &\times \overline{a_{A_i m_1^i \dots m_4^i}^{j_1 \dots j_4 L_R}(g_I^{A_i}) a_{B_i n_1^i \dots n_4^i}^{j_1 \dots j_4 L_R}(g_I^{B_i})} \delta_{m_1^i, -n_1^{i'}} \prod_{J=2}^4 \delta_{m_J^i, -m_J^{i'}} \delta_{n_J^i, -n_J^{i'}}, \end{aligned} \quad (35)$$

where  $C$  is a normalization factor.

If we now use the relation

$$\int dg_I a_{(r,s)m_1 \dots m_4}^{j_1 \dots j_4 L_R}(g_I) \overline{a_{(r,s)n_1 \dots n_4}^{j_1 \dots j_4 L_R}(g_I)} = |\sigma_{r,s}|^2 \prod_{I=1}^4 \frac{1}{d_{j_I}} \delta_{m_I, n_I}, \quad (36)$$

following from the commutation relation (23), to integrate away the  $B$  part, we get

$$\rho_A^{(n)}(g_I^1, \dots, g_I^n; g_I^1, \dots, g_I^n) = \left( \frac{\prod_{I=1}^4 d_{j_I}}{|\sigma_{r,s}|^2} \right)^n \prod_{i=1}^n a_{A_i m_1^i \dots m_4^i}^{j_1 \dots j_4 L_R}(g_I^i) \overline{a_{A_i m_1^i \dots m_4^i}^{j_1 \dots j_4 L_R}(g_I^i)} \delta_{m_1^i, m_1^i} \prod_{J=2}^4 \delta_{m_J^i, -m_J^{i'}} \delta_{m_J^i, -m_J^{i'}}, \quad (37)$$

where we have set the normalization factor  $C = (\prod_{I=1}^4 d_{j_I} / |\sigma_{r,s}|^2)^{2n}$ . We see that the normalized reduced density matrix we obtain is mixed, as a consequence of the relation  $m_1^i = m_1^i$  imposed by the first set of  $\delta$ 's and following from the property (36).

Remarkably, as it emerges from this simple example, any property about the rest of the graph traced away disappears: different completions of the same visible portions of the state lead to the same reduced density matrix. This is a direct consequence of the commutation relation (23) [which implies (36)]. This holographic property of our states remains valid also when tracing away a bigger graph in the bulk, exactly for the same mechanism. This means that, given a graph for the whole three-dimensional space foliation, if we choose the boundary of an arbitrary shell  $r$  and trace away all the rest of the graph, we end up with a mixed reduced density matrix which contains *no information* about the bulk degrees of freedom. The only entanglement that remains in the reduced density matrix is the one induced by the radial links of the closest shell. Let us reiterate the message: for these particular states, all the rest of the information about the remaining graph disappears

from the reduced density matrix, regardless of how big or intricate the rest of the graph is. This means that the entanglement entropy contribution comes uniquely from the entanglement with the closest shell, as expected by standard calculation in QFT on a sphere [61].

## B. Entanglement entropy

To compute the entanglement entropy we now need to diagonalize the reduced density matrix, that is, we need to compute the eigenvectors of (37). Let us consider the state

$$\Psi_A^{(n)}(n_1, g) = \left( \frac{\prod_{I=1}^4 d_{j_I}}{|\sigma_{r,s}|^2} \right)^{n/2} \prod_{i=1}^n \overline{a_{A_i n_1^i \dots n_4^i}^{j_1 \dots j_4 L_R}(g_I^i)} \prod_{J=2}^4 \delta_{n_J^i, -n_J^{i'}}, \quad (38)$$

which satisfies

$$\langle \Psi_A^{(n)}(n_1, g) | \Psi_A^{(n)}(n_1', g) \rangle = \prod_{i=1}^n \delta_{n_1^i, n_1^i}, \quad (39)$$

and compute

$$\begin{aligned} \int \prod_{i=1}^n dg_I \rho_{\text{red}}^{(n)}(g_I^1, \dots, g_I^n; g_I^1, \dots, g_I^n) \Psi_A^{(n)}(n_1, g) &= \left( \frac{\prod_{I=1}^4 d_{j_I}}{|\sigma_{r,s}|^2} \right)^{\frac{3}{2}n} \int \prod_{i=1}^n dg_I a_{A_i m_1^i \dots m_4^i}^{j_1 \dots j_4 L_R}(g_I^i) \overline{a_{A_i m_1^i \dots m_4^i}^{j_1 \dots j_4 L_R}(g_I^i)} \delta_{m_1^i, m_1^i} \\ &\times \prod_{J=2}^4 \delta_{m_J^i, -m_J^{i'}} \delta_{m_J^i, -m_J^{i'}} \overline{a_{A_i n_1^i \dots n_4^i}^{j_1 \dots j_4 L_R}(g_I^i)} \prod_{J=2}^4 \delta_{n_J^i, -n_J^{i'}} \\ &= \left( \frac{\prod_{I=1}^4 d_{j_I}}{|\sigma_{r,s}|^2} \right)^{\frac{n}{2}} \overline{a_{A_i m_1^i \dots m_4^i}^{j_1 \dots j_4 L_R}(g_I^i)} \prod_{J=2}^4 \delta_{m_J^i, -m_J^{i'}} \\ &= \Psi_A^{(n)}(m_1, g'), \end{aligned} \quad (40)$$

where we have used the relation (36) again. From the calculation above, we see that the states (38) are eigenstates of the shell reduced density matrix (37) with eigenvalue 1.

We can thus label all the eigenstates of the shell reduced density matrix using a graph basis and the notation  $\Psi_{r,s}^{(n)}(\Gamma_\alpha)$  to denote the states (38), where the structure of the given graph  $\Gamma_\alpha$  is encoded in the product of deltas  $\prod_{j=2}^4 \delta_{n_j^i, -n_j^{i'}}$ .

Given that this does not hold in general [10], it is instructive to show explicitly the orthogonality of the states  $\Psi_{r,s}^{(n)}(\Gamma_\alpha)$  for different graphs  $\Gamma_\alpha, \Gamma_{\alpha'}$ . We want to prove that

$$\langle \Psi_{r,s}^{(n)}(\Gamma_\alpha) | \Psi_{r,s}^{(n)}(\Gamma_{\alpha'}) \rangle = \delta_{\alpha,\alpha'} \prod_{i=1}^n \delta_{n_i^i, n_i^{i'}}. \quad (41)$$

To do so, for all  $\alpha \neq \alpha'$ , it is enough to consider one simple case, namely graphs with  $n_B = n_W = n/2 = 2$  (recall that all the graphs have always the same number of black and white vertices). These states are created by acting once with the refinement operators (8) and (9) on the seed state for the shell boundary.

A graph with four vertices arranged in a square. The top-left vertex is black and labeled 1. The top-right vertex is white and labeled 2'. The bottom-left vertex is white and labeled 3. The bottom-right vertex is black and labeled 4. Edges connect 1 to 2', 1 to 3, 2' to 4, and 3 to 4.

$$\quad (42)$$

where links 1 are the radial ones and links 2 are connected to the bulk of the shell. The two possible states with 4 vertices, when acting with the black (9) and white (8) refinement operators respectively, thus are

A graph with four vertices arranged in a square. The top-left vertex is white and labeled 1'''. The top-right vertex is black and labeled 1''. The bottom-left vertex is white and labeled 3'. The bottom-right vertex is black and labeled 3. Edges connect 1''' to 1'', 1''' to 3', 1'' to 3, and 3' to 3.

$$|\Psi_{r,1}^{(4)}\rangle : \quad (43)$$

A graph with four vertices arranged in a square. The top-left vertex is black and labeled 1''. The top-right vertex is white and labeled 1'''. The bottom-left vertex is white and labeled 3'. The bottom-right vertex is black and labeled 3. Edges connect 1'' to 1''', 1'' to 3', 1''' to 3, and 3' to 3.

$$|\Psi_{r,2}^{(4)}\rangle : \quad (44)$$

The scalar product between these two states gives

$$\begin{aligned} \langle \Psi_{r,2}^{(4)} | \Psi_{r,1}^{(4)} \rangle &= \int dg_1^1 dg_1^2 dg_1^3 dg_1^4 dg_1^1 dg_1^2 dg_1^3 dg_1^4 \\ &= \overline{a_{m_1^1 m_2^1 m_3^1 m_4^1}^{j_1 \dots j_4 L_R}(g_1^1) a_{m_2^2 m_3^2 m_4^2}^{j_1 \dots j_4 L_R}(g_1^2) a_{m_1^3 m_2^3 m_3^3 m_4^3}^{j_1 \dots j_4 L_R}(g_1^3) a_{m_4^4 m_3^4 m_2^4 m_1^4}^{j_1 \dots j_4 L_R}(g_1^4)} \\ &= \left( \prod_{i=1}^4 \frac{1}{d_{j_i}} \right)^4 (\delta_{m_1^1, n_1^1} \delta_{m_2^1, n_2^1} \delta_{m_3^1, n_3^1} \delta_{m_4^1, n_4^1} \delta_{m_1^2, n_1^2} \delta_{m_2^2, n_2^2} \delta_{m_3^2, n_3^2} \delta_{m_4^2, n_4^2} \\ &\quad + \delta_{m_1^1, n_1^1} \delta_{m_2^1, -n_2^1} \delta_{m_3^1, n_3^1} \delta_{m_4^1, n_4^1} \delta_{m_1^2, n_1^2} \delta_{m_2^2, -n_2^2} \delta_{m_3^2, n_3^2} \delta_{m_4^2, n_4^2}) \\ &\quad \times (\delta_{m_1^3, n_1^3} \delta_{m_2^3, n_2^3} \delta_{m_3^3, -n_3^3} \delta_{m_4^3, n_4^3} \delta_{m_1^4, n_1^4} \delta_{m_2^4, n_2^4} \delta_{m_3^4, n_3^4} \delta_{m_4^4, n_4^4} \\ &\quad + \delta_{m_1^3, n_1^3} \delta_{m_2^3, n_2^3} \delta_{m_3^3, -n_3^3} \delta_{m_4^3, n_4^3} \delta_{m_1^4, n_1^4} \delta_{m_2^4, -n_2^4} \delta_{m_3^4, n_3^4} \delta_{m_4^4, n_4^4}), \end{aligned}$$

where we have performed the Wick contractions and used (21). If we now expand this last expression, we end up with four products of  $\delta$ 's which all give rise to either one of the following relations,

$$\delta_{m_3^1, -m_3^2}, \quad \delta_{m_4^1, m_4^3}. \quad (45)$$

None of these two relations can be satisfied, though, due to the structure of the graph of the state (44). In fact, they

would both change the shell topology by creating two disconnected regions in the boundary graph. Therefore, the scalar product vanishes and the states (43) and (44) are orthogonal. Due to the local action of the refinement operators, similar relations follow when computing the scalar product between any two eigenstates at given  $n$  corresponding to the action of a sequence of refinement operators generating two different graphs. This implies that

all the eigenvectors  $\Psi_r^{(n)}(\Gamma_\alpha)$  of the total reduced density matrix (48) are orthogonal for  $\alpha \neq \alpha'$ . A similar calculation shows that when  $\alpha = \alpha'$  (41) is again satisfied.

This seemingly unremarkable results has a very important consequence, namely

$$\rho_{r,s}^{(n)}(\Gamma_\alpha)\Psi_{r,s}^{(n)}(\Gamma_{\alpha'}) = \begin{cases} \Psi_{r,s}^{(n)}(\Gamma_{\alpha'}) & \text{if } \alpha = \alpha' \\ 0 & \text{if } \alpha \neq \alpha', \end{cases} \quad (46)$$

which implies that we can diagonalize the reduced density matrix, since we have discovered its diagonal form in terms of graphs, even in the case in which the full sum over triangulations is kept. This result also shows that the computation of the entanglement entropy, at least in the particular corner of the Fock space that we are exploring, becomes a classic ‘‘counting of graphs’’ problem.

## V. SEMICLASSICALITY CONDITIONS

Our construction so far is general, in the sense that it can apply to any of the shells in the spacelike hypersurface of our foliation, and thus to generic spherically symmetric geometries. We now want to restrict our attention to a horizon 2-sphere cross section, i.e., we want to be able to interpret one of our shells as defining a spherical *horizon*, and for this we need to specify horizon boundary conditions. As we recalled in the introduction, in the standard LQG black hole entropy calculation these boundary conditions are specified by the notion of isolated horizon and they are implemented in the quantum theory through the imposition of the boundary conditions (1) (though most often this is already implemented at a classical level).

In the context of the GFT condensate for a spherically symmetric shell, imposition of the boundary condition (1) amounts to a relation between the flux variable  $E_1$  associated to the  $r$  radial link (in the convention adopted so far for the outer boundary of a shell) of a given vertex of the shell and the holonomy around it, constructed out of the group elements associated to the orthogonal remaining links. The latter correspond to the group elements  $g_2, g_3, g_4$  of the fundamental field operator  $\hat{\varphi}(g_I)$ . In fact, these are the group elements which have the geometrical interpretation of parallel transport from the center of a given tetrahedron to its faces lying on the horizon. The identification of the horizon shell could then be implemented through the construction of a second quantized curvature operator around a given puncture to be interpreted as the GFT holonomy operator around the radial link 1 of the corresponding vertex. However, given the fundamentally discrete nature of the GFT formalism and the lack of continuum manifold structures to be used as auxiliary tools to define quantum operators related to curvature, which generically involve correlation across several graph vertices and which correspond to intensive quantities from the GFT

many-body perspective, their precise definition for GFT condensates remains one of the main open technical challenges of this second quantization formalism.

Given this obstruction, we are not going to explicitly impose IH boundary conditions in operatorial terms. Instead, we are going to rely on a maximum entropy argument in order to characterise the most generic horizon shell geometry, as well as to capture some aspects of the semiclassical dynamics in our so far purely kinematical construction. Such treatment of the horizon semiclassical regime seems natural in light of the laws of black hole thermodynamics [4] and arguments for entropy bounds [62,63].

The isolated horizon formalism still plays a role in our entropy calculation. In fact, we check the compatibility of the single vertex Hilbert space degeneracy obtained through extremization of the global horizon entropy with the constraint (1), in the semiclassical limit. Moreover, we also require consistency with the thermodynamical properties of isolated horizons.

In addition to the notion of typicality, encoded in the maximum entropy argument, there are other geometrical requirements to be demanded in order to guarantee the validity of a semiclassical regime, namely the following:

- (1) large horizon shell boundary area;
- (2) small horizon shell bulk volume;
- (3) small fluctuations of the horizon shell geometrical operators.

This last requirement is guaranteed by the factorization property of the one-body geometrical operators (like area and volume, see Sec. II A) in the large  $n_{r,s}$  limit; from here on we thus take  $n_{r_0,s} \gg 1$ , with  $r_0$  denoting the horizon shell. It also follows rather generically from the condensate nature of our quantum states.

### A. Holographic aspects of GFT black hole condensates

Before proceeding with the black hole entropy calculation we need to identify the relevant degrees of freedom which contribute to it. In order to understand better the features of the quantum states we have constructed and of their reduced density matrix, we recall here some basic aspects of the holographic principle which is implemented by it in a specific manner.

We can identify two notions of holography: strong and weak [64]. In the strong version, the holographic principle becomes a fundamental feature of quantum gravity, asserting that all bulk degrees of freedom can be encoded on a boundary screen. This is the point of view adopted for instance in the AdS/CFT correspondence, when interpreted as an exact duality.

There are various arguments, however, to believe that in a background independent quantum gravity approach only a weaker notion of holography can survive [64]. In particular, in a thermodynamical context where interactions

between subsystems play a fundamental role, the causal structure of a black hole diagram strongly suggests that the only relevant degrees of freedom to account for the atomistic description beyond macroscopic variables are those living in the proximity of the horizon; in fact these are the only ones in causal contact with the exterior subsystem and relevant for the analysis of an external stationary observer. For these reasons, we are going to invoke a weak holographic principle in order to select the relevant reduced density matrix for the black hole entropy calculation.

Let us recall that in Sec. IV we modeled, before dynamical considerations were eventually implemented, the three-dimensional foliation of a spherically symmetric black hole geometry as a GFT pure condensate state factorized as a product over components of different shells. In particular, this means that refinement operators could act independently on different shells, suggesting that indeed strong holography is not realized at the microscopic level, at least in the kinematical setting. Selection of the microscopic degrees of freedom at the origin of the statistical mechanical nature of the Bekenstein-Hawking area entropy law thus requires us to trace out all bulk shells, both in the exterior and the interior of the horizon shell  $r_0$ .

Entanglement entropy calculations in QFT on a fixed classical background might lead to the expectation that it should be enough to trace out only part, but not all, of the bulk Hilbert space in order to obtain a result that scales with the area of the entangling surface [61]. However, this expectation is borne out of examples in which gravity is treated classically and entropy is associated to matter degrees of freedom, not gravitational ones. As we have argued above, in a general full quantum gravitational regime holography has probably a weaker nature.

At the same time, one could consider a subclass of the GFT condensate states we constructed in which nonlocal correlations between different shells are introduced by the imposition of some effective dynamics, which would allow an entropy counting compatible with a stronger form of holography. The extreme case of such a scenario would be physical states for which the refinement operators on a given shell are synchronized with those acting on all the other shells at the same time. In this case the number of boundary graphs would exhaust all the possible configurations of bulk graphs as well: Each boundary graph is in correspondence with just one graph in all the rest of the bulk and there is no extra degeneracy. Such states are in fact rather straightforward to construct.

Before proceeding, a few remarks are in order. At this stage, there is no indication that such a specific sector is the physically relevant one as this is ultimately dictated by the quantum equations of motion, even considered in some approximate form. The counting discussed in what follows, obtained by tracing over all the bulk vertices outside of the  $r_0$  shell, applies straightforwardly to this particular case of strong holography.

Let us also point out that implementation of the weak holographic principle does not automatically yield an area law for the entropy. In fact, *a priori* the horizon reduced density matrix could retain information about the rest of the bulk. It is therefore a nontrivial property of our construction the fact that, as seen above, by tracing out bulk degrees of freedom, the resulting reduced density matrix of the horizon is mixed but loses all the information about the bulk beyond its existence.

In addition, as we see in a moment, there is also a further holographic feature concerning the degeneracy associated to the space of wave functions for the single-vertex Hilbert space. This characteristic of our states is what we would expect from a causal barrier, suggesting that the foliation of space provided by our shell graphs could be naturally used to mimic a foliation by null surfaces. This does not obviously represent the most general case of foliation, but it suits very well the description of a black hole horizon. Moreover, if we were in a more general case where information about the bulk had not been washed away by the tracing operation, the implementation of weak holography for the horizon density matrix would have required some further restriction on the bulk wave functions, for instance coming from the dynamics. Due to the holographic nature of our particular states, such restriction is automatically included. What remains to be checked is the compatibility of these holographic states and their associated geometrical data with the imposition of the dynamics. This is far from trivial, as we have repeatedly stressed, and it is left for future work.

Finally, it is also remarkable that we could find an orthogonal set of eigenvectors for the horizon density matrix (48). This allows us to diagonalize it and then, due to the property (46), the calculation of the horizon von Neumann entropy can be performed precisely, without neglecting any entanglement contribution, and this corresponds exactly to a statistical counting *à la* Boltzmann. We have thus proven that the horizon entanglement entropy is the same as the horizon Boltzmann entropy. Let us now perform the counting.

## VI. COMBINATORIAL CONTRIBUTION: GRAPH COUNTING

Our states are defined via iterated action of refinement moves on the seed of each shell. The generic parametrization (12) makes use of a generic function of two sets ( $B, W$ ) of refinement operators. Written more explicitly,

$$\left( \sum_{n=0}^{\infty} \prod_s \prod_{m=1}^n (a_{r_0, B_s}^{n,m} \widehat{\mathcal{M}}_{r_0, B_s} + a_{r_0, W_s}^{n,m} \widehat{\mathcal{M}}_{r_0, W_s}) \right) |\tau\rangle. \quad (47)$$

The refinement move operators, together with the seed state, specify the general class of possible microscopic combinatorial structures, i.e., the specific portion of the



Fock space, that we are exploring. The parameters  $a$  control instead the relative weights of the different possible strings of refinement move operators that can be applied to the seed state. As we have said, we determine them following some global considerations, making use of a maximum entropy principle, appealing to typicality of the state given some mild general constraints.

Let us clarify this point further. From the perspective of macroscopic, large scale dynamics, making reference to the detailed combinatorial structure of the state does not seem a plausible approach. Instead, the state has to be determined as the most general state compatible with global constraints on topology, symmetry and semiclassicality. A statistical approach based on typicality and maximum ignorance seems to be more suitable, exploring as much as possible the space of states that we have described so far, and limiting the truncations to specific sectors of the GFT kinematical space to the ones that respect mild constraints associated to the physical regime we are interested in.

In Sec. IV B we have studied the properties of the reduced density matrix, showing a basis of states diagonalizing it. We showed how the problem of computing the entropy boils down to a (weighted) count of graphs, but we did not proceed further in the counting. We now resume the computation of the entropy by first evaluating the contribution arising from the graph proliferation.

The action of the refinement moves, via the Wick's theorem, generate a linear superposition of states possessing different combinatorial structure, according to each particular sequence of Wick contractions. Let  $G_n$  be the set of graphs that can be obtained via any sequence of  $n$  refinement moves per shell component starting from the given seed. The particular function in (12) then specifies a set of weights  $w_n(\Gamma)$ , the coefficients entering the diagonal of the reduced density matrix. The detailed dependence of these weights on the coefficients  $a$ 's is not necessary, but could be computed, in principle. Let us stress that the assumption that at each step a given horizon component gets refined with the same number of vertices as the other two is here made mainly to simplify the notation and it could be relaxed without affecting the final result of the entropy counting.

We have already shown that, for a given boundary graph, the reduced density matrix of the horizon takes the form (37), where  $A = r_0$  corresponds to the horizon shell, including all its three components. As argued, this form remains unchanged even when we trace away the whole bulk degrees of freedom (both interior and exterior ones). Therefore, we can write the total normalized reduced density matrix of the shell for a given number  $n$  of boundary vertices as

$$\rho_{\text{tot}, r_0}^{(n)} = \sum_{\alpha=1}^{\mathcal{N}} w_n(\Gamma_\alpha) \rho_{r_0}^{(n)}(\Gamma_\alpha), \quad (48)$$

where  $\mathcal{N} = \#G_n$  is the total number of shell graphs for a given number of vertices (which are  $2n$ , given the presence of two colors for the vertices), obtained through all the possible actions of the refinement operators. Here,  $\rho_{r_0}^{(n)}(\Gamma_\alpha)$  is the reduced density matrix in (37).

The presence of several graphs (at fixed topology) results into a combinatorial contribution to the entropy:

$$S_{\text{comb}} = - \sum_{\alpha=1}^{\mathcal{N}} w_n(\Gamma_\alpha) \log(w_n(\Gamma_\alpha)). \quad (49)$$

As explained in Sec. V, we determine the weights  $w_n$  [and then, implicitly, the function  $F_r$  in (12)] by maximizing this entropy. It is immediate to do so: the most disordered configuration is the one in which the weights are all equal,

$$w_n^{\text{max}}(\Gamma) = \frac{1}{\mathcal{N}}, \quad S_{\text{comb}}^{\text{max}} = \log(\mathcal{N}). \quad (50)$$

Therefore, the only thing left to determine is the size of the set of graphs that can be obtained from the given set of refinement moves.

The counting of the number of graphs generated by our refinement moves can be easily performed using familiar techniques. Let us focus on a given layer, for instance the outer boundary of the horizon shell. Following the convention adopted so far, let us suppress for a moment the edges of color 1, whose only role is to connect the horizon shell with the next one. They do not play a role in the counting since the state is completely determined by the specification of the combinatorial pattern of the gluings of edges of colors 2, 3, 4. This effectively reduces the counting problem to a counting of graphs obtained connecting 3-valent vertices (i.e., a lower dimensional problem, as should be expected).

At a first inspection, following the reasoning in [65], one would expect the counting of the boundary states to give us the Catalan numbers (for  $D = 2$ , in fact), as what we are looking at is the refinement of a melonic graph with the insertion of melons. However, some care is due in the case of the shell boundaries. In fact, we have only two moves at our disposal, inserting loops with colors (2,3) and (3,4) only [see (8) and (9)]. The insertion of loops with colors (2,4) would cause a crossing with an edge of color 1. Therefore the strategy of [65] has to be slightly adapted.

Notice that the refinement moves can be seen as the insertion of melons on links of color 2 and 4 only (this corresponds to the addition of tetrahedra incident on the same dual edge, i.e., the increase of the curvature around that edge). Therefore, our problem can be seen as the calculation of the number of ways in which we can insert, into a line of color 2 (for instance), a string of pairs of nodes connected according to our rules.

This is easily done using generating functions. Let  $u$  be a real (or complex) parameter such that the Taylor

coefficients of  $G(u)$  in 0, which we wish to determine, are the number of graphs obtained acting with our moves; i.e., the  $n$ th coefficient is the number of graphs with  $n$  pairs of nodes. The basic building blocks are 1PI graphs, which are then combined to get all the graphs.

The equation for the 1PI part can be obtained by inspection of the diagrammatics. We can obtain a recursion relation observing that, due to the peculiar connectivity, the only 1PI diagrams are the ones in which the first black vertex and the last white vertex are connected by a line of color 3. If this were not the case, one could cut the graph into two disconnected components removing the link between the white vertex connected with the first black vertex and the next black vertex. Let us call  $\Sigma(u)$  the function that generates the 1PI graphs. We have

$$G(u) = \frac{1}{1 + \Sigma(u)}. \quad (51)$$

However, it is easy to realize that  $\Sigma(u) = uG(u)$ , as the graphs that we would be counting are determined by all the possible ways of inserting  $n - 1$  pairs of nodes in the link that would connect the first and last node of the 1PI graph. The  $u$  prefactor keeps track of the initial pair to be inserted. Therefore,

$$G(u) = \frac{1}{1 + uG(u)}. \quad (52)$$

The solution, differentiable in 0, is

$$G(u) = \frac{-1 + \sqrt{1 + 4u}}{2u}, \quad (53)$$

which, as indeed expected, is the generating function for the Catalan numbers, namely

$$C_k = \frac{(2k)!}{(k+1)!k!}. \quad (54)$$

Therefore, the number of distinct graphs obtained with  $n$  refinement moves acting on  $l$  layers of the shell initial seed state is

$$\mathcal{N} = \left( \frac{(2n-2)!}{n!(n-1)!} \right)^l. \quad (55)$$

As explained above, in the horizon entropy counting we consider the most general case in which each layer is refined independently, i.e., we set  $l = 3$ . As it will be clear in a moment, considering a subclass of states in which the refinement proceeds in a more synchronized way among the different layers (and which therefore implements also a stronger form of holography) simply affects the numerical coefficient in front of the logarithmic corrections, but not the leading linear term.

Taking the large  $n$  limit of (55), in order to meet our semiclassicality requirements, by means of the Stirling formula we get

$$S_{\text{comb}}^{\text{max}} = \log(\mathcal{N}) = 2nl \log(2) - \frac{l}{2} \log(n). \quad (56)$$

It is interesting to note that this is not the first time that Catalan numbers enter the calculation of black hole entropy, see [66], albeit in a different context. We come back to this later.

## VII. FULL ENTROPY: FROM MACRO TO MICRO

The graph counting quantifies the contribution of the combinatorial degrees of freedom to the horizon entropy. This is not the entire entropy of the reduced state. There is also another, more geometrical, component which arises from the information attached to each vertex by the wave function: the degeneracy of the Hilbert space for a single vertex,  $\Delta(a)$ , for fixed values of the macroscopic quantities. This quantity measures the size of the space of wave functions compatible with our semiclassicality restrictions (and solution of the dynamics equations). Since the geometric and combinatorial components are independent, at this (kinematical) stage of the construction, the total horizon entropy is then

$$S(n, a) = \log(\mathcal{N}\Delta(a)) \\ = 2nl \log(2) + \log(\Delta(a)) - \frac{l}{2} \log(n). \quad (57)$$

This is the entropy of reduced matrix of the most typical state that we can construct with our special class of condensates, at fixed number of vertices.<sup>4</sup>

The next step is to maximize this entropy at fixed value of the classical area  $\mathcal{A}_{IH}$  of the horizon. Consider the function

$$\Sigma(n, a, \lambda) = S(n, a) + \lambda(\mathcal{A}_{IH} - 2an), \quad (58)$$

where  $\lambda$  is a Lagrange multiplier imposing the area constraint. Necessary conditions for the maximization of (58) are

$$\frac{\partial \Sigma}{\partial \lambda} = \mathcal{A}_{IH} - 2an = 0, \quad (59)$$

$$\frac{\partial \Sigma}{\partial n} \approx 2l \log(2) - 2\lambda a = 0, \quad (60)$$

<sup>4</sup>If the number of vertices is not fixed, an additional contribution should be added to include the effect of the dispersion in the number of vertices.

$$\frac{\partial \Sigma}{\partial a} = \frac{\Delta'(a)}{\Delta(a)} - 2n\lambda = 0, \quad (61)$$

where the prime indicates derivative with respect to  $a$ . In (60) we have dropped a term of order  $1/n$ , since we are in the large  $n$  regime, as the semiclassicality conditions of small fluctuations of the horizon geometrical properties imply. These are three equations for the values of the three quantities  $(n, a, \lambda)$  which maximize the entropy functional (58).

We could solve them explicitly only if we knew the expression the single vertex Hilbert space degeneracy  $\Delta$  as a function of  $a$ . In principle, its evaluation could be achieved by the imposition of the isolated horizon boundary condition, by computing the number of solutions to that equation compatible with the dynamics. However, both tasks, i.e., solving the isolated horizon boundary condition and the equations of motion defining the microscopic dynamics, are highly nontrivial and currently out of our reach, as we had already emphasized.

Alternatively, we could turn the argument around and use the three relations (59), (60), and (61) to get an explicit expression of  $a$  and  $\Delta(a)$  as functions of  $\mathcal{A}_{IH}$  and  $\lambda$ . At this point, then, we could require consistency with a further semiclassical property of a black hole horizon, namely its thermality, in order to determine the value of the Lagrange multiplier  $\lambda$  and thus remove the last ambiguity left in our entropy calculation. This gives a different twist to this discussion, turning it into the inference of how certain microscopic quantities should look if we want them to be compatible with macroscopic observations. Following this strategy, we obtain

$$a = \frac{l \log(2)}{\lambda}, \quad (62)$$

$$\Delta = c_0 \exp(2\lambda a n) = c_0 \exp(\lambda \mathcal{A}_{IH}), \quad (63)$$

where  $c_0$  is an integration constant left unspecified for the moment. The horizon entropy we derive is then

$$S(\mathcal{A}_{IH}) \approx 2\lambda \mathcal{A}_{IH} - \frac{l}{2} \log\left(\frac{\mathcal{A}_{IH}}{\ell_P^2}\right). \quad (64)$$

Notice that the solution (63) matches the expectation that the dimension of the Hilbert space for the wave function at fixed plaquette area should be finite once we implement the semiclassicality conditions listed in Sec. V. In particular, this would

- (1) exclude degenerate (zero-volume) configurations (which would be indeed incompatible with a reasonable semiclassical limit);
- (2) impose a finite volume of the shell, again consistent with the semiclassical intuition.

In other words, we should expect that whenever  $\Delta(a)$  diverges, we should not be worried about possible problems in the calculation of the entropy since the geometry would be highly problematic to be interpreted as a semiclassical one in the first place.

We can go further in our attempt of determine the properties of states that match the large scale behavior of classical gravity. Since the semiclassical limit involves large values of  $n$ , the area constraint (59) requires  $a$  to be small. In the limit of  $a \rightarrow 0$ , the IH boundary condition fixes the holonomy around each radial link to be flat; this can be achieved only if the spin labels of the tangent links are 0. Therefore, in the limit  $a \rightarrow 0$  the wave function should be a delta peaked on  $j_I = 0$ , which means  $\Delta(0) \sim 1$ . As soon as  $a > 0$ ,  $\Delta(a)$  should grow.<sup>5</sup> These expectations are matched by the Taylor expansion of the solution (63) around  $a = 0$ , if we fix the integration constant  $c_0 = 1$ , namely

$$\Delta(a) \sim 1 + 2\lambda a n, \quad (65)$$

for small  $a$ .

The free parameter  $\lambda$  can now be fixed by requiring consistency with the semiclassical thermodynamical condition relating the derivative of the entropy with respect to a local notion of energy at the horizon for fixed  $n$  to the (inverse) Unruh<sup>6</sup>  $1/T_U = \beta_U = 2\pi/(\ell_P^2 \kappa) = 2\pi\ell/\ell_P^2$ . In the previous expression,  $\ell$  is a local stationary observer proper distance and  $\kappa = 1/\ell$  is its surface gravity. For this purpose, we can use the local notion of energy introduced in [68] for isolated horizons, namely

$$\mathcal{E}_{IH} = \frac{\mathcal{A}_{IH}}{8\pi\ell}. \quad (66)$$

Dropping a subdominant term ( $\propto 1/\mathcal{A}_{IH}$ ), which becomes immaterial for large enough areas, we get

$$\beta_U = \frac{2\pi\ell}{\ell_P^2} = \frac{\partial S}{\partial \mathcal{E}_{IH}} = 8\pi\ell \frac{\partial S}{\partial \mathcal{A}_{IH}} \approx 16\pi\ell\lambda, \quad (67)$$

which leads, finally, to

$$\lambda \approx \frac{1}{8\ell_P^2}. \quad (68)$$

<sup>5</sup>Notice that these arguments about semiclassical limits are used only to constrain the functional dependence of  $\Delta(a)$  on  $a$ ; there is no requirement that  $j_I$  actually ever take the zero value, which may be problematic from the quantum geometric point of view.

<sup>6</sup>A geometric notion of temperature [39] can be associated to a quantum IH by demanding the Kubo-Martin-Schwinger condition [67] to be satisfied for a subalgebra of the holonomy-flux \* algebra of LQG. In the large IH area this notion of temperature coincides with the Unruh one.

Therefore, consistency with the IH thermodynamical properties yields an entropy

$$S(\mathcal{A}_{IH}) \approx \frac{\mathcal{A}_{IH}}{4\ell_P^2} - \frac{3}{2} \log\left(\frac{\mathcal{A}_{IH}}{\ell_P^2}\right) \quad (69)$$

whose leading term reproduces the Bekenstein-Hawking formula; the numerical coefficient in front of the subleading logarithmic correction comes from explicitly setting  $l = 3$  as argued above, and it matches the result of previous calculations in LQG [27,29,69] as well as those performed through CFT techniques [70].

### VIII. CONCLUSIONS AND FINAL REMARKS

We now briefly summarize the results obtained and observations made in this paper.

We have constructed, within the full quantum gravity formalism of group field theory, a class of quantum states that can be argued to describe (at least some key properties of) continuum spherically symmetric geometries. We have done this by making use of both its quantum geometric aspects, shared with loop quantum gravity, and its combinatorial aspects, shared with random tensor models. We have then imposed additional conditions on such states that support an interpretation of them as containing horizons, and thus describing black hole geometries. For such states, we have then identified the microscopic degrees of freedom contributing to the horizon entropy and computed the latter explicitly. Under appropriate semiclassicality restrictions and the assumption that the entropy is maximized, we have recovered both a general area law and the exact Bekenstein-Hawking value. The fact that our states are realistic quantum states in the full theory, involving also a highly nontrivial sum over graphs, including arbitrarily refined ones, is one main improvement over the existing derivations of the same results in the loop quantum gravity literature. Let us now mention a few additional interesting aspects of our analysis and results.

The entropy result (69) is completely independent on the Barbero-Immirzi parameter  $\gamma$ . This is a striking consequence of the GFT formalism in its Fock representation and the way in which it allows us to work with quantum gravity states. More precisely, the GFT fundamental field operators (2), encoding the state geometrical data in the collective wave function  $\sigma_r$  for a given shell, represent the key departing point from canonical LQG. It allows us to introduce a well-defined number operator in the theory and replace area eigenstates and eigenvalues for the quantum horizon with condensate states and area expectation values on a single-vertex state (which is the same for each fundamental block due to the condensate hypothesis). In this way, while  $\gamma$  still enters in the value of  $a$ , it completely disappears in the final expression of the entropy. Another way to understand the same important point is that this is a consequence of our choice of states in a

second quantized formalism (reasonably simple condensates), allowing us to do a microscopic counting of degrees of freedom, while not using eigenstates of the total area operator. The latter, customarily used in LQG, besides being dubious as corresponding to semiclassical black hole states, immediately introduce a dependence on the Barbero-Immirzi parameter, since this appears inevitably in the area spectrum. This also means that obtaining our results, including this intriguing independence from the Barbero-Immirzi parameter, has been facilitated greatly by the GFT formalism, with its convenient organization of spin-network states into a Fock space (with the mentioned existence of a number operator), and the tools it provides for handling sums over different triangulations/graphs. The reason for this independence from the Barbero-Immirzi parameter can also be traced back to the appearance of (63) (see the related discussion in the text). It is not just the area value that matters, but also the single-vertex density of states. The calculations that we have given above suggest that this should lead to terms compensating  $\gamma$  in the physical quantities. They also suggest that LQG calculations might be overlooking an essential step, signaled by the appearance of the Barbero-Immirzi parameter in expressions for quantities that should be independent of it, at least at the classical level. The evaluation of the expectation values of operators on properly defined semiclassical states (which are not necessarily eigenvectors of the area operator for some fixed graphs structure) seems to be a critical calculation, in this respect.

The graph basis provides, in this case, an orthonormal basis for the reduced density matrix. This, together with the holographic property of our states, allows us to compute the horizon entanglement entropy exactly and prove the result

$$\begin{aligned} S_{\text{von Neumann}} &\equiv -\text{Tr}(\rho_{\text{tot},r_0}^{(n)} \log(\rho_{\text{tot},r_0}^{(n)})) \\ &= \log(\mathcal{N}(n)) \equiv S_{\text{Boltzmann}}, \end{aligned} \quad (70)$$

which then reduces to a relatively simple counting exercise. As we have already mentioned, the particular class of states that we have chosen, obtained by what are very close to be melonic refinement of seed states, has a very specific growth rate with the number of vertices. This growth has also been considered in [66], in a different context and with a different picture in mind. Despite all the limitations that have been discussed, these results point at a specific behavior for physical quantum gravity states describing black holes. Together with the observation about the disappearance of the Barbero-Immirzi parameter, related to the density of physical states per vertex, these facts suggest that the superposition of microscopic kinematical configurations in the physical states cannot be neglected in such calculations, and it might also have to be severely constrained.

Following up on the last point, the dynamics has not yet been used, essentially. Our results can be seen as general,



therefore, only provided that the states we have used turn out to be good representatives of the true microstates of physical black holes, not only encoding in a more precise manner the various properties that we expect from such particular space-time geometries (at least semiclassically) and that we have imposed only implicitly, but also solving at least approximately the microscopic quantum dynamics of the theory.

Given the experience in the cosmological setting, one should expect that the imposition of the GFT equations of motion will result in some nonlinear (set of) effective equation(s) on the wave function and on the coefficients of the linear combinations defining our quantum states in terms of graphs. From the discussion above, it might be relatively easy, once even only a qualitative understanding of the structure of some of the solutions to the equations is achieved, to check the compatibility of the dynamics with macroscopic properties, as quantities as rate of growth of graphs with the number of quanta are already rather constrained. We leave this type of analysis for future work, though.

The maximum entropy principle that we are used, itself, may or may not be compatible with the full quantum dynamics. At this stage, we use it to fix the shape of the states given the available information and try to infer consequences about the nature of the states, but nothing more. It goes without saying that it would be even more significant if we could recover this reasoning as an end point of a calculation that starts from the microscopic equations, especially given the deep relations between gravity and thermodynamics.

### ACKNOWLEDGMENTS

Research at the Perimeter Institute for Theoretical Physics is supported in part by the Government of Canada through Natural Sciences and Engineering Research Council and by the Province of Ontario through Ministry of Research, Innovation and Science.

### APPENDIX: BASICS OF GFT FOCK SPACE

In this appendix we briefly summarize the necessary ideas about group field theories that are needed in the main body of the paper, in order to make the text as self-contained as possible. We closely follow [45]. A more thorough discussion can be found in [9,10].

In a nutshell, GFTs are a generalization of matrix models for two-dimensional quantum gravity. They are quantum field theories over group manifolds whose perturbative expansion generates Feynman diagrams whose combinatorial structure can be put in correspondence with triangulations of higher dimensional manifolds. Besides extending the combinatorial structure of the Feynman rules, they include additional data which can be used to store additional geometric information. In particular, these

data allow the construction of models which can be put in correspondence with LQG and spin foam models. While the formalism *per se* leaves considerable freedom for model building, we restrict the discussion to the simplest models that are relevant for 3 + 1 quantum gravity.

One possible presentation of GFTs is the one based on the Fock space representation, which starts from the idea that the basic building block of a theory, the elementary quantum excitation, is a single four-valent spin network vertex. The basic ingredients are group field ladder operators  $\hat{\varphi}, \hat{\varphi}^\dagger$ , defined over the group  $SU(2)^4$ . The argument  $g$  of one such field operator is then the tuple  $(g_1, g_2, g_3, g_4)$ .

A geometric interpretation is then attached to this type of data: each  $SU(2)$  argument is interpreted as the parallel transport, induced by a gauge connection, from a four-valent vertex to an end point of an edge emanating from it. For consistency with this interpretation, then, a gauge invariance property is enforced in terms of the invariance of the field operators under diagonal right action of  $SU(2)$ ,

$$\hat{\varphi}(gk_{\text{Diag}}) = \hat{\varphi}(g), \quad k_{\text{Diag}} = (k, k, k, k), \quad \forall k \in SU(2). \quad (\text{A1})$$

Notice that, with a slight abuse of notation, we refer to this sort of gauge invariance by using directly the  $SU(2)$  group elements, even when the diagonal action on  $SU(2)^4$  is intended.

The algebra of bosonic ladder operators is determined then by the commutation relations,

$$[\hat{\varphi}(g_v), \hat{\varphi}^\dagger(g'_v)] = \Delta_R(g_v, g'_v), \quad (\text{A2})$$

where we are using the following definition for the right-invariant Dirac delta distribution on  $SU(2)^4$ ,

$$\Delta_R(g_v, g'_v) = \int_{SU(2)} dh \prod_{I=1}^4 \delta(g_{v,I} h g_{v',I}^{-1}), \quad (\text{A3})$$

and  $\delta(g)$  denotes the ordinary delta function over a single copy of  $SU(2)$ .

The role of these ladder operators is then to create and annihilate objects with all the properties of spin-network vertices. Using the noncommutative Fourier transform on these basic operators gives rise to a representation in which the fundamental ladder operators correspond to elementary quanta labeled with the data of quantum tetrahedra.

Generic states in the Fock space defined by these representations, then, correspond to sets of quantum tetrahedra or spin network vertices variously arranged. Among these states there are the ones corresponding to closed connected spin networks, or, equivalently, triangulated three-dimensional manifolds. The properties of the states can be encoded into wave functions, as usual. For instance, a generic  $n$ -particle state can be written as

$$|\psi\rangle = \int_{(\text{SU}(2)^4)^n} \prod_{i=1}^n dg_{v_i} \left( \psi(g_{v_1}, \dots, g_{v_n}) \prod_{i=1}^n \hat{\varphi}^\dagger(g_{v_i}) \right) |0\rangle, \quad (\text{A4})$$

where  $|0\rangle$  is the Fock vacuum. The wave function  $\psi$  provides the bridge between GFT and the language of LQG [10]. In particular, by controlling the dependence on the group elements, the familiar wave functions of closed, gauge invariant spin network attached to graphs can be recovered.

This kinematic construction is completed by definition of suitable dynamics, which we do not describe in detail here. Typical choices for the equation of motion obeyed by a physical state  $|\Psi\rangle$  are

$$\frac{\delta \mathcal{S}}{\delta \hat{\varphi}(g)} [\hat{\varphi}, \hat{\varphi}^\dagger] |\Psi\rangle = 0 \quad (\text{A5})$$

where the action  $\mathcal{S}$  is designed in such a way that the Feynman expansion of the theory gives rise to diagrams that can be put in correspondence with four-dimensional simplicial complexes, decorated with appropriate geometric data. The various kernels appearing in the field monomials, then, can be adjusted to reproduce spin foam amplitudes.

The advantage of the Fock representation stems from the fact that the equation of motion (A5) does not generically single out states entirely contained in an eigenspace of the Fock number operator. The Fock representation of the theory makes it very easy to work with states which are superpositions of states with a different number of quanta, leveraging on the techniques familiar in the domains of quantum optics and condensed matter physics. The ability to do so is crucial if we want to obtain even an approximate description of the continuum regime following from (A5), where the notion of elementary quanta dissolves in the continuum soup.

- 
- [1] *Astrophysics of Black Holes: From Fundamental Aspects to Latest Developments*, edited by C. Bambi (Springer, Berlin/Heidelberg, 2016), Vol. 440.
- [2] L. Rezzolla, An introduction to astrophysical black holes and their dynamical production, *Lect. Notes Phys.* **905**, 1 (2016).
- [3] J. D. Bekenstein, Black holes and the second law, *Lett. Nuovo Cimento* **4**, 737 (1972).
- [4] J. M. Bardeen, B. Carter, and S. W. Hawking, The four laws of black hole mechanics, *Commun. Math. Phys.* **31**, 161 (1973).
- [5] J. D. Bekenstein, Black holes and entropy, *Phys. Rev. D* **7**, 2333 (1973).
- [6] S. W. Hawking, Black hole explosions, *Nature (London)* **248**, 30 (1974).
- [7] S. W. Hawking, Particle creation by black holes, *Commun. Math. Phys.* **43**, 199 (1975).
- [8] S. Carlip, Black hole thermodynamics, *Int. J. Mod. Phys. D* **23**, 1430023 (2014).
- [9] D. Oriti, Group field theory as the second quantization of loop quantum gravity, *Classical Quantum Gravity* **33**, 085005 (2016).
- [10] D. Oriti, <https://inspirehep.net/record/1312968/files/arXiv:1408.7112.pdf>.
- [11] R. Gurau and J. P. Ryan, Colored tensor models—a review, *SIGMA* **8**, 020 (2012).
- [12] C. Rovelli, *Quantum Gravity*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, 2004).
- [13] T. Thiemann, *Modern Canonical Quantum General Relativity* (Cambridge University Press, Cambridge, 2008).
- [14] A. Ashtekar, C. Beetle, and S. Fairhurst, Isolated horizons: A generalization of black hole mechanics, *Classical Quantum Gravity* **16**, L1 (1999).
- [15] A. Ashtekar, C. Beetle, and S. Fairhurst, Mechanics of isolated horizons, *Classical Quantum Gravity* **17**, 253 (2000).
- [16] J. Diaz-Polo and D. Pranzetti, Isolated horizons and black hole entropy in loop quantum gravity, *SIGMA* **8**, 048 (2012).
- [17] A. Perez, Black holes in loop quantum gravity, *Rep. Prog. Phys.* **80**, 126901 (2017).
- [18] A. Ashtekar, A. Corichi, and K. Krasnov, Isolated horizons: The classical phase space, *Adv. Theor. Math. Phys.* **3**, 419 (1999).
- [19] J. Engle, K. Noui, A. Perez, and D. Pranzetti, Black hole entropy from an SU(2)-invariant formulation of type-I isolated horizons, *Phys. Rev. D* **82**, 044050 (2010).
- [20] A. Ashtekar, J. C. Baez, and K. Krasnov, Quantum geometry of isolated horizons and black hole entropy, *Adv. Theor. Math. Phys.* **4**, 1 (2000).
- [21] K. A. Meissner, Black hole entropy in loop quantum gravity, *Classical Quantum Gravity* **21**, 5245 (2004).
- [22] M. Domagala and J. Lewandowski, Black hole entropy from quantum geometry, *Classical Quantum Gravity* **21**, 5233 (2004).
- [23] A. Ghosh and P. Mitra, An improved lower bound on black hole entropy in the quantum geometry approach, *Phys. Lett. B* **616**, 114 (2005).
- [24] A. Corichi, J. Diaz-Polo, and E. Fernandez-Borja, Black Hole Entropy Quantization, *Phys. Rev. Lett.* **98**, 181301 (2007).
- [25] A. Corichi, J. Diaz-Polo, and E. Fernandez-Borja, Quantum geometry and microscopic black hole entropy, *Classical Quantum Gravity* **24**, 243 (2007).
- [26] I. Agullo, J. F. Barbero, J. Diaz-Polo, E. Fernandez-Borja, and E. J. S. Villasenor, Black Hole State Counting in LQG: A Number Theoretical Approach, *Phys. Rev. Lett.* **100**, 211301 (2008).

- [27] R. K. Kaul and P. Majumdar, Logarithmic Correction to the Bekenstein-Hawking Entropy, *Phys. Rev. Lett.* **84**, 5255 (2000).
- [28] J. F. Barbero and E. J. S. Villasenor, On the computation of black hole entropy in loop quantum gravity, *Classical Quantum Gravity* **26**, 035017 (2009).
- [29] J. Engle, K. Noui, A. Perez, and D. Pranzetti, The SU(2) black hole entropy revisited, *J. High Energy Phys.* **05** (2011) 016.
- [30] A. Perez and D. Pranzetti, Static isolated horizons: SU(2) invariant phase space, quantization, and black hole entropy, *Entropy* **13**, 744 (2011).
- [31] E. Frodden, A. Perez, D. Pranzetti, and C. Röken, Modeling black holes with angular momentum in loop quantum gravity, *Gen. Relativ. Gravit.* **46**, 1828 (2014).
- [32] L. Smolin, Linking topological quantum field theory and nonperturbative quantum gravity, *J. Math. Phys.* **36**, 6417 (1995).
- [33] K. V. Krasnov, On quantum statistical mechanics of Schwarzschild black hole, *Gen. Relativ. Gravit.* **30**, 53 (1998).
- [34] C. Rovelli, Black Hole Entropy from Loop Quantum Gravity, *Phys. Rev. Lett.* **77**, 3288 (1996).
- [35] T. Jacobson, Renormalization and black hole entropy in loop quantum gravity, *Classical Quantum Gravity* **24**, 4875 (2007).
- [36] A. Ghosh and A. Perez, Black Hole Entropy and Isolated Horizons Thermodynamics, *Phys. Rev. Lett.* **107**, 241301 (2011); Erratum, *Phys. Rev. Lett.* **108**, 169901 (2012).
- [37] E. Frodden, M. Geiller, K. Noui, and A. Perez, Black hole entropy from complex Ashtekar variables, *Europhys. Lett.* **107**, 10005 (2014).
- [38] A. Ghosh, K. Noui, and A. Perez, Statistics, holography, and black hole entropy in loop quantum gravity, *Phys. Rev. D* **89**, 084069 (2014).
- [39] D. Pranzetti, Geometric temperature and entropy of quantum isolated horizons, *Phys. Rev. D* **89**, 104046 (2014).
- [40] N. Bodendorfer and Y. Neiman, Imaginary action, spin foam asymptotics and the trans-Planckian regime of loop quantum gravity, *Classical Quantum Gravity* **30**, 195018 (2013).
- [41] A. Ghosh and D. Pranzetti, CFT/gravity correspondence on the isolated horizon, *Nucl. Phys.* **B889**, 1 (2014).
- [42] J. B. Achour, A. Mouchet, and K. Noui, Analytic continuation of black hole entropy in loop quantum gravity, *J. High Energy Phys.* **06** (2015) 145.
- [43] J. B. Achour, K. Noui, and A. Perez, Analytic continuation of the rotating black hole state counting, *J. High Energy Phys.* **08** (2016) 149.
- [44] L. Freidel, A. Perez, and D. Pranzetti, Loop gravity string, *Phys. Rev. D* **95**, 106002 (2017).
- [45] D. Oriti, D. Pranzetti, J. P. Ryan, and L. Sindoni, Generalized quantum gravity condensates for homogeneous geometries and cosmology, *Classical Quantum Gravity* **32**, 235016 (2015).
- [46] S. Gielen, D. Oriti, and L. Sindoni, Cosmology from Group Field Theory Formalism for Quantum Gravity, *Phys. Rev. Lett.* **111**, 031301 (2013).
- [47] S. Gielen, D. Oriti, and L. Sindoni, Homogeneous cosmologies as group field theory condensates, *J. High Energy Phys.* **06** (2014) 013.
- [48] D. Oriti, L. Sindoni, and E. Wilson-Ewing, Emergent Friedmann dynamics with a quantum bounce from quantum gravity condensates, *Classical Quantum Gravity* **33**, 224001 (2016).
- [49] S. Gielen and L. Sindoni, Quantum cosmology from group field theory condensates: A review, *SIGMA* **12**, 082 (2016).
- [50] D. Oriti, D. Pranzetti, and L. Sindoni, Horizon Entropy from Quantum Gravity Condensates, *Phys. Rev. Lett.* **116**, 211301 (2016).
- [51] A. Baratin, S. Carrozza, D. Oriti, J. Ryan, and M. Smerlak, Melonic phase transition in group field theory, *Lett. Math. Phys.* **104**, 1003 (2014).
- [52] H. Sahlmann, Black hole horizons from within loop quantum gravity, *Phys. Rev. D* **84**, 044049 (2011).
- [53] D. Pranzetti and H. Sahlmann, Horizon entropy with loop quantum gravity methods, *Phys. Lett. B* **746**, 209 (2015).
- [54] H. Sahlmann and T. Thiemann, Chern-Simons theory, Stokes' theorem, and the Duflo map, *J. Geom. Phys.* **61**, 1104 (2011).
- [55] H. Sahlmann and T. Thiemann, Chern-Simons Expectation Values and Quantum Horizons from LQG and the Duflo Map, *Phys. Rev. Lett.* **108**, 111303 (2012).
- [56] K. Noui, A. Perez, and D. Pranzetti, Canonical quantization of noncommutative holonomies in 2 + 1 loop quantum gravity, *J. High Energy Phys.* **10** (2011) 036.
- [57] K. Noui, A. Perez, and D. Pranzetti, Noncommutative holonomies in 2 + 1 LQG and Kauffman's brackets, *J. Phys. Conf. Ser.* **360**, 012040 (2012).
- [58] D. Pranzetti, Turaev-Viro amplitudes from 2 + 1 loop quantum gravity, *Phys. Rev. D* **89**, 084058 (2014).
- [59] W. Donnelly, Entanglement entropy in loop quantum gravity, *Phys. Rev. D* **77**, 104006 (2008).
- [60] G. Chirco, F. M. Mele, D. Oriti, and P. Vitale, Fisher metric, geometric entanglement, and spin networks, *Phys. Rev. D* **97**, 046015 (2018).
- [61] S. N. Solodukhin, Entanglement entropy of black holes, *Living Rev. Relativity* **14**, 8 (2011).
- [62] J. D. Bekenstein, A universal upper bound on the entropy to energy ratio for bounded systems, *Phys. Rev. D* **23**, 287 (1981).
- [63] R. Bousso, A covariant entropy conjecture, *J. High Energy Phys.* **07** (1999) 004.
- [64] L. Smolin, The strong and weak holographic principles, *Nucl. Phys.* **B601**, 209 (2001).
- [65] V. Bonzom, R. Gurau, A. Riello, and V. Rivasseau, Critical behavior of colored tensor models in the large N limit, *Nucl. Phys.* **B853**, 174 (2011).
- [66] A. Davidson, Holographic shell model: Stack data structure inside black holes?, *Int. J. Mod. Phys.* **D23**, 1450041 (2014).
- [67] R. Haag, *Local Quantum Physics: Fields, Particles, Algebras* (Springer-Verlag, Berlin, Heidelberg, 1992).
- [68] E. Frodden, A. Ghosh, and A. Perez, Quasilocal first law for black hole thermodynamics, *Phys. Rev. D* **87**, 121503 (2013).
- [69] E. R. Livine and D. R. Terno, Quantum black holes: Entropy and entanglement on the horizon, *Nucl. Phys.* **B741**, 131 (2006).
- [70] S. Carlip, Logarithmic corrections to black hole entropy from the Cardy formula, *Classical Quantum Gravity* **17**, 4175 (2000).