

Hydrostatic equilibrium of stars without electroneutrality constraint

M. I. Krivoruchenko,^{1,2} D. K. Nadyozhin,^{1,3} and A. V. Yudin^{1,*}

¹*Institute for Theoretical and Experimental Physics, B. Chermushkinskaya 25, 117218 Moscow, Russia*

²*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Russia*

³*Kurchatov Institute, Akademika Kurchatova Pl. 1, 123182 Moscow, Russia*



(Received 26 February 2018; published 30 April 2018)

The general solution of hydrostatic equilibrium equations for a two-component fluid of ions and electrons without a local electroneutrality constraint is found in the framework of Newtonian gravity theory. In agreement with the Poincaré theorem on analyticity and in the context of Dyson's argument, the general solution is demonstrated to possess a fixed (essential) singularity in the gravitational constant G at $G = 0$. The regular component of the general solution can be determined by perturbation theory in G starting from a locally neutral solution. The nonperturbative component obtained using the method of Wentzel, Kramers and Brillouin is exponentially small in the inner layers of the star and grows rapidly in the outward direction. Near the surface of the star, both components are comparable in magnitude, and their nonlinear interplay determines the properties of an electro- or ionosphere. The stellar charge varies within the limits of -0.1 to 150 C per solar mass. The properties of electro- and ionospheres are exponentially sensitive to variations of the fluid densities in the central regions of the star. The general solutions of two exactly solvable stellar models without a local electroneutrality constraint are also presented.

DOI: [10.1103/PhysRevD.97.083016](https://doi.org/10.1103/PhysRevD.97.083016)

I. INTRODUCTION

Charged particles constitute an appreciable or dominant fraction of the matter in stars. The electromagnetic interaction between particles is stronger than the gravitational interaction by a factor of

$$\lambda_G = \frac{e^2}{Gm_u^2} \approx 1.25 \times 10^{36}, \quad (1.1)$$

where e denotes the proton charge in CGS units, G is the gravitational constant, and $m_u = 931.5 \text{ MeV}/c^2$ is the atomic mass unit, with c being the speed of light. Under the assumption of the complete screening of electrostatic interactions, gravitational forces should play the dominant role in determining stellar structure. The hydrostatic equilibrium (HE) equations are usually supplemented with a local electroneutrality constraint (LEC); however, it is not immediately clear how effective the screening is at compensating for the enormous difference in the strengths of the electromagnetic and gravitational forces.

Lebowitz and Lieb [1] showed that the LEC is required for the existence of extensive thermodynamic potentials of isolated systems. The stability of electrically neutral, isolated systems of charged particles in the thermodynamic limit was proven by Dyson and Lenard [2] and by Lieb and Thirring [3].

The thermodynamic equilibrium in an external potential field is determined by Gibbs' condition [4]:

$$\mu + V(x) = \text{const}, \quad (1.2)$$

where μ is the chemical potential and $V(x)$ is the external potential. A generalization of Eq. (1.2) for a multi-component fluid within the framework of general relativity theory (GRT) was given by O. Klein [5] and, in a broader context, by Kodama and Yamada [6] and Olson and Bailyn [7]. In the presence of gravity or other external fields, the chemical potentials of a multicomponent fluid acquire coordinate dependence, as already seen from Gibbs' condition, which is generally incompatible with the LEC.

As early as 1924, Rosseland [8] showed that in Newtonian gravity theory (NGT), the unconstrained HE equations for an ionized self-gravitating gas with a constant temperature admit solutions with similar distributions of ions and electrons. The charge density of the gas was found to be positive and λ_G times smaller than the particle number density. In the context of white dwarfs, the polarization of stars was discussed by Schatzman [9].

The smallness of the uncompensated charge density justifies the use of the LEC when describing effects not related to the electrostatic properties of stars.

Based on the variational energy principle of GRT, Olson and Bailyn [7] derived a complete system of unconstrained HE equations for a charged multicomponent fluid. In the stellar structure models discussed by Olson and Bailyn

* yudin@itep.ru

[10], Bally and Harrison [11], and Neslušan [12], the polarization effect leads to a stellar charge of $Q_s \sim eM_s/(m_u\lambda_G) \sim 100M_s/M_\odot$ C.

Rotondo *et al.* [13] and Belvedere *et al.* [14,15] showed recently that the components of a fluid have different spatial distributions, leading to the appearance of an electrosphere, the charge of which is sufficiently large to completely compensate for the positive charge of the star.

The deviations from local electroneutrality and, accordingly, the appearance of a large-scale electrostatic field in a star affects the diffusion and separation of chemical elements in the Sun [16] and in neutron stars [17]. The uncompensated stellar charge could influence possible mechanisms of supernova explosions [18–20].

Electrically charged stars are described in Refs. [8–15, 21] by particular solutions of the unconstrained HE equations. Some features of the general solution can be foreseen on the basis of simple qualitative considerations, as discussed below.

Let us consider a neutron star composed of neutrons, electrons and protons. Suppose that it is electrically neutral and in chemical equilibrium with respect to the weak interaction. The gravitational field of the star creates a potential well, in which a number of electrons ΔN_e can be confined, given that the energy of their electrostatic repulsion $\sim e^2\Delta N_e^2/R_s$ does not exceed the energy of their gravitational attraction to the star $\sim GM_s\Delta N_em_e/R_s$, where R_s and M_s are the radius and mass of the star, respectively, and m_e is the electron mass. This condition implies that $\Delta N_e \lesssim GM_sm_e/e^2 \sim 5 \times 10^{17}M_s/M_\odot$. The charge of a neutron star may vary from ~ -0.1 C to zero. An excess of electrons, however, would violate the assumption of chemical equilibrium. The relaxation of the neutron star to a new state through the inverse β decay is accompanied by the emission of energy in the form of neutrinos. The star then turns into a lower-energy state with the same electric charge. If gravitational stability is not lost during the relaxation process, then the above estimate remains valid. Considering that the number of electrons ΔN_e is many orders of magnitude smaller than the number of neutrons $M_\odot/m_u \approx 1.2 \times 10^{57}$, the neutron star remains gravitationally stable. Similar arguments lead to the conclusion that the neutron star is capable of confining $\Delta N_p \lesssim GM_sm_p/e^2 \sim 10^{21}M_s/M_\odot$ additional protons with a total charge not exceeding $Q_s \sim 150M_s/M_\odot$ C. The electric charge of a neutron star can vary from -0.1 to 150 C per solar mass. Similar considerations apply to white dwarfs.

The general solution of the unconstrained HE equations is expected to describe gravitationally stable configurations with a stellar charge of -0.1 to 150 C per solar mass.

A single-component self-gravitating fluid of hypothetical particles of mass m_f and charge ef , with $f \ll 1$, is discussed by S. Ray *et al.* [22]. A comparison of the gravitational force $\sim Gm_fM_s$ and the electrostatic repulsion force $\sim (ef)^2M_s/m_f$ acting on an elementary volume of the

fluid yields $f \sim m_f/(m_u\sqrt{\lambda_G})$ and a maximum stellar charge of $Q_s \sim efM_s/m_f \sim 10^{20}M_s/M_\odot$ C. In nature, stars are composed of multicomponent fluids and are subject to a $\sqrt{\lambda_G}$ times more stringent constraint on Q_s because of charge separation in the stellar electric fields.

Another fundamental property of the general solution follows from the Poincaré theorem on analyticity (see, e.g., Ref. [23]) and Dyson's argument [24].

The Poincaré theorem on analyticity is an intuitively clear theorem that in rough terms states that analyticity regions in the parameter space of ordinary differential equations (ODEs) are normally inherited by solutions of these equations. Analytic functions are determined by their singularities, so one can talk also about the singularities instead of the analyticity regions. It is not difficult to see that the unconstrained HE equations are singular at $G = 0$ or, equivalently, at $\lambda_G = \infty$. According to the Poincaré theorem, the general solution is thus expected to be singular at $\lambda_G = \infty$ as well. One of the unconstrained HE equations becomes an LEC in the limit of $\lambda_G = \infty$. It would be natural to consider the LEC solution as a zeroth-order approximation of the general solution in the framework of perturbation theory in the small parameter $1/\lambda_G$. The Poincaré theorem suggests, however, that as long as $\lambda_G = \infty$ is a singular point, such an expansion fails because the radius of convergence of the perturbation series is zero. Dyson's argument is interpreted similarly, with the same conclusions for convergence of the perturbation series.

The general solution to the unconstrained HE equations is therefore expected to be nonperturbative, which implies that the density distributions of the fluid components are singular at $\lambda_G = \infty$. Such solutions are typical of singularly perturbed ODE systems (see, e.g., [25]) whose order changes in the limit of a vanishing small parameter. The unconstrained HE equations belong to this class of singularly perturbed ODE systems, with the small parameter being $1/\lambda_G$.

In this paper, the properties of the general solution to the unconstrained HE equations are investigated within the NGT framework. The outline of the paper is as follows: Section II presents the unconstrained HE equations for a two-component fluid of ions and electrons described by polytropic equations of state (EoS). A unique regular solution at $\lambda_G = \infty$, which is a particular solution to these equations, is analyzed in detail. A regular solution can be constructed using perturbation theory in the small parameter $1/\lambda_G$, starting from the LEC solution. The solutions to the unconstrained HE equations that have previously been discussed in Refs. [8–17,21] are identified as such regular solutions. The regular solutions are investigated (i) for equal polytropic indices of ions and electrons, (ii) by expanding the solution in a power series in the stellar radius near the center, and (iii) by considering an expansion in $1/\lambda_G$ around the LEC solution. In Sec. III, the electron shell of a star is discussed. In Sec. IV, we consider an exactly solvable model with unit polytropic indices of ions and

electrons, which perfectly illustrates the non-perturbative nature of the general solution and clarifies the important role of the irregular part of the general solution in the formation of the electro- and ionospheres of stars. The mass, radius and charge of a star are found as functions of the central baryon number and charge densities. The exponential sensitivity of the electro- and ionospheric characteristics to the central charge density is established. A useful example of an exactly solvable model of a charged star consisting of a two-component incompressible fluid is further discussed in the Appendix. In Sec. V, we construct the general solution to the unconstrained HE equations for arbitrary polytropic indices. In the central stellar region, the irregular, non-perturbative part of the general solution can be found using perturbation theory of singular ODE systems (see, e.g., [25]). Such problems arise in the semiclassical limit of the Schrödinger equation, where the method of Wentzel, Kramers and Brillouin, commonly known as the WKB approximation, is proven to be effective. Perturbation theory applies everywhere, except in a thin subsurface layer of the star with a thickness of $\sim 1/\sqrt{\lambda_G}$. To match the internal solution to the outer electron or ion shell, numerical methods can be employed. The conclusions section summarizes the results.

II. REGULAR SOLUTION TO THE UNCONSTRAINED HYDROSTATIC EQUILIBRIUM EQUATIONS

We consider the problem of the polarization of a substance in a spherically symmetric star. The thermodynamic equilibrium equations for the chemical potentials of the ions, μ_i , and the electrons, μ_e , coincide with Gibbs' condition (1.2) and can be written as

$$\mu_i + m_i \varphi_G + Ze \varphi_E = \text{const}, \quad (2.1)$$

$$\mu_e + m_e \varphi_G - e \varphi_E = \text{const}, \quad (2.2)$$

where φ_G and φ_E are the gravitational and electrostatic potentials, eZ and $m_i = Am_u$ are the charge and mass of the ions. Bearing in mind possible applications to physics of white dwarfs, we set temperature equal to zero, and suppose the dominance of the contribution of electrons in thermodynamic functions. The chemical equilibrium with respect to the weak interaction is neglected. We also assume the ion and electron EoS are polytropic with $P \sim n^\gamma$ and attempt to reduce the problem to one known in the theory of polytropes (see, e.g., Ref. [26]).

A. Two-fluid model

The gravitational and electrostatic potentials satisfy the Poisson equations

$$\Delta \varphi_G = 4\pi G \rho_m, \quad (2.3)$$

$$\Delta \varphi_E = -4\pi \rho_e, \quad (2.4)$$

where the mass and charge densities are, respectively,

$$\rho_m = m_i n_i + m_e n_e, \quad (2.5)$$

$$\rho_e = Ze n_i - e n_e. \quad (2.6)$$

The radial derivative term of the Laplacian Δ is equal to

$$\Delta_r = \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \right).$$

By applying Δ to Eqs. (2.1) and (2.2) and constructing appropriate linear combinations of the chemical potentials, we obtain

$$\Delta_r (Z\mu_e + \mu_i) = -4\pi G (m_i + Zm_e) \rho_m, \quad (2.7)$$

$$\Delta_r (m_i \mu_e - m_e \mu_i) = -4\pi e (m_i + Zm_e) \rho_e. \quad (2.8)$$

According to Eqs. (2.1) and (2.2), the gradient $-\nabla(Z\mu_e + \mu_i)$ is equal to the gravitational force acting on an ion and Z electrons. The thermodynamic relationship $dP = nd\mu$ allows the gradients of the chemical potentials to be presented in terms of pressure gradients. As a special case, the hydrostatic limit of the Euler equation is recovered for an electroneutral fluid with $n_e = Zn_i$:

$$\frac{1}{\rho_m} \nabla(P_i + P_e) + \nabla \varphi_G = 0.$$

As mentioned above, we assume that $P_i \ll P_e$ holds everywhere.

First, we define the standard relations between pressures and concentrations from the theory of polytropes:

$$P_k = K_k n_k^{1+1/\eta_k}, \quad k = (i, e), \quad (2.9)$$

where K_k and η_k are fixed constants and η_k is called the polytropic index. The ion and electron components interact with each other only through the gravitational and electrostatic forces. No fundamental difficulties arise when considering the influence of the interactions of the components (see, e.g., [27]), although this consideration requires special attention. The dimensionless functions θ_k are defined as follows

$$n_k \equiv n_{k0} \theta_k^{\eta_k}, \quad (2.10)$$

where the n_{k0} are the concentrations at $r = 0$. Excluding the rest mass, the chemical potentials are equal to $\mu_k = \mu_{k0} \theta_k$, where $\mu_{k0} = K_k (1 + \eta_k) n_{k0}^{1/\eta_k}$. The functions $\theta_k = \theta_k(x)$ depend on a dimensionless coordinate x , which is related to r as follows:

$$r = r_0 x, \quad r_0^2 = \frac{Z\mu_{e0}}{4\pi G m_i (m_i + Zm_e) n_{i0}}. \quad (2.11)$$

These expressions allow Eqs. (2.7) and (2.8) to be rewritten in the following dimensionless form:

$$\Delta_x(\theta_e + \Lambda_i \theta_i) = -(\theta_i^{\eta_i} + \Lambda_m \Lambda_e \theta_e^{\eta_e}), \quad (2.12)$$

$$\Delta_x(\theta_e - \Lambda_m \Lambda_i \theta_i) = -\Lambda_G(\theta_i^{\eta_i} - \Lambda_e \theta_e^{\eta_e}), \quad (2.13)$$

with the initial conditions

$$\theta_k(0) = 1, \quad \theta'_k(0) = 0. \quad (2.14)$$

The Laplacian Δ_x that appears in Eqs. (2.12) and (2.13) acts on the coordinate x . The dimensionless Λ parameters are defined below.

The ratio between the densities n_{e0} and n_{i0} determines the parameter

$$\Lambda_e = \frac{n_{e0}}{Zn_{i0}} \approx 1, \quad (2.15)$$

which characterizes the deviation from electroneutrality in the center of the star. In the limit of $\Lambda_G = \infty$, Eqs. (2.13) and (2.14) require $\Lambda_e = 1$, which is the LEC at $r = 0$.

Λ_i characterizes the smallness of the ion pressure:

$$\Lambda_i = \frac{\mu_{i0}}{Z\mu_{e0}} \approx \frac{(1 + \eta_i)P_{i0}}{(1 + \eta_e)P_{e0}} \ll 1, \quad (2.16)$$

where P_{k0} is the pressure of the components at the center of the star. The last equality in (2.16) is valid to the same accuracy as the approximation $\Lambda_e \approx 1$. For $\eta_e \neq \eta_i$, the parameters Λ_e and Λ_i are independent.

The parameter Λ_m quantifies the smallness of the electron mass relative to the ion mass:

$$\Lambda_m = \frac{Zm_e}{m_i} = \frac{Zm_e}{Am_u} \approx 5.5 \times 10^{-4} \frac{Z}{A}. \quad (2.17)$$

The main parameter Λ_G is expressed as follows:

$$\Lambda_G = \frac{Z^2 e^2}{Gm_i^2} = \left(\frac{Z}{A}\right)^2 \lambda_G. \quad (2.18)$$

A high value of Λ_G plays the predominant role in specifying the nature of the problem.

Notably, Λ_e and Λ_i depend on the EoS and the stellar structure through n_{k0} and K_k , whereas Λ_m and Λ_G depend only on the chemical composition of the fluid (A and Z).

Equations (2.12)–(2.14) constitute the system of unconstrained HE equations in dimensionless form. This system defines a Cauchy problem with the initial conditions given in Eq. (2.14). In Eq. (2.13), the small parameter $1/\Lambda_G$ appears as a multiplier of the Laplacian; consequently,

Eqs. (2.12) and (2.13) belong to the class of singularly perturbed ODEs.

B. Two-fluid model with equal polytropic indices

The key aspect of the problem is best illustrated by first considering the case of identical polytropic indices $\eta_i = \eta_e \equiv \eta$. A particular solution to the system defined in Eqs. (2.12) and (2.13) can be represented as $\theta_i(x) = \theta_e(x) \equiv \theta(\tilde{x})$, with

$$\tilde{x} = x \sqrt{\frac{1 + \Lambda_m \Lambda_e}{1 + \Lambda_i}}, \quad (2.19)$$

where the function $\theta(\tilde{x})$ is a solution to the Lane-Emden equation

$$\Delta_{\tilde{x}} \theta(\tilde{x}) = -\theta^\eta(\tilde{x}) \quad (2.20)$$

with the standard initial conditions $\theta(0) = 1$ and $\theta'(0) = 0$. In the case of $\eta_i = \eta_e$, the parameters Λ_e and Λ_i are dependent:

$$\Lambda_i = \frac{K_i}{K_e \Lambda_e^{1/\eta} Z^{1+1/\eta}}. \quad (2.21)$$

The self-consistency of Eqs. (2.12) and (2.13) requires $\Lambda_e = \Lambda_e^{\text{reg}}$, where

$$\Lambda_e^{\text{reg}} = \frac{\Lambda_G(1 + \Lambda_i) - (1 - \Lambda_m \Lambda_i)}{\Lambda_G(1 + \Lambda_i) + \Lambda_m(1 - \Lambda_m \Lambda_i)}. \quad (2.22)$$

Equations (2.21) and (2.22) define Λ_i and Λ_e as functions of Λ_G .

In the model considered here, the ion and electron densities appear to be similar, as in the Rosseland model [8]. The parameter Λ_e differs from unity, so the LEC is violated at $O(1/\Lambda_G)$. The solution describes a uniformly polarized star with a positive electric charge.

The concentrations n_{i0} and n_{e0} are considered as input parameters of the model. One would expect the general solution to represent a two-parameter set of functions. The constraint $\Lambda_e = \Lambda_e^{\text{reg}}$ restricts the freedom to a one-parameter set. This seeming paradox is explained in Secs. IV and V. Here, we merely remark that an additional degree of freedom due to variations of Λ_e arises from the singular component of the general solution to Eqs. (2.12) and (2.13).

C. A power-series expansion near the center

A series expansion in powers of x is effective for solving Eqs. (2.12) and (2.13) for arbitrary polytropic indices, and it sheds further light on the origin of the constraint $\Lambda_e = \Lambda_e^{\text{reg}}$. Considering the initial conditions, we search for a solution in the following form:

$$\theta_k = 1 + \sum_{p=1}^{\infty} \beta_{kp} x^{2p}, \quad k = (i, e). \quad (2.23)$$

Terms with odd powers of x are absent because of the even parity of Δ_x . By substituting these expressions into Eqs. (2.12) and (2.13) and equating the coefficients at the same powers of x , we obtain an infinite system of algebraic equations.

The zeroth-order terms yield

$$6(\beta_{e1} + \Lambda_i \beta_{i1}) = -(1 + \Lambda_m \Lambda_e), \quad (2.24)$$

$$6(\beta_{e1} - \Lambda_m \Lambda_i \beta_{i1}) = -\Lambda_G(1 - \Lambda_e). \quad (2.25)$$

The left-hand side of Eq. (2.25) is of $O(1)$, so we require $\Lambda_e = 1 - \alpha_e/\Lambda_G$ with $\alpha_e = O(1)$. β_{e1} and β_{i1} can be found from Eqs. (2.24) and (2.25), whereas α_e remains a free parameter.

The $\sim x^2$ terms yield the equations

$$\begin{aligned} 20(\beta_{e2} + \Lambda_i \beta_{i2}) &= -(\beta_{i1} \eta_i + \Lambda_m \Lambda_e \beta_{e1} \eta_e), \\ 20(\beta_{e2} - \Lambda_m \Lambda_i \beta_{i2}) &= -\Lambda_G(\beta_{i1} \eta_i - \Lambda_e \beta_{e1} \eta_e). \end{aligned}$$

Again the boundedness of the expansion coefficients in last of the above equations yields

$$\beta_{i1} \eta_{i1} = \beta_{e1} \eta_{e1} + \frac{\gamma_1}{\Lambda_G}, \quad (2.26)$$

where $\gamma_1 = O(1)$ is an unknown parameter. This equality, with the use of Eqs. (2.24) and (2.25), allows one to determine the lowest-order parameters, including α_e , to an accuracy of $O(1/\Lambda_G)$:

$$\begin{aligned} \frac{\beta_{e1}}{\eta_i} = \frac{\beta_{i1}}{\eta_e} &= -\frac{1 + \Lambda_m}{6(\eta_i + \Lambda_i \eta_e)}, \\ \alpha_e &= (\eta_i - \Lambda_m \Lambda_i \eta_e) \frac{1 + \Lambda_m}{\eta_i + \Lambda_i \eta_e}. \end{aligned}$$

In the limit of $\eta_i = \eta_e$, α_e is consistent with Eq. (2.22). The higher-order terms in $1/\Lambda_G$ are determined, in turn, from the condition of the boundedness of the expansion coefficients at x^4 , etc. The regularity condition at $\Lambda_G = \infty$ thus imposes a constraint on the concentrations of ions and electrons in the center of the star. We are left with a set of one-parameter solutions.

The results of this subsection can be generalized to GRT [28].

D. A power-series expansion in G

The occurrence of a large parameter Λ_G in Eqs. (2.12) and (2.13) makes it possible to search for solutions throughout the entire x range using a power-series expansion:

$$\theta_k = \theta_{k0} + \theta_{k1} \Lambda_G^{-1} + O(\Lambda_G^{-2}), \quad (2.27)$$

$$\Lambda_e = \Lambda_{e0} + \Lambda_{e1} \Lambda_G^{-1} + O(\Lambda_G^{-2}). \quad (2.28)$$

To a zeroth-order approximation, the regularity of the right-hand side of Eq. (2.13) yields $\theta_{i0}^{\eta_i}(x) = \Lambda_{e0} \theta_{e0}^{\eta_e}(x)$. The initial conditions $\theta_{e0}(0) = \theta_{i0}(0) = 1$ imply that

$$\theta_{i0}^{\eta_i}(x) \equiv \theta_{e0}^{\eta_e}(x), \quad \Lambda_{e0} = 1. \quad (2.29)$$

Equation (2.12) yields

$$\begin{aligned} \Delta_x(\theta_{e0} + \Lambda_i \theta_{i0}) &= -(1 + \Lambda_m) \theta_{e0}^{\eta_e} \\ &= -(1 + \Lambda_m) \theta_{i0}^{\eta_i}, \end{aligned} \quad (2.30)$$

which is the generalized polytropic equation. We solve it for the initial conditions (2.14) in the interval from $x = 0$ up to the stellar surface at $x = x_b$, where $\theta_{e0}(x_b) = \theta_{i0}(x_b) = 0$.

The first-order terms in Eqs. (2.12) and (2.13) yield

$$\Delta_x(\theta_{e1} + \Lambda_i \theta_{i1}) = -\theta_{e0}^{\eta_e} \left(\eta_i \frac{\theta_{i1}}{\theta_{i0}} + \Lambda_m \eta_e \frac{\theta_{e1}}{\theta_{e0}} + \Lambda_m \Lambda_{e1} \right), \quad (2.31)$$

$$\Delta_x(\theta_{e0} - \Lambda_m \Lambda_i \theta_{i0}) = -\theta_{e0}^{\eta_e} \left(\eta_i \frac{\theta_{i1}}{\theta_{i0}} - \eta_e \frac{\theta_{e1}}{\theta_{e0}} - \Lambda_{e1} \right), \quad (2.32)$$

with the initial conditions $\theta_{k1}(0) = \theta'_{k1}(0) = 0$. Let us consider Eq. (2.32) for $x \rightarrow 0$. Considering the asymptotic behavior of θ_k (Eq. (2.27) for β_{k1} and α_e), we obtain the identity to zeroth order. Considering $\Lambda_{e1} = -\alpha_e$, Eq. (2.32) can be written as follows:

$$\frac{(1 + \Lambda_m) \Lambda_i}{\eta_i + \Lambda_i \eta_e} \Delta_x(\eta_e \theta_{e0} - \eta_i \theta_{i0}) = -\theta_{e0}^{\eta_e} \left(\eta_i \frac{\theta_{i1}}{\theta_{i0}} - \eta_e \frac{\theta_{e1}}{\theta_{e0}} \right).$$

For $\eta_i = \eta_e$, this equation is satisfied by the similar distribution $\theta_{i1} = \theta_{e1}$. Equation (2.31), when combined with Eq. (2.32), can also be written as follows:

$$\Delta_x(\theta_{e1} + \Lambda_i \theta_{i1} + \Lambda_m [\theta_{e0} - \Lambda_m \Lambda_i \theta_{i0}]) = -(1 + \Lambda_m) \theta_{e0}^{\eta_e} \eta_i \frac{\theta_{i1}}{\theta_{i0}}. \quad (2.33)$$

The zeroth- and first-order approximations are defined for all values of x . The region near the surface of the star should be treated separately, because for $\theta_{k0} \rightarrow 0$, the validity of the expansion (2.27) can be violated.

E. Global stellar parameters

The stellar radius (more precisely, the boundary of its baryon component) is defined as $R_s = r_0 x_b$, where r_0 is given by Eq. (2.11) and x_b is fixed by the condition

$\theta(x_b) = 0$ from the solution to Eqs. (2.12) and (2.13). The mass of the star takes the form

$$\begin{aligned} M_s &= \int_0^{R_s} 4\pi r^2 (n_i m_i + n_e m_e) dr \\ &= -4\pi r_0^3 n_{i0} m_i x^2 \frac{d}{dx} (\theta_e + \Lambda_i \theta_i)|_{x=x_b}, \end{aligned} \quad (2.34)$$

where Eq. (2.12) is used to proceed to the second line. The total stellar charge (at the boundary of the baryon component) is equal to

$$\begin{aligned} Q_s &= \int_0^{R_s} 4\pi r^2 (Z n_i - n_e) dr \\ &= -4\pi r_0^3 n_{i0} \frac{Ze}{\Lambda_G} x^2 \frac{d}{dx} (\theta_e - \Lambda_m \Lambda_i \theta_i)|_{x=x_b}, \end{aligned} \quad (2.35)$$

where we have used Eq. (2.13). The stellar charge can be estimated as

$$Q_s \sim \frac{Ze M_s}{A m_u \Lambda_G} \simeq 10^{21} e \frac{A}{Z} \frac{M_s}{M_\odot}, \quad (2.36)$$

where M_\odot is the mass of the Sun.

The total uncompensated electric charge of a star of one solar mass, on the boundary of its baryon component, is on the order of 100 C, in agreement with the estimates of Pikel'ner [21], Bally and Harrison [11] and Neslušan [12]. The maximum charges of neutron stars and white dwarfs are estimated to be 50 C and 500 C, respectively [10]. For comparison, one mole of ^{12}C contains $6N_A$ protons, whose total charge is 6×10^5 C. The total negative charge of the Earth at its surface is estimated to be $(4-5.7) \times 10^5$ C [29]. In absolute value, the solar charge is less than $(0.4-1) \times 10^{18}$ C [30].

III. ELECTROSPHERE

In Sec. II, we have seen that the self-consistent regular solution to the unconstrained HE equations describes a star with a positive charge. To move an electron from the surface of such a star to infinity, it is necessary to perform work against the forces of gravity and Coulomb attraction. In fact, a star can confine more electrons than it should be able to according to the regular solution. This possibility leads to the formation of an electrosphere that is quite independent of the internal structure of the star. We therefore discuss the electrosphere separately.

Under normal conditions, a metal is surrounded by a double charged layer, where an excess of positive charge at the subsurface is created by the atomic lattice and a negative charge outside is created by electrons. The thickness of the layer is equal to several Bohr radii, and the work function is several eV. An electrosphere surrounds the surface of hypothetical strange stars [31,32]. Within the

strange star, the positive charge of the baryon component is almost completely neutralized by electrons, whereas it is only partly neutralized in the subsurface. As a result, an electron shell forms above the surface, with a thickness exceeding hundreds of fm, and the chemical potential of electrons near the surface is several tens of MeV. The electrons also give rise to an electrostatic potential jump across the surface separating different phases of nuclear matter [33].

Earlier, we presented arguments for the existence of stars with a total charge of -0.1 to 150 C per solar mass, which are able to support either an electrosphere or an ionosphere. Here, we restrict ourselves to a discussion of the electrosphere. Section IV also describes the properties of the ionosphere.

A. A polytropic model

The regular solution explicitly yields a positive charge excess at $r = 0$. Furthermore, the balance of the gravitational and electrostatic forces ensures that the charge excess must be positive everywhere from the central region up to the surface. The electron component extends somewhat beyond the ion component and compensates for the stellar charge.

Applying the Laplacian to Eq. (2.2), we obtain

$$\Delta_r \mu_e = 4\pi e^2 n_e \left(1 - \frac{G m_e^2}{e^2} \right). \quad (3.1)$$

At the surface ($r = R_s$), an equilibrium condition holds:

$$-\frac{1}{n_e} \frac{dP_e}{dr} = \frac{e Q_s}{R_s^2} + \frac{G m_e M_s}{R_s^2}, \quad (3.2)$$

where M_s and Q_s are given by Eqs. (2.34) and (2.35).

Proceeding in the standard way, we define a dimensionless polytropic function based on the equation $n_e = n_{a0} \theta_a^{\eta_e}$, where n_{a0} is the electron density at the boundary of the ion component. The thickness of the envelope is assumed to be much smaller than R_s . This assumption will be checked at the end of the calculation. The Laplacian simplifies to $\Delta_r \rightarrow d^2/dr^2$. For convenience, we introduce the dimensionless parameter

$$\lambda_m = \frac{m_e}{m_u}. \quad (3.3)$$

The dimensionless variable y is defined by

$$r = R_s + r_a y, \quad r_a^2 = \frac{\mu_{a0}}{4\pi e^2 (1 - \lambda_m^2 / \lambda_G) n_{a0}}, \quad (3.4)$$

where μ_{a0} is the chemical potential of electrons at the surface. Thus, Eq. (3.1) becomes

$$\frac{d^2\theta_a}{dy^2} = \theta_a^{\eta_e}(y). \quad (3.5)$$

At the surface, $y = 0$ and $\theta_a(0) = 1$. Once the stellar structure equations are solved and the stellar mass, radius, and charge are determined, the initial condition for $\theta_a'(0)$ follows from Eq. (3.2). Considering Eq. (2.36), we write the charge in the form

$$Q_s = \frac{M_s}{m_u} \frac{eq_s}{\lambda_G}, \quad (3.6)$$

where $q_s = O(1)$ can be found by solving the unconstrained HE equations. We remark that $M_\odot/m_u \sim (\lambda_G/\alpha)^{3/2}$, where $\alpha = e^2/(\hbar c) \approx 1/137$ is the fine structure constant, and therefore, $Q_s \sim e\sqrt{\lambda_G}/\alpha^{3/2}$. The second initial condition can be written as

$$\left. \frac{d\theta_a}{dy} \right|_{y=0} = -(q_s + \lambda_m) \sqrt{\frac{P_{\text{un}}}{(1 + \eta_e)P_{a0}(\lambda_G - \lambda_m^2)}}, \quad (3.7)$$

where $P_{a0} = K_e n_{a0}^{1+1/\eta_e}$ is the pressure of the electrons at the boundary and P_{un} is the natural unit of pressure in the star:

$$P_{\text{un}} = \frac{GM_s^2}{4\pi R_s^4}.$$

Integrating Eq. (3.5) yields

$$\frac{d\theta_a}{dy} = -\sqrt{2(W_0 - V(\theta_a))}, \quad (3.8)$$

where

$$W_0 = \frac{\theta_a^2(0)}{2} - \frac{1}{1 + \eta_e},$$

$$V(\theta_a) = -\frac{\theta_a^{1+\eta_e}}{1 + \eta_e},$$

and $\theta_a'(0)$ is given by Eq. (3.7). A second integration yields

$$y = y_a - \frac{\theta_a}{\sqrt{2(W_0 - V(\theta_a))}} \times {}_2F_1\left(\frac{1}{2}, 1; \frac{2 + \eta_e}{1 + \eta_e}; \frac{-V(\theta_a)}{W_0 - V(\theta_a)}\right), \quad (3.9)$$

where ${}_2F_1$ is the Gauss hypergeometric function. Equation (3.9) implicitly specifies $\theta_a = \theta_a(y)$. With the initial condition $\theta_a(0) = 1$, the boundary of the electrosphere can be found from $\theta_a(y_a) = 0$ to give




FIG. 1. Electron shell thickness y_a in the nonrelativistic ($\eta_e = 3/2$) and relativistic ($\eta_e = 3$) cases versus the parameter W_0 .

$$y_a = \frac{1}{|\theta_a'(0)|} {}_2F_1\left(\frac{1}{2}, 1; \frac{2 + \eta_e}{1 + \eta_e}; \frac{2}{(1 + \eta_e)\theta_a^2(0)}\right).$$

Figure 1 shows the behavior of y_a as a function of W_0 . With a decrease in W_0 , the radial thickness of the envelope increases, and for $W_0 \rightarrow 0$, the radial thickness goes to infinity. However, the function y_a is still well defined at zero:

$$y_a(W_0 = 0) = -\frac{\sqrt{2(\eta_e + 1)}}{\eta_e - 1} < 0, \quad (3.10)$$

because of the irregularity of y_a for $W_0 = 0$. In comparison with the nonrelativistic case, the shell of relativistic electrons extends higher.

Equation (3.5) can be interpreted as the equation of motion of a point particle of unit mass in a potential $V(\theta_a)$, where y plays the role of a time variable. Equation (3.8) follows from the conservation of the total energy:

$$W = \frac{\theta_a^2}{2} + V(\theta_a).$$

There are three types of solutions. If the initial momentum is small, the particle moves to the left relative to its initial position $\theta_a(0) = 1$, stops at some point $0 < \theta_a < 1$, and then accelerates in the positive direction and goes to infinity. The total energy of the particle is negative: $W = W_0 < 0$. In the Thomas-Fermi theory of multielectron atoms, solutions of this type are called *compressed*; they terminate at $\theta_a' = 0$ for a finite electron density, for which the global electroneutrality of the atom holds [34]. In our problem, however, such solutions are not physical since the stars are, of course, isolated. The second type of solution corresponds to the case of zero energy: the particle arrives at the point $\theta_a = 0$ and stops there, such that $\theta_a' = 0$. In the Thomas-Fermi theory, such solutions describe neutral

atoms. In our case, they describe stars with a maximum negative total charge, i.e., $Q_{\text{tot}}^{\text{max}} < 0$. The third type of solution is obtained for $W_0 > 0$. The particle arrives at the point $\theta_a = 0$ with a finite velocity of $\theta'_a < 0$, and then the solution terminates. In the Thomas-Fermi theory, solutions of this type correspond to ionized atoms with a total positive charge. In our case, such a solution describes an electrosphere of a finite thickness y_a , and the total charge of the star can be negative, zero, or positive.

The existence of solutions, therefore, depends on the derivative of the chemical potential of the electrons at the boundary of the ion component. The absolute value of the derivative is constrained from below by the condition $W_0 \geq 0$, which imposes an upper limit on the number of electrons in the shell.

At the boundary of the ion component, the total positive charge of the star is equal to Q_s , and the charge of the electrosphere Q_e either partially, completely, or excessively compensates for Q_s . Considering Eqs. (3.5) and (3.8), the charge of the electrosphere can be written in the form

$$Q_e = -4\pi R_s^2 \int_0^{y_a} e n_e r_a dy \\ = 4\pi R_s^2 e n_{a0} r_a \left(\theta'_a(0) + \sqrt{\theta_a^2(0) - \frac{2}{1 + \eta_e}} \right). \quad (3.11)$$

Let κ be a parameter characterizing the screening, such that $Q_e = -\kappa Q_s$, where $0 \leq \kappa \leq \kappa_{\text{max}}$. In dimensionless units [see Eq. (3.6)], the charge of the electrosphere is equal to $q_e = -\kappa q_s$. In particular, for $\kappa = 1$, the screening is complete, and the total stellar charge vanishes. From Eq. (3.11), one can find

$$P_{a0} = \kappa q_s \frac{P_{\text{un}}}{\lambda_G} \left[q_s + \lambda_m - \frac{\kappa q_s}{2} (1 - \lambda_m^2 / \lambda_G) \right],$$

implying that the pressure in the envelope is λ_G times smaller than P_{un} . It is now convenient to rewrite Eq. (3.7) in the form

$$\left. \frac{d\theta_a}{dy} \right|_{y=0} = - \frac{q_s + \lambda_m}{\sqrt{\kappa q_s (1 + \eta_e) (1 - \lambda_m^2 / \lambda_G) (q_s + \lambda_m - \kappa q_s / 2 (1 - \lambda_m^2 / \lambda_G))}}. \quad (3.12)$$

The maximum negative charge of the envelope corresponds to $W_0 = 0$. From Eq. (3.12), one can find the critical value of the compensation parameter:

$$\kappa_{\text{max}} = \frac{1 + \lambda_m / q_s}{1 - \lambda_m^2 / \lambda_G} > 1. \quad (3.13)$$

Solutions with a total negative charge, $Q_{\text{tot}} = Q_s + Q_e = Q_s(1 - \kappa) < 0$, are admissible since an electrically neutral star has a gravitational field capable of binding electrons under the condition that the energy of their Coulomb repulsion does not exceed the energy of their gravitational attraction to the star. An estimate of $Q_s \sim 100$ C was obtained above. The maximum negative charge $Q_{\text{tot}}^{\text{max}}$ is proportional to the parameter $\lambda_m \ll 1$; thus, $Q_{\text{tot}}^{\text{max}} \sim -0.05$ C per solar mass.

The gravitational collapse of a charged single-component spherically symmetric fluid is discussed in Refs. [35,36].

The maximum charge of a neutron star is determined in Ref. [37] on the basis of the requirement of gravitational stability. We showed that the maximum charge can instead be determined from the requirement for the existence of a solution of finite extent for the electro- or ionosphere of the star. The latter requirement yields a limit that is $\sqrt{\lambda_G}$ times stronger than that obtained from the requirement of gravitational stability.

The positive charge excess of strange stars at the quark surface is found in Refs. [38,39] to be about 10^{20} C.

The electrosphere compensates this charge, although the detailed mechanism has not yet been studied. The particular values of the bag constant B used in Refs. [38,39] rule out the existence of massive neutron stars metastable against conversion to strange stars [40,41].

B. Electrically neutral stars

For $W_0 = 0$, Eqs. (3.9)–(3.10) yield

$$\theta_a^{\text{max}}(y) = \left(1 + \frac{y}{y_0} \right)^{-\frac{2}{\eta_e - 1}}, \quad (3.14)$$

where $y_0 = -y_a(W_0 = 0) > 0$. This solution extends to infinity, which means that an electron can be moved from the center of the star to infinity without doing work. In the case of a relativistic electron gas, $\eta_e = 3$, the solution decays rather slowly: $\theta_a^{\text{max}} \sim 1/y$. At low densities, the electrons become nonrelativistic, with $\eta_e = 3/2$, and the solution approaches $\theta_a^{\text{max}} \sim 1/y^4$. It is not difficult to directly verify that $\theta_a^{\text{max}}(y)$ is a solution to Eq. (3.5) with the proper boundary conditions.

In view of the approximate equality $\kappa_{\text{max}} \approx 1$, the electrosphere of a neutral star with $\kappa = 1$ is described by a solution close to $\theta_a^{\text{max}}(y)$. We return to Eq. (3.12) and set $\kappa = 1$. From this equation, by neglecting the small terms $\sim 1/\lambda_G$ and $\sim \lambda_m$, the derivative of $\theta_a(y)$ at $y = 0$ can be found to be

$$\left. \frac{d\theta}{dy} \right|_{y=0} \approx -\sqrt{\frac{2}{1+\eta_e}}(1+\xi), \quad (3.15)$$

where

$$\xi = \frac{1}{2} \left(\frac{\lambda_m}{q_s} \right)^2 \ll 1. \quad (3.16)$$

We are looking for a solution to Eq. (3.5) in the form $\theta_a = \theta_a^{\max} + \sigma$, where σ is a small correction. The initial conditions are of the form $\sigma(0) = 0$ and $\sigma'(0) = -\xi\sqrt{2/(1+\eta_e)}$. To the lowest order, σ satisfies the equation

$$\frac{d^2\sigma}{dy^2} = \frac{\eta_e\sigma}{(1+\frac{y}{y_0})^2}. \quad (3.17)$$

The solution to this equation is given by

$$\sigma(y) = \frac{2\xi}{3\eta_e + 1} \left[\left(1 + \frac{y}{y_0}\right)^{\frac{\eta_e+1}{\eta_e-1}} - \left(1 + \frac{y}{y_0}\right)^{\frac{2\eta_e}{\eta_e-1}} \right].$$

The dimensionless chemical potential of the electrosphere of a neutral star as a function of the distance to the surface is shown in Fig. 2 for $\eta_e = 3/2$ and in Fig. 3 for $\eta_e = 3$. The equilibrium conditions (2.1) and (2.2) are valid for nonrelativistic particles. For completeness, we also consider the relativistic case $\eta_e = 3$, which can be interesting for qualitative estimates. The total stellar charge is assumed to be zero ($\kappa = 1$), whereas the parameter $\xi = 1.5 \times 10^{-7}$ corresponds to $q_s = 1$. The dashed curves show the asymptotic solutions corresponding to Eq. (3.14), the




FIG. 2. Chemical potential θ_a of nonrelativistic electrons as a function of the distance from the surface of a neutral star. The dashed curve shows the limiting solution θ_a^{\max} to Eq. (3.14), the solid curve labeled E shows the exact solution to Eq. (3.9), and the solid curve labeled A shows the approximate solution $\theta_a^{\max} + \sigma$.




FIG. 3. Chemical potential θ_a of relativistic electrons as a function of the distance from the surface. All parameters and notation are the same as in Fig. 2.

solid curves labeled E show the exact solutions corresponding to Eq. (3.9), and the solid curves labeled A show the approximations $\theta_a^{\max} + \sigma$. The approximations reproduce the exact formula in Eq. (3.9) with reasonable accuracy. The thickness of the electrosphere can be estimated to be

$$\frac{y_a}{y_0} \approx \left(\frac{3\eta_e + 1}{2\xi} \right)^{\frac{\eta_e-1}{2(\eta_e+1)}} - 1. \quad (3.18)$$

An order-of-magnitude estimate of the thickness of the electrosphere $y_a r_a$ can be found from Eq. (3.4).

For nonrelativistic electrons with $\eta_e = 3/2$, the pressure and density are related by $P_e \sim \hbar^2 n_e^{5/3} / m_e$. Considering that $P_{a0} \sim P_{un} / \lambda_G$, we obtain

$$\begin{aligned} \frac{r_a}{R_s} &\sim \left[\left(\frac{a_B}{R_s} \right)^3 \frac{m_u}{M_s} \lambda_G \right]^{1/5} \\ &\approx 2.2 \times 10^{-16} \left(\frac{R_\odot}{R_s} \right)^{3/5} \left(\frac{M_\odot}{M_s} \right)^{1/5}, \end{aligned} \quad (3.19)$$

where $a_B = \hbar^2 / (e^2 m_e)$ is the Bohr radius. In comparison with the stellar radius, the thickness of the electrosphere is quite small, although it is not small on an atomic scale: $y_a r_a \sim 6 \times 10^4 a_B$ for an electrically neutral star of mass $M_s = M_\odot$ and radius $R_s = R_\odot$.

In the case of $\eta_e = 3$, the substitution of the relativistic relationship $P_e \sim \hbar c n_e^{4/3}$ into Eq. (3.4) yields

$$\frac{r_a}{R_s} \sim \sqrt{\frac{\lambda_G}{\alpha^{3/2} M_s}} \frac{m_u}{M_s} \approx 1.3 \times 10^{-9} \left(\frac{M_\odot}{M_s} \right)^{1/2}. \quad (3.20)$$

The thickness of the electrosphere of an electrically neutral neutron star with a radius of $R_s = 10$ km and a mass of $1.4 M_\odot$ can be estimated to be $y_a r_a \sim 1$ mm.

IV. EXACTLY SOLVABLE MODEL

As seen from the previous discussion, the regular solution describes a polarized star with a nonzero positive charge as measured at the ion surface. The properties of the electrosphere are uniquely determined by the internal structure of the star. The electrosphere ensures that the total stellar charge is generally not equal to zero. This feature is particularly evident in the model with equal polytropic indices considered in Sec. II.B. An electrosphere occurs provided that $\theta_e(x_b) \neq 0$, whereas for identical polytropic indices, the distributions vanish at the same point, $x = x_b$.

Earlier, we presented simple physical arguments in favor of the existence of stars with electric charges in the range of -0.1 to 150 C per solar mass. A regular solution corresponds to a single representative of this class. The regular solutions described in Refs. [8–15,21] and in Sec. II above do not exhaust all physical possibilities, nor are they general solutions to the unconstrained HE equations.

Fourth-order systems of ODEs, similar to Eqs. (2.12) and (2.13), depend on four initial conditions. In our case, two of the initial conditions constrain the derivatives of the chemical potentials [cf. Eq. (2.14)]. These conditions ensure smooth behavior of the solutions at $r = 0$ and cannot be modified. The remaining two initial conditions fix μ_i and μ_e and thereby define a two-parameter set of stellar configurations. As shown in Sec. II, the requirement of regularity imposes a constraint on the value of Λ_e , such that only a one-parameter set of stellar configurations survives.

Some of the solutions to Eqs. (2.12) and (2.13) appear to be lost with the imposition of a requirement for regularity in the neighborhood of $\Lambda_G = \infty$.

A. General properties of solutions

The analytic properties of solutions to ODE systems are determined by the Poincaré theorem on analyticity (see, e.g., [23]), which asserts, roughly speaking, that solutions to ODE systems, when they exist, are analytic functions of the initial coordinates and parameters in the region of analyticity of the ODEs. The origin of nonregular solutions can also be understood on the basis of Dyson's argument, which provides an effective qualitative criterion for the nonanalyticity of observables in terms of the system parameters [24]. From these perspectives, we discuss the analytic properties in the parameter Λ_G of the solutions to Eqs. (2.12) and (2.13).

1. Poincaré theorem on analyticity

Using the independent variables $\pi_k = \theta'_k$, Eqs. (2.12) and (2.13) can be reduced to a system of four first-order ODEs. In terms of $\Phi = (\pi_e, \theta_e, \pi_i, \theta_i)$, these equations take the form $\Phi' = \mathfrak{F}(x, \Phi, \Lambda)$, where \mathfrak{F} is a vector function

of x , Φ , and $\Lambda = (\Lambda_e, \Lambda_i, \Lambda_m, \Lambda_G)$. The initial conditions are $\Phi(x=0) = (0, 1, 0, 1)$.

The components of \mathfrak{F} , as can be easily seen, are analytic functions of Φ and Λ . The domain of analyticity of \mathfrak{F} is a Cartesian product of the complex planes of Φ and Λ without $\Phi = \infty$ and $\Lambda = \infty$. Given that the polytropic indices are not integers, $\theta_k = 0$ and $\theta_k = \infty$ are branch points of \mathfrak{F} .

According to the Poincaré theorem, the solutions, when they exist, depend analytically on $\Phi(0)$ and Λ in the domain of analyticity of $\mathfrak{F}(x, \Phi, \Lambda)$. From Eq. (2.14), we observe that the dimensionless chemical potentials at $x = 0$ are frozen, so that the initial values $\Phi(0)$ are not of interest. The essential information is contained in Λ_e and the dimensional quantities r_0 and n_{i0} . The analyticity theorem can be interpreted to mean that throughout the entire domain of their existence, solutions to Eqs. (2.12) and (2.13) are analytic in Λ for $\Lambda \neq \infty$. At infinity, the solutions are either regular or irregular.

The dependence on Λ_G is of particular interest because $\Lambda_G = \infty$ corresponds to the LEC. It would be natural to use the LEC solution as an initial approximation of the general solution to Eqs. (2.12) and (2.13) and to account for deviations from local electroneutrality by means of perturbation theory in $1/\Lambda_G \ll 1$. The analyticity theorem, however, asserts that the general solution is typically not regular at $\Lambda_G = \infty$. If this is so, the perturbation series in $1/\Lambda_G$ diverges because the radius of convergence, which is defined by the closest singularity on the complex Λ_G plane, is equal to zero.

The convergence of the perturbation series for solutions that are regular at $\Lambda_G = \infty$ can be determined by, e.g., the singularities of Λ_e and Λ_i as functions of Λ_G , as in the model of Sec. II.B. According to Eqs. (2.21) and (2.22), Λ_e , Λ_i , and the corresponding system of ODEs are not analytic for $|\Lambda_G| \sim 1$. If there are no singularities close to infinity, the point $\Lambda_G = \infty$ lies within the circle of convergence of the perturbation series centered at the physical value (2.18). In the model of Sec. II.B, one could obtain then the LEC solution in the form of a convergent series starting from the regular solution. The reciprocal statement is also true: starting from the LEC solution, one could obtain the regular solution in the form of a convergent series.

2. Dyson's argument

A physical argument leading to similar conclusions has been presented by Dyson [24]. In an extended context, the argument concerns the relationship between singular points in the parameter space and qualitative changes in the behavior of the system. As applied to quantum electrodynamics (QED), Dyson's argument leads to the well-known conclusions that $\alpha = 0$ is a singular point of physical observables, QED series in powers of the fine structure constant have radii of convergence equal to zero and therefore are asymptotic.

In our case, $\Lambda_G = \infty$ corresponds to $G = 0$. The physical picture for $G < 0$ undergoes obvious qualitative changes; namely, the attraction between particles is replaced by repulsion, and as a result, bound states (stars) no longer exist.

We also remark that the many-body Hamilton function for $G < 0$ is nonnegative, whereas for positive G it is not bounded from below.

It can be observed that the characteristic length given in Eq. (2.11) has branch points at $G = 0$ and ∞ . The stellar mass (2.34) and charge (2.35) inherit these singularities, which are further superimposed with the singularities of θ_k .

Dyson's argument indicates that the general solution to Eqs. (2.12) and (2.13) should be sought in the class of functions that depend on Λ_G in an irregular manner.

B. Two-fluid model with unit polytropic indices

The above statements can be illustrated and precisely formulated using the model of Sec. II. B with unit polytropic indices, which admits an explicit analytic solution. The adiabatic index of the model, $\gamma = 1 + 1/\eta = 2$, is

sufficiently close to that of an ideal gas model with $\gamma = 5/3$. In terms of the functions $\varphi_k \equiv x\theta_k(x)$, Eqs. (2.12) and (2.13) are reduced to a linear ODE system:

$$\varphi_e'' + \Lambda_i \varphi_i'' = -(\varphi_i + \Lambda_m \Lambda_e \varphi_e), \quad (4.1)$$

$$\varphi_e'' - \Lambda_m \Lambda_i \varphi_i'' = -\Lambda_G(\varphi_i - \Lambda_e \varphi_e), \quad (4.2)$$

where the prime denotes differentiation in x . We look for solutions of the form $\varphi_k = \alpha_k e^{\beta x}$. Equations (4.1) and (4.2) yield

$$(\alpha_e + \Lambda_i \alpha_i) \beta^2 = -(\alpha_i + \Lambda_m \Lambda_e \alpha_e), \quad (4.3)$$

$$(\alpha_e - \Lambda_m \Lambda_i \alpha_i) \beta^2 = -\Lambda_G(\alpha_i - \Lambda_e \alpha_e). \quad (4.4)$$

The algebraic system formed by Eqs. (4.3) and (4.4) has nontrivial solutions provided that the determinant is equal to zero. This condition gives rise to a quadratic equation in β^2 , whose solutions are

$$\begin{aligned} \beta_{\pm}^2 = & \frac{1}{2\Lambda_i(1+\Lambda_m)} \{ \Lambda_G(1+\Lambda_i\Lambda_e) - (1+\Lambda_m^2\Lambda_i\Lambda_e) \\ & \pm \sqrt{\Lambda_G^2(1+\Lambda_i\Lambda_e)^2 - 2\Lambda_G[(1-\Lambda_m\Lambda_i\Lambda_e)^2 - \Lambda_i\Lambda_e(1+\Lambda_m)^2] + (1+\Lambda_m^2\Lambda_i\Lambda_e)^2} \}. \end{aligned} \quad (4.5)$$

To the leading order in $1/\Lambda_G$, the values of β are given by

$$\beta_+^2 = \Lambda_G \frac{1 + \Lambda_i \Lambda_e}{\Lambda_i(1 + \Lambda_m)} + O(1), \quad (4.6)$$

$$\beta_-^2 = -\frac{\Lambda_e(1 + \Lambda_m)}{1 + \Lambda_i \Lambda_e} + O(\Lambda_G^{-1}). \quad (4.7)$$

The values $\pm\sqrt{\beta_+^2}$ are real and large, whereas the values $\pm\sqrt{\beta_-^2}$ are imaginary. We adopt the notation $\beta_1 = |\beta_+| > 0$ and $\beta_2 = |\beta_-| > 0$. The general solution has the form

$$\begin{aligned} \varphi_k(x) = & \tilde{\alpha}_{k1} \exp(-\beta_1 x) + \tilde{\alpha}_{k2} \exp(\beta_1 x) \\ & + \tilde{\alpha}_{k3} \sin(\beta_2 x) + \tilde{\alpha}_{k4} \cos(\beta_2 x). \end{aligned} \quad (4.8)$$

The initial conditions for θ_k lead to $\varphi_k(0) = 0$, $\varphi_k'(0) = 1$, and $\varphi_k''(0) = 0$. The general solution becomes

$$\varphi_k(x) = \alpha_{k1} \sinh(\beta_1 x) + \alpha_{k2} \sin(\beta_2 x), \quad (4.9)$$

with the constraint

$$\alpha_{k1}\beta_1 + \alpha_{k2}\beta_2 = 1. \quad (4.10)$$

By substituting $\beta^2 = \beta_1^2$ and $-\beta_2^2$ into Eq. (4.3) for α_{k1} and α_{k2} , respectively, and using Eq. (4.10), we can uniquely determine α_{k1} and α_{k2} and find the solution.

1. Closer inspection of analyticity in G

The analytic properties of $\varphi_k(x)$ in the complex Λ_G -plane can be clarified by transforming the system defined by Eqs. (4.1) and (4.2) into normal form with the independent variables φ_k and $\pi_k = \varphi_k'$. In terms of $\Phi \equiv (\pi_i, \varphi_i, \pi_e, \varphi_e)$, Eqs. (4.1) and (4.2) take the form

$$\Phi' = \mathfrak{A}\Phi, \quad (4.11)$$

where \mathfrak{A} is a real 4×4 matrix that depends linearly on the parameters $\Lambda = (\Lambda_e, \Lambda_i, \Lambda_m, \Lambda_G)$. The solution takes the form

$$\Phi(x) = \exp(\mathfrak{A}x)\Phi(0). \quad (4.12)$$

The series expansion of the exponential matrix converges, so the evolution operator $\mathfrak{D} = \exp(\mathfrak{A}x)$ is well defined. The matrix elements of \mathfrak{D} are analytic functions in Λ for $\Lambda \neq \infty$. A simple pole of \mathfrak{A} at $\Lambda = \infty$ is transformed into an essential fixed singularity of \mathfrak{D} , which does not depend on the initial values. The functions $\Phi(x)$ are, therefore,

analytic functions of Λ in a Cartesian product of the complex planes of Λ_e , Λ_i , Λ_m , and Λ_G . Equation (4.12) also shows that $\Phi(x)$ are analytic on the initial values $\Phi(0)$, in agreement with the Poincaré theorem on analyticity.

Let \mathbf{a}_n denote the eigenvalues of \mathfrak{A} that coincide with $\pm\beta_{\pm} \in \mathbb{C}^1$, and let \mathbf{r}_n and \mathbf{l}_n be their corresponding right and left eigenvectors satisfying $\mathfrak{A}\mathbf{r}_n = \mathbf{a}_n\mathbf{r}_n$ and $\mathbf{l}_n^T\mathfrak{A} = \mathbf{l}_n^T\mathbf{a}_n$. The eigenvectors are normalized according to $\mathbf{l}_n^T\mathbf{r}_m = \delta_{nm}$, where $n = 1, 2, 3, 4$. The evolution operator admits the representation

$$\mathfrak{D} = \sum_n \mathbf{r}_n \mathbf{l}_n^T \exp(\mathbf{a}_n x). \quad (4.13)$$

The eigenvalues \mathbf{a}_n are roots of the characteristic equation, so they have additional singularities compared with \mathfrak{A} . The analyticity of \mathfrak{D} implies that additional singularities cancel out with singularities of the projection operators $\mathbf{r}_n \mathbf{l}_n^T$.

A more explicit proof of this claim uses the Frobenius representation

$$\mathbf{r}_n \mathbf{l}_n^T = \prod_{m \neq n} \frac{\mathfrak{A} - \mathbf{a}_m}{\mathbf{a}_n - \mathbf{a}_m}. \quad (4.14)$$

We first note that the radicand in Eq. (4.5) vanishes for a complex-conjugate pair of Λ_G , where β_{\pm} have square-root branch points. The point $\Lambda_G = \infty$ is a simple pole of β_+^2 , a square-root branch point of β_+ , and an essentially singular point of $\varphi_k(x)$. $\Lambda_G = 0$ is a simple zero of β_+^2 and a square-root branch point of β_+ . The eigenvalues \mathbf{a}_n ($\pm\beta_{\pm}$) accordingly have four square-root branch points in the complex Λ_G plane. By passing around one of these points, we obtain simply a permutation of \mathbf{a}_n . The representation of \mathfrak{D} using Eq. (4.14) is explicitly invariant under these permutations, so the matrix elements of \mathfrak{D} do not inherit additional singularities of \mathbf{a}_n .

Consequently, the eigenvalues $\pm\beta_{\pm}$ and the projection operators of \mathfrak{A} have four singular points, whereas $\varphi_k(x)$ has a unique singular point $\Lambda_G = \infty$.

Given that $\varphi_k(x)$ is analytic in $\Lambda_G \in \mathbb{C}^1 \setminus \infty$, $\varphi_k(x)$ if not a constant must be singular at $\Lambda_G = \infty$. The existence of solutions independent of Λ_G is excluded by Eq. (4.2). Regular solutions at $\Lambda_G = \infty$ can be constructed by replacing the parameters, e.g., Λ_e and Λ_i , with functions of Λ_G , as in the model of Sec. II B. Solutions inherit the singularities of these functions, which makes it possible to construct a regular solution. The dependence of the parameters on Λ_G is a necessary condition for the existence of a regular solution in the neighborhood of $\Lambda_G = \infty$.

A qualitative analysis of Eqs. (4.1) and (4.2), developed above, reveals the mathematical reason why the parameters Λ_e and Λ_i are not independent in the regular solution. The analyticity in Λ_G follows from the representation given in Eq. (4.12). The requirement of regularity at $\Lambda_G = \infty$ would imply regularity in the extended complex Λ_G plane. In such

a case, $\varphi_k(x)$ must be a constant. By introducing a dependence of the parameters on Λ_G , particular, non-trivial, regular solutions at $\Lambda_G = \infty$ can be constructed, which can further be used to approximate the general solution.

2. Electro- and ionospheres

The first term in Eq. (4.9) depends exponentially on $\beta_1 \sim \sqrt{\Lambda_G}$. For $x \neq 0$, $\sinh(\beta_1 x)$ has no zeros. For the chemical potential to vanish at the boundary, the solution must contain the component $\sim \sin(\beta_2 x)$. The first term in Eq. (4.9) is maximal on the boundary of the star, where it should not exceed α_2 . The ratio α_1/α_2 is therefore exponentially small.

The solutions are recovered with exponential accuracy already for $\alpha_2\beta_2 = 1$. Substituting this expression into Eqs. (4.3) and (4.4), we obtain

$$(1 + \Lambda_i)\beta_2^2 = 1 + \Lambda_m\Lambda_e, \quad (4.15)$$

$$(1 - \Lambda_m\Lambda_i)\beta_2^2 = \Lambda_G(1 - \Lambda_e). \quad (4.16)$$

The self-consistency of the system leads to $\Lambda_e = \Lambda_e^{\text{reg}}$, where Λ_e^{reg} is given by Eq. (2.22). The corresponding β parameters (with an additional index 0) become

$$\beta_{10}^2 = \frac{\Lambda_G(1 + \Lambda_i) - (1 - \Lambda_m\Lambda_i)}{\Lambda_i(1 + \Lambda_m)}, \quad (4.17)$$

$$\beta_{20}^2 = \frac{\Lambda_G(1 + \Lambda_m)}{\Lambda_G(1 + \Lambda_i) + \Lambda_m(1 - \Lambda_m\Lambda_i)}. \quad (4.18)$$

Equations (2.22), (4.17) and (4.18) are exact to all orders in $1/\Lambda_G$. Λ_i is a function of Λ_G , according to Eq. (2.21). In the approximation $\alpha_1 = 0$, with exponential accuracy, the function $\theta(x)$ is given by

$$\theta(x) = \frac{\varphi(x)}{x} \approx \frac{\sin(\beta_{20}x)}{\beta_{20}x}, \quad (4.19)$$

in agreement with the analysis of Sec. II B for unit polytropic indices. The function $\theta(x)$ is regular for $\Lambda_G = \infty$ but inherits the singularities of β_{20} .

A variation in the electron density at $r = 0$ leads to a variation $\Delta\Lambda_e = \Lambda_e - \Lambda_e^{\text{reg}} \neq 0$, corresponding variations of Λ_i and β_{\pm} and, ultimately, to $\alpha_{k1} \neq 0$. We return to the general solution (4.9) and write the function θ_k in the form

$$\theta_k(x) = \alpha_{k1} \frac{\sinh(\beta_1 x)}{x} + \alpha_{k2} \frac{\sin(\beta_2 x)}{x}, \quad (4.20)$$

where, with account of Eq. (4.10),

$$\alpha_{k2} = \frac{1 - \alpha_{k1}\beta_1}{\beta_2}. \quad (4.21)$$

The regular solution (4.19) vanishes for $x_0 = \pi/\beta_{20}$. For $\alpha_{k1} \neq 0$, one of the functions $\theta_k(x)$ vanishes already at $x_b = x_0 + \Delta x$, whereas the second function is still finite. We write this condition in the form

$$\theta_i(x_b) = 0, \quad (4.22)$$

$$\theta_e(x_b) = \theta_{eb}, \quad (4.23)$$

where θ_{eb} is the value of the electron function at the boundary. We assume first that an electrosphere is formed.

As follows from Sec. III, the electron pressure at the boundary of the star is $P_{eb} \sim P_{un}/\Lambda_G$, where P_{un} is the characteristic pressure scale. The relation $P_e \sim n_e^{1+1/\eta_e} \sim \theta_e^{1+\eta_e}$ implies that $\theta_{eb} \sim \Lambda_G^{-1/(1+\eta_e)} \ll 1$. For the case under consideration, $\eta_e = 1$, and accordingly, $\theta_{eb} \sim 1/\sqrt{\Lambda_G}$. The typical scale of the electrosphere is $r_a/r_0 \sim \theta_{eb} \sim \Lambda_G^{-1/(1+\eta_e)}$, in agreement with the estimates given in Eqs. (3.19) and (3.20). Using Eq. (3.2) and the inequality $W_0 \geq 0$, we obtain

$$\theta_{eb} \leq \lambda_G^{-1/(1+\eta_e)} \left[\frac{(1+\eta_e)(q_s + \lambda_m)^2}{2\Lambda_e(1+\Lambda_m)(1-\lambda_m^2/\lambda_G)} \times \left(\frac{M_s}{4\pi r_0^3 n_{i0} m_i} \right)^2 \left(\frac{r_0}{R_s} \right)^4 \right]^{1/(1+\eta_e)}. \quad (4.24)$$

Equations (4.22) and (4.23) allow to express the unknown coefficients α_{k1} in terms of θ_{eb} . Given the relation in Eq. (4.21), the functions $\theta_k(x)$ are completely defined, but the parameters Δx and θ_{eb} on which these functions depend must be found. We first substitute α_{k1} into the first of Eqs. (4.3) and (4.4), setting $\beta^2 = \beta_1^2$, and then substitute α_{k2} , setting $\beta^2 = -\beta_2^2$. The two resulting equations are linear in θ_{eb} . By excluding θ_{eb} , we find

$$\frac{(1+\Lambda_i)\beta_2^2 - 1 - \Lambda_m\Lambda_e \sinh(\beta_1 x_b)}{\beta_2^2 - \Lambda_m\Lambda_e} \frac{\beta_1 x_b}{\beta_1 x_b} = \frac{(1+\Lambda_i)\beta_1^2 + 1 + \Lambda_m\Lambda_e \sin(\beta_2 x_b)}{\beta_1^2 + \Lambda_m\Lambda_e} \frac{\beta_2 x_b}{\beta_2 x_b}. \quad (4.25)$$

This equation defines x_b as a function of the parameters Λ_e , Λ_i , Λ_m , and Λ_G . Equation (4.4) is not used because for $\beta^2 = \beta_2^2$, it is not independent. Both parts of Eq. (4.25) define $\theta_{eb} \geq 0$; therefore, a solution can exist for $\beta_2^2 \geq \beta_{20}^2$ and $\Delta x \leq 0$. As noted above, $\theta_{eb} \sim 1/\beta_1 \sim 1/\sqrt{\Lambda_G}$, and so $\Delta x \sim r_a/r_0 \sim 1/\sqrt{\Lambda_G}$. Considering $\Delta x \sim 1/\beta_1 \ll 1$, the right-hand side of Eq. (4.25) can be expanded into a series in the neighborhood of $x_b = x_0$; as a result, we find

$$\Delta x = -\frac{1}{\beta_1} \mathcal{W}\left(\frac{\Delta\Lambda_e \exp(\pi\beta_1/\beta_2)}{2(1+\Lambda_i)}\right). \quad (4.26)$$

The displacement Δx is expressed in terms of the Lambert \mathcal{W} -function, which gives a solution to the equation $\mathcal{W}(x) \exp(\mathcal{W}(x)) = x$. Since the argument of the function is of order unity, the correction to Λ_e is exponentially small:

$$\Delta\Lambda_e \sim \exp(-O(\sqrt{\Lambda_G})).$$

For θ_{eb} , we obtain

$$\theta_{eb} = -\frac{\Delta x}{x_0} (1 + \Lambda_i). \quad (4.27)$$

In Eqs. (4.26) and (4.27), the $\sim \exp(-\pi\beta_1/\beta_2)$ terms and higher terms in $1/\sqrt{\Lambda_G}$ are omitted. At this level of accuracy, we can replace β_1 and β_2 with the expressions in Eqs. (4.17) and (4.18) everywhere, including exponents. The positive definiteness of θ_{eb} implies that $\Delta x < 0$ and $\Delta\Lambda_e > 0$.

For asymptotically large values of the argument of $\mathcal{W}(x)$, the expansion $\mathcal{W}(x) = \ln(x) - \ln(\ln(x)) + \dots$ is valid. A correction $\Delta\Lambda_e \sim 1/\sqrt{\Lambda_G}$ could lead to the estimate $\Delta x \sim \theta_{eb} \sim 1$, which contradicts the upper limit of Eq. (4.24). The set of admissible solutions, therefore, consists only of solutions with exponentially small deviations $\Delta\Lambda_e$.

Equations (4.1) and (4.2) also permit solutions corresponding to an ionosphere. Instead of Eqs. (4.22) and (4.23), we require

$$\theta_i(x_b) = \theta_{ib}, \quad (4.28)$$

$$\theta_e(x_b) = 0. \quad (4.29)$$

The equation for determining Δx is as follows:

$$\frac{(1+\Lambda_i)\beta_2^2 - 1 - \Lambda_m\Lambda_e \sinh(\beta_1 x_b)}{\beta_2^2\Lambda_i - 1} \frac{\beta_1 x_b}{\beta_1 x_b} = \frac{(1+\Lambda_i)\beta_1^2 + 1 + \Lambda_m\Lambda_e \sin(\beta_2 x_b)}{\beta_1^2\Lambda_i + 1} \frac{\beta_2 x_b}{\beta_2 x_b}. \quad (4.30)$$

Both parts are equal to $\theta_{ib} > 0$. The relationship between Δx , $\Delta\Lambda_e$ and θ_{ib} is established by the relations (4.26) and (4.27) with the substitutions $\theta_{eb} \rightarrow \theta_{ib}$, $\Delta\Lambda_e \rightarrow -\Delta\Lambda_e$ and $\Lambda_i \rightarrow 1/\Lambda_i$, while the other parameters remain unchanged. The condition $\theta_{ib} > 0$ implies that $\Delta x < 0$ and $\Delta\Lambda_e < 0$. An upper limit on θ_{ib} is found from the inequality $W_0 \geq 0$ and the balance between the pressure and the forces of gravitational attraction and electrostatic repulsion acting on the ions at the boundary of the electron component [cf. (3.2)]:

$$\theta_{ib} \leq \lambda_G^{-1/(1+\eta_i)} \left[\frac{(1+\eta_i)(-q_s + \lambda_m/\Lambda_m)^2}{2\Lambda_i(1+\Lambda_m)(1-1/\Lambda_G)} \right] \times \left(\frac{M_s}{4\pi r_0^3 n_{i0} m_i} \right)^2 \left(\frac{r_0}{R_s} \right)^4 \Big]^{1/(1+\eta_i)}. \quad (4.31)$$

An ionosphere exists provided that $q_s < \lambda_m/\Lambda_m = A/Z$.

For positive values of $\Delta\Lambda_e$, the substance in the center of the star is closer to the electrically neutral state than in the regular solution. On the surface of the baryon component, the bulk charge is positive, and the electrosphere compensates for this charge either partially, completely, or excessively. Suppression of the electron fraction increases the deviation from electroneutrality and corresponds to $\Delta\Lambda_e < 0$. Under the condition $q_s < A/Z$, the electron density vanishes when the ion density is still finite. On the surface of the electron component, the stellar charge is positive (by continuity) and increases with $|\Delta\Lambda_e|$. The ionosphere makes an additional contribution to this charge. $|\Delta\Lambda_e|$ can be increased until it violates (4.24) in the case of an electrosphere or (4.31) in the case of an ionosphere.

3. Charge-mass-radius relation

The general properties associated with a deviation from local electroneutrality are expected to be inherent in all realistic models. We thus consider the charge-mass-radius relation for stars. The mass and radius are expressed as follows:

$$\frac{M_s}{4\pi r_0^3 n_{i0} m_i} = \frac{\pi(1+\Lambda_i)}{\beta_2}, \quad (4.32)$$

$$\frac{R_s}{r_0} = \frac{\pi}{\beta_2}. \quad (4.33)$$

The violation of the LEC affects the mass and radius at higher orders of the expansion in $1/\sqrt{\Lambda_G}$. The correction to the radius is $\Delta R_s = r_0 \Delta x \sim r_0/\sqrt{\Lambda_G}$, so the mass-radius relation remains unchanged to within $\sim 1/\sqrt{\Lambda_G}$. The bulk charge of the star, q_s , and the electrosphere charge, q_e , are as follows:

$$\begin{aligned} \frac{Z}{A} q_s &= \frac{1-\Lambda_m\Lambda_i}{1+\Lambda_i} - \frac{(1+\Lambda_m)\Lambda_i}{(1+\Lambda_i)^2} \pi\Theta_{eb}, \\ \frac{Z}{A} q_e &= -\frac{1+\Lambda_m}{1+\Lambda_i} \left[-\sqrt{\left(1 - \frac{\Lambda_i}{1+\Lambda_i} \pi\Theta_{eb}\right)^2 - \frac{\Lambda_i}{1+\Lambda_i} \pi^2 \Theta_{eb}^2} \right. \\ &\quad \left. + \left(1 - \frac{\Lambda_i}{1+\Lambda_i} \pi\Theta_{eb}\right) \right], \end{aligned} \quad (4.34)$$

where $\Theta_{eb} = \beta_1 \theta_{eb}/\beta_2 \sim 1$. Both formulas are valid with an accuracy of $\sim 1/\sqrt{\Lambda_G}$. The existence of an electron shell

requires an excess of electrons relative to the solution $\theta_{eb} = \theta_{ib} = 0$. Accordingly, the second term in q_s is negative, and as Θ_{eb} increases, its contribution reduces the stellar charge. By equating the root expression to zero, we can obtain the maximum value of Θ_{eb} :

$$\Theta_{eb}^{\max} = \frac{(\sqrt{1+1/\Lambda_i} - 1)(1+\Lambda_i)}{\pi}. \quad (4.35)$$

This value corresponds to the upper limit in Eq. (4.24). For $\Theta_{eb} = \Theta_{eb}^{\max}$, the shell overcompensates for the positive charge of the star: $(q_s + q_e)^{\min} = -\lambda_m < 0$.

An ionosphere forms for a lower concentration of electrons. The bulk charge, q_s , measured at the boundary of the electron component and the ion envelope charge, q_e , are given by the expressions

$$\begin{aligned} \frac{Z}{A} q_s &= \frac{1-\Lambda_m\Lambda_i}{1+\Lambda_i} + \frac{(1+\Lambda_m)\Lambda_i}{(1+\Lambda_i)^2} \pi\Theta_{ib}, \\ \frac{Z}{A} q_e &= \frac{(1+\Lambda_m)\Lambda_i}{1+\Lambda_i} \left[-\sqrt{\left(1 - \frac{1}{1+\Lambda_i} \pi\Theta_{ib}\right)^2 - \frac{1}{1+\Lambda_i} \pi^2 \Theta_{ib}^2} \right. \\ &\quad \left. + \left(1 - \frac{1}{1+\Lambda_i} \pi\Theta_{ib}\right) \right], \end{aligned} \quad (4.36)$$

where $\Theta_{ib} = \beta_1 \theta_{ib}/\beta_2 \sim 1$. The charge q_s increases with Θ_{ib} . The radicand vanishes when Θ_{ib} takes its maximum value

$$\Theta_{ib}^{\max} = \frac{(\sqrt{1+\Lambda_i} - 1)(1+1/\Lambda_i)}{\pi}. \quad (4.37)$$

One can check that for $\Theta_{ib} = \Theta_{ib}^{\max}$, the inequality in Eq. (4.31) becomes an equality. The maximum total charge is equal to $(q_s + q_e)^{\max} = A/Z$. The condition $q_s < A/Z$ constrains the value of Θ_{ib} from above by $(1+\Lambda_i)/\pi$. However, this limit is weaker than $\Theta_{ib} \leq \Theta_{ib}^{\max}$.

Figure 4 plots the bulk charge of the star, q_s , and the total charge including the envelope, $q_s + q_e$, as functions of the difference $\Theta_{eb} - \Theta_{ib}$ in the interval $(-\Theta_{ib}^{\max}, \Theta_{eb}^{\max})$ for a mixture of electrons and protons, i.e., for $Z = A = 1$ and $\Lambda_i = \Lambda_m$. The region $\Theta_{eb} - \Theta_{ib} \sim 0$ is presented separately in Fig. 5 on an enlarged scale.

In summary, in the two-fluid model with unit polytropic indices, it is possible to explicitly construct the general solution describing charged stars with electro- and ionospheres. An electrosphere forms if the ion chemical potential vanishes at the boundary, $\theta_i(x_b) = 0$, whereas the electron chemical potential at the boundary is finite, $\theta_e(x_b) > 0$, according to Eqs. (4.22) and (4.23). An ionosphere forms in the opposite case; see Eqs. (4.28) and (4.29). The solution for the internal stellar structure




FIG. 4. The bulk charge q_s and the total charge $q_s + q_e$ as functions of the difference $\Theta_{eb} - \Theta_{ib}$ for a mixture of electrons and protons ($Z = A = 1$) and for $\Lambda_i = \Lambda_m$. Positive values of $\Theta_{eb} - \Theta_{ib}$ correspond to the formation of an electrosphere ($\Theta_{ib} = 0$), whereas negative values correspond to the formation of an ionosphere (proton envelope, $\Theta_{eb} = 0$). The region near zero is shown on an enlarged scale in Fig. 5.




FIG. 5. The bulk charge q_s and the total charge $q_s + q_e$ as functions of the difference $\Theta_{eb} - \Theta_{ib}$ on an enlarged scale. The notation is the same as in Fig. 4.

is unambiguously matched with the solution for the electro- or ionosphere. The guiding parameter of the problem is $\Delta\Lambda_e = \Lambda_e - \Lambda_e^{\text{reg}}$. The general solution also describes stars with zero total charge, but such stars are not locally neutral. In an exponentially small neighborhood of Λ_e^{reg} , the electron and ion densities can be varied without restrictions. The stellar envelope is sensitive to exponentially small deviations of Λ_e from Λ_e^{reg} . The same fact can be interpreted as indicating remarkable overall stellar stability: changes in the envelope, and, accordingly, in the total stellar charge (within acceptable limits) have an exponentially weak influence on the internal structure of a star.

V. GENERAL SOLUTION TO THE UNCONSTRAINED HYDROSTATIC EQUILIBRIUM EQUATIONS

From the analysis of the exactly solvable model, the specific nature of the problem becomes rather transparent: the general solution to Eqs. (2.12) and (2.13) contains a component that is irregular at $\Lambda_G = \infty$. In the model of Sec. IV, the regular component dominates up to the surface of the star, whereas the irregular component is exponentially small everywhere except for the subsurface layer $\Delta r \sim r_a \sim r_0/\sqrt{\Lambda_G}$, where the contributions of the two components are comparable. The exponential smallness of the irregular component makes perturbation theory appropriate for describing the inner layers of the star.

There are four independent parameters, Λ_e , Λ_i , Λ_m , and Λ_G , on which Eqs. (2.12) and (2.13) and their solutions depend. The regular solution is a three-parameter solution with the independent parameters Λ_i , Λ_m , and Λ_G . The function $\Lambda_e = \Lambda_e^{\text{reg}}$ is defined in Sec. II. C to $O(1/\Lambda_G)$. The general solution and the regular solution satisfy the equations with different values of Λ_e . We denote the variance by $\Delta\Lambda_e = \Lambda_e - \Lambda_e^{\text{reg}}$, $|\Delta\Lambda_e| \ll 1$. The parameter Λ_i is not modified since for the general case $\eta_e \neq \eta_i$ considered here, Λ_e and Λ_i are independent.

Let θ_{k0} denote the regular solution. The general solution is assumed to be of the form

$$\theta_k = \theta_{k0} + \chi_k, \quad (5.1)$$

where χ_k is a small correction. This representation is valid everywhere except within a thin subsurface layer. By substituting θ_k and θ_{k0} into Eqs. (2.12) and (2.13) and linearizing these equations in χ_k and $\Delta\Lambda_e$, we obtain

$$\Delta_x(\chi_e + \Lambda_i\chi_i) = -(\eta_i\theta_{i0}^{\eta_i-1}\chi_i + \Lambda_m\Lambda_{e0}\eta_e\theta_{e0}^{\eta_e-1}\chi_e + \Lambda_m\Delta\Lambda_e\theta_{e0}^{\eta_e}), \quad (5.2)$$

$$\Delta_x(\chi_e - \Lambda_m\Lambda_i\chi_i) = -\Lambda_G(\eta_i\theta_{i0}^{\eta_i-1}\chi_i - \Lambda_{e0}\eta_e\theta_{e0}^{\eta_e-1}\chi_e - \Delta\Lambda_e\theta_{e0}^{\eta_e}). \quad (5.3)$$

The initial conditions

$$\chi_k(0) = \chi'_k(0) = 0 \quad (5.4)$$

show that $\Delta\Lambda_e$ determines the overall scale of χ_k .

A. Irregular component in the WKB approximation

When both sides of Eq. (5.3) are divided by Λ_G , the Laplacian acquires the small coefficient $1/\Lambda_G$. A similar case occurs in the Schrödinger wave equation, in which the kinetic term with the Laplacian contains \hbar^2 . In the semiclassical limit, \hbar is small, which makes the similarity of Eqs. (5.2) and (5.3) to the semiclassical limit of the Schrödinger wave equation quite obvious. In quantum

mechanics, the Wentzel, Kramers, and Brillouin approximation is used to describe the semiclassical limit; this approach is known as the WKB method (see, e.g., [34]).

Following the general scheme, we look for a solution of the form

$$\chi_k^{\text{irr}}(x) = g_k(x) \exp(\sqrt{\Lambda_G} S(x)). \quad (5.5)$$

In the neighborhood of $\Lambda_G = \infty$, the functions g_k admit a series expansion in $1/\sqrt{\Lambda_G}$:

$$g_k(x) = g_{k0}(x) + \frac{g_{k1}(x)}{\sqrt{\Lambda_G}} + \frac{g_{k2}(x)}{\Lambda_G} + \dots \quad (5.6)$$

The exponential factor causes the point $\Lambda_G = \infty$ to be an essentially singular point of $\chi_k^{\text{irr}}(x)$. Applying Δ_x to $\chi_k^{\text{irr}}(x)$ yields

$$\begin{aligned} \Delta_x \chi_k^{\text{irr}} &= \exp(\sqrt{\Lambda_G} S) \\ &\times \left[\Lambda_G g_k(S')^2 + \sqrt{\Lambda_G} \frac{(x^2 g_k^2 S')'}{x^2 g_k} + \frac{(x^2 g_k')'}{x^2} \right]. \end{aligned} \quad (5.7)$$

The coefficients of powers of $\sqrt{\Lambda_G}$ are as follows:

$$\Lambda_G: g_{k0}(S')^2, \quad (5.8)$$

$$\sqrt{\Lambda_G}: g_{k1}(S')^2 + \frac{(x^2 g_{k0}^2 S')'}{x^2 g_{k0}}, \quad (5.9)$$

$$1: g_{k2}(S')^2 + \frac{(x^2 g_{k1}^2 S')'}{x^2 g_{k1}} + \frac{(x^2 g_{k0}')'}{x^2}, \quad (5.10)$$

where the common exponential factor is suppressed.

The Λ_G term in Eq. (5.2) yields the relation

$$g_{e0} + \Lambda_i g_{i0} = 0. \quad (5.11)$$

By substituting this relation into Eq. (5.3), we obtain with the same accuracy

$$S' = \pm \sqrt{\frac{\eta_i \theta_{i0}^{\eta_i-1} + \Lambda_i \Lambda_e^{\text{reg}} \eta_e \theta_{e0}^{\eta_e-1}}{\Lambda_i (1 + \Lambda_m)}}. \quad (5.12)$$

The exponent can be found by integrating Eq. (5.12). In the model with unit polytropic indices, this expression gives a simple linear dependence, $S \sim x$.

The $\sqrt{\Lambda_G}$ term in Eq. (5.2) leads to the relation

$$g_{e1} + \Lambda_i g_{i1} = 0. \quad (5.13)$$

Using this equality in Eqs. (5.3) and (5.12), we find that the second differential term in Eq. (5.9) vanishes. We thus obtain

$$x^2 g_{e0}^2 S' = \text{const.} \quad (5.14)$$

Near the center, g_{k0} behaves as $g_{k0} \sim 1/x$. Equation (5.12) leads to

$$S_{\pm}(x) = \pm \int_0^x \sqrt{\frac{\eta_i \theta_{i0}^{\eta_i-1}(x') + \Lambda_i \Lambda_e^{\text{reg}} \eta_e \theta_{e0}^{\eta_e-1}(x')}{\Lambda_i (1 + \Lambda_m)}} dx'.$$

The terms with S_{\pm} appear in the expression for $\chi_{k0}^{\text{irr}}(x)$ in a certain combination. The singularity at $x=0$ can be eliminated by taking the difference $\exp(\sqrt{\Lambda_G} S_+(x)) - \exp(\sqrt{\Lambda_G} S_-(x))$. Finally,

$$\chi_{k0}^{\text{irr}}(x) = C_{k0} \frac{\sinh(\sqrt{\Lambda_G} S_+(x))}{x \sqrt{S'_+(x)}} + O(\Lambda_G^{-\frac{1}{2}}), \quad (5.15)$$

where the C_{k0} are arbitrary constants.

The higher-order terms of the expansion can be found in a similar manner.

The functions $\chi_{k0}^{\text{irr}}(x)$ do not vanish at $x=0$. The normalization condition $\theta_k(0) = 1$ is restored when considering a particular solution of the inhomogeneous ODE system defined by Eqs. (5.2) and (5.3).

B. Correction to the irregular component

A particular solution to Eqs. (5.2) and (5.3) can be found from the original ODE system defined by Eqs. (2.12) and (2.13). We can construct the regular solution to this system, starting from the LEC solution, in the form of a series expansion in $1/\Lambda_G$. The functions $\theta_{k0}(x)$ and Λ_e^{reg} that are obtained in this way depend on the parameters Λ_i , Λ_m , and Λ_G .

Suppose that we are looking for the regular solution to Eqs. (2.12) and (2.13) with the initial conditions $\theta_k(0) \neq 1$ and $\theta'_k(0) = 0$ for $|\theta_k(0) - 1| \ll 1$. Let $\hat{\theta}_{k0}(x)$ denote the corresponding solutions, which depend on the parameters Λ_i , Λ_m , and Λ_G . The same parameters determine $\Lambda_e \equiv \Lambda_e^{\text{reg}}$.

It is not difficult to see that by rescaling x and $\hat{\theta}_{k0}(x)$, the problem can be reduced to the initial problem with the standard initial conditions, the rescaled variable \hat{x} ,

$$\hat{x} = x \sqrt{\frac{\theta_i^{\eta_i}(0)}{\theta_e^{\eta_e}(0)}},$$

and the new parameters Λ_e and Λ_i :

$$\hat{\Lambda}_e = \Lambda_e^{\text{reg}} \frac{\theta_e^{\eta_e}(0)}{\theta_i^{\eta_i}(0)}, \quad \hat{\Lambda}_i = \Lambda_i \frac{\theta_i(0)}{\theta_e(0)}.$$

By contrast, Λ_m is not modified. Obviously, $\hat{\Lambda}_e$ is a known function of $\hat{\Lambda}_i$, namely, $\hat{\Lambda}_e = \Lambda_e^{\text{reg}}(\hat{\Lambda}_i)$; thus,

$$\Lambda_e^{\text{reg}}(\Lambda_i) = \frac{\theta_i^{\eta_i}(0)}{\theta_e^{\eta_e}(0)} \Lambda_e^{\text{reg}} \left(\Lambda_i \frac{\theta_i(0)}{\theta_e(0)} \right).$$

Under the assumption of the smallness of $\Delta\theta_k = \theta_k(0) - 1$, we obtain

$$\begin{aligned} \Delta\Lambda_e &= \Lambda_e^{\text{reg}'} - \Lambda_e^{\text{reg}} = \Delta\theta_i \left(\eta_i \Lambda_e^{\text{reg}} + \Lambda_i \frac{\partial \Lambda_e^{\text{reg}}}{\partial \Lambda_i} \right) \\ &\quad - \Delta\theta_e \left(\eta_e \Lambda_e^{\text{reg}} + \Lambda_i \frac{\partial \Lambda_e^{\text{reg}}}{\partial \Lambda_i} \right). \end{aligned}$$

The solution to Eqs. (2.12) and (2.13) with the nonstandard boundary conditions can be written as follows:

$$\hat{\theta}_{k0}(x) = \theta_k(0)\theta_{k0} \left(x \sqrt{\frac{\theta_i^{\text{h}}(0)}{\theta_e(0)}}, \Lambda_i \frac{\theta_i(0)}{\theta_e(0)} \right). \quad (5.16)$$

The function

$$\chi_{k0} = \hat{\theta}_{k0} - \theta_{k0} \quad (5.17)$$

satisfies the linearized equations (5.2) and (5.3) for $\Delta\Lambda_e = \Lambda_e^{\text{reg}'} - \Lambda_e^{\text{reg}}$ and the initial conditions $\chi_{k0}(0) = \Delta\theta_k$. For small $\Delta\theta_k$, we can write

$$\begin{aligned} \chi_{k0}(x) &= \theta_{k0}(x)\Delta\theta_k + x \frac{\partial \theta_{k0}(x)}{\partial x} \frac{\eta_i \Delta\theta_i - \Delta\theta_e}{2} \\ &\quad + \Lambda_i \frac{\partial \theta_{k0}(x)}{\partial \Lambda_i} (\Delta\theta_i - \Delta\theta_e). \end{aligned} \quad (5.18)$$

$$\begin{aligned} \chi_i(x) &= \frac{C_{i0}}{x\sqrt{S'_+(x)}} \sinh \left(\sqrt{\Lambda_G} S_+(x) \right) - C_{i0} \sqrt{\Lambda_G S'_+(0)} \left[\theta_{i0}(x) + x \frac{\partial \theta_{i0}(x)}{\partial x} \frac{\eta_i + \Lambda_i}{2} + \Lambda_i \frac{\partial \theta_{i0}(x)}{\partial \Lambda_i} (1 + \Lambda_i) \right], \\ \chi_e(x) &= -\frac{C_{i0}\Lambda_i}{x\sqrt{S'_+(x)}} \sinh \left(\sqrt{\Lambda_G} S_+(x) \right) + C_{i0} \sqrt{\Lambda_G S'_+(0)} \left[\Lambda_i \theta_{e0}(x) - \frac{\partial \theta_{e0}(x)}{\partial x} \frac{\eta_i + \Lambda_i}{2} - \Lambda_i \frac{\partial \theta_{e0}(x)}{\partial \Lambda_i} (1 + \Lambda_i) \right]. \end{aligned}$$

In each of these expressions, the first term increases exponentially with x , dominates the second (regular) term for $x \gtrsim 1/\sqrt{\Lambda_G}$, and becomes comparable to the regular component θ_{k0} near the surface in the layer $|x - x_b| \sim \theta_{eb}$, where perturbation theory in χ_k is no longer applicable. The constant C_{i0} is exponentially small. This feature has been noted earlier in the analysis of the exactly solvable model of Sec. IV. The exponential smallness of the irregular solution in the inner layers of the star is obviously a common property. The functions $\chi_i(x)$ and $\chi_e(x)$ differ in sign, and the electron component is suppressed by approximately Λ_i .

The matching of θ_k and θ'_k with the electrosphere solution requires

$$\chi_e(x_b) \sim \theta_{eb}, \quad \chi'_e(x_b) \sim 1.$$

The first of the equations allows to fix C_{i0} . For $\theta_i(x_b) = 0$, Eq. (5.12) leads to $S'_+(x_b) \sim \theta_{eb}^{(\eta_e - 1)/2}$; the second equation then follows automatically. One can verify that for $\eta_e = \eta_i = 1$, χ_k^{irr} is analytic in $\Lambda_G \in \mathbb{C}^1 \setminus \infty$, with $\Lambda_G = \infty$

C. General solution

The irregular component of the general solution is given by the first term of the expansion in Eq. (5.6). Equation (5.15) is a solution to the homogeneous equation, whereas Eq. (5.18) is a solution to the nonhomogeneous equation. The general solution is defined by Eq. (5.1) with $\chi_k = \chi_k^{\text{irr}} + \chi_{k0}$. Among these functions, θ_{k0} and χ_{k0} are regular at $\Lambda_G = \infty$, whereas χ_k^{irr} is singular at $\Lambda_G = \infty$. The initial conditions (5.4) lead to the relation

$$C_{k0} \sqrt{\Lambda_G S'_+(0)} + \Delta\theta_k = 0, \quad (5.19)$$

where the C_{k0} are the constants that appear in Eq. (5.15). The derivative of χ_{k0} at $x = 0$ vanishes by construction. In the general case, $S_{\pm}(x) = xS'_{\pm}(0) + O(x^3)$; consequently, the derivative χ_k^{irr} also vanishes automatically at $x = 0$. As a consequence of Eq. (5.11), the coefficients C_{k0} satisfy

$$C_{e0} + \Lambda_i C_{i0} = 0.$$

By expressing the unknown coefficients in terms of C_{i0} , we can write

being a unique singular point of χ_k^{irr} , in agreement with the analysis of Sec. IV. B.

Standard numerical methods for solving ODE systems do not allow the separation of the regular solution from the irregular one due to the exponential smallness of the latter. A numerical solution to Eqs. (2.12) and (2.13) for $x \lesssim 1$ will inevitably contain an admixture of the irregular component, which is interpreted by the program as a machine zero. In the subsurface layer $|x - x_b| \sim \theta_{eb}$, due to the exponential growth of $\chi_k^{\text{irr}}(x)$, this machine zero is converted into a finite but random value. The stellar properties that are sensitive to $\Delta\Lambda_e$ cannot be determined using numerical methods. A change in the integration, e.g., of the grid step, is thus expected to result in uncontrolled changes in the stellar charge, the thickness of the electro- or ionosphere, and other quantities that depend on $\Delta\Lambda_e$. In view of these remarks, the numerical solutions of Refs. [13–15] are likely to be unstable.

At the same time, as the example of Eqs. (4.34) and (4.36) shows, relationships which do not depend on

$\Delta\Lambda_e$ are numerically stable. Each integration with a varying grid step leads to random values of q_s and θ_{eb} , but all points on the plane (q_s, θ_{eb}) must lie on a smooth curve. In the exactly solvable model of Sec. IV, the quantities q_s and θ_{eb} are related by a linear dependence.

VI. CONCLUSIONS

In this paper, we considered the unconstrained HE equations for stars in the absence of local electroneutrality. The condition of local electroneutrality is usually imposed on the EoS phenomenologically as an additional constraint (LEC). In our approach, the electrostatic interactions are considered *ab initio*; as a result, the LEC is satisfied approximately, with the uncompensated charge density being $1/\lambda_G \sim 10^{-36}$ of the particle number density. The smallness of the parameter $1/\lambda_G$ makes it possible to use the LEC to describe with a high accuracy properties that are not related to stellar electrostatics.

We found that the mass-radius relationship and the condition for gravitational stability for charged stars are not modified to an accuracy of $\sim \lambda_G^{-1/(1+\eta_e)}$. The admissible states of charged stars are determined by the condition that the electro- and ionospheres do not extend to infinity. This requirement provides lower and upper bounds on the total stellar charge of -0.1 and 150 C per solar mass, respectively.

The main system of unconstrained HE equations, Eqs. (II.12) and (II.13), belongs to the class of singularly perturbed ODEs. In such a system, the perturbation parameter (in our case, $1/\lambda_G$) appears as a coefficient of the higher derivative and thus plays the main role in specifying the nature of the problem. The use of conventional (regular) methods for solving ODEs via series expansion in a small parameter leads to the loss of certain solutions in this case. In the limit of $\lambda_G \rightarrow \infty$, Eq. (II.13) becomes a constraint on the particle concentrations, which we identify as an LEC.

The Poincaré theorem on analyticity and Dyson's argument indicate that the general solution to Eqs. (II.12)—(II.13) is singular for $\lambda_G = \infty$. A particular, regular solution can exist provided that one of the parameters, e.g., Λ_e , is replaced by a function of λ_G . Such a solution can be constructed as a formal power series in G ($\sim 1/\lambda_G$) starting from a locally neutral solution. Only these two types of solutions, namely, regular ones and locally neutral ones, have been discussed in the literature to date.

In this paper, the general solution to the HE problem was first constructed in NGT.

In Sec. IV, we presented an exactly solvable model of a two-component mixture of ions and electrons with unit polytropic indices. The general solution to the unconstrained HE equations appears to be irregular in the neighborhood of $\lambda_G = \infty$. The irregular component is exponentially small in the inner layers of the star.

In the general case, the WKB method can be used to construct the irregular component. Its magnitude is

comparable to that of the regular component in a subsurface layer of thickness $\sim \lambda_G^{-1/(1+\eta_e)}$, where these components interact non-linearly and determine the structure of the electro- or ionosphere. The properties of these envelopes are exponentially sensitive to variations in the charge density in the central regions of the star.

In general, stars are characterized by their mass, radius and charge. Nuclear reactions in the interior of a star cause variations in the parameters Λ_e , Λ_i , and Λ_m and thereby affect the bulk stellar charge, which, due to the conservation of electric charge, leads to a charge exchange between the envelope and the inner region.

The structure of the general solution to the unconstrained HE equations can be described as follows:

Starting from a locally neutral solution, one can construct a regular solution (θ_{k0} in Sec. V) in the form of a series in powers of G . On the basis of this solution, an irregular component can be constructed using the WKB method (χ_k^{irr} in Sec. V). This component is singular for $\lambda_G = \infty$. A correction to this irregular component is needed to satisfy the initial conditions (χ_{k0} in Sec. V). The sum of these three functions yields the general solution, excluding only a thin subsurface layer of thickness $\sim \lambda_G^{-1/(1+\eta_e)}$, where the magnitude of the irregular component becomes comparable to that of the regular component and the WKB method is no longer applicable. Numerical methods should then be used to match, within the subsurface layer, the general solution with the known solution for the electro- or ionosphere.

Extension of the formalism developed here to GRT is left as an open problem. An important direction of study is generalization to three or more component fluids in chemical equilibrium. Yet another interesting application is to find the general solution of the unconstrained HE equations for strange stars.

ACKNOWLEDGMENTS

The authors are grateful to S.I. Blinnikov and I.L. Iosilevsky for stimulating discussions. M.I.K. wishes to acknowledge M.D. Malykh for discussion of analytic properties of the solutions. The work is supported in part by the RFBR Grants No. 16-02-00228 and No. 18-02-00733.

APPENDIX: INCOMPRESSIBLE TWO-FLUID MODEL

As a second example of an exactly solvable, albeit somewhat exotic, model, we consider the case of $\eta_i = \eta_e = 0$, which corresponds to an incompressible fluid ($n = n_0 \theta^\eta \sim \text{const}$). Although the speed of sound in such a substance is infinite, this case is of methodological interest with regard to the restrictions on the parameter Λ_e in the regular solution, especially as $S_\pm(x) \rightarrow 0$ for $\eta_i = \eta_e = 0$. For an incompressible fluid, Eqs. (2.12) and (2.13) become

$$\Delta_x(\theta_e + \Lambda_i \theta_i) = -(1 + \Lambda_m \Lambda_e), \quad (\text{A1})$$

$$\Delta_x(\theta_e - \Lambda_m \Lambda_i \theta_i) = -\Lambda_G(1 - \Lambda_e). \quad (\text{A2})$$

Their solutions for the initial conditions $\theta_k(0) = 1$ and $\theta'_k(0) = 0$ have the form

$$\theta_i = 1 - \frac{(1 + \Lambda_m \Lambda_e) - \Lambda_G(1 - \Lambda_e)}{6\Lambda_i(1 + \Lambda_m)} x^2, \quad (\text{A3})$$

$$\theta_e = 1 - \frac{\Lambda_m(1 + \Lambda_m \Lambda_e) + \Lambda_G(1 - \Lambda_e)}{6(1 + \Lambda_m)} x^2. \quad (\text{A4})$$

In principle, these solutions describe a charged stellar configuration free from constraints on Λ_e . This freedom is due to the irregular nature of the solution for $\Lambda_G = \infty$. The requirement of regularity leads to the LEC $\Lambda_e = 1$, given that Λ_e is a parameter. If Λ_e is considered to be a function of Λ_G , then $\Lambda_e = 1 + O(1/\Lambda_G)$. The solutions (A3) and (A4) are valid in the inner region of the star. The outer region requires special consideration.

According to Eq. (A3), the ion concentration vanishes for

$$x_i = \sqrt{\frac{6\Lambda_i(1 + \Lambda_m)}{(1 + \Lambda_m \Lambda_e) - \Lambda_G(1 - \Lambda_e)}}. \quad (\text{A5})$$

If we require the electron concentration to vanish at the same point, $x_e = x_i$, then we obtain $\Lambda_e = \Lambda_e^{\text{reg}}$, where Λ_e^{reg} is given by Eq. (2.22). In this case, the star still has a charge. Suppose now that we do not require the concentrations to vanish simultaneously. Again, let us consider the outer part of the star, the electrosphere, where the ion concentration is zero. The equation for θ_e reduces to

$$(1 + \Lambda_m)\Delta_x(\theta_e) = \Lambda_e(\Lambda_G - \Lambda_m^2). \quad (\text{A6})$$

At the boundary $x = x_i$, the function θ_e and its first derivative are continuous. The solution to Eq. (A6) takes the form

$$\theta_{\text{ea}} = 1 - \frac{x_i^2(\Lambda_G + \Lambda_m)}{2(1 + \Lambda_m)} \left(1 - \frac{2x_i}{3x}\right) + \frac{\Lambda_e(\Lambda_G - \Lambda_m^2)}{6(1 + \Lambda_m)} x^2.$$

For an electrically neutral star, the boundary $x = x_e$, where $\theta_{\text{ea}}(x_e) = 0$, determines the total charge. The bulk charge is proportional to [cf. Eq. (2.35)]

$$Q_s \sim -x^2(\theta'_e - \Lambda_m \Lambda_i \theta'_i)|_0^{x_i}, \quad (\text{A7})$$

whereas the charge of the envelope is proportional to

$$Q_e \sim -\frac{\Lambda_G(1 + \Lambda_m)}{\Lambda_G - \Lambda_m^2} x^2 \theta'_{\text{ea}}|_{x_i}^{x_e}. \quad (\text{A8})$$

The condition $Q_s + Q_e = 0$ leads to the relation

$$x_e = x_i / \Lambda_e^{1/3}, \quad (\text{A9})$$

whereas the condition $\theta_{\text{ea}}(x_e) = 0$ leads to the following equation:

$$\begin{aligned} \Lambda_e(\Lambda_G + \Lambda_m) + \Lambda_i \Lambda_e^{1/3}(3\Lambda_G + 2\Lambda_m - \Lambda_m^2) \\ = \Lambda_G - 1 + 3\Lambda_i(\Lambda_G + \Lambda_m), \end{aligned} \quad (\text{A10})$$

whose solution, Λ_e , coincides with Λ_e^{reg} to an accuracy of $1/\Lambda_G$.

-
- [1] J. L. Lebowitz and E. H. Lieb, *Phys. Rev. Lett.* **22**, 631 (1969).
 - [2] F. J. Dyson and A. J. Lenard, *J. Math. Phys.* **8**, 423 (1967).
 - [3] E. H. Lieb and W. Thirring, *Phys. Rev. Lett.* **35**, 1116 (1975).
 - [4] L. D. Landau and E. M. Lifshitz, *Statistical Physics*, 3rd ed. (Butterworth-Heinemann, Washington, DC, 1980), Vol. 5.
 - [5] O. Klein, *Rev. Mod. Phys.* **21**, 531 (1949).
 - [6] T. Kodama and M. Yamada, *Prog. Theor. Phys.* **47**, 444 (1972).
 - [7] E. Olson and M. Bailyn, *Phys. Rev. D* **12**, 3030 (1975).
 - [8] S. Rosseland, *Mon. Not. R. Astron. Soc.* **84**, 525 (1924).
 - [9] E. L. Schatzman, *White Dwarfs* (North-Holland, Amsterdam, 1958).
 - [10] E. Olson and M. Bailyn, *Phys. Rev. D* **18**, 2175 (1978).
 - [11] J. Bally and E. R. Harrison, *Astrophys. J.* **220**, 743 (1978).
 - [12] L. Neslušan, *Astron. Astrophys.* **372**, 913 (2001).
 - [13] M. Rotondo, J. A. Rueda, R. Ruffini, and S.-S. Xue, *Phys. Lett. B* **701**, 667-671 (2011).
 - [14] R. Belvedere, D. Pugliese, J. A. Rueda, R. Ruffini, and S.-S. Xue, *Nucl. Phys.* **A883**, 1 (2012).
 - [15] R. Belvedere, J. A. Rueda, and R. Ruffini, *J. Korean Phys. Soc.* **65**, 897 (2014).
 - [16] A. B. Gorshkov and V. A. Baturin, *Astronomy Reports* **52**, 760 (2008).
 - [17] M. V. Beznogov and D. G. Yakovlev, *Phys. Rev. Lett.* **111**, 161101 (2013).
 - [18] D. K. Nadyozhin and V. S. Imshennik, *Int. J. Mod. Phys. A* **20**, 6597 (2005).
 - [19] M. I. Krivoruchenko, D. K. Nadyozhin, T. L. Razinkova, Yu. A. Simonov, M. A. Trusov, and A. V. Yudin, *Phys. At. Nucl.* **74**, 371 (2011).

- [20] D. K. Nadyozhin, in *Proceedings of the Conference Quark Phase Transition in Compact Objects and Multimessenger Astronomy: Neutrino Signals, Supernovae and Gamma-Ray Bursts*, 2015, the Special Astrophysical Observatory of Russian Academy of Sciences (SAO RAS) and Baksan Neutrino Observatory of Institute for Nuclear Research of Russian Academy of Sciences (BNO INR RAS), edited by V. V. Sokolov *et al.* (Sneg, Pyatigorsk 2016), p. 69.
- [21] S. B. Pikel'ner, *Fundamentals of Cosmic Electrodynamics* (Fizmatgiz, Moscow, 1961; NASA technical translation. TT F-175, 1964).
- [22] S. Ray, A. L. Espíndola, M. Malheiro, J. P. S. Lemos, and V. T. Zanchin, *Phys. Rev. D* **68**, 084004 (2003).
- [23] Y. Ilyashenko and S. Yakovenko, *Lectures on Analytic Differential Equations* (American Mathematical Society, Providence, 2008), Theorem 1.1.
- [24] F. J. Dyson, *Phys. Rev.* **85**, 631 (1952).
- [25] R. E. O'Malley, Jr, *Singular Perturbation Methods for Ordinary Differential Equations* (Springer-Verlag, New York, 1991).
- [26] G. P. Horedt, *Polytropes. Applications in Astrophysics and Related Fields* (Kluwer Academic Publishers, Boston, 2004).
- [27] I. L. Iosilevskiy, *J. Phys. A* **42**, 214008 (2009).
- [28] A. V. Yudin *et al.* (to be published).
- [29] G. Chavalier, *The Earth's Electrical Surface Potential. A Summary of Present Understanding* (California Institute for Human Science, Encinitas, 2007).
- [30] L. Iorio, *Gen. Relativ. Gravit.* **44**, 1753 (2012).
- [31] E. Witten, *Phys. Rev. D* **30**, 272 (1984).
- [32] J. Hu and R. X. Xu, *Astron. Astrophys.* **387**, 710 (2002).
- [33] M. I. Krivoruchenko and B. V. Martemyanov, *Yad. Fiz.* **58**, 536 (1995) [*Phys. At. Nucl.* **58**, 484 (1995)].
- [34] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics: Non-Relativistic Theory*, 3rd ed. (Pergamon Press, Oxford, 1977), Vol. 3.
- [35] J. D. Bekenstein, *Phys. Rev. D* **4**, 2185 (1971).
- [36] P. D. Lasky and A. W. C. Lun, *Phys. Rev. D* **75**, 104010 (2007).
- [37] C. R. Ghezzi, *Phys. Rev. D* **72**, 104017 (2005).
- [38] R. P. Negreiros, F. Weber, M. Malheiro, and V. Usov, *Phys. Rev. D* **80**, 083006 (2009).
- [39] J. D. V. Arbanil and M. Malheiro, *Phys. Rev. D* **92**, 084009 (2015).
- [40] M. I. Krivoruchenko, *Pis'ma Zh. Eksp. Teor. Fiz.* **46**, 5 (1987) [*JETP Lett.* **46**, 3 (1987)].
- [41] M. I. Krivoruchenko and B. V. Martemyanov, *Astrophys. J.* **378**, 628 (1991).