

Inflationary scale, reheating scale, and pre-BBN cosmology with scalar fieldsAlessandro Di Marco,^{*} Gianfranco Pradisi, and Paolo Cabella*University of Rome—Tor Vergata, Via della Ricerca Scientifica 1 INFN Sezione di Roma Tor Vergata, via della Ricerca Scientifica 1, 00133 Roma, Italy*

(Received 17 July 2018; published 13 December 2018)

In this paper, we discuss the constraints on the reheating temperature supposing an early postreheating cosmological phase dominated by one or more simple scalar fields produced from inflaton decay and decoupled from matter and radiation. In addition, we explore the combined effects of the reheating and nonstandard scalar field phases on the inflationary number of e -foldings.

DOI: [10.1103/PhysRevD.98.123511](https://doi.org/10.1103/PhysRevD.98.123511)**I. INTRODUCTION**

Before the hot big bang (HBB) epoch, our Universe likely experienced an early quantum gravity phase (at the so-called Planck scale) in which gravitational, strong, weak and electromagnetic interactions were unified in a single fundamental force [1]. Due to expansion and cooling, at lower (GUT) scales the gravitational interaction decoupled and the Universe entered an hypothetical phase where matter and radiation can be described in terms of a Grand-Unified gauge theory [2]. According to the inflationary paradigm, at a scale M_{inf} ($< 10^{16}$ GeV), after the spontaneous symmetry breaking to $SU(3) \times SU(2) \times U(1)$ (the gauge group of the Standard Model of particle physics), cosmological inflation is supposed to have taken place, in order to make the Universe almost flat, isotropic and homogeneous on large astronomical scales [3]. In the simplest version, the inflationary mechanism was driven by a scalar field, called inflaton, minimally coupled to gravity and probing an almost flat region (a false vacuum) of the corresponding effective scalar potential. At the end of inflation, where the potential steepens, the inflaton field falls in the global minimum of the potential, oscillates, decays, and “reheats” the Universe (see [4] for detailed studies on the mechanism and [5] for general constraints), giving rise to the standard HBB evolution characterized by an initial radiation dominated phase. However, this last step is not necessarily the unique possible scenario. Indeed, there is of course room for a peculiar evolution in the history of the Universe immediately after the reheating. In particular, the expansion of the Universe could have been submitted to additional phases where, for instance, it was driven by one (or more) new simple scalar species, before the radiation-dominated era and, especially, well before the big bang nucleosynthesis (BBN). Additional scalar fields, not necessarily directly interacting with the standard model

degrees of freedom (d.o.f.), are quite common in superstring theory with branes. They typically parametrize the brane positions along directions internal to the extra dimensions transverse to the branes. Since their energy density exhibits a modified dilution law, they can give rise to a nonstandard postreheating phase. Scenarios of this type have been recently introduced to study modification on relic abundances and decay rates of dark matter [6,7], as well as enhancements in the inflationary number of e -foldings [8,9]. In this paper, we consider nonstandard cosmologies inspired by string theory orientifold models [10] with, generically, multiple sterile scalar fields entering a nonstandard postreheating phase and we analyze in details the constraints put on the reheating temperature by the additional fields. As a consequence, we can derive more stringent model independent predictions about the number of e -folds during inflation. The paper is organized as follows. In Sec. II, we derive general expressions for the energy density in the case of nonstandard postreheating cosmological evolution, given by one or more scalar fields. In Sec. III, we discuss how the features of the new species affect the reheating scale. In particular, we derive an upper limit to the reheating temperature. In Sec. IV, we study the relation between reheating and postinflationary scalar fields and we calculate the inflationary number of e -foldings, also constrained by the maximum reheating temperature. In Sec. V, we add our conclusions and some discussions. In the Appendices, we show numerical examples of the consequences of the variation in the number of e -foldings on the inflationary predictions of n_s and r , for various selected inflaton potentials. In this manuscript, we use the particle natural units $c = \hbar = 1$, unless otherwise stated.

II. POSTINFLATIONARY SCALAR FIELDS AND COSMOLOGY

The cosmological history of the early Universe immediately after the reheating should be characterized by a

^{*}alessandro.di.marco@roma2.infn.it

radiation dominated era. In that phase, the corresponding evolution is well described by

$$H^2(T) \simeq \frac{1}{3M_{\text{Pl}}^2} \rho_{\text{rad}}(T), \quad \rho_{\text{rad}}(T) = \frac{\pi^2}{30} g_E(T) T^4, \quad (1)$$

where H denotes the Hubble rate, M_{Pl} is the reduced Planck mass, ρ_{rad} is the radiation energy density and T indicates the temperature scale of the universe at a given (radiation-dominated) epoch. Finally, g_E is the effective number of relativistic d.o.f. turning out to be

$$g_E(T) = \sum_b g_b \left(\frac{T_b}{T} \right)^4 + \frac{7}{8} \sum_f g_f \left(\frac{T_f}{T} \right)^4, \quad (2)$$

where b and f label contributions from bosonic and fermionic d.o.f., respectively, and T_b and T_f indicate the corresponding temperatures. In this section, we would like to analyze a modification of the evolution of the early Universe after the reheating phase, realized through the presence of a set of scalar fields $\phi_i (i = 1, \dots, k)$. They are assumed to dominate at different time scales until radiation becomes the most relevant component, well before the BBN era [6]. The last assumption is crucial in order to not spoil the theoretical successes related to the prediction of light element abundances (see [7]). Therefore, the total energy density after the inflaton decay can be assumed to be

$$\rho(T) = \rho_{\text{rad}}(T) + \sum_{i=1}^k \rho_{\phi_i}(T). \quad (3)$$

We introduce the scalar fields in such a way that, for $i > j$, ρ_{ϕ_i} hierarchically dominates at higher temperatures over ρ_{ϕ_j} when the temperature decreases. All the scalar fields that are supposed to be completely decoupled from each other and from matter and radiation fields can be described as perfect fluids diluting faster than radiation. In this respect, the dynamics is encoded in

$$\dot{\rho}_{\phi_i} + 3H\rho_{\phi_i}(1 + w_i) = 0, \quad (4)$$

where $w_i = w_{\phi_i}$ is the equation of state (EoS) parameter of the field i . Integrating this equation, one finds

$$\rho_{\phi_i}(T) = \rho_{\phi_i}(T_i) \left(\frac{a(T_i)}{a(T)} \right)^{4+n_i}, \quad n_i = 3w_i - 1, \quad (5)$$

where the index n_i , the ‘‘dilution’’ coefficient, is understood to satisfy the conditions

$$n_i > 0, \quad n_i < n_{i+1}. \quad (6)$$

T_i can be conveniently identified with the transition temperature at which the contribution of the energy density of ϕ_i becomes subdominant with respect to the one of ϕ_{i-1} . In other words, the scalar fields are such that

$$\rho_{\phi_i} > \rho_{\phi_{i-1}} \quad \text{for } T > T_i \quad (7)$$

$$\rho_{\phi_i} = \rho_{\phi_{i-1}} \quad \text{for } T = T_i \quad (8)$$

$$\rho_{\phi_i} < \rho_{\phi_{i-1}} \quad \text{for } T < T_i. \quad (9)$$

Using the conservation of the ‘‘comoving’’ entropy density,

$$g_S(T) a^3(T) T^3 = g_S(T_i) a^3(T_i) T_i^3, \quad (10)$$

with g_S defined by

$$g_S(T) = \sum_b g_b \left(\frac{T_b}{T} \right)^3 + \frac{7}{8} \sum_f g_f \left(\frac{T_f}{T} \right)^3, \quad (11)$$

the effective number of relativistic d.o.f. associated with entropy, the energy density of the various fields at a temperature T can be expressed in terms of the transition temperatures T_i [6,9]

$$\rho_{\phi_i}(T) = \rho_{\phi_i}(T_i) \left(\frac{g_S(T)}{g_S(T_i)} \right)^{\frac{4+n_i}{3}} \left(\frac{T}{T_i} \right)^{4+n_i}. \quad (12)$$

For the first scalar field ϕ_1 , by definition, the transition temperature is such that its energy density is identical to the one of the radiation fluid, so that

$$\rho_{\phi_1}(T_1) = \rho_{\text{rad}}(T_1) = \frac{\pi^2}{30} g_E(T_1) T_1^4. \quad (13)$$

The second scalar field ϕ_2 is subdominant compared to ϕ_1 below the temperature T_2 . Using Eq. (12) and observing that T_2 is the transition temperature at which $\rho_{\phi_2}(T_2) = \rho_{\phi_1}(T_2)$, one gets

$$\rho_{\phi_2}(T) = \rho_{\phi_1}(T_1) \left(\frac{T_2 g_S^{1/3}(T_2)}{T_1 g_S^{1/3}(T_1)} \right)^{4+n_1} \left(\frac{T g_S^{1/3}(T)}{T_2 g_S^{1/3}(T_2)} \right)^{4+n_2} \quad (14)$$

This equation tells us that the energy density of the scalar field ϕ_2 depends on the ratio between the two scales T_1 and T_2 , where the ϕ_1 -dominance occurs. In the same way, we can derive the analogous expression for the other scalar fields ϕ_i . The general expression for the energy density carried by ϕ_i turns out to be

$$\rho_{\phi_i}(T) = \rho_{\phi_1}(T_1) \prod_{j=1}^{i-1} \left(\frac{T_{j+1} g_S^{1/3}(T_{j+1})}{T_j g_S^{1/3}(T_j)} \right)^{4+n_j} \times \left(\frac{T g_S^{1/3}(T)}{T_i g_S^{1/3}(T_i)} \right)^{4+n_i}, \quad i \geq 2. \quad (15)$$

Inserted in Eq. (3), the previous expressions provide the total energy density dominating the expansion of the

Universe after the standard reheating phase, up to the beginning of the radiation-dominated epoch. In particular, the Hubble rate acquires the compact form

$$H^2(T) \simeq \frac{1}{3M_{\text{Pl}}^2} \rho_{\phi_1}(T_1) \sum_{i=1}^k f_i(n_i, T, T_1, \dots, T_i), \quad (16)$$

where f_i can be extracted by the previous equations.

III. NATURE OF THE SCALAR FIELDS AND REHEATING TEMPERATURE

In the previous section, we have discussed a modified postreheating scenario, where several component species in the form of noninteracting (decoupled) scalar fields, are added to the relativistic plasma. Even if we do not specify their nature, it should be underlined that these kind of components are quite common both in scalar modifications of general relativity and in theories with extra dimensions. In particular, in orientifold superstring models compactified to four dimensions and equipped with D-branes, the presence of additional scalars is an almost ubiquitous phenomenon [10]. Indeed, the D-brane action is the sum of a DBI term and a Wess-Zumino term, generalizations of the familiar mass and charge terms of a particle action. The dynamical fluctuations of the D-branes in the transverse directions correspond to d.o.f. that are described by scalar fields. Their coupling to the four-dimensional metric is induced on D-branes by the embedding inside the ten-dimensional space-time, and gives rise typically to a warp factor depending on the internal coordinates and to additional couplings entering the DBI action (disformal terms, see [9] and references therein). The most important point, however, is that these scalar fields always interact with the inflaton, that can thus decay into them and the remaining components of the standard reheating fluid after inflation. In this paper, we neglect the interactions of the scalar fields with matter longitudinal to the D-branes. From Eq. (15), we can easily argue that $\rho(T)$ increases with temperature, reaching the maximum value at $T = T_{\text{reh}}$. For instance, in the case of a single additional scalar field ϕ_1 , with a transition-to-radiation temperature T_1 , one has

$$\rho_{\phi_1}(T) = \rho_{\phi_1}(T_1) \left(\frac{g_S(T)}{g_S(T_1)} \right)^{\frac{4+n_1}{3}} \left(\frac{T}{T_1} \right)^{4+n_1}. \quad (17)$$

It is clear that there must be an upper bound to this energy density for $T = T_{\text{reh}}$. At this stage, whatever the nature of ϕ_1 , the energy density cannot assume arbitrary values, since it is at least limited by the presence of the Planck scale, M_{Pl} . In other words, we have to introduce a maximum scale M (with $M \leq M_{\text{Pl}}$) such that

$$\rho_{\phi_1}(T_{\text{reh}}) \leq M^4, \quad (18)$$

corresponding to an upper limit to the production scale of ϕ_1 . As a consequence, it turns out also to be an upper limit to the reheating temperature, once we set the scale M . Since T_1 is the transition-to-radiation temperature, i.e.,

$$\rho_{\phi_1}(T_1) = \rho_{\text{rad}}(T_1), \quad (19)$$

using

$$g_E(T) \sim g_S(T) \sim 100 \quad \text{for } T > T_{\text{QCD}}, \quad (20)$$

(where $T_{\text{QCD}} > 150$ MeV is the QCD phase transition scale), the reheating temperature must satisfy the condition

$$T_{\text{reh}} \leq \alpha_1 M \left(\frac{T_1}{M} \right)^{\frac{n_1}{4+n_1}}, \quad \alpha_1 = \left(\frac{30}{\pi^2 g_E} \right)^{\frac{1}{4+n_1}}, \quad (21)$$

with a resulting upper limit

$$T_{\text{reh}}^{\text{max}} = \alpha_1 M \left(\frac{T_1}{M} \right)^{\frac{n_1}{4+n_1}}. \quad (22)$$

In general, the scale M could be the Planck scale M_{Pl} but also, for instance, the GUT scale M_{GUT} or a lower scale of the order of the string scale, M_s , that is unconstrained in orientifolds [10, 11]. In particular, if the field ϕ_1 is supposed to be produced by the inflaton decay or during the reheating phase, we can also assume $M = M_{\text{inf}}$, where M_{inf} is the inflationary scale. It is interesting to analyze how the upper limit on T_{reh} varies with the scale M . In Fig. 1, we plot the behavior of $T_{\text{reh}}^{\text{max}}$ as a function of the model parameter $n = n_1$ for given values of the scale M . The transition-to-radiation temperature is chosen to be $T_1 \sim 10^4$ GeV. As expected, the maximum reheating temperature is larger for larger values of M , while it decreases with the model parameter n . The region below each curve representing $T_{\text{reh}}^{\text{max}}(n)$ describes the




FIG. 1. The maximum reheating temperature $T_{\text{reh}}^{\text{max}}$ as a function of the parameter n for $T_1 = 10^4$ GeV. The maximum reheating temperature becomes larger as M increases, while it decreases with n .




FIG. 2. The maximum reheating temperature $T_{\text{reh}}^{\text{max}}$ as a function of the parameter n for different values of the transition-to-radiation temperature and an inflationary scale $M_{\text{inf}} \sim 10^{15}$ GeV. The maximum reheating temperature becomes larger as the scale T_1 increases.

possible reheating temperatures compatible with the chosen bound M . For example, for $n = 2$ and $M = M_{\text{Pl}}$, we might have $T_{\text{reh}} \leq 10^{13}$ GeV, while for $M = M_{\text{inf}}$ we might only have $T_{\text{reh}} \leq 10^{11}$ GeV. For $n = 4$, a reheating temperature of the order of 10^9 GeV is compatible with $M = M_{\text{inf}}$ and *a fortiori* with Planckian or GUT bounds. In Fig. 2, we fix the scale M to the inflationary scale ($\sim 10^{15}$ GeV) and plot the behavior of $T_{\text{reh}}^{\text{max}}(n)$ for different values of the transition-to-radiation temperature. It happens that $T_{\text{reh}}^{\text{max}}(n)$ becomes smaller and smaller, for a fixed n , as the transition temperature decreases. For example, with $n = 2$ and $T_1 \sim 10^7$ GeV, we get $T_{\text{reh}} \leq 10^{12}$ GeV, while with $n = 4$ $T_{\text{reh}} \leq 10^{11}$. We postpone the discussion of the case with more scalar fields to Sec. V.

IV. INFLATIONARY e -FOLDINGS, REHEATING AND PRE-BBN SCALAR FIELDS

In the case of standard postreheating radiation dominated Universe, the inflationary number of e -foldings N_* has been calculated and used in many works [5]. In this section, we would like to discuss how this number changes in the presence of a nonstandard postinflationary scenario. As shown in [9], in the nonstandard case N_* acquires an additional e -folds term $\Delta N(\phi_i, T_{\text{reh}})$, that depends on the reheating temperature and on the features of the additional decoupled scalar fields discussed in Sec. II. Thus, we may write

$$N_* = \xi_* - \frac{1}{3(1+w_{\text{reh}})} \ln\left(\frac{\rho_{\text{end}}}{\rho_{\text{reh}}}\right) + \frac{1}{4} \ln\left(\frac{V_*^2}{M_{\text{pl}}^4 \rho_{\text{reh}}}\right) + \Delta N(\phi_i, T_{\text{reh}}), \quad (23)$$

where ρ_{end} is the energy density at the end of inflation, ρ_{reh} is the energy density when the reheating is completely realized, w_{reh} is the mean value of the EoS parameter of the reheating fluid, while $V_* = M_{\text{inf}}^4$ is the inflationary energy density. In Eq. (23),

$$\xi_* = -\ln\left(\frac{k_*}{a_0 H_0}\right) + \ln\left(\frac{T_0}{H_0}\right) + c, \quad (24)$$

with

$$c = -\frac{1}{12} \ln g_{\text{reh}} + \frac{1}{4} \ln\left(\frac{1}{9}\right) + \ln\left(\frac{43}{11}\right)^{\frac{1}{3}} \left(\frac{\pi^2}{30}\right)^{\frac{1}{4}}, \quad (25)$$

where k_* is the pivot scale for testing the cosmological parameters, a_0 and H_0 are the scale factor and the Hubble rate at the current epoch, respectively, T_0 is the CMB photon temperature while g_{reh} denotes the effective number of relativistic d.o.f. at the end of reheating [we are using $g_E(T_{\text{reh}}) = g_S(T_{\text{reh}}) = g_{\text{reh}}$ because of Eq. (20)]. Assuming $k_* = 0.002$ Mpc $^{-1}$, $H_0 = 1.75 \times 10^{-42}$ GeV, $T_0 = 2.3 \times 10^{-13}$ GeV and $g_{\text{reh}} \sim 100$, we get $\xi_* \sim 64$ and $c \sim 0.77$. The additional term comes out to be

$$\Delta N(\phi_i, T_{\text{reh}}) = \frac{1}{4} \ln \eta(n_i, T_i, T_{\text{reh}}), \quad (26)$$

where η is the ratio of the total energy density to the energy density of radiation at the reheating temperature,

$$\eta = 1 + \frac{\sum_i \rho \phi_i(T_{\text{reh}})}{\rho_{\text{rad}}(T_{\text{reh}})}. \quad (27)$$

Using Eq. (15) and expressing the radiation energy density in terms of T_1 ,

$$\rho_{\text{rad}}(T_{\text{reh}}) = \rho_{\text{rad}}(T_1) \frac{g_E(T_{\text{reh}})}{g_E(T_1)} \left(\frac{T_{\text{reh}}}{T_1}\right)^4, \quad (28)$$

we can write

$$\eta = 1 + \frac{g_E(T_1)}{g_E(T_{\text{reh}})} \left(\frac{T_1}{T_{\text{reh}}}\right)^4 \left\{ \left[\frac{T_{\text{reh}} g_S^{1/3}(T_{\text{reh}})}{T_1 g_S^{1/3}(T_1)} \right]^{4+n_1} + \sum_{i=2}^k \prod_{j=1}^{i-1} \left[\frac{T_{\text{reh}} g_S^{1/3}(T_{\text{reh}})}{T_i g_S^{1/3}(T_i)} \right]^{4+n_i} \left[\frac{T_{j+1} g_S^{1/3}(T_{j+1})}{T_j g_S^{1/3}(T_j)} \right]^{4+n_j} \right\}. \quad (29)$$

It should be noticed that the more scalar fields we have, the larger the parameter η is. Moreover, N_* is inflationary-model dependent due to the presence of the potential function in the second and third contributions of Eq. (23). However, by assuming $\rho_{\text{end}} \sim M_{\text{inf}}^4$, converting ρ_{reh} in T_{reh} and neglecting some small numerical factors, N_* can also be written as

$$N_* \sim \xi_* - \frac{1 - 3w_{\text{reh}}}{3(1 + w_{\text{reh}})} \ln\left(\frac{M_{\text{inf}}}{T_{\text{reh}}}\right) + \ln\left(\frac{M_{\text{inf}}}{M_{\text{Pl}}}\right) + \frac{1}{3(1 + w_{\text{reh}})} \ln \eta. \quad (30)$$

We can distinguish three main contributions. The first,

$$A(w_{\text{reh}}, T_{\text{reh}}) = \frac{1 - 3w_{\text{reh}}}{3(1 + w_{\text{reh}})} \ln \frac{M_{\text{inf}}}{T_{\text{reh}}}, \quad (31)$$

is entirely related to the reheating phase, the second involves the ratio between the Planck scale and the inflationary scale while the last one is due to the fraction of energy carried by the scalar fields, namely to the η factor. Let us provide a simple example considering a single scalar field postreheating dominance. By using Eq. (20), the general expression Eq. (29) turns out to be

$$\eta = 1 + \left(\frac{T_1}{T_{\text{reh}}}\right)^4 \left(\frac{T_{\text{reh}}}{T_1}\right)^{4+n_1} \simeq \left(\frac{T_{\text{reh}}}{T_1}\right)^{n_1}, \quad (32)$$

and therefore

$$\Delta N(\phi_1, T_{\text{reh}}) = \frac{n_1}{3(1 + w_{\text{reh}})} \ln\left(\frac{T_{\text{reh}}}{T_1}\right), \quad (33)$$

which, for the trivial $w_{\text{reh}} = 0$ case, results in

$$\Delta N(\phi_1, T_{\text{reh}}) = \frac{n_1}{3} \ln\left(\frac{T_{\text{reh}}}{T_1}\right). \quad (34)$$

The reheating and the η terms are strongly correlated. Indeed, in Sec. II, we have shown that the reheating temperature is constrained by an upper bound dependent on a scale M , by the transition-to-radiation temperature T_1 and by the dilution coefficient $n = n_1$. As a consequence, we have a lower bound on the reheating contribution in Eq. (31). Using the bound in Eq. (21), we get

$$A(w_{\text{reh}}, T_{\text{reh}}) \geq \frac{1 - 3w_{\text{reh}}}{3(1 + w_{\text{reh}})} \ln \frac{M_{\text{inf}}}{\alpha_1 M} \left(\frac{M}{T_1}\right)^{\frac{n_1}{4+n_1}}. \quad (35)$$

In Fig. 3, we report the quantity $\Delta N(\phi_1, T_{\text{reh}})$ as a function of the transition-to-radiation temperature for some values of n_1 , assuming $w_{\text{reh}} = 0$ and a reheating temperature $T_{\text{reh}} \sim 10^9$ GeV. It should be noticed that for $n = 4$ and $T_1 \sim 10^4$ GeV, we would obtain more than 15 extra e -folds, while for a larger $T_1 \sim 10^6$ GeV, we would have $\Delta N \sim 9$. In Fig. 4, we plot the complete result for the variable N_* as a function of the reheating equation of state parameter w_{reh} for $n = 1, 2, 3, 4$, assuming $T_1 \sim 10^4$ GeV. In general, the value of N_* increases with w_{reh} , as expected by the expression in Eq. (30). For $n = 2$ and $w_{\text{reh}} = 0$, we get $N_* \sim 59$, while $N_* \sim 67$ for $n = 4$ and $w_{\text{reh}} = 0$. In the




FIG. 3. Number of extra e -folds with $w_{\text{reh}} = 0$ and $T_{\text{reh}} \sim 10^9$ GeV. We have chosen this temperature because it is compatible with all values of n from 1 to 4 and with all the transition temperatures $T_1 > 10^4$ GeV, as seen in Sec. II.




FIG. 4. The inflationary number of e -folds N_* in a nonstandard postreheating cosmology as a function of w_{reh} . N_* increases with the value of the EoS parameter. The growth of N_* is also decreasing with n .

next section, we briefly discuss the multifield cases. The obtained results have nontrivial consequences on the theoretical predictions of the underlying inflationary models. The reason is that one usually infers the values of the two main inflationary parameters, the scalar spectral index n_s and the tensor-to-scalar ratio r , assuming an N_* in the range between 50 and 60. Therefore, if we considered a different N_* we could have new predictions to compare with the current experimental bounds [12]. In the Appendices, we will briefly examine how the nontrivial values of N_* affect some paradigmatic inflationary models.

V. SUMMARY AND DISCUSSION

Inflation should have taken place at very high-energy scales. The accelerated expansion was followed by a

“reheat” stage that produced the standard model radiation fluid and the observable large comoving entropy of the Universe. However, available data do not guarantee that the mentioned scenario is the correct one. For instance, a viable alternative is to have one (or more) additional field(s) that dominates the energy budget of the Universe at different phases after the reheating epoch. In particular, many authors have recently considered the inclusion of new scalar fields (quintessence, scalar field decoupled from matter and radiation or even scalar fields or moduli coupled to gravity) to approach some problems related to dark matter relics abundances or to the number of inflationary e -folds (see [6–9] and references therein). The relation between scalar decaying particle and black hole formation in GUT cosmology was also studied in the past [13]. In general, nonstandard cosmological histories before the BBN are interesting possibilities whose signatures could be tested in the near future, for instance by gravity-waves experiments (see [14] for details).

In this paper, extending the approach of [9], we have described a postreheating era dominated by a collection of simple scalar fields ϕ_i , ($i = 1, \dots, k$) completely decoupled from standard model matter and radiation. Their presence is described in terms of perfect fluids with energy densities that scale as $\rho_i \sim a^{-(4+n_i)}$, $n_i > 0$. Each ϕ_i dominates at different times. In particular, ϕ_1 is the field connecting the nonstandard part of the postreheating phase to the radiation-dominated era. In this scenario, it is mandatory to assume that the transition to radiation occurs well before the BBN, in order not to ruin the theoretical predictions about light element abundances [6]. In Eqs. (3) and (15), the general expressions related to the total energy density and to the energy density of a single field ϕ_i have been derived, with the proviso of absence of entropy variation. The changes in the Hubble rate during the multifield driven evolution are regulated by Eqs. (15) and (16). In Sec. III, we observed that the energy density after reheating must be at most Planckian. As a consequence, there exists an upper limit to the reheating temperature, as shown in Eq. (22) and illustrated in Fig. 1 for different choices of the limiting scale and a transition temperature to HBB $\sim 10^4$ GeV. The upper bound depends on T_1 and also on the indices n_i , as shown in Fig. 2.

Let us take a closer look to the multifield case, already mentioned in Sec. III. Of course, the upper bound on T_{reh} is always present, but it depends on the intermediate temperatures T_i . For instance, in the presence of two scalar fields, with ϕ_2 dominating at higher temperature $T > T_2$ on ϕ_1 , the condition becomes $\rho_{\phi_2} \leq M^4$ at $T = T_{\text{reh}}$. As a consequence, one gets

$$T_{\text{reh}} < \alpha_2 M \left(\frac{T_1^{n_1} T_2^{n_2 - n_1}}{M^{n_2}} \right)^{\frac{1}{4+n_2}} \quad (36)$$

where $\alpha_2 = (30/\pi^2 g_E)^{1/4+n_2}$ (by assumption $n_2 > n_1$). Using, for instance, $n_1 = 1$, $n_2 = 2$, $T_1 \sim 10^4$ GeV and $T_2 \sim 10^6$ GeV one has $T_{\text{reh}} < 10^{12}$ GeV for $M < M_{\text{inf}}$, while $T_{\text{reh}} < 10^{13.7}$ GeV for $M < M_{\text{Pl}}$. Note that in the first case the value of $T_{\text{reh}}^{\text{max}}$ is very close to the one found in the presence of a single scalar field with $n_1 = 2$ at a transition-to-radiation temperature $T_1 \sim 10^7$ GeV (see Sec. II). The upper bound can obviously be computed for any number k of scalar fields and it turns out to depend on $2k$ parameters, the T_i temperatures and the n_i dilution coefficients.

A nonstandard cosmological epoch after reheating gives also rise to an extra term in the general expression of the inflationary number of e -foldings, N_* [see Eqs. (23) and (30)]. The upper bound on the energy density of the k -th scalar field leads to an additional constraint on the contribution to N_* coming from the reheating phase, as shown in Eq. (35). As a result, we found the possibility of having an inflationary number of e -foldings well beyond 60, as shown in Fig. 4. The higher is the number of scalar fields, the larger is the correction ΔN to N_* , since the ratio of the total energy density to the radiation density at T_{reh} is larger. For instance, with two scalar fields and using Eq. (20), Eq. (29) provides

$$\eta \simeq 1 + \left(\frac{T_1}{T_{\text{reh}}} \right)^4 \left(\frac{T_{\text{reh}}}{T_1} \right)^{4+n_1} + \left(\frac{T_1}{T_{\text{reh}}} \right)^4 \left(\frac{T_2}{T_1} \right)^{4+n_1} \left(\frac{T_{\text{reh}}}{T_2} \right)^{4+n_2}. \quad (37)$$

By choosing the same data as after Eq. (36) and $w_{\text{reh}} = 0$, $T_{\text{reh}} \sim 10^{13}$ GeV, one gets $\eta \sim 10^{16}$, $\Delta N(\phi_1, \phi_2) \sim 12$ and $N_* \sim 70$. As expected, a nonstandard postreheating phase produces a variety of enhancements in the inflationary number of e -foldings, depending on the number of additional scalar fields and on the details of their dilution properties. Enhancements affect the theoretical predictions of the inflationary models, mainly in the bottom right portion of the familiar (n_s, r) plane. In [9], Maharana and Zavala have studied the functions $n_s(N_*)$ and $r(N_*)$. In Appendices A and B, we report some results for typical classes of inflationary models, extending the range of parameters provided in [9]. We deserve an extended analysis to a future publication [15].

ACKNOWLEDGMENTS

This work was supported in part by the MIUR-PRIN Contract No. 2015MP2CX4 “Non-perturbative Aspects Of Gauge Theories And String.” P. C. would also like to thank the company “L’isola che non c’è S.r.l.” for its support.

APPENDIX A: MONOMIAL POTENTIALS

The first class of inflationary models we are going to analyze are those characterized by single monomial potentials of the form

$$V(\varphi) = \lambda_p \varphi^p, \quad \lambda_p = M_{\text{inf}}^4 M_{\text{Pl}}^{-p}. \quad (\text{A1})$$

In this class of models, the inflaton field, in order to drive inflation, must exhibit a super-Planckian variation $\Delta\phi > M_{\text{Pl}}$. Historically, the most known scenarios are the ones with $p = 2$ and $p = 4$, that were introduced by Linde [3]. Monomial potentials naturally occur also in superstring compactifications, where they are called ‘‘axion monodromy’’ [16]. In these models, one uses a cover of the compactification manifold, with branes wrapping suitable internal cycles. As a result, even though the manifold is compact, the wrapping of branes around certain cycles weakly breaks the original shift symmetry, allowing for closed-string axions with super-Planckian excursions and the suppression of dangerous higher-dimensional operators. The involved inflationary potentials come out precisely of the form $V(\varphi) \sim \varphi^p$ with $p = 2/5, 2/3, 1$ or $4/3$. The slow-roll parameters give rise to standard theoretical predictions for the spectral index and the tensor-to-scalar ratio in terms of the inflationary number of e -foldings:

TABLE I. Inflationary predictions for monomial potentials in nonstandard postreheating cosmology. We assume a single scalar field with $T_1 \sim 10^4$ GeV, $n_1 = 4$ and $T_{\text{reh}} \sim 10^9$ GeV, giving $N_* = 67$.

Model parameter p	$n_s(N_*)$	$r(N_*)$
Axion model $p = 2/5$	0.9821	0.0238
Axion model $p = 2/3$	0.9801	0.0398
Axion model $p = 1$	0.9776	0.0597
Axion model $p = 4/3$	0.9751	0.0796
Linde model $p = 2$	0.9701	0.1194
Linde model $p = 4$	0.9552	0.2388

TABLE II. Inflationary predictions for monomial potentials in nonstandard postreheating cosmology. We assume two scalar fields with $T_1 \sim 10^4$ GeV, $n_1 = 1$, $T_2 \sim 10^6$ GeV, $n_2 = 2$ and $T_{\text{reh}} \sim 10^{13}$ GeV, giving $N_* = 70$.

Model parameter p	$n_s(N_*)$	$r(N_*)$
Axion model $p = 2/5$	0.9830	0.0229
Axion model $p = 2/3$	0.9810	0.0381
Axion model $p = 1$	0.9785	0.0571
Axion model $p = 4/3$	0.9762	0.0762
Linde model $p = 2$	0.9714	0.1143
Linde model $p = 4$	0.9571	0.2286

$$n_s \sim 1 - \frac{p+2}{2N_*}, \quad r = \frac{4p}{N_*}. \quad (\text{A2})$$

It should be noticed that both n_s and r depend on the model parameter p . In Tables I and II, we report the theoretical predictions for some scenarios related to monomial potentials, assuming two possible nonstandard postreheating data.

APPENDIX B: EXPONENTIAL POTENTIALS

The second class of models we would like to consider is that of exponential potentials of the form

$$V(\varphi) \sim M_{\text{inf}}^4 (1 - e^{-b\varphi}), \quad b = \sqrt{\frac{2}{3\alpha}}, \quad (\text{B1})$$

where α is a free parameter. These potentials arise in many contexts. Important examples are the well-known Starobinsky model ($\alpha = 1$), the Goncharov-Linde model ($\alpha = 1/9$) and the Higgs Inflation model ($\alpha = \sqrt{2/3}$) [17]. More recently, the so-called α -attractor models of inflation [18] have also been considered, that fall in the same class of Eq. (B1). Furthermore, other very interesting examples come out in superstring-inspired scenarios, like Kähler moduli inflation, poly-instanton inflation, and fiber inflation [19]. At first order, the theoretical predictions of this class of models result

$$n_s \sim 1 - \frac{2}{N_*}, \quad r \sim \frac{12\alpha}{N_*^2}. \quad (\text{B2})$$

In this case, the scalar spectral index does not depend on the value of α . Therefore, for $N_* = 67$, one has $n_s \sim 0.9701$, while for $N_* = 70$, $n_s = 0.9714$, independently on α . On the contrary, the tensor-to-scalar ratio depends on α as shown in Table III, where we report its values for some choices of the parameters.

TABLE III. The tensor-to-scalar ratio for some exponential potential models depending on α . In the first column, we assume $N_1 = 67$ (related to the case with a single scalar field and $T_1 \sim 10^4$ GeV, $n_1 = 4$, $T_{\text{reh}} \sim 10^9$ GeV). In the second column, we assume $N_2 = 70$ (related to a pair of scalar fields characterized by $T_1 \sim 10^4$ GeV, $n_1 = 1$, $T_2 \sim 10^6$ GeV, $n_2 = 2$ and $T_{\text{reh}} \sim 10^{13}$ GeV.)

Model parameter α	$r(N_1)$	$r(N_2)$
Starobinsky $\alpha = 1$	2.7×10^{-3}	2.4×10^{-3}
Fiber inflation $\alpha = 2$	5.3×10^{-3}	4.9×10^{-3}
Goncharov-Linde $\alpha = 1/9$	2.9×10^{-4}	2.7×10^{-4}
Poly-instanton $\alpha = 3 \times 10^{-3}$	8.0×10^{-6}	7.3×10^{-6}
Kähler moduli $\alpha = 3 \times 10^{-8}$	8.0×10^{-11}	7.3×10^{-11}

- [1] A. D. Linde, Particle physics and inflationary cosmology, *Contemp. Concepts Phys.* **5**, 1 (1990); D. Bailin and A. Love, *Cosmology in Gauge Field Theory and String Theory* (IOP, Bristol, 2004), p. 313; V. Mukhanov, *Physical Foundations of Cosmology* (Cambridge University Press, Cambridge, England, 2005), p. 421; V. A. Rubakov and D. S. Gorbunov, *Introduction to the Theory of the Early Universe: Hot Big Bang Theory* (World Scientific, Singapore, 2011); D. S. Gorbunov and V. A. Rubakov, *Introduction to the Theory of the Early Universe: Cosmological Perturbations and Inflationary Theory* (World Scientific, Hackensack, 2011), p. 489; S. Weinberg, *Cosmology* (Oxford University Press, New York, 2008), p. 593.
- [2] H. Georgi and S. L. Glashow, Unity of All Elementary Particle Forces, *Phys. Rev. Lett.* **32**, 438 (1974); P. Langacker, Grand Unified Theories and Proton Decay, *Phys. Rep.* **72**, 185 (1981); W. de Boer, Grand unified theories and supersymmetry in particle physics and cosmology, *Prog. Part. Nucl. Phys.* **33**, 201 (1994).
- [3] A. H. Guth, The inflationary universe: A possible solution to the horizon and flatness problems, *Phys. Rev. D* **23**, 347 (1981); A. D. Linde, A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems, *Phys. Lett.* **108B**, 389 (1982); A. Albrecht and P. J. Steinhardt, Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking, *Phys. Rev. Lett.* **48**, 1220 (1982); S. W. Hawking and I. G. Moss, Supercooled phase transitions in the very early universe, *Phys. Lett.* **110B**, 35 (1982); A. D. Linde, Chaotic inflation, *Phys. Lett.* **129B**, 177 (1983); A. Vilenkin, Creation of universes from nothing, *Phys. Lett.* **117B**, 25 (1982); The birth of inflationary universes, *Phys. Rev. D* **27**, 2848 (1983); Quantum creation of universes, *Phys. Rev. D* **30**, 509 (1984).
- [4] M. S. Turner, Coherent scalar field oscillations in an expanding universe, *Phys. Rev. D* **28**, 1243 (1983); A. Albrecht, P. J. Steinhardt, M. S. Turner, and F. Wilczek, Reheating an Inflationary Universe, *Phys. Rev. Lett.* **48**, 1437 (1982); A. D. Dolgov and A. D. Linde, Baryon asymmetry in inflationary universe, *Phys. Lett.* **116B**, 329 (1982); L. F. Abbott, E. Farhi, and M. B. Wise, Particle production in the new inflationary cosmology, *Phys. Lett.* **117B**, 29 (1982); E. W. Kolb and M. S. Turner, Grand unified theories and the origin of the baryon asymmetry, *Annu. Rev. Nucl. Part. Sci.* **33**, 645 (1983); M. Morikawa and M. Sasaki, Entropy production in the inflationary universe, *Prog. Theor. Phys.* **72**, 782 (1984); Entropy production in an expanding universe, *Phys. Lett.* **165B**, 59 (1985); A. D. Dolgov and D. P. Kirilova, On particle creation by a time dependent scalar field, *Yad. Fiz.* **51**, 273 (1990) [*Sov. J. Nucl. Phys.* **51**, 172 (1990)]; J. H. Traschen and R. H. Brandenberger, Particle production during out-of-equilibrium phase transitions, *Phys. Rev. D* **42**, 2491 (1990); L. Kofman, A. D. Linde, and A. A. Starobinsky, Reheating After Inflation, *Phys. Rev. Lett.* **73**, 3195 (1994); Y. Shtanov, J. H. Traschen, and R. H. Brandenberger, Universe reheating after inflation, *Phys. Rev. D* **51**, 5438 (1995); L. Kofman, A. D. Linde, and A. A. Starobinsky, Nonthermal Phase Transitions After Inflation, *Phys. Rev. Lett.* **76**, 1011 (1996); S. Y. Khlebnikov and I. I. Tkachev, Classical Decay of Inflaton, *Phys. Rev. Lett.* **77**, 219 (1996); I. I. Tkachev, Phase transitions at preheating, *Phys. Lett. B* **376**, 35 (1996); L. Kofman, A. D. Linde, and A. A. Starobinsky, Towards the theory of reheating after inflation, *Phys. Rev. D* **56**, 3258 (1997); S. Khlebnikov, L. Kofman, A. D. Linde, and I. Tkachev, First Order Nonthermal Phase Transition After Preheating, *Phys. Rev. Lett.* **81**, 2012 (1998); G. N. Felder, L. Kofman, and A. D. Linde, Instant preheating, *Phys. Rev. D* **59**, 123523 (1999); P. B. Greene and L. Kofman, Preheating of fermions, *Phys. Lett. B* **448**, 6 (1999); R. Allahverdi, R. Brandenberger, F. Y. Cyr-Racine, and A. Mazumdar, Reheating in inflationary cosmology: Theory and applications, *Annu. Rev. Nucl. Part. Sci.* **60**, 27 (2010).
- [5] J. Martin and C. Ringeval, First CMB constraints on the inflationary reheating temperature, *Phys. Rev. D* **82**, 023511 (2010); J. B. Munoz and M. Kamionkowski, Equation-of-state parameter for reheating, *Phys. Rev. D* **91**, 043521 (2015); J. L. Cook, E. Dimastrogiovanni, D. A. Easson, and L. M. Krauss, Reheating predictions in single field inflation, *J. Cosmol. Astropart. Phys.* **04** (2015) 047; V. Domcke and J. Heisig, Constraints on the reheating temperature from sizable tensor modes, *Phys. Rev. D* **92**, 103515 (2015); K. D. Lozanov and M. A. Amin, The Equation of State and Duration to Radiation Domination after Inflation, *Phys. Rev. Lett.* **119**, 061301 (2017); J. Martin, C. Ringeval, and V. Vennin, Observing Inflationary Reheating, *Phys. Rev. Lett.* **114**, 081303 (2015); Information gain on reheating: The one bit milestone, *Phys. Rev. D* **93**, 103532 (2016); R. J. Hardwick, V. Vennin, K. Koyama, and D. Wands, Constraining curvaton reheating, *J. Cosmol. Astropart. Phys.* **08** (2016) 042; M. Eshaghi, M. Zarei, N. Riazi, and A. Kiasatpour, CMB and reheating constraints to α -attractor inflationary models, *Phys. Rev. D* **93**, 123517 (2016); Y. Ueno and K. Yamamoto, Constraints on α -attractor inflation and reheating, *Phys. Rev. D* **93**, 083524 (2016); A. Di Marco, P. Cabella, and N. Vittorio, Constraining the general reheating phase in the α -attractor inflationary cosmology, *Phys. Rev. D* **95**, 103502 (2017); N. Rashidi and K. Nozari, α -attractor and reheating in a model with non-canonical scalar fields, *Int. J. Mod. Phys. D* **27**, 1850076 (2018); P. Cabella, A. Di Marco, and G. Pradisi, Fibre inflation and reheating, *Phys. Rev. D* **95**, 123528 (2017); M. Drewes, J. U. Kang, and U. R. Mun, CMB constraints on the inflaton couplings and reheating temperature in α -attractor inflation, *J. High Energy Phys.* **11** (2017) 072; S. Bhattacharya, K. Dutta, and A. Maharana, Constrains on Khler moduli inflation from reheating, *Phys. Rev. D* **96**, 083522 (2017); *Phys. Rev. D* **96**, 109901(A) (2017).
- [6] F. D’Eramo, N. Fernandez, and S. Profumo, When the universe expands too fast: Relentless dark matter, *J. Cosmol. Astropart. Phys.* **05** (2017) 012; Some authors studied the possibilities that massive particles can also decay at very low temperature altering the abundances of the light elements. Some very useful references are M. Kawasaki, K. Kohri, and N. Sugiyama, Cosmological Constraints on Late Time Entropy Production, *Phys. Rev. Lett.* **82**, 4168 (1999); MeV scale reheating temperature and thermalization of neutrino background, *Phys. Rev. D* **62**, 023506 (2000).

- [7] P. Salati, Quintessence and the relic density of neutralinos, *Phys. Lett. B* **571**, 121 (2003); R. Catena, N. Fornengo, A. Masiero, M. Pietroni, and F. Rosati, Dark matter relic abundance and scalar-tensor dark energy, *Phys. Rev. D* **70**, 063519 (2004); A. B. Lahanas, N. E. Mavromatos, and D. V. Nanopoulos, Dilaton and off-shell (non-critical string) effects in Boltzmann equation for species abundances, *PMC Phys. A* **1**, 2 (2007); A. Arbey and F. Mahmoudi, SUSY constraints from relic density: High sensitivity to pre-BBN expansion rate, *Phys. Lett. B* **669**, 46 (2008); R. Catena, N. Fornengo, M. Pato, L. Pieri, and A. Masiero, Thermal Relics in modified cosmologies: Bounds on evolution histories of the early universe and cosmological boosts for PAMELA, *Phys. Rev. D* **81**, 123522 (2010); C. Pallis, Cold dark matter in non-standard cosmologies, PAMELA, ATIC and Fermi LAT, *Nucl. Phys.* **B831**, 217 (2010); G. B. Gelmini, J. H. Huh, and T. Rehagen, Asymmetric dark matter annihilation as a test of non-standard cosmologies, *J. Cosmol. Astropart. Phys.* **08** (2013) 003; T. Rehagen and G. B. Gelmini, Effects of kination and scalar-tensor cosmologies on sterile neutrinos, *J. Cosmol. Astropart. Phys.* **06** (2014) 044; H. Iminiyaz and X. Chen, Relic abundance of asymmetric dark matter in quintessence, *Astropart. Phys.* **54**, 125 (2014); M. T. Meehan and I. B. Whittingham, Asymmetric dark matter in braneworld cosmology, *J. Cosmol. Astropart. Phys.* **06** (2014) 018; Dark matter relic density in Gauss-Bonnet braneworld cosmology, *J. Cosmol. Astropart. Phys.* **12** (2014) 034; Dark matter relic density in scalar-tensor gravity revisited, *J. Cosmol. Astropart. Phys.* **12** (2015) 011; S. z. Wang, H. Iminiyaz, and M. Mamat, Asymmetric dark matter and the scalar-tensor model, *Int. J. Mod. Phys. A* **31**, 1650021 (2016); B. Dutta, E. Jimenez, and I. Zavala, Dark matter Relics and the expansion rate in scalar-tensor theories, *J. Cosmol. Astropart. Phys.* **06** (2017) 032; D-brane disformal coupling and thermal dark matter, *Phys. Rev. D* **96**, 103506 (2017); T. Koivisto, D. Wills, and I. Zavala, Dark D-brane cosmology, *J. Cosmol. Astropart. Phys.* **06** (2014) 036; L. Aparicio, M. Cicoli, B. Dutta, F. Muia, and F. Quevedo, Light higgsino dark matter from non-thermal cosmology, *J. High Energy Phys.* **11** (2016) 038; E. Hardy, Higgs portal dark matter in non-standard cosmological histories, *J. High Energy Phys.* **06** (2018) 043; N. Bernal, C. Cosme, and T. Tenkanen, Phenomenology of Self-Interacting Dark Matter in a Matter-Dominated Universe, [arXiv:1803.08064](https://arxiv.org/abs/1803.08064); N. Bernal, C. Cosme, T. Tenkanen, and V. Vaskonen, Scalar singlet dark matter in non-standard cosmologies, [arXiv:1806.11122](https://arxiv.org/abs/1806.11122).
- [8] K. Dutta and A. Maharana, Inflationary constraints on modulus dominated cosmology, *Phys. Rev. D* **91**, 043503 (2015); K. Das, K. Dutta, and A. Maharana, Inflationary predictions and Moduli masses, *Phys. Lett. B* **751**, 195 (2015); M. Cicoli, K. Dutta, A. Maharana, and F. Quevedo, Moduli vacuum misalignment and precise predictions in string inflation, *J. Cosmol. Astropart. Phys.* **08** (2016) 006; Note that a change in the number of e -folds could also be possible in quintessential α -attractor inflation, where the post inflationary phase is dominated by the kinetic energy of the inflaton field itself, as recently shown in Y. Akrami, R. Kallosh, A. Linde, and V. Vardanyan, Dark energy, α -attractors, and large-scale structure surveys, *J. Cosmol. Astropart. Phys.* **06** (2018) 041.
- [9] A. Maharana and I. Zavala, Post-inflationary scalar tensor cosmology and inflationary parameters, *Phys. Rev. D* **97**, 123518 (2018).
- [10] For reviews see: C. Angelantonj and A. Sagnotti, Open strings, *Phys. Rep.* **371**, 1 (2002); Erratum, *Phys. Rep.* **376**, 407(E) (2003); R. Blumenhagen, B. Kors, D. Lust, and S. Stieberger, Four-dimensional string compactifications with D-branes, orientifolds and fluxes, *Phys. Rep.* **445**, 1 (2007); E. Dudas, Theory and phenomenology of type I strings and M theory, *Classical Quantum Gravity* **17**, R41 (2000).
- [11] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, New dimensions at a millimeter to a Fermi and superstrings at a TeV, *Phys. Lett. B* **436**, 257 (1998); I. Antoniadis, E. Dudas, and A. Sagnotti, Supersymmetry breaking, open strings and M theory, *Nucl. Phys.* **B544**, 469 (1999); K. R. Dienes, E. Dudas, and T. Gherghetta, TeV scale GUTs, [arXiv:hep-ph/9807522](https://arxiv.org/abs/hep-ph/9807522); C. P. Bachas, Unification with low string scale, *J. High Energy Phys.* **11** (1998) 023.
- [12] P. A. R. Ade *et al.* (Planck Collaboration), Planck 2015 results. XIII. Cosmological parameters, *Astron. Astrophys.* **594**, A13 (2016); Planck 2015 results. XX. Constraints on inflation, *Astron. Astrophys.* **594**, A20 (2016); P. A. R. Ade *et al.* (BICEP2 and Keck Array Collaborations), Improved Constraints on Cosmology and Foregrounds from BICEP2 and Keck Array Cosmic Microwave Background Data with Inclusion of 95 GHz Band, *Phys. Rev. Lett.* **116**, 031302 (2016).
- [13] M. Y. Khlopov and A. G. Polnarev, Primordial black holes as a cosmological test of grand unification, *Phys. Lett.* **97B**, 383 (1980); A. G. Polnarev and M. Yu. Khlopov, Primordial black holes and the ERA of superheavy particle dominance in the early universe, *Astron. Zh.* **58**, 706 (1981) *Sov. Astron.* **25**, 406 (1981); The ERA of superheavy particle dominance and big bang nucleosynthesis, *Astron. Zh.* **59**, 15 (1982) [*Sov. Astron.* **26**, 9 (1982)]; Dust-like stages in the early universe and constraints on the primordial black-hole spectrum, *Astron. Zh.* **59**, 639 (1982) [*Sov. Astron.* **26**, 391 (1982)]; Grand unification cosmology and the parameters of a neutrino dominated universe, *Pis'ma Astron. Zh.* **9**, 323 (1983) [*Sov. Astron. Lett.* **9**, 171 (1983)]; A. G. Polnarev and M. Yu. Khlopov, Cosmology, primordial black holes, and supermassive particles, *Usp. Fiz. Nauk* **145**, 369 (1985) [*Sov. Phys. Usp.* **28**, 213 (1985)]; M. Khlopov, B. A. Malomed, and I. B. Zeldovich, Gravitational instability of scalar fields and formation of primordial black holes, *Mon. Not. R. Astron. Soc.* **215**, 575 (1985); M. Yu. Khlopov and V. M. Chechetkin, Antiprotons in the Universe as a cosmological test of grand unification, *Fiz. Elem. Chastits At. Yadra* **18**, 627 (1987) [*Sov. J. Part. Nucl.* **18**, 267 (1987)].
- [14] Y. Cui, M. Lewicki, D. E. Morrissey, and J. D. Wells, Cosmic archaeology with gravitational waves from cosmic strings, *Phys. Rev. D* **97**, 123505 (2018); G. Barenboim and W. I. Park, Gravitational waves from first order phase transitions as a probe of an early matter domination era and its inverse problem, *Phys. Lett. B* **759**, 430 (2016); M. Artymowski, O. Czerwinska, Z. Lalak, and M. Lewicki, Gravitational wave signals and cosmological consequences of gravitational reheating, *J. Cosmol. Astropart. Phys.* **04**

- (2018) 046; D. J. H. Chung and P. Zhou, Gravity waves as a probe of Hubble expansion rate during an electroweak scale phase transition, *Phys. Rev. D* **82**, 024027 (2010).
- [15] P. Cabella, A. Di Marco, and G. Pradisi (to be published).
- [16] E. Silverstein and A. Westphal, Monodromy in the CMB: Gravity waves and string inflation, *Phys. Rev. D* **78**, 106003 (2008); L. McAllister, E. Silverstein, and A. Westphal, Gravity waves and linear inflation from axion monodromy, *Phys. Rev. D* **82**, 046003 (2010); L. McAllister, E. Silverstein, A. Westphal, and T. Wrase, The powers of monodromy, *J. High Energy Phys.* **09** (2014) 123.
- [17] A. A. Starobinsky, A new type of isotropic cosmological models without singularity, *Phys. Lett.* **91B**, 99 (1980); B. Whitt, Fourth order gravity as general relativity plus matter, *Phys. Lett.* **145B**, 176 (1984); A. Vilenkin, Classical and quantum cosmology of the Starobinsky inflationary model, *Phys. Rev. D* **32**, 2511 (1985); L. A. Kofman, A. D. Linde, and A. A. Starobinsky, Inflationary universe generated by the combined action of a scalar field and gravitational vacuum polarization, *Phys. Lett.* **157B**, 361 (1985); M. B. Mijic, M. S. Morris, and W. M. Suen, The R^2 cosmology: Inflation without a phase transition, *Phys. Rev. D* **34**, 2934 (1986); A. S. Goncharov and A. D. Linde, Chaotic inflation of the universe in supergravity, *Zh. Eksp. Teor. Fiz.* **86**, 1594 (1984) [*Sov. Phys. JETP* **59**, 930 (1984)]; Chaotic inflation in supergravity, *Phys. Lett.* **139B**, 27 (1984); F. L. Bezrukov and M. Shaposhnikov, The standard model Higgs boson as the inflaton, *Phys. Lett. B* **659**, 703 (2008).
- [18] R. Kallosh and A. Linde, Universality class in conformal inflation, *J. Cosmol. Astropart. Phys.* **07** (2013) 002; Multi-field conformal cosmological attractors, *J. Cosmol. Astropart. Phys.* **12** (2013) 006; S. Ferrara, R. Kallosh, A. Linde, and M. Porrati, Minimal supergravity models of inflation, *Phys. Rev. D* **88**, 085038 (2013); R. Kallosh and A. Linde, Superconformal generalizations of the Starobinsky model, *J. Cosmol. Astropart. Phys.* **06** (2013) 028; R. Kallosh, A. Linde, and D. Roest, Superconformal inflationary α -attractors, *J. High Energy Phys.* **11** (2013) 198; M. Galante, R. Kallosh, A. Linde, and D. Roest, Unity of Cosmological Inflation Attractors, *Phys. Rev. Lett.* **114**, 141302 (2015); R. Kallosh and A. Linde, Cosmological attractors and asymptotic freedom of the inflaton field, *J. Cosmol. Astropart. Phys.* **06** (2016) 047; See also P. Carrilho, D. Mulryne, J. Ronayne, and T. Tenkanen, Attractor behaviour in multifield inflation, *J. Cosmol. Astropart. Phys.* **06** (2018) 032; L. Järv, A. Racioppi, and T. Tenkanen, Palatini side of inflationary attractors, *Phys. Rev. D* **97**, 083513 (2018) for more recent developments.
- [19] J. P. Conlon and F. Quevedo, Kahler moduli inflation, *J. High Energy Phys.* **01** (2006) 146; A. Maharana, M. Rummel, and Y. Sumitomo, Accidental Kähler moduli inflation, *J. Cosmol. Astropart. Phys.* **09** (2015) 040; M. Cicoli, F. G. Pedro, and G. Tasinato, Poly-instanton Inflation, *J. Cosmol. Astropart. Phys.* **12** (2011) 022; M. Cicoli, C. P. Burgess, and F. Quevedo, Fibre inflation: Observable gravity waves from IIB string compactifications, *J. Cosmol. Astropart. Phys.* **03** (2009) 013; C. P. Burgess, M. Cicoli, S. de Alwis, and F. Quevedo, Robust inflation from fibrous strings, *J. Cosmol. Astropart. Phys.* **05** (2016) 032; M. Cicoli, F. Muia, and P. Shukla, Global embedding of fibre inflation models, *J. High Energy Phys.* **11** (2016) 182; B. J. Broy, D. Ciupke, F. G. Pedro, and A. Westphal, Starobinsky-type inflation from α' -corrections, *J. Cosmol. Astropart. Phys.* **01** (2016) 001; M. Cicoli, D. Ciupke, S. de Alwis, and F. Muia, α' inflation: Moduli stabilisation and observable tensors from higher derivatives, *J. High Energy Phys.* **09** (2016) 026.