

Dirac equation in exotic spacetimes

Javier Faba García

*Departamento de Física Teórica, Universidad Complutense de Madrid, Plaza de Ciencias,
1 28040 Madrid, Spain*

Carlos Sabín

Instituto de Física Fundamental, CSIC, Serrano, 113-bis, 28006 Madrid, Spain

 (Received 13 November 2018; published 14 January 2019)

We find solutions of the Dirac equation in curved spacetime. In particular, we consider 1 + 1 dimensional sections of several exotic metrics: the Alcubierre metric, which describes a scenario that allows faster-than-light velocity; the Gödel metric, which describes a universe containing closed timelike curves; and the Kerr metric, which corresponds to the metric of a rotating pointlike source, for instance a rotating black hole. Moreover, we also show that the techniques that we use in these cases can be extended to nonstatic metrics.

DOI: [10.1103/PhysRevD.99.025008](https://doi.org/10.1103/PhysRevD.99.025008)

I. INTRODUCTION

In 1928, Paul Dirac formulated a special relativistic version of the wave equation in quantum mechanics ($\hbar = 1$, $c = 1$):

$$i\gamma^\mu \partial_\mu \psi - m\psi = 0,$$

where, in the standard representation,

$$\gamma^0 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad \gamma^1 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$$

$$\gamma^2 = \begin{pmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix} \quad \gamma^3 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

This equation was born as an attempt of linearization of the Klein-Gordon equation, which comes from the application of the Einstein relation $E^2 = p^2 + m^2$ to the Schrödinger equation Hamiltonian. The Dirac equation was a turning point in physics. Joining quantum mechanics and relativity, this equation predicts the spin of the electron, the existence of antimatter etc., and it is considered as the natural transition between relativistic quantum mechanics and quantum field theory [1]. Moreover, a renewed interest in the Dirac equation has come from the field of quantum simulations, which has enabled the experimental observation of some of its more interesting features in tabletop

experiments [2–8]. Nevertheless, even though the general solutions of this equation are well known, this is not the case of the Dirac equation in curved spacetime. In general, obtaining solutions in curved space is more complicated. For a detailed analysis of the 3 + 1 Dirac equation in Riemannian spacetimes, see for instance [9]. Some particular 1 + 1 D cases can be found, for instance, in [10,11].

In this work, the method explained by one of us in [11] will be applied to obtain solutions of the Dirac equation in curved spacetime for different exotic metrics in 1 + 1 dimensional sections, with the help of known solutions in Minkowski spacetime. If we take a 1 + 1 dimensional section of the spacetime, the Dirac equations reduces to [10]

$$i\left(\partial_t + \frac{\dot{\Omega}}{2\Omega}\right)\psi = -i\sigma_x\left(\partial_x + \frac{\Omega'}{2\Omega}\right)\psi + \sigma_z\Omega m\psi,$$

where Ω is the conformal factor—note that any metric in 1 + 1 dimensions allow a coordinate change such that it acquires the form $ds^2 = \Omega^2(-dt^2 + dx^2)$ —the dot and prime stand for time and spatial derivatives, respectively, and σ_x and σ_z are the Pauli matrices:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Note that if $\Omega = 1$, we obtain the well-known Dirac equation in 1 + 1 dimensional flat spacetime (see, for instance [12]). This equation has been studied in detail, including striking properties such as Zitterbewegung or Klein paradox [13,14]. Given Ω , we could try to find solutions of this partial differential equation system. However, that procedure could be very involved, depending

on the particular form of Ω . The method presented in [11] allows to find the solution of this partial differential equation system through a coordinate change and an existing solution in Minkowski spacetime. Thus, the method will be applied to several exotic metrics of interest, such as the Alcubierre metric, which allows faster-than-light (FTL) velocity; to the Gödel metric, which allows closed timelike curves (CTC); and the Kerr metric, which describes the spacetime geometry generated by a rotating body (rotating black holes). Furthermore, we will see that, although in [11] $\dot{\Omega} = 0$ (static spacetime) is an apparent requirement for the validity of the method, in fact the method is also valid for nonstatic spacetimes.

Our results might be of interest in the context of quantum simulation of the Dirac equation. Several ideas have been proposed for simulating the Dirac equation in Minkowski spacetime for 1 + 1 dimensional sections: e.g., experiments have been already realized with trapped ions [2,5] and Bose-Einstein condensates [7,8], while realistic proposals exist with superconducting circuits [15]. Also, some ideas have been proposed for simulating the Dirac equation in curved spacetime for 1 + 1 dimensional sections: e.g., through a mapping between this equation and a multiphoton quantum Rabi model [16]. However, since we obtain solutions in curved spacetime through transformations of Minkowski solutions, the results obtained in this work could be helpful for simulations of the Dirac equation in nontrivial spacetimes with currently existing setups.

The structure of the paper is the following. We start in Sec. II by recalling the procedure for obtaining solutions of the Dirac equation in curved spacetimes out of flat-spacetime solutions, first introduced in [11]. Then, we proceed to apply it in Sec. III to the aforementioned curved spacetimes of interest. Finally, we introduce a generalization of the method to nonstatic spacetimes in Sec. IV, which might be useful for further applications. We conclude in Sec. V with a summary of our results.

II. METHOD

We summarize here the procedure to obtain solutions of the Dirac equation in curved spacetimes, first discussed in [11]. In general, the Dirac equation in curved spacetime is obtained by replacing the standard partial derivatives in the Minkowski equation by the corresponding covariant ones, where the affine connection would carry the dependence on the particular form of the metric. In 1 + 1 D, using the conformally flat form of the metric and with a little algebra (see the details in [10]), we can get

$$i\left(\partial_t + \frac{\dot{\Omega}}{2\Omega}\right)\psi = -i\sigma_x\left(\partial_x + \frac{\Omega'}{2\Omega}\right)\psi. \quad (1)$$

For the static case, in which the conformal factor is time-independent $\dot{\Omega} = 0$, the equation is

$$i\partial_t\psi = -i\sigma_x\left(\partial_x + \frac{\Omega'}{2\Omega}\right)\psi. \quad (2)$$

If we make the transformation $\psi = \Omega^{-1/2}\phi$, Eq. (2) becomes

$$i\partial_t\phi = -i\sigma_x\partial_x\phi$$

which corresponds to the Dirac equation for a massless particle in Minkowski spacetime, whose solution is well known (see, for instance, [17]). Thus, using this transformation and the conformal factor of the 1 + 1 dimensional metric, we can find analytical solutions for the Dirac equation in curved spacetime. So, for a given metric, we have to follow this three-step procedure:

- (1) Find a change of coordinates $(t, x) \rightarrow (\bar{t}, \bar{x})$ such that, in the new coordinates, the metric acquires the form $ds^2 = \Omega^2(-d\bar{t}^2 + d\bar{x}^2)$.
- (2) Using those new coordinates (\bar{t}, \bar{x}) , apply $\psi(\bar{t}, \bar{x}) = \Omega^{-1/2}(\bar{x})\phi(\bar{t}, \bar{x})$, where $\phi(\bar{t}, \bar{x})$ is the solution of the Minkowski spacetime Dirac equation.
- (3) Once having $\psi(\bar{t}, \bar{x})$, apply the coordinate change $(\bar{t}, \bar{x}) \rightarrow (t, x)$ to finally find the solution of the curved spacetime Dirac equation in (t, x) coordinates.

Please note that, in spite of its apparent simplicity, in general it might not be necessarily straightforward to perform steps (1) and (3). To write the metric in a conformally flat form, it might be needed to express the change of coordinates in a differential form, which could generate an equation that might still not be straightforward to solve, as we will see in Sec. III B.

This method was applied in [11] to find solutions of the Dirac equation in a 1 + 1 dimensional section of a traversable wormhole spacetime. Let us now consider other examples of interest.

III. RESULTS

A. Gödel and Alcubierre metric

Both metrics will be discussed in the same section because of their similarity.

In 1994, Alcubierre proposed a metric that, in principle, allows FTL motion [18]. Note that, in general relativity, FTL speed is only forbidden locally. This is not as exotic as it might seem at first glance: for instance, the expansion of the Universe can make that two distant galaxies move at FTL speed between them, while each one is moving locally inside its light cone. The opposite might be possible too: if the spacetime were contracting fast enough, each galaxy were moving near the speed of light locally (inside its light cone) in opposite directions, but globally both were getting closer. With these considerations in mind, Alcubierre's idea is simple: to create, in the front of an object, a spacetime contraction, and in the back, a spacetime dilation. Thus, the contraction will pull the object forward, and the dilation will push the object forward too. Locally the object will be inside its light cone, but due to this spacetime manipulation,

it would move FTL—as compared with c , the speed of light in flat-spacetime vacuum.

The Alcubierre metric inside the “bubble” created by the space-temporal contraction/dilation, under the limit $\sigma \rightarrow \infty$ [18], and taking a spatial section $y = y_0$ and $z = z_0$ (with constant y_0, z_0), acquires the form [18,19]

$$ds^2 = -(1 - v_s^2)dt^2 + dx^2 - 2v_s dx dt. \quad (3)$$

As expected, Eq. (3) becomes the Minkowski metric when $v_s = 0$. Now we make a coordinate change $(t, x) \rightarrow (\bar{t}, \bar{x})$, such that

$$\begin{aligned} dt &= d\bar{t} \\ dx &= d\bar{x} + v_s d\bar{t}. \end{aligned}$$

The Alcubierre metric, using those new coordinates, acquires the form $ds^2 = -d\bar{t}^2 + d\bar{x}^2$, which is simply the Minkowski metric, so the conformal factor will be $\Omega^2 = 1$, and the solution of the curved spacetime Dirac equation, $\psi(\bar{t}, \bar{x})$, will be equal to the Minkowski spacetime solution, $\phi(\bar{t}, \bar{x})$. Now, due to the big similarity—regarding to the application of our techniques—between the Alcubierre and Gödel metric, we first discuss the latter case and then we compare both results. It is relevant to mention that, although the Dirac equation in a Gödel universe has already been considered in the study of tachyon’s stability [20], our objective is to find explicit analytical solutions for this equation in the Gödel’s background, using our procedure.

Gödel, in 1949, found a solution of the Einstein equations [21] corresponding to a homogeneous mass distribution that rotates at each point of the space [22]. That distribution of matter causes unusual effects, such as the existence of CTC. In cylindrical coordinates [22], the metric is given by the following expression:

$$\begin{aligned} ds^2 &= dt^2 - \frac{dr^2}{1 + \left(\frac{r}{2a}\right)^2} - r^2 \left(1 - \left(\frac{r}{2a}\right)^2\right) d\phi^2 - dz^2 \\ &+ \frac{2r^2}{a\sqrt{2}} dt d\phi, \end{aligned} \quad (4)$$

where a is a parameter with units of length, which represents a characteristic distance. In particular, $r_G = 2a$ represents the critical radius from which CTC can exist [22].

Now, taking a radial section with $\phi = \phi_0$ and $z = z_0$ [19], the Gödel metric becomes

$$ds^2 = dt^2 - \frac{1}{1 + \left(\frac{r}{2a}\right)^2} dr^2.$$

By making the coordinate change $(t, r) \rightarrow (\bar{t}, \bar{r})$, with

$$\begin{aligned} d\bar{r}^2 &= \frac{1}{1 + \left(\frac{r}{2a}\right)^2} dr^2 \\ d\bar{t}^2 &= dt^2 \end{aligned} \quad (5)$$

the previous metric transforms into the Minkowski metric $ds^2 = d\bar{t}^2 - d\bar{r}^2$. Furthermore, we can find the relationship

between the (t, r) and (\bar{t}, \bar{r}) coordinates by performing the following integration:

$$\bar{r}(r) = \int \frac{dr}{\sqrt{1 + \left(\frac{r}{2a}\right)^2}}$$

whose solution—setting $\bar{r}(r = 0) = 0$ —is

$$\bar{r}(r) = 2a \sinh^{-1} \left(\frac{r}{2a} \right).$$

With these new coordinates, the conformal factor becomes $\Omega^2 = 1$, and the Minkowski spacetime solution will be the same as the curved spacetime solution, as in the case of the Alcubierre metric. Thus, the only difference between the Gödel and the Alcubierre metric is the relationship between the original coordinates and the new coordinates.

Now, if we assume that the wave function has a Gaussian initial form:

$$\phi(\bar{x}, 0) = N e^{-\frac{(\bar{x}-\bar{x}_0)^2}{\sigma^2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix},$$

where \bar{x} is the conformally flat coordinate and N is a normalization constant, the solution for the Dirac equation in Minkowski’s spacetime will be [11,17]

$$\phi(\bar{x}, \bar{t}) = N e^{-\frac{(\bar{x}-\bar{x}_0)^2}{\sigma^2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}. \quad (6)$$

Since $\psi = \phi$, to find the curved spacetime solution we only have to apply the coordinate change for each metric. In the Alcubierre case we have

$$\phi(\bar{x}(x), \bar{t}(t)) = N e^{-\frac{(x-x_0)^2}{\sigma^2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad (7)$$

while, for the Gödel metric

$$\phi(\bar{r}(r), \bar{t}(t)) = N e^{-\frac{(r-(2a\sinh^{-1}(r/(2a)-\bar{x}_0))^2}{\sigma^2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}. \quad (8)$$

Both solutions are represented in Fig. 1.

For (a), (b) and (c), as the parameter v_s increases, the slope of the Gaussian wave packet’s “trajectory” decreases. In other words, the particle’s velocity increases as v_s is bigger. In fact, if we take a look to the expression (7), we can see its equivalence to the solution of the Minkowski spacetime Dirac equation (6) if we substitute t with $(1 + v_s)t$. This means that, physically, the particle moves with a FTL velocity $\bar{c} = 1 + v_s$, as expected for a massless particle since this is precisely the speed of light in this spacetime.

For (d), (e) and (f), we can see that the particle velocity is no longer constant. This variation is inversely proportional




FIG. 1. Spacetime diagrams for the probability density of Dirac equation solutions in (a)–(c) Alcubierre metric with (a) $v_s = 0$ (equivalent to Minkowski spacetime), (b) $v_s = 0.5$ (subluminal spacetime bubble), (c) $v_s = 2$ (superluminal spacetime bubble); (d)–(f) Gödel metric with (d) $a = 2$, (*e) $a = 1$, (f) $a = 0.5$. Please note that the wave function is not normalized.

to the value of a , as expected, since in the limit $a \rightarrow \infty$ the Gödel metric (4) becomes the Minkowski metric in cylindrical coordinates. In addition, the slope decreases with the propagation of the particle, implying FTL velocity. Of course, in general the existence of CTC—which is the most genuine feature of Gödel spacetime—is intimately linked with the possibility of FTL (see, for instance, [23]).

B. Rotating black hole

The main motivation to study the Kerr metric is the fact that it describes the spacetime geometry of a rotating body, so it can be used to analyze the physics of black holes in a more realistic scenario than the one provided by the Schwarzschild metric. Using Boyer-Lindquist coordinates (t, r, θ, ϕ) , the Kerr metric is [24,25]

$$ds^2 = -\left(1 - \frac{2Mr}{\Sigma}\right) dt^2 - \frac{4Marsin^2\theta}{\Sigma} dt d\theta + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2 + \left(r^2 + a^2 + \frac{2Ma^2r\sin^2\theta}{\Sigma}\right) \sin^2\theta d\phi^2,$$

where

$$\Sigma(r, \theta) = r^2 + a^2 \cos^2\theta \quad \Delta(r) = r^2 + a^2 - 2Mr.$$

If we take a radial section ($\theta = \theta_0, \phi = \phi_0$), we obtain [19]

$$ds^2 = -\left(1 - \frac{2Mr}{\Sigma(r, \theta_0)}\right) dt^2 + \frac{\Sigma(r, \theta_0)}{\Delta(r)} dr^2,$$

where $\Sigma(r, \theta_0)$ is a function of the coordinate r [for simplicity, $\Sigma(r, \theta_0) = \Sigma(r)$]. So, if we make the coordinate change $(t, r) \rightarrow (\bar{t}, \bar{x})$ with

$$\begin{aligned} \frac{\Sigma(r)}{\Delta(r)} dr^2 &= \left(1 - \frac{2Mr}{\Sigma(r)}\right) d\bar{x}^2 \\ dt^2 &= d\bar{t}^2 \end{aligned} \quad (9)$$

the metric acquires the form

$$ds^2 = \Omega^2(r)(-d\bar{t}^2 + d\bar{x}^2)$$

with

$$\Omega^2(r) = 1 - \frac{2Mr}{\Sigma(r)}.$$

Due to the conditions (9), r will be a function of \bar{x} , whereas $t = \bar{t}$, so the conformal factor $\Omega(r)$ will be a function of $r(\bar{x})$:

$$\Omega^2(r(\bar{x})) = 1 - \frac{2Mr(\bar{x})}{\Sigma(r(\bar{x}))}. \quad (10)$$

To find the conformal factor $\Omega(r(\bar{x}))$ we need to obtain $r(\bar{x})$. Using the first equation in (9) and, after doing some algebraic manipulations, we obtain that $\bar{x}(r)$ is given by

$$\bar{x}(r) = \int dr \Sigma(r) \sqrt{\frac{1}{\Delta(r)(\Sigma(r) - 2Mr)}} + C. \quad (11)$$

In general, solving this integral is not straightforward. Later, we will analyze some particular cases in which the integral can be solved. For now, let us just assume that $r(\bar{x})$ is known. Thus, the conformal factor is given by Eq. (10) and, therefore, the relation between the solution to Dirac equation in curved spacetime (ψ) and flat spacetime (ϕ) will be

$$\begin{aligned} |\psi(\bar{x}, \bar{t})|^2 &= |\Omega^{-1/2}(r(\bar{x}))|^2 |\phi(\bar{x}, \bar{t})|^2 \\ &= \left| \sqrt{\frac{\Sigma(r(\bar{x}))}{\Sigma(r(\bar{x})) - 2Mr(\bar{x})}} \right| |\phi(\bar{x}, \bar{t})|^2. \end{aligned} \quad (12)$$

All the wave function properties caused by the spacetime curvature are contained in the factor $\Omega^{-1}(r(\bar{x}))$, so it is interesting to analyze this function. First, there exist regions in which the probability density of the wave function in curved spacetime becomes infinite. It happens when the

following condition is fulfilled—we relax the notation and write r instead of $r(\bar{x})$:

$$\Sigma(r) - 2Mr = 0. \quad (13)$$

This condition is satisfied whenever $r = r_{\pm}$, with

$$r_{\pm} = M \pm \sqrt{M^2 - a^2 \cos^2 \theta_0}, \quad (14)$$

which are the well-known apparent singularities where the temporal component of the metric changes sign, defining the ergosphere [26]. On the other hand, there exist points in which the wave function is null, independently of the form of the solution in Minkowski's spacetime. This will happen whenever $\Omega^{-1}(r) = 0$, that is, when

$$\Sigma(r) = 0 \rightarrow r^2 + a^2 \cos^2 \theta_0 = 0,$$

which will be satisfied when at least one of those conditions apply:

- (1) $r = 0$, $a = 0$ (no rotation).
- (2) $r = 0$, $\theta_0 = \frac{\pi}{2}$.

Finally, there is a region of the space in which the conformal factor is an imaginary number. This occurs if

$$\Sigma(r) - 2Mr < 0.$$

This second-degree inequality is verified for $r_- < r < r_+$, namely, within the ergosphere.

1. Radial section with $\theta_0 = 0$

Let us consider $\theta_0 = 0$. Then $\Sigma(r) = r^2 + a^2$, $\Delta(r) = \Sigma - 2Mr$, and the integral becomes

$$\bar{x}(r) = \int dr \frac{\Sigma(r)}{\Delta(r)} + C = \int dr \frac{r^2 + a^2}{r^2 + a^2 - 2Mr} + C$$

whose solution is

$$\begin{aligned} \bar{x}(r) &= \frac{2M^2}{\sqrt{a^2 - M^2}} \tan^{-1} \left(\frac{r - M}{\sqrt{a^2 - M^2}} \right) \\ &+ M \log(a^2 - 2Mr + r^2) + r + C. \end{aligned}$$

Bearing in mind that, for a rotating black hole, $M^2 > a^2$ is satisfied (otherwise, it would not be a black hole, but a highly rotating body [27]), and using the relationship $\tan^{-1}(ix) = i \tanh^{-1}(x)$, we can write the solution as

$$\begin{aligned} \bar{x}(r) &= \frac{2M^2}{\sqrt{M^2 - a^2}} \tanh^{-1} \left(\frac{M - r}{\sqrt{M^2 - a^2}} \right) \\ &+ M \log(a^2 - 2Mr + r^2) + r + C. \end{aligned} \quad (15)$$

The constant of integration can be chosen in such a way that the constant imaginary part of Eq. (15) cancels out in each of the three spacetime regions defined by the apparent singularities r_{\pm} . Thus,




FIG. 2. $\bar{x}(r)$ for different values of the parameter a . In the absence of rotation ($a = 0$), the singular points are in $r = 0$ and $r = 2M$ (Schwarzschild's black hole). As the rotation increases, both points ($r = r_{\pm}$) come closer to $r = M$.

$$C = \begin{cases} -i \frac{\pi M^2}{\sqrt{M^2 - a^2}} & \text{if } r < r_- \\ -i M \pi & \text{if } r_- < r < r_+ \\ -i \frac{\pi M^2}{\sqrt{M^2 - a^2}} & \text{if } r_+ < r. \end{cases}$$

In Fig. 2, we plot $\bar{x}(r)$ for different values of M and a .

Finally, assuming again that our particle is characterized by a Gaussian wave packet, the solution in Kerr spacetime is

$$\begin{aligned} |\psi(\bar{x}(r), t)|^2 &= \sqrt{\frac{\Sigma(r)}{\Sigma(r) - 2Mr}} |\phi(\bar{x}(r), t)|^2 \\ &= 2N^2 \sqrt{\frac{\Sigma(r)}{\Sigma(r) - 2Mr}} e^{-\frac{2(t - (\bar{x}(r) - \bar{x}_0))^2}{\sigma^2}}. \end{aligned}$$

This solution is represented in Fig. 3 for several values of a and M .

Naturally, when we set $M = a = 0$, the metric becomes the Minkowski one, $\Omega^2 = 1$, and the expression (15) reduces to $\bar{x}(r) = r$. In other words, the solution corresponds to a free particle in flat spacetime, as it is observed in Fig. 3(a). If we slightly increase the value of M (with $a = 0$, thus we are in Schwarzschild spacetime), we see that the free solution is slightly disturbed, increasing the probability density near the horizon, as can be seen in Fig. 3(b). As the mass increases, the probability density accumulates nearby the horizon, placed in $r = 2M$. In addition, we see that the width of the wave packet is drastically reduced (the particle is more localized in space), and the particle's speed diminishes, vanishing in the horizon. Finally, if we set the mass to a constant value and we change a [Figs. 3(e)–3(h)], we see that, as a increases, the inner horizon is generated, around which, again, the probability density accumulates in the same




FIG. 3. Dirac equation solution for Kerr metric and different values of the parameters a and M . (a)–(d), $a = 0$ (no rotation, Schwarzschild metric) and M ranging from 0 to 2. The horizon is at $r = 2M$. (e)–(h) $M = 2$ and a ranging from 0 to 2, two singularities appear according to Eq. (14) with $\theta_0 = 0$ giving rise to an ergosphere which splits the full spacetime into three separate regions. Notice that as a increases r_+ , r_- move from $2M$ and 0, respectively, to M , where they would merge in the limit $M = a$. Please note that the wave function is not normalized.

way as before. In addition, in the region between both singular points—the ergosphere—we see how the particle tends to be expelled towards the interior or exterior horizon.

IV. NONSTATIC SPACETIMES

So far, we have only analyzed static metrics, that is, $\dot{\Omega} = 0$. We now show that our techniques are also valid

for the case of nonstatic metrics ($\dot{\Omega} \neq 0$). Performing the substitution $\psi = \Omega^{-1/2}\phi$ in Eq. (1) and with a little algebra, we find

$$i\left(\partial_t + \frac{\dot{\Omega}}{2\Omega}\right)(\Omega^{-1/2}\phi) = -i\sigma_x\left(\partial_x + \frac{\Omega'}{2\Omega}\right)(\Omega^{-1/2}\phi)$$

$$\partial_t(\Omega^{-1/2}\phi) + \frac{\dot{\Omega}}{2\Omega}\Omega^{-1/2}\phi = -\sigma_x\left[\partial_x(\Omega^{-1/2}\phi) + \frac{\Omega'}{2\Omega}\Omega^{-1/2}\phi\right].$$

Notice that

$$\partial_t(\Omega^{-1/2}\phi) = -\frac{1}{2}\Omega^{-3/2}\dot{\Omega}\phi + \Omega^{-1/2}\partial_t\phi. \quad (16)$$

Therefore the terms not containing $\partial_t\phi$ cancel out. The same thing happens with the spatial partial derivative: the terms without $\partial_x\phi$ cancel out. Thus, putting everything together we recover the equation:

$$i\partial_t\phi = -i\sigma_x\partial_x\phi$$

which is the Dirac equation for Minkowski spacetime. Therefore, our techniques could be also applied to nonstatic spacetimes, such as Friedmann-Lemaître-Robertson-Walker spacetime, as in [10].

V. CONCLUSIONS

Following [11], we have shown that is possible to obtain solutions of the Dirac equation in curved spacetime by means of a transformation $\psi(\bar{x}, \bar{t}) = \Omega^{-1/2}(\bar{x})\phi(\bar{x}, \bar{t})$, where $\psi(\bar{x}, \bar{t})$ is the curved spacetime solution, $\phi(\bar{x}, \bar{t})$ is the Minkowski spacetime solution and $\Omega(\bar{x})$ is the conformal factor. Then, in order to be able to apply this transformation, it is necessary to perform first a coordinate

change $(x, t) \rightarrow (\bar{x}, \bar{t})$, such that in the new coordinates, the metric is conformally flat. We have applied this technique to three different metrics: 1 + 1 dimensional sections of Alcubierre, Gödel and Kerr metrics. For Alcubierre and Gödel, the conformal factor turns out to be $\Omega^2 = 1$, so the only difference between both solutions is the coordinate transformation employed in each case. Setting a Gaussian solution in Minkowski spacetime, and after undoing the coordinate transformation, we obtain the solutions (7) and (8), for Alcubierre and Gödel spacetimes, respectively. In this way, we have been able to analyze the FTL behavior of massless particles in these exotic spacetimes. In the case of the Kerr metric, the conformal factor is nontrivial $\Omega^2(r) = 1 - \frac{2Mr}{\Sigma(r)}$. We find an analytical solution in a spatial section with $\theta_0 = 0$. We have discussed the dependence of this solution on the black-hole mass and rotation, as well as the behavior of the particle near the singularities and within the ergosphere. Finally, we have shown that our technique is valid also for nonstatic spacetimes.

Our results might, in principle, be useful in the context of quantum simulations of the Dirac equation. In particular, following the spirit of [11], notice that our technique entails that curved spacetimes can be encoded into a transformation realized onto a flat-spacetime Dirac wave packet. Therefore, in principle, our results could be tested in already existing quantum simulators of the Dirac equation in flat spacetime.

ACKNOWLEDGMENTS

C. S. has received financial support through the Junior Leader Postdoctoral Fellowship Programme from “la Caixa” Banking Foundation and Fundación General CSIC (ComFuturo Programme).

-
- [1] B. Thaller, *The Dirac Equation* (Springer-Verlag, Berlin-Heidelberg, 1992).
 - [2] R. Gerritsma, G. Kirchmair, F. Zähringer, E. Solano, R. Blatt, and C. F. Roos, *Nature (London)* **463**, 68 (2010).
 - [3] L. Lamata, J. León, T. Schätz, and E. Solano, *Phys. Rev. Lett.* **98**, 253005 (2007).
 - [4] J. Casanova, J. J. García-Ripoll, R. Gerritsma, C. F. Roos, and E. Solano, *Phys. Rev. A* **82**, 020101 (2010).
 - [5] R. Gerritsma, B. P. Lanyon, G. Kirchmair, F. Zähringer, C. Hempel, J. Casanova, J. J. García-Ripoll, E. Solano, R. Blatt, and C. F. Roos *Phys. Rev. Lett.* **106**, 060503 (2011).
 - [6] C. Sabín, J. Casanova, J. J. García-Ripoll, L. Lamata, E. Solano, and J. León, *Phys. Rev. A* **85**, 052301 (2012).
 - [7] T. Salger, C. Grossert, S. Kling, and M. Weitz, *Phys. Rev. Lett.* **107**, 240401 (2011).
 - [8] L. J. LeBlanc, M. C. Beeler, K. Jiménez-García, A. R. Perry, S. Sugawa, R. A. Williams, and I. B. Spielman, *New J. Phys.* **15**, 073011 (2013).
 - [9] M. Gasperini, *Theory of Gravitational Interactions*, 2nd ed. (Springer, Berlin, 2017).
 - [10] C. Koke, C. Noh, and D. Angelakis, *Ann. Phys. (Amsterdam)* **374**, 162 (2016).
 - [11] C. Sabín, *Sci. Rep.* **7**, 40346 (2017).
 - [12] H. Nitta, T. Kudo, and H. Minowa, *Am. J. Phys.* **67**, 966 (1999).
 - [13] F. M. Toyama, Y. Nogami, and F. A. B. Coutinho, *J. Phys. A* **30**, 2585 (1997).

- [14] V. Alonso and S. De Vincenzo, *Phys. Scr.* **61**, 396 (2000).
- [15] J. S. Pedernales, R. di Candia, D. Ballester, and E. Solano, *New J. Phys.* **15**, 055008 (2013).
- [16] J. S. Pedernales, M. Beau, S. M. Pittman, I. L. Egusquiza, L. Lamata, E. Solano, and A. del Campo, *Phys. Rev. Lett.* **120**, 160403 (2018).
- [17] B. Thaller, *Advanced Visual Quantum Mechanics* (Springer, New York, 2005).
- [18] M. Alcubierre, *Classical Quantum Gravity* **11**, L73 (1994).
- [19] C. Sabín, *New J. Phys.* **20**, 053028 (2018).
- [20] R. A. Konoplya, *Phys. Lett. B* **706**, 451 (2012).
- [21] K. Gödel, *Rev. Mod. Phys.* **21**, 447 (1949).
- [22] E. Kajari, R. Walser, W. P. Schleich, and A. Delgado, *Gen. Relativ. Gravit.* **36**, 2289 (2004).
- [23] C. Mallery, G. Khanna, and R. H. Price, *Classical Quantum Gravity* **35**, 175020 (2018).
- [24] R. P. Kerr, *Phys. Rev. Lett.* **11**, 237 (1963).
- [25] C. Bambi, *Rev. Mod. Phys.* **89**, 025001 (2017).
- [26] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, England, 1973).
- [27] H. Stephani, *Relativity. An Introduction to Special and General Relativity*, 3rd ed. (Cambridge University Press, Cambridge, England, 2004).