

Vortices of $SO(2)$ gauged Skyrmons in $2 + 1$ dimensions

Francisco Navarro-Lérida¹ and D. H. Tchrakian^{2,3}

¹*Departamento de Física Teórica and IPARCOS, Ciencias Físicas, Universidad Complutense de Madrid, E-28040 Madrid, Spain*

²*School of Theoretical Physics, Dublin Institute for Advanced Studies, 10 Burlington Road, Dublin DO4 C932, Ireland*

³*Department of Computer Science, Maynooth University, Maynooth, Ireland*



(Received 14 December 2018; published 15 February 2019)

Vortices of the $SO(2)$ gauged planar Skyrme model with (a) only Maxwell, (b) only Chern-Simons, and (c) both Maxwell and Chern-Simons dynamics are studied systematically. In cases (a) and (b), where both models feature a single parameter λ (the coupling of the potential term), the dependence of the energy on λ is analyzed. It is shown that the plots of the energy vs λ feature discontinuities and branches. In case (c), the emphasis is on the evolution of the topological charge, taking noninteger values. Throughout, the properties studied are contrasted with those of the corresponding Abelian Higgs models.

DOI: [10.1103/PhysRevD.99.045007](https://doi.org/10.1103/PhysRevD.99.045007)

I. INTRODUCTION

Solitons of the gauged Skyrme model, Skyrmons, present much more complex properties than their Higgs counterparts, e.g., the Abelian Higgs vortices in $2 + 1$ dimensions [1], monopoles in $3 + 1$ dimensions [2,3] and monopoles in $D + 1$ dimensions [4]. One reason for this is that the topological charge densities of gauged Higgs systems can be systematically constructed with a top down approach [4] by subjecting Chern-Pontryagin densities to dimensional descent, while the topological charge densities of gauged Skyrme models, as defined in Ref. [5], can be constructed instead only with a bottom up approach.¹

By gauged Skyrmon, we understand the soliton of a $SO(N)$ gauged $O(D + 1)$ sigma model² in $D + 1$ dimensional spacetime, or on \mathbb{R}^D , with $D \geq N \geq 2$, supporting topologically stable solitons. In the case at hand, this is the $SO(2)$ gauged $O(3)$ sigma model. It is understood that the vacuum of the gauged model is the same as the vacuum of the gauge-decoupled model, such that the gauge symmetry

¹Indeed this bottom up approach can be applied also to Higgs field systems [6]. In that case, the bottom up definition coincides with the top down definition of the topological charge only for the Higgs models on \mathbb{R}^2 .

²The most general Lagrangian of a Skyrme model, whether gauged or ungauged, consists of a potential term plus quadratic, quartic, and up to $2D$ -atic kinetic terms. This is to satisfy the Derrick scaling requirement.

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

is not broken in the vacuum. This is in stark contrast to (all) Higgs models, whose dynamics explicitly displays (gauge) symmetry breaking.

Our purpose is to investigate the detailed properties of $SO(2)$ gauged Skyrmons whose dynamics is controlled by a Maxwell (M) term, or a Chern-Simons (CS) term, or both (MCS). The influence of the CS dynamics in the model incorporating both Maxwell and Chern-Simons terms, was studied recently³ in (Sec. IV of) Ref. [9]. There the emphasis was laid on the (unusual) dependence of the energy on the electric charge and the angular momentum. Here, by contrast, our attention will be focused on the evolution of the topological charge, which we recently considered in [10] both for $2 + 1$ - and $3 + 1$ -dimensional $SO(2)$ gauged Skyrmons.

Topologically stable vortices of the $SO(2)$ gauged planar Skyrmon with Maxwell dynamics were first given by Schroers [11] and later in Ref. [12]. The vortices constructed in [11] were self-dual, while the vortices presented in [12]⁴ were solutions of second-order equations for a model which in the gauge-decoupling limit reduced to the (ungauged) planar Skyrme model. The definition of the topological charge proposed in [11] was subsequently extended to higher dimensions in Ref. [5]. (See also Appendix B of [13], and references therein.) Employing the Chern-Simons term as an alternative for the Maxwell

³Similar models were used in Refs. [7,8], with various self-interaction potentials, where the solutions were constructed subject to no symmetries, where the effect of the Chern-Simons term on the interaction of the solitons was investigated.

⁴An interesting question studied in [12] was the dependence of the topological lower bound on the gauge coupling constant.

dynamics was considered in Refs. [14–16], in analogy with its Higgs counterpart studied in Refs. [17,18].

In the present work, we consider (a) the model whose gauge dynamics is described by the Maxwell term, namely the model studied in Ref. [11], (b) the model whose gauge dynamics is described by the Chern-Simons term, studied in Refs. [14–16], and (c) the model whose gauge dynamics is described both by the Maxwell and the Chern-Simons terms, e.g., as in [9], where the emphasis was different.

We have two distinct but not unrelated objectives. The *first* objective is to reveal the peculiar dependence of the energy on a parameter in the model, that is typical of gauged Skyrmons. This has been studied in the case of the $SO(3)$ gauged Skyrmion on \mathbb{R}^3 , namely the $O(4)$ sigma model, where the energy was plotted against the coupling of the (quartic kinetic) Skyrme term. It was found in [19,20] and in [21] that the energy profile exhibited discontinuities and branches. This contrasts with the corresponding Higgs model, namely the $SO(3)$ gauged Georgi-Glashow model supporting the monopole [2,3], where the energy profile plotted against the coupling strength of the Higgs self-interaction potential is shown [22] to be continuous and monotonic and, as it happens, flattens out for large Higgs coupling. The situation in the case of the $SO(2)$ gauged Higgs model on \mathbb{R}^2 , namely the Abelian Higgs model supporting vortices, is qualitatively similar to the three-dimensional case. It is shown in Ref. [23] that energy profiles plotted against the Higgs coupling are continuous and monotonic and increase logarithmically for large Higgs coupling. The corresponding study for any $SO(2)$ gauged Skyrmion in $2+1$ dimensions, either with Maxwell [11], or with Chern-Simons [14–16] dynamics, has not been carried out. This is our first objective. Our analysis here will confirm that also in the two dimensional case, some energy profiles exhibit discontinuities and branches.

Our second objective, which from the viewpoint of dynamics is the dominant one, is to track the evolution of the topological charge in a given gauged Skyrme model. (The relationship of the energy with the electric charge and the spin are already studied in Ref. [9].) The gauge-decoupled Skyrmion has integer “baryon number” which is calculated as the volume integral of the winding number density. Introduction of the $SO(2)$ gauge field alters the definition of the topological charge density, and further, in the presence of Chern-Simons dynamics the integral of the altered topological charge density departs from the integer value. This mechanism is predicated on the presence of the electric component of the gauge potential, which in turn results from the presence of the Chern-Simons term. (Not surprisingly, this mechanism is also responsible for the unusual energy/electric charge and energy/angular momentum dependence, which was the subject of investigation in Ref. [9].)

The most general Lagrangian considered is

$$\begin{aligned} \mathcal{L}_{(\lambda)} = & -\frac{1}{4}\beta F_{\mu\nu}^2 + \kappa\epsilon^{\lambda\mu\nu}A_\lambda F_{\mu\nu} - \frac{1}{8}\tau|D_{[\mu}\phi^a D_{\nu]}\phi^b|^2 \\ & + \frac{1}{2}\eta^2|D_\mu\phi^a|^2 - \eta^4V[\phi]. \end{aligned} \quad (1)$$

In achieving our first objective, we employ the two models: (a) with Maxwell term only, i.e., with $\kappa = \tau = 0$ and (b) with Chern-Simons term only, i.e., with $\beta = \tau = 0$. We study the energy profiles plotted against the coupling strength of the potential term, with special emphasis to the first-order self-dual solutions in each case. The models considered do not contain quartic kinetic “Skyrme” terms since inclusion of these would prevent the saturation of the topological lower bound, and the study of first-order solutions is the main feature of these models.

To achieve our second objective, we employ the model (c) incorporating both Maxwell and Chern-Simons terms, *i.e.*, with nonvanishing couplings β and κ in (1). This is the simplest system which enables the tracking of the topological charge leading to its annihilation. We have however incorporated the quartic Skyrme kinetic term as well with coupling τ . The first reason is that the model in question should, in the gauge decoupling limit, support the ungauged Skyrmion characterized by the “baryon number” prior to gauging. This necessitates the inclusion of the quartic Skyrme term. It turns out also, that the presence of this term renders the numerical computations simpler. It is important to note that to achieve this aim, the potential $V[\phi]$ is chosen suitably as explained in Appendix C.

Our study of (a) the gauged planar Skyrmion with Maxwell dynamics is presented in Sec. II, where we have employed two different potentials. The first one is the potential employed in Ref. [11] which allows the construction of self-dual solutions. The second model we study employs the “pion mass,” where solutions to the second-order equations are constructed. The numerical integrations here are simpler since the decay is exponential. In Sec. III, we study (b) the model with (only) Chern-Simons dynamics, where the new choice of potentials arises. In both Secs. II and III, we have studied the solutions of the second-order equations, in addition to those of the first-order equations. In Sec. IV, the combined Maxwell–Chern-Simons model (c) is studied, where there are only solutions to the second-order equations. Throughout, the said properties are studied quantitatively using numerical methods. A summary of our results is given in Sec. V. The topological charge densities are defined in Appendix A, and the resulting topological lower bounds on the energy densities (Belavin inequalities) for Maxwell dynamics and for Chern-Simons dynamics are presented in Appendixes B and C, respectively. Since all the features revealed contrast starkly with the corresponding features in Higgs models, and since the crucial role is played by the topological

charges and Belavin inequalities, we have inserted subsections in each Appendix exposing the (familiar) definitions for the corresponding Higgs models against which their Skyrme counterparts are contrasted.

Before proceeding to the study of the three types of models (a), (b), and (c), we state the results of symmetry imposition, including the topological and global charges, which are common to all types of models (a), (b), and (c).

A. Imposition of symmetry, topological, and the global charges

Since the field content of the models studied in the next three sections is the same one, we state the results of the imposition of axial symmetry and the resulting charges here in advance. The definitions of the topological charges prior to the imposition of symmetry, are given in Appendix A. For the definition of the familiar global charges, we refer to Ref. [9].

1. Imposition of symmetry

The planar Skyrme model, namely the $O(3)$ sigma model, is described by the Skyrme scalar $\phi^a = (\phi^\alpha, \phi^3)$ subject to the constraint $|\phi^a|^2 = 1$. The axially symmetric *Ansatz* is

$$\phi^\alpha = \sin f(r)n^\alpha, \quad \phi^3 = \cos f(r), \quad n^\alpha = \begin{pmatrix} \cos n\theta \\ \sin n\theta \end{pmatrix}, \quad (2)$$

where r is the radial coordinate, θ is the azimuthal angle, and n is the winding number of the Skyrme scalar.

Imposition of symmetry on the $SO(2)$ (Abelian) gauge field $A_\mu = (A_i, A_0)$ $\mu = i, 0$, is achieved by the *Ansatz*

$$A_i = \left(\frac{a(r) - n}{r} \right) (\epsilon \hat{x})_i \Rightarrow F_{ij} = -\frac{a'}{r} \epsilon_{ij}, \quad (3)$$

$$A_0 = b(r) \Rightarrow F_{i0} = b'. \quad (4)$$

The prescription of gauging employed is

$$D_\mu \phi^\alpha = \partial_\mu \phi^\alpha + A_\mu (\epsilon \phi)^\alpha, \quad (\epsilon \phi)^\alpha = \epsilon^{\alpha\beta} \phi^\beta, \quad (5)$$

$$D_\mu \phi^3 = \partial_\mu \phi^3, \quad (6)$$

and the covariant derivatives of ϕ^a are

$$D_i \phi^\alpha = (\sin f)' \hat{x}_i n^\alpha - \frac{a \sin f}{r} (\epsilon \hat{x})_i (\epsilon n)^\alpha, \quad (7)$$

$$D_0 \phi^\alpha = b \sin f (\epsilon n)^\alpha, \quad (7)$$

$$D_i \phi^3 = (\cos f)' \hat{x}_i, \quad D_0 \phi^3 = 0. \quad (8)$$

Anticipating the usual choice of boundary conditions (A2) in the Appendix, the corresponding one-dimensional conditions read

$$\lim_{r \rightarrow 0} f(r) = \pi, \quad \lim_{r \rightarrow \infty} f(r) = 0, \quad (9)$$

consistent with the finiteness of the energy, as well as with regularity at the origin. This choice of boundary values means that we are excluding the choice of symmetry-breaking potentials.⁵

Concerning the boundary values of the function $a(r)$ (3) at the origin, regularity requires that

$$\lim_{r \rightarrow 0} a(r) = n. \quad (10)$$

As for $a(\infty)$, the boundary values of the function $a(r)$ at infinity, this will be left free. This is in stark contrast with usual⁶ Abelian Higgs model, where $a(\infty) = 0$.

2. Topological charge

Calculating the integral of ϱ_0 defined by (A5), subject to (2) and (3), gives $\int \varrho_0 d^2x = -8\pi n$. We, thus, denote the topological charge density of the ungauged Skyrmion as

$$q_0 = -\frac{1}{8\pi} \int \varrho_0 d^2x = n, \quad (11)$$

which is the ‘‘baryon number’’ n .

After gauging, the topological charge is given by the density (A10) and (A11), and the resulting topological charge is

$$q = -\frac{1}{8\pi} \left(\int \varrho_0 d^2x + 2\epsilon_{ij} \int \partial_i [(\phi^3 - v)A_j] d^2x \right). \quad (12)$$

When the choice $v = 1$ is made, which as we have seen in Appendix B is obligatory in the case when the Abelian dynamics is controlled by the Maxwell term only, the topological charge equals the ‘‘baryon number’’ n .

With all other values of v , the topological charge departs from n . As seen in Appendix C, the choice of $v \neq 1$ is viable in models featuring a Chern-Simons term, and the results obtained in the generic case are qualitatively the same. Here, we have made the choice $v = 0$ throughout, as this is the simplest option.

⁵Such potentials lead to a magnetic flux, and such models were considered in [24], investigating the relation of the magnetic flux and spin. Recently symmetry-breaking potentials in $2 + 1$ dimensions were employed in [25], where minimal energy multi-solitons and their interactions were considered. This is not in the scope of the present work.

⁶The usual Abelian Higgs (AH) model is the $p = 1$ AH model, for which $a(\infty) = 0$ is forced by the requirement of finite energy. By contrast, all other AH models with $p \geq 2$ can support finite energy solutions [9] for $a(\infty) \neq 0$.

With $v = 0$ the topological charge is

$$q = -\frac{1}{8\pi} \left(\int \varrho_0 d^2x + 2\varepsilon_{ij} \int \partial_i [(\phi^3) A_j] d^2x \right), \quad (13)$$

where the contribution of the second integral is not necessarily vanishing.

Evaluating (13) subject to the symmetries (2) and (3) gives rise to

$$q = \frac{1}{2} (n + a_\infty), \quad (14)$$

where $a_\infty = a(\infty)$. The expression (14) enables tracking the evolution of the topological charge (“baryon number”) by considering solutions with evolving values of a_∞ , i.e., where $a_\infty \neq 0$. It is important here to stress that this is a strictly Skyrme theoretic mechanism, and is excluded by Higgs theoretic dynamics. Notably, $a_\infty = 0$ for the usual ($p = 1$) Abelian Higgs (AH) model.⁷

3. Global charges

We will also consider the energy E and the angular momentum J of these solutions. Their corresponding expressions can be derived from the 00 and 0 φ components of the stress-energy tensor, respectively. Anticipating the model studied in Sec. IV, the reduced one-dimensional energy density of that model is

$$\begin{aligned} \mathcal{E} = & \frac{1}{2} \left(\frac{a^2}{r^2} + b^2 \right) + \tau \left(\frac{a^2}{r^2} + b^2 \right) f'^2 \sin^2 f \\ & + \frac{1}{2} \eta^2 \left[f'^2 + \left(\frac{a^2}{r^2} + b^2 \right) \sin^2 f \right] \\ & + \frac{1}{32} \lambda \left(\frac{\eta^3}{\kappa} \right)^2 \sin^2 f \cos^2 f. \end{aligned} \quad (16)$$

The reduced angular momentum density is

⁷It is, however, the case that vortices of the $p \geq 2$ AH models, alluded to in footnote 3 above, with $a_\infty \neq 0$ exist. This fact does not lead to the evolution of the topological charge (the magnetic vortex number μ). To see this we recall the expression for the topological charge of the p -AH model, which is descended [4] from the $2p$ -th Chern-Pontryagin density. In terms of the radial functions $a(r)$ and $h(r)$ parametrizing the radially symmetric $U(1)$ field and the Higgs field $\phi^\alpha = hn^\alpha$, in the notation of Sec. IA 1 above, the expression for the topological charge corresponding to (14) is

$$\begin{aligned} \mu = & \frac{1}{2\pi} \int \frac{d}{dr} [(1 - h^2)^{2p-1} a] dr = [(1 - h^2)^{2p-1} a]_{r=0}^{r=\infty} \\ = & a(0) = n, \end{aligned} \quad (15)$$

since $h(\infty) = 1$. The topological charge of a p -Abelian Higgs model cannot take any other value but the vortex number n .

$$\mathcal{J} = a'b' + \eta^2 ab \sin^2 f + 2\tau ab f'^2 \sin^2 f, \quad (17)$$

and the electric charge density is

$$\rho_e = -\frac{1}{r} (rb')' - 4\kappa \frac{a'}{r}, \quad (18)$$

derived from the Gauss Law equation.

The definitions of both angular momentum and electric charge depend on the 0th component of the Abelian connection A_0 , which in turn depends on the presence of the Chern-Simons term.

The energy, the angular momentum and the electric charge are then given by

$$E = 2\pi \int_0^\infty r \mathcal{E} dr, \quad (19)$$

$$J = 2\pi \int_0^\infty r \mathcal{J} dr = 4\pi\kappa (a_\infty^2 - n^2). \quad (20)$$

$$Q_e = 2\pi \int_0^\infty r \rho_e dr = 8\pi\kappa (n - a_\infty). \quad (21)$$

In contrast to our conclusion in Sec. IA 2 above, where we excluded the possibility of evolving topological charge of the Higgs model analogues, the situation here relating to (20) and (21) is different. This is because the definitions of \mathcal{J} and of ρ_e are independent of the scalar field content, i.e., Skyrme or Higgs. Thus, as long as the Higgs model employed is a p -AH model with $p \geq 2$, which supports solutions with $a_\infty \neq 0$ like a Skyrme model, the evolutions of the spin J , (20), and electric charge Q_e , (21), can be tracked. In Ref. [9], this was carried out in detail and the corresponding evolutions of the energy with respect to Q and J were tracked.

II. $SO(2)$ GAUGED SKYRMIONS ON \mathbb{R}^2 WITH MAXWELL DYNAMICS

The Lagrangian and the static Hamiltonian of the $SO(2)$ gauged planar Skyrme model with Maxwell dynamics are

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu}^2 + \frac{1}{2} \eta^2 |D_\mu \phi^a|^2 - \eta^4 V[\phi^3], \quad \mu = 0, i, \quad (22)$$

$$\mathcal{H} = \frac{1}{4} F_{ij}^2 + \frac{1}{2} \eta^2 |D_i \phi^a|^2 + \eta^4 V[\phi^3], \quad i = 1, 2. \quad (23)$$

Two potential functions $V[\phi^3]$ in (22) and (23) will be considered.

Subjecting the static Hamiltonian (23) to the symmetry (2) and (3), the resulting one-dimensional energy density functional is

$$H = \frac{1}{2} \left[\frac{a'^2}{r} + \eta^2 \left(r f'^2 + \frac{a^2 \sin^2 f}{4r} \right) \right] + \eta^4 V[\cos f]. \quad (24)$$

We study the vortices resulting from the use of the two potentials

$$V = \frac{1}{2} \lambda (1 - \phi^3)^2, \quad (25)$$

$$V = \lambda' (1 - \phi^3). \quad (26)$$

The potential (25), with $\lambda = 1$ leads to the saturation of the topological lower bound of the energy. After imposition of symmetry on (B8) and (B9), the two first-order ODE's are

$$f' = -\frac{a \sin f}{r}, \quad (27)$$

$$\frac{a'}{r} = \eta^2 (1 - \cos f). \quad (28)$$

This is the case studied in Ref. [11]. Here we will consider vortices of models also with $\lambda \neq 1$, by solving the second-order equations of motion. We will seek to find the energy profiles vs the real and positive parameter λ . These profiles display discontinuities and branches.

The reason for considering the other potential, (26), is that it is a rather natural choice resembling the ‘‘pion mass’’ potential of the three-dimensional Skyrme model. (The topological lower bound established (B7) for the potential (25) remains valid in this case.) Clearly with this choice of potential, only solutions to the second-order equations of motion can be sought. A distinguishing feature of these solutions is that they decay exponentially, in contrast to the power decay in the case of potential (25).

A common feature between the solutions of the models with potentials (25) and (26) is, that in both cases the topological charge equals the winding number n , as seen from (12) with $v = 1$.

We are specially interested here in solutions with the ranges of λ which are subject to the lower bounds (B11)–(B12). Following Ref. [26], one can estimate numerically the relative magnitude of the energy per unit vorticity n .

A. The solutions

Assuming power behavior in the $r \ll 1$ region, we find

$$f(r) = \pi + f_n r^{|n|} + o(r^{|n|+2}), \quad (29)$$

$$a(r) = n + a_2 r^2 + o(r^4), \quad (30)$$

where n is a nonzero integer.

At infinity, the value $f(\infty) = 0$ is fixed by (9). How $f(r)$ decays at infinity depends on the potential employed.

a1: The choice of potential $V = \frac{1}{2} \lambda (1 - \phi^3)^2$, (25), leads to a power decay, given by




FIG. 1. E vs λ for (M) vortices with $n = 1, 2, 3$, $\eta = 1/\sqrt{2}$ for the quadratic potential.

$$f(r) = \sqrt{\frac{2}{\lambda}} \frac{1}{r} + \dots, \quad (31)$$

$$a(r) = \frac{a_1}{r^{\sqrt{1+2/\lambda}-1}} + \dots, \quad (32)$$

for $a_\infty = 0$, and

$$f(r) = \frac{f_1}{r^{a_\infty}} + \dots, \quad (33)$$

$$a(r) = a_\infty + \frac{1}{4} \eta^2 \frac{f_1^2}{a_\infty - 1} \frac{1}{r^{2(a_\infty-1)}} + \dots, \quad (34)$$

for $a_\infty \neq 0$. For the quadratic potential (25) our numerical resolution is lower. We have obtained solutions for $\lambda < 1$ which have $a_\infty = 0$ and self-dual solution for $\lambda = 1$ which have $a_\infty \neq 0$ [11]. The dependence of the energy on λ for $\lambda \leq 1$ is shown in Fig. 1. We see that in that region the interaction between vortices is attractive. For $\lambda > 1$, the numerical schemes seem to indicate that the solutions might not exist, since their accuracy degenerates quite fast as higher resolution is required. Those solutions do exist if a Skyrme term is added to the theory, though. One can produce solutions with nonvanishing a_∞ for any value of λ when the Skyrme term is present, but as soon as one tries to remove it the convergence of the numerical schemes ceases, indicating that the solutions in the limit without the Skyrme term might not exist for $\lambda > 1$.

a2: The choice of potential $V = \lambda' (1 - \phi^3)$, (26), results in the exponential decay, given by

$$f(r) = \frac{f_1}{\sqrt{r}} e^{-\eta\sqrt{\lambda}r} + \dots, \quad (35)$$

$$a(r) = a_\infty + \frac{a_\infty f_1^2}{4\lambda' r} e^{-2\eta\sqrt{\lambda}r} + \dots. \quad (36)$$




FIG. 2. Left panel: E vs λ' for (M) vortices with $n = 1, 2, 3$, $\eta = 1/\sqrt{2}$ for the ‘pion-mass’ potential. Right panel: a_∞ vs λ' for the same vortices.

In both (34) and (36), the value of a_∞ , on the other hand, is generally fixed by the numerical process.

In Fig. 2 (left), we show the energy per vorticity E/n for the vortices with the pion-mass potential (26). We observe there is a limiting value of the coupling constant of the potential given by

$$\lambda' = \frac{n-1}{n}. \quad (37)$$

For $n = 1$, the limiting value coincides with the minimal value of λ' (zero) but for $n > 1$ it corresponds to the maximal value of λ' . For these solutions a_∞ ranges for 0 to n . The dependence of a_∞ on λ' is exhibited in Fig. 2 (right).

III. $SO(2)$ GAUGED SKYRMIONS WITH CHERN-SIMONS DYNAMICS IN 2+1 DIMENSIONS

Apart from its intrinsic interest, the main motivation for considering models whose Abelian field dynamics is described by Chern-Simons term in the Lagrangian, is that it enables the choice of a potential in which the constant v may take values $v \neq 1$. This potential is given in Appendix C by (C8) and the topological lower bound resulting from the Belavin inequality (C9) remains valid for any value of v . In the present work, we opt for the simplest such value $v = 0$. As a result, the topological charge defined by (A10) in Appendix A, with $v = 0$, results in the expression (14) for the topological charge.

Equation (14) enables the tracking the evolution of the topological charge, by tracking the evolving values of a_∞ that parametrize the solutions. We will find that for the model in this section the values of a_∞ do not result in the dissipation of the topological charge, so we defer this question to the study of the model with both Chern-Simons and Maxwell dynamics in the next section. In the present

section, we restrict our considerations to the intrinsic properties of the solutions.

The Lagrangian and its static Hamiltonian studied in this section are defined by (C1) and (C4), respectively, in both of which the potential $V[\phi]$ is given by (C12). The static (energy) density employed in this model is (C4), which is seen from (C9) to be bounded from below by the topological charge. The static Hamiltonian is

$$\begin{aligned} \mathcal{H}_{(\lambda)} = & 4 \left(\frac{\kappa}{\eta} \right)^2 \frac{F_{ij}^2}{|\phi^\alpha|^2} + \frac{1}{2} \eta^2 |D_i \phi^\alpha|^2 \\ & + \frac{\lambda}{32} \left(\frac{\eta^3}{\kappa} \right)^2 |\phi^\alpha|^2 (\phi^3 - v)^2. \end{aligned} \quad (38)$$

What is different here, contrasting with the Maxwell case above, is that the value of the real constant v is not restricted. This is obvious since the potential term in (38) would vanish at infinity for any value of v since $|\phi^\alpha|^2(\infty) = 0$.

Another striking difference here, contrasting with the Maxwell case above, is that in this case the second-order equations of motion are solved by the first-order equations (C10) and (C11) for any value of v for the self-dual case $\lambda = 1$. As seen from (A10), only when the choice $v = 1$ is made, is the topological charge equal to the winding number n . For all other values of $v \neq 1$ the topological charge of the gauged system departs from n and depends on a_∞ .

After imposition of symmetry using the Ansatz (2) and (3), (38) reduces to the one-dimensional density

$$\begin{aligned} H_{(\lambda)} = & 8 \left(\frac{\kappa}{\eta} \right)^2 \frac{a^2}{r \sin^2 f} + \frac{1}{2} \eta^2 \left(r f'^2 + \frac{a^2 \sin^2 f}{4r} \right) \\ & + \frac{\lambda}{32} \left(\frac{\eta^3}{\kappa} \right)^2 r \sin^2 f (\cos f - v)^2. \end{aligned} \quad (39)$$




FIG. 3. Left panel: E vs a_∞ for (CS) vortices with $n = 1, 2, 3, 4, \eta = 1, \kappa = 1$ and $\lambda = 1$. Right panel: b_∞ vs a_∞ for (CS) vortices with $n = 4, \eta = 1, \kappa = 1$ and $\lambda = 1$.

When $\lambda = 1$ the second-order equations of motion are solved by the first-order equations

$$f' = \frac{a \sin f}{r}, \quad (40)$$

$$\kappa a' = -2rU \sin^2 f, \quad (41)$$

where the function U in (41), given by (C7), reduces to

$$U = \frac{1}{8} \left(\frac{\eta}{\kappa} \right) (\cos f - v). \quad (42)$$

These self-dual solutions were constructed in [16]. Here, we seek solutions to the second-order equations, with $\lambda \neq 1$.

We will restrict our quantitative considerations to the models with $v = 0$ since this is the simplest case of $v \neq 1$, this being the case of interest from the viewpoint of tracking the evolution of the topological charge. The model with $v = 1$ has been studied in Ref. [27].⁸

Proceeding with the study of the $v = 0$ model, the expansions at the origin for this solutions read

$$a(r) = n + \frac{\eta^2 b_0 f_n^2}{8(n+1)\kappa} r^{2(n+1)} + \dots, \quad (43)$$

$$b(r) = b_0 + \frac{\eta^2 f_n^2}{8\kappa} r^{2n} + \dots, \quad (44)$$

⁸The emphasis there was in the dependence of the energy per unit vortex number n on the parameter λ . In comparison with the Abelian Higgs model, studied in Ref. [26], the results were qualitatively similar. Unlike the latter, however, where the solutions had boundary value $a_\infty = 0$, in this model a continuous family of solutions parametrized by $a_\infty \neq 0$ was constructed.

$$f(r) = \pi + f_n r^n + \dots. \quad (45)$$

Concerning the asymptotic behavior, the situation is more complicated. We have to distinguish between two cases: $a_\infty > n$ and $a_\infty < n$ ($a_\infty = n$ is a singular case). For $a_\infty > n$, we have a power decay in a $1/r$ resulting in

$$a(r) = a_\infty - \frac{1}{128} \frac{\lambda}{a_\infty - 1} \frac{\eta^6}{\kappa^3} f_1^2 \frac{1}{r^{2(a_\infty - 1)}} + \dots, \quad (46)$$

$$b(r) = \frac{1}{16} \frac{\eta^3}{\kappa^2} \lambda - \frac{1}{8\kappa} \eta^2 f_1^2 \frac{1}{r^{2a_\infty}} + \dots, \quad (47)$$

$$f(r) = \frac{f_1}{r^{a_\infty}} + \dots. \quad (48)$$

For $a_\infty < n$, the functions decay exponentially. The corresponding expressions are quite involved so we will not reproduce them here.

This change in the asymptotic behavior is reflected on the dependence of the energy on a_∞ . In Fig. 3 (left), we represent the energy E vs the parameter a_∞ for $SO(2)$ gauged Skyrmons with CS coupling constant $\kappa = 1, \lambda = 1$ and vorticities $n = 1, 2, 3, 4$. For $a_\infty > n$ the energy becomes $E = 2\pi(n + a_\infty)$, namely, it saturates the topological lower bound. Whereas for $a_\infty < n$ the dependence of the energy on a_∞ is different. Such solutions, which may be considered to be sphalerons, were not sought in [27] in the case of the model with $v = 1$.

Although the transitions from $a_\infty < n$ to $a_\infty > n$ look continuous at the level of the energy (with change in the slope at $a_\infty = n$), it is not so in fact. If one looks at the quantity b_∞ , we can observe a jump at $a_\infty = n$. We show this in Fig. 3 (right), for the set of solutions with $n = 4$




FIG. 4. E vs λ for (CS) vortices with $n = 1, 2, 3$, $\eta = 1$, $\kappa = 1$ and $a_\infty = 0$.

shown in Fig. 3 (left). Notice that above $a_\infty = n$, b_∞ is constant [and equal to the leading term of Eq. (47)].

We finally show the dependence of the energy on λ . That behavior depends on the value of a_∞ chosen. For $a_\infty < 1$, we have the standard pattern where the curves for several vorticities do not cross each other, resulting in an attractive phase (see Fig. 4 for $a_\infty = 0$). However, for $a_\infty > 1$ solutions with $n < a_\infty$ end up crossing other curves. We see that the situation here, for the pure Chern-Simons model with $v = 0$ is much less transparent than in the corresponding model with $v = 1$ [27], and in the corresponding Higgs model [26]. We do not pursue this line further.

IV. $SO(2)$ GAUGED SKYRMIONS WITH MAXWELL-CHERN-SIMONS DYNAMICS IN 2+1 DIMENSIONS

While in the previous two sections our attention was focused on the energy profiles vs λ , the dimensionless constant parametrizing the coupling of the potential, here instead we focus on the evolution of the topological charge in a given Maxwell-Chern-Simons-Skyrme theory. The topological lower bound for this model follows immediately from the topological lower bound (C9) given in Appendix C for the model with Chern-Simons dynamics only, since adding the Maxwell term to the Lagrangian does not invalidate the ensuing topological lower bound (C9). In particular as elsewhere in this work, we will select the simplest option $v \neq 1$, namely $v = 0$.

As can be seen from the Lagrangian (C1), it does not contain a quartic (“Skyrme”) kinetic term. We add such a term nonetheless, so that in the gauge decoupling limit a topological lower bound persists. The soliton of the ungauged system is stabilized by the “baryon number,”

the departure of the topological charge from which is what we aim to track.

We will find out, in fact, that the evolution of the solutions can result in the total annihilation of the topological charge.

The model is described by the Lagrangian

$$\begin{aligned} \mathcal{L}_{(\lambda)} = & -\frac{1}{4}F_{\mu\nu}^2 + \kappa e^{\lambda\mu\nu} A_\lambda F_{\mu\nu} - \frac{1}{8}\tau |D_{[\mu}\phi^a D_{\nu]}\phi^b|^2 \\ & + \frac{1}{2}\eta^2 |D_\mu\phi^a|^2 - \eta^4 V[\phi], \end{aligned} \quad (49)$$

in which the potential $V[\phi]$ is that given by (C8), with $v = 0$.

This choice of $V[\phi]$ is predicated by our desire to avail of the topological lower bound (C9), where the charge density is that given by (A10), or, by (A8). This topological lower bound is not invalidated by adding the Maxwell term to the Lagrangian (C1) which results in the addition of a positive definite contribution to the static density (C2).

While in the absence of the Maxwell term the electric component of the Abelian potential A_0 could be solved using the Gauss Law equation, (C3) in Appendix C, here this is not possible. So in the imposition of symmetry, the function $b(r)$ in (4) appears explicitly in the one-dimensional reduced Lagrangian, which results in

$$\begin{aligned} -r^{-1}L = & \frac{1}{2}\left(\frac{a'^2}{r^2} - b'^2\right) - \frac{2\kappa}{r}[(ab' - ba') - nb'] \\ & + \tau\left(\frac{a^2}{r^2} - b^2\right)f'^2 \sin^2 f \\ & + \frac{1}{2}\eta^2\left[f'^2 + \left(\frac{a^2}{r^2} - b^2\right)\sin^2 f\right] \\ & + \frac{1}{32}\lambda\left(\frac{\eta^3}{\kappa}\right)^2 \sin^2 f \cos^2 f, \end{aligned} \quad (50)$$

in which the potential term is multiplied by the real positive parameter λ , in order to allow for generic values of the potential coupling constant.

The second-order equations of (50) are solve subject to the boundary values

$$\lim_{r \rightarrow 0} f(r) = \pi, \quad \lim_{r \rightarrow 0} a(r) = n, \quad \lim_{r \rightarrow 0} b'(r) = 0, \quad (51)$$

$$\lim_{r \rightarrow \infty} f(r) = 0, \quad \lim_{r \rightarrow \infty} a(r) = a_\infty, \quad \lim_{r \rightarrow \infty} b(r) = b_\infty, \quad (52)$$

where a_∞ is not necessarily zero and b_∞ is a free parameter that allows us to vary the electric charge of the solutions within a concrete model (i.e., choice of the parameters in the Lagrangian). Notice that a_∞ is numerically related to b_∞ .




FIG. 5. b_∞ vs a_∞ for (MCS) vortices with $n = 1$, $\lambda = 1.6$, $\eta = 1$, $\kappa = 1$, and $\tau = 1$.

A. Numerical results

The field equations resulting from (50) cannot be integrated analytically and one has to resort to numerical methods to carry out that task. Given a concrete theory (e.g., fixed κ , τ , η , and λ) and a fixed winding number n , there is just one free parameter to be varied. As said above we use b_∞ as such a parameter, a_∞ being numerically related to b_∞ . In Fig. 5 we show the relation between b_∞ and a_∞ for a generic solution ($n = 1$, $\lambda = 1.6$, $\eta = 1$, $\kappa = 1$, and $\tau = 1$). We observe that b_∞ is bounded while a_∞ seems not to be.⁹ That indicates that there exist solutions with arbitrarily large (positive or negative) values of the topological charge q . The same holds for the electric charge Q_e and the angular momentum J . Concerning the energy E , it diverges as $|a_\infty|$ gets larger and larger. In fact, we know already from our results in Ref. [9] that due to the dependence between a_∞ and b_∞ , the dependence of the mass/energy E on the global charges is nonstandard, namely the dependence on the electric charge Q_e and the angular momentum J displayed, respectively (left and right panels), in Fig. 6.

Let us analyze the effect of the topological charge q on the energy E . In Fig. 7 (left), we represent the energy per unit winding number n vs the topological charge for three values of n for the same parameters as in Fig. 5. For small values of the Skyrme constant τ the minimum of the energy occurs at values of the topological charge around the winding number. However, when higher values of τ are considered the minimum of the energy occurs at values clearly different from the winding number. This is shown in

⁹We have found certain regions in the parameter space where a_∞ could not be increased or decreased arbitrarily. However, it was not clear whether that fact was due to the actual absence of solutions or to a difficulty of the methods used to find them.

Fig. 7 (right), where the minimum of the curve for $n = 1$, $\tau = 10$ is located at $q \approx 0.556$. In the limit of large τ the minimum of the energy occurs at $q = n/2$, as shown in Fig. 8. So the minimum takes place within the range $q \in [n/2, n]$. Concerning the stability of these solutions, one would be tempted to state that most stable solution would correspond to that with less energy, which according to Fig. 7 (right) does not possess integer topological charge, in general.

V. SUMMARY AND DISCUSSION

We have made a systematic study of $SO(2)$ gauged planar Skyrmions in $2 + 1$ dimensions. The models in question feature (a) Maxwell (only), (b) Chern-Simons (only) and (c) (both) Maxwell and Chern-Simons dynamics. The studies in (a) and (b) are aimed at exposing discontinuities and branchings in the energy profiles of such solutions, which were encountered in the solutions to the $SO(3)$ gauged $O(4)$ Skyrme model on \mathbb{R}^3 . These were shown in Refs. [19,20,21] and are expected to be typical features of all gauged sigma model, in contrast with (gauged) Higgs models where energy profiles are always continuous and exhibit monotonic behavior.

In case (a), pertaining to Maxwell dynamics (only), we have studied the model (a1) with potential (25) that supports self-dual solutions of first-order equations, and the model (a2) with a ‘‘pion mass’’ potential which has solutions to the second-order equations only. The energy profiles of both models exhibit nonmonotonic features.

The solutions we found in the model (a1) are characterized by $a_\infty = 0$, for the coupling constant $\lambda < 1$, $\lambda = 1$ corresponding to the self-dual solution. However, when the model is augmented with the quartic (Skyrme) kinetic term ($\tau > \neq 0$), we find solutions with any λ and $a_\infty \neq 0$. Not surprisingly the quartic term, which has a smoothing effect, becomes singular in the limit $\tau \rightarrow 0$. The energy profiles for $\tau = 0$ are given in Fig. 1, which while they do not exhibit branches and look smooth, they do stop at $\lambda = 1$.

The solutions of the model (a2) with potential (26) exhibit more marked discontinuities in the energy profiles, which range from zero to some finite value of the coupling λ' as shown in Fig. 2 (left) for $n = 1, 2, 3$. The behavior of these profiles is typified by the value of a_∞ , which emerges from the numerical process. These latter are exhibited in Fig. 2 (right). The end point of each of these profiles is observed to coincide with the value of the coupling

$$\lambda' = \frac{n-1}{n}.$$

In case (b), pertaining to Chern-Simons dynamics (only), we have studied only one model. This model, typified by a potential term exhibiting no constant, i.e., $v = 0$ in (38). This choice was made in anticipation of selecting the desired model (c), subsequently studied in Sec. III. The




FIG. 6. Left panel: Energy E vs electric charge Q_e for (MCS) vortices with $n = 1$, $\lambda = 1.6$, $\eta = 1$, $\tau = 1$, and $\kappa = 1, 5, 10$. Right panel: Energy E vs angular momentum J for the same vortices.




FIG. 7. Left panel: E vs q for (MCS) vortices with $n = 1, 2, 3$, $\lambda = 1.6$, $\eta = 1$, $\kappa = 1$, and $\tau = 1$. Right panel: Same for (MCS) vortices with $n = 1$, $\lambda = 1.6$, $\eta = 1$, $\kappa = 1$, and $\tau = 0, 1, 5, 10$.

solutions here present a more complex pattern, occasioned by two different asymptotic behaviors characterized by $a_\infty > n$ and $a_\infty < n$. When $a_\infty > n$, the functions decay with a power behavior, while when $a_\infty < n$ they decay exponentially, solutions with $a_\infty = n$ being singular. It is noteworthy to point out that the exponentially decaying solutions with $a_\infty > n$, the energy has the constant value

$$E = 2\pi(n + a_\infty),$$

as seen from Fig. 3 (left). While the transition from the $a_\infty < n$ to the $a_\infty > n$ solution looks smooth in this plot, it

is in fact discontinuous, as seen from Fig. 3 (right) showing the relation of a_∞ and b_∞ for $n = 4$, where the jump at $n = 4$ is seen.

Concerning the energy profiles vs the coupling λ in this case, (b), we have plotted these in Fig. 4 for the solutions $a_\infty < n$. These E/n profiles are smooth and they do not cross at the self-duality point $\lambda = 1$, hence the system describes solutions exclusively in the attractive phase. In this respect, these solutions differ from those of the vortices of the Abelian Higgs model, where both attractive and repulsive phases occur, and also, from vortices [27] of the Chern-Simons-Skyrme model with $v = 1$ rather than the




FIG. 8. E vs q for (MCS) vortices with $n = 1, 2, 3$, $\lambda = 1.6$, $\eta = 1$, $\kappa = 1$, and $\tau = 1000$.

$v = 0$ model studied here. The situation is different for the $a_\infty > n$ solutions, in which case curves of different vorticity do seem to cross, or at least meet. This completes our first task, namely the investigation of discontinuities and branchings in the energy profiles of gauged (planar) Skyrmions, employing models featuring only Maxwell and only Chern-Simons dynamics, in turn, and, in the absence of the quartic Skyrme kinetic term.

In carrying out our second task, namely the tracking of the topological charge with changing energy, we proceeded to the third case (c). The system studied contains both Maxwell and Chern-Simons dynamics.¹⁰ Such a system was studied in Sec. IV of Ref. [9], albeit with a different (Skyrme) potential term. There, the emphasis was on the nonstandard dependence of the global charges, electric charge and spin, while here, the emphasis is on the evolution of the topological charge and its dependence on the energy. The choice of potential in the present work, is motivated by our desire to influence the topological charge. As can be seen from the definition (A10) of topological charge, if $v = 1$ as, e.g., in the “pion mass” potential, the integral of the charge density will yield the winding number n , as in the gauge-decoupled case. Here, we have adopted a potential featuring the value $v = 0$, which does not force the topological charge to take the value n . This potential is inspired by Chern-Simons dynamics, arrived at via the Belavin inequalities presented in Appendix C.

In case (c) here, we have included also the quartic (Skyrme) kinetic term with coupling strength τ , since we

¹⁰Such systems, (c), studied here and in [9], do not result in energy profiles exhibiting discontinuities and singularities. The solutions are even smoother, when the quartic kinetic Skyrme term is added.

require that our model support solitons [28] in the gauge-decoupling limit.

The mechanism giving rise to the features found in [9], and here, is the same one, namely, the effect of the Chern-Simons dynamics. Technically, this is due to the intertwining of the magnetic and electric functions a and b present in the Chern-Simons density, the asymptotic values a_∞ and b_∞ of which characterize the solutions in any given model. In the present case, the relation between a_∞ and b_∞ is displayed in Fig. 5, which corresponds to Fig. 6 in Ref. [9]. The latter result in the nonstandard dependence of the energy on the global charges featured in Figs. 7 and in 8 in [9]. The model studied in [9] featured the “pion-mass” potential, while the potential used here is (C8) (with $v = 0$) employed in (49). For completeness, we reproduce these results for the system studied here, in Figs. 6 (left) and 6 (right).

The most interesting result in the present work is the dependence of the topological charge on the energy. In Figs. 7 (left), 7 (right), and 8, we observe that the topological charge evolves, taking values different from the “baryon number” n , or the winding number, due to the effect of the gauge field. We observe that in the absence of the quartic kinetic (Skyrme) term ($\tau = 0$) the energetically favored states occur at the value of the topological charge coinciding with the winding number. When the Skyrme term is switched on, the energetically favored states appear at smaller values, down to one half of the winding number for large τ . Also in the limit of large τ , we find a clear asymmetry between positive and negative values of the topological charge q . For large positive values of q the energy becomes independent of the winding number n , whereas for large negative values of q the gap in energy between curves with consecutive values of n does not depend on n ; i.e., the curves are equispaced.

The present work is entirely devoted to the study of gauged Skyrmions in $2 + 1$ dimensions. Of special interest is the influence of Chern-Simons dynamics which results in the nonstandard behavior of the energy on global charges, as reported in [9]. Going further, in the present work we investigate the effect of the Chern-Simons dynamics on the evolution of the topological charge. We expect that these effects are not limited to the $2 + 1$ -dimensional Skyrmions, but they feature also in higher dimensions.

In a recent article [10], $SO(2)$ gauged Skyrmions in both $2 + 1$ and $3 + 1$ dimensions were studied. It was found there that in the $3 + 1$ -dimensional case, where there is no usual¹¹ definition of a Chern-Simons density, these new effects were not observed. In the context of $3 + 1$ dimensions, however, there do exist Chern-Simons densities, namely, the Higgs–Chern-Simons densities (also defined

¹¹Skyrme–Chern-Simons densities, proposed in [13], can be defined in even dimensions, but a systematic (numerical) study of this has not been attempted to date.

in [13]), whose dynamics does result in the nonstandard relation of energy and global charges. This was demonstrated for the $SO(5)$ and $SU(3)$ Yang-Mills–Higgs monopoles in Refs. [29,30]. It is important in this connection, that Higgs–Chern-Simons dynamics cannot lead to the evolution of topological charge observed here for gauged Skyrmions influenced by Chern-Simons dynamics.

ACKNOWLEDGMENTS

We thank Valery Rubakov for valuable comments. We are grateful to Eugen Radu for substantial discussions and collaboration at the early stage of this work. We thank Yasha Shnir for bringing some interesting points to our attention and pointing out some relevant references. F.N.-L. thanks the Dublin Institute of Advanced Studies for hospitality during the performance of parts of this work.

APPENDIX A: TOPOLOGICAL CHARGE OF $SO(2)$ GAUGED $O(3)$ SKYRME SYSTEM ON \mathbb{R}^2

The definition of the topological charge is independent of the dynamics of the gauge field. The definitions given in this Appendix, apply to the following Appendixes B and C, pertaining, respectively, to Maxwell and Chern-Simons dynamics.

The topological charge for the $SO(N)$ gauged $O(D+1)$ Skyrme system on \mathbb{R}^D , with $2 \leq N \leq D$, is given in [5] and is reviewed in Appendix B of Ref. [13]. In the case $D=2$, this definition coincides with that of Ref. [11]. Here, we revisit the $D=2$ case for completeness and with more detail.

The $O(3)$ sigma model(s) are described by the scalar field ϕ^a , $a = \alpha, 3$; $\alpha = 1, 2$, which is subject to the constraint

$$|\phi^a|^2 = 1. \quad (\text{A1})$$

From the outset, we restrict our attention strictly to topologically stable Skyrme solutions, which are characterized by the asymptotics,

$$\lim_{r \rightarrow 0} \phi^3 = \mp 1, \quad \lim_{r \rightarrow \infty} \phi^3 = \pm 1. \quad (\text{A2})$$

Moreover, in what follows, we will restrict to the upper signs in (A2), for simplicity.

The prescription of gauging employed is

$$D_i \phi^\alpha = \partial_i \phi^\alpha + A_i (\varepsilon \phi)^\alpha, \quad (\text{A3})$$

$$D_i \phi^3 = \partial_i \phi^3, \quad (\text{A4})$$

such that the component ϕ^3 is gauge invariant.

The winding number density prior to gauging is

$$q_0 = \varepsilon_{ij} \varepsilon^{abc} \partial_i \phi^a \partial_j \phi^b \phi^c, \quad (\text{A5})$$

which is essentially total divergence but is gauge variant, while the density

$$q_G = \varepsilon_{ij} \varepsilon^{abc} D_i \phi^a D_j \phi^b \phi^c, \quad (\text{A6})$$

is gauge invariant but is not a total divergence. Thus, neither qualifies as a topological charge density.

However, (A5) and (A6) are related through

$$q_G = q_0 + 2\varepsilon_{ij} \partial_i (\phi^3 A_j) - \varepsilon_{ij} \phi^3 F_{ij}. \quad (\text{A7})$$

Collecting the two gauge invariant terms together, and the two total divergence terms together, we propose two equivalent definitions for a topological charge density

$$\tilde{q} = q_0 + 2\varepsilon_{ij} \partial_i [(\phi^3) A_j] \quad (\text{A8})$$

$$= q_G + \varepsilon_{ij} (\phi^3) F_{ij}. \quad (\text{A9})$$

The first line, (A8), is manifestly a total divergence while the second line, (A9) is manifestly gauge invariant. Both are gauge invariant and total divergence, and are candidates for topological charge densities.

It is clear that the definitions (A8) and (A9) can be extended by adding to (or subtracting from) each, the first Chern-Pontryagin density¹² $\varepsilon_{ij} F_{ij}$, which is both total divergence and gauge invariant. Subtracting $\varepsilon_{ij} F_{ij} = 2\varepsilon_{ij} \partial_i A_j$, with a real constant coefficient v , results in the most general definition of the topological charge

$$q = q_0 + 2\varepsilon_{ij} \partial_i [(\phi^3 - v) A_j] \quad (\text{A10})$$

$$= q_G + \varepsilon_{ij} (\phi^3 - v) F_{ij}. \quad (\text{A11})$$

Our considerations are restricted strictly to Skyrme asymptotics (A2), mostly to the upper signs there, throughout. For the purpose of establishing Belavin inequalities, definition (A11) for the topological charge will be employed, while (A10) will be employed to evaluate the topological charge itself.

1. Topological charge of $SO(2)$ gauged Higgs system on \mathbb{R}^2

Here, we return to the definition of the topological charge of the $SO(2)$ gauged Higgs scalar, in a manner analogous to what was described above for the $O(3)$ Skyrme system. The systematic way to achieve this is to subject a Chern-Pontryagin (CP) density to dimensional descent to the desired dimension, in this case \mathbb{R}^2 . (See [4] and references therein.)

This alternative approach was proposed in [6] for the Higgs systems on \mathbb{R}^2 and \mathbb{R}^3 . Here, we present only the case in \mathbb{R}^2 .

¹²This can be done in every even-dimension $D = 2n$ with the n th Chern-Pontryagin density.

The covariant derivative of the complex Higgs scalar $\varphi = \phi^1 + i\phi^2$, written for the real doublet ϕ^a , $a = 1, 2$ is

$$D_i\phi^a = \partial_i\phi^a + A_i(\varepsilon\phi)^a. \quad (\text{A12})$$

In analogy with (A5) and (A6), we define the winding number density prior to gauging

$$q_0 = \varepsilon_{ij}\varepsilon^{ab}\partial_i\phi^a\partial_j\phi^b, \quad (\text{A13})$$

which is essentially total divergence but is gauge variant, and its covariantized version,

$$q_G = \varepsilon_{ij}\varepsilon^{ab}D_i\phi^aD_j\phi^b, \quad (\text{A14})$$

which is gauge invariant but is not a total divergence. Thus, neither qualifies as a topological charge density.

The relation between q_0 and q_G , analogous with (A7) is

$$q_G = q_0 - \varepsilon_{ij}\partial_i(|\phi^a|^2A_j) + \frac{1}{2}\varepsilon_{ij}F_{ij}|\phi^a|^2, \quad (\text{A15})$$

from which we deduce the two definitions of the topological charge density

$$\tilde{q} = q_0 - \varepsilon_{ij}\partial_i[|\phi^a|^2A_j] \quad (\text{A16})$$

$$= q_G - \frac{1}{2}\varepsilon_{ij}|\phi^a|^2F_{ij}. \quad (\text{A17})$$

analogous with (A8)–(A17).

Finally, we add η^2 times the 1st CP density $\frac{1}{2}\varepsilon_{ij}F_{ij}$ to state the final result

$$q^{(n)} = q_0 + \varepsilon_{ij}\partial_i[(\eta^2 - |\phi^a|^2)A_j] \quad (\text{A18})$$

$$= q_G + \frac{1}{2}\varepsilon_{ij}(\eta^2 - |\phi^a|^2)F_{ij}. \quad (\text{A19})$$

Here, the real constant η has dimension $[L^{-1}]$ like the Higgs scalar, unlike the dimensionless constant v above, (A10) and (A11), for the Skyrme scalar.

APPENDIX B: TOPOLOGICAL LOWER BOUND FOR $SO(2)$ GAUGED SKYRMIONS IN 2 + 1 DIMENSIONS WITH MAXWELL DYNAMICS

The static Hamiltonian considered is the $SO(2)$ gauged planar Skyrme scalar with Maxwell dynamics,

$$\mathcal{H}_{(1)} = \frac{1}{4}F_{ij}^2 + \frac{1}{2}\eta^2|D_i\phi^a|^2 + \eta^4V[\phi^3], \quad (\text{B1})$$

where η is a constant¹³ with dimension L^{-1} and $V[\phi^3]$ is the (dimensionless) potential, which will be determined after

¹³ η is introduced for the purpose of keeping track of the dimensions of each term in all the $O(3)$ sigma models considered, and is set equal to a given value in the numerical calculations.

requiring that (B1) is bounded from below by either the topological charge density (A9) or (A11).

The inequality giving rise to the q_G term in (A9) [or (A11)] is

$$|D_i\phi^a - \varepsilon_{ij}\varepsilon^{abc}D_j\phi^b\phi^c|^2 \geq 0 \Rightarrow |D_i\phi^a|^2 \geq q_G, \quad (\text{B2})$$

and to reproduce the other term in (B1), we consider the inequality

$$(\varepsilon_{ij}F_{ij} - \eta^2U)^2 \geq 0 \Rightarrow F_{ij}^2 + \frac{1}{2}\eta^4U^2 \geq \eta^2\varepsilon_{ij}F_{ij}U, \quad (\text{B3})$$

for some real gauge invariant function $U[\phi^3]$ to be determined via the inequalities (B2) and (B3), which in turn imply the following Belavin inequality,

$$\frac{1}{4}F_{ij}^2 + \frac{1}{2}\eta^2|D_i\phi^a|^2 + \frac{1}{8}\eta^4U^2 \geq \frac{1}{2}\eta^2\left[q_G + \frac{1}{2}\varepsilon_{ij}F_{ij}U\right]. \quad (\text{B4})$$

Requiring that the right-hand side of (B4) to be identified with the topological charge density (A11) fixes $U[\phi^3]$ as

$$U = 2(\phi^3 - v). \quad (\text{B5})$$

Finally, identifying the left-hand side of (B4) with the static Hamiltonian (B1) fixes the potential $V = \frac{1}{8}U^2$,

$$V[\phi^3] = \frac{1}{2}(\phi^3 - v)^2. \quad (\text{B6})$$

So far, the value of the constant v is not fixed. It is clear that with (the upper sign of) Skyrme asymptotics (A2), the energy integral resulting from the left-hand side of (B4) will diverge except when one chooses $v = 1$.

The topological charge is conveniently evaluated using the expression (A10), which in this case will get its value from the integral of the first term q_0 only, since the second term vanishes at infinity by virtue of (A2) (upper sign). Thus, the topological charge of these vortices is equal to the “baryon number” of the planar Skyrme in the gauge-decoupled limit.

The lower bound on the static Hamiltonian (B1) is now

$$\mathcal{H}_{(1)} = \frac{1}{4}F_{ij}^2 + \frac{1}{2}\eta^2|D_i\phi^a|^2 + \frac{1}{2}\eta^4(\phi^3 - 1)^2 \geq \frac{1}{2}\eta^2q, \quad (\text{B7})$$

which is saturated when both inequalities (B2) and (B3) are themselves saturated, resulting in the two first-order equations,

$$D_i\phi^a = \varepsilon_{ij}\varepsilon^{abc}D_j\phi^b\phi^c, \quad (\text{B8})$$

$$F_{ij} = \eta^2 \varepsilon_{ij} (\phi^3 - 1), \quad (\text{B9})$$

which solve the second-order equations.

This lower bound is not invalidated if the potential (B6) is replaced by another potential which is everywhere (within the acceptable range of values [$\phi^3(0) = -1, \phi^3(\infty) = 1$]) larger than (B6). A natural example is the ‘‘pion mass’’ potential $V = 1 - \phi^3$.

More interesting, however, is the choice

$$V[\phi^3] = \frac{1}{2} \lambda (\phi^3 - 1)^2, \quad \lambda > 0. \quad (\text{B10})$$

Denoting the resulting static Hamiltonian defined in terms of the potential (B10) as $\mathcal{H}_{(\lambda)}$, the lower bound corresponding to (B7) now reads

$$\mathcal{H}_{(\lambda)} > \frac{1}{2} \eta^2 Q, \quad \lambda > 1, \quad (\text{B11})$$

$$> \frac{1}{2} \eta^2 \lambda Q, \quad \lambda < 1, \quad (\text{B12})$$

following the analysis given in Ref. [26] for the Abelian Higgs model.

1. Topological lower bound for the Maxwell-Higgs case

Our reason for considering the Higgs vortices analogous with the Skyrme vortices is to emphasize the marked difference between these. We have in mind here the usual Abelian Higgs model whose static Hamiltonian is

$$\mathcal{H} = \frac{1}{4} F_{ij}^2 + \frac{1}{2} |D_i \phi^a|^2 + \frac{1}{8} (\eta^2 - |\phi^a|^2)^2, \quad (\text{B13})$$

where the constant¹⁴ η is the VEV of the Higgs field.

The inequalities analogous with (B2) and (B3) in this case are

$$|D_i \phi^a - \varepsilon_{ij} \varepsilon^{ab} D_j \phi^b|^2 \geq 0 \Rightarrow |D_i \phi^a|^2 \geq Q_G, \quad (\text{B14})$$

$$(\varepsilon_{ij} F_{ij} - U)^2 \geq 0 \Rightarrow 2F_{ij}^2 + U^2 \geq 2\varepsilon_{ij} F_{ij} U. \quad (\text{B15})$$

Combining (B14) and (B15), and insisting that the resulting Belavin inequality be

$$\frac{1}{4} F_{ij}^2 + \frac{1}{2} |D_i \phi^a|^2 + \frac{1}{8} (\eta^2 - |\phi^a|^2)^2 \geq \frac{1}{2} Q^{(n)}, \quad (\text{B16})$$

$Q^{(n)}$ given by (A19) results in

$$U = (\eta^2 - |\phi^a|^2). \quad (\text{B17})$$

¹⁴ η in Higgs models is not employed as an artifact like in the sigma models.

It is clear that setting $\eta = 0$, namely by opting for the definition (A17) for the topological charge, the potential $V = U^2$ results in diverging energy since the covariant constancy of the Higgs field implies that its magnitude vanishes at infinity.

APPENDIX C: TOPOLOGICAL LOWER BOUND FOR $SO(2)$ GAUGED SKYRMIONS IN 2+1 DIMENSIONS WITH CHERN-SIMONS DYNAMICS

The topological lower bound on the energy of the static solutions to the Lagrangian,

$$\mathcal{L}_{(1)} = \kappa \varepsilon^{\lambda\mu\nu} A_\lambda F_{\mu\nu} + \frac{1}{2} \eta^2 |D_\mu \phi^a|^2 - \eta^4 V[\phi], \quad (\text{C1})$$

$$\mu = 0, i \quad i = 1, 2,$$

are sought. Here, κ is the coupling strength of the Chern-Simons (CS) term and like η has dimension [L^{-1}]. The gauging prescription is that of the Maxwell case, (5) and (6), such that the static Hamiltonian $2T_{00}$ of (C1) is expressed as

$$\mathcal{H} = \frac{1}{2} \eta^2 (|D_i \phi^a|^2 + A_0^2 |\phi^a|^2) + \eta^4 V[\phi], \quad (\text{C2})$$

$$a = \alpha, 3; \quad \alpha = 1, 2.$$

The Gauss Law equation (0-component of the Maxwell equation) is

$$A_0 = -\frac{2\kappa}{\eta^2 |\phi^a|^2} \varepsilon_{ij} F_{ij}. \quad (\text{C3})$$

Substituting (C3) in (C2), we have the final form of the static Hamiltonian density:

$$\mathcal{H}_{(1)} = 4 \left(\frac{\kappa}{\eta} \right)^2 \left(\frac{F_{ij}^2}{|\phi^a|^2} \right) + \frac{1}{2} \eta^2 |D_i \phi^a|^2 + \eta^4 V[\phi], \quad (\text{C4})$$

$$a = \alpha, 3; \quad \alpha = 1, 2.$$

So far, the potential $V[\phi]$ is not specified. This can be done once the Belavin inequalities that give the lower bound on the energy, namely on the integral of (C4), are stated. This depends on the definition of the topological charge density.

The inequality giving rise to the Q_G term in (A9) [or (A11)] is of course (B2), again. To account for the term $F_{ij}^2/|\phi^a|^2$ in (C4), consider the inequality

$$\left(\frac{\kappa}{\eta |\phi^a|} \varepsilon_{ij} F_{ij} - \eta^2 |\phi^a| U \right)^2 \geq 0$$

$$\Rightarrow 2 \left(\frac{\kappa}{\eta} \right)^2 \left(\frac{F_{ij}^2}{|\phi^a|^2} \right) + \eta^4 |\phi^a|^2 U^2 \geq 2\kappa \eta \varepsilon_{ij} F_{ij} U. \quad (\text{C5})$$

Combining (B2) and (C5), we have the Belavin inequality:

$$4\left(\frac{\kappa}{\eta}\right)^2 \frac{F_{ij}^2}{|\phi^\alpha|^2} + \frac{1}{2}\eta^2 |D_i \phi^a|^2 + 2\eta^4 |\phi^\alpha|^2 U^2 \geq \frac{1}{2}\eta^2 \left[\varrho_G + 8\left(\frac{\kappa}{\eta}\right) \varepsilon_{ij} F_{ij} U \right]. \quad (\text{C6})$$

Requiring that the expression in the square brackets on the right-hand side of (C6) be the topological charge q , given by (A11), fixes U as

$$U = \frac{1}{8} \left(\frac{\eta}{\kappa}\right) (\phi^3 - v), \quad (\text{C7})$$

and identifying the left-hand side of (A11) with the static Hamiltonian density (C4) results in the following potential $V = 2|\phi^\alpha|^2 U^2$,

$$V[\phi] = \frac{1}{32} \left(\frac{\eta}{\kappa}\right)^2 |\phi^\alpha|^2 (\phi^3 - v)^2. \quad (\text{C8})$$

So far, the value of the constant v in (C8) is not fixed. In contrast with the Maxwell dynamics above, here the energy integral will not diverge if the choice $v = 1$ is not made. This is seen immediately by inspecting (C8), which must decay at infinity since

$$\lim_{r \rightarrow \infty} |\phi^\alpha|^2 = 0,$$

for any value of v . The question as to what interesting choices for v can be made depends on the detailed analysis of the solutions.

What is important here is that when calculating the topological charge by evaluating the integral of (A10), the result will not be limited to the winding number, namely the integral of ϱ_0 , except when the choice $v = 1$ is made. This is in contrast to the case of Maxwell dynamics considered above.

The final result is the Belavin inequality,

$$\mathcal{H}_{(1)} = 4\left(\frac{\kappa}{\eta}\right)^2 \frac{F_{ij}^2}{|\phi^\alpha|^2} + \frac{1}{2}\eta^2 |D_i \phi^a|^2 + \frac{1}{32} \left(\frac{\eta^3}{\kappa}\right)^2 |\phi^\alpha|^2 (\phi^3 - v)^2 \geq \frac{1}{2}\eta^2 q, \quad (\text{C9})$$

which is saturated if the two inequalities (B2) and (C5) are saturated, i.e., when the first-order equations,

$$\varepsilon_{ij} D_i \phi^a = \varepsilon^{abc} D_j \phi^b \phi^c, \quad (\text{C10})$$

$$8\kappa^2 \varepsilon_{ij} F_{ij} = \eta^4 |\phi^\alpha|^2 (\phi^3 - v), \quad (\text{C11})$$

are satisfied, which solve the second-order equations.

Again, as in the Maxwell case above, the lower bound (C9) is not invalidated if the potential (C8) is replaced by

$$V[\phi] = \frac{1}{32} \lambda \left(\frac{\eta}{\kappa}\right)^2 |\phi^\alpha|^2 (\phi^3 - v)^2, \quad \lambda > 0. \quad (\text{C12})$$

Denoting the static Hamiltonian defined in terms of the potential (C12) as $\mathcal{H}_{(\lambda)}$, the lower bound corresponding to (C9) now reads

$$\begin{aligned} \mathcal{H}_{(\lambda)} &> \frac{1}{2}\eta^2 q, & \lambda > 1, \\ &> \frac{1}{2}\eta^2 \lambda q, & \lambda < 1, \end{aligned}$$

formally the same as (B11) and (B12).

1. Topological lower bound for the Chern-Simons–Higgs case

Again, we consider the Higgs analogue of the Chern-Simons–Skyrme vortices discussed above, to emphasize the essential difference between them.

Starting with the Lagrangian

$$\begin{aligned} \mathcal{L} &= \kappa \varepsilon^{\lambda\mu\nu} A_\lambda F_{\mu\nu} + \frac{1}{2}\eta^2 |D_\mu \phi^a|^2 - V[|\phi^\alpha|^2], \\ \mu &= 0, i & i = 1, 2, & a = 1, 2, \end{aligned} \quad (\text{C13})$$

one finds from the Gauss Law equation that

$$A_0 = -\frac{2\kappa}{|\phi^\alpha|^2} \varepsilon_{ij} F_{ij}, \quad (\text{C14})$$

and thus the static Hamiltonian can be expressed as

$$\mathcal{H} = 4\kappa^2 \left(\frac{F_{ij}^2}{|\phi^\alpha|^2}\right) + \frac{1}{2} |D_i \phi^a|^2 + V. \quad (\text{C15})$$

Here, in addition to the inequality (B14), one employs

$$\begin{aligned} \left(\frac{\kappa}{|\phi^\alpha|} \varepsilon_{ij} F_{ij} - |\phi^\alpha| U\right)^2 &\geq 0 \\ \Rightarrow 2\kappa^2 \left(\frac{F_{ij}^2}{|\phi^\alpha|^2}\right) + |\phi^\alpha|^2 U^2 &\geq 2\kappa \varepsilon_{ij} F_{ij} U. \end{aligned} \quad (\text{C16})$$

Combining (B14) and (C16), one ends up with the Belavin inequality,

$$\begin{aligned} 4\kappa^2 \frac{F_{ij}^2}{|\phi^\alpha|^2} + \frac{1}{2} |D_i \phi^a|^2 + 2|\phi^\alpha|^2 U^2 \\ \geq \frac{1}{2}\eta^2 \left[\varrho_G + 8\left(\frac{\kappa}{\eta}\right) \varepsilon_{ij} F_{ij} U \right], \end{aligned} \quad (\text{C17})$$

and insisting that the right-hand side of (C17) be equal to the topological charge density (A11) fixes U as

$$U = \frac{1}{16\kappa} (\eta^2 - |\phi^\alpha|^2). \quad (\text{C18})$$

It follows that the potential V is now

$$V[|\phi^a|^2] = \frac{2}{16^2 \kappa^2} |\phi^a|^2 (\eta^2 - |\phi^a|^2). \quad (\text{C19})$$

This is known from the work of [17,18].

What must be emphasized here is that setting $\eta = 0$ results in divergent energy since, at infinity, $|\phi^a|^2 \rightarrow \eta^2$, in contrast with the Skyrme case above where $|\phi^a|^2 \rightarrow 0$, which allows the choice $v = 0$ there.

-
- [1] H. B. Nielsen and P. Olesen, *Nucl. Phys.* **B61**, 45 (1973).
 [2] G. 't Hooft, *Nucl. Phys.* **B79**, 276 (1974).
 [3] A. M. Polyakov, *Pis'ma Zh. Eksp. Teor. Fiz.* **20**, 430 (1974) [*JETP Lett.* **20**, 194 (1974)].
 [4] T. Tchrakian, *J. Phys. A* **44**, 343001 (2011).
 [5] D. H. Tchrakian, *Lett. Math. Phys.* **40**, 191 (1997).
 [6] T. Tchrakian, in *From Integrable Models to Gauge Theories*, edited by V. G. Gurzadyan *et al.* (World Scientific, Singapore, 2002), p. 283–291.
 [7] A. Samoilenka and Y. Shnir, *Phys. Rev. D* **95**, 045002 (2017).
 [8] A. Samoilenka and Y. Shnir, *Phys. Rev. D* **93**, 065018 (2016).
 [9] F. Navarro-Lérida, E. Radu, and D. H. Tchrakian, *Phys. Rev. D* **95**, 085016 (2017).
 [10] F. Navarro-Lérida, E. Radu, and D. H. Tchrakian, arXiv:1811.09535.
 [11] B. J. Schroers, *Phys. Lett. B* **356**, 291 (1995).
 [12] J. Gladikowski, B. M. A. G. Piette, and B. J. Schroers, *Phys. Rev. D* **53**, 844 (1996).
 [13] D. H. Tchrakian, *J. Phys. A* **48**, 375401 (2015).
 [14] P. K. Ghosh and S. K. Ghosh, *Phys. Lett. B* **366**, 199 (1996).
 [15] K. Kimm, K.-M. Lee, and T. Lee, *Phys. Rev. D* **53**, 4436 (1996).
 [16] K. Arthur, D. H. Tchrakian, and Y.-S. Yang, *Phys. Rev. D* **54**, 5245 (1996).
 [17] J. Hong, Y. Kim, and P. Y. Pac, *Phys. Rev. Lett.* **64**, 2230 (1990).
 [18] R. Jackiw and E. J. Weinberg, *Phys. Rev. Lett.* **64**, 2234 (1990).
 [19] Y. Brihaye, B. Kleihaus, and D. H. Tchrakian, *J. Math. Phys. (N.Y.)* **40**, 1136 (1999).
 [20] Y. Brihaye and D. H. Tchrakian, *Nonlinearity* **11**, 891 (1998).
 [21] B. Kleihaus, D. H. Tchrakian, and F. Zimmerschied, *J. Math. Phys. (N.Y.)* **41**, 816 (2000).
 [22] T. W. Kirkman and C. K. Zachos, *Phys. Rev. D* **24**, 999 (1981).
 [23] J. Burzlaff and F. Navarro-Lérida, *Phys. Rev. D* **82**, 125033 (2010).
 [24] A. Y. Loginov, *Izv. Vuz. Fiz.* **2014**, 103 (2014) [*Russ. Phys. J.* **57**, 1732 (2015)]; A. Y. Loginov, *Zh. Eksp. Teor. Fiz.* **145**, 250 (2014) [*J. Exp. Theor. Phys.* **118**, 217 (2014)].
 [25] A. Samoilenka and Y. Shnir, *Phys. Rev. D* **97**, 045004 (2018).
 [26] L. Jacobs and C. Rebbi, *Phys. Rev. B* **19**, 4486 (1979).
 [27] K. Arthur, *Phys. Lett. B* **385**, 181 (1996).
 [28] R. A. Battye, S. Krusch, and P. M. Sutcliffe, *Phys. Lett. B* **626**, 120 (2005).
 [29] F. Navarro-Lérida, E. Radu, and D. H. Tchrakian, *Int. J. Mod. Phys. A* **29**, 1450149 (2014).
 [30] F. Navarro-Lérida and D. H. Tchrakian, *Int. J. Mod. Phys. A* **30**, 1550079 (2015).