

Adsorption of two-dimensional polymers with two- and three-body self-interactionsNathann T. Rodrigues^{✉*} and Tiago J. Oliveira[†]*Departamento de Física, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais, Brazil*Thomas Prellberg[‡]*School of Mathematical Sciences, Queen Mary University of London, Mile End Road, London, E1 4NS, United Kingdom*Aleksander L. Owczarek[§]*School of Mathematics and Statistics, University of Melbourne, Victoria 3010, Australia*

(Received 3 November 2019; published 27 December 2019)

Using extensive Monte Carlo simulations, we investigate the surface adsorption of self-avoiding trails on the triangular lattice with two- and three-body on-site monomer-monomer interactions. In the parameter space of two-body, three-body, and surface interaction strengths, the phase diagram displays four phases: swollen (coil), globule, crystal, and adsorbed. For small values of the surface interaction, we confirm the presence of swollen, globule, and crystal bulk phases. For sufficiently large values of the surface interaction, the system is in an adsorbed state, and the adsorption transition can be continuous or discontinuous, depending on the bulk phase. As such, the phase diagram contains a rich phase structure with transition surfaces that meet in multicritical lines joining in a single special multicritical point. The adsorbed phase displays two distinct regions with different characteristics, dominated by either single- or double-layer adsorbed ground states. Interestingly, we find that there is no finite-temperature phase transition between these two regions though rather a smooth crossover.

DOI: [10.1103/PhysRevE.100.062504](https://doi.org/10.1103/PhysRevE.100.062504)**I. INTRODUCTION**

Understanding the behavior of self-interacting single polymer chains in a solvent is of fundamental interest from both the theoretical and experimental point of view [1,2]. In general, one can find a rich phase structure. In a good solvent, a polymer exists in a swollen globule state, whereas in a bad solvent the polymer collapses into a partially dense amorphous globule or presents in a fully dense crystal-like structure.

The study of a variety of lattice models of polymers has played a fundamental role in elucidating these phases and phase transitions between them. For example, the coil-globule transition between a swollen coil and a collapsed globule at the so-called θ point has been analyzed in both two and three dimensions [3–5], with theoretically predicted critical exponents [6–8] associated with the transition confirmed by simulations and experiment.

More generally, interacting self-avoiding walks and trails with competing two-body and three-body interactions give rise to phase diagrams, which, in addition to the swollen phase, show both collapsed globule and dense crystal-like phases, with all three phases meeting at a multicritical point [9–11]. A schematic phase diagram is shown in Fig. 1.

Adsorption of a polymer chain onto a surface is another fundamental problem in polymer physics [12–20]. If a polymer is tethered with one end to a sticky surface, then increasing the strength of the surface attraction induces an adsorption transition from a desorbed bulk state to an adsorbed state in which the polymer is bound to the surface. If the polymer is in a swollen bulk phase, then one speaks of a “normal” surface transition, whereas if the polymer in bulk is in the critical θ -state, one speaks of a “special” surface transition, each of which is associated with its own set of critical surface exponents [21]. If the polymer adsorbs from a collapsed coil, one finds a surface-attached globule (which is absent in two-dimensional models, in which the adsorbing surface is modeled by a line).

Recently, there has been renewed interest in the polymer adsorption transition due to the fact that numerical studies pointed at possible nonuniversal behavior of the normal adsorption transition [22–25]. Of interest here is the fact that these works show that the determination of these exponents is rather subtle due to the presence of strong finite-size effects, despite the fact that the phases themselves could be identified in a robust manner.

In this paper, we consider the adsorption transition of a polymer chain with both two-body and three-body interactions. The lattice model considered here is one of self-avoiding trails with on-site interactions on the triangular lattice, which has been studied previously [9]. We now study this model in a half-plane with the trails restricted to start at the origin, and adsorption mediated by on-site attractive interactions at vertices in the boundary line. In addition to the three

*nathan.rodrigues@ufv.br

†tiago@ufv.br

‡t.prellberg@qmul.ac.uk

§owczarek@unimelb.edu.au




FIG. 1. Schematic phase diagram for a single polymer chain in a solvent with two-body and three-body interactions, showing the presence of coil, globule, and fully dense phases. The coil-globule transition is critical, whereas the coil-dense transition is first-order. In two dimensions, the globule-dense transition is expected to be continuous.

bulk phases, we find an adsorbed phase. All phases meet each other pairwise at transition surfaces, which meet each other in multicritical lines joining in a single special multicritical point. The analysis is made somewhat more difficult by the fact that the adsorbed phase shows two distinct regions with different characteristics, dominated by either single or double layer adsorbed ground states.

We point out that an analogous scenario has been analyzed using the bond-fluctuation model, which can be seen as a self-avoiding lattice path model in which step lengths different from unity are allowed [26], with a certain set of short-range interactions added. The interplay of interaction and adsorption in such a model has been studied in two [27] and three [28] dimensions.

This paper is organized as follows. In Sec. II we define the model and the quantities used to investigate its thermodynamics properties. Some details on the Monte Carlo methods and simulations are given in Sec. III. Sections IV and V present results, respectively, for the boundary planes and for slices of the three-dimensional parameter space. A summary of the full phase diagram is presented in Sec. VI.

II. MODEL AND QUANTITIES OF INTEREST

A self-avoiding trail (SAT) is defined as a path on a regular lattice, composed of a finite collection of adjacent vertices on the sites of the lattice that are linked by steps along the edges of the lattice, with the restriction that all edges have to be unique. We identify the vertices of this path as monomers and the edges as a sequence of bonds connecting such monomers, in a way that no closed loop is formed. Thereby, the polymer excluded volume interaction is introduced in this model by the restriction of one bond per edge, while more than one monomer can be placed at the same site. In a lattice of




FIG. 2. Illustration of a self-avoiding trail with $n = 22$ steps on the triangular lattice. The solid black circle denotes the origin where the trail is tethered. The Boltzmann weights associated with different site configurations are also indicated.

coordination number q , the maximal number of monomers at one site is $\lfloor q/2 \rfloor$.

Here we are interested in investigating transitions among bulk and adsorbed phases of polymers modeled as self-attracting SATs defined on the half-plane consisting of a triangular lattice limited by a horizontal surface (see Fig. 2). As the triangular lattice has coordination number $q = 6$, bulk sites can be visited by up to three monomers. The coordination number of sites on the surface is 4, so surface sites can have at most two monomers. Self-attraction is included in the SATs by associating energies $-\epsilon_2 \leq 0$ to doubly visited (regardless they are in surface or bulk) and $-\epsilon_3 \leq 0$ to triply visited sites. Moreover, a polymer-surface interaction is introduced by assigning an energy $-\epsilon_s \leq 0$ to each monomer lying on the surface. A configuration for this system is illustrated in Fig. 2. The trail is tethered to the origin, which is located on the surface.

To write down the partition function of the system, we associate Boltzmann weights $\omega_2 = e^{\beta\epsilon_2}$ and $\omega_3 = e^{\beta\epsilon_3}$ to each doubly and triply visited site, respectively, and $\kappa = e^{\beta\epsilon_s}$ to each monomer on the surface, with $\beta = 1/(k_B T)$, where k_B is the Boltzmann constant and T the temperature. The partition function is then given by

$$Z_n(\omega_2, \omega_3, \kappa) = \sum_{m_2, m_3, m_s} C_{\{m_2, m_3, m_s\}}^{(n)} \omega_2^{m_2} \omega_3^{m_3} \kappa^{m_s}, \quad (1)$$

where $C_{\{m_2, m_3, m_s\}}^{(n)}$ is the number of n -step lattice trails with m_2 (m_3) doubly (triply) visited sites and m_s monomers on the surface. The expected value of any thermodynamic quantity Q is then defined by the average

$$\langle Q \rangle(\omega_2, \omega_3, \kappa) = \frac{1}{Z_n} \sum_{\psi_n} \omega_2^{m_2(\psi_n)} \omega_3^{m_3(\psi_n)} \kappa^{m_s(\psi_n)} Q(\psi_n), \quad (2)$$

where the sum runs over all n -step trails ψ_n . We are particularly interested in the energy contribution per step $e_j^{(n)}$ for doubly, triply, and surface contacts:

$$e_j^{(n)} = \frac{\langle m_j \rangle}{n} = \frac{1}{n Z_n} \sum_{m_2, m_3, m_s} m_j C_{\{m_2, m_3, m_s\}}^{(n)} \omega_2^{m_2} \omega_3^{m_3} \kappa^{m_s}, \quad (3)$$

where $j = 2, 3, s$.

A key quantity to identify possible phase transitions is the fluctuation $c_j^{(n)}$ of these energy contributions:

$$c_j^{(n)}(\omega_2, \omega_3, \kappa) = \frac{\langle m_j^2 \rangle - \langle m_j \rangle^2}{n}. \quad (4)$$

Close to a continuous bulk transition the fluctuations $c_2^{(n)}$ and $c_3^{(n)}$ of doubly and triply visited sites, respectively, are expected to follow the scaling law

$$c_i^{(n)} \sim n^{2\phi_b - 1} h(\tau n^{\phi_b}), \quad (5)$$

where ϕ_b is the crossover exponent related to bulk transitions, $\tau \equiv T - T_c$ is the temperature relative to the bulk transition temperature T_c , and $h(\cdot)$ is a scaling function. Scaling around the bulk transition is discussed in [2], see also [3,29]. Further analysis can be found in [30].

The scaling theory around the adsorption transition was developed in [31]. For an adsorption transition, the internal surface energy $e_s^{(n)}$ is the order parameter. Close to the adsorption point the surface energy is expected to behave as

$$e_s^{(n)} \sim n^{\phi_s - 1} f(\tau n^{1/\delta}), \quad (6)$$

where $\tau \equiv T - T_a$ is the temperature relative to the adsorption temperature T_a , ϕ_s is a critical exponent, $1/\delta$ is the crossover exponent associated with the adsorption transition, and $f(\cdot)$ is a scaling function.

To analyze the phase transitions, we need to introduce several quantities. The exponent $1/\delta$ can be obtained from the quantity

$$\Gamma_n = \frac{\langle m_s^2 \rangle - \langle m_s \rangle^2}{\langle m_s \rangle}, \quad (7)$$

whose maximum in curves of Γ_n as function of T behaves as

$$\Gamma_{n,\max} \sim n^{1/\delta}. \quad (8)$$

Although we will not be concerned with a careful study of critical exponents here, in some cases estimates for such exponents are very important to locate the transition points. In particular, the crossover exponents ϕ_b and $1/\delta$ are crucial if we want to determine $T_{x,\infty}$ from finite-size estimates $T_{x,n}$, for $x = c$ or a , since

$$T_{x,n} \simeq T_{x,\infty} + n^{-\psi}, \quad (9)$$

where ψ is equal to ϕ_b (when $x = c$) or $1/\delta$ (when $x = a$).

Some metric quantities, such as the mean squared end-to-end distance, R_n^2 , are also important in the study of both bulk and surface transitions. For our simulations we created a triangular lattice by augmenting a square lattice with diagonals, and a simple linear transform implies that the parallel and perpendicular components of R_n^2 with respect to the surface are given by

$$R_{\perp,n}^2(\omega_2, \omega_3, \kappa) = \frac{3}{4} \langle y_n^2 \rangle, \quad (10)$$

$$R_{\parallel,n}^2(\omega_2, \omega_3, \kappa) = \langle x_n^2 \rangle + \frac{1}{4} \langle y_n^2 \rangle + \langle x_n y_n \rangle. \quad (11)$$

Close to the adsorption such quantities are expected to follow

$$R_{\perp/\parallel,n}^2 \sim n^{2\nu_{\perp/\parallel}} g(\tau n^{1/\delta}), \quad (12)$$

where $g(\cdot)$ is a scaling function and $\nu_{\perp/\parallel}$ are the Flory exponents. Clearly the scaling of $R_n^2 = R_{\perp,n}^2 + R_{\parallel,n}^2$ is dominated

by the scaling of the larger quantity on the right-hand side, and thus has an associated Flory exponent $\nu = \max(\nu_{\perp}, \nu_{\parallel})$. In nonadsorbed (bulk) phases, such exponents are equal (i.e., $\nu_{\parallel} = \nu_{\perp}$) in the thermodynamic limit, but in finite systems one finds effective exponent estimates $\nu_{\perp/\parallel,n}$ for which $\nu_{\parallel,n} < \nu_{\perp,n}$. In adsorbed phases, on the other hand, the trails behave as quasi-one-dimensional walks, so that $\nu_{\parallel,n} \rightarrow 1$ and $\nu_{\perp,n} \rightarrow 0$. Therefore, in finite trails these exponents cross at some temperature $T_{a,n}$, located in between the nonadsorbed and adsorbed phases. These crossing temperatures can thus be used as a finite-size estimate of the adsorption transition temperature.

We can also use the exponent ν to locate bulk transitions such as the coil-globule transition. In two dimensions, in the coil phase ν assumes the value of $3/4$ [32], while in the globule phase a value $\nu = 1/2$ [32] is expected. For finite systems, however, we have that $\nu_{\text{coil},n} \rightarrow 3/4$ from below and $\nu_{\text{globule},n} \rightarrow 1/2$ from above as the length n increases, so that curves of ν_n against T for different values of n cross at some intermediate temperature, which can be taken as an estimate of $T_{c,n}$. At criticality, we expect to find $\nu = 4/7$ [7]. In a similar fashion, we can locate the collapse transition by using the scaling of the partition function

$$Z_n \sim \mu^n n^{\gamma - 1}, \quad (13)$$

where μ is the connective constant and γ the entropic exponent. Similarly to ν , curves of $\gamma \times T$ for different lengths n should cross each other in between the coil and globule phases, giving also estimates of $T_{c,n}$.

III. NUMERICAL SIMULATIONS

To build up the full phase diagram of the model for a broad range of parameters, we sample SATs with the flatPERM algorithm [33]. Similarly to the pruned-enriched Rosenbluth method (PERM) [34], flatPERM is an improvement of the classical Rosenbluth algorithm. In both flatPERM and PERM, a SAT is stochastically grown with pruning and enrichment strategies intended to prevent its attrition and the appearance of rare configurations with small or extremely large statistical weights. In the flatPERM, the pruning and enrichment processes are implemented independently of the thermodynamic parameters of the system, so that the method gives us an estimate of the density of states $C_{\{m_2, m_3, m_s\}}^{(n)}$. PERM, on the other hand, gives us an estimate of the partition function Z_n for a given set of parameters.

The dimensionality of the density of states (i.e., the number of energy parameters in the model, which is three here) is the main factor limiting the trails' sizes studied with flatPERM. For instance, by not fixing any of the model parameters (ω_2 , ω_3 , and κ), in principle we can numerically obtain the full density of states and from this the entire thermodynamic behavior of the model. It turns out, however, that with this three-parameter approach we are not able to obtain accurate estimates of $C_{\{m_2, m_3, m_s\}}^{(n)}$ for lengths larger than $n \approx 64$ steps in a reasonable amount of time. We can overcome this limitation by fixing one of the parameters and running a two-parameter flatPERM simulation for several slices of the full parameter space. In the following, we will use this strategy to build up the finite-size phase diagram of the model for trails with up

to $n = 128$ steps. To extrapolate from these relatively short finite-size results to the phase diagram in the thermodynamic limit, we perform one-parameter flatPERM simulations (by keeping two parameters fixed) for lengths up to $n = 1024$ for values of particular interest. We also ran PERM simulations for $n \leq 4096$ with all parameters fixed. In all cases, results from averages of at least of 10^9 trails are presented.

IV. RESULTS FOR THE BOUNDARY PLANES

As implicit in the model definition above, we are interested here in investigating only the most physically interesting situation where all interactions are attractive. This means that we will explore the three-parameter space $(\omega_2, \omega_3, \kappa)$ where the three Boltzmann weights are larger than or equal 1. To begin, it is interesting to analyze the three planes forming the boundary of such a space. Namely, the planes $(\omega_2, \omega_3, 1)$, $(1, \omega_3, \kappa)$, and $(\omega_2, 1, \kappa)$. These planes correspond to the absence of surface, two-body, and three-body interactions, respectively.

A. Plane $(\omega_2, \omega_3, 1)$

First we investigate the case where the surface is inert, that is, it does not interact with the polymer. This should be directly comparable to the bulk case previously studied without a surface [9]. To determine the thermodynamic behavior of this plane, we analyze the density of states $\sum_{m_s} C_{(m_2, m_3, m_s)}$ obtained from a two-parameter flatPERM simulation at $\kappa = 1$ for $n = 128$. We calculate the covariance matrix

$$\begin{bmatrix} \langle m_2^2 \rangle - \langle m_2 \rangle^2 & \langle m_2 m_3 \rangle - \langle m_2 \rangle \langle m_3 \rangle \\ \langle m_3 m_2 \rangle - \langle m_3 \rangle \langle m_2 \rangle & \langle m_3^2 \rangle - \langle m_3 \rangle^2 \end{bmatrix}, \quad (14)$$

whose maximum eigenvalue λ is a measure of the fluctuations, which can be plotted against ω_2 and ω_3 as a density plot. In general, local maxima in such fluctuation maps indicate the possible existence of phase boundaries there, so that we can associate lines of suitably defined maxima with pseudotransition lines, for the finite system. Then, by considering other quantities such as the distributions of monomer contacts, we can identify the phases present in each region of the fluctuation map, and also whether the pseudotransition lines can be associated with discontinuous or continuous phase transitions. This method allows us to build a “finite-size phase diagram,” which, in most cases, captures the main features of the actual phase diagram in the thermodynamic limit. We caution that this method does not necessarily pick up higher-order phase transitions and would also identify smooth crossovers as transitions. Hence scaling in length needs to be performed to conclusively identify the presence of phase transitions.

The fluctuation map along with approximate transition lines (i.e., the finite-size phase diagram) for the plane $(\omega_2, \omega_3, 1)$ is shown in Fig. 3. Distributions of doubly and triply visited sites [not shown] reveal, for small values of ω_2 and ω_3 , the presence of configurations typically dominated by a high number of singly visited sites, characteristic of the swollen coil phase. In the region of small ω_3 and large ω_2 , we find configurations with a large number of doubly visited sites, which may be identified as the trails’ collapsed globule phase.




FIG. 3. Fluctuation map for the plane $(\omega_2, \omega_3, 1)$. The lighter (darker) colors indicate regions of larger (smaller) fluctuations. The lower (higher) solid lines are approximations for the continuous coil-globule and crystal-globule transition lines, while the dashed line is the discontinuous coil-crystal transition line.

Conversely, for small ω_2 and large ω_3 the trails are dominated by triply visited sites. In this region the configurations are maximally dense, so that we will refer to this phase as a crystal phase [9]. (More correctly, one should think of this phase as crystal-like, as it has a nonvanishing entropy.)

Along the line of maximal fluctuation separating the coil-globule and globule-crystal phases the distributions have a single peak, indicating the existence of continuous transitions there. Between the coil and crystal phases, on the other hand, the distributions of triply visited sites are double-peaked, suggesting that the coil-crystal transition is discontinuous. It is important to remark that it is difficult to accurately determine the transition lines when the regions of maximal fluctuation separating different phases become close to each other. For instance, in Fig. 3, we can identify only a region where the coil-crystal line change to a globule-crystal one and due to the proximity of this region with the coil-globule line, it is reasonable to infer that exists a multicritical point there, where the three lines meet. As prefigured above, such phase behavior is very similar to the one reported in [9], for the case where the surface is absent, where indeed such lines meet at a multicritical point. This is quite expected since the presence of an inert surface should not change the thermodynamic behavior of such systems.

To verify the reliability of these results for small trails (with $n = 128$ steps), we perform a finite-size analysis considering trails with lengths up to $n = 1024$. The bulk transitions are investigated, in most cases, through the fluctuations c_j in the number of doubly ($j = 2$) and triply ($j = 3$) visited sites. As an example, curves of c_3 versus ω_3 are depicted in Fig. 4(a), for $\omega_2 = 1$ and different lengths. From the maximum in such curves, $c_{3, \max}^{(n)}$, and the scaling behavior from Eq. (5), we estimate the crossover exponent ϕ_b , which is slightly larger than 1 in this case, consistently with the strong increase




FIG. 4. (a) Fluctuation in the number of triply visited sites c_3 versus ω_3 , for $\omega_2 = \kappa = 1$ and different lengths. The open black circles indicate the maximum of each curve. In the insert the values of $\omega_{3,n}$ at the maxima are plotted against $1/n$. The dashed line is a linear fit used in the extrapolation. (b) Effective exponents ν_n as function of ω_2 , for $\omega_3 = \kappa = 1$ and several lengths. The dashed vertical line indicates the asymptotic value estimated from the crossing points.

of fluctuations observed in Fig. 4(a). Then we identify the values of ω_3 at each maximum as the finite-size transition parameter and by assuming that they follow Eq. (9), with the ϕ_b just estimated, we extrapolate such values to obtain the transition point in the thermodynamic limit [see the insertion in Fig. 4(a)]. For $\omega_2 = 1$, this yields $\omega_3 = 6.91(5)$, which is on the locus of the coil-crystal transition. Similar to the findings for $n = 128$ around the transition point the distribution of the number of triply visited sites still exhibits a bimodal behavior for the long trails, which together with the large ϕ_b exponent indicates that such transition is most likely discontinuous.

By increasing ω_2 one still finds a very similar behavior until $\omega_2 \lesssim \frac{5}{3}$, confirming the existence of a line of discontinuous coil-crystal transition in the phase diagram (see Fig. 5). Exactly at $\omega_2 = \frac{5}{3}$, we observe a continuous transition at $\omega_3 = 8.37(4)$. All these results are consistent with those reported by Doukas *et al.* [9], where a discontinuous coil-crystal transition ending at a multicritical point located at $(\omega_2, \omega_3) = (\frac{5}{3}, \frac{25}{3})$ was found.




FIG. 5. Phase diagram for the plane $(\omega_2, \omega_3, 1)$. The blue circles indicate the continuous coil-globule transition, the magenta stars the continuous globule-crystal transition, and the cyan triangles the discontinuous coil-crystal transition. The red square is a multicritical point. In all cases the lines are interpolations.

From Fig. 3 and also from the results in Doukas *et al.* [9], we expect to find a coil-globule transition for (relatively) small ω_3 . Following a procedure similar to the one above for the coil-crystal transition, the exponent ϕ_b was estimated from the scaling behavior of $c_{2,\max}^{(n)}$. For example, for $\omega_3 = 1$, one obtains $\phi_b = 0.53(4)$, which is a bit larger than the value for the θ class in two dimensions: $\phi_{b,\theta} = 3/7$ [7]. However, as extensively reported elsewhere [35–37], estimates of ϕ_b from the bulk fluctuations are usually hampered by strong finite-size effects. For instance, by extrapolating the finite-size transition points, $\omega_{2,n}$, obtained from the maximum of $c_2^{(n)}$ curves for $\omega_3 = 1$ using $\phi_{b,\theta}$, as done in Doukas *et al.* [9], we find $\omega_2 \approx 2.6$ in agreement with the value found in [9]. However, curves of $\omega_{2,n}$ versus $n^{-\phi_b}$ do not display a good linear behavior neither for $\phi_b = \phi_{b,\theta}$ nor for $\phi_b = 0.53$, indicating that such estimate of ω_2 is not so reliable. Other evidence of this is the fact that if one uses the maximum in the fluctuations in the number of trimers, instead of dimers [that is, $c_3^{(n)}$ rather than $c_2^{(n)}$] to determine $\omega_{2,n}$, an extrapolated value $\omega_2 \gtrsim 3$ is obtained (for $\omega_3 = 1$).

To obtain a more accurate estimate of the location of the coil-globule transition, we study the crossing of curves of effective exponents ν_n and γ_n versus the appropriate parameter. Fig. 4(b) shows an example of this for $\omega_3 = 1$, where one sees that curves for a broad range of lengths cross each other in a narrow interval of ω_2 , demonstrating that finite-size effects are very small in this measure. In fact, by using the crossing points for pairs of successive lengths as estimates of $\omega_{2,n}$, one observes only a mild dependence with n . Extrapolating such values using Eq. (9), with the exponent $\phi_{b,\theta}$, we obtain $\omega_2 = 3.12(8)$. A very similar value is obtained from an analysis of the crossing points of effective entropic γ exponents, where finite-size corrections are also weak. The same scenario discussed here for $\omega_3 = 1$ is found for larger values of ω_3 , along the entire coil-globule transition line. Since the results coming from the effective exponents are more accurate than those for $c_2^{(n)}$ and somewhat agree with those from $c_3^{(n)}$, we use them




FIG. 6. Fluctuation map for the plane $(1, \omega_3, \kappa)$. The lighter (darker) colors indicate regions of larger (smaller) fluctuations. The solid (black) line is the continuous coil-adsorbed line, while the slanted (indigo) and vertical (blue) dashed lines are the discontinuous crystal-adsorbed and coil-crystal transition lines, respectively.

to determine the correct coil-globule transition line, which is shown in Fig. 5. We notice that the location of this line differs considerably from the one reported in Doukas *et al.* [9]. Regardless of the numerical difference, both line estimates end at the same multicritical point, limiting similar regions of the phase diagram. Moreover, our analysis of exponents confirms the claim in [9] that this is a tricritical θ line, since the values of ν and γ at the crossing points are found to be reasonably close to the θ exponents.

Finally, for large values of ω_3 we find a continuous crystal-globule transition line [see Fig. 5], which was determined following the same procedure employed above using the maxima of fluctuations. We notice that, similarly to the coil-crystal transition, here results from $c_2^{(n)}$ and $c_3^{(n)}$ are consistent. However, once again the locus of our transition line is shifted from that reported in [9], resulting from differing finite-size analyses. Importantly, similar to the phase diagram from [9] our line seems to start at the multicritical point and extends to large values of ω , as also suggested by the fluctuation map for $n = 128$. Actually, despite their quantitative differences, the entire “finite-size diagram” from the fluctuation map in Fig. 3 is consistent with the phase diagram given here in Fig. 5.

B. Plane $(1, \omega_3, \kappa)$

Now we investigate the case where the two-body monomer-monomer interaction is absent, so that $\omega_2 = 1$. Similar to the plane $(\omega_2, \omega_3, 1)$, we will start analyzing the properties of the fluctuation map and the respective finite-size phase diagram, for $n = 128$. This is shown in Fig. 6, where one clearly sees three regions separated by stripes of large fluctuations. The distributions of number of monomers reveal that such regions are associated with the phases: coil (for small κ and ω_3), crystal (for small κ and large ω_3),




FIG. 7. Phase diagram for the plane $(1, \omega_3, \kappa)$. The cyan triangles indicate the discontinuous coil-crystal transition, the indigo circles the discontinuous crystal-adsorbed transition, and the black stars the continuous coil-adsorbed transition. The red circle represents the critical-end-point. All lines are guide-to-eye.

and adsorbed (for large κ). This last one is characterized by a high number of contacts with surface. An analysis of the modality of the distribution for the number of trimers points out that the coil-crystal and crystal-adsorbed phases are separated by discontinuous transitions, while a continuous transition line seems to exist between coil and adsorbed phases. As it is well known [14], when the strength of the polymer-surface interaction is not so large, the bulk transitions should not be affected by such interaction. This means that the discontinuous coil-crystal transition is expected to happen at a straight line parallel to the κ axis in the (κ, ω_3) space. This important observation facilitates the determination of the phase boundaries in the finite-size diagram. The maxima of fluctuations in Fig. 6 suggest that the three transition lines meet at single point, which turns out to be a critical end point (CEP).

Let us now explore the phase behavior in the thermodynamic limit. To determine and confirm the existence of the (bulk) coil-crystal transition line, we follow the same procedures described in the previous subsection. This yields an approximately vertical (κ -independent) line in the phase diagram, located at $\omega_3 \approx 6.93$ (see Fig. 7). This coil-crystal coexistence line ends at $\kappa \approx 2.08$, above which it gives place to a crystal-adsorbed discontinuous transition. The crystal-adsorbed coexistence line, which was determined in the same way of the coil-crystal one, increases monotonically with ω_3 , as seen in Fig. 7.

To analyze the (presumed continuous) coil-adsorbed transition, one starts calculating Γ_n [Eq. (7)] and using the maxima in their curves to estimate the crossover exponent $1/\delta$ from the scaling relation 8. We always find $1/\delta \approx 1/2$ along this line, in agreement with the expected value for the adsorption transition in two dimensions [38]. Then, we use the position of the maxima of Γ_n as the finite-size transition estimate κ_n and extrapolate them according to Eq. (9) with $1/\delta = 1/2$. An




FIG. 8. (a) Γ_n for different lengths as function of κ , for $\omega_2 = \omega_3 = 1$. The circles indicate the maximum in each curve. The insertion shows the values of κ_n at the maxima against $n^{-1/2}$. The dashed line is a linear fit used in extrapolation. (b) Effective Flory exponents $\nu_{\parallel,n}$ (dashed lines) and $\nu_{\perp,n}$ (solid lines) versus of κ , for $\omega_2 = 1$ and $\omega_3 = 2$ and several lengths.

example of this is shown in Fig. 8(a), for $\omega_3 = \omega_2 = 1$, for which one obtains $\kappa = 2.113(5)$.

The other accurate way of locating the adsorption transition is through the crossing of the Flory exponents ν_{\parallel} and ν_{\perp} [24]. Figure 8(b) presents an example for $\omega_3 = 2$. The crossing points for different lengths serve also as good estimates of κ_n and their extrapolation [with Eq. (9) and $1/\delta = 1/2$] yields an estimate for the thermodynamic value for the adsorption point. Though this method returns results similar to those from fluctuations, it has weaker finite-size effects, being so more reliable to determine the adsorption transition line. Such a line is also depicted in the phase diagram shown in Fig. 7, where one sees that it ends at the point where the coexistence lines meet, confirming the existence of a CEP in the diagram at $(\omega_3, \kappa) \approx (6.93, 2.08)$, as suggested by the finite-size fluctuation map.

C. Plane $(\omega_2, 1, \kappa)$

Next we discuss the case where trimers at a given site do not interact, so $\omega_3 = 1$. The fluctuation map and the finite-size




FIG. 9. (a) Fluctuation map for the plane $(\omega_2, 1, \kappa)$. The vertical blue line indicates the coil-globule transition. The approximately horizontal line to the left (right) of the coil globule transition is the black (green) line being the coil-adsorbed (globule-adsorbed) transition line. The transitions in all these lines seem to be continuous. While the adsorbed phase is a single phase (there are no finite temperature phase transitions) it has two regions where the ground state differs. Illustrations of the two different ground state configurations are shown in (b) for the Ad₁ region and (c) Ad₂ region.

transition lines are displayed in Fig. 9(a). Using the same methodology previously discussed, we find for small values of κ indications of a continuous coil-globule transition. For large values of κ one has signatures of two adsorbed regions. The ordinary adsorbed region (characterized, in the ground state, by a single line of monomers lying on the surface [see Fig. 9(b)]) is found for small ω_2 . For large ω_2 , on the other hand, we observe an adsorbed region which, in the ground state, is featured by a bilayer of doubly visited sites, as illustrated in Fig. 9(c). Hereafter, we will refer to these monolayer and bilayer regions as Ad₁ and Ad₂ regions, respectively. Importantly below our analysis indicates that there are no finite-temperature transitions between them, so they are *not* thermodynamically stably different phases.

Our finite-size analysis indicates that both bulk (coil and globule) phases adsorb through a continuous transition. While the globule adsorbs to the Ad₂ region, the coil phase adsorbs to either an Ad₁ or Ad₂ region, depending on the value of ω_2 . The globule adsorption to the Ad₂ region can be understood, from an energetic “cost” point of view since both feature a relatively large proportion of doubly visited sites. By locating the phase boundaries in the finite-size phase diagram, we found indications of a possible multicritical point located where all the three continuous lines, coil-globule, coil-adsorbed, and globule-adsorbed meet.

In the fluctuation map, we see a clear region with large fluctuation between the regions Ad₁ and Ad₂, suggesting the possible existence of an Ad₁-Ad₂ transition. As already remarked, however, from the finite behavior alone we cannot determine whether this is a phase transition or simply a smooth crossover. To check this, we performed PERM calculations for trails with up to $n = 4096$ steps, for several values of κ and $\omega_2 = 2.4$. We chose this value of ω_2 since here the coil adsorbs to an Ad₂ region at the transition. Moreover, for very large values of κ the adsorbed phase becomes Ad₁.




FIG. 10. (a) Γ_n as function of κ for $\omega_3 = 1$, $\omega_2 = 2.4$, and several lengths. (b) Fluctuation in the number of doubly visited sites $c_2^{(n)}$ versus κ for the same parameters as in (a).

Figures 10(a) and 10(b) present Γ_n and $c_2^{(n)}$, respectively, from such simulations, verifying this scenario. There is a single peak in Γ_n around $\kappa \approx 2$, where the coil-adsorbed transition is expected. There is no indication of an Ad_1 - Ad_2 transition, but rather evidence for a smooth crossover.

From a physical point of view, one can understand the two regions simply as two one-dimensional layers, and short-range interactions in one-dimensional systems are not able to induce any finite-temperature phase transition, so that there can only be a smooth crossover between these regions.

Therefore, by using the same methods employed in previous subsections, we obtain the phase diagram depicted in Fig. 11. Once again, it is consistent with our findings from the fluctuation map, with continuous coil-globule, coil-adsorbed, and globule-adsorbed transition lines. Similarly to the coil-crystal transition found in the plane $(1, \omega_3, \kappa)$, an approximately vertical (κ -independent) straight line is found separating the coil and the globule phases, as expected. This line, located at $\omega_2 \approx 3.2$, starts at $\kappa = 1$ and ends at a multicritical point located at $\kappa = 1.96(2)$, where it meets the coil-adsorbed and globule-adsorbed lines. It is interesting to note in Fig. 11 that the globule-adsorbed line appears for lower values of κ than those at the crystal-adsorbed line (see Fig. 7). This shows that trails in the globule phase adsorb easier than those in the




FIG. 11. Phase diagram for the plane $(\omega_2, 1, \kappa)$. The black stars indicate the continuous coil-adsorbed transition, the blue circles the θ -like coil-globule transition, and the green diamonds the continuous globule-adsorbed transition. The multicritical point is denoted by the red circle. All lines are guide-to-eye.

crystal phase. One way to understand this is that due to the presence of the Ad_2 region in the plane $(\omega_2, 1, \kappa)$ the globule phase adsorbs to a state in which the trails can retain their large number of doubly visited sites.

V. THREE-PARAMETER MODEL

To get some insight on the phase behavior in the full three-parameter space, it is instructive to begin by synthesizing the information from the three boundary planes discussed in the previous section. By simply putting these together one finds a picture as depicted in Fig. 12, which indicates the regions where each of the four thermodynamic phases (coil, globule,




FIG. 12. Summary of the phases diagrams for the three boundary planes of three parameter space plotted together. The blue squares are the coil-globule transition lines estimated here. The open cyan squares are the points of discontinuous coil-crystal transitions, while the open green triangles the discontinuous crystal-adsorbed transition. The black circles denote the continuous coil-adsorbed transitions and the dark-green triangles the critical globule-adsorbed transition. The magenta diamonds indicates the continuous globule-crystal transition. The red filled circles represent the CEP and the two multicritical points.




FIG. 13. Fluctuation maps in spaces $(\omega_2, \omega_3, 2)$ and $(\omega_2, \omega_3, 3)$ in (a) and (b), respectively, with transition lines indicated. In both panels solid and dashed lines indicate the presence of unimodal and bimodal energy distributions near the transition, respectively.

crystal, and the adsorbed) found above appears in the full phase space and how they are separated. Of course one is assuming no new phases appear. So for instance, the continuous coil-adsorbed transition lines in the planes $(\kappa, \omega_2, 1)$ and $(\kappa, 1, \omega_3)$ strongly suggest that a critical surface exists separating these phases. Similarly, the tricritical coil-globule lines in the planes $(\kappa, \omega_2, 1)$ and $(\omega_2, \omega_3, 1)$ indicate that such phases are separated by a tricritical (possibly θ) surface. Moreover, the discontinuous transition lines between the coil and crystal phases [in the planes $(\kappa, 1, \omega_3)$ and $(\omega_2, \omega_3, 1)$] provide strong evidence on the existence of a coil-crystal coexistence surface.

As remarked above the full three-parameter flatPERM simulations could only be undertaken for length less than $n \approx 64$ to a reasonable degree of convergence. As such we investigated the full phase diagram based on the finite-size behavior of two-parameter slices of the three-parameter space, for $n = 128$. We remark that even such two-parameter flatPERM approach has demanded extensive simulations. As already shown in the previous section, these finite-size phase diagrams (built up from the fluctuation maps) capture the correct thermodynamic properties of the system, though quantitatively incorrect in most cases. Indeed we found no evidence of new phases, beyond the four ones already discussed. Moreover, since there is only a crossover between the adsorbed Ad_1 rich and Ad_2 rich regions, no transition line separating them will be shown, even if we indicate their locations in the figures.

A. κ slices

Let us start analyzing slices for κ fixed. We recall that for $\kappa = 1$ [i.e., the plane $(\omega_2, \omega_3, 1)$ considered in Sec. IV A] the three bulk phases (coil, globule, and crystal) are present in the system, as shows Figs. 3, 5, and 12. Since, for small values of κ , the bulk transitions are not expected to be affected by the presence of an interacting surface, the same diagram of $\kappa = 1$ is expected in this region. In fact, as demonstrated in Fig. 13(a), the finite-size phase diagram for $\kappa = 2$ is practically identical to the one for $\kappa = 1$ shown in Fig. 3. This strongly indicates that the transition lines between the three bulk phases give rise to three surfaces: a tricritical coil-globule, a critical crystal-globule, and a coexistence

coil-crystal, consistently with the predictions above based on Fig. 12. Moreover, the multicritical point found in the case $\kappa = 1$ gives rise to a multicritical line, where the three surfaces above meet. Since this line is related to bulk phases only, we will refer to it as the bulk multicritical line (BML). Note that these bulk transition surfaces, as well as the BML are all expected to be κ -independent, i.e., perpendicular to the (ω_2, ω_3) plane. This is indeed confirmed by the quantitative agreement between the finite-size transition lines in Figs. 3 and 13(a) for $\kappa = 1$ and 2, respectively, as well as for intermediate κ slices (not shown).

All these bulk transition surfaces (and the BML, as well) are expected to end when they meet the adsorption transition surfaces. Figures 7, 11, and 12 indicate that the coil-adsorbed and the globule-adsorbed critical surfaces in the thermodynamic limit change mildly with ω_2 and ω_3 , being located close to $\kappa = 2$. The same thing happens in the diagrams for $n = 128$, as seen in Figs. 6 and 9(a), where such surfaces are in the region $2 \lesssim \kappa \lesssim 3$. Therefore, it is hard to observe these adsorption surfaces in planes of fixed κ . Anyhow, as shown in Fig. 13(b), for the $\kappa = 3$ slice, the only bulk phase appearing in this diagram is the crystal one, confirming that we are above the coil-adsorbed and globule-adsorbed surfaces. Interestingly, in the region for small ω_2 (where the coil phase was observed for smaller κ) one finds the adsorbed phase is Ad_1 rich, while for large values of ω_2 the adsorbed phase is Ad_2 rich.

At finite length $n = 128$, as shown in the diagram for $\kappa = 3$ [Fig. 13(b)], the crystal-adsorbed transition is still found to be discontinuous for small ω_2 , as evidenced by a bimodal energy distribution. As discussed above, there is only one adsorbed phase, so that one would expect this transition to remain discontinuous for all values of ω_2 . Unfortunately, we are unable to detect a bimodal distribution for large ω_2 but believe that this is simply a finite size effect.

B. ω_2 slices

Now we investigate planes for ω_2 fixed. As already demonstrated in Sec. IV B, for $\omega_2 = 1$ three phases are present in the diagram: adsorbed, coil, and crystal (see Fig. 6, for $n = 128$). Figure 14(a) displays the behavior for $\omega_2 = 1.5$, which is the same for $\omega_2 = 1$, confirming the existence of a continuous coil-adsorbed transition surface, as well as of discontinuous coil-crystal and crystal-adsorbed ones. Moreover, since the former critical surface seems to end at its junction with the latter two coexistence ones, they shall form a line of critical-end-points (a CEP line) there.

The slice $\omega_2 = 2$ is shown in Fig. 14(b), where a different thermodynamic behavior is observed. According to Figs. 3 and 13, in such plane, with κ small, one should observe transitions from coil to globule and then from globule to crystal by increasing ω_3 . It turns out, however, that the fluctuation map displayed in Fig. 14(b) fails in capturing the coil-globule transition, although for very small ω_3 the trails have the characteristics of the coil phase. This caveat is certainly due to the short lengths considered. For large values of κ , the expected adsorbed phase is observed.

We note that for the crystal-adsorbed transition at $\omega_2 = 2.0$ we were unable to detect a bimodal distribution. While




FIG. 14. Fluctuation maps in spaces $(1.5, \omega_3, \kappa)$ and $(2.0, \omega_3, \kappa)$ in (a) and (b), respectively, with transition lines indicated. In all panels solid and dashed lines indicate the presence of unimodal and bimodal energy distributions near the transition, respectively.

consistent with the numerical evidence in the κ slices we reiterate that we expect this to be a finite-size effect.

For fixed ω_2 , the globule-crystal transition line meets the adsorbed phase at a dense-adsorbed multicritical point, which in the full phase diagram extends to a dense-adsorbed multicritical (DAM) line. Such a line seems also to join the CEP and the BML lines at a special multicritical point.

C. ω_3 slices

We now present the results for planes with ω_3 fixed. Fig. 15(a) presents the diagram for the $\omega_3 = 8$ slice, which is qualitatively identical to the one for $\omega_3 = 1$, see Fig. 9(a); slices for intermediate values of ω_3 (not shown) display the same behavior. This gives additional confirmation of the existence of a tricritical coil-globule and of critical coil-adsorbed and globule-adsorbed surfaces. These three surfaces meet, yielding a third multicritical line in the three-parameter space, which we will refer to as the collapsed-adsorbed multicritical (CAM) line. We expect that this CAM line also ends at the special multicritical point where the DAM, CEP, and BML lines meet.




FIG. 15. Fluctuation maps in spaces $(\omega_2, 8, \kappa)$ and $(\omega_2, 12, \kappa)$ in (a) and (b), respectively, with transition lines indicated. In both panels solid and dashed lines indicate the presence of unimodal and bimodal energy distributions near the transition, respectively.




FIG. 16. Qualitative representation of the full phase diagram, presenting the four phases found (regarding the regions Ad₁ and Ad₂ simply as the adsorbed phase), the critical-end-point (CEP) line, as well as the bulk (BML), collapsed-adsorbed (CAM), and dense-adsorbed (DAM) multicritical lines. The special multicritical point is denoted by a circle.

As expected from Figs. 3, 6, and 12, for large values of ω_3 the crystal phase appears in place of the coil phase for small values of ω_2 and κ . For $\omega_3 = 12$, this is demonstrated in Fig. 15(b). We are unable to detect a bimodal energy distribution along the whole crystal-adsorbed transition line, but attribute this to finite-size effects as discussed above.

VI. SUMMARY: FULL PHASE DIAGRAM

Our simulation results for self-attracting self-avoiding trails living on the triangular lattice and attractively interacting with a surface reveals a very rich thermodynamic behavior, with three bulk phases (coil, globule, and crystal) and an adsorbed phase with two regions related to different ground states (the ordinary monolayer Ad₁ rich region and a bilayer Ad₂ rich region). The coil phase exists in a bounded part of the full phase diagram where the three Boltzmann weights ω_2 , ω_3 , and κ are small. The crystal phase appears in the region where κ and ω_2 are relatively small, but ω_3 is large. Similarly, the globule phase exists for small κ and large ω_2 . Finally, the adsorbed phase appear in the region of large κ . These features are summarized in the sketch of the full phase diagram depicted in Fig. 16, where no distinction is made between both adsorbed regions since we determined that there is only a smooth crossover between them.

Three bulk transition surfaces are present in the system, for small κ : the continuous coil-globule (possibly in θ class [9]) and globule-crystal surfaces and a discontinuous coil-crystal surface. These three surfaces meet at the multicritical BML line, which seems to end close to $\kappa = 2$. We remark that the coordinates of such line in the plane (ω_2, ω_3) with $\kappa = 1$, i.e., in the absence of the surface interaction, is exactly known to be $(\omega_2, \omega_3) = (\frac{5}{3}, \frac{25}{3})$ [9]. Hence, by assuming that the bulk transitions are not affected by the surface interaction when κ is small, the BML line should end at a special multicritical point located at $(\omega_2 = \frac{5}{3}, \omega_3 = \frac{25}{3}, \kappa \approx 2)$.

Beyond the three bulk surfaces, there are also several adsorption transition surfaces in the full phase diagram, which separate the bulk phases from the adsorbed ones. The coil-adsorbed and the globule-adsorbed are continuous, and they

meet the tricritical coil-globule surface at the multicritical CAM line. Part of the critical coil-adsorbed surface ends at the coexistence surfaces separating the crystal phase from the coil and adsorbed ones, so that a CEP line exists there. Moreover, the globule-crystal, globule-adsorbed, and crystal-adsorbed surfaces meet at another multicritical DAM line. Therefore, there are three multicritical (BML, CAM, and DAM), and a CEP line in the phase diagram. Our results for finite trails indicate that all these lines meet at the special multicritical point discussed above.

ACKNOWLEDGMENTS

T.J.O. and N.T.R. acknowledge support of the Brazilian agencies CNPq, CAPES, and FAPEMIG, and the use of the computing cluster of Universidade Federal de Viçosa. N.T.R. thanks Queen Mary University London for hosting while this work was begun, and gratefully acknowledges use of the university's HPC cluster. Financial support from the Australian Research Council via its Discovery Projects scheme (DP160103562) is acknowledged by A.L.O.

-
- [1] J. des Cloizeaux and G. Jannink, *Polymers in Solution: Their Modelling and Structure* (Clarendon, Oxford, 1990).
- [2] P.-G. De Gennes, *Scaling Concepts in Polymer Physics* (Cornell University Press, Ithaca, NY, 1979).
- [3] P.-G. De Gennes, *J. Physique Lett.* **36**, 55 (1975).
- [4] M. Stephen, *Phys. Lett. A* **53**, 363 (1975).
- [5] B. Duplantier, *J. Phys. France* **43**, 991 (1982).
- [6] B. Duplantier, *Europhys. Lett.* **1**, 491 (1986).
- [7] B. Duplantier and H. Saleur, *Phys. Rev. Lett.* **59**, 539 (1987).
- [8] B. Duplantier, *J. Chem. Phys.* **86**, 4233 (1987).
- [9] J. Doukas, A. L. Owczarek, and T. Prellberg, *Phys. Rev. E* **82**, 031103 (2010).
- [10] A. Bedini, A. L. Owczarek, and T. Prellberg, *Phys. Rev. E* **86**, 011123 (2012).
- [11] A. Bedini, A. L. Owczarek, and T. Prellberg, *J. Phys. A: Math. Theor.* **49**, 214001 (2016).
- [12] E. Eisenriegler, K. Kremer, and K. Binder, *J. Chem. Phys.* **77**, 6296 (1982).
- [13] K. De' Bell and T. Lookman, *Rev. Mod. Phys.* **65**, 87 (1993).
- [14] T. Vrbová and S. G. Whittington, *J. Phys. A: Math. Gen.* **29**, 6253 (1996).
- [15] T. Vrbová and S. G. Whittington, *J. Phys. A: Math. Gen.* **31**, 3989 (1998).
- [16] T. Vrbová and K. Procházka, *J. Phys. A: Math. Gen.* **32**, 5469 (1999).
- [17] P. Grassberger, *J. Phys. A: Math. Gen.* **38**, 323 (2005).
- [18] A. L. Owczarek, A. Rechnitzer, J. Krawczyk, and T. Prellberg, *J. Phys. A: Math. Theor.* **40**, 13257 (2007).
- [19] M.-B. Luo, *J. Chem. Phys.* **128**, 044912 (2008).
- [20] L. I. Klushin, A. A. Polotsky, H.-P. Hsu, D. A. Markelov, K. Binder, and A. M. Skvortsov, *Phys. Rev. E* **87**, 022604 (2013).
- [21] N. T. Rodrigues, T. Prellberg, and A. L. Owczarek, *Phys. Rev. E* **100**, 022121 (2019).
- [22] J. A. Plascak, P. H. L. Martins, and M. Bachmann, *Phys. Rev. E* **95**, 050501(R) (2017).
- [23] P. H. L. Martins, J. A. Plascak, and M. Bachmann, *J. Chem. Phys.* **148**, 204901 (2018).
- [24] C. J. Bradly, A. L. Owczarek, and T. Prellberg, *Phys. Rev. E* **97**, 022503 (2018).
- [25] C. J. Bradly, A. L. Owczarek, and T. Prellberg, *Phys. Rev. E* **98**, 062141 (2018).
- [26] I. Carmesin and K. Kremer, *Macromolecules* **21**, 2819 (1988).
- [27] Ch. Sandhya, K. H. Bindu, K. P. N. Murthy, and V. S. S. Sastry, *AIP Conf. Proc.* **1349**, 117 (2011).
- [28] J. Luettmer-Strathmann, F. Rampf, W. Paul, and K. Binder, *J. Chem. Phys.* **128**, 064903 (2008).
- [29] M. A. Moore, *J. Phys. A: Math. Gen.* **10**, 305 (1977).
- [30] R. Brak, A. L. Owczarek, and T. Prellberg, *J. Phys. A: Math. Gen.* **26**, 4565 (1993).
- [31] P.-G. De Gennes, *J. Phys. France* **37**, 1445 (1976).
- [32] P. J. Flory, *Principles of Polymer Chemistry* (Cornell University Press, Ithaca, NY, 1971).
- [33] T. Prellberg and J. Krawczyk, *Phys. Rev. Lett.* **92**, 120602 (2004).
- [34] P. Grassberger, *Phys. Rev. E* **56**, 3682 (1997).
- [35] J. H. Lee, S.-Y. Kim, and J. Lee, *J. Chem. Phys.* **135**, 204102 (2011).
- [36] I. Chang and H. Meirovitch, *Phys. Rev. E* **48**, 3656 (1993).
- [37] H. Meirovitch and H. A. Lim, *J. Chem. Phys.* **91**, 2544 (1989).
- [38] T. W. Burkhardt, E. Eisenriegler, and I. Guim, *Nuc. Phys. B* **316**, 559 (1989).